Structure of the connexin-43 gap junction channel reveals a closed sieve-like molecular gate

Chao Qi[1,2,#], Silvia Acosta-Gutierrez[3,#], Pia Lavriha[1,2], Alaa Othman[4], Diego Lopez-Pigozzi[5,6], Erva Bayraktar[6], Dina Schuster[1,2,4], Paola Picotti[4], Nicola Zamboni[4], Mario Bortolozzi[5,6], Francesco L. Gervasio[7,8,9*] and Volodymyr M. Korkhov[1,2*]

[1] Institute of Molecular Biology and Biophysics, ETH Zurich, Switzerland

[2] Laboratory of Biomolecular Research, Paul Scherrer Institute, Villigen, Switzerland

[3] Institute for the Physics of Living Systems, Institute of Structural and Molecular Biology, and Department of Chemistry. University College London, London, UK

[4] Institute of Molecular Systems Biology, ETH Zurich, Switzerland

[5] Department of Physics and Astronomy "G. Galilei", University of Padova, Padua, Italy

[6] Veneto Institute of Molecular Medicine (VIMM), Padua, Italy

[7] Department of Chemistry, University College London, WC1E 6BT London, UK

[8] School of Pharmaceutical Sciences, University of Geneva, CH-1211 Geneva, Switzerland

[9] ISPSO, University of Geneva, CH-1211 Geneva, Switzerland

* Corresponding authors: francesco.gervasio@unige.ch, volodymyr.korkhov@psi.ch

# These authors contributed equally

Abstract

Gap junction channels (GJCs) mediate intercellular communication by connecting two
neighboring cells and enabling direct exchange of ions and small molecules. Cell coupling via
connexin-43 (Cx43) GJCs is important in a wide range of cellular processes in health and
disease [1-3], yet the structural basis of Cx43 function and regulation has not been determined
until now. Here we describe the structure of a human Cx43 GJC solved by cryo-EM and single
particle analysis at 2.26 Å resolution. The pore region of Cx43 GJC features several lipid-like
densities per Cx43 monomer, located close to a putative lateral access site at the monomer
boundary. We found a previously undescribed closed sieve-like gate on the cytosolic side of
the pore, formed by the N-terminal domain and the transmembrane helix 2 of Cx43 and
stabilized by a small molecule. Structures of the Cx43 GJC and hemichannels in nanodiscs
confirm the sieve-like gate arrangement. The features of the Cx43 GJC and hemichannel cryo-
EM maps and the channel properties revealed by molecular dynamics simulations suggest that
the captured states of Cx43 are closed.

Introduction

Gap junction (GJ) mediated intercellular communication is one of the major pathways of information exchange between the cells. GJs are specialized regions of the plasma membrane at the cell-cell interface that link two adjacent cells and establish their metabolic and electrical coupling [4]. Connexins are the building blocks of the GJ channels (GJCs) which belong to a group of large-pore channels. This group includes a number of structurally related (innexins, pannexins, LRRC8) and unrelated proteins (CALHM) [5]. A total of 21 connexin genes have been identified in the human genome [6]. The 43 kDa connexin-43 (Cx43, gene name *GJA1*) was identified as a major constituent of rat heart GJs in 1987 [7], and it is arguably one of the most extensively studied connexins. Like all connexin homologues, Cx43 monomers assemble into hexameric hemichannels (HCs), also known as connexons. HCs that reach the plasma membrane of one cell interact with their counterparts on the neighboring cell, forming GJCs, typically organized into hexagonal arrays at the intercellular interface [8]. GJCs enable direct metabolic and electric coupling between the cells, facilitating the passage of ions, small molecules, metabolites, peptides and other cellular components below a size threshold of approximately 1.5 kDa [9]. GJCs formed by Cx43 are crucial for a wide range of physiological processes, from propagation of heart action potentials [3] to maintenance of neuro-glial syncytium [2] and skin wound healing [1]. The clinical importance of Cx43 is highlighted by mutations linked to several genetic disorders, such as oculodentodigital dysplasia (ODDD) [10-12], hypoplastic left heart syndrome 1 [13], Hallermann-Streiff syndrome [14], atrioventricular septal defect 3 [13], etc., and by recognition of the protein as a drug target for treatment of cancer, skin wounds, eye injury and inflammation and cardiac arrhythmias (reviewed by Laird and Lampe [9]).

Much of what we know today about the molecular biology, electrophysiological properties and regulation of connexin gap junctions has been derived from the studies on Cx43 (reviewed in [15]). Early attempts to characterize the structure of Cx43 using cryo-electron microscopy (cryo-EM) of 2D crystals produced low resolution reconstructions [16]. More recently, several homologous connexin GJCs and HCs have been structurally characterized at high resolution, using X-ray crystallography and cryo-EM. The structures of Cx26 [17] and Cx46/50 GJCs [18,19], together with the recent structure of Cx31.3 HC [20] provided deep insights into the shared structural features of the connexin channels. These structures hint at the role of the N-terminal domain (NTD) in molecular gating of GJCs and HCs, with several conformations of the NTD observed in different connexin homologues [18-20]. Additionally, biochemical evidence points to the roles in channel gating played by the link between Cx43 intracellular loop and the C-terminal region [21,22]. However, despite the availability of this evidence, the molecular determinants of intracellular connexin channel gating remain unclear. To shed light on the structural basis of Cx43 gating, we set out to determine its structure by cryo-EM and to analyse its dynamics with molecular dynamics (MD) simulations.

Results

Structures of Cx43 GJC and HC in detergent micelles and in nanodiscs. Our Cx43 expression system was tested using electrophysiology in HeLa cells (Figure S1A-D). The experiments confirmed the functionality of Cx43 GJCs expressed in transfected cells. For large-scale protein expression, Cx43 featuring a C-terminal HRV3C-YFP-twinStrep tag, was expressed in adherent mammalian cells (HEK293F) using transient transfection method. The produced protein was purified using affinity and size exclusion chromatography in digitonin (Figure S2A-C) and flash frozen on cryo-EM grids. The grids were subjected to single particle cryo-EM analysis (Figure S3), yielding the final 3D reconstruction at 2.26 Å resolution (Figure 1A, Figure S3-4).

The overall architecture of the protein complex resembles that observed with other connexin
GJCs, with two hemichannels (connexons) from adjacent plasma membrane regions coupled
to form a full GJC (Figure 1A-B). The exterior of the channel is decorated by several ordered
lipid-like density elements (Figure 1A-B), reminiscent of the previously observed lipids bound
at the protein-bilayer interface in other connexins [19]. In the case of Cx43, the lipid-like densities
appear to decorate the protein-lipid bilayer interface at both the inner and the outer leaflet of
the membrane. Comparison of the Cx43 monomers to the available structures of Cx26 and
Cx46 GJCs and the Cx31.3 HC revealed a major difference in the NTD arrangement in these
channels (Figure 1C-D). The conformation of the NTD in the Cx31.3 HC structure appears to
be the closest to that in Cx43 GJC. Analysis of the tryptic peptides revealed that most of the
protein lacks the residue M1 (Figure S5), and thus the model was built starting with G2.

To ascertain that the conformation of Cx43 GJC is not induced by the detergent present in the
sample we removed the detergent and reconstituted the protein into MSP2N2 nanodiscs, using
1-palmitoyl-2-oleoyl-glycero-3-phosphocholine (POPC) as a mimic for the native lipid
environment. The reconstituted protein was subjected to the same imaging and analysis
workflow as Cx43 GJC in digitonin. The 2D classes of the MSP2N2-reconstituted Cx43
showed features consistent with a mixture of GJCs and HC. Processing of the corresponding
particles resulted in 3D reconstructions of the Cx43 GJC at 2.95 Å resolution (Figure S6, S8)
and HC in nanodiscs at 3.98 Å resolution (Figure 1E-F, Figure S7, S8). The GJCs in detergent
and in nanodiscs are nearly identical, with an RMSD of 0.97 Å between the aligned Cx43
monomers (Figure 1G, Figure S9A). Although the differences between the HC in nanodisc and
the GJC are more pronounced (Figure S9A), the conformation of the cytosolic region and the
NTD is highly conserved (Figure 1G).

Sieve-like gate conformation in Cx43. The reconstructed gate region of Cx43 shows several
narrow openings connecting the pore vestibule and the pore interior: a single ~6-7 Å wide
central gate (CG) and six adjacent peripheral gates (PG) of similar dimensions (Figure 2A-B).
This sieve-like arrangement of the gate region is distinct from any of the previously observed
states of connexin GJCs or HCs. This particular gate conformation is established through an
interplay of two structural elements: the NTD and the TM2 (Figure 2C-D). The NTD of Cx43
is arranged near parallel to the membrane plane, in a centre-oriented conformation (Figure 2C).
The HC of Cx31.3 assembles in a similar manner, with a well-ordered NTD resolved by cryo-
EM (Figure 2C-D). However, the TM2 region of Cx31.3 forms a tight seal with the NTD
(Figure 2E). In contrast, the TM2 of Cx43 is shifted away from the pore centre, creating six
PG openings (Figure 2F). This arrangement of the NTD appears to be stabilized by a small
molecule: a density element likely corresponding to a bound small molecule is present within
the PG region, wedged between the adjacent NTDs (Figure 2G). Thus, although the
conformation of the Cx43 gate is sieve-like, the PG is blocked by a yet unidentified small
molecule (which we refer to as the PG lipid, or PGL).

In addition to the PGL molecules stabilizing the NTD arrangement, the cryo-EM density of
Cx43 GJC features several well resolved densities inside the pore region (lipid-1 and lipid-2,
Figure 2H). These elements likely correspond to bound sterol molecules, such as cholesterol
co-purified with the protein from the mammalian cells or cholesterol hemisuccinate (CHS)
added to the solubilization mixture during protein extraction from the membrane. Similar
densities are present in the Cx43 GJC reconstruction in nanodiscs, in the absence of detergent
molecules (Figure S10A).

Identification of the lipid-like molecules bound to Cx43 is a significant challenge. Despite the
high overall resolution of the 3D reconstruction of the Cx43 GJC (2.24 Å resolution in
detergent and 2.95 Å in nanodisc; Figure S3, S6), the cryo-EM map features in regions
corresponding to the lipid densities are insufficient to assign the lipid identity unambiguously.

To determine the identities of these lipids experimentally, we performed lipidomic analysis of
the organic extracts prepared from the purified Cx43 samples and compared them to the mock
control (extracts of prepared from eluates of the purification procedure using cells that do not
overexpress the protein as starting material). The results showed that several types of
phospholipids, as well as cholesterol were present in both samples. The only lipid-like
compound specifically enriched in the purified sample was dehydroepiandrosterone (DHEA;
Figure S10B). DHEA is a most abundantly expressed neurosteroid known to modulate ligand-
gated ion channels, such as GABA$_A$ [23] or NMDA and AMPA receptors [24]. Although we are
presently not able to conclusively state that DHEA corresponds to any of the densities present
in our reconstruction, its enrichment in the sample is an interesting observation that deserves
detailed future investigations.

Cx43 junction interface. As in other connexins, the Cx43 GJC interface is formed by two
extracellular loops, ECL1 and ECL2 (Figure 3A-B). The portion of the ECL1 comprising
residues N55-Q58 makes contacts with two Cx43 monomers from the opposite membrane,
with N55-Q58 of one monomer and Q58'-P59' of the neighboring monomer within 4 Å
distance (Figure 3C, *top left*). The ECL2 region that directly participates in sealing the junction
includes residues P193-D197. Comparisons of the junction-forming loops in Cx43 to those in
Cx46 (Figure 2C, *middle*) and Cx26 (Figure 2C, *right*) show that each protein uses a similar
pattern of inter-HC interaction within each GJC. It is noteworthy that both the amino acid
sequences (Figure 3D) and the interaction networks (Figure 3C) are similar in the ECL1 for
Cx43 and Cx46, and in the ECL2 for Cx46 and Cx26. The differences in these two loops, and
especially in ECL2, underlie the inability of Cx43 (which has been grouped into a docking
group 2) to engage in heterotypic interactions with group 1 connexins (such as Cx46 and Cx26)
[25]. Our structure is thus in line with the recognized consensus motif necessary for heterotypic
complementarity of connexin GJCs [26].

Conformation of the Cx43 hemichannel. Although the two Cx43 GJC structures (in detergent
micelles and in nanodiscs) are almost identical, the Cx43 HC shows a notable difference in the
ECL1-2 conformation (Figure 3E-F). The resolution of the HC reconstruction does not allow
us to make definitive statements about the orientations of individual side chains, but it is clear
that both loops engaging in junction formation move inward upon docking of the two HCs
(Figure 3F). This conformational change likely involves intra- and intermolecular
cooperativity. The ECL1 and ECL2 are linked via several disulphide bonds, and thus the
rearrangement within each molecule would require concerted movement of the whole
extracellular domain (ECD). The ECD movement within one monomer is likely cooperatively
coupled to the neighboring chains within the Cx43 HC.

Disease-linked mutations in Cx43. A number of disease-linked mutations associated with
Cx43 can be mapped directly to three regions of interest revealed by our structures: (i) the GJC
intra-pore lipid-like site, (ii) the ECD, and (iii) the gating region (Figure 4A). The interior of
the channel features two lipid-like densities (Figure 2H, 4B; Figure S4E, S10A). The observed
position of lipid 1 is consistent with the previously found phospholipid-like densities inside the
pore of the Cx31.3 HC [20]. Lipid 2 inserts into the pocket formed by TM1 and TM2, parallel
with the NTD. Unlike Cx31.3, where elongated densities within the pore region could be
interpreted as hydrophobic tails of bound phospholipids, the density in the Cx43 GJC appears
consistent with that of a sterol (Figure 2H, 4B). The presence of a small hydrophobic small
molecule in Cx43 at this site suggests a potential mechanism of lipid-based Cx43 regulation,
whereby binding of a lipid could directly influence the conformation of the gating elements of
the protein (such as NTD). An effect of a cholesterol analogue 7-ketocholesterol on Cx43
permeability has been observed previously [27], and such an effect may be mediated via the
binding sites within the GJC (or HC) pore (Figure 4B; Figure S10A). The presence nearby of

several residues known to be linked to ODDD when mutated (S27P, I31M[28], S86Y[29], L90V[10])
points to a potential functional significance of the intra-pore lipid binding site.

Two known mutations linked to ODDD are located in the Cx43 ECD: P59 in ECL1 [30] and
H194 in ECL2 [31]. As suggested by the Cx43 GJC structures, these residues are located in
conserved regions where any substitution can be expected to cause disruption of the contacts
critical for junction formation (Figure 4C).

Multiple mutations in Cx43 associated with ODDD are located in the N-terminus (G2V, D3N,
L7V, L11I/P, Y17S, S18P), the TM1 region (G21R, G22E, K23T) and the TM2 region
proximal to the gate (V96M/E/A, Y98C, K102N) [11]. In some cases, the mutations reduce the
ability of Cx43 to form the gap junction plaques at the plasma membrane, as is the case for
Y17S or G21R (as well as L90V, located at the pore lipid binding site) [32]. The mutations tend
to have deleterious effects on the permeability of Cx43 to ions and small molecules [11].
Interestingly, a G8V amino acid substitution has been linked to palmoplantar keratoderma and
congenital alopecia 1 (PPKCA1). This mutant can form functional gap junctions and has
enhanced HC activity [33]. Mapping these sites on the structure of Cx43 allows us to gain insights
into the possible mechanisms that underlie the associated disorders (Figure 4D-E). For
example, channel blockage or hyperpermeability due to mutations in or around the gating
region may link Cx43 to diseases.

To test the functional consequences of mutations in the gating region, and in particular in the
PG region, we introduced several single residue substitutions in the NTD (A6F, G8F, G8V)
and TM2 (R101W) of Cx43. The mutations were designed to introduce bulky hydrophobic
side-chains into the locations that may perturb the integrity of the PG: either obstructing the
PG or preventing the PGL from binding to it, which could in turn influence the GJC transport.
We tested the ability of the mutated proteins to form functionally active GJCs in fluorescence
recovery after photobleaching (FRAP) assays in HEK293F cells expressing the mutant
proteins. Cx43 and the mutants could be expressed and showed expression patterns consistent
with gap junction formation between the neighbouring cells (Figure S11). Each of the tested
mutant variants of Cx43 retained the ability to transport a small molecule substrate,
sulforhodamin-101 (Figure S12). As noted above, the PPKCA1-linked mutant G8V has been
shown to form hyperactive HCs, and thus the lack of its effect on GJC function is not
unexpected. None of the other mutations were sufficient to disrupt the basic function of Cx43
GJCs in transporting the large substrates. Thus, although we can visualize the locations of the
disease-linked mutations, the underlying causes of the disease phenotypes will require more
detailed future investigations. Within the scope of this study, one can conclude that
introduction of bulky residues in the positions 6, 8 and 101 does not affect the transport function
of Cx43.

Molecular dynamics simulations of Cx43 GJC. Based on our model of Cx43, the dimensions
of the CG are likely incompatible with the translocation of a larger molecule, such as ATP,
cAMP or IP3 (typical substrates of Cx43 GJC-mediated transport). To assess the permeability
of the gate to ions, we performed MD simulations. We performed four independent molecular
dynamics simulations of the Cx43 GJC embedded in a double POPC bilayer solvated in 150
mM KCl (Figure 5A). Given the ambiguity in the lipid identity and binding mode, no potential
lipid surrogates were included inside the pore or NTD region. After equilibration (detailed in
Materials and Methods) the dodecameric structure was stable in all simulations, as indicated
by the small root mean square deviation (RMSD, Figure 5B; Figure S13).

As noted above, the CG pore has a solvent accessible radius very close to the effective hydrated
radius of K^+ (~6.6 Å) and Cl^- (~7.2 Å). The NTD regions of Cx43 are flexible and can move
laterally and vertically (Movie S1), allowing the passage of ions. The charge distribution inside
Cx43 resembles that of the other GJCs [18,19,34] with a positive electrostatic potential in the NTD

region of both HCs within a channel and a negative/neutral region near the HC interface (Figure
5C). When no transmembrane potential is applied (0 mV, Figure 5D), Cl^- ions accumulate in
the NTD regions of Cx43 GJC (negative free-energy in the NTD region) as previously
observed for C31.3 [34]. At 0 mV, there is a of ~2 kcal/mol barrier for cations (K^+) while no
barrier is found for anions. Cations get trapped between the two NTD barriers, and no full
translocations are observed during the simulation time at 0 mV because the gates remain
closed. Upon application of transmembrane voltage, the barrier on one HC lowers and full
translocations are observed (Figure 5D; detailed balance is provided in Table S2).

Interestingly, the GJC did not fully open in any of our simulations, i.e., a conformation with
all NTDs within the channel symmetrically moving to open the pore was not established at any
point. However, voltage-induced fluctuations of the individual NTDs allow the ions to pass.
To illustrate this effect we calculated the angle formed by the NTD and TM1 for all of the
chains in each of simulation (Figure 6A-D). We assessed the difference between the open
conformation (θ ~25°), the initial cryo-EM structure in the closed conformation (θ ~75°) and
the angle after equilibration (θ ~55°) in all of our simulations (Figure 6B). At 0 mV the NTD
fluctuates in different intermediate positions but is never fully open (Figure 6C). Only at high
applied transmembrane voltages the NTD fully folds inwards, thus opening the channel (Figure
6D). This is an important observation, taking into account the six-fold symmetry of the
connexin channels: although our understanding of the channel structure implies a symmetrical
arrangement of all its subunits, the gating events under physiological conditions, and under the
MD conditions as shown here, may involve asymmetrical conformational changes within the
NTD or other regions of the GJC or HC.

The MD simulations confirm that the structure is closed and semi-permeable to ions, even in
the absence of the PGL molecules occupying the gating region. Ion conductance only occurs
upon voltage application and is gated by the NTD regions folding inwards toward an open-like
conformation. The open NTD conformations of individual Cx43 monomers forming under high
voltage conditions in our MD simulations are similar to the open NTD conformations
previously reported using experimental approaches [17,18].

Discussion

The structures of Cx43 we have obtained by cryo-EM reveal a novel conformation of a
connexin channel (both GJC and HC) featuring a closed sieve-like molecular gate. The state is
distinct from the previously observed structures of connexin GJCs and HCs, adding an
important missing component to our understanding of connexin-mediated intercellular
communication. With these structures in mind, we can now establish the existence of several
structurally defined gating substates of the connexin channels: (i) a fully open gate, (ii) a semi-
permeable central gate, and (iii) a closed sieve-like gate (Figure 7). The gating mechanisms of
connexin HCs and GJC are complex and involve multiple components. The available evidence
indicates that connexin channels are gated by membrane potential (V_m), by transjunctional
potential (V_t), as well as by a range of chemical agents such as pH, Ca^{2+} and organic molecules
[35]. For example, a ball-and-chain model has been proposed for Cx26, whereby at low pH the
NTDs extend deep into the pore and occlude the substrate translocation pathway [36]. This
conformation is distinct from the one captured by our cryo-EM reconstructions, and it is
possible that under certain conditions the Cx43 NTD may transition from a sieve-like
conformation to a low pH Cx26-like state.

The cytosolic surface of the gate in our structures is positively charge, indicating that specific
interaction with anions may be relevant to some of the gating events. As Cx43 is capable of
cation and anion permeation [37], the presence of the positively charged surface in this region
does not indicate ion selectivity. The differences in the electrostatic properties of the gates and

the pore regions in different connexin channels may reflect the differences in their selectivity
for ions and small molecules (Figure S14). However, it is also likely that the unstructured
regions in Cx43 and other connexins, including the intracellular loop and the C-terminal
domain, contribute to the electrostatic potential, substrate selectivity and regulation of these
channels. These domains are known to play important roles in connexin channel regulation [38].
However, as these regions are unresolved in any of the available 3D reconstructions of
connexin channels, the structural basis of their action remains to be determined.

The observation of lipid-like molecules bound inside the pore is intriguing. The use of
detergents and lipids is currently a prerequisite for structural analysis of membrane proteins,
and it is possible that under solubilization conditions the abundance of added detergent and
lipid (CHS) forces the protein to interact with these molecules. Nevertheless, these interactions
may correspond to the naturally occurring interactions with the endogenous lipids present in
the lipid bilayer of the cell. The presence of such molecules could have important implications
for HC or GJC assembly, substrate permeation and molecular gating. It remains to be
determined whether Cx43 or other connexin channels harbour intra-pore lipids in the cellular
membranes *in vivo*.

Distinct gating modes have been shown in the literature for both Cx43 HCs and GJCs, with
fast transitions from an open state to a state of residual conductance, and slow transitions from
an open to a fully closed state [35]. The structures of Cx43 described here likely represent the
fully closed states, observed previously by electrophysiological recordings. This state could be
stabilized by a yet unknown agent that occludes the PG opening formed by the arrangement of
the NTDs and TM2 regions observed in our structures. Detailed investigations of Cx43 and
other connexin channels will be required to determine whether the sieve-like gate conformation
is a common feature among different GJCs and HCs, and to pinpoint the physiological
circumstances that establish this gating conformation.

Acknowledgements

We thank Emiliya Poghossian (EM Facility, PSI) and Miroslav Peterek (ScopeM, ETH Zurich) for expert support in cryo-EM data collection. We also thank Spencer Bliven and Marc Caubet-Serrabou (PSI) for the support in high performance computing. The work was supported by a grant from Horten Foundation, and by the Swiss National Science Foundation grant 184951 (VMK). FLG and SAG thank the CSCS and PRACE for supercomputing resources (projects pr126 and s1107).

Competing interests

Authors declare that they have no competing interests.

Data and materials availability

The atomic coordinates and structure factors have been deposited in the Protein Data Bank (7Z1T, 7Z22, 7Z23); the density maps have been deposited in the Electron Microscopy Data Bank (EMD-14452, EMD-14455, EMD-14456). The mass spectrometry data will be deposited at ProteomeXChange via PRIDE. All other data are available in the main text or the supplementary materials.

Author contributions

C.Q. planned and performed the experiments, analyzed the data, wrote the manuscript, S.A.G. planned, performed and analyzed the simulations, co-wrote the manuscript, P.L. performed the experiments, co-wrote the manuscript, A.O. performed the experiments, analyzed the data, D.L.P. performed the experiments, analyzed the data, E.B. performed the experiments, analyzed the data, D.S. performed the experiments, analysed the data, co-wrote the manuscript, P.P. provided the reagents, equipment and expertise for mass spectrometry experiments, N.Z. provided the reagents, equipment and expertise for lipidomics analysis, M.B. planned the experiments, analyzed the data, co-wrote the manuscript, F.L.G. planned and analyzed the simulations, co-wrote the manuscript, V.M.K. planned and performed the experiments, analyzed the data, wrote the manuscript.

341
342
Figure 1. Structure of human Cx43 gap junction channel. A-B, Cryo-EM density map and
model of Cx43 GJC solved by cryo-EM at 2.26 Å resolution. The individual Cx43 monomers
in each HC within the GJC are coloured blue, pink, grey, green, salmon and orange. Grey
densities correspond to the detergent micelle and the bound sterol-like molecules. **C**, The
position of a single Cx43 monomer (orange) within a GJC (represented as a surface of
juxtaposed Cx43 monomers in two distinct membrane regions, white and grey). **D**, Alignment
of the monomers of Cx43, Cx26 (PDB ID: 2zw3), Cx46 (PDB ID: 7jkc) and Cx31.3 (PDB ID:
6l3t) shows the distinct orientations of the N-terminal domain ("NTD") helix. Individual TM
domains, extracellular loops 1 and 2 ("ECL1-2"), relative positions of the intracellular loop
("ICL") and the C-terminus ("C-term") are indicated. **E-F**, Same as A-B, for the Cx43 HC in
MSP2N2 nanodiscs at 3.98 Å resolution. **G**, Alignment of three indicated structures shows that
the conformation of the gating region is highly conserved. The models are shown as ribbons;
one of the aligned monomers in each of the structures is represented as cartoon, to clearly
indicate the monomer boundaries.
357

358
359
Figure 2. The Cx43 gate adopts a sieve-like conformation. A-B, Pore analysis using HOLE
reveals a constriction of the pore at the gate. Only a central opening and one of the peripheral
openings within one HC of Cx43 GJC in digitonin is shown. The distance is calculated from
the center of the GJC pore to a point outside of the channel. Central gate ("CG") and peripheral
gate ("PG") regions are indicated in the panel. **C-D**, Comparison of the Cx43 in digitonin with
the Cx31.3 HC shows that the peripheral gate opening is created by the particular arrangement
of the NTD and by adjustment of the TM2. **E-F**, The distinct NTD/TM2 arrangement results
in a single pore opening in the structure of Cx31.3 (indicated with an asterisk, E), contrary to
Cx43 (F). **G**, A view of the Cx43 gating region from the cytosol reveals the location of the PG
lipids ("PGL") stabilizing the NTD arrangement, shown as isolated PGL densities (cyan). **H**,
A slab view of the gating region parallel to the membrane plane shows the relative arrangement
of the PGLs and the intra-pore lipids ("Lipid-1" and "Lipid-2"; orange).
372

373
374
Figure 3. Extracellular domain of Cx43 GJC and HC. A-B, The lateral views of Cx43 GJC
in digitonin, indicating the positions of the extracellular loops 1 and 2. **C**, Views of the
extracellular loops ECL1 (top) and ECL2 (bottom), for Cx43 (left), Cx46, (middle; PDB ID:
7jkc), and Cx26 (right; PDB ID: 2zw3). The residues of one monomer in each of the structures,
directly involved in junction formation, are coloured by element with carbon atoms as
orange/yellow (Cx43), brown/pink (Cx46) and teal/cyan (Cx26). The residues of the
neighbouring connexin monomers within 4 Å distance are shown coloured with white carbon
atoms. Dotted lines indicate electrostatic contacts, calculated in PyMol. **D**, Sequence alignment
of the complete ECL1 and ECL2 regions of the three proteins shown in C. The black lines
indicate the GJC interface residues, as shown in C. **E**, Alignment of the Cx43 GJC structures
in digitonin micelles and in nanodiscs. Cα atoms of the interface residues shown in C are
represented as spheres (the ribbon and spheres of Cx43 in nanodiscs is coloured black). **F**,
Same as E, for a comparison between Cx43 GJC in digitonin and Cx43 HC in nanodiscs. The
arrow indicates the movement of the two loops that accompanies GJC formation. The ribbon
and spheres of Cx43 HC are light grey.
390

Figure 4. Locations of the disease-linked mutations in the Cx43 GJC structure. A, Disease-linked mutations mapped on the sequence of Cx43. The amino acid residues proximal to the intrapore lipid densities ("Pore lipid sites") are shown as white dots. The residues at the GJC interface are shown as yellow dots ("ECD"). The residues of the gating region ("Gate") are shown as blue (NTD), green (TM1) and violet (TM2) dots. The sequence elements not resolved in our 3D reconstruction are indicated with a dashed line. **B**, Two sterol-like density elements (Lipid-1 and -2) are located within the pore region of Cx43 GJC (white). The indicated disease-linked mutations are located within close distance of the pore lipid densities. **C**, A view of the extracellular loops forming the junction, indicating two residues known to be linked with ODDD, P59 and H194 (side chains shown as sticks; one of the monomers in the GJC is coloured yellow). **D**, A view of the gating region with the highlighted disease-linked mutations, colored as in A. **E**, A similar representation of the disease-linked mutations, with Cα atoms shown as spheres, illustrating the contribution of the mutants to gate formation.

407
408
Figure 5. Molecular dynamics simulations of the Cx43 GJC. A, Molecular dynamics
simulation box. **B**, C_α root-mean-square-deviation from the cryo-EM structure during the
different molecular dynamics simulations. **C**, The surface representation of Cx43 coloured
according to the calculated electrostatic potential; the slab view shows the properties of the
pore. **D**, Free energy experienced by the K$^+$ (blue) and Cl$^-$ (red) while permeating the pore with
different applied transmembrane potential. The dotted lines correspond to the maximum free
energy barrier for anions (red) and cations (blue); black dotted line corresponds to the baseline.
416

417
418
419
Figure 6. Molecular dynamics simulations reveal voltage-dependent partial opening of
the gate. A, Views of a GJC, with twelve chain names referred to in c-d indicated. **B**,
Superimposition of a single chain depicted as cartoon for the closed structure of Cx43 (NTD
coloured in black), a semipermeable (rare) state extracted from the molecular dynamics
simulation (NTD coloured red) and the open structure of Cx46 (PDB ID: 7jjp, NTD coloured
blue). The angle θ corresponds to the calculated angle of the NTD domain with respect to TM1.
C, Population density of the NTD opening angle, θ, for the different chains in the GJC during
MD at 0 mV. **D**, The same analysis as in c, for the independent simulations with different
applied transmembrane voltages, for chains A to F (*top*) and for chains G to L (*bottom*).
429

Figure 7. A structure-based view of connexin gating states. A, The side view of a fully open gate, as observed in Cx26 GJC (PDB ID: 2zw3); the gate-forming regions of the protein, NTD (red) and TM2 (blue), are shown as cylinders. Similar arrangement of the gating elements has been observed in Cx46/Cx50. This conformation of the gate is permeable to a wide range of substrates, including small molecules and ions. **B**, The semi-permeable central gate is featured in the structure of Cx31.3 HC (PDB ID: 6l3t). In this conformation the gate is likely selective for ions (Cx31.3 has been shown to have a preference for anions), based on the dimensions of the gate. **C**, The sieve-like closed gate is featured in the Cx43 structures (here, the Cx43 in digitonin is shown as an example). **D-F**, Same as A-C, viewed from the cytosolic side of the channel. The black surface corresponds to the NTD and TM2 regions; the white surface corresponds to the rest of the protein. **G-I**, A schematic representation of the three gating states of connexin channels.

References

Churko, J. M. & Laird, D. W. Gap junction remodeling in skin repair following wounding and disease. *Physiology (Bethesda)* **28**, 190-198, doi:10.1152/physiol.00058.2012 (2013).

Liang, Z. *et al.* The Multifaceted Role of Astrocyte Connexin 43 in Ischemic Stroke Through Forming Hemichannels and Gap Junctions. *Front Neurol* **11**, 703, doi:10.3389/fneur.2020.00703 (2020).

Poelzing, S. & Rosenbaum, D. S. Altered connexin43 expression produces arrhythmia substrate in heart failure. *Am J Physiol Heart Circ Physiol* **287**, H1762-1770, doi:10.1152/ajpheart.00346.2004 (2004).

Rodriguez-Sinovas, A., Sanchez, J. A., Valls-Lacalle, L., Consegal, M. & Ferreira-Gonzalez, I. Connexins in the Heart: Regulation, Function and Involvement in Cardiac Disease. *Int J Mol Sci* **22**, doi:10.3390/ijms22094413 (2021).

Syrjanen, J., Michalski, K., Kawate, T. & Furukawa, H. On the molecular nature of large-pore channels. *J Mol Biol*, 166994, doi:10.1016/j.jmb.2021.166994 (2021).

Srinivas, M., Verselis, V. K. & White, T. W. Human diseases associated with connexin mutations. *Biochim Biophys Acta Biomembr* **1860**, 192-201, doi:10.1016/j.bbamem.2017.04.024 (2018).

Beyer, E. C., Paul, D. L. & Goodenough, D. A. Connexin43: a protein from rat heart homologous to a gap junction protein from liver. *J Cell Biol* **105**, 2621-2629, doi:10.1083/jcb.105.6.2621 (1987).

Robertson, J. D. The Occurrence of a Subunit Pattern in the Unit Membranes of Club Endings in Mauthner Cell Synapses in Goldfish Brains. *J Cell Biol* **19**, 201-221, doi:10.1083/jcb.19.1.201 (1963).

Laird, D. W. & Lampe, P. D. Therapeutic strategies targeting connexins. *Nat Rev Drug Discov* **17**, 905-921, doi:10.1038/nrd.2018.138 (2018).

Paznekas, W. A. *et al.* Connexin 43 (GJA1) mutations cause the pleiotropic phenotype of oculodentodigital dysplasia. *Am J Hum Genet* **72**, 408-418, doi:10.1086/346090 (2003).

Paznekas, W. A. *et al.* GJA1 mutations, variants, and connexin 43 dysfunction as it relates to the oculodentodigital dysplasia phenotype. *Hum Mutat* **30**, 724-733, doi:10.1002/humu.20958 (2009).

Wiest, T. *et al.* Clinical and genetic variability of oculodentodigital dysplasia. *Clin Genet* **70**, 71-72, doi:10.1111/j.1399-0004.2006.00631.x (2006).

Dasgupta, C. *et al.* Identification of connexin43 (alpha1) gap junction gene mutations in patients with hypoplastic left heart syndrome by denaturing gradient gel electrophoresis (DGGE). *Mutat Res* **479**, 173-186, doi:10.1016/s0027-5107(01)00160-9 (2001).

Pizzuti, A. *et al.* A homozygous GJA1 gene mutation causes a Hallermann-Streiff/ODDD spectrum phenotype. *Hum Mutat* **23**, 286, doi:10.1002/humu.9220 (2004).

Ribeiro-Rodrigues, T. M., Martins-Marques, T., Morel, S., Kwak, B. R. & Girao, H. Role of connexin 43 in different forms of intercellular communication - gap junctions, extracellular vesicles and tunnelling nanotubes. *J Cell Sci* **130**, 3619-3630, doi:10.1242/jcs.200667 (2017).

16   Unger, V. M., Kumar, N. M., Gilula, N. B. & Yeager, M. Three-dimensional structure of
a recombinant gap junction membrane channel. *Science* **283**, 1176-1180,
doi:10.1126/science.283.5405.1176 (1999).
17   Maeda, S. *et al.* Structure of the connexin 26 gap junction channel at 3.5 A
resolution. *Nature* **458**, 597-602, doi:10.1038/nature07869 (2009).
18   Myers, J. B. *et al.* Structure of native lens connexin 46/50 intercellular channels by
cryo-EM. *Nature* **564**, 372-377, doi:10.1038/s41586-018-0786-7 (2018).
19   Flores, J. A. *et al.* Connexin-46/50 in a dynamic lipid environment resolved by
CryoEM at 1.9 A. *Nat Commun* **11**, 4331, doi:10.1038/s41467-020-18120-5 (2020).
20   Lee, H. J. *et al.* Cryo-EM structure of human Cx31.3/GJC3 connexin hemichannel. *Sci*
Adv **6**, eaba4996, doi:10.1126/sciadv.aba4996 (2020).
21   Xu, Q. *et al.* Gating of connexin 43 gap junctions by a cytoplasmic loop calmodulin
binding domain. *Am J Physiol Cell Physiol* **302**, C1548-1556,
doi:10.1152/ajpcell.00319.2011 (2012).
22   Ponsaerts, R. *et al.* Intramolecular loop/tail interactions are essential for connexin
43-hemichannel activity. *FASEB J* **24**, 4378-4395, doi:10.1096/fj.09-153007 (2010).
23   Gartside, S. E., Griffith, N. C., Kaura, V. & Ingram, C. D. The neurosteroid
dehydroepiandrosterone (DHEA) and its metabolites alter 5-HT neuronal activity via
modulation of GABAA receptors. *J Psychopharmacol* **24**, 1717-1724,
doi:10.1177/0269881109105836 (2010).
24   Kimonides, V. G., Khatibi, N. H., Svendsen, C. N., Sofroniew, M. V. & Herbert, J.
Dehydroepiandrosterone (DHEA) and DHEA-sulfate (DHEAS) protect hippocampal
neurons against excitatory amino acid-induced neurotoxicity. *Proc Natl Acad Sci U S*
A **95**, 1852-1857, doi:10.1073/pnas.95.4.1852 (1998).
25   Bai, D., Yue, B. & Aoyama, H. Crucial motifs and residues in the extracellular loops
influence the formation and specificity of connexin docking. *Biochim Biophys Acta*
Biomembr **1860**, 9-21, doi:10.1016/j.bbamem.2017.07.003 (2018).
26   Koval, M., Molina, S. A. & Burt, J. M. Mix and match: investigating heteromeric and
heterotypic gap junction channels in model systems and native tissues. *FEBS Lett*
588, 1193-1204, doi:10.1016/j.febslet.2014.02.025 (2014).
27   Girao, H., Catarino, S. & Pereira, P. 7-Ketocholesterol modulates intercellular
communication through gap-junction in bovine lens epithelial cells. *Cell Commun*
Signal **2**, 2, doi:10.1186/1478-811X-2-2 (2004).
28   Richardson, R., Donnai, D., Meire, F. & Dixon, M. J. Expression of Gja1 correlates with
the phenotype observed in oculodentodigital syndrome/type III syndactyly. *J Med*
Genet **41**, 60-67, doi:10.1136/jmg.2003.012005 (2004).
29   Jamsheer, A. *et al.* Three novel GJA1 missense substitutions resulting in oculo-dento-
digital dysplasia (ODDD) - further extension of the mutational spectrum. *Gene* **539**,
157-161, doi:10.1016/j.gene.2014.01.066 (2014).
30   Vasconcellos, J. P. *et al.* A novel mutation in the GJA1 gene in a family with
oculodentodigital dysplasia. *Arch Ophthalmol* **123**, 1422-1426,
doi:10.1001/archopht.123.10.1422 (2005).
31   Vitiello, C. *et al.* A novel GJA1 mutation causes oculodentodigital dysplasia without
syndactyly. *Am J Med Genet A* **133A**, 58-60, doi:10.1002/ajmg.a.30554 (2005).
32   Lai, A. *et al.* Oculodentodigital dysplasia connexin43 mutations result in non-
functional connexin hemichannels and gap junctions in C6 glioma cells. *J Cell Sci* **119**,
532-541, doi:10.1242/jcs.02770 (2006).

33 Wang, H. *et al.* Exome sequencing reveals mutation in GJA1 as a cause of
keratoderma-hypotrichosis-leukonychia totalis syndrome. *Hum Mol Genet* **24**, 243-
250, doi:10.1093/hmg/ddu442 (2015).
34 Hyuk-Joon Lee, H. J., Jaekyung Hyun, Bumhan Ryu, Kunwoong Park, Hyun-Ho Lim,
Jejoong Yoo, Jae-Sung Woo. Cryo-EM structure of human Cx31.3/GJC3 connexin
hemichannel. *Science Advances* **6** (2020).
35 Bukauskas, F. F. & Verselis, V. K. Gap junction channel gating. *Biochim Biophys Acta*
1662, 42-60, doi:10.1016/j.bbamem.2004.01.008 (2004).
36 Khan, A. K. *et al.* A Steric "Ball-and-Chain" Mechanism for pH-Mediated Regulation of
Gap Junction Channels. *Cell Rep* **31**, 107482, doi:10.1016/j.celrep.2020.03.046
(2020).
37 Wang, H. Z. & Veenstra, R. D. Monovalent ion selectivity sequences of the rat
connexin43 gap junction channel. *J Gen Physiol* **109**, 491-507,
doi:10.1085/jgp.109.4.491 (1997).
38 D'Hondt, C., Iyyathurai, J., Himpens, B., Leybaert, L. & Bultynck, G. Cx43-hemichannel
function and regulation in physiology and pathophysiology: insights from the bovine
corneal endothelial cell system and beyond. *Front Physiol* **5**, 348,
doi:10.3389/fphys.2014.00348 (2014).
556