Structure of the connexin-43 gap junction channel in a putative closed state

Chao Qi[1,2,#], Silvia Acosta-Gutierrez[3,#], Pia Lavriha[1,2], Alaa Othman[4], Diego Lopez-Pigozzi[5,6], Erva Bayraktar[6], Dina Schuster[1,2,4], Paola Picotti[4], Nicola Zamboni[4], Mario Bortolozzi[5,6], Francesco L. Gervasio[7,8,9*] and Volodymyr M. Korkhov[1,2*]

[1] Institute of Molecular Biology and Biophysics, ETH Zurich, Switzerland

[2] Laboratory of Biomolecular Research, Paul Scherrer Institute, Villigen, Switzerland

[3] Institute for the Physics of Living Systems, Institute of Structural and Molecular Biology, and Department of Chemistry. University College London, London, UK

[4] Institute of Molecular Systems Biology, ETH Zurich, Switzerland

[5] Department of Physics and Astronomy "G. Galilei", University of Padova, Padua, Italy

[6] Veneto Institute of Molecular Medicine (VIMM), Padua, Italy

[7] Department of Chemistry, University College London, WC1E 6BT London, UK

[8] School of Pharmaceutical Sciences, University of Geneva, CH-1211 Geneva, Switzerland

[9] ISPSO, University of Geneva, CH-1211 Geneva, Switzerland

* Corresponding authors: francesco.gervasio@unige.ch, volodymyr.korkhov@psi.ch

# These authors contributed equally

Abstract

Gap junction channels (GJCs) mediate intercellular communication by connecting two neighboring cells and enabling direct exchange of ions and small molecules. Cell coupling via connexin-43 (Cx43) GJCs is important in a wide range of cellular processes in health and disease [1-3], yet the structural basis of Cx43 function and regulation has not been determined until now. Here we describe the structure of a human Cx43 GJC solved by cryo-EM and single particle analysis at 2.26 Å resolution. The pore region of Cx43 GJC features several lipid-like densities per Cx43 monomer, located close to a putative lateral access site at the monomer boundary. We found a previously undescribed conformation on the cytosolic side of the pore, formed by the N-terminal domain and the transmembrane helix 2 of Cx43 and stabilized by a small molecule. Structures of the Cx43 GJC and hemichannels in nanodiscs reveal a similar gate arrangement. The features of the Cx43 GJC and hemichannel cryo-EM maps and the channel properties revealed by molecular dynamics simulations suggest that the captured states of Cx43 are consistent with a closed state.

Introduction

Gap junction (GJ) mediated intercellular communication is one of the major pathways of information exchange between the cells. GJs are specialized regions of the plasma membrane at the cell-cell interface that link two adjacent cells and establish their metabolic and electrical coupling [4]. Connexins are the building blocks of the GJ channels (GJCs) which belong to a group of large-pore channels. This group includes a number of structurally related (innexins, pannexins, LRRC8) and unrelated proteins (CALHM) [5]. A total of 21 connexin genes have been identified in the human genome [6]. The 43 kDa connexin-43 (Cx43, gene name *GJA1*) was identified as a major constituent of rat heart GJs in 1987 [7], and it is arguably one of the most extensively studied connexins. Like all connexin homologues, Cx43 monomers assemble into hexameric hemichannels (HCs), also known as connexons. HCs that reach the plasma membrane of one cell may interact with their counterparts on the neighboring cell, forming GJCs, typically organized into hexagonal arrays at the intercellular interface [8]. GJCs enable direct metabolic and electric coupling between the cells, facilitating the passage of ions, small molecules, metabolites, peptides and other cellular components below a size threshold of approximately 1.5 kDa [9]. GJCs formed by Cx43 are crucial for a wide range of physiological processes, from propagation of heart action potentials [3] to maintenance of neuro-glial syncytium [2] and skin wound healing [1]. The clinical importance of Cx43 is highlighted by mutations linked to several genetic disorders, such as oculodentodigital dysplasia (ODDD) [10-12], hypoplastic left heart syndrome 1 [13], Hallermann-Streiff syndrome [14], atrioventricular septal defect 3 [13], etc., and by recognition of the protein as a drug target for treatment of cancer, skin wounds, eye injury and inflammation and cardiac arrhythmias (reviewed by Laird and Lampe [9]).

Much of what we know today about the molecular biology, electrophysiological properties and regulation of connexin gap junctions has been derived from the studies on Cx43 (reviewed in [15]). Early attempts to characterize the structure of Cx43 using cryo-electron microscopy (cryo-EM) of 2D crystals produced low resolution reconstructions [16]. More recently, several homologous connexin GJCs and HCs have been structurally characterized at high resolution, using X-ray crystallography and cryo-EM. The structures of Cx26 [17] and Cx46/50 GJCs [18,19], together with the recent structure of Cx31.3 HC [20] provided deep insights into the shared structural features of the connexin channels. These structures hint at the role of the N-terminal domain (NTD) in molecular gating of GJCs and HCs, with several conformations of the NTD observed in different connexin homologues [18-20]. Additionally, biochemical evidence points to the roles in channel gating played by the link between Cx43 intracellular loop and the C-terminal region [21,22]. However, despite the availability of this evidence, the molecular determinants of intracellular connexin channel gating remain unclear. To shed light on the structural basis of Cx43 gating, we set out to determine its structure by cryo-EM and to analyse its dynamics with molecular dynamics (MD) simulations.

Results

Structures of Cx43 GJC and HC in detergent micelles and in nanodiscs. Our Cx43 expression system was tested using electrophysiology in HeLa cells (Extended Data Fig. 1a-d). The experiments confirmed the functionality of Cx43 GJCs expressed in transfected cells. For large-scale protein expression, Cx43 featuring a C-terminal HRV3C-YFP-twinStrep tag, was expressed in adherent mammalian cells (HEK293F) using transient transfection method. The produced protein was purified using affinity and size exclusion chromatography in digitonin (Extended Data Fig. 2a-c) and flash frozen on cryo-EM grids. The coomassie-stained SDS PAGE gel bands of the purified protein (Extended Data Fig. 2c) are consistent with the western blot analysis (Extended Data Fig. 2d) of the expressed untagged Cx43. The grids were

subjected to single particle cryo-EM analysis (Extended Data Fig. 3), yielding the final 3D
reconstruction at 2.26 Å resolution (Fig. 1a, Extended Data Fig. 3-4).

The overall architecture of the protein complex resembles that observed with other connexin
GJCs, with two hemichannels (connexons) from adjacent plasma membrane regions coupled
to form a full GJC (Fig. 1a-b). The exterior of the channel is decorated by several ordered
detergent and/or lipid-like density elements (Fig. 1a-b), reminiscent of the previously observed
lipids bound at the protein-bilayer interface in other connexins [19]. In the case of Cx43, the lipid-
like densities appear to decorate the protein-lipid bilayer interface at both the inner and the
outer leaflet of the membrane. Comparison of the Cx43 monomers to the available structures
of Cx26 and Cx46 GJCs and the Cx31.3 HC revealed a major difference in the NTD
arrangement in these channels (Fig. 1c-d). The conformation of the NTD in the Cx31.3 HC
structure appears to be the closest to that in Cx43 GJC. Analysis of the tryptic peptides revealed
that most of the protein lacks the residue M1 (Extended Data Fig. 5), and thus the model was
built starting with G2.

To ascertain that the conformation of Cx43 GJC is not induced by the detergent present in the
sample, e.g. through detergent binding at specific sites on the protein surface, we removed the
detergent and reconstituted the protein into MSP2N2 nanodiscs, using 1-palmitoyl-2-oleoyl-
glycero-3-phosphocholine (POPC) as a mimic for the native lipid environment. The
reconstituted protein was subjected to the same imaging and analysis workflow as Cx43 GJC
in digitonin. The 2D classes of the MSP2N2-reconstituted Cx43 showed features consistent
with a mixture of GJCs and HC. Processing of the corresponding particles resulted in 3D
reconstructions of the Cx43 GJC at 2.95 Å resolution (Extended Data Fig. 6, 8) and HC in
nanodiscs at 3.98 Å resolution (Fig. 1e-f, Extended Data Fig. 7, 8). The GJCs in detergent and
in nanodiscs are nearly identical, with an RMSD of 0.97 Å between the aligned Cx43
monomers (Fig. 1g, Extended Data Fig. 9a). Although the differences between the HC in
nanodisc and the GJC are more pronounced (Extended Data Fig. 9a), the conformation of the
cytosolic region and the NTD is highly conserved (Fig. 1g).

Conformation of the Cx43 gate region. The reconstructed gate region of Cx43 shows several
narrow openings connecting the pore vestibule and the pore interior: a single ~6-7 Å wide
central opening and six adjacent openings of similar dimensions (Fig. 2a-b). This arrangement
of the gate region is distinct from any of the previously observed states of connexin GJCs or
HCs. This particular gate conformation is established through an interplay of two structural
elements: the NTD and the TM2 (Fig. 2c-d). The NTD of Cx43 is arranged near parallel to the
membrane plane, in a centre-oriented conformation (Fig. 2c). The HC of Cx31.3 assembles in
a similar manner, with a well-ordered NTD resolved by cryo-EM (Fig. 2c-d). However, the
TM2 region of Cx31.3 forms a tight seal with the NTD (Fig. 2e). In contrast, the TM2 of Cx43
is shifted away from the pore centre, creating six openings (Fig. 2f). Alignment of the
protomers of Cx43 and Cx31.3 showed an RMSD of 1.78 Å (using cealign in PyMol). In
contrast, the RMSD value for the region corresponding to the NTD and TM2 was 5.675 Å
(calculated using rms_cur in PyMol, with the residue range selections of 2-47 and 2-45 for the
aligned protomers of Cx43 and Cx31.3, respectively), further confirming that these two regions
differ substantially between Cx43 and Cx31.3. The arrangement of the Cx43 NTD appears to
be stabilized by a small molecule: a density element likely corresponding to a bound small
molecule is present within this region (referred to as the "NTD lipid site"), wedged between
the adjacent NTDs (Fig. 2g). Thus, although the conformation of the Cx43 gate features an
opening, this site is blocked by a yet unidentified small molecule (which we refer to as "lipid-
N").

While a direct comparison of the Cx43 gating regions of the Cx43 and Cx31.3 HCs is possible
(Extended Data Fig. 9c), the recently determined 3D reconstruction of the Cx26 mutant N176Y

was not accompanied by a new atomic model [23]. Instead the study reporting this reconstruction
compared the density map with an HC model from a Cx26 GJC structure (PDB ID: 5ERA) [24].
As this model lacks the NTD, any comparisons of the gating region in the Cx43 HC with that
in the Cx26 HC are presently limited (Extended Data Fig. 9d).

In addition to the lipid-N molecules stabilizing the NTD arrangement, the cryo-EM density of
Cx43 GJC features several well resolved densities inside the pore region (lipid-1 and lipid-2,
Fig. 2h). These elements likely correspond to bound sterol molecules, such as cholesterol co-
purified with the protein from the mammalian cells or cholesterol hemisuccinate (CHS) added
to the solubilization mixture during protein extraction from the membrane. Similar densities
are present in the Cx43 GJC reconstruction in nanodiscs, in the absence of detergent molecules
(Extended Data Fig. 10a). Additionally, it is noteworthy that the annular lipid densities are
conserved in both the detergent-solubilized and the nanodisc-reconstitued Cx43 GJCs,
indicating that the protein-lipid interface of Cx43 features sites where ordered lipid molecules
may bind (Extended Data Fig. 10b). The functional significance of the ordered annular lipids
for the channel activity and for GJ plaque assembly remains to be carefully investigated.

Identification of the lipid-like molecules bound to Cx43 is a significant challenge. Despite the
high overall resolution of the 3D reconstruction of the Cx43 GJC (2.26 Å resolution in
detergent and 2.95 Å in nanodisc; Extended Data Fig. 3, 6), the cryo-EM map features in
regions corresponding to the lipid densities are insufficient to assign the lipid identity
unambiguously. To determine the identities of these lipids experimentally, we performed
lipidomic analysis of the organic extracts prepared from the purified Cx43 samples and
compared them to the mock control (extracts of prepared from eluates of the purification
procedure using cells that do not overexpress the protein as starting material). The results
showed that several types of phospholipids, as well as cholesterol were present in both samples.
The only lipid-like compound specifically enriched in the purified sample was
dehydroepiandrosterone (DHEA; Extended Data Fig. 10c). DHEA is a most abundantly
expressed neurosteroid known to modulate ligand-gated ion channels, such as $GABA_A$ [25] or
NMDA and AMPA receptors [26]. Although we are presently not able to conclusively state that
DHEA corresponds to any of the densities present in our reconstruction, its enrichment in the
sample is an interesting observation that deserves detailed future investigations. While the
density corresponding to lipid-N may correspond to DHEA (Extended Data Fig. 4e), it is also
possible that this density corresponds to a larger molecule that is only partially ordered.

Cx43 junction interface. As in other connexins, the Cx43 GJC interface is formed by two
extracellular loops, ECL1 and ECL2 (Fig. 3a-b). The portion of the ECL1 comprising residues
N55-Q58 makes contacts with two Cx43 monomers from the opposite membrane, with N55-
Q58 of one monomer and Q58'-P59' of the neighboring monomer within 4 Å distance (Fig.
3c, *top left*). The ECL2 region that directly participates in sealing the junction includes residues
P193-D197. Comparisons of the junction-forming loops in Cx43 to those in Cx46 (Fig. 2c,
middle) and Cx26 (Fig. 2c, *right*) show that each protein uses a similar pattern of inter-HC
interaction within each GJC. It is noteworthy that both the amino acid sequences (Fig. 3d) and
the interaction networks (Fig. 3c) are similar in the ECL1 for Cx43 and Cx46, and in the ECL2
for Cx46 and Cx26. The differences in these two loops, and especially in ECL2, underlie the
inability of Cx43 (which has been grouped into a docking group 2) to engage in heterotypic
interactions with group 1 connexins (such as Cx46 and Cx26) [27]. Our structure is thus in line
with the recognized consensus motif necessary for heterotypic complementarity of connexin
GJCs [28].

Conformation of the Cx43 hemichannel. Although the two Cx43 GJC structures (in detergent
micelles and in nanodiscs) are almost identical, the Cx43 HC shows a notable difference in the
ECL1-2 conformation (Fig. 3e-f). The resolution of the HC reconstruction does not allow us to
make definitive statements about the orientations of individual side chains, but it is clear that

both loops engaging in junction formation move inward upon docking of the two HCs (Fig.
3f). This conformational change likely involves intra- and intermolecular cooperativity. The
ECL1 and ECL2 are linked via several disulphide bonds, and thus the rearrangement within
each molecule would require concerted movement of the whole extracellular domain (ECD).
The ECD movement within one monomer is likely cooperatively coupled to the neighboring
chains within the Cx43 HC.

Disease-linked mutations in Cx43. A number of disease-linked mutations associated with
Cx43 can be mapped directly to three regions of interest revealed by our structures: (i) the GJC
intra-pore lipid-like site, (ii) the ECD, and (iii) the gating region (Fig. 4a). The interior of the
channel features two lipid-like densities (Fig. 2h, 4b; Extended Data Fig. 4e, 10a). The
observed position of lipid 1 is consistent with the previously found phospholipid-like densities
inside the pore of the Cx31.3 HC [20]. Lipid 2 inserts into the pocket formed by TM1 and TM2,
parallel with the NTD. Unlike Cx31.3, where elongated densities within the pore region could
be interpreted as hydrophobic tails of bound phospholipids, the density in the Cx43 GJC
appears consistent with that of a sterol (Fig. 2h, 4b). The presence of a small hydrophobic small
molecule in Cx43 at this site suggests a potential mechanism of lipid-based Cx43 regulation,
whereby binding of a lipid could directly influence the conformation of the gating elements of
the protein (such as NTD). An effect of a cholesterol analogue 7-ketocholesterol on Cx43
permeability has been observed previously [29], and such an effect may be mediated via the
binding sites within the GJC (or HC) pore (Fig. 4b; Extended Data Fig. 10a). The presence
nearby of several residues known to be linked to ODDD when mutated (S27P, I31M[30], S86Y[31],
L90V[10]) points to a potential functional significance of the intra-pore lipid binding site.

Two known mutations linked to ODDD are located in the Cx43 ECD: P59 in ECL1 [32] and
H194 in ECL2 [33]. As suggested by the Cx43 GJC structures, these residues are located in
conserved regions where any substitution can be expected to cause disruption of the contacts
critical for junction formation (Fig. 4c).

Multiple mutations in Cx43 associated with ODDD are located in the N-terminus (G2V, D3N,
L7V, L11I/P, Y17S, S18P), the TM1 region (G21R, G22E, K23T) and the TM2 region
proximal to the gate (V96M/E/A, Y98C, K102N) [11]. In some cases, the mutations reduce the
ability of Cx43 to form the gap junction plaques at the plasma membrane, as is the case for
Y17S or G21R (as well as L90V, located at the pore lipid binding site) [34]. The mutations tend
to have deleterious effects on the permeability of Cx43 to ions and small molecules [11]. The
mutations G2V, D3N, W4A and L7V were shown to eliminate the function of Cx43 GJCs [35].
Interestingly, a G8V amino acid substitution has been linked to palmoplantar keratoderma and
congenital alopecia 1 (PPKCA1). This mutant can form functional gap junctions and has
enhanced HC activity [36]. Mapping these sites on the structure of Cx43 allows us to gain insights
into the possible mechanisms that underlie the associated disorders (Fig. 4d-e). For example,
channel blockage or hyperpermeability due to mutations in or around the gating region may
link Cx43 to diseases.

Molecular dynamics simulations of Cx43 GJC. Based on our model of Cx43, the dimensions
of the pore opening are likely incompatible with the translocation of a larger molecule, such as
ATP, cAMP or IP3 (typical substrates of Cx43 GJC-mediated transport). To assess the
permeability of the gate to ions, we performed MD simulations. We performed eighteen
independent molecular dynamics simulations of the Cx43 GJC embedded in a double POPC
bilayer solvated in 150 mM KCl. Given the ambiguity in the lipid identity and binding mode,
in the first set of simulations, no potential lipid surrogates were included inside the pore or
NTD region. After equilibration (detailed in Materials and Methods) the dodecameric structure
was stable in all simulations, as indicated by the small root mean square deviation (RMSD,
Extended Data Fig. 11 a,d).

The pore opening has a solvent-accessible radius of ~6Å (Figure 5c) very close to the effective
hydrated radius of K^+ (~6.6 Å) and Cl^- (~7.2 Å). This makes it the most narrow pore opening
observed for a connexin channel to date (a comparison of the pore openings in the cryo-EM
structures of connexin channels is shown in Extended Data Fig. 12). The NTD regions of Cx43
are flexible (Extended Data Fig. 13) and can move laterally and vertically (Movie 1), allowing
the passage of ions. The charge distribution inside Cx43 resembles that of the other GJCs [18,19,37]
with a positive electrostatic potential in the NTD region of both HCs within a channel and a
negative/neutral region near the HC interface (Fig. 5c). When no transmembrane potential is
applied (0 mV, Fig. 5b), Cl^- ions accumulate in the NTD regions of Cx43 GJC (anion density
peaks in the NTD region) as previously observed for C31.3 [37]. At 0 mV, there is an entry
barrier of ~3.38 kcal/mol for cations (K^+) in the NTD region of both HCs (Figure 5d) which is
slightly higher than previously reported PMFs values for other homomeric GJCs (Cx46/50) [18].
Conversely, anions can permeate the NTD region barrierless (Figure 5d), but they face a higher
barrier ~3.84 kcal/mol below this region where the electrostatic potential of the channel is
negative, as observed for other GJC [18]. Because the NTD domains are flexible, but they do not
fully fold inwards, most ions enter and exit the same HC (Extended Data Table 2). Application
of a transjunctional voltage lowers the barrier on one HC increasing the number of permeation
events. Very few full transjunctional permeation events were observed during the simulations
with any applied voltage.

As mentioned before, the GJC did not fully open in any of our simulations, i.e., a conformation
with all NTDs within the channel symmetrically moving to open the pore was not established
at any point. To illustrate this effect, we calculated the root-mean-square deviation for each
NTD domain in all simulations (Extended Data Fig. 13). Only at high applied transjunctional
voltages only one NTD domain adopted a fully open folded inward conformation (Extended
Data Fig. 13).

The MD simulations in the absence of a bound ligand reveal an intermediate, metastable state
in which ions can permeate from both HCs with similar free-energy barriers and full
transjunctional permeations are very rare events. The GJC selectivity and conductance
properties are modulated by complex mechanisms involving both the steric aperture and the
unique pattern of electrostatic features. In our case, the electrostatic properties of the channel
are similar to those published for the Cx46/50 channels [18], but the steric contribution of the
NTD region is significantly higher, increasing the barrier for K^+, resulting in smaller $\Delta\Delta G$
differences than those observed for the open channel (Cx46/50). Other MD studies have shown
similar energy barriers for ions [18], leading to entry events but very rare full translocations on
the molecular dynamics timescale. Therefore, in the absence of the ligand the channel may
exhibit a low residual conductance, as described experimentally [38].

Despite not knowing the precise identity of the lipid-N, we performed nine MD simulations of
the Cx43 HC with a DHEA molecule as a lipid-N surrogate bound between the adjacent chains
(Fig. 5e, Extended Data Fig. 14a,b, Movie 2), to understand whether the channel is closed in
the presence of the lipid molecule. As expected, even when applying high transmembrane
voltage the ion density for both species, K^+ and Cl^-, in the NTD region remained zero and no
permeation events were observed (Fig. 5e, Extended Data Fig. 14c). Hence, the presence of the
bound lipid-N molecule seals the channel by rigidifying the NTD domains.

Discussion

The structures of Cx43 we have obtained by cryo-EM reveal a novel conformation of a
connexin channel (both GJC and HC) featuring a closed molecular gate. The state is distinct
from the previously observed structures of connexin GJCs and HCs, adding an important
missing component to our understanding of connexin-mediated intercellular communication.

With these structures in mind, we can now establish the existence of several structurally
defined gating substates of the connexin channels: (i) a fully open gate, (ii) a semi-permeable
central gate, and (iii) a closed gate (Fig. 6). The gating mechanisms of connexin HCs and GJC
are complex and involve multiple components. The available evidence indicates that connexin
channels are gated by membrane potential (V_m), by transjunctional potential (V_j), as well as by
a range of chemical agents such as pH, Ca^{2+} and organic molecules [38]. For example, a ball-and-
chain model has been proposed for Cx26, whereby at low pH the NTDs extend deep into the
pore and occlude the substrate translocation pathway [39]. This conformation is distinct from the
one captured by our cryo-EM reconstructions, and it is possible that under certain conditions
the Cx43 NTD may transition from the observed conformation to a low pH Cx26-like state. It
is worth mentioning that for the Cx43 HCs the open probability has been shown to be very low
[40], and thus a closed conformation such as the one described here may represent the
predominant state of the channel.

The cytosolic surface of the gate in our structures is positively charged, indicating that specific
interaction with anions may be relevant to some of the gating events. As Cx43 is capable of
cation and anion permeation [41], the presence of the positively charged surface in this region
does not indicate ion selectivity. The differences in the electrostatic properties of the gates and
the pore regions in different connexin channels may reflect the differences in their selectivity
for ions and small molecules (Extended Data Fig. 12). However, it is also likely that the
unstructured regions in Cx43 and other connexins, including the intracellular loop and the C-
terminal domain, contribute to the electrostatic potential, substrate selectivity and regulation of
these channels. These domains are known to play important roles in connexin channel
regulation [42]. However, as these regions are unresolved in any of the available 3D
reconstructions of connexin channels, the structural basis of their action remains to be
determined.

The observation of lipid-like molecules bound inside the pore is intriguing. The use of
detergents and lipids is currently a prerequisite for structural analysis of membrane proteins,
and it is possible that under solubilization conditions the abundance of added detergent and
lipid (CHS) forces the protein to interact with these molecules. Nevertheless, these interactions
may correspond to the naturally occurring interactions with the endogenous lipids present in
the lipid bilayer of the cell. The presence of such molecules could have important implications
for HC or GJC assembly, substrate permeation and molecular gating. It remains to be
determined whether Cx43 or other connexin channels harbour intra-pore lipids in the cellular
membranes *in vivo*.

Distinct gating modes have been shown in the literature for both Cx43 HCs and GJCs, with
fast transitions from an open state to a state of residual conductance, and slow transitions from
an open to a fully closed state [38]. The structures of Cx43 described here may represent the
closed states, based on the cryo-EM- and MD simulation-based evidence. The closed state
could be stabilized by a yet unknown agent that occupies the NTD lipid site formed by the
arrangement of the NTDs and TM2 regions observed in our structures. It is worth noting that
the majority of Cx43 channels in a GJ plaque are closed, with only a small fraction of the
channels active [43]. Detailed investigations of Cx43 and other connexin channels will be
required to determine whether the gate conformation observed here is a common feature among
different GJCs and HCs, and to pinpoint the physiological circumstances that establish this
gating conformation. Finally, capturing the structures of Cx43 channels in distinct
conformations under a wide range of physiologically relevant conditions will be required to
establish its mechanism of action, gating and regulation.

Acknowledgements

We thank Emiliya Poghossian (EM Facility, PSI) and Miroslav Peterek (ScopeM, ETH Zurich) for expert support in cryo-EM data collection. We also thank Spencer Bliven and Marc Caubet-Serrabou (PSI) for the support in high performance computing. The work was supported by a grant from Horten Foundation, and by the Swiss National Science Foundation grant 184951 (VMK). FLG and SAG thank the CSCS and PRACE for supercomputing resources (projects pr126 and s1107).

Competing interests

Authors declare that they have no competing interests.

Data and materials availability

The atomic coordinates and structure factors have been deposited in the Protein Data Bank (7Z1T, 7Z22, 7Z23); the density maps have been deposited in the Electron Microscopy Data Bank (EMD-14452, EMD-14455, EMD-14456). The mass spectrometry data have been deposited at ProteomeXChange via PRIDE (PXD033824). The MD trajectories will be uploaded to Zenodo. All other data are available in the main text or the supplementary materials.

Author contributions

C.Q. planned and performed the experiments, analyzed the data, wrote the manuscript, S.A.G. planned, performed and analyzed the simulations, co-wrote the manuscript, P.L. performed the experiments, co-wrote the manuscript, A.O. performed the experiments, analyzed the data, D.L.P. performed the experiments, analyzed the data, E.B. performed the experiments, analyzed the data, D.S. performed the experiments, analysed the data, co-wrote the manuscript, P.P. provided the reagents, equipment and expertise for mass spectrometry experiments, N.Z. provided the reagents, equipment and expertise for lipidomics analysis, M.B. planned the experiments, analyzed the data, co-wrote the manuscript, F.L.G. planned and analyzed the simulations, co-wrote the manuscript, V.M.K. planned and performed the experiments, analyzed the data, wrote the manuscript.

362
363 **Figures**
364

365
366
Figure 1. Structure of human Cx43 gap junction channel. a-b, Cryo-EM density map and
model of Cx43 GJC solved by cryo-EM at 2.26 Å resolution. The individual Cx43 monomers
in each HC within the GJC are coloured blue, pink, grey, green, salmon and orange. Grey
densities correspond to the detergent micelle and the bound sterol-like molecules. **c**, The
position of a single Cx43 monomer (orange) within a GJC (represented as a surface of
juxtaposed Cx43 monomers in two distinct membrane regions, white and grey). **d**, Alignment
of the monomers of Cx43, Cx26 (PDB ID: 2zw3), Cx46 (PDB ID: 7jkc) and Cx31.3 (PDB ID:
6l3t) shows the distinct orientations of the N-terminal domain ("NTD") helix. Individual TM
domains, extracellular loops 1 and 2 ("ECL1-2"), relative positions of the intracellular loop
("ICL") and the C-terminus ("C-term") are indicated. **e-f**, Same as a-b, for the Cx43 HC in
MSP2N2 nanodiscs at 3.98 Å resolution. **g**, Alignment of three indicated structures shows that
the conformation of the gating region is highly conserved. The models are shown as ribbons;
one of the aligned monomers in each of the structures is represented as cartoon, to clearly
indicate the monomer boundaries.
381
382

Figure 2. The Cx43 gate adopts a closed conformation. a-b, Analysis of the pore opening dimensions using HOLE reveals a constriction of the pore in the gate region. Only a central opening and one of the six peripheral openings within one HC of Cx43 GJC in digitonin is shown. The distance is calculated from the center of the GJC pore to a point outside of the channel. Central and peripheral openings are coloured blue and cyan, respectively. **c-d**, Comparison of the Cx43 in digitonin with the Cx31.3 HC shows that the peripheral opening is created by the particular arrangement of the NTD and by adjustment of the TM2. **e-f**, The distinct NTD/TM2 arrangement results in a single pore opening in the structure of Cx31.3 (indicated with an asterisk, e), contrary to Cx43 (f). **g**, A view of the Cx43 gating region from the cytosol reveals the location of the "lipid-N" molecules stabilizing the NTD arrangement, shown as isolated densities (cyan). **h**, A slab view of the gating region parallel to the membrane plane shows the relative arrangement of the lipid-N and the intra-pore lipid densities ("lipid-1" and lipid-2"; orange).

Figure 3. Extracellular domain of Cx43 GJC and HC. a–b, The lateral views of Cx43 GJC in digitonin, indicating the positions of the extracellular loops 1 and 2. **c**, Views of the extracellular loops ECL1 (top) and ECL2 (bottom), for Cx43 (left), Cx46, (middle; PDB ID: 7jkc), and Cx26 (right; PDB ID: 2zw3). The residues of one monomer in each of the structures, directly involved in junction formation, are coloured by element with carbon atoms as orange/yellow (Cx43), brown/pink (Cx46) and teal/cyan (Cx26). The residues of the neighbouring connexin monomers within 4 Å distance are shown coloured with white carbon atoms. Dotted lines indicate electrostatic contacts, calculated in PyMol. **d**, Sequence alignment of the complete ECL1 and ECL2 regions of the three proteins shown in c. The black lines indicate the GJC interface residues, as shown in c. **e**, Alignment of the Cx43 GJC structures in digitonin micelles and in nanodiscs. Cα atoms of the interface residues shown in C are represented as spheres (the ribbon and spheres of Cx43 in nanodiscs is coloured black). **f**, Same as e, for a comparison between Cx43 GJC in digitonin and Cx43 HC in nanodiscs. The arrow indicates the movement of the two loops that accompanies GJC formation. The ribbon and spheres of Cx43 HC are light grey. The grey numbers in brackets indicate displacement of selected Cα atoms (residues P59, Q58, P193, P194).

418
419
Figure 4. Locations of the disease-linked mutations in the Cx43 GJC structure. a, Disease-
linked mutations mapped on the sequence of Cx43. The amino acid residues proximal to the
intrapore lipid densities ("Pore lipid sites") are shown as white dots. The residues at the GJC
interface are shown as yellow dots ("ECD"). The residues of the gating region ("Gate region")
are shown as blue (NTD), green (TM1) and violet (TM2) dots. The sequence elements not
resolved in our 3D reconstruction are indicated with a dashed line. **b**, Two sterol-like density
elements (lipid-1 and -2) are located within the pore region of Cx43 GJC (white). The indicated
disease-linked mutations are located within close distance of the pore lipid densities. **c**, A view
of the extracellular loops forming the junction, indicating two residues known to be linked with
ODDD, P59 and H194 (side chains shown as sticks; one of the monomers in the GJC is
coloured yellow). **d**, A view of the gating region with the highlighted disease-linked mutations,
colored as in a. **e**, A similar representation of the disease-linked mutations, with Cα atoms
shown as spheres, illustrating the contribution of the mutants to gate formation.
433

434
435
Figure 5. Molecular dynamics simulations of the Cx43 GJC and HC. a, Cartoon
representation of the double-bilayer system including the GJC (cartoon) and ions (vdw
spheres). Lipidic membrane and water residues have been removed for clarity. The direction
of the applied constant electric field, E_z, is indicated with an arrow. **b**, Ion density (ρ) profiles
(average and fluctuations) along the diffusion axis of the GJC coloured in red for anions and
blue for cations, for the simulated applied transjunctional voltages. **c**, The solvent-accessible
radius (and fluctuations during MD) along the diffusion axis of the GJC. The dotted lines
correspond to the minimum radius and the position of the NTD regions. The surface
representation of Cx43 is coloured according to the calculated electrostatic potential; the slab
view shows the properties of the pore. **d**, Average free energy experienced (at least two
simulation replicates per panel) by the K^+ (blue) and Cl^- (red) while permeating the pore with
different applied transmembrane potential (from right to left: 0 mV, -100 mV, -200 mV and -
500 mV). The dotted lines correspond to the maximum free energy barrier for anions (red) and
cations (blue). **e**, Cartoon representation of the lipid-bound system including the HC (cartoon)
and ions (vdw spheres). Lipidic membrane and water residues have been removed for clarity.
The direction of the applied constant electric field, E_z, is indicated with an arrow. **f**, Ion density

452   (ρ) profiles (average and fluctuations) along the diffusion axis of the HC coloured in red for
453   anions and blue for cations, for the simulated applied transmembrane voltages.
454

Open gate   Semi-permeable gate   Closed gate

Figure 6. A structure-based view of connexin gating states. a, The side view of a fully open gate, as observed in Cx26 GJC (PDB ID: 2zw3); the gate-forming regions of the protein, NTD (red) and TM2 (blue), are shown as cylinders. Similar arrangement of the gating elements has been observed in Cx46/Cx50. This conformation of the gate is permeable to a wide range of substrates, including small molecules and ions. **b**, The semi-permeable central gate is featured in the structure of Cx31.3 HC (PDB ID: 6l3t). In this conformation the gate is likely selective for ions (Cx31.3 has been shown to have a preference for anions), based on the dimensions of the gate. **c**, The closed gate is featured in the Cx43 structures (here, the Cx43 in digitonin is shown as an example). **d-f**, Same as a-c, viewed from the cytosolic side of the channel. The gray surface corresponds to the NTD and TM2 regions; the white surface corresponds to the rest of the protein; lipid-N molecules in f are shown as black spheres, using a model of DHEA manually placed into the six NTD lipid site regions for illustration purposes (the exact identity of the lipid-N molecule is unknown).

References

Churko, J. M. & Laird, D. W. Gap junction remodeling in skin repair following wounding and disease. *Physiology (Bethesda)* **28**, 190-198 (2013). https://doi.org:10.1152/physiol.00058.2012

Liang, Z. *et al.* The Multifaceted Role of Astrocyte Connexin 43 in Ischemic Stroke Through Forming Hemichannels and Gap Junctions. *Front Neurol* **11**, 703 (2020). https://doi.org:10.3389/fneur.2020.00703

Poelzing, S. & Rosenbaum, D. S. Altered connexin43 expression produces arrhythmia substrate in heart failure. *Am J Physiol Heart Circ Physiol* **287**, H1762-1770 (2004). https://doi.org:10.1152/ajpheart.00346.2004

Rodriguez-Sinovas, A., Sanchez, J. A., Valls-Lacalle, L., Consegal, M. & Ferreira-Gonzalez, I. Connexins in the Heart: Regulation, Function and Involvement in Cardiac Disease. *Int J Mol Sci* **22** (2021). https://doi.org:10.3390/ijms22094413

Syrjanen, J., Michalski, K., Kawate, T. & Furukawa, H. On the molecular nature of large-pore channels. *J Mol Biol*, 166994 (2021). https://doi.org:10.1016/j.jmb.2021.166994

Srinivas, M., Verselis, V. K. & White, T. W. Human diseases associated with connexin mutations. *Biochim Biophys Acta Biomembr* **1860**, 192-201 (2018). https://doi.org:10.1016/j.bbamem.2017.04.024

Beyer, E. C., Paul, D. L. & Goodenough, D. A. Connexin43: a protein from rat heart homologous to a gap junction protein from liver. *J Cell Biol* **105**, 2621-2629 (1987). https://doi.org:10.1083/jcb.105.6.2621

Robertson, J. D. The Occurrence of a Subunit Pattern in the Unit Membranes of Club Endings in Mauthner Cell Synapses in Goldfish Brains. *J Cell Biol* **19**, 201-221 (1963). https://doi.org:10.1083/jcb.19.1.201

Laird, D. W. & Lampe, P. D. Therapeutic strategies targeting connexins. *Nat Rev Drug Discov* **17**, 905-921 (2018). https://doi.org:10.1038/nrd.2018.138

Paznekas, W. A. *et al.* Connexin 43 (GJA1) mutations cause the pleiotropic phenotype of oculodentodigital dysplasia. *Am J Hum Genet* **72**, 408-418 (2003). https://doi.org:10.1086/346090

Paznekas, W. A. *et al.* GJA1 mutations, variants, and connexin 43 dysfunction as it relates to the oculodentodigital dysplasia phenotype. *Hum Mutat* **30**, 724-733 (2009). https://doi.org:10.1002/humu.20958

Wiest, T. *et al.* Clinical and genetic variability of oculodentodigital dysplasia. *Clin Genet* **70**, 71-72 (2006). https://doi.org:10.1111/j.1399-0004.2006.00631.x

Dasgupta, C. *et al.* Identification of connexin43 (alpha1) gap junction gene mutations in patients with hypoplastic left heart syndrome by denaturing gradient gel electrophoresis (DGGE). *Mutat Res* **479**, 173-186 (2001). https://doi.org:10.1016/s0027-5107(01)00160-9

Pizzuti, A. *et al.* A homozygous GJA1 gene mutation causes a Hallermann-Streiff/ODDD spectrum phenotype. *Hum Mutat* **23**, 286 (2004). https://doi.org:10.1002/humu.9220

Ribeiro-Rodrigues, T. M., Martins-Marques, T., Morel, S., Kwak, B. R. & Girao, H. Role of connexin 43 in different forms of intercellular communication - gap junctions, extracellular vesicles and tunnelling nanotubes. *J Cell Sci* **130**, 3619-3630 (2017). https://doi.org:10.1242/jcs.200667

16 Unger, V. M., Kumar, N. M., Gilula, N. B. & Yeager, M. Three-dimensional structure of
a recombinant gap junction membrane channel. *Science* **283**, 1176-1180 (1999).
https://doi.org:10.1126/science.283.5405.1176
17 Maeda, S. *et al.* Structure of the connexin 26 gap junction channel at 3.5 A
resolution. *Nature* **458**, 597-602 (2009). https://doi.org:10.1038/nature07869
18 Myers, J. B. *et al.* Structure of native lens connexin 46/50 intercellular channels by
cryo-EM. *Nature* **564**, 372-377 (2018). https://doi.org:10.1038/s41586-018-0786-7
19 Flores, J. A. *et al.* Connexin-46/50 in a dynamic lipid environment resolved by
CryoEM at 1.9 A. *Nat Commun* **11**, 4331 (2020). https://doi.org:10.1038/s41467-020-
18120-5
20 Lee, H. J. *et al.* Cryo-EM structure of human Cx31.3/GJC3 connexin hemichannel. *Sci
Adv* **6**, eaba4996 (2020). https://doi.org:10.1126/sciadv.aba4996
21 Xu, Q. *et al.* Gating of connexin 43 gap junctions by a cytoplasmic loop calmodulin
binding domain. *Am J Physiol Cell Physiol* **302**, C1548-1556 (2012).
https://doi.org:10.1152/ajpcell.00319.2011
22 Ponsaerts, R. *et al.* Intramolecular loop/tail interactions are essential for connexin
43-hemichannel activity. *FASEB J* **24**, 4378-4395 (2010).
https://doi.org:10.1096/fj.09-153007
23 Khan, A. K., Jagielnicki, M., Bennett, B. C., Purdy, M. D. & Yeager, M. Cryo-EM
structure of an open conformation of a gap junction hemichannel in lipid bilayer
nanodiscs. *Structure* **29**, 1040-1047 e1043 (2021).
https://doi.org:10.1016/j.str.2021.05.010
24 Bennett, B. C. *et al.* An electrostatic mechanism for Ca(2+)-mediated regulation of
gap junction channels. *Nat Commun* **7**, 8770 (2016).
https://doi.org:10.1038/ncomms9770
25 Gartside, S. E., Griffith, N. C., Kaura, V. & Ingram, C. D. The neurosteroid
dehydroepiandrosterone (DHEA) and its metabolites alter 5-HT neuronal activity via
modulation of GABAA receptors. *J Psychopharmacol* **24**, 1717-1724 (2010).
https://doi.org:10.1177/0269881109105836
26 Kimonides, V. G., Khatibi, N. H., Svendsen, C. N., Sofroniew, M. V. & Herbert, J.
Dehydroepiandrosterone (DHEA) and DHEA-sulfate (DHEAS) protect hippocampal
neurons against excitatory amino acid-induced neurotoxicity. *Proc Natl Acad Sci U S
A* **95**, 1852-1857 (1998). https://doi.org:10.1073/pnas.95.4.1852
27 Bai, D., Yue, B. & Aoyama, H. Crucial motifs and residues in the extracellular loops
influence the formation and specificity of connexin docking. *Biochim Biophys Acta
Biomembr* **1860**, 9-21 (2018). https://doi.org:10.1016/j.bbamem.2017.07.003
28 Koval, M., Molina, S. A. & Burt, J. M. Mix and match: investigating heteromeric and
heterotypic gap junction channels in model systems and native tissues. *FEBS Lett*
588, 1193-1204 (2014). https://doi.org:10.1016/j.febslet.2014.02.025
29 Girao, H., Catarino, S. & Pereira, P. 7-Ketocholesterol modulates intercellular
communication through gap-junction in bovine lens epithelial cells. *Cell Commun
Signal* **2**, 2 (2004). https://doi.org:10.1186/1478-811X-2-2
30 Richardson, R., Donnai, D., Meire, F. & Dixon, M. J. Expression of Gja1 correlates with
the phenotype observed in oculodentodigital syndrome/type III syndactyly. *J Med
Genet* **41**, 60-67 (2004). https://doi.org:10.1136/jmg.2003.012005
31 Jamsheer, A. *et al.* Three novel GJA1 missense substitutions resulting in oculo-dento-
digital dysplasia (ODDD) - further extension of the mutational spectrum. *Gene* **539**,
157-161 (2014). https://doi.org:10.1016/j.gene.2014.01.066

32   Vasconcellos, J. P. *et al.* A novel mutation in the GJA1 gene in a family with
oculodentodigital dysplasia. *Arch Ophthalmol* **123**, 1422-1426 (2005).
https://doi.org:10.1001/archopht.123.10.1422
33   Vitiello, C. *et al.* A novel GJA1 mutation causes oculodentodigital dysplasia without
syndactyly. *Am J Med Genet A* **133A**, 58-60 (2005).
https://doi.org:10.1002/ajmg.a.30554
34   Lai, A. *et al.* Oculodentodigital dysplasia connexin43 mutations result in non-
functional connexin hemichannels and gap junctions in C6 glioma cells. *J Cell Sci* **119**,
532-541 (2006). https://doi.org:10.1242/jcs.02770
35   Shao, Q. *et al.* Structure and functional studies of N-terminal Cx43 mutants linked to
oculodentodigital dysplasia. *Mol Biol Cell* **23**, 3312-3321 (2012).
https://doi.org:10.1091/mbc.E12-02-0128
36   Wang, H. *et al.* Exome sequencing reveals mutation in GJA1 as a cause of
keratoderma-hypotrichosis-leukonychia totalis syndrome. *Hum Mol Genet* **24**, 243-
250 (2015). https://doi.org:10.1093/hmg/ddu442
37   Hyuk-Joon Lee, H. J., Jaekyung Hyun, Bumhan Ryu, Kunwoong Park, Hyun-Ho Lim,
Jejoong Yoo, Jae-Sung Woo. Cryo-EM structure of human Cx31.3/GJC3 connexin
hemichannel. *Science Advances* **6** (2020).
38   Bukauskas, F. F. & Verselis, V. K. Gap junction channel gating. *Biochim Biophys Acta*
1662, 42-60 (2004). https://doi.org:10.1016/j.bbamem.2004.01.008
39   Khan, A. K. *et al.* A Steric "Ball-and-Chain" Mechanism for pH-Mediated Regulation of
Gap Junction Channels. *Cell Rep* **31**, 107482 (2020).
https://doi.org:10.1016/j.celrep.2020.03.046
40   Contreras, J. E., Saez, J. C., Bukauskas, F. F. & Bennett, M. V. Gating and regulation of
connexin 43 (Cx43) hemichannels. *Proc Natl Acad Sci U S A* **100**, 11388-11393 (2003).
https://doi.org:10.1073/pnas.1434298100
41   Wang, H. Z. & Veenstra, R. D. Monovalent ion selectivity sequences of the rat
connexin43 gap junction channel. *J Gen Physiol* **109**, 491-507 (1997).
https://doi.org:10.1085/jgp.109.4.491
42   D'Hondt, C., Iyyathurai, J., Himpens, B., Leybaert, L. & Bultynck, G. Cx43-hemichannel
function and regulation in physiology and pathophysiology: insights from the bovine
corneal endothelial cell system and beyond. *Front Physiol* **5**, 348 (2014).
https://doi.org:10.3389/fphys.2014.00348
43   Bukauskas, F. F. *et al.* Clustering of connexin 43-enhanced green fluorescent protein
gap junction channels and functional coupling in living cells. *Proc Natl Acad Sci U S A*
97, 2556-2561 (2000). https://doi.org:10.1073/pnas.050588497
603