Dynamic interaction of BcsD subunit in type I bacterial cellulose synthase

Tatsuya Kondo[1], Yui Nakamura[2], Shingo Nojima[2], Min Yao[2], Tomoya Imai[1]

[1] Research Institute for Sustainable Humanosphere (RISH), Kyoto University, Uji, Kyoto 611-0011,

Japan.

[2] Faculty of Advanced Life Science, Hokkaido University, N10W Kita-Ku, Sapporo 060-0810, Japan.

For correspondence:

Tomoya Imai, imai.tomoya.4c@kyoto-u.ac.jp; Research Institute for Sustainable Humanosphere, Kyoto

University, Uji, Kyoto 611-0011, Japan.

Min Yao, yao@castor.sci.hokudai.ac.jp; Faculty of Advanced Life Science, Hokkaido University,

N10W Kita-Ku, Sapporo 060-0810, Japan.

Running title: Dynamic interaction of BcsD in type I bacterial cellulose synthase

Keywords: cellulose, cellulose synthase, membrane enzyme, protein complex, protein-protein

interaction, polysaccharide synthesis

Abstract

Cellulose is a promising biological material for supporting sustainable human life. This natural polymer

is synthesized by cellulose synthase, a protein complex in the cell membrane. Cellulose synthase in

bacteria is a hetero-subunit complex, and its subunit organization varies widely depending on the

species. In the type I bacterial cellulose synthase complex, the BcsD (bacterial cellulose synthase D)

protein is believed to play an important role in producing cellulose with long slender fiber morphology

and high crystallinity, given the phenotype of the *bcsD*-deficient mutant and the specific existence of the

type I operon in bacterial species synthesizing crystalline cellulose microfibrils such as Acetobacter. In

this study, we successfully established a heterogeneously expressed Bcs protein in Escherichia coli as

an experimental model and conducted biochemical studies for the BcsD protein and the other three

major subunits of bacterial cellulose synthase, BcsA, BcsB, and BcsC. It has been shown that the BcsD

protein interacts with the functionally required minimal subunits of the BcsAB complex, as well as the

BcsC protein. Furthermore, it was shown that BcsD interacts with the BcsAB complex in two modes:

direct protein-protein interactions and indirect interactions through the product cellulose. The former and

latter modes represent the basal and active states of the type I bacterial cellulose synthase, respectively.

This dynamic behavior of the BcsD protein is important for the type I bacterial cellulose synthase

complex to regulate the crystallization process of cellulose.

Introduction

Cellulose is a natural polymer, whose molecular structure is composed of a homopolymer of glucose with a $\beta1\rightarrow4$ linkage, which is very simple. However, cellulose molecules are insoluble in water, and the apparent elementary structure of cellulose is an aggregation of many cellulose molecules in long crystalline fibers. This structural feature of cellulose underlies its structure-sustaining property, which many cellulose-producing organisms rely on.

This polymer assembly is produced by the enzyme "cellulose synthase." In many known cases, *in vivo* cellulose synthase is a protein complex with several polypeptide subunits. The most important subunit in this complex is glycosyltransferase, which is called CesA. The first structural biological study of the CesA protein was reported in 2013 for cellulose synthase from the bacterium *Rhodobacter sphaeroides* (1,2), which has stimulated many biochemical studies on cellulose synthase.

Given its simplicity, bacterial cellulose synthase (Bcs) has been a model in many studies on cellulose synthase. Particularly, Bcs from bacteria belonging to the genus *Acetobacter*, some of which are currently named *Gluconacetobacter* or *Komagataeibacter* (the genus name "*Acetobacter*" is used for these bacteria in this study), have been used extensively in studies. Although it is now known that many other bacteria produce cellulose, the function and structure of cellulose appears to vary with species (3). For example, phosphoethanolamine cellulose has been reported in *Escherichia coli*, and this natural cellulose derivative is important for interacting with the host organism (4).

Among cellulose-synthesizing bacteria, *Acetobacter*, which has a gene cluster or operon categorized as

type I (3), is characteristic of synthesizing highly crystalline cellulose microfibrils (5) as well as plant

cellulose, a major natural source of cellulose. Previous studies have reported that the Bcs-complex in

Acetobacter is composed of at least four subunits (6,7) – BcsA (bacterial CesA, catalytic subunit), BcsB

(auxiliary subunit having a carbohydrate-binding domain and flavodoxin domain), BcsC (secretion of a

glucan chain through the outer membrane), and BcsD (crystallization subunit). In addition, endoglucanase

CMCax (8,9) (glycoside hydrolase family 8 (GH8)), unknown function protein Ccp (8,10) (also named as

BcsH (3)), and β-glucosidase bglAx (GH3) (11) are considered to be functionally involved in the complex.

Given that the cooperation of these subunits produces highly crystalline cellulose microfibrils, the spatial

organization of these subunits in the cell is important for understanding the cellulose biosynthesis mechanism

as a protein function (12).

In these subunits, the BcsD protein is found specifically in bacterial species producing highly crystalline

cellulose microfibrils such as *Acetobacter* and *Asaia* (3,13,14), which possess a type I Bcs-operon (3). The

bcsD-deficient mutant of *Acetobacter* showed a reduction in cellulose production, as well as a lack of ability

to synthesize crystalline cellulose microfibrils (6,7,15). Sunagawa *et al.* also demonstrated that BcsD

interacts with Ccp (BcsH) in a pull-down assay (16). Given that the BcsD protein contains glucan chains

inside its octamer structure (15), it has been proposed that BcsD is functionally located downstream of the

BcsAB complex (the cellulose-polymerizing core of the Bcs-complex (1,6,17,18)) (19,20). Detailed

information on how these functional units work together is key to understanding the mechanism by which

cellulose synthase can bundle the synthesized glucan chains for spinning a long slender crystalline fiber at

room temperature in an aqueous environment.

In this study, we analyzed protein-protein interactions in the type I Bcs-complex using the recombinant

BcsABCD protein with the *E. coli* expression system and clarified the interaction of BcsD with BcsAB

complex as well as BcsC protein. One of the advantages of using recombinant proteins is that mutant proteins

are easily available (21). Owing to this advantage, this study sheds light on the interaction between BcsAB

and BcsD and demonstrates that there are two modes for this interaction: direct protein-protein interactions

and indirect interactions through the cellulose molecules. We hypothesized that such bimodal interactions

between BcsAB and BcsD proteins could be key to controlling cellulose synthesis dynamically *in vivo*.

Results

Production of endogenous *bcsA* gene-deficient *E. coli*

The expression of the *Acetobacter* Bcs complex in *E. coli* cells might lead to contamination by endogenous

E. coli Bcs (22).We then attempted to delete *bcsA* in the *E. coli* host, which encodes the catalytic subunit of

Bcs. The deletion of the *bcsA* gene in *E. coli* BL21, a strain expressing the Bcs recombinant protein in this

study, was performed using the Red/ET recombination system (Gene Bridges GmdH, Germany), in which

the *bcsA* gene was replaced by the FRT-PGK-gb2-neo-FRT cassette with homology arms of the *bcsA*-flanking

region using homologous recombination (Fig. 1A). This genetically modified BL21 (BL21-GM) strain was

confirmed by colony direct polymerase chain reaction (PCR) to contain the neomycin-resistance gene

cassette (FRT-PGK-gb2-neo-FRT cassette) (Fig. 1B). When amplified with primer #2 (antibiotic gene

cassette-specific) and #3 (*E. coli*-specific), a band of approximately 1,500 bp was found in BL21-GM, while

no specific product band was found in wild-type BL21, indicating the insertion of the neomycin-resistance

gene cassette specifically in BL21-GM. However, when amplified with primers #1 and #3 (either *E. coli*-

specific), wild-type BL21 and BL21-GM produced a band corresponding to the *bcsA* gene (2,818 bp) and

the inserted gene cassette (1,731 bp), respectively. These results confirmed the successful deletion of the

bcsA gene of BL21 in "BL21-GM."

Next, we measured the *in vivo* cellulose synthesis activity of the prepared BL21-GM (Fig. 1C) by partially

using our previous method (18): diguanylate cyclase (DGC) fused to thioredoxin ($_{Trx}$DGC) was expressed in

BL21 and BL21-GM to synthesize c-di-GMP to activate endogeneous bacterial cellulose synthase. The

amounts of cellulose synthesized were approximately 2.3 and 0.7 μg/mL of culture in BL21 and BL21-GM,

respectively. Although weak cellulose-synthesizing activity was observed, we used BL21-GM as a suitable

host to express the Bcs subunits with minimal impact on the endogenous Bcs-complex of *E. coli*.

Characterization of BsCel5A

Cellulase is an important tool used in this study to scrutinize the role of cellulose molecules in the subunit-

subunit interaction of the Bcs-complex. In this study, we used a bacterial endo-β-1,4-glucanase from *Bacillus*

subtilis (BsCel5A), whose structure has been previously reported (23). The full length (FL) of this cellulase

is a two-domain protein, in which the catalytic domain of GH5 is linked to the carbohydrate binding module

of family 3 (CBM3). In this study, the catalytic domain alone (CD) was also used to maximize access to the

soluble substrate. FL and CD of BsCel5A were expressed as glutathione-*S*-transferase (GST)-fused proteins

and purified with glutathione affinity resin, as confirmed by SDS-PAGE (Fig. S1A, left panel); (Fig. S1A,

left panel, white arrow). Activity-deficient mutant of E169Q/E257Q in FL (BsCel5A_FLm) and CD

(BsCel5A_CDm) was also used for a negative control protein in the cellulase treatment experiment.

The binding activity of the purified BsCel5A to phosphoric acid-swollen cellulose (PASC) was surveyed

to confirm correct folding of the expressed proteins. The results showed that all the BsCel5A proteins used

in this study retained their binding activity to PASC (Fig. S1A, right panel), indicating that all of them were

properly folded despite sequence modification. The enzymatic activity of BsCel5A was surveyed to

determine whether this cellulase could function at low temperature and neutral pH, which is required for

analyzing the Bcs-complex. Both BsCel5A_FL and BsCel5A_CD hydrolyzed barley β-glucan, carboxyl

methyl cellulose (CMC), and PASC at 4℃ and pH 7.5 (Fig. S1B). For Avicel, full-length BsCel5A showed

higher activity than the others, stressing the function of CBM to bind solid cellulose.

Substrate specificity of BsCel5A was investigated using cello-oligosaccharides with various degrees of

polymerization (DP). BsCel5A hydrolyzed cellotetraose, cellopentaose, and cellohexaose but not cellobiose

or cellotriose (Fig. S1C). The final reaction products of BsCel5A for substrates longer than triose were

cellobiose and cellotriose, and the smallest substrate was cellotetraose (Fig. S1D).

Summarizing these, both BsCel5A_FL and BsCel5A_CD were able to hydrolyze cellulose of low

crystallinity at 4℃ and pH 7.5, and cellulose molecules with DP higher than 3 were the substrates. This

information was considered when interpreting the data.

In vivo cellulose synthase activity of recombinant Bcs-complex

This study analyzed heterogeneously expressed Bcs proteins in *E. coli* as an experimental model. We then

checked if these heterogeneously expressed proteins were functional by using CESEC (cellulose-

synthesizing *E. coli*), a platform for estimating the cellulose-synthesizing activity with the co-expression of

diguanylate cyclase (DGC) (18). All the Bcs-complexes expressed in DGC-expressing BL21-GM

synthesized a significant amount of cellulose (Fig. 2A), indicating that each of the Bcs-complexes expressed

in this study are functional, regardless of whether BcsD is tagged. In addition, neither BcsAB nor the

BcsABCD complex with the activity-deficient mutant of the catalytic subunit BcsA (D333N or W373A)

showed cellulose-synthesizing activity, consistent with our previous study (21). These results support the

idea that the Bcs proteins expressed in this study reflect the state in function and are suitable for an

experimental model.

Despite the lower cellulose production shown in Fig. 2A, we decided to express Bcs proteins without co-

expression of DGC for subsequent purification because higher expression levels of Bcs proteins were

detected by western blotting (WB), as shown in Fig. 2B. Given that the cellulose quantification in this study

used strong acid treatment with acetic acid and nitric acid, we think that highly crystalline cellulose was

majorly considered in the data presented in Fig. 2A. Then it cannot be ruled out that the Bcs-complex

expressed in *E. coli* without DGC expression synthesizes cellulose.

The complex formation of BcsAB, BcsC, and BcsD

Given that BcsD holds the glucan chains in its channel (15), it is functionally located downstream of the

BcsAB complex, the core of cellulose molecule synthesis, indicating that BcsD operates in the periplasmic

space to assemble cellulose molecules extruded from the BcsAB complex. Furthermore, the BcsC protein

has a β-barrel porin-like structure at its C-terminus and is supposed to secrete cellulose through the outer

membrane (24). The results from several studies suggest that BcsAB, BscC, and BcsD work together in a

coordinated manner to spin bacterial cellulose. We tried to co-express BcsABCD to purify the whole Bcs-

complex using a hexa-histidine tag attached to the C-terminus of BcsD (BcsD$_{H6}$), as the tag at the C-terminus

does not disturb cellulose-synthesizing activity (16). After the optimization of the conditions (Fig. S2), all

four subunits were successfully purified together by immobilized metal affinity chromatography (IMAC)

(Fig. 3). This is the first successful purification of the BcsABCD complex ever reported. However, BcsC was

found to be mostly insoluble in detergent (*n*-dodecyl-β-D-maltoside, DDM, in this study) and tended to

degrade or aggregate during the purification process (Fig. 3A, BcsC antibody strip). We did not decipher

BcsC behavior in this Bcs-complex purification experiment.

Analysis of the interaction between BcsAB and BcsD by cellulase treatment

As described above, co-expression of BcsABCD$_{H6}$ in *E. coli* allowed the BcsABD-complex to be purified.

Given that BcsABCD$_{H6}$ expressed in *E. coli* is functional as shown in Fig. 2A, we speculated that a glucan

chain may link BcsAB and BcsD during this purification. This hypothesis was tested by checking whether

the digestion of cellulose resulted in the dissociation of BcsAB from BcsD$_{H6}$ during IMAC purification.

BsCel5A cellulase was applied to BcsABCD$_{H6}$ bound to IMAC resin prior to washing and elution. The

amounts of BcsA, BcsB, and BcsD proteins in the eluate were measured by semi-quantitative WB (Fig. 4A);

the reliability of quantification in our WB was confirmed, as shown in Fig. S3. The signal intensities of BcsA

and BcsB were normalized to that of BcsD in the eluate for comparison between samples. Notably, the

amounts of BcsA and BcsB bound to BcsD were decreased by BsCel5A treatment (Fig. 4B), indicating that

the BcsAB complex was detached from BcsD by cellulose digestion. This trend was clearer when

BsCel5A_CD was used as cellulase.

To rule out the possibility that the observed detachment of BcsAB was due to non-specific interaction of

the added BsCel5A protein, the activity-deficient mutant of BsCel5A (E169Q/E257Q) was applied to

BcsABCD$_{H6}$ bound to the metal affinity resin (Fig. 4, C and D). As a result of purification, no decrease in

the signal intensity of BcsA and BcsB was observed in the eluate, indicating that cellulose hydrolysis actually

caused detachment of BcsAB from BcsD during purification. In summary, the BcsAB complex indirectly

interacts with BcsD through the glucan chain.

Although cellulase treatment allowed the BcsAB complex to be detached from BcsD during purification,

as demonstrated above, a substantial portion of the BcsA and BcsB proteins remained with BcsD in this

purification process (Fig. 4B). We supposed that a direct interaction between BcsAB and BcsD is also found

in addition to the indirect interaction. To test this hypothesis, we attempted to purify the activity-deficient

mutants of BcsABCD$_{H6}$, BcsA^{D333N}BCD$_{H6}$, and BcsA^{W373A}BCD$_{H6}$, neither of which synthesize cellulose, as

shown in Fig. 2A. As a result, these mutants allowed BcsA and BcsB to be purified together with BcsD and

the wild type (Fig. 5A and B). Notably, cellulase treatment did not reduce the amount of BcsAB purified

together with BcsD for either $BcsA^{D333N}BCD_{H6}$ or $BcsA^{W373A}BCD_{H6}$ (Fig. 5C, 5D, and 5E). These results

support the existence of a direct interaction between BcsAB and BcsD when cellulose is not synthesized.

Compiling these cellulase treatment experiments, it was demonstrated that there are two modes of interaction

between BcsAB and BcsD: direct and indirect modes through cellulose molecules.

Quaternary structure of the Bcs-complex

Despite the demonstration of two modes for the interaction between BcsAB and BcsD, the quaternary

structure of the entire Bcs-complex remains to be elucidated. We then performed pull-down assays to further

explore the subunit-subunit interactions in the BcsABCD complex by fully utilizing recombinant Bcs

proteins. The assay was designed based on the results from previous studies (Fig. 6A): (i) BcsAB forms a

complex in the inner membrane (1), (ii) BcsC is an outer membrane protein (24,25), (iii) BcsD protein is

located in the periplasmic space (15,20), and (iv) BcsD interacts with the BcsAB complex (this study).

First, the interaction between BcsAB and BcsD, which has been demonstrated above, was tested by pull-

down with a bait of Streptag(II)-fused BcsD ($BcsD_{StrepII}$) and a prey of crude BcsAB without tags (DDM-

solubilized cell membrane of *E. coli* expressing BcsAB). As expected, BcsAB bound to $BcsD_{StrepII}$ (Fig. 6B).

The pull-down assay yielded a result consistent with our purification experiment of the BcsABD-complex

and supported our hypothesis for the interaction between BcsAB and BcsD.

Given the successful pull-down assay with BcsAB and BcsD$_{StrepII}$, we next performed a pull-down assay

to survey the BcsC-involved interaction in the complex (see Fig. 6A). For this purpose, TPR tetratricopeptide-

repeat (TPR) domain of the BcsC protein (BcsC_TPR) was used based on two assumptions: (i) the role of

the TPR domain is protein-protein interaction (26), and (ii) the full-length BcsC protein is unstable, as shown

in our purification. Pull-down by BcsD$_{StrepII}$ trapped $_{H6}$BcsC_TPR$_{H6}$, which has hexa-histidine tags at both

the N- and C-termini of BcsC_TPR, in a dose-dependent manner (Fig. 6C), clearly showing that these two

proteins interact with each other. Furthermore, pull-down by the purified $_{H6}$BcsC_TPR$_{H6}$ (bait) with the crude

BcsAB (prey) also supported their interaction with each other (Fig. 6D), although it appeared less striking

than the case with $_{H6}$BcsC_TPR$_{H6}$ and BcsD$_{StrepII}$ (Fig. 6C).

Finally, we performed a pull-down assay using all subunits in the complex: crude BcsAB, purified

$_{H6}$BcsC_TPR$_{H6}$, and purified BcsD$_{StrepII}$ (Fig. 6E). Streptactin beads trapped BcsA, BcsB, and $_{H6}$BcsC_TPR$_{H6}$

together with BcsD$_{StrepII}$, although BcsA was only faintly detected by WB. It is then summarized that the

BcsD protein can simultaneously interact with BcsA, BcsB, and the TPR region of BcsC.

Discussion

Bacterial cellulose synthase is a heterosubunit complex. Primary evidence for this hypothesis is the

existence of gene clusters or operons encoding cellulose synthase-related genes (3,6,7,11). However, this

hypothetical model has not been tested intensively, owing to the lack of studies on the protein. Two recent

reports have addressed this question. Sunagawa et al. demonstrated an interaction between BcsD and BcsH

(Ccp) proteins using a pull-down assay (16). Another prominent study was the X-ray crystallographic

analysis of the BcsAB protein complex by Morgan et al., which clarified that the BcsA and BcsB proteins

closely interact with each other to polymerize and translocate the glucan chain from the cytosol to the

extracellular side (1), as confirmed also in *E.coli* (27). In this study, we analyzed the type I Bcs complex and

demonstrated that the BcsD protein interacted with the BcsAB complex (Fig. 3, 4, 5, and Fig. 6B) and the

TPR domain of BcsC (Fig. 6C). Furthermore, it was shown that BcsC_TPR interacted with the BcsAB

complex, although the result was not as clear as the interaction between BcsC_TPR and BcsD (Fig. 6D).

Finally, a pull-down assay using $BcsD_{StrepII}$ with a mixture of crude BcsAB and purified $_{H6}BcsC\_TPR_{H6}$

indicated that BcsD binds to both BcsAB and BcsC_TPR simultaneously, suggesting that these four subunits

interact with each other (Fig. 6E). Our study, together with previous studies, experimentally demonstrated

that the subunits of BcsA, BcsB, BcsC, BcsD, and BcsH are included in the type I Bcs-complex.

Interestingly, this study showed that the interaction of BcsD with BcsAB has two modes: direct and indirect

interactions through the polymerized glucan chain (Fig. 4). Direct interaction between the BcsAB complex

and BcsD protein was observed for both wild type and catalytically deficient mutant cellulose synthase,

whereas the interaction via cellulose was observed only for the wild type (Fig. 5). We then propose a two-

state model for the type I Bcs complex: basal and active states, which are represented by the direct interaction

(Fig. 7, left) and indirect interaction modes (Fig. 7, right), respectively.

BcsD protein in this basal state is statically fixed on the BcsAB complex for lying in wait for a glucan

chain synthesized by the BcsAB complex. Once cellulose synthesis is initiated, the synthesized glucan chain

is guided into the octameric ring of BcsD to crystallize cellulose molecules, as proposed previously

(7,15,16,20,28), which is a shift from the resting state to the active state of the complex. In the active state

(Fig. 7, right), the BcsD protein is detached from the BcsAB complex but tethered to BcsAB by the glucan

chain extruded from BcsAB. Considering the substrate specificity of BsCel5A (Fig. S1), BcsD can be

physically separated from BcsB by a glucan chain with a length of four or more glucose residues (Fig. 7).

In this two-state model, the active state of the Bcs complex allows BcsD to be launched toward the outer

membrane, while BcsD in the basal state may be fixed statically on BcsB. The TPR domain of the BcsC

protein may control diffusible BcsD in the active state in the periplasmic space by gentle but significant

interactions. Given that the C-terminal domain of BcsC is an outer membrane β-barrel channel (Fig. S4) (24),

the TPR domain of BcsC may escort BcsD carrying glucan chains toward the outer membrane channel in

BcsC to spin the glucan chains to the extracellular side. The following questions about this model will be

whether (i) BcsD protein could dock to the outer membrane channel part of the BcsC protein and (ii) the

interaction between BcsC and BcsD could be state-dependent. These questions will be addressed in future

studies to enhance our understanding of the Bcs-complex function.

Our study demonstrated the intermolecular interactions of subunits in the BcsABCD complex to update

our knowledge of the quaternary structure of the type I Bcs complex. Experiments in this study were

performed using recombinant proteins heterogeneously expressed in *E. coli*; thus, further studies will be

necessary to test the hypothesis proposed in this study, the two modes for BcsAB-BcsD interaction: the basal

and the active state. Nonetheless, the data obtained in this study were intensively tested using mutagenesis

experiments of proteins, and they were consistent with each other. We believe that our hypothesis in this

study is sufficiently reliable, at least for starting further discussion about the mechanism of Bcs with regard

to protein function. In this model, BcsD is not statically fixed in the Bcs-complex but dynamically interacts

with the other subunits in the complex (Fig. 7). Such a dynamic behavior may be involved in the

crystallization function of cellulose molecules, as well as twisting in bacterial cellulose microfibrils, as

reported previously (16). The putative state control with BcsD, as shown in Fig. 7, is the regulation of the

quaternary structure of bacterial cellulose synthase. Next, we need to further study the original *Acetobacter*

cells and observe the whole Bcs-complex at higher resolution with cryo-EM under controlled two-state

conditions.

Experimental procedures

Chemicals

All chemicals used in this study were purchased from FUJIFILM Wako Pure Chemicals Co. Ltd. (Osaka,

Japan), Nacalai Tesque Co. Ltd. (Kyoto, Japan), and Sigma-Aldrich Inc. (MO, USA), unless otherwise

indicated.

Cloning

Bcs genes (*bcsA*, *bcsB*, *bcsC*, and *bcsD*) were prepared from genomic DNA extracted from cultured cells

of *Komagataeibacter sucrofermentans* JCM9730, which was provided by the RIKEN Bio Resource Center

(Tsukuba, Japan) and known by the strain name BPR2001 (29,30). The endo-β1,4-glucanase gene from

Bacillus subtilis strain 168 (*bscel5A*) (23) was prepared from the genomic DNA provided by the American

Type Type Culture Collection (ATCC23857D). The dgc gene (*dgc*) was identical to that used in our previous study

(18): tDGC and DGC from *Thermotoga maritima* MSB8 with a point mutation of R158A and truncation of

the N-terminal 81 amino acid residues from V1 to V81 (31). The gene for green fluorescent protein UV

(GFPuv) (32) was prepared from the pGFPuv plasmid purchased from Takara Bio, Co. Ltd. (Shiga, Japan).

The cloning of these genes was performed by PCR using PrimeStarGXL (Takara Bio Co. Ltd.). Site-

directed mutagenesis was performed by QuikChange II (Agilent Inc., CA, USA) or the equivalent PCR-based

method. The sequence of the PCR-amplified DNA was checked by the dye-terminator method in a subcloning

vector (pGEM-T easy (Promega Inc., WI, USA) or pTAC-2 (BioDynamics Laboratory Co. Ltd., Tokyo,

Japan)), or final expression vector, which was performed at either of the Uji DNA-sequencing core (Kyoto

University Research Administration Office) with an ABI Prism 310 Genetic Analyzer (Applied Biosystems

Inc., MA, USA), or Eurofins Genomics Inc. (Tokyo, Japan).

Construction of the expression vectors

The protein expression vectors used in this study are summarized in Table 1. Expression vectors for BcsAB

(BcsAB, BcsA^{D333N}B, and BcsA^{W373A}B) were identical to those used in previous studies (18,21). The *bcsAB*

gene in the operon form was inserted between the EcoRI and KpnI sites in pQE-80L (Qiagen, Hilden,

Germany), and the upstream of the start codon was the original ribosomal binding site of *K. sucrofermentans*

JCM9730.

Expression vectors for BcsABCD (BcsABCD, BcsABC$_{H6}$D, BcsABCD$_{H6}$, BcsA^{D333N}BCD$_{H6}$,

BcsA^{W373A}BCD$_{H6}$, and BcsABCD$_{GFPuvH6}$) proteins were constructed from the vectors for BcsAB by inserting

the *bcsABCD* gene in the operon form between the EcoRI and HindIII sites in pQE-80L. The ribosomal

binding site in these BcsABCD expression vectors is exactly the same as that in the expression vectors of

BcsAB, one of *K. sucrofermentans* JCM9730. DNA encoding the tag (hexa-histidine or GFPuv) was added

to *bcsD* during the PCR-amplification amplification.

The expression vectors for the TPR region of BcsC (BcsC_TPR: C18-G848) and BcsD were prepared for

the pull-down assay. DNA encoding hexa-histidine tag was added to the 3′-end of the gene for BcsC_TPR

during PCR amplification and inserted into the NdeI and XhoI sites in pET-28b (Merck Corp.., Darmstadt,

Germany) to express BcsC_TPR protein with dual hexa-histidine tag at both N- and C- terminal –

$_{H6}$BcsC_TPR$_{H6}$. DNA encoding Streptag (II) was added to the 3′-end of the gene for BcsD and inserted

between the NcoI and HindIII sites in pET28b to express BcsD with C-terminal Streptag(II)–BcsD$_{StrepII}$.

The gene for tDGC was inserted between the NdeI and HindIII sites in pBAD33et32, a customized plasmid

DNA in this study, to express the thioredoxin-fused tDGC protein to count the cellulose-synthesizing activity

of the recombinant BcsAB and BcsABCD complex in our previous studies (18,21). The detailed procedure

for preparing pBAD33et32 is described in Supplementary Information. Thioredoxin gene (*trx*) in

pBAD33et32 was copied from pET32a (Merck Corp.) to replace the His-patch thioredoxin used in our

previous study (18), which was derived from pBAD202-TOPO (Invitrogen Inc., MA, USA).

For the bacterial endoglucanase gene *bscel5A*, the putative signal peptide sequence (M1–A28) was

removed from the full-length construct. Full-length (A29–N499) and catalytic domain (A29–T332) genes

were inserted between BamHI and XhoI sites in pGEX4T-3 (Cytiva Inc., MA, USA) to express the N-

terminal GST-fused endoglucanase BsCel5A.

Deletion of *bcsA* gene from *Escherichia coli* BL21

In this study, we used genetically modified *E. coli* BL21 to express Bcs proteins, which lack the whole

bcsA gene (Fig. 1A). This gene manipulation was performed using the Red/ET recombination system (Gene

Bridges GmbH, Heidelberg, Germany), according to the manufacturer's specifications. The detailed

procedure is described in Supplementary Information.

Briefly, *E. coli* BL21 strain (purchased from New England Labs Inc., MA, USA) was transformed with

the pRedET plasmid carrying the genes for DNA-recombination proteins and selected by tetracycline

resistance. The second transformation was followed by introducing linear DNA of neomycin-resistant gene

flanked by the homology arm sequences corresponding to the upstream and downstream of *bcsA* gene

(CP010816.1); the recombinant proteins were expressed with arabinose-induction for disrupting *bcsA* gene,

and subsequently the pRedET plasmid was expelled out of the cell by culturing at 37°C. Finally, the *bcsA-*

disrupted BL21 transformant was selected for kanamycin resistance and tetracycline sensitivity. Gene

modification was confirmed by direct PCR (Fig. 1B). We named this genetically modified *E. coli* BL21 strain

as BL21-GM. Chemically competent cells of BL21-GM were then prepared by treating the cultured cells in

a rubidium-based buffer according to the protocol on the website of the Center for Molecular Biology and

Genetics at Mie University (https://www.gene.mie-u.ac.jp/Protocol/Original/Transform-Comp-Cell-

Freeze.html, in Japanese) (33).

***In vivo* cellulose synthase activity**

The endogenous cellulose-synthesizing activity of *E. coli* and cellulose-synthesizing activity of the

heterogeneously expressed Bcs-complex in *E. coli* was estimated by expressing DGC protein to activate

bacterial cellulose synthase in the cell by the produced c-di-GMP (18). In this study, the pBAD33et32-tDGC

plasmid was used to express the DGC. The obtained *E. coli* transformants were pre-cultured in LB medium

including the proper antibiotics at 37°C overnight, and then 0.1 mL of the pre-culture was inoculated into 10

mL of ZYP-5052 medium with 50 μg/mL chloramphenicol (100 μg/mL ampicillin when expressing Bcs-

complex) and 0.001% L-arabinose. Cells were grown at 20°C in an orbital shaker at 180 rpm for 24 h and the

cultured cells were collected by centrifugation at $2,000 \times g$ for 20 min. The amount of cellulose in the pellet

was quantified using the anthrone-sulfuric acid method (34) as described below.

Two milliliters of AN-reagent (a mixture of glacial acetic acid and nitric acid with volumetric ratio of 8:1)

were added to the pellet and then heated at 105°C for 30 min. The remaining cellulose pellets were washed

with 10 mL of water once by centrifugation at $2,000 \times g$ for 10 min, and then 1 mL of 72% (v/v) H_2SO_4 was

added to the pellets and incubated for 1 h at room temperature. The solution (0.1 mL) was mixed with water

(0.4 mL), and then, 1 mL of fresh anthrone reagent (0.2% w/v anthrone in concentrated H_2SO_4) was added

and mixed. The solution was heated to 90°C for 15 min and allowed to stand until it reached room temperature

(approximately 15 min). The absorbance of the solution was measured at both 625 and 700 nm, and the

absorbance at 625 nm, from which the absorbance at 700 nm was subtracted, was used for quantification for

better reproducibility. Calibration was performed with phosphoric acid-swollen cellulose (PASC) prepared

using a previously reported protocol (35).

Purification and characterization of BsCel5A

Expression of GST-fused BsCel5A endoglucanase was performed in *E. coli* BL21 (not GM strain): all the

four constructs ((BsCel5A_FL, BsCel5A_CD, BsCel5A_FL^m, and BsCel5A_CD^m) were expressed and

purified in the same manner.

The transformant of BL21 with pGEX-BsCel5A were inoculated in 10 mL LB containing 100 μg/mL

ampicillin in a 100 mL flask and grown at 37°C with orbital shaking at 220 rpm overnight. The cells were

harvested by centrifugation at 3,000 × g for 20 min, suspended in 1 L of LB containing 100 μg/mL ampicillin

in a 5 L baffle flask, and cultured at 37°C until OD$_{600}$ reached 0.6. The culture medium was cooled in ice

water for 10 min, and then IPTG was added to 0.1 mM and incubated at 20°C for 20 h at 160 rpm with

shaking.

The cells were then collected at 3,000 × g for 20 min, washed with 40 mL of 1 × phosphate-buffered saline

(PBS), suspended in 40 mL of 1 × PBS containing 1 mM phenylmethylsulfonyl fluoride (PMSF), and

sonicated on ice for ~5 min (exposure: 10 s, rest: 50 s; until slightly translucent). The cell lysate clarified by

20,000 × *g* for 30 min at 2°C was applied to 5 mL of Glutathione Sepharose 4B (Cytiva Inc,), washed with

20 mL of 1 × PBS, and eluted with 5 c.v. of 0.1 M Tris-HCl pH 8.0, 0.5 M NaCl, 20 mM reduced-glutathione.

The eluted fraction was buffer exchanged with 1 × PBS using Vivaspin 6 (MWCO 10 kDa; Sartorius AG,

Germany).

To remove the GST tag fused to the N-terminus, 100 units of thrombin (Merck Corp.) was added to 10 mg

of $_{GST}$BsCel5A and incubated at 20°C for two days. Thrombin was then removed by adding 1 mL of

Benzamidine Sepharose 6 B (Cytiva Inc.), and the cleaved GST was removed by adding 5 mL of Glutathione

Sepharose 4 B. A solution of PMSF was added to the obtained BsCel5A protein at a final concentration of 1

mM to inactivate a small amount of residual thrombin.

The recombinant BsCel5A proteins purified were characterized with regard to the following points: (i)

binding activity to cellulose, (ii) glycoside hydrolase activity, and (iii) substrate specificity. The detailed

experimental procedure is described in Supplementary Information.

Purification of BcsABCD complex

All the protein purification experiments in this study were performed at low temperature (~4°C). The

composition of the protease inhibitor mix (PI) used for protein purification was as follows: 0.1 mM PMSF, 1

mM benzamidine-HCl, 5 μM bestatin, 10 μM leupeptin, 1 μM pepstatin A.

The transformant of *E. coli* BL21-GM with pQE-BcsABCD$_{H6}$ was inoculated in 10 mL of LB containing

100 μg/mL ampicillin in a 100 mL flask and grown at 37°C with shaking at 220 rpm overnight. The cells

were harvested by centrifugation at 3,000 × g for 20 min, resuspended in 1 L of ZYP-5052 containing 100

μg/mL ampicillin and 0.0005% (v/v) antifoam SI (Fujifim Wako Chemicals Co. Ltd., Japan) in a 5 L baffle

flask, and incubated at 20°C for 22 h with an orbital shaker at 100 rpm. The cultured cells were collected at

3,000 × g for 20 min and suspended in 50 mL of buffer A (20 mM sodium phosphate buffer (pH 7.5), 300

mM NaCl, 8% glycerol, 5 mM MgCl$_2$, 5 mM cellobiose) with PI, prior to cell lysis by two passes of high-

pressure homogenizer (French Press, Thermo Fisher Scientific Inc., MA, USA) at a pressure of 15,000 psi.

The lysate clarified by centrifugation at 5,000 × g for 10 min and ultracentrifugation at 150,000 × g for 30

min at 2°C. The obtained cell membranes were resuspended in buffer A with PI and NaCl concentrations

increased to 1 M and were ultracentrifuged again under the same conditions. Finally, the cell membrane was

suspended in buffer A with PI to approximately 10 mg/mL (estimated by the DC protein assay, Bio-Rad Inc.,

CA, USA). The collected cell membrane was stored at -80°C until use in experiments.

Ten milliliters of the cell membrane suspension were mixed with n-dodecyl-β-D-maltopyranoside (DDM)

at a final concentration of 0.5% (w/v) and shaken gently for 1 h at 4°C for solubilizing Bcs-complex.

Supernatant of the following ultracentrifugation at 150,000 × g for 30 min at 2°C was collected as solubilized

fraction, and imidazole stock solution (2 M, pH 8.0) was added to achieve a final concentration of 10 mM,

prior to loading to 0.4 mL a His-Select Cobalt Affinity Gel (Sigma-Aldrich Inc.), which was equilibrated

with 4 mL of wash buffer (20 mM imidazole and 0.025% (w/v) DDM in buffer A) in advance. After loading

the solubilized protein onto the affinity resin in a plastic open column, the resin in the column was washed

with 3 mL of wash buffer and finally eluted with elution buffer (200 mM imidazole and 0.025% (w/v) DDM

in buffer A).

Cellulase treatment of Bcs-complex

Cellulase treatment of BcsABCD$_{H6}$ was performed using a cobalt affinity gel. The same procedure was

followed for the binding step of BcsABCD$_{H6}$ on the affinity gel. After all the solubilized fraction passed

through the column, 0.8 mL of 1.5 μM BsCel5A solution in wash buffer (20 mM imidazole and 0.025% (w/v)

DDM in buffer A) was loaded to the column. The column exit was plugged when most of the cellulase

solution passed, and then the gel was incubated at 10°C for 3 h with the remaining cellulase solution. After

unplugging, the affinity gel in the column was washed with 2 mL of wash buffer and eluted with elution

buffer (200 mM imidazole and 0.025% (w/v) DDM in buffer A).

Protein preparation for pull-down assay: BcsC_TPR, BcsD, and crude BcsAB

The pull-down assay in this study used purified BcsC_TPR, purified BcsD, and crude BcsAB proteins,

each of which was prepared as described below.

The transformant of *E. coli* BL21(DE3) with pET-$_{H6}$BcsC_TPR$_{H6}$ was inoculated in 10 mL of LB

containing 50 μg/mL kanamycin in a 100 mL flask and grown at 37°C with shaking at 220 rpm overnight.

The cells were harvested by centrifugation at 3,000 × *g* for 20 min, and resuspended in 1 L of ZYP-5052 (36)

containing 200 μg/mL kanamycin and 0.0005%(v/v) antifoam SI in a 5 L baffle flask, and incubated at 20°C

for 24 h in a shaking incubator at 100 rpm. The cultured cells were collected at 3,000 × *g* for 20 min, and

washed with 40 mL of 1 × PBS prior to resuspension in 100 mL of 10 mM imidazole in buffer B (20 mM

sodium phosphate buffer (pH 7.5), 500 mM NaCl, 8% glycerol) with PI, and sonicated on ice for ~5 min

(exposure: 10 s, rest: 50 s; until slightly translucent). The cell lysate was clarified by centrifugation at 40,000

× g for 30 min at 2°C and applied to 4 mL a His-Select Cobalt Affinity Gel (Sigma-Aldrich Inc.), followed

by washing with 10 mM imidazole in buffer B, and eluted 250 mM imidazole in buffer B. Finally, the buffer

of eluted fraction including $_{H6}BcsC\_TPR_{H6}$ protein was replaced with buffer B using PD-10 desalting column

(Cytiva Inc.).

The BcsD protein was expressed in $E.\ coli$ BL21(DE3) with pET-BcsD$_{StrepII}$. The transformant was

inoculated in 10 mL of LB containing 50 μg/mL kanamycin in a 100 mL flask and grown at 37°C with

shaking at 220 rpm overnight. The cells were harvested by centrifugation at 3,000 × g, 20 min, resuspended

in 1 L of LB containing 50 μg/mL kanamycin in a 5 L baffle flask and cultured at 37°C until OD$_{600}$ was ~0.6.

The culture medium was cooled in ice water for 30 min, and IPTG (0.5 mM) was added and incubated at

25°C for 20 h at 120 rpm. Cells were collected at 3,000 × g for 20 min, suspended in 30 mL of buffer C (20

mM Tris-HCl (pH 7.5), 150 mM NaCl, and 1 mM EDTA) with PI, and sonicated on ice for ~10 min

(exposure: 10 s, rest: 50 s; until slightly translucent). The cell lysate was centrifuged at 20,000 × g for 30 min

at 2°C, applied to 2 mL of Streptactin Sepharose HP (Cytiva Inc.), washed with 20 mL of buffer C, and eluted

with 4 mM d-desthiobiotin in buffer C. Finally, buffer of eluted fraction was replaced by buffer C using

Amicon Ultra-4 MWCO 50 kDa (Merck Corp.).

The crude protein of BcsAB was prepared using the transformant $E.\ coli$ BL21-GM with pQE carrying the

bcsAB gene. The procedure to isolate the cell membrane, from culture to ultracentrifugation, was same as in

the case of BcsABCD$_{H6}$ as described previously, except for the buffer to suspend the cells and cell membrane:

buffer D (20 mM sodium phosphate buffer (pH 7.5), 100 mM NaCl, 8% glycerol, 5 mM MgCl$_2$, and 5 mM

cellobiose) with PI was used. The obtained cell membrane was suspended in buffer D with PI at

approximately 10 mg/mL (estimated by the DC protein assay, Bio-Rad Inc.). Finally, crude BcsAB protein

was prepared by solubilizing 7 mL of the cell membrane fraction with 0.5% (w/v) DDM for 1 h and

centrifugation at 150,000 ×*g* for 30 min at 2℃.

Pull-down assays

For the pull-down by BcsD$_{StrepII}$ with the crude BcsAB, purified BcsD$_{StrepII}$ (0.2 mg) was added to 7 mL of

crude BcsAB, incubated for 3 h at 4 ℃, and then loaded to 0.1 mL of Streptactin Sepharose HP. The resin

was washed with 1 mL of 0.025% (w/v) DDM in buffer D, and finally the bound protein was eluted with 4

mM desthiobiotin in buffer D with 0.025% (w/v) DDM.

For the pull-down assay, BcsD$_{StrepII}$ with $_{H6}$BcsC_TPR$_{H6}$, BcsD$_{StrepII}$ (0.1 mg) were incubated with

$_{H6}$BcsC_TPR$_{H6}$ (0, 0.1, 0.3, 0.5 mg) in 3 mL of buffer D (–/–) (buffer D without MgCl$_2$ and cellobiose) for 3

h at 4 ℃, and then loaded to 0.1 mL of Streptactin sepharose HP. After washing with buffer D (–/–), the

bound protein was eluted with 1 mL 4 mM *d*-desthiobiotin in buffer D (–/–).

For the pull-down assay by $_{H6}$BcsC_TPR$_{H6}$ with BcsAB, purified $_{H6}$BcsC_TPR$_{H6}$ (0.2 mg) was mixed with

crude BcsAB (7 mL), and further NaCl stock solution was added to set 500 mM NaCl in the final solution.

475 After incubation for 1 h at 4 °C, the sample was loaded to 0.2 mL a His-Select Cobalt Affinity Gel and washed

476 with 2 mL of 20 mM imidazole in buffer B (+/+) (buffer B with 5 mM $MgCl_2$ and 5 mM cellobiose) with

477 0.025% (w/v) DDM. The bound protein was eluted with 200 mM imidazole in buffer B containing 0.025%

478 (w/v) DDM.

479 Finally, we performed a pull-down assay using $BcsD_{StrepII}$, purified BcsC_TPR; crude BcsAB. $BcsD_{StrepII}$

480 (0.2 mg) and $_{H6}BcsC\_TPR_{H6}$ (0.2 mg) were mixed with crude BcsAB (7 mL of DDM extract); NaCl stock

481 solution (5 M) was added to a final concentration of 500 mM NaCl. After incubation for 3 h at 4 °C, the

482 sample solution was loaded to 0.1 mL of Streptactin Sepharose HP. After washing with 1 mL of buffer B

483 (+/+) containing 0.025% (w/v) DDM, the bound protein was eluted with 4 mM d-desthiobiotin in buffer B

484 (+/+) containing 0.025% (w/v) DDM.

485 Prey protein without bait protein was used as a negative control for the pull-down assay using the same

486 procedures. The eluted fraction was analyzed by SDS-PAGE, followed by WB as described later.

488 **Western blotting (WB) analysis**

489 Rabbit polyclonal antibodies against BcsA, BcsB, BcsC, and BcsD proteins were generated with the

490 synthetic peptide corresponding to a part of the C-terminus of BcsA (18,20,21,37), the C-terminus of BcsB

491 (this study), the C-terminus of BcsC (37), and the loop between the β3 and β4-strand (20,37). The dilution

492 rate was 10,000-fold for anti-BcsA, anti-BcsB, and anti-BcsC, whereas anti-BcsD was diluted 20,000-fold.

493 Each of the antibodies against the peptide tag were diluted as follows: anti-Strep tag II (15 ng/mL; ab180957,

Abcam Corp., Cambridge, UK), anti-His5 (0.1 µg/mL; 34660, Qiagen N.V., Germany), and anti-Thioredoxin

(1 µg/mL; ab139677, Abcam Corp.).

Horseradish peroxidase (HRP)-conjugated anti-rabbit IgG (W401B; Promega Inc.) and HRP-conjugated

anti-mouse IgG (7076S; Cell Signaling Technology Inc., MA, USA) were diluted 20,000-fold and used as

the secondary antibodies. ECL-Prime (Cytiva Inc.) was used for chemical luminescence with HRP.

For the WB with anti-BcsD, anti-Strep, anti-His, and anti-thioredoxin, the samples for SDS-PAGE were

heat-treated at 100°C for 3 min after mixing with the sample buffer, while the samples to visualize by the

other antibodies were incubated in the sample buffer for several minutes at room temperature. The samples

were analyzed by SDS-PAGE with a precast polyacrylamide gel (SuperSep Ace 5-20%, Fujifim Wako

Chemicals Co. Ltd.), and blotting from the gel to PVDF membrane (Immobilon-P, 0.45 µm-ϕ, Cytiva Inc.)

was done by a constant current mode at 150 mA/10 cm^2 for 30 min. Blocking of the membrane was performed

with 0.3% skim milk and 0.02% NaN$_3$ in tris-buffered saline (TBS) containing 0.05% Tween-20 (TBS-T),

for the WB with anti-Bcs proteins, while blocking was performed with 2.5% BSA, 0.1% gelatin, 0.02%

Tween-20, 0.04% NaN$_3$ in PBS at 4°C overnight or at room temperature for 0.5–1 h, for either of anti-Strep-

tag II, anti-His5, or anti-thioredoxin. The membranes were briefly rinsed with TBS-T and then subjected to

the primary antibody reaction in blocking buffer for 1–2 h at room temperature. After removing the antibody

solution, the membrane was briefly rinsed and washed with TBS-T thrice for 5 min, followed by a secondary

antibody reaction in TBS-T for 30–45 min at room temperature. The membrane was briefly rinsed and

washed with TBS-T three times for 15 min each time. The protein bands were then visualized by casting the

ECL prime solution on the blotted PVDF membrane (0.2 mL/9 cm^2 membrane), and digitally recorded with

AE-9300H EZ-Capture MG (ATTO Co. Ltd., Japan).

Quantitative analysis was performed using ImageJ (version 1.53a) to quantify the band signal intensity,

calculated as relative values to the negative control, and the relative signal intensities of BcsA and BcsB were

normalized to the signal intensity of BcsD. For BcsA; two major bands were reproducibly observed near the

75-kDa and 250-kDa marker (Fig. 2B, top row), as well as in our previous studies (18,37), which are probably

BcsA monomer and BcsA-including oligomer, respectively (the estimated molecular weight for the BcsA

monomer is 84 kDa). Given these observations, the band intensity for the BcsA-containing oligomer was

also considered in addition to the monomer band intensity in subsequent analyses.

Data availability

All data are contained within this article and available from the corresponding author on reasonable

request.

Supporting information

This article contains supporting information.

Acknowledgments

The authors thank Dr. Kenji Tajima of Hokkaido University for his advice. The authors thank Professor

Ryoichi Yamaji and Dr. Yasuyuki Kobayashi at Osaka Prefecture University for their advice regarding the

pull-down assay.

Author contributions

T. I., and M. Y. conceptualization; T. K., T. I., and M. Y. methodology; T. K. formal analysis, T. K., N.

Y., and S. N. investigation; T. I. resources; T. K., and T. I. data curation; T. K., and T. I. writing-original

draft; T. K., M. Y., and T. I. writing-review and editing; T. K., and T. I. visualization; T. I., and M. Y.

supervision; T. K., M. Y., and T. I. project administration; T. I. funding acquisition.

Funding and additional information

This work was supported by JPSP KAKENHI (grant numbers 15H04530 and 19H00950), Exploratory

Research on Humanosphere Science (RISH, Kyoto University), and JST CREST (grant number:

JPMJCR13B2).

Conflict of interest

The authors declare that they have no conflicts of interest with the contents of this article.

References

1.   Morgan, J. L. W., Strumillo, J., and Zimmer, J. (2013) Crystallographic snapshot of cellulose

553   synthesis and membrane translocation. *Nature* **493**, 181-186

554   2.   Morgan, J. L. W., McNamara, J. T., and Zimmer, J. (2014) Mechanism of activation of

555   bacterial cellulose synthase by cyclic di-GMP. *Nat. Struct. Mol. Biol.* **21**, 489-496

556   3.   Römling, U., and Galperin, M. Y. (2015) Bacterial cellulose biosynthesis: Diversity of

557   operons, subunits, products, and functions. *Trends Microbiol.* **23**, 545-557

558   4.   Thongsomboon, W., Serra, D. O., Possling, A., Hadjineophytou, C., Hengge, R., and Cegelski,

559   L. (2018) Phosphoethanolamine cellulose: A naturally produced chemically modified

560   cellulose. *Science* **359**, 334-338

561   5.   Tajima, K., Imai, T., Yui, T., Yao, M., and Saxena, I. (2022) Cellulose-synthesizing

562   machinery in bacteria. *Cellulose* **29**, 2755-2777

563   6.   Wong, H. C., Fear, A. L., Calhoon, R. D., Eichinger, G. H., Mayer, R., Amikam, D., Benziman,

564   M., Gelfand, D. H., Meade, J. H., Emerick, A. W., Bruner, R., Ben-Bassat, A., and Tal, R.

565   (1990) Genetic organization of the cellulose synthase operon in *Acetobacter xylinum*. *Proc.*

566   *Natl. Acad. Sci. U.S.A.* **87**, 8130-8134

567   7.   Saxena, I. M., Kudlicka, K., Okuda, K., and Brown Jr, R. M. (1994) Characterization of

568   genes in the cellulose-synthesizing operon (acs operon) of *Acetobacter xylinum*:

569   Implications for cellulose crystallization. *J. Bacteriol.* **176**, 5735-5752

8.   Standal, R., Iversen, T. G., Coucheron, D. H., Fjaervik, E., Blatny, J. M., and Valla, S. (1994)

A new gene required for cellulose production and a gene encoding cellulolytic activity in

Acetobacter xylinum are colocalized with the bcs operon. *J. Bacteriol.* **176**, 665-672

9.   Kawano, S., Tajima, K., Kono, H., Erata, T., Munekata, M., and Takai, M. (2002) Effects of

endogenous endo-β-1,4-glucanase on cellulose biosynthesis in *Acetobacter xylinum*

ATCC23769. *J. Biosci. Bioeng.* **94**, 275-281

10. Tajima, K., Nakajima, K., Yamashita, H., Shiba, T., Munekata, M., and Takai, M. (2001)

Cloning and sequencing of the beta-glucosidase gene from *Acetobacter xylinum* ATCC

23769. *DNA Res.* **8**, 263-269

11. Tonouchi, N., Tahara, N., Kojima, Y., Nakai, T., Sakai, F., Hayashi, T., Tsuchida, T., and

Yoshinaga, F. (1997) A beta-glucosidase gene downstream of the cellulose synthase operon

in cellulose-producing acetobacter. *Biosci. Biotechnol. Biochem.* **61**, 1789-1790

12. McNamara, J. T., Morgan, J. L. W., and Zimmer, J. (2015) A Molecular Description of

Cellulose Biosynthesis. *Annu. Rev. Biochem.* **84**, 895-921

13. Kumagai, A., Mizuno, M., Kato, N., Nozaki, K., Togawa, E., Yamanaka, S., Okuda, K.,

Saxena, I. M., and Amano, Y. (2011) Ultrafine Cellulose Fibers Produced by *Asaia*

bogorensis, an Acetic Acid Bacterium. *Biomacromolecules* **12**, 2815-2821

14. Penttilä, P. A., Imai, T., Capron, M., Mizuno, M., Amano, Y., Schweins, R., and Sugiyama,

J. (2018) Multimethod approach to understand the assembly of cellulose fibrils in the

589   biosynthesis of bacterial cellulose. *Cellulose*

590   15.   Hu, S. Q., Gao, Y. G., Tajima, K., Sunagawa, N., Zhou, Y., Kawano, S., Fujiwara, T., Yoda,

591   T., Shimura, D., Satoh, Y., Munekata, M., Tanaka, I., and Yao, M. (2010) Structure of

592   bacterial cellulose synthase subunit D octamer with four inner passageways. *Proc. Natl.*

593   *Acad. Sci. U.S.A.* **107**, 17957-17961

594   16.   Sunagawa, N., Fujiwara, T., Yoda, T., Kawano, S., Satoh, Y., Yao, M., Tajima, K., and Dairi,

595   T. (2013) Cellulose complementing factor (Ccp) is a new member of the cellulose synthase

596   complex (terminal complex) in Acetobacter xylinum. *J. Biosci. Bioeng.* **115**, 607-612

597   17.   Omadjela, O., Narahari, A., Strumillo, J., Melida, H., Mazur, O., Bulone, V., and Zimmer, J.

598   (2013) BcsA and BcsB form the catalytically active core of bacterial cellulose synthase

599   sufficient for in vitro cellulose synthesis. *Proc. Natl. Acad. Sci. U.S.A.* **110**, 17856-17861

600   18.   Imai, T., Sun, S.-j., Horikawa, Y., Wada, M., and Sugiyama, J. (2014) Functional

601   reconstitution of cellulose synthase in *Escherichia coli. Biomacromolecules* **15**, 4206–4213

602   19.   Du, J., Vepachedu, V., Cho, S. H., Kumar, M., and Nixon, B. T. (2016) Structure of the

603   cellulose synthase complex of *Gluconacetobacter hansenii* at 23.4 A resolution. *PLoS ONE*

604   **11**, e0155886

605   20.   Sun, S.-j., Imai, T., Sugiyama, J., and Kimura, S. (2017) CesA protein is included in the

606   terminal complex of Acetobacter. *Cellulose* **24**, 2017–2027

607   21.   Sun, S.-j., Horikawa, Y., Wada, M., Sugiyama, J., and Imai, T. (2016) Site-directed

608   mutagenesis of bacterial cellulose synthase highlights sulfur–arene interaction as key to

609   catalysis. *Carbohydr. Res.* **434**, 99–106

610   22.   Acheson, J. F., Ho, R. Y., Goularte, N. F., Cegelski, L., and Zimmer, J. (2021) Molecular

611   organization of the *E. coli* cellulose synthase macrocomplex. *Nat. Struct. Mol. Biol.* **28**, 310-

612   318

613   23.   Santos, C. R., Paiva, J. H., Sforça, M. L., Neves, J. L., Navarro, R. Z., Cota, J., Akao, P. K.,

614   Hoffmam, Z. B., Meza, A. N., Smetana, J. H., Nogueira, M. L., Polikarpov, I., Xavier-Neto,

615   J., Squina, F. M., Ward, R. J., Ruller, R., Zeri, A. C., and Murakami, M. T. (2012) Dissecting

616   structure-function-stability relationships of a thermostable GH5-CBM3 cellulase from

617   *Bacillus subtilis* 168. *Biochem. J.* **441**, 95-104

618   24.   Acheson, J. F., Derewenda, Z. S., and Zimmer, J. (2019) Architecture of the Cellulose

619   Synthase Outer Membrane Channel and Its Association with the Periplasmic TPR Domain.

620   *Structure* **27**, 1855-1861.e3

621   25.   Nojima, S., Fujishima, A., Kato, K., Ohuchi, K., Shimizu, N., Yonezawa, K., Tajima, K., and

622   Yao, M. (2017) Crystal structure of the flexible tandem repeat domain of bacterial cellulose

623   synthesis subunit. *Sci. Rep.* **7**, 13018

624   26.   Perez-Riba, A., and Itzhaki, L. S. (2019) The tetratricopeptide-repeat motif is a versatile

625   platform that enables diverse modes of molecular recognition. *Curr. Opin. Struct. Biol.* **54**,

626   43-49

27.   Abidi, W., Zouhir, S., Caleechurn, M., Roche, S., and Krasteva, P. V. (2021) Architecture and

regulation of an enterobacterial cellulose secretion system. *Sci. Adv.* **7**, eabd8049

28.   Uto, T., Ikeda, Y., Sunagawa, N., Tajima, K., Yao, M., and Yui, T. (2021) Molecular Dynamics

Simulation of Cellulose Synthase Subunit D Octamer with Cellulose Chains from Acetic

Acid Bacteria: Insight into Dynamic Behaviors and Thermodynamics on Substrate

Recognition. *J.Chem.Theory Comput.* **17**, 488-496

29.   Toyosaki, H., Naritomi, T., Seto, A., Matsuoka, M., Tsuchida, T., and Yoshinaga, F. (1995)

Screening of Bacterial Cellulose-producing Acetobacter Strains Suitable for Agitated

Culture. *Biosci. Biotechnol. Biochem.* **59**, 1498-1502

30.   Nakai, T., Moriya, A., Tonouchi, N., Tsuchida, T., Yoshinaga, F., Horinouchi, S., Sone, Y.,

Mori, H., Sakai, F., and Hayashi, T. (1998) Control of expression by the cellulose synthase

(bcsA) promoter region from *Acetobacter xylinum* BPR 2001. *Gene* **213**, 93-100

31.   Rao, F., Pasunooti, S., Ng, Y., Zhuo, W., Lim, L., Liu, A. W., and Liang, Z. X. (2009)

Enzymatic synthesis of c-di-GMP using a thermophilic diguanylate cyclase. *Anal. Biochem.*

389, 138-142

32.   Crameri, A., Whitehorn, E. A., Tate, E., and Stemmer, W. P. C. (1996) Improved green

fluorescent protein by molecular evolution using DNA shuffling. *Nat. Biotechnol.* **14**, 315-

319

33.   https://www.gene.mie-u.ac.jp/Protocol/Original/Transform-Comp-Cell-Freeze.html,   the

latest access date: April 7 2022

34.   Updegraff, D. M. (1969) Semimicro determination of cellulose inbiological materials. *Anal.*

Biochem. **32**, 420-424

35.   Stahlberg, J., Johansson, G., and Pettersson, G. (1993) *Trichoderma reesei* has no true exo-

cellulase: All intact and truncated cellulase produce new reducing end groups on cellulose.

Biochim. Biophys. Acta - Gen. Subj. **1157**, 107-113

36.   Studier, F. W. (2005) Protein production by auto-induction in high density shaking cultures.

Protein Expr. Purif. **41**, 207-234

37.   Hashimoto, A., Shimono, K., Horikawa, Y., Ichikawa, T., Wada, M., Imai, T., and Sugiyama,

J. (2011) Extraction of cellulose-synthesizing activity of *Gluconacetobacter xylinus* by

alkylmaltoside. *Carbohydr. Res.* **346**, 2760–2768

Abbreviations

The abbreviations used are: Bcs, bacterial cellulose synthase; CD, catalytic domain; c.v., column

volume; c-di-GMP, cyclic-di-guanosine monophosphate; DDM, *n*-β-D-maltopyranoside; DGC,

diguanylate cyclase; DP, degree of polymerization;. FL, full length; GH, glycoside hydrolase; IMAC,

immobilized metal affinity chromatography; PASC, phosphoric acid swollen cellulose; TBS, tris-

buffered saline; TBS-T, tris-buffered saline with Tween-20; TPR, tetratricopeptide repeat

 Figure legends

Figure 1. Deletion of the *E. coli* endogenous *bcsA* gene

A Schematic diagram of the experiment for deleting the endogenous *bcsA* gene using the Red/ET

recombination system. The primer sequences used are listed in Table S1. **B** Electrophoresis pattern of colony-

direct PCR using two pairs of primers. M: Molecular weight marker (Marker 6 (Nippon Gene Co. Ltd., Tokyo,

Japan)); Δ*bcsA*: *bcsA*-deleted BL21; WT: wild-type BL21. **C** Quantification of endogenous cellulose

synthesis in BL21-GM and BL21 cells. The CESEC system (18), which uses DGC expression to synthesize

c-di-GMP to activate bacterial cellulose synthase, was used to count the amount of cellulose that used DGC

expression (bottom). The expression of DGC was confirmed by WB using an anti-Trx antibody (top). The

experiments were performed in triplicate ($n = 3$) and are represented as mean ± standard error of the mean

(SEM), together with each of the data values. Statistical tests (Student's t-test) were performed using the

built-in function in Microsoft Excel (TTEST).

Figure 2. *in vivo* cellulose-synthesizing activity of recombinant Bcs-complex

A Cellulose-synthesizing activity of the recombinant Bcs-complexes tested in this study: the amount of

cellulose synthesized by the CESEC system (18) with BL21-GM ($\Delta bcsA$) is shown for each recombinant

protein. All experiments were performed in triplicate ($n = 3$) and are represented as mean $\pm$ SEM together

with the significance of difference: * for $p < 0.05$, ** for $p < 0.01$, *** for $p < 0.001$, and *n.s.* for not significant

($p > 0.1$) as a result of Student's t-test, which was performed using the built-in function of Microsoft Excel

(TTEST). **B** Western blotting (WB) was performed for the CESEC experiment to check the expression of

BcsA, BcsB, BcsC, BcsD, and $_{TRX}$DGC. Cultures with and without DGC expression were tested. For BcsA,

the data are shown separately for the oligomer and monomer bands. Due to the limited number of wells in

the SDS-PAGE gel, the gel was divided into two parts for electrophoresis.

Figure 3. Purification of BcsABCD$_{H6}$

A Western blotting (WB) analysis of the purification process: M, membrane fraction; S, solubilized fraction;

F, IMAC unbound fraction; E, IMAC-eluted fraction. Three protein samples, wild-type (BcsABCD$_{H6}$),

D333N on BcsA (BcsA^{D333N}BCD$_{H6}$), and W373A on BcsA (BcsA^{W373A}BCD$_{H6}$), were analyzed using the

antibodies indicated. **B** WB analyses of the eluate from three independent purification experiments for each

of the three protein samples.

Approximately 2 µg of the protein was loaded onto the gel for WB with BcsA, BcsB, and BcsC antibodies,

and 1 µg of the protein was added to the gel for WB with BcsD antibody. For BcsA, the data are shown

separately for the oligomer and monomer bands.

Figure 4. Effect of cellulase treatment to the purification of Bcs-complex

A, Western blot images of IMAC-eluted fractions of BcsABCD$_{H6}$ after treatment with wild-type cellulase

(BsCel5A_FL and BsCel5A_CD). Coomassie Brilliant Blue (CBB)-stained images were used as loading

controls. **B** Band intensity in western blot analysis of **A**. Error bars indicate SEM, and the significance of the

difference as a result of Student's *t*-test, which was performed using the built-in function of Microsoft Excel

(TTEST), is indicated by the symbols: ** for $p < 0.01$, *** for $p < 0.001$, and *n.s.* for not significant ($p >$

0.1). **C**, Western blot images of IMAC-eluted fractions of BcsABCD$_{H6}$ after cellulase mutant (BsCel5A_FLm

and BsCel5A_CDm) treatment. CBB-stained images were visualized using BcsD as a loading control. **D**,

Band intensity in western blot analysis of **C**. Error bars indicate SEM, and Student's *t*-test, which was

performed using the built-in function of Microsoft Excel (TTEST), showed no significant difference ($p >$

0.1; eight independent replicates).

For BcsA, the data are shown separately for the oligomer band, monomer band, and their sum. Each elution

sample (~0.2 mg/mL) was added to an SDS-PAGE gel in 10 μL (anti-BcsA and anti-BcsB) and 1 μL (anti-

BcsD and CBB stain), respectively. Elution samples for CBB staining were loaded into gel wells 10-fold

more than those for WB (BcsD lane).

Figure 5. Effect of cellulase treatment on the purification of mutant Bcs-complex

A Western blot images of IMAC-eluted fractions of $BcsABCD_{H6}$, $BcsA^{D333N}BCD_{H6}$, and $BcsA^{W373A}BCD_{H6}$.

CBB-stained images were visualized using BcsD as a loading control. **B** Band intensity in western blot

analysis of **A**. An error bar indicates SEM, and the significance of the difference as a result of Student's t-

test, which was performed using the built-in function of Microsoft Excel (TTEST), is indicated by the

symbols † for $p < 0.1$, ** for $p < 0.01$, *** for $p < 0.001$, and *n.s.* for not significant ($p > 0.1$). **C**, Western

blot images of IMAC-eluted fractions of $BcsA^{D333N}BCD_{H6}$ and $BcsA^{W373A}BCD_{H6}$ after treatment with wild-

type cellulase (BsCel5A_FL and BsCel5A_CD). **D** and **E** Band intensity on western blot analysis of **C**: D for

$BcsA^{D333N}BCD$ and E for $BcsA^{W373A}BCD$. An error bar indicates SEM, and the significance of the difference

as a result of Student's t-test, which was performed using the built-in function of Microsoft Excel (TTEST),

showed no significant difference between any dataset.

All experiments were performed with four independent replicates ($n = 4$). For BcsA, the data are shown

separately for the oligomer band, monomer band, and their sum. Each elution sample (~0.2 mg/mL) was

added to an SDS-PAGE gel in 10 µL (anti-BcsA and anti-BcsB) and 1 µL (anti-BcsD and CBB stain),

respectively. Elution samples for CBB staining were loaded into gel wells 10-fold more than those for WB

(BcsD lane).

Figure 6. Pull-down assay with the Bcs subunits

A Schematic diagram showing the partners (bait and prey proteins) in a pull-down assay for each subunit of

BcsABCD. Alphabets on the cartoon indicate panels B–D (the result of pull-down experiments), by which

the subunit-subunit interaction was surveyed. **B** Western blot images of the pull-down assay using BcsD$_{StrepII}$

(bait) with crude BcsAB (prey). The pull-down fraction was analyzed by western blot using anti-BcsA, anti-

BcsB, and anti-Streptag antibodies. **C** Western blot images of the pull-down assay using BcsD$_{StrepII}$ (bait)

with purified $_{H6}$BcsC_TPR$_{H6}$ (prey). The pull-down fraction was analyzed by CBB staining and western

blotting using anti-His5 and anti-Streptag antibodies. **D** Western blot images of the pull-down assay using

$_{H6}$BcsC_TPR$_{H6}$ (bait) with crude BcsAB (prey). The pull-down fraction was analyzed by western blot using

anti-BcsA, anti-BcsB, and anti-His5 antibodies. **E** Western blot images of the pull-down assay using

BcsD$_{StrepII}$ (bait) with crude BcsAB (prey_1) and purified $_{H6}$BcsC_TPR$_{H6}$ (prey_2). The pull-down fraction

was analyzed by western blot using anti-BcsA, anti-BcsB, anti-His5, and anti-Streptag antibodies. Input:

sample immediately after mixing; pull-down: elution fraction after pull-down. All the detailed methods are

described in the "Experimental Procedures" section of the text.

Figure 7. The hypothetical two-state model for the type I Bcs-complex

Hypothetical two-state model for the Bcs-complex: basal state (left) and active state (right). In this model,

BcsD plays a key role in the periplasmic space to control the state of the complex separately from the state

of the BcsA enzyme (1,2). In the active state (right), BcsD draws the glucan chains in its octameric ring and

brings those chains to the channel pore of BcsC in the outer membrane like a "button whirligig'. The TPR

domain in BcsC may escort the BcsD protein to correctly transfer the glucan chains in BcsD to the BcsC

channel. In the basal state (left), BcsD docks with the BcsB complex given that BcsB is located in the

periplasmic space (1) as well as BcsD (12,20). In the model, BcsD in the basal state also docks with BcsC

without interfering with BcsB and is ready to draw the glucan chains in it once BcsAB starts synthesizing

and extruding out the glucan chain. Thus, BcsD in this state may be a standby state for effectively transferring

glucan chains into the extracellular space through the BcsC channel once BcsA is activated. Subunits that

were not assessed in this study are indicated in gray.

 Table 1. Experimental resources to express the proteins in this study

Material	Source	Description	Reference
Bacterial cellulose synthase			
pQE-BcsAB (GOI* was inserted between EcoRI and KpnI site in pQE-80L vector (Qiagen Inc.))	LC428282 (*Komagataeibacter sucrofermentans* JCM9730)	Expression vector for BcsAB protein without tags	Imai et al. 2014
pQE-BcsA^{D333N}B (GOI* was inserted between EcoRI and KpnI site in pQE-80L vector (Qiagen Inc.))	LC428282 (*Komagataeibacter sucrofermentans* JCM9730)	Expression vector for BcsAB protein with a catalytic-deficient point mutation of D333N, without tags	Sun et al., 2016
pQE-BcsA^{W373A}B (GOI* was inserted between EcoRI and KpnI site in pQE-80L vector (Qiagen Inc.))	LC428282 (*Komagataeibacter sucrofermentans* JCM9730)	Expression vector for BcsAB protein with a catalytic-deficient point mutation of W373A, without tags	Sun et al., 2016
pQE-BcsABCD (GOI was inserted between EcoRI and HindIII site in pQE-80L vector (Qiagen Inc.))	LC687564 (*Komagataeibacter sucrofermentans* JCM9730)	Expression vector for BcsABCD complex without tags	This study
pQE-BcsABCD$_{H6}$ (GOI was inserted between EcoRI and HindIII site in pQE-80L vector (Qiagen Inc.))	LC687564 (*Komagataeibacter sucrofermentans* JCM9730)	Expression vector for BcsABCD complex with hexahistidine-tag at the C-terminal of BcsD protein	This study
pQE-BcsABC$_{H6}$D (GOI was inserted between EcoRI and HindIII site in pQE-80L vector (Qiagen Inc.))	LC687564 (*Komagataeibacter sucrofermentans* JCM9730)	Expression vector for BcsABCD complex with hexahistidine-tag at the N-terminal of BcsD protein	This study
pQE-BcsABCD$_{GFPuvH6}$ (GOI was inserted between EcoRI and HindIII site in	LC687564 (*Komagataeibacter sucrofermentans*	Expression vector for BcsABCD complex with GFPuv followed by hexahistidine-tag at the C-terminal	This study

pQE-80L vector (Qiagen Inc.))	JCM9730)	of BcsD protein	
pQE-BcsA^{D333N}BCD$_{H6}$ (GOI was inserted between EcoRI and HindIII site in pQE-80L vector (Qiagen Inc.))	LC687564 (*Komagataeibacter sucrofermentans* JCM9730)	Expression vector for BcsABCD complex with a catalytic-deficient point mutation of D333N introduced in BcsA; hexahistidine-tag is fused to the C-terminal of BcsD protein	This study
pQE-BcsA^{W373A}BCD$_{H6}$ (GOI was inserted between EcoRI and HindIII site in pQE-80L vector (Qiagen Inc.))	LC687564 (*Komagataeibacter sucrofermentans* JCM9730)	Expression vector for BcsABCD complex with a catalytic-deficient point mutation of W373A introduced in BcsA; hexahistidine-tag is fused to the C-terminal of BcsD protein	This study
pET-$_{H6}$BcsC_TPR$_{H6}$ (GOI was inserted between NdeI and XhoI site in pET28b (Novagene Inc.))	LC687564 (*Komagataeibacter sucrofermentans* JCM9730)	Expression vector for TPR domain of BcsC protein (Cys18 – Gly848): hexahistidine-tag is fused to both N- and C-terminal of the protein	This study
pET-BcsD$_{StrepII}$ (GOI was inserted between NcoI and HindIII site in pET28b vector (Novagen Inc.))	LC687564 (*Komagataeibacter sucrofermentans* JCM9730)	Expression vector for BcsD protein: Streptag(II) is fused to the C-terminal of BcsD protein	This study

		Diguanylate cyclase (DGC)	
pBAD33et32-tDGC (Self-made vector in the previous study)	NP_229585 (*Thermotoga maritima* MSB8)	Expression vector of diguanylate cyclase with N-terminal truncation (K82 – S248) and a point mutation (R158A mutation): thioredoxin was fused to Nterminal end of the protein	This study

		Cellulase	
pGEX-BsCel5A_FL (GOI was inserted between BamHI and XhoI site in	Z29076 (*Bacillus subtilis* strain 168)	Expression vector for full-length endoglucanase protein from *Bacillus subtilis* (BsCel5A):	This study

pGEX4T-3 (GE Healthcare Inc.))		signal-peptide sequence (from Met1 to Ala28) was removed; GST-tag is fused to the N-terminal of BsCel5A protein, which consists of glycosylhydrolase 5 domain (GH-5 domain, Ala29 – Thr332) and carbohydrate-binding module (CBM-3, Ile354 – Asn499)	
pGEX-BsCel5A_CD (GOI was inserted between BamHI and XhoI site in pGEX4T-3 (GE Healthcare Inc.))	Z29076 (*Bacillus subtilis* strain 168)	Expression vector for catalytic domain (A29 – T332) of BsCel5A: GST-tag is fused to the N-terminal of the protein, which lacks the signal peptide sequence (Met1 – Ala28)	This study
pGEX-BsCel5A_FLᵐ (GOI was inserted between BamHI and XhoI site in pGEX4T-3 (GE Healthcare Inc.))	Z29076 (*Bacillus subtilis* strain 168)	Expression vector for full-length BsCel5A with activity-deficient double mutation (E169Q/E257Q): GST-tag is fused to the N-terminal of the protein, which lacks the signal peptide sequence (Met1 – Ala28)	This study
pGEX-BsCel5A_CDᵐ (GOI was inserted between BamHI and XhoI site in pGEX4T-3 (GE Healthcare Inc.))	Z29076 (*Bacillus subtilis* strain 168)	Expression vector for catalytic domain (A29 – T332) of BsCel5A with activity-deficient double mutation (E169Q/E257Q): GST-tag is fused to the N-terminal of the protein, which lacks the signal peptide sequence (Met1 – Ala28)	This study

768 * GOI: gene of interest

Figure 1

A

WT (BL21)

2,818 bp

#1 *bcsA* *bcsB*

#3

FRT-PGK-gb2-neo-FRT

Franking Region 1   Franking region 2

Frankign region 1: 5'- ACAGCTGGCTCAGCATTGATCAGTGGTTGCGTTGCTGGCGTCGCCTGCGT-3'
Franking region 2: 5'- CGGCTGAAGAGATACTGACGCTGGCGAACTGGTGCCTGTTGAACTATTCC-3'

ΔbcsA (BL21-GM1)

#1 #2

FRT-PGK-gb2-neo-FRT   *bcsB*

#3

1,505 bp

1,731 bp

Primers
#1: 5'-AAGAGATACTGACGCTGGCGAACTGG-3'
#2: 5'-TCAGAAGAACTCGTCAAGAAGGC-3'
#3: 5'-ACAGCTGGCTCAGCATTGATCAGTGG-3'

B

primer #2/#3   primer #1/#3

(bp) M ΔbcsA WT ΔbcsA WT M

4,260 — | | — 4,260
3,470 — | | — 3,470
2,690 — | | — 2,690

1,880 — | | — 1,880
1,490 — | | — 1,490

930 — | | — 930

C

WB: αTrx ($_{TRX}$DGC)

arabinose: − + − +

(m/kDa) 50 — 37 —

Amount of synthesized cellulose (μg/ml of culture)

$p < 0.01$

$_{TRX}$DGC: − + − +

BL21-GM   BL21

Figure 2

Figure 3

A

B

Figure 4

Figure 5

Figure 6

Figure 7