1

Dynamic interaction of bacterial cellulose synthase subunit D (BcsD) in type I bacterial cellulose synthase

Tatsuya Kondo[1], Yui Nakamura[2], Shingo Nojima[2], Min Yao[2*], Tomoya Imai[1*]

1 Research Institute for Sustainable Humanosphere (RISH), Kyoto University, Uji, Kyoto 611-0011, Japan.

2 Faculty of Advanced Life Science, Hokkaido University, N10W Kita-ku, Sapporo 060-0810, Japan.

*Corresponding authors:

Tomoya Imai, imai.tomoya.4c@kyoto-u.ac.jp; Research Institute for Sustainable Humanosphere, Kyoto University, Uji, Kyoto 611-0011, Japan.

Min Yao, yao@castor.sci.hokudai.ac.jp; Faculty of Advanced Life Science, Hokkaido University, N10W Kita-ku, Sapporo 060-0810, Japan.

Email: imai.tomoya.4c@kyoto-u.ac.jp

Author Contributions: T. I. and M. Y. conceptualization; T. K., T. I., and M. Y. methodology; T. K. formal analysis, T. K., N. Y., and S. N. investigation; T. I. resources; T. K. and T. I. data curation; T. K. and T. I. writing-original draft; T. K., M. Y., and T. I. writing-review and editing; T. K. and T. I. visualization; T. I. and M. Y. supervision; T. K., M. Y., and T. I. project administration; T. I. funding acquisition.

Competing Interest Statement: The authors declare that they have no conflicts of interest regarding the content of this article.

Classification: Biological Sciences

Keywords: cellulose, cellulose synthase, membrane enzyme, protein complex, polymer solid formation

This PDF file includes:

      Main Text
      Figures 1 to 7
      Tables 1

Abstract

Cellulose is a promising biological material for supporting sustainable human life. This natural polymer is synthesized by cellulose synthase, a protein complex in the cell membrane. Cellulose synthase in bacteria is a hetero-subunit complex, and its subunit organization varies widely

depending on the species. In the type I bacterial cellulose synthase complex, the BcsD (bacterial
cellulose synthase D) protein is believed to play an important role in producing cellulose with long
slender fiber morphology and high crystallinity, given the phenotype of the bcsD-deficient mutant
and the specific existence of the type I operon in bacterial species synthesizing crystalline
cellulose microfibrils such as Acetobacter. In this study, we successfully established a
heterogeneously expressed Bcs protein in Escherichia coli as an experimental model and
conducted biochemical studies for the BcsD protein and the other three major subunits of
bacterial cellulose synthase, BcsA, BcsB, and BcsC. It has been shown that the BcsD protein
interacts with the functionally required minimal subunits of the BcsAB complex, as well as the
BcsC protein. Furthermore, it was shown that BcsD interacts with the BcsAB complex in two
modes: direct protein-protein interactions and indirect interactions through the product cellulose.
The former and latter modes represent the basal and active states of the type I bacterial cellulose
synthase, respectively. This dynamic behavior of the BcsD protein is important for the type I
bacterial cellulose synthase complex to regulate the crystallization process of cellulose.

Significance Statement

Cellulose synthase is a membrane enzyme and its function is not only polymerization of glucose
residues, but also bundling the polymerized glucan chains to form a slender crystalline fiber. This
is a fascinating mechanism considering that long water-insoluble cellulose chains are actively
assembled in aqueous solvent at room temperature. To clarify this mechanism as the protein
function, this study analyzed bacterial cellulose synthase using protein purification experiments.
57

58   **Main Text**
59
Introduction

Cellulose is a natural polymer, whose molecular structure is composed of a homopolymer of
glucose with a β1→4 linkage, which is very simple. However, cellulose molecules are insoluble
in water, and the apparent elementary structure of cellulose is an aggregation of many cellulose
molecules in long crystalline fibers. This structural feature of cellulose underlies its structure-
sustaining property, which many cellulose-producing organisms rely on.

This polymer assembly is produced by the enzyme "cellulose synthase." In many known cases,
in vivo cellulose synthase is a protein complex with several polypeptide subunits. The most
important subunit in this complex is glycosyltransferase, which is called CesA. The first
structural biological study of the CesA protein was reported in 2013 for cellulose synthase from
the bacterium *Rhodobacter sphaeroides* (1, 2), which has stimulated many biochemical studies on
cellulose synthase.

Given its simplicity, bacterial cellulose synthase (Bcs) has been a model in many studies on
cellulose synthase. Particularly, Bcs from bacteria belonging to the genus *Acetobacter*, some of
which are currently named as *Gluconacetobacter* or *Komagataeibacter* (the genus name
"*Acetobacter*" is used for these bacteria in this study), has been used extensively in studies.
Although it is now known that many other bacteria produce cellulose, the function and structure
of cellulose appears to vary with species (3). For example, phosphoethanolamine cellulose has
been reported in *Escherichia coli*, and this natural cellulose derivative is important for interacting
with the host organism (4).

Among cellulose-synthesizing bacteria, *Acetobacter*, which has a gene cluster or operon
categorized as type I (3), is characteristic of synthesizing highly crystalline cellulose microfibrils
(5) as well as plant cellulose, a major natural source of cellulose. Previous studies have reported
that the Bcs-complex in *Acetobacter* is composed of at least four subunits (6, 7) – BcsA (bacterial
CesA, catalytic subunit), BcsB (auxiliary subunit having a carbohydrate-binding domain and
flavodoxin domain), BcsC (secretion of a glucan chain through the outer membrane), and BcsD
(crystallization subunit). In addition, endoglucanase CMCax (8, 9) (glycoside hydrolase family 8
(GH8)), unknown function protein Ccp (8, 10) (also named as BcsH (3)), and β-glucosidase
bglAx (GH3) (11) are considered to be functionally involved in the complex. Given that the
cooperation of these subunits produces highly crystalline cellulose microfibrils, the spatial
organization of these subunits in the cell is important for understanding the cellulose biosynthesis
mechanism as a protein function (12).

In these subunits, the BcsD protein is found specifically in bacterial species producing highly
crystalline cellulose microfibrils such as *Acetobacter* and *Asaia* (3, 13, 14), which possess a type I
Bcs-operon (3). The *bcsD*-deficient mutant of *Acetobacter* showed a reduction in cellulose
production, as well as a lack of ability to synthesize crystalline cellulose microfibrils (6, 7, 15).
Sunagawa *et al.* also demonstrated that BcsD interacts with Ccp (BcsH) in a pull-down assay
(16). Given that the BcsD protein contains glucan chains inside its octamer structure (15), it has
been proposed that BcsD is functionally located downstream of the BcsAB complex (the
cellulose-polymerizing core of the Bcs-complex (1, 6, 17, 18)) (19, 20). Detailed information on
how these functional units work together is key to understanding the mechanism by which
cellulose synthase can bundle the synthesized glucan chains for spinning a long slender
crystalline fiber at room temperature in an aqueous environment.

In this study, we analyzed protein-protein interactions in the type I Bcs-complex using the
recombinant BcsABCD protein with the *E. coli* expression system and clarified the interaction of
BcsD with BcsAB complex as well as BcsC protein. One of the advantages of using recombinant
proteins is that mutant proteins are easily available (21). Owing to this advantage, this study
sheds light on the interaction between BcsAB and BcsD and demonstrates that there are two
modes for this interaction: direct protein-protein interactions and indirect interactions through the
cellulose molecules. We hypothesized that such bimodal interactions between BcsAB and BcsD
proteins could be key to controlling cellulose synthesis dynamically *in vivo*.

Results

Production of endogenous *bcsA* gene-deficient *E. coli*

The expression of the *Acetobacter* Bcs complex in *E. coli* cells might lead to contamination by
endogenous *E. coli* Bcs (22).We then attempted to delete *bcsA* in the *E. coli* host, which encodes
the catalytic subunit of Bcs. The deletion of the *bcsA* gene in *E. coli* BL21, a strain expressing the
Bcs recombinant protein in this study, was performed using the Red/ET recombination system
(Gene Bridges GmdH, Germany), in which the *bcsA* gene was replaced by the FRT-PGK-gb2-
neo-FRT cassette with homology arms of the *bcsA*-flanking region using homologous
recombination (Fig. 1A). This genetically modified BL21 (BL21-GM) strain was confirmed by
colony direct polymerase chain reaction (PCR) to contain the neomycin-resistance gene cassette
(FRT-PGK-gb2-neo-FRT cassette) (Fig. 1B). When amplified with primer #2 (antibiotic gene
cassette-specific) and #3 (*E. coli*-specific), a band of approximately 1,500 bp was found in BL21-
GM, while no specific product band was found in wild-type BL21, indicating the insertion of the
neomycin-resistance gene cassette specifically in BL21-GM. However, when amplified with
primers #1 and #3 (either *E. coli*-specific), wild-type BL21 and BL21-GM produced a band
corresponding to the *bcsA* gene (2,818 bp) and the inserted gene cassette (1,731 bp), respectively.
These results confirmed the successful deletion of the *bcsA* gene of BL21 in "BL21-GM."

Next, we measured the *in vivo* cellulose synthesis activity of the prepared BL21-GM (Fig. 1C)
by partially using our previous method (18): diguanylate cyclase (DGC) fused to thioredoxin
($_{Trx}$DGC) was expressed in BL21 and BL21-GM to synthesize c-di-GMP to activate endogeneous
bacterial cellulose synthase. The amounts of cellulose synthesized were approximately 2.3 and
0.7 μg/mL of culture in BL21 and BL21-GM, respectively. Although weak cellulose-synthesizing
activity was observed, we used BL21-GM as a suitable host to express the Bcs subunits with
minimal impact on the endogenous Bcs-complex of *E. coli*.

Characterization of BsCel5A

Cellulase is an important tool used in this study to scrutinize the role of cellulose molecules in
the subunit-subunit interaction of the Bcs-complex. In this study, we used a bacterial endo-β-1,4-
glucanase from *Bacillus subtilis* (BsCel5A), whose structure has been previously reported (23).
The full length (FL) of this cellulase is a two-domain protein, in which the catalytic domain of
GH5 is linked to the carbohydrate binding module of family 3 (CBM3). In this study, the catalytic
domain alone (CD) was also used to maximize access to the soluble substrate. FL and CD of
BsCel5A were expressed as glutathione-*S*-transferase (GST)-fused proteins and purified with
glutathione affinity resin, as confirmed by SDS-PAGE (Fig. S1A, left panel); (Fig. S1A, left

panel, white arrow). Activity-deficient mutant of E169Q/E257Q in FL (BsCel5A_FLm) and CD
(BsCel5A_CDm) was also used for a negative control protein in the cellulase treatment
experiment.

The binding activity of the purified BsCel5A to phosphoric acid-swollen cellulose (PASC) was
surveyed to confirm correct folding of the expressed proteins. The results showed that all the
BsCel5A proteins used in this study retained their binding activity to PASC (Fig. S1A, right
panel), indicating that all of them were properly folded despite sequence modification. The
enzymatic activity of BsCel5A was surveyed to determine whether this cellulase could function at
low temperature and neutral pH, which is required for analyzing the Bcs-complex. Both
BsCel5A_FL and BsCel5A_CD hydrolyzed barley β-glucan, carboxyl methyl cellulose (CMC),
and PASC at 4°C and pH 7.5 (Fig. S1B). For Avicel, full-length BsCel5A showed higher activity
than the others, stressing the function of CBM to bind solid cellulose.

Substrate specificity of BsCel5A was investigated using cello-oligosaccharides with various
degrees of polymerization (DP). BsCel5A hydrolyzed cellotetraose, cellopentaose, and
cellohexaose but not cellobiose or cellotriose (Fig. S1C). The final reaction products of BsCel5A
for substrates longer than triose were cellobiose and cellotriose, and the smallest substrate was
cellotetraose (Fig. S1D).

Summarizing these, both BsCel5A_FL and BsCel5A_CD were able to hydrolyze cellulose of
low crystallinity at 4°C and pH 7.5, and cellulose molecules with DP higher than 3 were the
substrates. This information was considered when interpreting the data.

In vivo cellulose synthase activity of recombinant Bcs-complex

This study analyzed heterogeneously expressed Bcs proteins in *E. coli* as an experimental
model. We then checked if these heterogeneously expressed proteins were functional by using
CESEC (cellulose-synthesizing *E. coli*), a platform for estimating the cellulose-synthesizing
activity with the co-expression of diguanylate cyclase (DGC) (18). All the Bcs-complexes
expressed in DGC-expressing BL21-GM synthesized a significant amount of cellulose (Fig. 2A),
indicating that each of the Bcs-complexes expressed in this study are functional, regardless of
whether BcsD is tagged. In addition, neither BcsAB nor the BcsABCD complex with the activity-
deficient mutant of the catalytic subunit BcsA (D333N or W373A) showed cellulose-synthesizing
activity, consistent with our previous study (21). These results support the idea that the Bcs
proteins expressed in this study reflect the state in function and are suitable for an experimental
model.

Despite the lower cellulose production shown in Fig. 2A, we decided to express Bcs proteins
without co-expression of DGC for subsequent purification because higher expression levels of
Bcs proteins were detected by western blotting (WB), as shown in Fig. 2B. Given that the
cellulose quantification in this study used strong acid treatment with acetic acid and nitric acid,
we think that highly crystalline cellulose was majorly considered in the data presented in Fig. 2A.
Then it cannot be ruled out that the Bcs-complex expressed in *E. coli* without DGC expression
synthesizes cellulose.

The complex formation of BcsAB, BcsC, and BcsD

Given that BcsD holds the glucan chains in its channel (15), it is functionally located
downstream of the BcsAB complex, the core of cellulose molecule synthesis, indicating that
BcsD operates in the periplasmic space to assemble cellulose molecules extruded from the
BcsAB complex. Furthermore, the BcsC protein has a β-barrel porin-like structure at its C-
terminus and is supposed to secrete cellulose through the outer membrane (24). The results from
several studies suggest that BcsAB, BscC, and BcsD work together in a coordinated manner to
spin bacterial cellulose. We tried to co-express BcsABCD to purify the whole Bcs-complex using
a hexa-histidine tag attached to the C-terminus of BcsD (BcsD$_{H6}$), as the tag at the C-terminus
does not disturb cellulose-synthesizing activity (16). After the optimization of the conditions (Fig.
S2), all four subunits were successfully purified together by immobilized metal affinity
chromatography (IMAC) (Fig. 3). This is the first successful purification of the BcsABCD
complex ever reported. However, BcsC was found to be mostly insoluble in detergent (*n*-dodecyl-
β-D-maltoside, DDM, in this study) and tended to degrade or aggregate during the purification
process (Fig. 3A, BcsC antibody strip). We did not decipher BcsC behavior in this Bcs-complex
purification experiment.

Analysis of the interaction between BcsAB and BcsD by cellulase treatment

As described above, co-expression of BcsABCD$_{H6}$ in *E. coli* allowed the BcsABD-complex to
be purified. Given that BcsABCD$_{H6}$ expressed in *E. coli* is functional as shown in Fig. 2A, we
speculated that a glucan chain may link BcsAB and BcsD during this purification. This
hypothesis was tested by checking whether the digestion of cellulose resulted in the dissociation
of BcsAB from BcsD$_{H6}$ during IMAC purification. BsCel5A cellulase was applied to
BcsABCD$_{H6}$ bound to IMAC resin prior to washing and elution. The amounts of BcsA, BcsB,
and BcsD proteins in the eluate were measured by semi-quantitative WB (Fig. 4A); the reliability
of quantification in our WB was confirmed, as shown in Fig. S3. The signal intensities of BcsA
and BcsB were normalized to that of BcsD in the eluate for comparison between samples.
Notably, the amounts of BcsA and BcsB bound to BcsD were decreased by BsCel5A treatment
(Fig. 4B), indicating that the BcsAB complex was detached from BcsD by cellulose digestion.
This trend was clearer when BsCel5A_CD was used as cellulase.

To rule out the possibility that the observed detachment of BcsAB was due to non-specific
interaction of the added BsCel5A protein, the activity-deficient mutant of BsCel5A
(E169Q/E257Q) was applied to BcsABCD$_{H6}$ bound to the metal affinity resin (Fig. 4, C and D).
As a result of purification, no decrease in the signal intensity of BcsA and BcsB was observed in
the eluate, indicating that cellulose hydrolysis actually caused detachment of BcsAB from BcsD
during purification. In summary, the BcsAB complex indirectly interacts with BcsD through the
glucan chain.

Although cellulase treatment allowed the BcsAB complex to be detached from BcsD during
purification, as demonstrated above, a substantial portion of the BcsA and BcsB proteins
remained with BcsD in this purification process (Fig. 4B). We supposed that a direct interaction
between BcsAB and BcsD is also found in addition to the indirect interaction. To test this
hypothesis, we attempted to purify the activity-deficient mutants of BcsABCD$_{H6}$,
BcsA^{D333N}BCD$_{H6}$, and BcsA^{W373A}BCD$_{H6}$, neither of which synthesize cellulose, as shown in Fig.
2A. As a result, these mutants allowed BcsA and BcsB to be purified together with BcsD and the
wild type (Fig. 5A and B). Notably, cellulase treatment did not reduce the amount of BcsAB
purified together with BcsD for either BcsA^{D333N}BCD$_{H6}$ or BcsA^{W373A}BCD$_{H6}$ (Fig. 5C, 5D, and

5E). These results support the existence of a direct interaction between BcsAB and BcsD when
cellulose is not synthesized. Compiling these cellulase treatment experiments, it was
demonstrated that there are two modes of interaction between BcsAB and BcsD: direct and
indirect modes through cellulose molecules.

Quaternary structure of the Bcs-complex

Despite the demonstration of two modes for the interaction between BcsAB and BcsD, the
quaternary structure of the entire Bcs-complex remains to be elucidated. We then performed pull-
down assays to further explore the subunit-subunit interactions in the BcsABCD complex by fully
utilizing recombinant Bcs proteins. The assay was designed based on the results from previous
studies (Fig. 6A): (i) BcsAB forms a complex in the inner membrane (1), (ii) BcsC is an outer
membrane protein (24, 25), (iii) BcsD protein is located in the periplasmic space (15, 20), and (iv)
BcsD interacts with the BcsAB complex (this study).

First, the interaction between BcsAB and BcsD, which has been demonstrated above, was
tested by pull-down with a bait of Streptag(II)-fused BcsD (BcsD$_{StrepII}$) and a prey of crude
BcsAB without tags (DDM-solubilized cell membrane of *E. coli* expressing BcsAB). As
expected, BcsAB bound to BcsD$_{StrepII}$ (Fig. 6B). The pull-down assay yielded a result consistent
with our purification experiment of the BcsABD-complex and supported our hypothesis for the
interaction between BcsAB and BcsD.

Given the successful pull-down assay with BcsAB and BcsD$_{StrepII}$, we next performed a pull-
down assay to survey the BcsC-involved interaction in the complex (see Fig. 6A). For this
purpose, TPR tetratricopeptide-repeat (TPR) domain of the BcsC protein (BcsC_TPR) was used
based on two assumptions: (i) the role of the TPR domain is protein-protein interaction (26), and
(ii) the full-length BcsC protein is unstable, as shown in our purification. Pull-down by BcsD$_{StrepII}$
trapped $_{H6}$BcsC_TPR$_{H6}$, which has hexa-histidine tags at both the N- and C-termini of
BcsC_TPR, in a dose-dependent manner (Fig. 6C), clearly showing that these two proteins
interact with each other. Furthermore, pull-down by the purified $_{H6}$BcsC_TPR$_{H6}$ (bait) with the
crude BcsAB (prey) also supported their interaction with each other (Fig. 6D), although it
appeared less striking than the case with $_{H6}$BcsC_TPR$_{H6}$ and BcsD$_{StrepII}$ (Fig. 6C).

Finally, we performed a pull-down assay using all subunits in the complex: crude BcsAB,
purified $_{H6}$BcsC_TPR$_{H6}$, and purified BcsD$_{StrepII}$ (Fig. 6E). Streptactin beads trapped BcsA, BcsB,
and $_{H6}$BcsC_TPR$_{H6}$ together with BcsD$_{StrepII}$, although BcsA was only faintly detected by WB. It
is then summarized that the BcsD protein can simultaneously interact with BcsA, BcsB, and the
TPR region of BcsC.

Discussion

Bacterial cellulose synthase is a heterosubunit complex. Primary evidence for this hypothesis is
the existence of gene clusters or operons encoding cellulose synthase-related genes (3, 6, 7, 11).
However, this hypothetical model has not been tested intensively, owing to the lack of studies on
the protein. Two recent reports have addressed this question. Sunagawa et al. demonstrated an
interaction between BcsD and BcsH (Ccp) proteins using a pull-down assay (16). Another
prominent study was the X-ray crystallographic analysis of the BcsAB protein complex by

Morgan et al., which clarified that the BcsA and BcsB proteins closely interact with each other to
polymerize and translocate the glucan chain from the cytosol to the extracellular side (1), as
confirmed also in *E.coli* (27). In this study, we analyzed the type I Bcs complex and demonstrated
that the BcsD protein interacted with the BcsAB complex (Fig. 3, 4, 5, and Fig. 6B) and the TPR
domain of BcsC (Fig. 6C). Furthermore, it was shown that BcsC_TPR interacted with the BcsAB
complex, although the result was not as clear as the interaction between BcsC_TPR and BcsD
(Fig. 6D). Finally, a pull-down assay using BcsD$_{StrepII}$ with a mixture of crude BcsAB and
purified $_{H6}$BcsC_TPR$_{H6}$ indicated that BcsD binds to both BcsAB and BcsC_TPR
simultaneously, suggesting that these four subunits interact with each other (Fig. 6E). Our study,
together with previous studies, experimentally demonstrated that the subunits of BcsA, BcsB,
BcsC, BcsD, and BcsH are included in the type I Bcs-complex.

Interestingly, this study showed that the interaction of BcsD with BcsAB has two modes: direct
and indirect interactions through the polymerized glucan chain (Fig. 4). Direct interaction
between the BcsAB complex and BcsD protein was observed for both wild type and catalytically
deficient mutant cellulose synthase, whereas the interaction via cellulose was observed only for
the wild type (Fig. 5). We then propose a two-state model for the type I Bcs complex: basal and
active states, which are represented by the direct interaction (Fig. 7, left) and indirect interaction
modes (Fig. 7, right), respectively.

BcsD protein in this basal state is statically fixed on the BcsAB complex for lying in wait for a
glucan chain synthesized by the BcsAB complex. Once cellulose synthesis is initiated, the
synthesized glucan chain is guided into the octameric ring of BcsD to crystallize cellulose
molecules, as proposed previously (7, 15, 16, 20, 28), which is a shift from the resting state to the
active state of the complex. In the active state (Fig. 7, right), the BcsD protein is detached from
the BcsAB complex but tethered to BcsAB by the glucan chain extruded from BcsAB.
Considering the substrate specificity of BsCel5A (Fig. S1), BcsD can be physically separated
from BcsB by a glucan chain with a length of four or more glucose residues (Fig. 7).

In this two-state model, the active state of the Bcs complex allows BcsD to be launched toward
the outer membrane, while BcsD in the basal state may be fixed statically on BcsB. The TPR
domain of the BcsC protein may control diffusible BcsD in the active state in the periplasmic
space by gentle but significant interactions. Given that the C-terminal domain of BcsC is an outer
membrane β-barrel channel (Fig. S4) (24), the TPR domain of BcsC may escort BcsD carrying
glucan chains toward the outer membrane channel in BcsC to spin the glucan chains to the
extracellular side. The following questions about this model will be whether (i) BcsD protein
could dock to the outer membrane channel part of the BcsC protein and (ii) the interaction
between BcsC and BcsD could be state-dependent. These questions will be addressed in future
studies to enhance our understanding of the Bcs-complex function.

Our study demonstrated the intermolecular interactions of subunits in the BcsABCD complex
to update our knowledge of the quaternary structure of the type I Bcs complex. Experiments in
this study were performed using recombinant proteins heterogeneously expressed in *E. coli*; thus,
further studies will be necessary to test the hypothesis proposed in this study, the two modes for
BcsAB-BcsD interaction: the basal and the active state. Nonetheless, the data obtained in this
study were intensively tested using mutagenesis experiments of proteins, and they were consistent
with each other. We believe that our hypothesis in this study is sufficiently reliable, at least for
starting further discussion about the mechanism of Bcs with regard to protein function. In this
model, BcsD is not statically fixed in the Bcs-complex but dynamically interacts with the other
subunits in the complex (Fig. 7). Such a dynamic behavior may be involved in the crystallization

function of cellulose molecules, as well as twisting in bacterial cellulose microfibrils, as reported
previously (16). The putative state control with BcsD, as shown in Fig. 7, is the regulation of the
quaternary structure of bacterial cellulose synthase. Next, we need to further study the original
Acetobacter cells and observe the whole Bcs-complex at higher resolution with cryo-EM under
controlled two-state conditions.

326
327
Materials and Methods
329
Chemicals
331 All chemicals used in this study were purchased from FUJIFILM Wako Pure Chemicals Co.
332 Ltd. (Osaka, Japan), Nacalai Tesque Co. Ltd. (Kyoto, Japan), and Sigma-Aldrich Inc. (MO,
333 USA), unless otherwise indicated.

Cloning

Bcs genes (*bcsA*, *bcsB*, *bcsC*, and *bcsD*) were prepared from genomic DNA extracted from
cultured cells of *Komagataeibacter sucrofermentans* JCM9730, which was provided by the
RIKEN Bio Resource Center (Tsukuba, Japan) and known by the strain name BPR2001 (29, 30).
The endo-β1,4-glucanase gene from *Bacillus subtilis* strain 168 (*bscel5A*) (23) was prepared from
the genomic DNA provided by the American Type Culture Collection (ATCC23857D). The dgc
gene (*dgc*) was identical to that used in our previous study (18): tDGC and DGC from
Thermotoga maritima MSB8 with a point mutation of R158A and truncation of the N-terminal 81
amino acid residues from V1 to V81 (31). The gene for green fluorescent protein UV (GFPuv)
(32) was prepared from the pGFPuv plasmid purchased from Takara Bio, Co. Ltd. (Shiga, Japan).

The cloning of these genes was performed by PCR using PrimeStarGXL (Takara Bio Co. Ltd.).
Site-directed mutagenesis was performed by QuikChange II (Agilent Inc., CA, USA) or the
equivalent PCR-based method. The sequence of the PCR-amplified DNA was checked by the
dye-terminator method in a subcloning vector (pGEM-T easy (Promega Inc., WI, USA) or pTAC-
2 (BioDynamics Laboratory Co. Ltd., Tokyo, Japan)), or final expression vector, which was
performed at either of the Uji DNA-sequencing core (Kyoto University Research Administration
Office) with an ABI Prism 310 Genetic Analyzer (Applied Biosystems Inc., MA, USA), or
Eurofins Genomics Inc. (Tokyo, Japan).

Construction of the expression vectors

The protein expression vectors used in this study are summarized in Table 1. Expression
vectors for BcsAB (BcsWTB, BcsA^{D333N}B, and BcsA^{W373A}B) were identical to those used in
previous studies (18, 21). The *bcsAB* gene in the operon form was inserted between the EcoRI
and KpnI sites in pQE-80L (Qiagen, Hilden, Germany), and the upstream of the start codon was
the original ribosomal binding site of *K. sucrofermentans* JCM9730.

Expression vectors for BcsABCD (BcsABCD, BcsABC$_{H6}$D, BcsABCD$_{H6}$, BcsA^{D333N}BCD$_{H6}$,
BcsA^{W373A}BCD$_{H6}$, and BcsABCD$_{GFPuvH6}$) proteins were constructed from the vectors for BcsAB
by inserting the *bcsABCD* gene in the operon form between the EcoRI and HindIII sites in pQE-

80L. The ribosomal binding site in these BcsABCD expression vectors is exactly the same as that
in the expression vectors of BcsAB, one of *K. sucrofermentans* JCM9730. DNA encoding the tag
(hexa-histidine or GFPuv) was added to *bcsD* during the PCR-amplification amplification.

The expression vectors for the TPR region of BcsC (BcsC_TPR: C18-G848) and BcsD were
prepared for the pull-down assay. DNA encoding hexa-histidine tag was added to the 3′-end of
the gene for BcsC_TPR during PCR amplification and inserted into the NdeI and XhoI sites in
pET-28b (Merck Corp.., Darmstadt, Germany) to express BcsC_TPR protein with dual hexa-
histidine tag at both N- and C- terminal – $_{H6}$BcsC_TPR$_{H6}$. DNA encoding Streptag (II) was added
to the 3′-end of the gene for BcsD and inserted between the NcoI and HindIII sites in pET28b to
express BcsD with C-terminal Streptag(II)–BcsD$_{StrepII}$.

The gene for tDGC was inserted between the NdeI and HindIII sites in pBAD33et32, a
customized plasmid DNA in this study, to express the thioredoxin-fused tDGC protein to count
the cellulose-synthesizing activity of the recombinant BcsAB and BcsABCD complex in our
previous studies (18, 21). The detailed procedure for preparing pBAD33et32 is described in
Supplementary Information. Thioredoxin gene (*trx*) in pBAD33et32 was copied from pET32a
(Merck Corp.) to replace the His-patch thioredoxin used in our previous study (18), which was
derived from pBAD202-TOPO (Invitrogen Inc., MA, USA).

For the bacterial endoglucanase gene *bscel5A*, the putative signal peptide sequence (M1–A28)
was removed from the full-length construct. Full-length (A29–N499) and catalytic domain (A29–
T332) genes were inserted between BamHI and XhoI sites in pGEX4T-3 (Cytiva Inc., MA, USA)
to express the N-terminal GST-fused endoglucanase BsCel5A.

Deletion of *bcsA* gene from *Escherichia coli* BL21

In this study, we used genetically modified *E. coli* BL21 to express Bcs proteins, which lack
the whole *bcsA* gene (Fig. 1A). This gene manipulation was performed using the Red/ET
recombination system (Gene Bridges GmbH, Heidelberg, Germany), according to the
manufacturer's specifications. The detailed procedure is described in Supplementary Information.

Briefly, *E. coli* BL21 strain (purchased from New England Labs Inc., MA, USA) was
transformed with the pRedET plasmid carrying the genes for DNA-recombination proteins and
selected by tetracycline resistance. The second transformation was followed by introducing linear
DNA of neomycin-resistant gene flanked by the homology arm sequences corresponding to the
upstream and downstream of *bcsA* gene (CP010816.1); the recombinant proteins were expressed
with arabinose-induction for disrupting *bcsA* gene, and subsequently the pRedET plasmid was
expelled out of the cell by culturing at 37°C. Finally, the *bcsA-disrupted* BL21 transformant was
selected for kanamycin resistance and tetracycline sensitivity. Gene modification was confirmed
by direct PCR (Fig. 1B). We named this genetically modified *E. coli* BL21 strain as BL21-GM.
Chemically competent cells of BL21-GM were then prepared by treating the cultured cells in a
rubidium-based buffer according to the protocol on the website of the Center for Molecular
Biology and Genetics at Mie University (https://www.gene.mie-
u.ac.jp/Protocol/Original/Transform-Comp-Cell-Freeze.html, in Japanese) (33).

In vivo cellulose synthase activity

The endogenous cellulose-synthesizing activity of *E. coli* and cellulose-synthesizing activity of
the heterogeneously expressed Bcs-complex in *E. coli* was estimated by expressing DGC protein
to activate bacterial cellulose synthase in the cell by the produced c-di-GMP (18). In this study,
the pBAD33et32-tDGC plasmid was used to express the DGC. The obtained *E. coli*
transformants were pre-cultured in LB medium including the proper antibiotics at 37°C
overnight, and then 0.1 mL of the pre-culture was inoculated into 10 mL of ZYP-5052 medium
with 50 μg/mL chloramphenicol (100 μg/mL ampicillin when expressing Bcs-complex) and
0.001% L-arabinose. Cells were grown at 20°C in an orbital shaker at 180 rpm for 24 h and the
cultured cells were collected by centrifugation at 2,000 × *g* for 20 min. The amount of cellulose in
the pellet was quantified using the anthrone-sulfuric acid method (34) as described below.

Two milliliters of AN-reagent (a mixture of glacial acetic acid and nitric acid with volumetric
ratio of 8:1) were added to the pellet and then heated at 105°C for 30 min. The remaining
cellulose pellets were washed with 10 mL of water once by centrifugation at 2,000 × *g* for 10
min, and then 1 mL of 72% (v/v) H_2SO_4 was added to the pellets and incubated for 1 h at room
temperature. The solution (0.1 mL) was mixed with water (0.4 mL), and then, 1 mL of fresh
anthrone reagent (0.2% w/v anthrone in concentrated H_2SO_4) was added and mixed. The solution
was heated to 90°C for 15 min and allowed to stand until it reached room temperature
(approximately 15 min). The absorbance of the solution was measured at both 625 and 700 nm,
and the absorbance at 625 nm, from which the absorbance at 700 nm was subtracted, was used for
quantification for better reproducibility. Calibration was performed with phosphoric acid-swollen
cellulose (PASC) prepared using a previously reported protocol (35).

Purification and characterization of BsCel5A

Expression of GST-fused BsCel5A endoglucanase was performed in *E. coli* BL21 (not GM
strain): all the four constructs ((BsCel5A_FL, BsCel5A_CD, BsCel5A_FL^m, and BsCel5A_CD^m)
were expressed and purified in the same manner.

The transformant of BL21 with pGEX-BsCel5A were inoculated in 10 mL LB containing 100
μg/mL ampicillin in a 100 mL flask and grown at 37°C with orbital shaking at 220 rpm overnight.
The cells were harvested by centrifugation at 3,000 × *g* for 20 min, suspended in 1 L of LB
containing 100 μg/mL ampicillin in a 5 L baffle flask, and cultured at 37°C until OD_{600} reached
0.6. The culture medium was cooled in ice water for 10 min, and then IPTG was added to 0.1 mM
and incubated at 20°C for 20 h at 160 rpm with shaking.

The cells were then collected at 3,000 × *g* for 20 min, washed with 40 mL of 1 × phosphate-
buffered saline (PBS), suspended in 40 mL of 1 × PBS containing 1 mM phenylmethylsulfonyl
fluoride (PMSF), and sonicated on ice for ~5 min (exposure: 10 s, rest: 50 s; until slightly
translucent). The cell lysate clarified by 20,000 × *g* for 30 min at 2°C was applied to 5 mL of
Glutathione Sepharose 4B (Cytiva Inc,), washed with 20 mL of 1 × PBS, and eluted with 5 c.v. of
0.1 M Tris-HCl pH 8.0, 0.5 M NaCl, 20 mM reduced-glutathione. The eluted fraction was buffer
exchanged with 1 × PBS using Vivaspin 6 (MWCO 10 kDa; Sartorius AG, Germany).

To remove the GST tag fused to the N-terminus, 100 units of thrombin (Merck Corp.) was
added to 10 mg of $_{GST}$BsCel5A and incubated at 20°C for two days. Thrombin was then removed
by adding 1 mL of Benzamidine Sepharose 6 B (Cytiva Inc.), and the cleaved GST was removed
by adding 5 mL of Glutathione Sepharose 4 B. A solution of PMSF was added to the obtained

BsCel5A protein at a final concentration of 1 mM to inactivate a small amount of residual
thrombin.

The recombinant BsCel5A proteins purified were characterized with regard to the following
points: (i) binding activity to cellulose, (ii) glycoside hydrolase activity, and (iii) substrate
specificity. The detailed experimental procedure is described in Supplementary Information.

Purification of BcsABCD complex

All the protein purification experiments in this study were performed at low temperature
(~4°C). The composition of the protease inhibitor mix (PI) used for protein purification was as
follows: 0.1 mM PMSF, 1 mM benzamidine-HCl, 5 μM bestatin, 10 μM leupeptin, 1 μM
pepstatin A.

The transformant of *E. coli* BL21-GM with pQE-BcsABCD$_{H6}$ was inoculated in 10 mL of LB
containing 100 μg/mL ampicillin in a 100 mL flask and grown at 37°C with shaking at 220 rpm
overnight. The cells were harvested by centrifugation at $3,000 \times g$ for 20 min, resuspended in 1 L
of ZYP-5052 containing 100 μg/mL ampicillin and 0.0005% (v/v) antifoam SI (Fujifim Wako
Chemicals Co. Ltd., Japan) in a 5 L baffle flask, and incubated at 20°C for 22 h with an orbital
shaker at 100 rpm. The cultured cells were collected at $3,000 \times g$ for 20 min and suspended in 50
mL of buffer A (20 mM sodium phosphate buffer (pH 7.5), 300 mM NaCl, 8% glycerol, 5 mM
MgCl$_2$, 5 mM cellobiose) with PI, prior to cell lysis by two passes of high-pressure homogenizer
(French Press, Thermo Fisher Scientific Inc., MA, USA) at a pressure of 15,000 psi. The lysate
clarified by centrifugation at $5,000 \times g$ for 10 min and ultracentrifugation at $150,000 \times g$ for 30
min at 2°C. The obtained cell membranes were resuspended in buffer A with PI and NaCl
concentrations increased to 1 M and were ultracentrifuged again under the same conditions.
Finally, the cell membrane was suspended in buffer A with PI to approximately 10 mg/mL
(estimated by the DC protein assay, Bio-Rad Inc., CA, USA). The collected cell membrane was
stored at -80°C until use in experiments.

Ten milliliters of the cell membrane suspension were mixed with *n*-dodecyl-β-D-
maltopyranoside (DDM) at a final concentration of 0.5% (w/v) and shaken gently for 1 h at 4°C
for solubilizing Bcs-complex. Supernatant of the following ultracentrifugation at $150,000 \times g$ for
30 min at 2°C was collected as solubilized fraction, and imidazole stock solution (2 M, pH 8.0)
was added to achieve a final concentration of 10 mM, prior to loading to 0.4 mL a His-Select
Cobalt Affinity Gel (Sigma-Aldrich Inc.), which was equilibrated with 4 mL of wash buffer (20
mM imidazole and 0.025% (w/v) DDM in buffer A) in advance. After loading the solubilized
protein onto the affinity resin in a plastic open column, the resin in the column was washed with 3
mL of wash buffer and finally eluted with elution buffer (200 mM imidazole and 0.025% (w/v)
DDM in buffer A).

Cellulase treatment of Bcs-complex

Cellulase treatment of BcsABCD$_{H6}$ was performed using a cobalt affinity gel. The same
procedure was followed for the binding step of BcsABCD$_{H6}$ on the affinity gel. After all the
solubilized fraction passed through the column, 0.8 mL of 1.5 μM BsCel5A solution in wash
buffer (20 mM imidazole and 0.025% (w/v) DDM in buffer A) was loaded to the column. The

column exit was plugged when most of the cellulase solution passed, and then the gel was
incubated at 10°C for 3 h with the remaining cellulase solution. After unplugging, the affinity gel
in the column was washed with 2 mL of wash buffer and eluted with elution buffer (200 mM
imidazole and 0.025% (w/v) DDM in buffer A).

Protein preparation for pull-down assay: BcsC_TPR, BcsD, and crude BcsAB

The pull-down assay in this study used purified BcsC_TPR, purified BcsD, and crude BcsAB
proteins, each of which was prepared as described below.

The transformant of *E. coli* BL21(DE3) with pET-$_{H6}$BcsC_TPR$_{H6}$ was inoculated in 10 mL of
LB containing 50 μg/mL kanamycin in a 100 mL flask and grown at 37°C with shaking at 220
rpm overnight. The cells were harvested by centrifugation at $3,000 \times g$ for 20 min, and
resuspended in 1 L of ZYP-5052 (36) containing 200 μg/mL kanamycin and 0.0005%(v/v)
antifoam SI in a 5 L baffle flask, and incubated at 20°C for 24 h in a shaking incubator at 100
rpm. The cultured cells were collected at $3,000 \times g$ for 20 min, and washed with 40 mL of $1 \times$
PBS prior to resuspension in 100 mL of 10 mM imidazole in buffer B (20 mM sodium phosphate
buffer (pH 7.5), 500 mM NaCl, 8% glycerol) with PI, and sonicated on ice for ~5 min (exposure:
10 s, rest: 50 s; until slightly translucent). The cell lysate was clarified by centrifugation at 40,000
$\times g$ for 30 min at 2°C and applied to 4 mL a His-Select Cobalt Affinity Gel (Sigma-Aldrich Inc.),
followed by washing with 10 mM imidazole in buffer B, and eluted 250 mM imidazole in buffer
B. Finally, the buffer of eluted fraction including $_{H6}$BcsC_TPR$_{H6}$ protein was replaced with buffer
B using PD-10 desalting column (Cytiva Inc.).

The BcsD protein was expressed in *E. coli* BL21(DE3) with pET-BcsD$_{StrepII}$. The transformant
was inoculated in 10 mL of LB containing 50 μg/mL kanamycin in a 100 mL flask and grown at
37°C with shaking at 220 rpm overnight. The cells were harvested by centrifugation at $3,000 \times g$,
20 min, resuspended in 1 L of LB containing 50 μg/mL kanamycin in a 5 L baffle flask and
cultured at 37°C until OD$_{600}$ was ~0.6. The culture medium was cooled in ice water for 30 min,
and IPTG (0.5 mM) was added and incubated at 25°C for 20 h at 120 rpm. Cells were collected at
$3,000 \times g$ for 20 min, suspended in 30 mL of buffer C (20 mM Tris-HCl (pH 7.5), 150 mM NaCl,
and 1 mM EDTA) with PI, and sonicated on ice for ~10 min (exposure: 10 s, rest: 50 s; until
slightly translucent). The cell lysate was centrifuged at $20,000 \times g$ for 30 min at 2°C, applied to 2
mL of Streptactin Sepharose HP (Cytiva Inc.), washed with 20 mL of buffer C, and eluted with 4
mM *d*-desthiobiotin in buffer C. Finally, buffer of eluted fraction was replaced by buffer C using
Amicon Ultra-4 MWCO 50 kDa (Merck Corp.).

The crude protein of BcsAB was prepared using the transformant *E. coli* BL21-GM with pQE
carrying the *bcsAB* gene. The procedure to isolate the cell membrane, from culture to
ultracentrifugation, was same as in the case of BcsABCD$_{H6}$ as described previously, except for
the buffer to suspend the cells and cell membrane: buffer D (20 mM sodium phosphate buffer (pH
7.5), 100 mM NaCl, 8% glycerol, 5 mM MgCl$_2$, and 5 mM cellobiose) with PI was used. The
obtained cell membrane was suspended in buffer D with PI at approximately 10 mg/mL
(estimated by the DC protein assay, Bio-Rad Inc.). Finally, crude BcsAB protein was prepared by
solubilizing 7 mL of the cell membrane fraction with 0.5% (w/v) DDM for 1 h and centrifugation
at $150,000 \times g$ for 30 min at 2°C.

Pull-down assays

For the pull-down by BcsD$_{StrepII}$ with the crude BcsAB, purified BcsD$_{StrepII}$ (0.2 mg) was added to 7 mL of crude BcsAB, incubated for 3 h at 4 °C, and then loaded to 0.1 mL of Streptactin Sepharose HP. The resin was washed with 1 mL of 0.025% (w/v) DDM in buffer D, and finally the bound protein was eluted with 4 mM desthiobiotin in buffer D with 0.025% (w/v) DDM.

For the pull-down assay, BcsD$_{StrepII}$ with $_{H6}$BcsC_TPR$_{H6}$, BcsD$_{StrepII}$ (0.1 mg) were incubated with $_{H6}$BcsC_TPR$_{H6}$ (0, 0.1, 0.3, 0.5 mg) in 3 mL of buffer D (–/–) (buffer D without MgCl$_2$ and cellobiose) for 3 h at 4 °C, and then loaded to 0.1 mL of Streptactin sepharose HP. After washing with buffer D (–/–), the bound protein was eluted with 1 mL 4 mM d-desthiobiotin in buffer D (–/–).

For the pull-down assay by $_{H6}$BcsC_TPR$_{H6}$ with BcsAB, purified $_{H6}$BcsC_TPR$_{H6}$ (0.2 mg) was mixed with crude BcsAB (7 mL), and further NaCl stock solution was added to set 500 mM NaCl in the final solution. After incubation for 1 h at 4 °C, the sample was loaded to 0.2 mL a His-Select Cobalt Affinity Gel and washed with 2 mL of 20 mM imidazole in buffer B (+/+) (buffer B with 5 mM MgCl$_2$ and 5 mM cellobiose) with 0.025% (w/v) DDM. The bound protein was eluted with 200 mM imidazole in buffer B containing 0.025% (w/v) DDM.

Finally, we performed a pull-down assay using BcsD$_{StrepII}$, purified BcsC_TPR; crude BcsAB. BcsD$_{StrepII}$ (0.2 mg) and $_{H6}$BcsC_TPR$_{H6}$ (0.2 mg) were mixed with crude BcsAB (7 mL of DDM extract); NaCl stock solution (5 M) was added to a final concentration of 500 mM NaCl. After incubation for 3 h at 4 °C, the sample solution was loaded to 0.1 mL of Streptactin Sepharose HP. After washing with 1 mL of buffer B (+/+) containing 0.025% (w/v) DDM, the bound protein was eluted with 4 mM d-desthiobiotin in buffer B (+/+) containing 0.025% (w/v) DDM.

Prey protein without bait protein was used as a negative control for the pull-down assay using the same procedures. The eluted fraction was analyzed by SDS-PAGE, followed by WB as described later.

Western blotting (WB) analysis

Rabbit polyclonal antibodies against BcsA, BcsB, BcsC, and BcsD proteins were generated with the synthetic peptide corresponding to a part of the C-terminus of BcsA (18, 20, 21, 37), the C-terminus of BcsB (this study), the C-terminus of BcsC (37), and the loop between the β3 and β4-strand (20, 37). The dilution rate was 10,000-fold for anti-BcsA, anti-BcsB, and anti-BcsC, whereas anti-BcsD was diluted 20,000-fold. Each of the antibodies against the peptide tag were diluted as follows: anti-Strep tag II (15 ng/mL; ab180957, Abcam Corp., Cambridge, UK), anti-His5 (0.1 µg/mL; 34660, Qiagen N.V., Germany), and anti-Thioredoxin (1 µg/mL; ab139677, Abcam Corp.).

Horseradish peroxidase (HRP)-conjugated anti-rabbit IgG (W401B; Promega Inc.) and HRP-conjugated anti-mouse IgG (7076S; Cell Signaling Technology Inc., MA, USA) were diluted 20,000-fold and used as the secondary antibodies. ECL-Prime (Cytiva Inc.) was used for chemical luminescence with HRP.

For the WB with anti-BcsD, anti-Strep, anti-His, and anti-thioredoxin, the samples for SDS-PAGE were heat-treated at 100°C for 3 min after mixing with the sample buffer, while the

samples to visualize by the other antibodies were incubated in the sample buffer for several
minutes at room temperature. The samples were analyzed by SDS-PAGE with a precast
polyacrylamide gel (SuperSep Ace 5-20%, Fujifim Wako Chemicals Co. Ltd.), and blotting from
the gel to PVDF membrane (Immobilon-P, 0.45 μm-φ, Cytiva Inc.) was done by a constant
current mode at 150 mA/10 cm^2 for 30 min. Blocking of the membrane was performed with 0.3%
skim milk and 0.02% NaN$_3$ in tris-buffered saline (TBS) containing 0.05% Tween-20 (TBS-T),
for the WB with anti-Bcs proteins, while blocking was performed with 2.5% BSA, 0.1% gelatin,
0.02% Tween-20, 0.04% NaN$_3$ in PBS at 4°C overnight or at room temperature for 0.5–1 h, for
either of anti-Strep-tag II, anti-His5, or anti-thioredoxin. The membranes were briefly rinsed with
TBS-T and then subjected to the primary antibody reaction in blocking buffer for 1–2 h at room
temperature. After removing the antibody solution, the membrane was briefly rinsed and washed
with TBS-T thrice for 5 min, followed by a secondary antibody reaction in TBS-T for 30–45 min
at room temperature. The membrane was briefly rinsed and washed with TBS-T three times for
15 min each time. The protein bands were then visualized by casting the ECL prime solution on
the blotted PVDF membrane (0.2 mL/9 cm^2 membrane), and digitally recorded with AE-9300H
EZ-Capture MG (ATTO Co. Ltd., Japan).

Quantitative analysis was performed using ImageJ (version 1.53a) to quantify the band signal
intensity, calculated as relative values to the negative control, and the relative signal intensities of
BcsA and BcsB were normalized to the signal intensity of BcsD. For BcsA; two major bands
were reproducibly observed near the 75-kDa and 250-kDa marker (Fig. 2B, top row), as well as
in our previous studies (18, 37), which are probably BcsA monomer and BcsA-including
oligomer, respectively (the estimated molecular weight for the BcsA monomer is 84 kDa). Given
these observations, the band intensity for the BcsA-containing oligomer was also considered in
addition to the monomer band intensity in subsequent analyses.

598
599
600 **Acknowledgments**
601

The authors thank Dr. Kenji Tajima of Hokkaido University for his advice. The authors thank
Professor Ryoichi Yamaji and Dr. Yasuyuki Kobayashi at Osaka Prefecture University for their
advice regarding the pull-down assay. This work was supported by JPSP KAKENHI (grant
numbers 15H04530 and 19H00950 to T. I.) and Exploratory Research on Humanosphere Science
(RISH, Kyoto University). We would like to thank Prof. Junji Sugiyama for JST CREST (grant
number: JPMJCR13B2).

608
609
References

1. J. L. W. Morgan, J. Strumillo, J. Zimmer, Crystallographic snapshot of cellulose synthesis
and membrane translocation. *Nature* **493**, 181-186 (2013).
2. J. L. W. Morgan, J. T. McNamara, J. Zimmer, Mechanism of activation of bacterial
cellulose synthase by cyclic di-GMP. *Nat. Struct. Mol. Biol.* **21**, 489-496 (2014).
3. U. Römling, M. Y. Galperin, Bacterial cellulose biosynthesis: Diversity of operons,
subunits, products, and functions. *Trends Microbiol.* **23**, 545-557 (2015).
4. W. Thongsomboon *et al.*, Phosphoethanolamine cellulose: A naturally produced
chemically modified cellulose. *Science* **359**, 334-338 (2018).
5. K. Tajima, T. Imai, T. Yui, M. Yao, I. Saxena, Cellulose-synthesizing machinery in bacteria.
Cellulose (2021). https://doi.org/10.1007/s10570-021-04225-7

6. H. C. Wong *et al.*, Genetic organization of the cellulose synthase operon in Acetobacter
xylinum. *Proc. Natl. Acad. Sci.U.S.A.* **87**, 8130-8134 (1990).
7. I. M. Saxena, K. Kudlicka, K. Okuda, R. M. Brown Jr, Characterization of genes in the
cellulose-synthesizing operon (acs operon) of Acetobacter xylinum: Implications for
cellulose crystallization. *J. Bacteriol.* **176**, 5735-5752 (1994).
8. R. Standal *et al.*, A new gene required for cellulose production and a gene encoding
cellulolytic activity in Acetobacter xylinum are colocalized with the bcs operon. *J.*
Bacteriol. **176**, 665-672 (1994).
9. S. Kawano *et al.*, Effects of endogenous endo-β-1,4-glucanase on cellulose biosynthesis in
Acetobacter xylinum ATCC23769. *J. Biosci. Bioeng.* **94**, 275-281 (2002).
10. K. Tajima *et al.*, Cloning and sequencing of the beta-glucosidase gene from *Acetobacter*
xylinum ATCC 23769. *DNA Res.* **8**, 263-269 (2001).
11. N. Tonouchi *et al.*, A beta-glucosidase gene downstream of the cellulose synthase operon
in cellulose-producing acetobacter. *Biosci. Biotechnol. Biochem.* **61**, 1789-1790 (1997).
12. J. T. McNamara, J. L. W. Morgan, J. Zimmer (2015) A molecular description of cellulose
biosynthesis. in *Annu. Rev. Biochem.* (Annual Reviews Inc.), pp 895-921.
13. A. Kumagai *et al.*, Ultrafine Cellulose Fibers Produced by Asaia bogorensis, an Acetic Acid
Bacterium. *Biomacromolecules* **12**, 2815-2821 (2011).
14. P. A. Penttilä *et al.*, Multimethod approach to understand the assembly of cellulose fibrils
in the biosynthesis of bacterial cellulose. *Cellulose* (2018).
15. S. Q. Hu *et al.*, Structure of bacterial cellulose synthase subunit D octamer with four inner
passageways. *Proc. Natl. Acad. Sci.U.S.A.* **107**, 17957-17961 (2010).
16. N. Sunagawa *et al.*, Cellulose complementing factor (Ccp) is a new member of the cellulose
synthase complex (terminal complex) in *Acetobacter xylinum*. *J. Biosci. Bioeng.* **115**, 607-
612 (2013).
17. O. Omadjela *et al.*, BcsA and BcsB form the catalytically active core of bacterial cellulose
synthase sufficient for in vitro cellulose synthesis. *Proc. Natl. Acad. Sci.U.S.A.* **110**, 17856-
17861 (2013).
18. T. Imai, S.-j. Sun, Y. Horikawa, M. Wada, J. Sugiyama, Functional reconstitution of
cellulose synthase in *Escherichia coli. Biomacromolecules* **15**, 4206–4213 (2014).
19. J. Du, V. Vepachedu, S. H. Cho, M. Kumar, B. T. Nixon, Structure of the cellulose synthase
complex of *Gluconacetobacter hansenii* at 23.4 Å resolution. *PLoS ONE* **11**, e0155886
(2016).
20. S.-j. Sun, T. Imai, J. Sugiyama, S. Kimura, CesA protein is included in the terminal
complex of Acetobacter. *Cellulose* **24**, 2017–2027 (2017).
21. S.-j. Sun, Y. Horikawa, M. Wada, J. Sugiyama, T. Imai, Site-directed mutagenesis of
bacterial cellulose synthase highlights sulfur–arene interaction as key to catalysis.
Carbohydr. Res. **434**, 99–106 (2016).
22. J. F. Acheson, R. Y. Ho, N. F. Goularte, L. Cegelski, J. Zimmer, Molecular organization of
the E. coli cellulose synthase macrocomplex. *Nat. Struct. Mol. Biol.* **28**, 310-318 (2021).
23. C. R. Santos *et al.*, Dissecting structure-function-stability relationships of a thermostable
GH5-CBM3 cellulase from *Bacillus subtilis* 168. *Biochem. Journal* **441**, 95-104 (2012).
24. J. F. Acheson, Z. S. Derewenda, J. Zimmer, Architecture of the Cellulose Synthase Outer
Membrane Channel and Its Association with the Periplasmic TPR Domain. *Structure* **27**,
1855-1861 (2019).
25. S. Nojima *et al.*, Crystal structure of the flexible tandem repeat domain of bacterial
cellulose synthesis subunit. *Sci. Rep.* **7**, 13018 (2017). https://doi.org/10.1038/s41598-017-
12530-0
26. A. Perez-Riba, L. S. Itzhaki, The tetratricopeptide-repeat motif is a versatile platform that

enables diverse modes of molecular recognition. *Curr. Opin. Struct. Biol.* **54**, 43-49 (2019).

27. W. Abidi, S. Zouhir, M. Caleechurn, S. Roche, P. V. Krasteva, Architecture and regulation
of an enterobacterial cellulose secretion system. *Sci. Adv.* **7**, eabd8049 (2021).

28. T. Uto *et al.*, Molecular Dynamics Simulation of Cellulose Synthase Subunit D Octamer
with Cellulose Chains from Acetic Acid Bacteria: Insight into Dynamic Behaviors and
Thermodynamics on Substrate Recognition. *J.Chem. Theory Comput.* **17**, 488-496 (2021).

29. H. Toyosaki *et al.*, Screening of Bacterial Cellulose-producing Acetobacter Strains Suitable
for Agitated Culture. *Biosci. Biotechnol. Biochem.* **59**, 1498-1502 (1995).

30. T. Nakai *et al.*, Control of expression by the cellulose synthase (*bcsA*) promoter region
from *Acetobacter xylinum* BPR 2001. *Gene* **213**, 93-100 (1998).

31. F. Rao *et al.*, Enzymatic synthesis of c-di-GMP using a thermophilic diguanylate cyclase.
Analytical Biochemistry **389**, 138-142 (2009).

32. A. Crameri, E. A. Whitehorn, E. Tate, W. P. C. Stemmer, Improved green fluorescent
protein by molecular evolution using DNA shuffling. *Nat. Biotechnol.* **14**, 315-319 (1996).

33. https://www.gene.mie-u.ac.jp/Protocol/Original/Transform-Comp-Cell-Freeze.html, the
latest access date: Mar 26 2022

34. D. M. Updegraff, Semimicro determination of cellulose inbiological materials. *Anal.*
Biochem. **32**, 420-424 (1969).

35. J. Stahlberg, G. Johansson, G. Pettersson, Trichoderma reesei has no true exo-cellulase: All
intact and truncated cellulase produce new reducing end groups on cellulose. *Biochim.*
Biophys. Acta Gen. Subj. **1157**, 107-113 (1993).

36. F. W. Studier, Protein production by auto-induction in high density shaking cultures.
Protein Expr. Purif. **41**, 207-234 (2005).

37. A. Hashimoto *et al.*, Extraction of cellulose-synthesizing activity of *Gluconacetobacter*
xylinus by alkylmaltoside. *Carbohydr. Res.* **346**, 2760–2768 (2011).

695
696

697 **Figures and Tables**

698
699

Figure 1. Deletion of the *E. coli* endogenous *bcsA* gene

A Schematic diagram of the experiment for deleting the endogenous *bcsA* gene using the Red/ET
recombination system. The primer sequences used are listed in Table S1. **B** Electrophoresis
pattern of colony-direct PCR using two pairs of primers. M: Molecular weight marker (Marker 6
(Nippon Gene Co. Ltd., Tokyo, Japan)); Δ*bcsA*: *bcsA*-deleted BL21; WT: wild-type BL21. **C**
Quantification of endogenous cellulose synthesis in BL21-GM and BL21 cells. The CESEC
system (18), which uses DGC expression to synthesize c-di-GMP to activate bacterial cellulose
synthase, was used to count the amount of cellulose that used DGC expression (bottom). The
expression of DGC was confirmed by WB using an anti-Trx antibody (top). The experiments
were performed in triplicate ($n = 3$) and are represented as mean ± standard error of the mean
(SEM), together with each of the data values. Statistical tests (Student's t-test) were performed
using the built-in function in Microsoft Excel (TTEST).

Figure 2. *in vivo* cellulose-synthesizing activity of recombinant Bcs-complex

A Cellulose-synthesizing activity of the recombinant Bcs-complexes tested in this study: the amount of cellulose synthesized by the CESEC system (18) with BL21-GM ($\Delta bcsA$) is shown for each recombinant protein. All experiments were performed in triplicate ($n = 3$) and are represented as mean ± SEM together with the significance of difference: * for $p < 0.05$, ** for $p < 0.01$, *** for $p < 0.001$, and *n.s.* for not significant ($p > 0.1$) as a result of Student's t-test, which was performed using the built-in function of Microsoft Excel (TTEST). **B** Western blotting (WB) was performed for the CESEC experiment to check the expression of BcsA, BcsB, BcsC, BcsD, and $_{TRX}$DGC. Cultures with and without DGC expression were tested. For BcsA, the data are shown separately for the oligomer and monomer bands. Due to the limited number of wells in the SDS-PAGE gel, the gel was divided into two parts for electrophoresis.

Figure 3. Purification of BcsABCD_{H6}

A Western blotting (WB) analysis of the purification process: M, membrane fraction; S,
solubilized fraction; F, IMAC unbound fraction; E, IMAC-eluted fraction. Three protein samples,
wild-type (BcsABCD_{H6}), D333N on BcsA (BcsA^{D333N}BCD_{H6}), and W373A on BcsA
(BcsA^{W373A}BCD_{H6}), were analyzed using the antibodies indicated. **B** WB analyses of the eluate
from three independent purification experiments for each of the three protein samples.

Approximately 2 µg of the protein was loaded onto the gel for WB with BcsA, BcsB, and BcsC
antibodies, and 1 µg of the protein was added to the gel for WB with BcsD antibody. For BcsA,
the data are shown separately for the oligomer and monomer bands.

Figure 4. Effect of cellulase treatment to the purification of Bcs-complex

A, Western blot images of IMAC-eluted fractions of $BcsWTBCD_{H6}$ after treatment with wild-type cellulase (BsCel5A_FL and BsCel5A_CD). Coomassie Brilliant Blue (CBB)-stained images were used as loading controls. **B** Band intensity in western blot analysis of **A**. Error bars indicate standard error (SE), and the significance of the difference as a result of Student's t-test, which was performed using the built-in function of Microsoft Excel (TTEST), is indicated by the symbols: ** for $p < 0.01$, *** for $p < 0.001$, and *n.s.* for not significant ($p > 0.1$). **C**, Western blot images of IMAC-eluted fractions of $BcsWTBCD_{H6}$ after cellulase mutant (BsCel5A_FLm and BsCel5A_CDm) treatment. CBB-stained images were visualized using BcsD as a loading control. **D**, Band intensity in western blot analysis of **C**. Error bars indicate SEM, and Student's t-test, which was performed using the built-in function of Microsoft Excel (TTEST), showed no significant difference ($p > 0.1$; eight independent replicates).

For BcsA, the data are shown separately for the oligomer band, monomer band, and their sum. Each elution sample (~0.2 mg/mL) was added to an SDS-PAGE gel in 10 µL (anti-BcsA and anti-BcsB) and 1 µL (anti-BcsD and CBB stain), respectively. Elution samples for CBB staining were loaded into gel wells 10-fold more than those for WB (BcsD lane).

Figure 5. Effect of cellulase treatment on the purification of mutant Bcs-complex

A Western blot images of IMAC-eluted fractions of BcsWTBCD$_{H6}$, BcsA^{D333N}BCD$_{H6}$, and BcsA^{W373A}BCD$_{H6}$. CBB-stained images were visualized using BcsD as a loading control. **B** Band intensity in western blot analysis of **A**. An error bar indicates SEM, and the significance of the difference as a result of Student's *t*-test, which was performed using the built-in function of Microsoft Excel (TTEST), is indicated by the symbols † for $p < 0.1$, ** for $p < 0.01$, *** for $p < 0.001$, and *n.s.* for not significant ($p > 0.1$). **C**, Western blot images of IMAC-eluted fractions of BcsA^{D333N}BCD$_{H6}$ and BcsA^{W373A}BCD$_{H6}$ after treatment with wild-type cellulase (BsCel5A_FL and BsCel5A_CD). **D** and **E** Band intensity on western blot analysis of **C**: D for BcsA^{D333N}BCD and E for BcsA^{W373A}BCD. An error bar indicates SEM, and the significance of the difference as a result of Student's *t*-test, which was performed using the built-in function of Microsoft Excel (TTEST), showed no significant difference between any dataset.

All experiments were performed with four independent replicates ($n = 4$). For BcsA, the data are shown separately for the oligomer band, monomer band, and their sum. Each elution sample (~0.2 mg/mL) was added to an SDS-PAGE gel in 10 µL (anti-BcsA and anti-BcsB) and 1 µL (anti-BcsD and CBB stain), respectively. Elution samples for CBB staining were loaded into gel wells 10-fold more than those for WB (BcsD lane).

Figure 6. Pull-down assay with the Bcs subunits

A Schematic diagram showing the partners (bait and prey proteins) in a pull-down assay for each subunit of BcsABCD. Alphabets on the cartoon indicate panels B–D (the result of pull-down experiments), by which the subunit-subunit interaction was surveyed. **B** Western blot images of the pull-down assay using BcsD$_{StrepII}$ (bait) with crude BcsAB (prey). The pull-down fraction was analyzed by western blot using anti-BcsA, anti-BcsB, and anti-Streptag antibodies. **C** Western blot images of the pull-down assay using BcsD$_{StrepII}$ (bait) with purified $_{H6}$BcsC_TPR$_{H6}$ (prey). The pull-down fraction was analyzed by CBB staining and western blotting using anti-His5 and anti-Streptag antibodies. **D** Western blot images of the pull-down assay using $_{H6}$BcsC_TPR$_{H6}$ (bait) with crude BcsAB (prey). The pull-down fraction was analyzed by western blot using anti-BcsA, anti-BcsB, and anti-His5 antibodies. **E** Western blot images of the pull-down assay using BcsD$_{StrepII}$ (bait) with crude BcsAB (prey_1) and purified $_{H6}$BcsC_TPR$_{H6}$ (prey_2). The pull-down fraction was analyzed by western blot using anti-BcsA, anti-BcsB, anti-His5, and anti-Streptag antibodies. Input: sample immediately after mixing; pull-down: elution fraction after pull-down. All the detailed methods are described in the "Experimental Procedures" section of the text.

Figure 7. The hypothetical two-state model for the type I Bcs-complex

Hypothetical two-state model for the Bcs-complex: basal state (left) and active state (right). In this model, BcsD plays a key role in the periplasmic space to control the state of the complex separately from the state of the BcsA enzyme (1, 2). In the active state (right), BcsD draws the glucan chains in its octameric ring and brings those chains to the channel pore of BcsC in the outer membrane like a "button whirligig'. The TPR domain in BcsC may escort the BcsD protein to correctly transfer the glucan chains in BcsD to the BcsC channel. In the basal state (left), BcsD docks with the BcsB complex given that BcsB is located in the periplasmic space (1) as well as BcsD (12, 20). In the model, BcsD in the basal state also docks with BcsC without interfering with BcsB and is ready to draw the glucan chains in it once BcsAB starts synthesizing and extruding out the glucan chain. Thus, BcsD in this state may be a standby state for effectively transferring glucan chains into the extracellular space through the BcsC channel once BcsA is activated. Subunits that were not assessed in this study are indicated in gray.

808 Table 1. Experimental resources to express the proteins in this study

Material	Source	Description	Reference
Bacterial cellulose synthase			
pQE-BcsAB (GOI* was inserted between EcoRI and KpnI site in pQE-80L vector (Qiagen Inc.))	LC428282 (*Komagataeibacter sucrofermentans* JCM9730)	Expression vector for BcsAB protein without tags	Imai et al. 2014
pQE-BcsA^{D333N}B (GOI* was inserted between EcoRI and KpnI site in pQE-80L vector (Qiagen Inc.))	LC428282 (*Komagataeibacter sucrofermentans* JCM9730)	Expression vector for BcsAB protein with a catalytic-deficient point mutation of D333N, without tags	Sun et al., 2016
pQE-BcsA^{W373A}B (GOI* was inserted between EcoRI and KpnI site in pQE-80L vector (Qiagen Inc.))	LC428282 (*Komagataeibacter sucrofermentans* JCM9730)	Expression vector for BcsAB protein with a catalytic-deficient point mutation of W373A, without tags	Sun et al., 2016
pQE-BcsABCD (GOI was inserted between EcoRI and HindIII site in pQE-80L vector (Qiagen Inc.))	LC687564 (*Komagataeibacter sucrofermentans* JCM9730)	Expression vector for BcsABCD complex without tags	This study
pQE-BcsABCD$_{H6}$ (GOI was inserted between EcoRI and HindIII site in pQE-80L vector (Qiagen Inc.))	LC687564 (*Komagataeibacter sucrofermentans* JCM9730)	Expression vector for BcsABCD complex with hexahistidine-tag at the C-terminal of BcsD protein	This study
pQE-BcsABC $_{H6}$D (GOI was inserted between EcoRI and HindIII site in pQE-80L vector (Qiagen Inc.))	LC687564 (*Komagataeibacter sucrofermentans* JCM9730)	Expression vector for BcsABCD complex with hexahistidine-tag at the N-terminal of BcsD protein	This study
pQE-BcsABCD$_{GFPuvH6}$ (GOI was inserted between EcoRI and HindIII site in pQE-80L vector (Qiagen Inc.))	LC687564 (*Komagataeibacter sucrofermentans* JCM9730)	Expression vector for BcsABCD complex with GFPuv followed by hexahistidine-tag at the C-terminal of BcsD protein	This study
pQE-BcsA^{D333N}BCD$_{H6}$ (GOI was inserted between EcoRI and HindIII site in	LC687564 (*Komagataeibacter*	Expression vector for BcsABCD complex with a catalytic-deficient point mutation of D333N	This study

pQE-80L vector (Qiagen Inc.))	*sucrofermentans* JCM9730)	introduced in BcsA; hexahistidine-tag is fused to the C-terminal of BcsD protein	
pQE-BcsA^{W373A}BCD_{H6} (GOI was inserted between EcoRI and HindIII site in pQE-80L vector (Qiagen Inc.))	LC687564 (*Komagataeibacter sucrofermentans* JCM9730)	Expression vector for BcsABCD complex with a catalytic-deficient point mutation of W373A introduced in BcsA; hexahistidine-tag is fused to the C-terminal of BcsD protein	This study
pET-_{H6}BcsC_TPR_{H6} (GOI was inserted between NdeI and XhoI site in pET28b (Novagene Inc.))	LC687564 (*Komagataeibacter sucrofermentans* JCM9730)	Expression vector for TPR domain of BcsC protein (Cys18 – Gly848): hexahistidine-tag is fused to both N- and C-terminal of the protein	This study
pET-BcsD_{StrepII} (GOI was inserted between NcoI and HindIII site in pET28b vector (Novagen Inc.))	LC687564 (*Komagataeibacter sucrofermentans* JCM9730)	Expression vector for BcsD protein: Streptag(II) is fused to the C-terminal of BcsD protein	This study

Diguanylate cyclase (DGC)			
pBAD33et32-tDGC (Self-made vector in the previous study)	NP_229585 (*Thermotoga maritima* MSB8)	Expression vector of diguanylate cyclase with N-terminal truncation (K82 – S248) and a point mutation (R158A mutation): thioredoxin was fused to Nterminal end of the protein	This study

Cellulase			
pGEX-BsCel5A_FL (GOI was inserted between BamHI and XhoI site in pGEX4T-3 (GE Healthcare Inc.))	Z29076 (*Bacillus subtilis* strain 168)	Expression vector for full-length endoglucanase protein from *Bacillus subtilis* (BsCel5A): signal-peptide sequence (from Met1 to Ala28) was removed; GST-tag is fused to the N-terminal of BsCel5A protein, which consists of glycosylhydrolase 5 domain (GH-5 domain, Ala29 – Thr332) and carbohydrate-binding module (CBM-3, Ile354 – Asn499)	This study
pGEX-BsCel5A_CD (GOI was inserted between BamHI and XhoI site in pGEX4T-3 (GE Healthcare Inc.))	Z29076 (*Bacillus subtilis* strain 168)	Expression vector for catalytic domain (A29 – T332) of BsCel5A: GST-tag is fused to the N-terminal of the protein, which lacks the signal peptide sequence (Met1 – Ala28)	This study

pGEX-BsCel5A_FLᵐ (GOI was inserted between BamHI and XhoI site in pGEX4T-3 (GE Healthcare Inc.))	*Z29076 (Bacillus subtilis* strain 168)	Expression vector for full-length BsCel5A with activity-deficient double mutation (E169Q/E257Q): GST-tag is fused to the N-terminal of the protein, which lacks the signal peptide sequence (Met1 – Ala28)	This study
pGEX-BsCel5A_CDᵐ (GOI was inserted between BamHI and XhoI site in pGEX4T-3 (GE Healthcare Inc.))	*Z29076 (Bacillus subtilis* strain 168)	Expression vector for catalytic domain (A29 – T332) of BsCel5A with activity-deficient double mutation (E169Q/E257Q): GST-tag is fused to the N-terminal of the protein, which lacks the signal peptide sequence (Met1 – Ala28)	This study

811 * GOI: gene of interest