Transcriptomic profiling implicates PAF1 in both active and repressive immune regulatory networks

Matthew W. Kenaston[1], Oanh H. Pham[1], Marine J. Petit[1,#a], and Priya S. Shah[1,2*]

[1] Department of Microbiology and Molecular Genetics, University of California, Davis, Davis, California, USA.

[2] Department of Chemical Engineering, University of California, Davis, Davis, California, USA.

* Corresponding author

E-mail: prsshah@ucdavis.edu

[#a] Current Address: University of Glasgow MRC-Centre for Virus Research, Glasgow, United Kingdom

Running Title: PAF1 and Immune Gene Expression

Abstract

Background: Sitting at the interface of gene expression and host-pathogen interaction, polymerase associated factor 1 complex (PAF1C) is a rising player in the innate immune response. The complex localizes to the nucleus and associates with chromatin to modulate RNA polymerase II (RNAPII) elongation of gene transcripts. Performing this function at both proximal and distal regulatory elements, PAF1C interacts with many host factors across such sites, along with several microbial proteins during infection. Therefore, translating the ubiquity of PAF1C into specific impacts on immune gene expression remains especially relevant.

Results: Advancing past work, we treat PAF1 knockout cells with a slate of immune stimuli to identify key trends in PAF1-dependent gene expression with broad analytical depth. From our transcriptomic data, we confirm PAF1 is an activator of traditional immune response pathways as well as other cellular pathways correlated with pathogen defense. With this model, we employ computational approaches to refine how PAF1 may contribute to both gene activation and suppression. Specifically focusing on transcriptional motifs, we predict gene regulatory elements strongly associated with PAF1, including those implicated in an immune response. Overall, our results suggest PAF1 is potentially involved in innate immunity at several distinct axes of regulation.

Conclusions: By identifying PAF1-dependent gene expression across several pathogenic contexts, we confirm PAF1C to be a key mediator of innate immunity. Combining these transcriptomic profiles with potential regulatory networks corroborates the previously identified functions of PAF1C. With this, we foster new avenues for its study

41 as a regulator of innate immunity, and our results will serve as a basis for targeted study

42 of PAF1C in future validation studies.

43 **Keywords:** polymerase associated factor 1 complex (PAF1C), innate immunity, network

44 analysis, gene expression, transcriptomics, regulatory motifs

Background

A robust innate immune response relies on the early detection of pathogen-associated

molecular patterns (PAMPs) by host pattern recognition receptors. This sensing triggers

various signaling pathways culminating in responsive gene expression meant to prevent

or combat infection. The polymerase associated factor 1 complex (PAF1C) is increasingly

implicated in such a response at the level of gene expression—resulting from its key roles

in transcriptional elongation, chromatin remodeling, and histone modification. The primary

members of this nuclear complex are PAF1, LEO1, CTR9 and CDC73 [1]. First

discovered as a homolog in yeast [2], PAF1C serves as a scaffold for recruitment of

histone- and RNA polymerase II (RNAPII)-modifying enzymes [3–7]. At these sites,

PAF1C activity regulates the release of a paused RNAPII at promoters, permitting

productive elongation of the transcript [8–10].

Interestingly, PAF1C occupation of certain super-enhancers in cancer cells results in

negative regulation of associated genes [11]. As such, loss of PAF1C in certain contexts

can activate gene expression. Other work has attributed enhancer occupation to PAF1C's

modulation of RNAPII elongation rates rather than direct suppression [12]. Though the

exact mechanisms behind these divergent functions are not fully understood, PAF1C is

poised to extensively modulate gene expression depending on the genes and regulatory

elements present.

PAF1C regulation of gene expression impacts the ability of cells to mount an antimicrobial

response. Specifically, PAF1 depletion has been shown to disrupt interferon-related

signaling and lead to increases in viral replication for both dengue virus and influenza A

virus (IAV) [13–15]. These same studies found such viruses target PAF1 to abrogate its

activation of an immune response, and, recently, similar antagonism has been discovered

in bacteria [16]. PAF1 and various complex members also act as restriction factors for

HIV-1 [17–19]. Others have shown PAF1C to increase tumor necrosis factor (TNF)

expression as part of the inflammatory response [20]. Thus, PAF1C is emerging as an

influential factor in maintaining host innate immunity, though its scope of regulation

remains imprecise.

Here, we computationally interrogate the extent and conservation of PAF1C-mediated

gene expression by treating PAF1 knockout (PAF1 KO) cells with a diverse set of immune

stimuli, including well-recognized PAMPs. Building on previous work, we confirm PAF1

promotes an antiviral response. We further identify the complex as not only a modulator

of innate immunity, but a suppressor of mechanisms that induce a proviral state in cells.

By identifying a conserved set of genes that are PAF1-dependent, we construct various

models to predict potential transcription factors and regulatory elements that may interact

with PAF1C to activate various innate immune pathways. Ultimately, our results

strengthen the model of PAF1C as a crucial player in the host immune response while

providing the necessary foundation for future work.

Results

Shared and distinct immune response pathways in A549 cells

To broaden our identification of PAF1-dependent gene expression, we sought to activate multiple immune pathways. This way, we could differentiate a core set of genes correlated to PAF1 while preserving any nuance for a particular stress response (Table 1). Based on our work using a dsRNA mimic in PAF1 KO cells, we hypothesized that genes both upstream and downstream of type I interferon (IFN-I) signaling were PAF1-dependent [14]. PAF1 KO and non-targeting control (ncgRNA) cells were previously developed in A549 lung epithelial cells, which have been used extensively for the study of the innate immune response [21–23]. We have shown PAF1 Rescue cells recapitulate PAF1-dependent gene expression [14]. Therefore, we opted to use ncgRNA cells as a universal control that has the potential to enable comparisons across multiple gene knockouts in the future.

Both cell lines were treated with the selected stimuli for three hours and subjected to 3' Tag-sequencing (Fig. 1A). Though PAF1 nuclear localization may be marginally disrupted in A549 cells under poly I:C-induced stress, this trend was not observed for LPS or IFN-β (Fig. S1). To capture all the variability of our processed reads, cell genotype (PAF1 KO, ncgRNA) and immune stimuli were treated as distinct conditions within the computational design of our differential gene expression analysis. This allowed for numerous direct comparisons to be made while accounting for inherent differences amongst those factors.

Stimulus	Type	Target	Effect	Reference
poly I:C	dsRNA	RIG-I, MDA5, TLR3	IFN-I production	(Gantier et al., 2007)
poly dA:dT	dsDNA	cytosolic DNA sensors (CDS)	IFN-I production, inflammasome formation	(Unterholzner, 2013); (Hornung et al., 2009)
Lipopolysaccharide (LPS)	G$^{(-)}$ bacteria	TLR4, caspases	Inflammatory cytokine production, inflammasome formation	(Rosadini & Kagan, 2017); (Demon et al., 2014)
Tumor necrosis factor alpha (TNF-α)	inflammatory cytokine	TNFR pathway	Inflammation, apoptosis, NF-κB related pathways	(Kalliolias & Ivashkiv, 2016)
Interferon-beta (IFN-β)	IFN-I	IFNAR1/2, JAK-STAT cascade	interferon-stimulated gene (ISG) expression	(Mesev et al., 2019)

105 **Table 1. Summary of innate immune stimuli.**

A

B

Log$_2$FC (Stimulation/Mock)

+ 2 1 0 -1 -2 -

poly I:C
poly dA:dT
LPS
TNF-α
IFN-β

IRF9
GEM
IFITM3
IFI16
OAS1

DDX60
VGF
KCNJ8
CASP1
COL3A1
PTAFR

BPIFB1
BPIFA1
PXDN

SLC22A13
SLC15A2
CD226
JAK3

PAK3
IL33
PLAC8
PTPRD
SLAMF7
SYK
ITM2A

CD68
STYK1
CD40
ETS1
CXCL5

DLL1
IL1R2
SLFN13
ITGB2

CCL5
OASL
IFIT2
IFNL2
IFNL1
IFNL3
IFIT1
IRF7
IFI44L
CD274
ISG15
IFIT3
IFIH1
DDX58
MX1
IFI44
IL1A
CCL2
IL32
PI3

C

PATHWAY	RANKS	NES (p.adj < 0.05)
IFN-ALPHA/BETA SIGNALING		
IFN-GAMMA SIGNALING		
INTERFERON SIGNALING		
PEPTIDE LIGAND BINDING RECEPTORS		
DDX58-IFIH1 INDUCTION OF IFN-ALPHA/BETA		
IL-4 AND IL-13 SIGNALING		
CLASS A1 RHODOPSIN LIKE RECEPTORS		
ANTIVIRAL MECHANISM BY IFN STIMULATED GENES		
CYTOKINE SIGNALING IN IMMUNE SYSTEM		
poly I:C SIGNALING BY INTERLEUKINS		

PATHWAY	RANKS	NES
INTERFERON SIGNALING		
IFN-GAMMA SIGNALING		
ANTIVIRAL MECHANISM BY IFN STIMULATED GENES		
DDX58-IFIH1 INDUCTION OF IFN-ALPHA/BETA		
CLASS A1 RHODOPSIN LIKE RECEPTORS		
CYTOKINE SIGNALING IN IMMUNE SYSTEM		
IL-4 AND IL-13 SIGNALING		
SIGNALING BY INTERLEUKINS		
G ALPHA-I SIGNALLING EVENTS		
poly dA:dT GPCR LIGAND BINDING		

PATHWAY	RANKS	NES
INTERFERON SIGNALING		
CYTOKINE SIGNALING IN IMMUNE SYSTEM		
SIGNALING BY INTERLEUKINS		
G ALPHA-I SIGNALLING EVENTS		
GPCR LIGAND BINDING		
TLR4 CASCADE		
TOLL-LIKE RECEPTOR CASCADES		
SIGNALING BY GPCR		
INNATE IMMUNE SYSTEM		
LPS ADAPTIVE IMMUNE SYSTEM		

PATHWAY	RANKS	NES
INTERFERON SIGNALING		
CYTOKINE SIGNALING IN IMMUNE SYSTEM		
SIGNALING BY INTERLEUKINS		
G ALPHA-I SIGNALLING EVENTS		
TOLL-LIKE RECEPTOR CASCADES		
GPCR LIGAND BINDING		
C-TYPE LECTIN RECEPTORS CLRS		
EXTRACELLULAR MATRIX ORGANIZATION		
INNATE IMMUNE SYSTEM		
TNF-α ADAPTIVE IMMUNE SYSTEM		

PATHWAY	RANKS	NES
IFN-ALPHA/BETA SIGNALING		
TRAF3-DEPENDENT IRF-ACTIVATION PATHWAY		
INTERLEUKIN10 SIGNALING		
INTERFERON SIGNALING		
NICOTINAMIDE SALVAGING		
ANTIVIRAL MECHANISM BY IFN STIMULATED GENES		
INTERFERON-GAMMA SIGNALING		
IFN-β CYTOKINE SIGNALING IN IMMUNE SYSTEM		

0   8000   0   1   2

Figure 1. Stimuli activate shared and distinct pathways of the innate immune response. (A) Across 3 biological replicates, ncgRNA (grey) and PAF1 KO (teal) cells were subject to a 3-hour stimulation by the treatments described. Following, RNA was extracted, purified, and sequenced. Aligned reads underwent DESeq2 differential gene expression analysis. (B) A heatmap of log2 foldchanges (ncgRNA stimulation / ncgRNA mock) for immune response genes (GO:0002376) confirms immune activation. Hierarchal clustering using Ward's method identified notable clusters of genes for labeling. (C) GSEA was performed on DEGs in ncgRNA cells for each stimulation, wherein genes were ranked by a weighted statistic for log2 foldchange and adjusted p-value. The normalized enrichment scores (NES) for, at most, the top 10 enriched Reactome pathways were plotted for each stimulus (p.adj < 0.05). The full GSEA output is available in Table S2A.

We first analyzed the data by focusing on the ncgRNA control cells treated with various stimuli to confirm that these treatments induced immune response gene expression compared to mock-treated cells. We found each stimulus activated typical genes associated with the immune response (GO:0002376) (Fig. 1B, Table S1A). The inflammatory stimuli (TNF-α, LPS) appeared to activate some genes distinct from the interferon-related stimuli (IFN-β, poly dA:dT, poly I:C), and vice versa. These results align with principal component analysis (PCA), which shows two axes of distinct immune activation (Fig. S2A). Poly I:C was the strongest stimulus as measured by number of genes induced. The same trends are reiterated upon identifying the top differentially expressed genes (DEGs) (|log2 fold-change| > 1.5, p-value < 0.05) for each stimulus (Fig. S2B) and validating specific gene expression by qRT-PCR (Fig. S2C).

We next implemented gene set enrichment analysis (GSEA) to measure induction of the

immune response in an unbiased manner, using Reactome gene sets encompassing a

broad number of cellular pathways. With this, we confirmed all stimuli activated known

immune response pathways (Fig. 1C). Corroborating previous observations, our

inflammatory stimuli induced some unique pathways, as did the pathways activating the

interferon response. Specifically, broad GSEA revealed each stimulus as a potent driver

of interferon-related pathways, including interferon alpha-beta and gamma (Fig. 1C).

Given their mediation of the inflammatory response, TNF-α and LPS were more enriched

for pathways related to the adaptive immune response and TLR signaling cascades

compared to their counterparts (Fig. 1C, Table S2A). Overall, our stimuli are broad and

effective inducers of the immune response.

PAF1 KO results in dysregulation of immune gene expression

To discern whether PAF1-mediated gene expression is differentiated across immune

stimuli, we compared PAF1 KO cells to the control ncgRNA cells for every stimulus. We

performed PCA on immune genes while excluding batch effects and variance attributed

to differences previously found amongst the stimuli. In doing so, we found immune gene

expression in PAF1 KO cells is distinct from that of ncgRNA cells, accounting for 72% of

the remaining variance (Fig. 2A).

Figure 2. PAF1 KO cells exhibit a distinct immune gene expression profile. (A) PCA

shows distinct gene expression profile of PAF1 KO relative to ncgRNA. The counts for

immune genes (GO:0002376) were corrected with a variance-stabilizing transformation

via DESeq2, and the variability of biological replicates and immune activation were

removed as batch effects with ComBat. PCA was performed on the resulting counts, with

the top 2 principal components being plotted. Cell lines are represented by shape (circle:

ncgRNA, triangle: PAF1 KO) and stimuli are dictated by color. (B) Volcano plots show

differential gene expression of immune genes in PAF1 KO relative to ncgRNA across

stimuli. Contrasts were performed within the DESeq2 pipeline. The resulting log2

foldchanges and –log10 p-values are plotted for immune genes, with the top-10 positive

and negative log2 foldchanges being highlighted/labeled. Cutoffs are set at log2

foldchange > 1.5 or < -1.5, and p-value < 0.05.

For each comparison of PAF1 KO to ncgRNA, we identified the top immune system DEGs

(GO:0002376, |log2 foldchange| > 1.5, p-value < 0.05). Across stimuli, we found several

immune genes which are both upregulated and downregulated in PAF1 KO relative to

ncgRNA (Fig. 2B, Fig. S3). cGAS, a well-recognized cytosolic DNA sensor [24], had

significantly lower expression in PAF1 KO cells compared to ncgRNA cells for all stimuli

(Fig. 2B, Fig. S3, Table S1B). We found similar trends for the expression of

PYCARD/ASC, a key activator of inflammatory caspases for subsequent maturation of

IL-1β [25] and mediator of NF-κB related apoptosis [26]. The same was also found for

IRF8 and LRRK1, both of which are regulators of NF-κB activation for various immune

pathways [27, 28].

Similar to previously published studies indicating a negative regulatory role for PAF1, we

also identified genes whose expression increased following PAF1 KO. In general, these

genes were more variable in terms of immune modulation. Several of them are known to

repress the immune response under various cellular or pathogenic contexts—including

IL33, S1PR1, and MECOM [29–33](Fig. 2B, Fig. S3). However, other upregulated genes,

such as OAS2, PXDN, and TLR4, are predominantly mediators of immune activation and

antimicrobial defense [34–36]. TLR4 gene expression is also suppressed by ZNF160 [37],

a gene with consistent downregulation in PAF1 KO (Fig. 2B, Table S1B). Thus, some of

these effects may be secondary to a direct effect by PAF1 on the gene of interest. Overall,

the apparent upregulation and downregulation of gene expression following PAF1 KO

further reinforces the model of PAF1 suppressing some gene expression mechanisms

while likely inducing others.

Immune pathway analysis of PAF1-dependent gene expression

Next, we applied an increasingly targeted approach to identify how altered gene

expression may modulate the innate immune response. First, we chose to ascertain

whether immune induction in PAF1 KO was reduced compared to ncgRNA control cells.

When comparing stimuli against the mock-treatment for both cell lines, we found an

overall decrease in immune gene expression for PAF1 KO (Fig. 3A). Though these

changes were relatively modest, they were consistent and resulted in strong statistical

significance for each comparison. Importantly, some genes happened to be more induced

in PAF1 KO (Fig. 3A), which aligns with our previous analysis of individual DEGs (Fig.

2B). Thus, PAF1 is evidently required for robust immune gene expression regardless of

downstream effect, yet likely fills suppressive roles as well.

Figure 3. PAF1 modulates the expression of genes related to immune mechanisms

and host machinery. (A) The log2 foldchanges of immune genes (GO: 0002376)

upregulated (log2 foldchange > 2) following each stimulus in ncgRNA cells (relative to the

mock) are plotted. These genes are mapped to their counterpart log2 foldchanges in

PAF1 KO cells. A paired Wilcoxon signed rank test with Bonferroni correction was

performed to assess significant changes between the distributions of foldchanges. (B)

GSEA was performed on DEGs from PAF1 KO (relative to ncgRNA) for each stimulus,

wherein genes were ranked by a weighted statistic for log2 foldchange and adjusted p-

value. Significantly enriched Reactome pathways (p.adj < 0.05) were analyzed for overlap

amongst stimuli and the results shown on an upset plot. The number of pathways that

overlap across all stimuli is highlighted in yellow. Filled dots indicate which stimuli shared

enrichment for the given number of pathways. (C-D) GSEA random walks with the running

total enrichment score (ES) were plotted for Reactome pathways found to be negatively

(C) or positively (D) enriched across stimuli. Colors differentiate stimuli on each plot.

Numbers listed in the upper-right corner of plots indicate the size of the gene set shown.

The full GSEA output is available in Table S2B.

Seeking to validate this result, we again used GSEA to identify pathways that are PAF1-

dependent. We performed GSEA on our DEGs, evaluating pathway overlap to ascertain

which phenotypes may be conserved or differentiated across stimuli. Most significantly

enriched pathways (p.adj < 0.05) were shared across stimuli, totaling to 103 pathways

(Fig. 3B). We hypothesized that an antimicrobial effect of PAF1 could derive from

positively regulated genes that are antimicrobial, or negatively regulated genes that are

promicrobial. Therefore, we investigated these pathways for those that were

downregulated and antimicrobial, or upregulated and promicrobial. Conserved

downregulated pathways included the antiviral mechanism by IFN-stimulated genes as

well as SUMOylation (Fig. 3C). Both pathways are predisposed to contribute to a

defensive state in cells if active [38–42]. We also found negative enrichment of pathways

associated with the nuclear pore complex (Table S2B), including disassembly (Fig. 3C).

Therefore, PAF1 may facilitate innate immune defense beyond traditional pathways

alone.

Expanding on the potentially suppressive role of PAF1, we also identified upregulated

pathways in PAF1 KO. Strong positive enrichment is evident for both laminin and integrin

cell surface interactions (Fig. 3D, Table S2B), both correlated with sites prone to pathogen

adhesion and entry [43–47]. Similarly, ER to Golgi anterograde transport was also

positively enriched (Fig. 3D), a process coopted by some viruses for egress [48]. This

suggests PAF1 activates certain defensive mechanisms in tandem with suppressing

pathways prone to microbial hijacking.

To see if this trend could be validated for specific viruses, we next considered whether

PAF1 is suppressive of viral host dependency factors, as shown with flaviviruses

previously [14]. Across stimuli, GSEA revealed that *Flavivirus* host-dependency factors

were induced following PAF1 KO (Fig. 4A-B), suggesting PAF1 may be suppressive of

those factors typically. For Influenza A virus (IAV), this trend was present yet reduced

across stimuli (Fig. 4A, C). However, the Reactome pathway for NS1-mediated effects on

host pathways was significantly downregulated for all stimuli in PAF1 KO (Table S2B),

which is expected because NS1 targets PAF1 for antagonism [13]. We also assessed

host dependency factors for HIV and SARS-CoV-2 given their implication as targets of

PAF1-mediated restriction [19, 49]. Like IAV, there was a strong effect under poly I:C

stimulation for these viruses, but not the other stimuli (Fig. 4A, D-E). Hence, the extent of

PAF1 restriction by suppression of dependency factors may depend on the type and

degree of immune activation in the first place.

Figure 4. PAF1 KO shows increased expression of virus host-dependency factors.

GSEA was performed on DEGs in PAF1 KO relative to ncgRNA for each stimulation using

custom gene sets describing host-dependency factors for various viruses. Colors indicate

stimuli type and transparency indicates adjusted p-values for each test. (A) NES values

are plotted with error bars representing log2 of the expected error for the standard

deviation of the P-value logarithm. The size of each gene set is provided in parentheses.

GSEA random walks as summarized in (A) are shown for *Flaviviruses* (B), Influenza A

Virus (C), HIV (D), and SARS-CoV-2 (E).

Considering these deviations in PAF1-dependency across stimuli, we hypothesized there

could be stimulus-specific gene expression that was also PAF1-dependent. Many of

these stimulus-specific genes were attributed to different degrees of immune activation

relative to the mock. Yet, we still identified some immune genes which were consistently

activated across stimuli in ncgRNA cells but were stimulus dependent in PAF1 KO (Fig.

S4A-B). For example, REC8 was significantly downregulated under nucleic acid mimic

stimuli (poly I:C, poly dA:dT) in PAF1 KO cells, but not for inflammatory stimuli (LPS, TNF-

α) (Fig. S4A). Interestingly, REC8 promotes innate immune signaling via stabilization of

MAVS and STING [50], connecting back to our previously identified dysregulation of the

cGAS-STING pathway (Fig. 2B). We also compared upstream PAMPs (poly I:C, poly

dA:dT, LPS) to downstream activators (TNF-α, IFN-β), but found few genes strongly

induced across stimuli in ncgRNA cells, yet differentially expressed across stimuli in PAF1

KO (Fig. S4B). Moreover, none were strongly associated with an immune response. Thus,

though PAF1C can function in a stimulus-dependent manner [51–53], this does not

appear to be the central driving factor for innate immune modulation.

Predicted transcriptional motifs underlying PAF1-dependent immune regulation

Since most of the holistic trends we identified were shared across stimuli, we sought to understand how this may translate to consistency across PAF1-dependent expression for individual genes. To that end, we queried all significant DEGs (p.adj < 0.05) from our individual comparisons and aggregated them to identify overlaps. There were also numerous genes associated with only some stimuli but not others, reinforcing that other inputs also contribute to the expression of PAF1-dependent genes. Despite this variability, we discovered 50 downregulated genes and 104 upregulated genes shared across all stimuli (Fig. 5A).

Figure 5. Promoters of conserved PAF1-dependent genes are associated with immune-related transcription factors. (A) Venn diagrams show overlapping DEGs (p.adj < 0.05) from comparison of PAF1 KO versus ncgRNA for each stimulus. The diagram on the left shows downregulated genes and the one to the right shows upregulated genes. Overlaps having greater than 10 genes were labeled. (B-C) Core genes (marked at the center of Venn diagrams) were subjected to TFBS enrichment analysis with DAVID. The fold enrichment (FE) of the top 15 associated TFBS for downregulated (B) and upregulated (C) genes are shown. Emphasis (larger size and labeling) was given to those with greater than 0.5 difference in FE compared to the

opposing gene list (downregulated versus upregulated (B), and vice versa (C)). Statistical

significance for enrichment was calculated with the Bonferroni method and is shown by

shape (circle: p.adj < 0.05, triangle: p.adj > 0.05). TFBS associated with the immune

response and significantly enriched have bolded and colored labels.

We performed gene ontology (GO) enrichment analyses on these core genes, but we did

not identify any sets that were significantly enriched. PAF1C is known to interact with

several transcription factors during the recruitment of RNAPII [7, 51, 54, 55]. Thus, we

extended our approach to focus on regulatory elements associated with core genes.

Taking the downregulated genes conserved across stimuli, we identified significant

enrichment for specific transcription factor binding sites (TFBS) associated with the

promoters of those genes (Fig. S5A). Many of the top hits were implicated in the innate

immune response, including c-Rel, ATF2/c-Jun, RELA/NF-κB p65, IRF1, AP-1, and

EGR3 (Fig. 5B, Table S3A). Supporting the model in which PAF1 is an activator at these

sites, there was comparatively less enrichment amongst upregulated genes for these

same TFBS. Specifically, c-Rel, ATF2/c-Jun, and EGR3 all displayed greater than 2-fold

enrichment for downregulated genes. However, their upregulated-associated

counterparts were approximately less than or equal to 1-fold enrichment (no enrichment)

(Fig. 5B, Table S3A). This same analytical approach was performed to identify the top

TFBS associations for upregulated genes. Yet, this resulted in enrichments of smaller

magnitude and less often associated with immune-related TFBS (Fig. 5C). What's more,

these top associations were often outperformed by their downregulated-associated

counterparts. Taken together, PAF1-dependent genes have promoters associated with

known transcription factors, including those involved with regulation of innate immunity.

We next investigated enhancers associated with our genes because PAF1 could occupy

these sites for activation or suppression [8, 9, 11, 12]. We extracted all the genomic

regions of enhancers associated with our core genes (Fig. S5A, Table S5B). Mapping the

identified sites onto the human genome, we evaluated if PAF1-associated enhancer

occupancy was distinct between upregulated and downregulated genes. Strikingly,

upregulated and downregulated genes shared little overlap between their respective

enhancers (Fig. 6A). When this was tested with a Jaccard statistic, we could only find a

single region of overlap, resulting in an extremely low score of 5.55×10^{-5}. Our

downregulated genes had enhancers that were more widely distributed than the

upregulated genes (Fig. S5B). Additionally, for the number of genes in between the

enhancer and the gene of interest, upregulated-associated enhancers were found to be

closer on average (two-tailed T-test, p value < 0.001) (Fig. S5C).

Figure 6. Predicted enhancers associated with PAF1-dependency and the innate

immune response. (A) Enhancers associated with the core PAF1-dependent

upregulated (orange) and down-regulated (magenta) genes are displayed on the human

genome. Enhancer chromosomal regions were extracted and plotted in 0.5 Mb bins. The

extent these regions overlapped was calculated with a Jaccard statistic. (B) Volcano plots

show results of relative enrichment analysis on extracted enhancers. Significantly

enriched motifs (E < 0.05, enrichment ratio > 2) are colored. The top 10 enrichment ratios

for known motifs are highlighted for each set. (C) Venn diagram shows overlap of enriched

motifs associated with upregulated and downregulated genes. (D) Manhattan plot

displays significant (p.adj < 0.05) motifs associated with GO-BP immune response terms

(GO: 0006955). Motifs were submitted to GOMo and enriched for correlated terms,

correcting for false-discovery rate with Benjamini-Hochberg method. Color indicates

whether the enriched motif was associated with downregulated (magenta) or upregulated

genes (orange) alone, or if it was shared between them (grey). The analytical pipeline for

these analyses is further clarified in Figure S3A.

Although the genomic regions differed, we also considered whether motifs in these

enhancers differed as well. For both upregulated and downregulated genes, we

discovered several significantly enriched motifs (p.adj < 0.05, log2 fold enrichment > 1)

(Fig. 6B-C, Table S3C). Primarily, top motifs for both groups were often assorted zinc-

finger binding motifs or those associated with thyroid hormone receptor-related factors

(Fig. 6B). Approximately one third of these motifs were shared between upregulated and

downregulated genes (Fig. 6C), suggesting these motifs are not necessarily driving the

direction of PAF1 regulation for upregulated and downregulated genes. Importantly, there

were unique enhancer motifs for both gene sets, so PAF1's differential gene regulation

may be due to these motifs.

To extend this predictive model, we evaluated all significantly enriched motifs for

enrichment of GO biological processes (GO-BP) related to the innate immune response

(GO: 0006955) (Fig. S5A). Implementing GOMo (Gene Ontology for Motifs), we evaluated

whether potential gene targets associated with our enhancer regions were enriched for

GO-BPs. We discovered a subset of our motifs were significantly enriched (p.adj < 0.05)

for immune-related terms, including innate immunity and the defense response to virus

(Fig. 6D, S3D Table). Contrary to our TFBS promoter analysis (Fig. 5B), the enhancers

associated with upregulated genes comprised most of the motifs related to the immune

response (Fig. 6D). Although motifs associated with downregulated genes or with both

sets were less often enriched for GO-BPs, a large fraction still had known immune gene

targets (Fig. S6). This illustrates how PAF1 activity at different regulatory elements could

be highly variable in regard to innate immunity.

Discussion

Expanding the cellular role of PAF1C, here we explore the extent of PAF1-dependent

gene expression as it relates to the immune response. Not only do we reinforce results

from previous studies, but we delineate potential mechanisms that determine and

contribute to the degree and type of PAF1-mediation. We show that PAF1 promotes gene

expression across a broad range of genes in response to diverse immune stimuli. We

also provide additional support that PAF1 can negatively regulate some genes, including

proviral host factors. We show these trends in gene expression correlate with known

immune response motifs at the promoter and enhancer level. With this work, we refine

the role of PAF1C in responding to microbial stress and identify potential immune

regulatory networks to be investigated in subsequent studies.

Our work also contributes to reconciling the recently described, divergent functions of

PAF1C. From one perspective, PAF1C allows the proximal release of a paused RNAPII

from genes, resulting in activation [8]. Yet, the complex can also be a negative regulator

of this process [9], maintaining the paused status of RNAPII, especially at super-

enhancers in cancer cell lines [11]. Recent work has combined these models by

suggesting loss of PAF1C changes the distribution and rates of RNAPII transcription [12],

delineating between proximal and distal genomic sites. Building on these models, we

show that PAF1 activation is indeed correlated with promoter elements while suppression

is more often associated with enhancer sites. Experimental validation of these trends in

future work will be pivotal to fully integrating the two models of PAF1C.

At the level of immune gene expression, PAF1C-mediated activation is apparent and

aligns with past work. The diminished expression of cGAS and PYCARD/ASC across all

stimuli (Fig. 2B), for example, mirrors our group's past work in finding cGAS expression

to be reduced following PAF1 knockout in poly I:C-stimulated cells [14]. Considering both

cGAS and ASC are involved in dsDNA-mediated immune activation and exhibit cross-

regulation [56], it is particularly interesting to see their shared downregulation. Also, we

detected consistently increased expression of IFI16 in PAF1 KO (Table S1B), which can

function cooperatively with cGAS [57], suggesting its overexpression may reflect

additional dysregulation of this system. Given the cGAS-STING pathway and formation

of the AIM2 inflammasome are activated by diverse pathogens regardless of dsDNA

genome status [58, 59], PAF1 mediation of their expression could be protective against

cellular vulnerability. This is confirmed when PAF1 KO displays a loss in gene expression

for the antiviral mechanism by IFN-stimulated genes (Fig. 3).

Though loss of canonical pathways for immune defense is evident, we found additional

cellular pathways significantly affected by PAF1 and still linked to immune defense,

though less directly. Decreased expression of SUMOylation pathways in our PAF1 KO

cells (Fig. 3B) is especially interesting considering all PAF1C members are a target of

SUMO to activate the interferon response and counteract IAV [60]. This suggests PAF1C

may rely on a potential positive feedback mechanism to increase its mediation of an

antimicrobial response. We also highlight decreased expression of genes related to

nuclear pore complex maintenance, for their expression has been correlated with immune

effector import into the nucleus [61–63]. For this reason, nuclear pore complexes are

often targeted by various viruses for degradation [64–68]. Since PAF1 likely facilitates

immune gene expression reliant on the translocation of cytoplasmic factors into the

nucleus, maintaining nuclear pore complexes could support innate immunity. Then again,

restructuring of these complexes could also benefit viruses that rely on the nucleus for

genome replication and immune evasion [69, 70]. Establishing how these trends in gene

expression translate to cellular outcomes will require knockout or knockdown of factors

related to these pathways and subsequent validation.

Beyond this, our prediction of transcription factors at promoter elements further validates

PAF1C as an activator at promoter-proximal sites. Downregulated genes were heavily

associated with immune response promoter motifs, including NF-κB components like c-

Rel and p65/RelA and coregulator c-Jun/AP-1 (Fig. 5B). This may explain why PAF1-

dependent genes were often correlated with the immune response through NF-κB, which

plays multiple roles in cytokine expression [71–73]. Moreover, these same promoter-

associated motifs were far less enriched in our upregulated genes. Since we associate

downregulated genes with PAF1-mediated activation, this result suggests PAF1C triggers

this immune response via proximal release of RNAPII. As previously stated, the complex

has already been connected to activation at proximal sites [8], so our work fits well into

this model of PAF1-dependent gene expression.

Our findings for upregulated immune genes in PAF1 KO are more complex, but they still

recapitulate a suppressive role for PAF1C. Though there were some genes found to be

immune activators, several others were poised to act in immunosuppressive roles (Fig.

2B). Additionally, PAF1 was found to be suppressive for proviral mechanisms often

subject to viral hijacking (Fig. 3C). As such, loss or antagonism of PAF1 may abrogate

the immune response by RNAPII release from certain transcription sites. As we had

previously shown [14], this could explain why we find positive enrichment for flavivirus

host-dependency factors in PAF1 KO (Fig. 4). This effect was diminished for other viruses

across weaker immune stimuli. Hence, PAF1 could be required for maintaining a proviral

microenvironment for some viruses, but perhaps this only occurs when certain immune

pathways are activated during infection.

When we analyzed enhancer motifs associated with our PAF1-dependent genes, our

results indicated PAF1C-mediation at these sites may be variable, depending on several

factors. For one, a large fraction of enriched enhancer motifs was found to be shared

between upregulated and downregulated genes (Fig. 6C). This occurs despite them

occupying exclusive genomic regions, suggesting genomic location itself is a stronger

determinant of PAF1 functionality at certain sites. Since PAF1C extensively regulates

chromatin modification and remodeling [74–78], these roles could be driving such a trend.

However, we found a large subset of enhancer motifs exclusively associated with

upregulated genes. Several of them were immune-related (Fig. 6), suggesting PAF1

represses certain immune genes through enhancer occupancy. Enhancers associated

with upregulated genes also had a tighter genomic distribution and were typically closer

to expressed genes compared to those downregulated associated (Fig. S4). This could

imply that upregulated genes are regulated by super-enhancers, which are often

considered to be motifs in close genetic proximity [79]. Given PAF1C occupancy of super-

enhancers has been associated with suppression previously [11], our results support

PAF1C restricting gene expression by pausing RNAPII at such sites.

This begs the question as to why we observe enrichment for enhancers associated with

downregulated genes and those shared by both gene sets. When we map all motifs to

their respective target genes, there is a strong overlap between upregulated and

downregulated immune gene targets (Fig. S6). Thus, PAF1C likely induces positive

regulation of innate immunity by occupying promoters, in addition to some enhancers.

Meanwhile, the complex could further bolster immunity by repressing immunosuppressive

genes via enhancer occupancy or activating negative regulators upstream. It suggests a

sort of "tug-of-war" between PAF1C acting at proximal versus distal regulatory elements.

With this complexity in mind, PAF1C may shift between predominantly active or

suppressive roles depending on the context of cell perturbation. This harkens back to our

analysis of host-dependency factors, wherein the type of stimuli contributed to the extent

of PAF1-mediated suppression. Regardless of the exact mechanisms, our results

highlight how models of PAF1C as either an activator or suppressor are not inherently

conflicting but may exhibit cross-regulation in the context of innate immunity. It should be

noted that PAF1C-mediated suppression of genes has largely been observed in cancer

cell lines [9, 11]. Studying if these dual regulatory modes of PAF1C exist in primary

immune cells and other epithelial cell lines will be valuable.

These results provide the foundation to further investigate PAF1C occupancy of

transcriptional sites in the context of innate immunity. Identifying such sites with ATAC-

Seq [80] and ChIP-Seq [81] in PAF1 KO cells would reveal whether our predicted motifs

are transcriptionally accessible during the immune response. Also, corroborating our

results by studying PAF1C overexpression and monitoring for inverted effects would

further strengthen the trends identified here. Strikingly, a concurrent study has already

validated aspects of our computational predictions, showing PAF1 to be supportive of NF-

κB-dependent gene expression in a siRNA knockdown model [16]. Such work reiterates

how our contribution is a critical starting point for future studies focusing on PAF1C and

the immune response.

Conclusions

In summary, we clarify the extent of PAF1C-mediated gene expression and identify

notable trends that reaffirm its roles in both activation and suppression of immune gene

expression. We show strong correlations between gene expression and known

transcriptional motifs. This provides basis for future elucidation of molecular and cellular

mechanisms underlying PAF1C function during an immune response. This is especially

important since PAF1C can be coopted through host-pathogen interactions to disrupt
innate immunity.

Materials and Methods

Cells. ncgRNA and PAF1 KO cell lines were generated by CRISPR/Cas9 in A549 lung
epithelial cells as previously described [14]. All cell lines were cultured in Dulbecco's
modified Eagle's medium (DMEM, Gibco ThermoFisher) with 10 % fetal bovine serum
(FBS, Gibco ThermoFisher) at 37ºC. *Mycoplasma spp.* PCR testing was done monthly.

Sample preparation. ncgRNA and PAF1 KO cells were plated at a density of 6.0×10^5
cells per well in 6-well plates and incubated at 37ºC for 24 hours. Cells were then
stimulated with poly I:C (2.0 µg/mL, Tocris), poly dA:dT (1.0 µg/mL, InvivoGen), LPS (1.0
µg/mL, eBioscience), IFN-β (50 U/mL, R&D Systems), and TNF-α (100 ng/mL, Gibco
Thermofisher) at the given concentrations. For poly I:C and poly dA:dT, transfection was
performed with Lipofectamine 2000 (ThermoFisher) per manufacturer instruction. After 3
hours post-stimulation, all samples were collected with Trizol reagent (Ambion) and
stored at -80ºC. Total cellular RNA was extracted and purified with the RNeasy mini kit
(Qiagen). At the sequencing facility, RNA samples were prepared with the QuantSeq3'
mRNA-Seq Library kit for Illumina (Lexogen GmbH), following a standard protocol [82].
The resulting cDNA libraries were sequenced on an Illumina NextSeq500 system with
approximately 4 million reads per sample.

Quantitative reverse transcription-PCR (qRT-PCR). Following RNA extraction, sample cDNA was synthesized with the iScript kit (BioRad). Quantitative real time PCR was performed with iTaq SYBR Green premix (BioRad) in 96-well plate format, with data collection and analysis conducted on a LightCycler 480 (Roche). All Ct values were normalized against GAPDH expression. Quantification of gene expression was calculated by the ΔΔCt method, as previously described [83]. Primer sequences are provided in Table S4.

Immunofluorescence microscopy. Parental A549 cells were fixed with 4% paraformaldehyde following 3-hour treatment with poly I:C (2.0 µg/mL, Tocris), LPS (1.0 µg/mL, eBioscience), and IFN-β (50 U/mL, R&D Systems) at the given concentrations. Cells were permeabilized with 0.25% Triton X-100 and subsequently blocked with 5% goat serum in PBS. Samples were incubated in primary and secondary antibodies for 1 hour each at 25⁰C. Cell nuclei were stained with Hoechst (Invitrogen). PAF1 primary antibody was diluted at 1:500 (Atlas antibodies, HPA043637). AlexaFluor555 and Hoechst33342, both at 1:1000 dilution, detected PAF1 and Hoechst, respectively. Epifluorescence microscopy images were produced with a Nikon ECLIPSE Ti2 Series microscope using 40x objective. StarDist (Weigert 2018) was used to detect nuclei of cells for measurement of PAF1 fluorescent intensity. All other measurements were performed within Fiji interface (Schindelin 2012).

Differential gene expression analysis. Read data was evaluated for sequence quality and sample contamination with FastQC (v0.11.9) [84]. Reads were aligned against the human reference genome (build 38) with STAR aligner (v2.5.2b) [85]. FeatureCounts (v2.0.3) calculated read counts for each gene and built a corresponding matrix for all

samples [86]. Prior to differential gene expression (DGE) analysis, genes with mean counts less than 1 were excluded. DGE analysis was performed with DESeq2 (v1.34.0) [87], wherein the design accounted for the biological replicate number, type of immune stimuli, and cell type (PAF1 KO, ncgRNA). Direct contrasts between different conditions and cell types were performed after normalization of read counts. Batch effects were removed with ComBat function of the sva package (v3.42.0) [88]. Significant changes in specific genes (p.adj < 0.05) were identified after adjusting for false discovery rate using the Benjamini-Hochberg method.

Gene set enrichment analysis. Gene-set enrichment analysis (GSEA) was performed with fgsea [89], using Reactome gene sets [90] downloaded from the Molecular Signatures Database [91]. Custom GSEA gene sets for were curated manually from publicly available data and are listed in Table S5. All genes were included in the random walk and ranked by a DESeq2 weighted statistic (stat) for fold change and p-value. Each run performed 10,000 iterations across gene sets, which included sets of 10 to 500 genes for targeted assessment and 50 to 1000 genes for broad assessment. Significant changes in specific gene sets (p.adj < 0.05) were identified after adjusting for false discovery rate using the Benjamini-Hochberg method. Normalized enrichment scores (NES) represent the enrichment score (ES) following normalization to the mean enrichment of random samples of the same gene set size.

Regulatory elements analysis. The Database for Annotation, Visualization and Integrated Discovery (DAVID) (Knowledgebase v2021q4) [92] was used to assess for statistical enrichment of promoter-associated transcription factor binding sites (TFBS). This was done according to the UCSC Genome Browser database of TFBS conserved

tracks, which are experimentally-determined sites [93]. DEGs were inputted and enriched

for TFBS with EASE scoring, which produces a modified Fisher's Exact test p-value [94],

correcting for false discovery rate using the Bonferroni method.

Enhancers of DEGs were aggregated with GeneALaCart/GeneHancer as synchronized

with GeneCards (v5.7) [95]. Mapping to human genome was done with chromPlot

(v1.22.0) [96] and a Jaccard overlap statistic for upregulated-associated versus

downregulated-associated enhancers was calculated with the valr package (v0.6.4)[97].

Extracted enhancer regions were filtered as "Enhancer" and "Enhancer/Promoter" and

met the criteria of existing in A549 lung epithelial cells and having a GeneCards

confidence score > 10 before enrichment. Then, MEME Suite (v4.5.1) program Simple

Enrichment Analysis (SEA) [98] identified relatively enriched motifs in the enhancer

regions of DEGs. Gene Ontology for Motifs (GOMo) [99] identified which of these

enhancer motifs were associated with GO-BP terms. Both MEME Suite programs

corrected for false discovery rates using the Benjamini-Hochberg method. Regulatory

relationships between transcription factors and target genes were extracted from the

Transcriptional Regulatory Relationships Unraveled by Sentence-based Text Mining

(TRRUST, v2) database [100].

Declarations

Ethics approval and consent to participate

Not applicable

Consent for publication

Not applicable

Availability of data and materials

RNA-sequencing data are available in the ArrayExpress database under accession number E-MTAB-11620. The Sequence Read Archive (SRA) accession number is PRJEB52223. The basic code for the read alignment, differential gene expression analysis, and gene set enrichment analysis is available at:

https://github.com/mwkenast/PAF1KO

Competing interests

The authors declare that they have no competing interests.

Funding

Funding was provided by the University of California, Davis and the W. M. Keck Foundation.

Authors' contributions

MWK and PSS conceived of the work and designed experiments. MWK, OHP, and MJP performed experiments. MWK conducted bioinformatic analyses. MWK and PSS wrote the manuscript. PSS secured funding for work. All authors read and approved the final manuscript.

Acknowledgements

602   We would like to thank members of the Shah Lab for constructive criticism and generative

603   discussions. Sequencing was carried out at the DNA Technologies and Expression

604   Analysis Cores at the UC Davis Genome Center, supported by NIH Shared

605   Instrumentation Grant 1S10OD010786-01.

References

1. Mueller CL, Jaehning JA. Ctr9, Rtf1, and Leo1 are components of the Paf1/RNA polymerase II complex. Mol Cell Biol. 2002;22:1971–80.

2. Shi X, Finkelstein A, Wolf AJ, Wade PA, Burton ZF, Jaehning JA. Paf1p, an RNA polymerase II-associated factor in Saccharomyces cerevisiae, may have both positive and negative roles in transcription. Mol Cell Biol. 1996;16:669–76.

3. Costa PJ, Arndt KM. Synthetic lethal interactions suggest a role for the Saccharomyces cerevisiae Rtf1 protein in transcription elongation. Genetics. 2000;156:535–47.

4. Betz JL, Chang M, Washburn TM, Porter SE, Mueller CL, Jaehning JA. Phenotypic analysis of Paf1/RNA polymerase II complex mutations reveals connections to cell cycle regulation, protein synthesis, and lipid and nucleic acid metabolism. Mol Genet Genomics MGG. 2002;268:272–85.

5. Squazzo SL, Costa PJ, Lindstrom DL, Kumer KE, Simic R, Jennings JL, et al. The Paf1 complex physically and functionally associates with transcription elongation factors in vivo. EMBO J. 2002;21:1764–74.

6. Rondón AG, Gallardo M, García-Rubio M, Aguilera A. Molecular evidence indicating that the yeast PAF complex is required for transcription elongation. EMBO Rep. 2004;5:47–53.

7. Francette AM, Tripplehorn SA, Arndt KM. The Paf1 Complex: A Keystone of Nuclear Regulation Operating at the Interface of Transcription and Chromatin. J Mol Biol. 2021;433:166979.

8. Yu M, Yang W, Ni T, Tang Z, Nakadai T, Zhu J, et al. RNA polymerase II-associated

factor 1 regulates the release and phosphorylation of paused RNA polymerase II.

Science. 2015;350:1383–6.

9. Chen FX, Woodfin AR, Gardini A, Rickels RA, Marshall SA, Smith ER, et al. PAF1, a

Molecular Regulator of Promoter-Proximal Pausing by RNA Polymerase II. Cell.

2015;162:1003–15.

10. van den Heuvel D, Spruijt CG, González-Prieto R, Kragten A, Paulsen MT, Zhou D,

et al. A CSB-PAF1C axis restores processive transcription elongation after DNA

damage repair. Nat Commun. 2021;12:1342.

11. Chen FX, Xie P, Collings CK, Cao K, Aoi Y, Marshall SA, et al. PAF1 regulation of

promoter-proximal pause release via enhancer activation. Science. 2017;357:1294–8.

12. Hou L, Wang Y, Liu Y, Zhang N, Shamovsky I, Nudler E, et al. Paf1C regulates RNA

polymerase II progression by modulating elongation rate. Proc Natl Acad Sci.

2019;116:14583–92.

13. Marazzi I, Ho JSY, Kim J, Manicassamy B, Dewell S, Albrecht RA, et al.

Suppression of the antiviral response by an influenza histone mimic. Nature.

2012;483:428–33.

14. Petit MJ, Kenaston MW, Pham OH, Nagainis AA, Fishburn AT, Shah PS. Nuclear

dengue virus NS5 antagonizes expression of PAF1-dependent immune response

genes. PLOS Pathog. 2021;17:e1010100.

15. Shah PS, Link N, Jang GM, Sharp PP, Zhu T, Swaney DL, et al. Comparative

Flavivirus-Host Protein Interaction Mapping Reveals Mechanisms of Dengue and Zika

Virus Pathogenesis. Cell. 2018;175:1931-1945.e18.

16. Fischer NL, Boyer MA, Bradley WP, Spruce LA, Fazelinia H, Shin S. A Coxiella

burnetii effector interacts with the host PAF1 complex and suppresses the innate

immune response. 2022;:2022.04.20.488957.

17. Abdel-Mohsen M, Wang C, Strain MC, Lada SM, Deng X, Cockerham LR, et al.

Select host restriction factors are associated with HIV persistence during antiretroviral

therapy. AIDS Lond Engl. 2015;29:411–20.

18. Raposo RAS, Abdel-Mohsen M, Bilska M, Montefiori DC, Nixon DF, Pillai SK.

Effects of Cellular Activation on Anti-HIV-1 Restriction Factor Expression Profile in

Primary Cells. J Virol. 2013;87:11924–9.

19. Liu L, Oliveira NM, Cheney KM, Pade C, Dreja H, Bergin A-MH, et al. A whole

genome screen for HIV restriction factors. Retrovirology. 2011;8:94.

20. Parnas O, Jovanovic M, Eisenhaure TM, Herbst RH, Dixit A, Ye CJ, et al. A

Genome-wide CRISPR Screen in Primary Immune Cells to Dissect Regulatory

Networks. Cell. 2015;162:675–86.

21. Geiss GK, Salvatore M, Tumpey TM, Carter VS, Wang X, Basler CF, et al. Cellular

transcriptional profiling in influenza A virus-infected lung epithelial cells: The role of the

nonstructural NS1 protein in the evasion of the host innate defense and its potential

contribution to pandemic influenza. Proc Natl Acad Sci. 2002;99:10736–41.

22. Muñoz-Jordán JL, Sánchez-Burgos GG, Laurent-Rolle M, García-Sastre A.

Inhibition of interferon signaling by dengue virus. Proc Natl Acad Sci. 2003;100:14333–

8.

23. Yamada T, Sato S, Sotoyama Y, Orba Y, Sawa H, Yamauchi H, et al. RIG-I triggers

a signaling-abortive anti-SARS-CoV-2 defense in human lung cells. Nat Immunol.

2021;22:820–8.

24. Hopfner K-P, Hornung V. Molecular mechanisms and cellular functions of cGAS–

STING signalling. Nat Rev Mol Cell Biol. 2020;21:501–21.

25. Martinon F, Burns K, Tschopp J. The inflammasome: a molecular platform triggering

activation of inflammatory caspases and processing of proIL-beta. Mol Cell.

2002;10:417–26.

26. Masumoto J, Dowds TA, Schaner P, Chen FF, Ogura Y, Li M, et al. ASC is an

activating adaptor for NF-kappa B and caspase-8-dependent apoptosis. Biochem

Biophys Res Commun. 2003;303:69–73.

27. Simon PS, Sharman SK, Lu C, Yang D, Paschall AV, Tulachan SS, et al. The NF-κB

p65 and p50 homodimer cooperate with IRF8 to activate iNOS transcription. BMC

Cancer. 2015;15:770.

28. Morimoto K, Baba Y, Shinohara H, Kang S, Nojima S, Kimura T, et al. LRRK1 is

critical in the regulation of B-cell responses and CARMA1-dependent NF-κB activation.

Sci Rep. 2016;6:25738.

29. Buonamici S, Li D, Mikhail FM, Sassano A, Platanias LC, Colamonici O, et al. EVI1

abrogates interferon-alpha response by selectively blocking PML induction. J Biol

Chem. 2005;280:428–36.

30. Liang Y, Yi P, Yuan DMK, Jie Z, Kwota Z, Soong L, et al. IL-33 induces

immunosuppressive neutrophils via a type 2 innate lymphoid cell/IL-13/STAT6 axis and

protects the liver against injury in LCMV infection-induced viral hepatitis. Cell Mol

Immunol. 2019;16:126–37.

31. Nascimento DC, Melo PH, Piñeros AR, Ferreira RG, Colón DF, Donate PB, et al. IL-

33 contributes to sepsis-induced long-term immunosuppression by expanding the

regulatory T cell population. Nat Commun. 2017;8:14919.

32. Priceman SJ, Shen S, Wang L, Deng J, Yue C, Kujawski M, et al. S1PR1 Is Crucial

for Accumulation of Regulatory T Cells in Tumors via STAT3. Cell Rep. 2014;6:992–9.

33. Teijaro JR, Studer S, Leaf N, Kiosses WB, Nguyen N, Matsuki K, et al. S1PR1-

mediated IFNAR1 degradation modulates plasmacytoid dendritic cell interferon-α

autoamplification. Proc Natl Acad Sci U S A. 2016;113:1351–6.

34. Mozzi A, Pontremoli C, Forni D, Clerici M, Pozzoli U, Bresolin N, et al. OASes and

STING: adaptive evolution in concert. Genome Biol Evol. 2015;7:1016–32.

35. Shi R, Cao Z, Li H, Graw J, Zhang G, Thannickal VJ, et al. Peroxidasin contributes

to lung host defense by direct binding and killing of gram-negative bacteria. PLoS

Pathog. 2018;14:e1007026.

36. Kawai T, Akira S. The role of pattern-recognition receptors in innate immunity:

update on Toll-like receptors. Nat Immunol. 2010;11:373–84.

37. Takahashi K, Sugi Y, Hosono A, Kaminogawa S. Epigenetic regulation of TLR4

gene expression in intestinal epithelial cells for the maintenance of intestinal

homeostasis. J Immunol Baltim Md 1950. 2009;183:6522–9.

38. McNab F, Mayer-Barber K, Sher A, Wack A, O'Garra A. Type I interferons in

infectious disease. Nat Rev Immunol. 2015;15:87–103.

39. Cohen S, Au S, Panté N. How viruses access the nucleus. Biochim Biophys Acta

BBA - Mol Cell Res. 2011;1813:1634–45.

40. Ding X, Wang A, Ma X, Demarque M, Jin W, Xin H, et al. Protein SUMOylation Is

Required for Regulatory T Cell Expansion and Function. Cell Rep. 2016;16:1055–66.

41. Garaude J, Farrás R, Bossis G, Charni S, Piechaczyk M, Hipskind RA, et al.

SUMOylation regulates the transcriptional activity of JunB in T lymphocytes. J Immunol

Baltim Md 1950. 2008;180:5983–90.

42. Hannoun Z, Maarifi G, Chelbi-Alix MK. The implication of SUMO in intrinsic and

innate immunity. Cytokine Growth Factor Rev. 2016;29:3–16.

43. Singh B, Fleury C, Jalalvand F, Riesbeck K. Human pathogens utilize host

extracellular matrix proteins laminin and collagen for adhesion and invasion of the host.

FEMS Microbiol Rev. 2012;36:1122–80.

44. Paulsson M, Riesbeck K. How bacteria hack the matrix and dodge the bullets of

immunity. Eur Respir Rev. 2018;27.

45. Ulanova M, Gravelle S, Barnes R. The role of epithelial integrin receptors in

recognition of pulmonary pathogens. J Innate Immun. 2009;1:4–17.

46. Isberg RR, Tran Van Nhieu G. Binding and internalization of microorganisms by

integrin receptors. Trends Microbiol. 1994;2:10–4.

47. Feire AL, Koss H, Compton T. Cellular integrins function as entry receptors for

human cytomegalovirus via a highly conserved disintegrin-like domain. Proc Natl Acad

Sci. 2004;101:15470–5.

48. Ravindran MS, Bagchi P, Cunningham CN, Tsai B. Opportunistic intruders: how

viruses orchestrate ER functions to infect cells. Nat Rev Microbiol. 2016;14:407–20.

49. Vishnubalaji R, Shaath H, Alajez NM. Protein Coding and Long Noncoding RNA

(lncRNA) Transcriptional Landscape in SARS-CoV-2 Infected Bronchial Epithelial Cells

Highlight a Role for Interferon and Inflammatory Response. Genes. 2020;11:760.

50. Chen S, Liu Q, Zhang L, Ma J, Xue B, Li H, et al. The role of REC8 in the innate

immune response to viral infection. J Virol. 2022. https://doi.org/10.1128/jvi.02175-21.

51. Kim W, Bennett EJ, Huttlin EL, Guo A, Li J, Possemato A, et al. Systematic and

Quantitative Assessment of the Ubiquitin-Modified Proteome. Mol Cell. 2011;44:325–40.

52. Wu J, Xu W. Histone H3R17me2a mark recruits human RNA Polymerase-

Associated Factor 1 Complex to activate transcription. Proc Natl Acad Sci.

2012;109:5675–80.

53. Zeng H, Xu W. Ctr9, a key subunit of PAFc, affects global estrogen signaling and

drives ERα-positive breast tumorigenesis. Genes Dev. 2015;29:2153–67.

54. Jaenicke LA, Eyss B von, Carstensen A, Wolf E, Xu W, Greifenberg AK, et al.

Ubiquitin-Dependent Turnover of MYC Antagonizes MYC/PAF1C Complex

Accumulation to Drive Transcriptional Elongation. Mol Cell. 2016;61:54–67.

55. He N, Chan CK, Sobhian B, Chou S, Xue Y, Liu M, et al. Human Polymerase-

Associated Factor complex (PAFc) connects the Super Elongation Complex (SEC) to

RNA polymerase II on chromatin. Proc Natl Acad Sci. 2011;108:E636–45.

56. Wan D, Jiang W, Hao J. Research Advances in How the cGAS-STING Pathway

Controls the Cellular Inflammatory Response. Front Immunol. 2020;11.

57. Almine JF, O'Hare CAJ, Dunphy G, Haga IR, Naik RJ, Atrih A, et al. IFI16 and

cGAS cooperate in the activation of STING during DNA sensing in human

keratinocytes. Nat Commun. 2017;8:14392.

58. Cheng Z, Dai T, He X, Zhang Z, Xie F, Wang S, et al. The interactions between

cGAS-STING pathway and pathogens. Signal Transduct Target Ther. 2020;5:91.

59. Rathinam VAK, Jiang Z, Waggoner SN, Sharma S, Cole LE, Waggoner L, et al. The

AIM2 inflammasome is essential for host defense against cytosolic bacteria and DNA

viruses. Nat Immunol. 2010;11:395–402.

60. Domingues P, Golebiowski F, Tatham MH, Lopes AM, Taggart A, Hay RT, et al.

Global Reprogramming of Host SUMOylation during Influenza Virus Infection. Cell Rep.

2015;13:1467–80.

61. Fagerlund R, Kinnunen L, Köhler M, Julkunen I, Melén K. NF-{kappa}B is

transported into the nucleus by importin {alpha}3 and importin {alpha}4. J Biol Chem.

2005;280:15942–51.

62. Yi S, Chen Y, Wen L, Yang L, Cui G. Downregulation of nucleoporin 88 and 214

induced by oridonin may protect   OCIM2 acute erythroleukemia cells from

apoptosis through regulation of nucleocytoplasmic   transport of NF-κB. Int J Mol

Med. 2012;30:877–83.

63. Meyer T, Vinkemeier U. Nucleocytoplasmic shuttling of STAT transcription factors.

Eur J Biochem. 2004;271:4606–12.

64. Gustin KE, Sarnow P. Effects of poliovirus infection on nucleo-cytoplasmic

trafficking and nuclear pore complex composition. EMBO J. 2001;20:240–9.

65. Park N, Katikaneni P, Skern T, Gustin KE. Differential Targeting of Nuclear Pore

Complex Proteins in Poliovirus-Infected Cells. J Virol. 2008;82:1647–55.

66. Satterly N, Tsai P-L, van Deursen J, Nussenzveig DR, Wang Y, Faria PA, et al.

Influenza virus targets the mRNA export machinery and the nuclear pore complex. Proc

Natl Acad Sci. 2007;104:1853–8.

67. Miorin L, Kehrer T, Sanchez-Aparicio MT, Zhang K, Cohen P, Patel RS, et al.

SARS-CoV-2 Orf6 hijacks Nup98 to block STAT nuclear import and antagonize

interferon signaling. Proc Natl Acad Sci. 2020;117:28344–54.

68. De Jesús-González LA, Cervantes-Salazar M, Reyes-Ruiz JM, Osuna-Ramos JF,

Farfán-Morales CN, Palacios-Rápalo SN, et al. The Nuclear Pore Complex: A Target for

NS3 Protease of Dengue and Zika Viruses. Viruses. 2020;12:583.

69. Le Sage V, Mouland AJ. Viral Subversion of the Nuclear Pore Complex. Viruses.

2013;5:2019–42.

70. Kobiler O, Drayman N, Butin-Israeli V, Oppenheim A. Virus strategies for passing

the nuclear envelope barrier. Nucleus. 2012;3:526–39.

71. Pfeffer LM. The Role of Nuclear Factor κB in the Interferon Response. J Interferon

Cytokine Res. 2011;31:553–9.

72. Liu T, Zhang L, Joo D, Sun S-C. NF-κB signaling in inflammation. Signal Transduct

Target Ther. 2017;2:1–9.

73. Jin J, Hu H, Li HS, Yu J, Xiao Y, Brittain GC, et al. Noncanonical NF-κB Pathway

Controls the Production of Type I Interferons in Antiviral Innate Immunity. Immunity.

2014;40:342–54.

74. Zhu B, Mandal SS, Pham A-D, Zheng Y, Erdjument-Bromage H, Batra SK, et al.

The human PAF complex coordinates transcription with events downstream of RNA

synthesis. Genes Dev. 2005;19:1668–73.

75. Kim J, Roeder RG. Direct Bre1-Paf1 Complex Interactions and RING Finger-independent Bre1-Rad6 Interactions Mediate Histone H2B Ubiquitylation in Yeast *. J Biol Chem. 2009;284:20582–92.

76. Simic R, Lindstrom DL, Tran HG, Roinick KL, Costa PJ, Johnson AD, et al. Chromatin remodeling protein Chd1 interacts with transcription elongation factors and localizes to transcribed genes. EMBO J. 2003;22:1846–56.

77. Dey P, Ponnusamy MP, Deb S, Batra SK. Human RNA Polymerase II-Association Factor 1 (hPaf1/PD2) Regulates Histone Methylation and Chromatin Remodeling in Pancreatic Cancer. PLOS ONE. 2011;6:e26926.

78. Kim J, Guermah M, Roeder RG. The human PAF1 complex acts in chromatin transcription elongation both independently and cooperatively with SII/TFIIS. Cell. 2010;140:491–503.

79. Pott S, Lieb JD. What are super-enhancers? Nat Genet. 2015;47:8–12.

80. Buenrostro J, Wu B, Chang H, Greenleaf W. ATAC-seq: A Method for Assaying Chromatin Accessibility Genome-Wide. Curr Protoc Mol Biol Ed Frederick M Ausubel Al. 2015;109:21.29.1-21.29.9.

81. Furey TS. ChIP-seq and Beyond: new and improved methodologies to detect and characterize protein-DNA interactions. Nat Rev Genet. 2012;13:840–52.

82. Moll P, Ante M, Seitz A, Reda T. QuantSeq 3′ mRNA sequencing for RNA quantification. Nat Methods. 2014;11:i–iii.

83. Livak KJ, Schmittgen TD. Analysis of relative gene expression data using real-time quantitative PCR and the 2(-Delta Delta C(T)) Method. Methods San Diego Calif. 2001;25:402–8.

84. Babraham Bioinformatics - FastQC A Quality Control tool for High Throughput

Sequence Data. https://www.bioinformatics.babraham.ac.uk/projects/fastqc/. Accessed

13 Feb 2022.

85. Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, et al. STAR:

ultrafast universal RNA-seq aligner. Bioinformatics. 2013;29:15–21.

86. Liao Y, Smyth GK, Shi W. featureCounts: an efficient general purpose program for

assigning sequence reads to genomic features. Bioinforma Oxf Engl. 2014;30:923–30.

87. Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion

for RNA-seq data with DESeq2. Genome Biol. 2014;15:550.

88. Leek JT, Johnson WE, Parker HS, Jaffe AE, Storey JD. The sva package for

removing batch effects and other unwanted variation in high-throughput experiments.

Bioinformatics. 2012;28:882–3.

89. Korotkevich G, Sukhov V, Budin N, Shpak B, Artyomov MN, Sergushichev A. Fast

gene set enrichment analysis. 2021;:060012.

90. Croft D, O'Kelly G, Wu G, Haw R, Gillespie M, Matthews L, et al. Reactome: a

database of reactions, pathways and biological processes. Nucleic Acids Res. 2011;39

Database issue:D691–7.

91. Liberzon A, Subramanian A, Pinchback R, Thorvaldsdóttir H, Tamayo P, Mesirov

JP. Molecular signatures database (MSigDB) 3.0. Bioinformatics. 2011;27:1739–40.

92. Huang DW, Sherman BT, Lempicki RA. Bioinformatics enrichment tools: paths

toward the comprehensive functional analysis of large gene lists. Nucleic Acids Res.

2009;37:1–13.

93. Navarro Gonzalez J, Zweig AS, Speir ML, Schmelter D, Rosenbloom KR, Raney BJ,

et al. The UCSC Genome Browser database: 2021 update. Nucleic Acids Res.

2021;49:D1046–57.

94. Sherman BT, Hao M, Qiu J, Jiao X, Baseler MW, Lane HC, et al. DAVID: a web

server for functional enrichment analysis and functional annotation of gene lists

(2021 update). Nucleic Acids Res. 2022;:gkac194.

95. Fishilevich S, Nudel R, Rappaport N, Hadar R, Plaschkes I, Iny Stein T, et al.

GeneHancer: genome-wide integration of enhancers and target genes in GeneCards.

Database. 2017;2017:bax028.

96. Oróstica KY, Verdugo RA. chromPlot: visualization of genomic data in chromosomal

context. Bioinforma Oxf Engl. 2016;32:2366–8.

97. Riemondy KA, Sheridan RM, Gillen A, Yu Y, Bennett CG, Hesselberth JR. valr:

Reproducible genome interval analysis in R. F1000Research. 2017;6:1025.

98. Bailey TL, Grant CE. SEA: Simple Enrichment Analysis of motifs.

2021;:2021.08.23.457422.

99. Buske FA, Bodén M, Bauer DC, Bailey TL. Assigning roles to DNA regulatory motifs

using comparative genomics. Bioinformatics. 2010;26:860–6.

100. Han H, Cho J-W, Lee S, Yun A, Kim H, Bae D, et al. TRRUST v2: an expanded

reference database of human and mouse transcriptional regulatory interactions. Nucleic

Acids Res. 2018;46:D380–6.

Supplementary Material

S1 Figure. Immune stress marginally affects PAF1 nuclear localization.

(A) Immunofluorescence microscopy shows how selected immune stimuli do not significantly disrupt the nuclear localization of PAF1. A549 cells were stimulated for 3 hours with the given stimuli before being fixed and then stained with PAF1 antibody (red) and Hoechst (blue) for marking nuclei. Images were produced with epifluorescence microscopy. All scale bars represent 50 μM. (B) Microscopy images were assessed quantitatively for PAF1 nuclear localization. Cell nuclei were selected for with StarDist, using Hoechst staining as the training control. Grey mean area was used to measure intensity of PAF1 signal in these nuclei (as represented by each point), relative to the average of the total grey mean area for each image. These values were plotted as a

distribution. Statistical significance was determined using an unpaired two-tailed

Student's t-test.

S2 Figure. Further characterization of immune activation by stimuli.

(A) PCA shows distinct gene expression profile of stimulations relative to mock across

both ncgRNA and PAF1 KO cell line. The counts for all genes were corrected with a

variance-stabilizing transformation via DESeq2, and only the variability of biological

replicates was removed as a batch effect with ComBat. PCA was performed on the

resulting counts, with the top 2 principal components being plotted. (B) Volcano plots

show activation of the immune response in ncgRNA cell line following each stimulation.

Comparisons were performed within the DESeq2 pipeline. The resulting log2 foldchanges

and –log10 p-values are plotted for all genes, where top immune response genes are

labeled. Cutoffs are set at log2 foldchange > 1.5 or < -1.5, and p-value < 0.05. (C) qRT-

PCR analysis validates activation of the immune response in ncgRNA and PAF1 KO cells.

Fold changes for ISG15 and CCL20 were calculated using the ΔΔCt method for three

technical replicates, being normalized against GAPDH expression. Error bars indicate

standard deviation across three biological replicates.

S3 Figure. Top PAF1-dependent genes across stimuli.

Heatmap shows relative changes in gene expression as log2 foldchanges of DEGs from

PAF1 KO versus ncgRNA contrasts of Figure 2. Colors are desaturated for genes with an

adjusted p-value > 0.05.

Supplementary Figure 4

S4 Figure. PAF1 alters gene expression unique to some immune stimuli.

Dot plots show PAF1-dependent genes include variability across the type of stimulation.

(A) The log2 foldchanges and adjusted p-values for PAF1 KO versus ncgRNA cells were

averaged across interferon (poly I:C, poly dA:dT) and inflammatory (LPS, TNF-α) stimuli.

The average log2 foldchanges were plotted on two axes. Density plots are shown for all

genes with log2 foldchanges greater than 1 or less than -1. Significant genes (p.adj <

0.05) are plotted as larger, bolded points and notable genes are labeled. Genes with

strong induction (avg. log2 fold-change > 1) in ncgRNA cells are bolded. (B) The same

analytical approach discussed above is applied to PAF1 KO DEGs for upstream (poly I:C,

poly dA:dT, LPS) versus downstream (TNF-α, IFN-β) stimuli.

Supplementary Figure 5

S5 Figure. PAF1-associated enhancer analysis pipeline and quality control.

(A) Summarization of pipeline for promoter and enhancer motif analyses. (B) Distribution

of distances of enhancer motifs from target upregulated (orange) or downregulated

(magenta) gene. (C) Distribution of the number of genes in between enhancer motif and

target upregulated (orange) or downregulated (magenta) gene.

S6 Figure. Transcription factor predicted targets network. Identified enhancer motifs

associated with known transcription factors map to immune genes (GO:0002376).

Significantly enriched enhancer motifs (E < 0.05, enrichment ratio > 2) were screened

against the TRRUST v2 database for target genes and the type of interaction. Non-

mapping motifs include those that were either not transcription factors or had no known

associations in the database. Target genes with consistent differential expression across

stimuli for PAF1 KO versus ncgRNA (average log2 foldchange > 1 or < -1) were colored

accordingly (red = downregulated, blue = upregulated, grey = did not meet criteria or

insufficient reads).

Table S1. DESeq2 outputs.

Table S2. GSEA outputs.

Table S3. Associated Promoter and Enhancer Motifs.

Table S4. qRT-PCR primers.

Table S5. Host-dependency factors.