Dysregulated *H19/Igf2* expression disrupts cardiac-placental axis during development of Silver Russell Syndrome-like mouse models

Suhee Chang[1], Diana Fulmer[1,2], Stella K. Hur[1], Joanne L. Thorvaldsen[1], Li Li[1,2], Yemin Lan[1], Eric A. Rhon-Calderon[1], Xiaowen Chen[2], Jonathan A. Epstein[1,2], Marisa S. Bartolomei[1]

[1]Department of Cell and Developmental Biology, Epigenetics Institute, Perelman School of Medicine, University of Pennsylvania, Philadelphia, PA, United States

[2]Penn Cardiovascular Institute, Perelman School of Medicine, University of Pennsylvania, Philadelphia, PA, United States

[*Keywords*: *H19*; *Igf2*; SRS; cardiac development; cardiac-placental axis; VSD; ECM; endothelial cells]

Corresponding author: bartolom@pennmedicine.upenn.edu

Abstract

Dysregulation of the imprinted *H19/IGF2* locus can lead to Silver-Russell Syndrome (SRS) in

humans. However, the mechanism of how abnormal *H19/IGF2* expression contributes to various SRS

phenotypes remains unclear, largely due to incomplete understanding of the developmental functions of

these two genes. We previously generated a mouse model with humanized *H19/IGF2* ICR (*hIC1*) on the

paternal allele that exhibited *H19/Igf2* dysregulation together with SRS-like growth restriction and

perinatal lethality. Here we dissect the role of *H19* and *Igf2* in cardiac and placental development utilizing

multiple mouse models with varying levels of *H19* and *Igf2*. We report severe cardiac defects such as

ventricular septal defects (VSDs) and thinned myocardium, placental anomalies including thrombosis and

vascular malformations, together with growth restriction in mouse embryos that correlated with the extent

of *H19/Igf2* dysregulation. Transcriptomic analysis using cardiac endothelial cells of these mouse models

shows that *H19/Igf2* dysregulation disrupts pathways related to extracellular matrix (ECM) and

proliferation of endothelial cells. Our work links the heart and placenta through regulation by *H19* and

Igf2, demonstrating that accurate dosage of both *H19* and *Igf2* is critical for normal embryonic

development, especially related to the cardiac-placental axis.

Introduction

Genomic imprinting is a mammalian-specific phenomenon where a small number of genes are

expressed in an allele-specific manner. Functionally, imprinted genes have central roles in development

and growth in both humans and mice (Barlow & Bartolomei, 2014). Additionally, proper gene dosage of

most imprinted genes is essential for normal development. Human chromosome 11 and the orthologous

region on mouse chromosome 7 harbor two jointly controlled growth regulators with opposing functions;

H19 long noncoding RNA (lncRNA) and Insulin-like Growth Factor 2 (*IGF2/Igf2*). These two imprinted

genes share an imprinting control region (ICR), a *cis*-regulatory element located between two genes,

which is essential for their allele-specific expression, as well as tissue-specific enhancers located

downstream of *H19* (Chang & Bartolomei, 2020). The *H19/IGF2* ICR, which is designated as IC1 in

humans, binds CTCF on the maternal allele, forming an insulator and enabling *H19* exclusive access to

the shared enhancers (Figure 1A). On the paternal allele, the *H19/IGF2* ICR is methylated, which

prevents CTCF from binding and an insulator from forming. Consequently, *IGF2* usurps the shared

enhancers and *H19* is repressed on the paternal allele. Ultimately, allele-specific ICR methylation

facilitates monoallelic expression of *H19* and *IGF2* with *H19* expressed from the maternal allele and

IGF2 expressed from the paternal allele.

Dysregulation of the *H19/IGF2* cluster is associated with two growth disorders, Beckwith-

Wiedemann Syndrome (BWS) and Silver-Russell Syndrome (SRS). In contrast to overgrowth observed

for BWS, SRS is characterized by intrauterine growth restriction resulting in small for gestational age

(SGA) births. Other symptoms of SRS vary widely among patients and include hemihypotrophy,

cognitive impairment, relative macrocephaly, and fifth-finger clinodactyly (Wakeling et al., 2017).

Approximately 50% of patients with SRS exhibit IC1 hypomethylation (Eggermann, Buiting, & Temple,

2011). Lack of methylation may allow the formation of an ectopic insulator on paternal IC1, which likely

explains why this class of SRS individuals has biallelic *H19* expression and greatly diminished *IGF2*

expression (Abi Habib et al., 2017; Gicquel et al., 2005). Importantly, ICR mutations in the mouse that

were generated to study imprinted gene regulation provided critical information suggesting a role for *H19*

and *IGF2* in SRS. For example, mutating CpGs at CTCF sites on the paternal *H19/Igf2* ICR resulted in

the loss of ICR methylation, activation of paternal *H19* and reduced *Igf2* expression (Engel, West,

Felsenfeld, & Bartolomei, 2004). This mutation led to restricted embryonic growth, which phenocopies

SRS. Nevertheless, although we and others successfully modeled a subset of SRS mutations in the mouse,

not all mutations were translatable because the mouse *H19/Igf2* ICR lacks extensive sequence

conservation with human IC1. Thus, a mouse model with human IC1 sequence substituted for the

endogenous mouse *H19/Igf2* ICR was generated (*H19^{hIC1}*, shortened as *hIC1*; Hur et al., 2016) to model

human mutations more precisely (Freschi et al., 2021, 2018). Consistent with expectations, maternally

transmitted *hIC1* successfully maintained insulator function, suggesting the possibility to model human

IC1 mutations endogenously in mice upon maternal transmission. In contrast, paternally transmitted *hIC1*

showed loss of methylation and formation of an ectopic insulator. Consequently, paternal *hIC1*

transmission caused elevated *H19* expression and *Igf2* depletion together with growth restriction and

embryonic lethality (Hur et al., 2016). Although the epigenetic defects and growth restriction of these

mice model SRS, the epigenetic defects in human are mosaic (Soellner et al., 2019), with a subset of cells

showing normal methylation patterns, providing a potential explanation for why SRS individuals are

viable.

The mechanism by which *H19/IGF2* expression dysregulation causes SRS phenotypes is

unknown, largely because the function of these two genes during development is incompletely

understood. *IGF2* is a well-described growth factor promoting fetoplacental growth, which functions in

an endocrine/paracrine manner through binding to IGF/Insulin receptors (Harris & Westwood, 2012).

Decreased *IGF2* levels in patients with SRS suggests that IGF2 contributes to restricted growth in these

patients (Abi Habib et al., 2017; Begemann et al., 2015; Gicquel et al., 2005). Consistently, in mice,

paternal-specific deletion of *Igf2* resulted in loss of *Igf2* expression and pre- and postnatal growth

restriction (DeChiara, Robertson, & Efstratiadis, 1991; Haley et al., 2012). In contrast, the exact role of

H19 remains unclear. Previous studies in mouse suggested that *H19* lncRNA is a precursor for microRNA

(miR)-675, which regulates *Igf1r* expression (Keniry et al., 2012), and that *H19* represses *Igf2* expression

in trans (Gabory et al., 2009). As a result, *H19* has been largely overlooked and suggested to be an

occasional regulator of *Igf2* expression. Nevertheless, a previously described mouse model with *H19*

overexpression without changes in *Igf2* expression showed embryonic growth restriction (Drewell et al.,

2000), suggesting that *H19* mediates growth suppression independently from *Igf2*. Here, we report severe

developmental defects of the heart and placenta in mouse models with dysregulated *H19/Igf2* expression.

Embryos with the paternal *hIC1* showed atrioventricular (AV) cushion defects in the heart coupled with

ventricular septal defects (VSDs), and extremely thinned myocardial walls. Combined with placental

anomalies, the cardiac defects most likely contribute to the lethality of these mice (Kochilas, Li, Jin,

Buck, & Epstein, 1999; Snider & Conway, 2011). Normalization of *H19* rescued neither the lethality nor

growth restriction, although we observed a minimal rescue of the earlier growth and resorption frequency.

Ultimately, modifying both *H19* and *Igf2* expression was necessary to rescue most phenotypes.

Transcriptomic analysis of embryonic cardiac endothelial cells identified several key signaling pathways

that are affected by the dysregulated *H19* and *Igf2,* and are potentially responsible for the paternal *hIC1-*

related cardiac defects. This work emphasizes the importance of accurate dosage of *H19* and *Igf2*

expression in normal cardiac and placental development, disruption of which can lead to SRS-like

pathologies.

99
100

Figure 1. *H19/Igf2* cluster and mouse models utilized in this study.

(A) A schematic representation of the wild-type *H19/Igf2* cluster in mouse. The maternally-expressed *H19* and the paternally-expressed *Igf2* genes are shown in red and blue, respectively. Black lollipops on the paternal allele represent DNA methylation. The maternal ICR is bound to CTCF proteins (pink hexagons) at CTCF binding sites, forming an insulator that blocks the maternal *Igf2* promoter from the shared enhancers (green circles). These enhancers interact with the *H19* promoter on the maternal allele and *Igf2* promoter on the paternal allele. (B) Schematic of the mouse endogenous *H19/Igf2* ICR, *H19^hIC1* (shortened as *hIC1*; Hur et al. 2016), *H19^Δ2.8kb-H19* (shortened as *ΔH19*)*, and *H19^Δ3.8kb-5'H19* (shortened as *Δ3.8*; Thorvaldsen et al. 2002, 2006) alleles. Restriction site locations (kb) are relative to the *H19* transcription start site. Gray lines on the ICR represent conserved CTCF binding sequences.

Results

Mouse models with genetic modifications that perturb *H19* and *Igf2* to different extents were used

to address the phenotypic consequences of abnormal *H19* and *Igf2* levels (Figure 1B). *hIC1* refers to the

humanized *H19^hIC1* allele that substitutes the endogenous mouse *H19/Igf2* ICR with the corresponding

human IC1 sequence, which was initially generated to study human BWS and SRS mutations in the

mouse (Freschi et al., 2018; Hur et al., 2016). Paternal transmission of *hIC1* [+/*hIC1*] was previously

reported to increase *H19* and greatly diminish *Igf2* expression and resulted in embryonic lethality.

Cardiac and placental defects in +/*hIC1* embryos

Our initial experiments examined the developmental phenotype of +/*hIC1* embryos to determine

how abnormal *H19/Igf2* expression resulted in dramatic growth defects and lethality. As previously

reported (Hur et al., 2016), +/*hIC1* embryos showed severe growth restriction (note that for heterozygous

embryos, maternal allele is written first, Figure 2A). The growth restriction appeared as early as E11.5

and was greatly exaggerated by the end of gestation. Although E18.5 +/*hIC1* embryos were observed

alive and weighed approximately 40% of their wild-type littermates, +/*hIC1* neonates were perinatally

lethal, with no live pups found on the day of birth. To ascertain the source of lethality, we first examined

lungs from dead +/*hIC1* neonates. The lungs floated in water, demonstrating that +/*hIC1* pups respired

after birth (Borensztein et al., 2012). Additionally, +/*hIC1* neonates did not have a cleft palate, but no

milk spots were found, indicating a lack of feeding.

Histological analyses were performed throughout development on major organs where *H19* and

Igf2 are highly expressed. In contrast to the liver where the paternal *hIC1* did not lead to significant

abnormal morphology (data not shown), severe developmental defects were found in the +/*hIC1* heart

and placenta. Cardiac defects in +/*hIC1* embryos were observed as early as E12.5, where the superior and

inferior endocardial cushions failed to fuse into a common AV cushion (Figure 2C). The cushion defect

preceded incomplete interventricular septum (IVS) formation. At E15.5, a severe perimembranous

ventricular septal defect (VSD) was observed in all 5 +/*hIC1* hearts that were evaluated (Figure 2C.

Importantly, this congenital heart defect resembles malformations reported in several SRS patients with

IC1 hypomethylation (Ghanim et al., 2013), although the prevalence is unclear and may reflect the degree

of mosaicism in SRS. Additionally, E15.5 hearts showed an extremely thin myocardium (Figure 2C).

Both VSD and thinned ventricular walls persisted in E17.5 +/*hIC1* hearts (Figure 2B, C). Additionally, 6

out of 16 +/*hIC1* hearts in the late gestation group (E15.5 to P0) had bicuspid pulmonary valve (BPV), a

rare cardiac defect in which the pulmonary valve only develops two cusps as opposed to the normal

tricuspid structure (Figure 2D and Supplemental Figure 1A). These results demonstrate that paternal *hIC1*

transmission results in variably penetrant cardiac phenotypes. Notably, atrioventricular valvuloseptal

morphogenesis, the fusion of the AV cushion and nascent septa during cardiogenesis, is required for

proper cardiac septation (Eisenberg & Markwald, 1995). Segmentation of the heart into four separate

chambers is required to establish distinct pulmonary and systemic blood flow during heart development

and is required to prevent the mixing of oxygenated and deoxygenated blood. We speculate that the

failure of *hIC1* mutants to establish complete ventricular septation could have led to a reduced ability to

provide oxygen and nutrient rich blood to the rest of the developing body (Savolainen, Foley, & Elmore,

2009; Spicer, Hsu, Co-Vu, Anderson, & Fricker, 2014).

Figure 2. Growth anomalies and cardiac defects of *+/hIC1* embryos.

(A) Fetal weight of the wild-type (blue) and *+/hIC1* (red) embryos at E11.5, E12.5, E14.5, E17.5 and E18.5. Each data point represents an average weight of each genotype from one litter. 8 litters for E11.5, 10 litters for E12.5, 7 litters for E14.5, 31 litters for E17.5, 4 litters for E18.5 are presented. (B) Quantification of ventricular wall thickness (μm), measured from E17.5 wild-type and *+/hIC1* hearts. Each data point represents an individual conceptus. 5 wild-type and 8 *+/hIC1* embryos from four different litters were examined. (C) Representative cross-sections of wild-type and *+/hIC1* embryonic hearts at E12.5, E15.5 and E17.5, stained with hematoxylin and eosin. Note the lack of fusion between AV cushions at E12.5, the VSD at E15.5 and E17.5 in *+/hIC1* hearts (yellow arrowheads).

The boxed regions of E15.5 and E17.5 images are enlarged at the bottom of the figure, to show where the ventricular wall thickness is measured. Scale bars = 500 µm. (D) A representative image of pulmonary valves in E17.5 wild-type and *+/hIC1* hearts. The *+/hIC1* right pulmonary cusp is enlarged (yellow arrowhead) and there is no cusp in the anterior position, in contrast to the tricuspid structure in the wild-type heart. Scale bars = 500 µm. Statistics used are (A) multiple paired t-test and (B) multiple unpaired t-test with **$P < 0.01$, ***$P < 0.001$, ****$P < 0.0001$ and ns = not significant.

Another major organ with high *H19/Igf2* expression, which forms early in development, is the

placenta. Multiple developmental defects were observed in *+/hIC1* placentas. As previously described

(Hur et al., 2016), *+/hIC1* placentas were growth restricted throughout development (Figure 3A),

although the fetal to placental weight ratio was not affected through E15.5 (Supplemental Figure 2A).

However, at E17.5, the fetal to placental weight ratio was lower in *+/hIC1* conceptuses, indicating that

the fetal growth restriction was more severe than the placental growth restriction as the concepti neared

term (Figure 3B). In addition to placental undergrowth, the junctional to labyrinth zone ratio was

increased in *+/hIC1* placentas (Figure 3C), suggesting that the growth of the labyrinth layer, where the

maternal-fetal exchange occurs, was more affected. Moreover, *H19* overexpression was exaggerated in

the labyrinth in E17.5 *+/hIC1* placentas, while the *Igf2* depletion was consistent throughout the whole

placenta (Figure 3D), indicating that *H19* overexpression contributed disproportionately to the phenotype

of growth restriction in the labyrinth. Large thrombi were observed in the labyrinth zone of these *+/hIC1*

placentas (Figure 3F), in a male-skewed manner (Figure 3E). As the thrombi could be formed due to

defective vasculature structures, wild-type and *+/hIC1* placentas were stained for CD34, a marker for the

fetoplacental endothelial cells that line the microvessels in the labyrinth layer (Figure 3G). Vessels in the

+/hIC1 labyrinth were highly dilated and had decreased density, which was confirmed by quantifying the

stained areas on the images (Figure 3H and Supplemental Figure 2B). The reduced labyrinth layer and

defective microvascular expansion likely compromised the ability of *+/hIC1* placentas to supply nutrients

and oxygen to the fetus. These results support the hypothesis that the abnormal growth of *+/hIC1*

 embryos may be explained by failure in multiple organs, especially the heart and placenta, which are

 developmentally linked (Barak, Hemberger, & Sucov, 2019).

Figure 3. Placental defects of *+/hIC1* embryos.

(A) Placental weight of the wild-type (blue) and *+/hIC1* (red) samples at E11.5, E14.5, E17.5 and E18.5. Each data point represents an average weight of each genotype from one litter. 3 litters for E11.5, 6 litters for E12.5, 7 litters for E14.5, 22 litters for E17.5, 4 litters for E18.5 are presented. (B) Fetal to placental weight ratios in E17.5 wild-type and *+/hIC1* samples (mean ± SD). 40 wild-type and

43 +/*hIC1* embryos from 22 litters are presented. (C) Junctional zone (Jz) to labyrinth (Lb) ratio in

E17.5 wild-type and +/*hIC1* placentas. (D) Relative total expression of *H19* and *Igf2* in E17.5 wild-

type and +/*hIC1* placentas and labyrinths (mean ± SD). (E) Number of wild-type, male and female

+/*hIC1* placentas with thrombi observed. (F) Representative cross-sections of E12.5, E15.5 and

E17.5 wild-type and +/*hIC1* placentas stained with hematoxylin and eosin. Black arrowhead indicates

a large thrombus formed in the +/*hIC1* labyrinth. Scale bars = 1mm. (G) Representative images of

CD34 immunostaining counterstained with hematoxylin on E17.5 wild-type and +/*hIC1* placental

sections. Black arrowheads indicate thrombi in the +/*hIC1* labyrinth. Scale bars = 1mm. (H)

Quantification of the microvessel density in E17.5 wild-type and +/*hIC1* placentas. 11 wild-type, 12

+/*hIC1* male, 10 +/*hIC1* female placentas from 8 litters were quantified. (B, C, D, H) Each data point

represents an individual conceptus from different litters. Statistics used are (A) multiple paired t-test,

(B) unpaired Student's t-test, (C, D, H) one-way ANOVA with Tukey's multiple comparisons test, (E)

Fisher's exact test. *$P < 0.05$, **$P < 0.01$, ***$P < 0.001$, ****$P < 0.0001$, ns = not significant.

Normalizing *H19* expression partially rescues paternal *hIC1* defects

It has been previously reported that *Igf2* null mice are viable (DeChiara et al., 1991). Thus, we

hypothesize that *H19* overexpression combined with a loss of *Igf2* expression is the molecular contributor

to the lethality of paternal *hIC1* transmission. To examine if reduced *H19* expression would rescue the

paternal *hIC1* transmission phenotypes, we generated a mouse model with deletion of the *H19*

transcription unit [*H19$^{\Delta2.8kb-H19}$*; shortened as *ΔH19*] (Figure 1B, Supplemental Figure 3A and B).

Consistent with previous reports, maternal deletion of *H19* [*ΔH19/+*] led to an absence of *H19* expression

and tissue-specific minimal activation of maternal *Igf2* (Supplemental Figure 3C). These mice are viable

and fertile, regardless of whether the deletion is maternally or paternally transmitted, although maternal

transmission is associated with increased fetal weight from E14.5 and onwards (Supplemental Figure 3D

and data not shown).

Heterozygous *ΔH19* females were mated with heterozygous *hIC1* males to generate *ΔH19/hIC1*

embryos. These embryos were expected to have lower *H19* expression compared to +/*hIC1* embryos, as

the maternal *H19* expression was silenced (Figure 4A). Among four possible genotypes from this

breeding, +/*hIC1* embryos constituted approximately 15% per litter at E17.5, as opposed to the expected

Mendelian ratio of 25%. In contrast, *ΔH19/hIC1* embryos comprised approximately 30% per litter at

E17.5, indicating partial rescue of the resorption frequency by maternal *H19* deletion (Figure 4B and

Supplemental Figure 4A). However, *ΔH19/hIC1* embryos still exhibited perinatal lethality, with no live

pups observed on the day of birth. With respect to growth restriction, the maternal *ΔH19* allele partially

rescued the phenotype. At E11.5, *ΔH19/hIC1* fetal weight was not significantly different from wild-type

littermates (Figure 4C). However, at late gestation (E17.5), although *ΔH19/hIC1* fetuses had a significant

increase in fetal weight compared to the *+/hIC1* fetuses, *ΔH19/hIC1* fetuses remained significantly

smaller compared to wild-type. As perinatal lethality was still observed, conceptuses were analyzed

histologically to characterize their developmental defects. The AV cushion defect persisted in all

examined E13.5 *ΔH19/hIC1* embryos, and 50% of the examined E17.5 *ΔH19/hIC1* hearts showed either

perimembranous or muscular VSDs (Figure 4D). Thrombi were still present in approximately 50% of

ΔH19/hIC1 placentas (Figure 4E), and placental weight remained significantly lower compared to wild-

type littermates (Supplemental Figure 4B). Of note, none of these histological defects were observed in

ΔH19/+ embryonic hearts and placentas (Figure 4D and data not shown; 3 *ΔH19/+* hearts each from

E13.5, E15.5 and E17.5 concepti, and 20 E17.5 *ΔH19/+* placentas were examined). E17.5 *ΔH19/hIC1*

tissues demonstrated wild-type levels of *H19* expression, while the *Igf2* expression remained markedly

lower than wild-type (Figure 4F and Supplemental Figure 4C). From these results, we conclude that

restoring *H19* expression is not sufficient to rescue completely the lethality and developmental defects

upon paternal transmission of *hIC1*. Thus, phenotypes are likely caused by abnormal expression of both

H19 and *Igf2*.

Figure 4. Normalizing *H19* expression through the maternal deletion of *H19*.

(A) A schematic representation of the rescue breeding between *ΔH19* heterozygous female and *hIC1/+* male mice is depicted. *ΔH19/hIC1* embryos are expected to express *H19* only from the paternal allele, and maternally express *Igf2* in a tissue-specific manner. (B) Ratio of *+/hIC1* and *ΔH19/hIC1* embryos observed in E11.5 and E17.5 litters (8 E11.5 litters and 15 E17.5 litters with litter size larger than 5 pups were examined). Each data point represents one litter. (C) Relative fetal weights of wild-type, *ΔH19/+, +/hIC1* and *ΔH19/hIC1* embryos at E11.5 and E17.5, normalized to the

average body weight of the wild-type littermates (mean ± SD). (D) Representative cross-sections of wild-type, *ΔH19/+* and *ΔH19/hIC1* embryonic hearts at E13.5 and E17.5, stained with hematoxylin and eosin. Note the cushion defect at E13.5 and the VSD at E17.5 in *ΔH19/hIC1* hearts (yellow arrows). Scale bars = 500 µm. (E) (Top) Representative cross-section of E17.5 *ΔH19/hIC1* placenta stained with hematoxylin and eosin. Black arrowhead indicates a thrombus. Scale bar = 1mm. (Bottom) Number of the wild-type, male and female *+/hIC1* placentas with thrombi observed. (F) Relative total expression of *H19* and *Igf2* in E17.5 wild-type, *ΔH19/+, +/hIC1* and *ΔH19/hIC1* hearts (mean ± SD). (C, F) Each data point represents an individual conceptus from different litters. Statistics used are (B, C, F) one-way ANOVA with Tukey's multiple comparisons test and (E) Fisher's exact test. *$P < 0.05$, **$P < 0.01$, ***$P < 0.001$, ****$P < 0.0001$, ns = not significant.

Paternal *hIC1* defects are rescued by deletion of the maternal *H19/Igf2* ICR

205   Finally, we utilized a previously published mouse model with a 3.8-kb deletion spanning the

206 *H19/Igf2* ICR [*H19^{Δ3.8kb-5'H19}*; shortened as *Δ3.8*] to modulate both *H19* and *Igf2* expression (Thorvaldsen

207 et al., 2006, 2002). Absence of the maternal *H19/Igf2* ICR activates the maternal *Igf2* allele and reduces

208 *H19* expression. Thus, *Igf2* expression from the maternal allele in *Δ3.8/hIC1* embryos was expected to

209 restore *Igf2* levels (Figure 5A).

210   Crosses between heterozygous *Δ3.8* females and heterozygous *hIC1* males produced the expected

211 Mendelian ratio of offspring with *Δ3.8/hIC1* mice appearing fully viable. Both fetal and placental weights

212 were not significantly different between *Δ3.8/hIC1* and wild-type concepti at E17.5 (Figure 5B and

213 Supplemental Figure 5A), demonstrating full rescue of the lethality and growth restriction. However, a

214 subset of *Δ3.8/+* and *Δ3.8/hIC1* embryonic hearts (3 out of 6 *Δ3.8/+* hearts and 2 out of 5 *Δ3.8/hIC1*

215 hearts) had VSDs, although the lesions in the IVS were smaller than those found in *+/hIC1* hearts (Figure

216 5C). While *Igf2* expression in the E17.5 *Δ3.8/hIC1* hearts was restored to wild-type levels, normalization

217 of *H19* expression varied among embryos. Although not statistically significant in heart, *Δ3.8/hIC1*

218 embryos tended to have higher *H19* expression compared to the wild-type littermates (Figure 5D and

219 Supplemental Figure 5B). These results suggest that the physiological levels of *H19* and *Igf2* expression

220 are critical for normal cardiac development, and the variability in *H19* rescue could help to explain the

221   varying penetrance of the cardiac phenotype. No thrombi were detected in the *Δ3.8/hIC1* placentas, and

222   the placental morphology was normal with the junctional to labyrinth zone ratio not significantly different

223   compared to wild-type (Supplemental Figure 5C). In sum, restoring both *H19* and *Igf2* to near wild-type

224   levels was necessary for the full rescue of the most severe pathologies of paternal *hIC1* transmission.

Figure 5. Restoring *H19* and *Igf2* expression utilizing maternal *H19/Igf2* ICR deletion.
(A) A schematic representation of the offspring produced when *Δ3.8* heterozygous female and *hIC1/+* male mice are mated is depicted. *Δ3.8/hIC1* embryos are expected to show activation of maternal *Igf2* expression as well as paternal *H19* expression. (B) Relative fetal weights of wild-type, *Δ3.8/+, +/hIC1,* and *Δ3.8/hIC1* embryos at E11.5 and E17.5, normalized to the average body weight of wild-type littermates (mean ± SD). 2 E11.5 and 10 E17.5 litters are presented. (C) Representative cross-sections of E17.5 *Δ3.8/+* and E18.5 *Δ3.8/hIC1* embryonic hearts with VSDs (yellow arrows), stained

with hematoxylin and eosin. Sections from wild-type littermates are shown together for comparison. Scale bars = 500 µm. (D) Relative total expression of *H19* and *Igf2* in E17.5 wild-type, *Δ3.8/+*, *+/hIC1,* and *Δ3.8/hIC1* hearts (mean ± SD). (B, D) Each data point represents an individual conceptus from different litters. One-way ANOVA with Tukey's multiple comparisons test was used with *$P < 0.05$, ***$P < 0.001$, ****$P < 0.0001$ and ns = not significant.

Transcriptomic analysis of cardiac endothelial cells with various *H19/Igf2* expression

Severe cardiac phenotypes associated with paternal *hIC1* transmission prompted us to question

the mechanism of how dysregulated *H19/Igf2* causes such developmental defects. The paternal *hIC1-*

associated cardiac phenotypes were observed as early as E12.5 when AV cushion fusion is delayed in

developing hearts (Figure 2B), making E12.5 an optimal time point to identify the key pathways of valve

development and cardiac septation that are disrupted by *H19/Igf2* dysregulation. Endothelial and

endothelial-derived cells comprise the majority population in the AV cushion and majority non-myocyte

population of the ventricular septum at E12.5 (Von Gise & Pu, 2012). Moreover, both *H19* and *Igf2* are

strongly expressed in the endocardial layer of developing heart (García-Padilla, Domínguez, Aránega, &

Franco, 2019). Therefore, transcriptomic analysis was performed on cardiac endothelial cells of each

mutant.

CD31+ cardiac endothelial cells from E12.5 wild-type, *+/hIC1*, *ΔH19/+, ΔH19/hIC1, Δ3.8/+,*

and *Δ3.8/hIC1* embryos were collected for RNA sequencing. First, we confirmed that these 6 groups

show gradual alteration of *H19/Igf2* expression (Figure 6A), which enabled us to generate multiple

comparisons relative to *H19* and *Igf2* levels and potentially enabling attribution of phenotypes to *H19* or

Igf2. Notably, *H19/Igf2* expression was indistinguishable in the wild-type and *Δ3.8/hIC1* samples.

Additionally, there were no sex-specific differences in *H19/Igf2* expression across all the groups.

To elucidate candidate genes and cellular processes in conferring the paternal *hIC1*-specific

cardiac phenotypes, we compared *+/hIC1* and wild-type samples. This comparison resulted in 224

significant differentially expressed genes (DEGs) (Figure 6B, left). Gene ontology (GO) analysis revealed

that pathways related to proliferation and remodeling of endothelial and epithelial cells are enriched in

these DEGs (Figure 6B, right) (Mi, Muruganujan, Ebert, Huang, & Thomas, 2019). Notably, genes

involved in extracellular matrix (ECM) and specifically, collagen matrix production, were differentially

enriched in E12.5 *+/hIC1* samples; consistent with increased collagen in the E17.5 *+/hIC1* hearts, as

confirmed through immunofluorescence staining (Figure 6C, Supplemental Figure 6A and B). Thus,

differential gene expression in the presence of paternal *hIC1* was evidenced by E12.5 and correlated with

histological consequences persisting through gestation.

Because cardiac phenotypes were indistinguishable between males and females, we next used

same-sex comparisons between *+/hIC1* and wild-type to identify the *hIC1*-specific DEGs that are

significant in both males and females (Figure 6D, left and Supplemental Figure 6C). The 25 genes that

overlapped between male and female *+/hIC1*-specific DEGs included *Col14a1,* which was upregulated in

+/hIC1, emphasizing the importance of collagen-related ECM in *hIC1*-associated cardiac defects. The

expression pattern of these 25 genes across all samples clustered *+/hIC1* and *ΔH19/hIC1*, the two groups

with the most severe cardiac defects and perinatal lethality (Figure 6D, right).

To separate the effect of *Igf2* depletion from that of *H19* overexpression, we utilized *ΔH19/hIC1*

samples. Consistent with previous observations from E17.5 hearts (Figure 4F), *ΔH19/hIC1* endothelial

cells exhibited low *Igf2,* but *H19* expression was not significantly different from wild-type (Supplemental

Figure 7A). Thus, *ΔH19/hIC1* samples were compared to *+/hIC1* to clarify the sole effect of *H19*

overexpression. Here, 46 DEGs were identified (Supplemental Figure 7B), which were enriched in

vascular endothelial cell proliferation pathways (Supplemental Figure 7C). Surprisingly, although

physiologically similar to *+/hIC1,* the *ΔH19/hIC1* transcriptome was quite different from that of *+/hIC1*.

Compared to wild-type, *ΔH19/hIC1* only had 23 DEGs (Supplemental Figure 7A), in contrast to *+/hIC1*

showing more than 200 DEGs compared to wild-type (Figure 6B, left). This result underscores the

overwhelming effect of increased *H19* in transcriptomic regulation.

We then analyzed *Δ3.8/hIC1* samples to identify cellular processes that are required to rescue the

paternal *hIC1*-associated lethality, as *Δ3.8/hIC1* mice are fully viable with occasionally observed VSD.

While *Δ3.8/hIC1* had only two DEGs compared to wild-type (Figure 6E), 487 genes were differentially

expressed in *Δ3.8/hIC1* compared to *+/hIC1* (Supplemental Figure 8A). Within these DEGs, we wanted

to clarify the genes that were likely involved in restoring viability. The 487 DEGs between *+/hIC1* and

Δ3.8/hIC1 were compared to the DEGs between *+/hIC1* and wild-type to filter genes that are commonly

affected in both comparisons (Figure 6F, left). GO analysis showed that the 116 overlapping genes are

associated with endothelial/epithelial cell proliferation and remodeling (Supplemental Figure 8B),

emphasizing the importance of the proper regulation of these pathways in the rescued viability of

Δ3.8/hIC1 embryos. The expression pattern of these 116 genes clustered *+/hIC1* and *ΔH19/hIC1*,

implicating these genes in the perinatal lethality characteristic of these two groups (Figure 6F, right).

Figure 6. Transcriptomic analysis of E12.5 cardiac endothelial cells from wild-type and mutant embryos.

(A) A gradient *H19* and *Igf2* expression levels are depicted in E12.5 wild-type, *+/hIC1, ⊿H19/+,* *⊿H19/hIC1, ⊿3.8/+* and *⊿3.8/hIC1* cardiac endothelial cells. M: Male, F: Female. (B) (Left) Comparison between *+/hIC1* and the wild-type samples with a volcano plot shows 224 DEGs between *+/hIC1* and the wild-type samples. (Right) GO pathways that are enriched for the 224 DEGs. (C) Immunofluorescence staining for MF20 (red) and collagen (green) on E17.5 wild-type and *+/hIC1* hearts. Images on the right are enlarged from the boxed area of images on the left. Scale bars = 100 µm. (D) Expression pattern of 25 genes that are differentially expressed in both male and female *+/hIC1* samples is compared to wild-type. (E) A volcano plot represents 2 DEGs between *⊿3.8/hIC1* and the wild-type samples. (F) Expression pattern of 116 genes that are commonly differentially expressed in *+/hIC1* compared to the wild-type and *⊿3.8/hIC1* samples.

Discussion

In this study, we report that the overexpression of *H19* combined with *Igf2* depletion leads to

severe morphological defects in the heart and placenta, which are likely to be involved with the perinatal

lethality and restricted growth of SRS mouse models. Genetically correcting *H19* was not sufficient to

fully rescue the developmental defects, indicating that the SRS-like phenotypes of paternal *hIC1*

transmission are not solely attributable to *H19* overexpression. Unexpectedly, although moderately

adjusting both *H19* and *Igf2* rescued the lethality, septal defects persisted in some of the *⊿3.8/hIC1*

embryos, suggesting that cardiac development is extremely sensitive to the dosage of *H19* and *Igf2*. Our

transcriptomic profiling of cardiac endothelial cells with various levels of *H19* and *Igf2* expression

uncovers critical pathways driven by *H19* and *Igf2* that are important for cardiac structure formation. The

result suggests that the regulation of ECM and proliferation of endothelial cells are tightly regulated by

H19 and *Igf2,* and potentially responsible for the paternal *hIC1*-associated cardiac defects.

The function of *H19* in cardiac development is understudied even though the expression of *H19* is

robust in the developing endocardium and epicardium throughout gestation (García-Padilla et al., 2019).

Abnormal *H19/Igf2* expression in *+/hIC1* hearts disrupted AV cushion fusion, ventricular septation and

valve formation processes with variable penetrance. These events require properly established ECM,

which accommodates endothelial-mesenchymal transition, cell proliferation, and cell-cell adhesion in

developing hearts (Kruithof, Krawitz, & Gaussin, 2007; Sullivan & Black, 2013; Von Gise & Pu, 2012).

In adult murine hearts, where *H19* has been well implicated in cardiac fibrosis and remodeling (Greco et

al., 2016; Hobuß et al., 2020; Lee et al., 2011; Wang, Sun, & Sun, 2021), *H19* overexpression led to

increased ECM and fibrosis markers after myocardial injury, while deleting *H19* resulted in

downregulated ECM genes (Choong et al., 2019). In our *+/hIC1* endothelial cells, the expression of

several key ECM genes such as *Periostin* (*Postn*) (Snider, 2009; Sullivan, 2013), *Col14a1*, and *Adamts17*

(Hubmacher, 2015) was significantly upregulated compared to wild-type cells. Combined with increased

collagen in E17.5 *+/hIC1* hearts (Figure 6C), we hypothesize that failing to establish proper ECM

contributes significantly to the *+/hIC1*-associated cardiac defects during development. Additionally,

regulators of cell proliferation, including *E2f5, Trabd2b,* and *Septin4,* and genes associated with

inflammation such as *Sele* and *Cd200* were also differentially expressed in *+/hIC1* samples. Overall, this

work provides some hints regarding the potential mechanism underlying how increased *H19* expression

disrupts normal cardiac development.

To identify the genes whose expression is solely affected by *H19,* we compared *+/hIC1,*

ΔH19/hIC1, and *ΔH19/+* endothelial cells. Among 224 genes that are differentially expressed in *+/hIC1*

endothelial cells relative to wild-type, 15 genes are also differentially expressed in *ΔH19/hIC1* samples,

suggesting that these 15 genes were affected by the loss of *Igf2* rather than increased *H19.* Among the

remaining 209 DEGs that are not altered in *ΔH19/hIC1* samples, only *Fgf10* and *H19* were also

differentially expressed in *ΔH19/+* endothelial cells compared to wild-type. *Fgf10,* a key regulator of

cardiac fibroblast development, mediates communication between cardiac progenitor cells and regulates

cardiac myocyte proliferation (Hubert, Payan, & Rochais, 2018; Vega-Hernández, Kovacs, de Langhe, &

Ornitz, 2011). Consistently, *Fgf10* null mouse embryos showed abnormal cardiac morphology with

reduced heart size and thinned ventricular wall (Rochais et al., 2014; Vega-Hernández et al., 2011). In our

samples, *Fgf10* is upregulated when *H19* is deleted and downregulated upon *H19* overexpression, linking

H19 and *Fgf10* closely in the context of cardiac development. Additionally, Gene Set Enrichment

Analysis (GSEA) revealed that the set of imprinted genes (BRIDEAU_IMPRINTED_GENES) including

Cdkn1c, Dlk1 and *Gatm* was differentially enriched in *+/hIC1* samples (Supplemental Figure 9A)

(Mootha et al., 2003; Subramanian et al., 2005). The same set of genes was also differentially enriched in

ΔH19/+ samples (Supplemental Figure 9B), but not *ΔH19/hIC1* (Supplemental Figure 9C), underscoring

a role for *H19* as a master regulator of the imprinted gene network (IGN) (Gabory, Jammes, & Dandolo,

2010).

In contrast to the less well understood role for *H19*, *Igf2* is the main growth factor in the

developing ventricular wall. Similar to *H19, Igf2* is highly expressed in the developing cardiac

endocardium and epicardium from early gestation (Shen et al., 2015) before ventricular septation is

completed (Savolainen et al., 2009). Deletion of *Igf2* and its receptors caused decreased cardiomyocyte

proliferation and ventricular hypoplasia, suggesting that *Igf2* is the major regulator of ventricular wall

thickening (Li et al., 2011; Shen et al., 2015). Additionally, the interventricular septum is comprised of

both mesenchymal and muscular components (Penny & Vick, 2011; Spicer et al., 2014), and a reduction

in cardiomyocyte proliferation can lead to VSD (Snider & Conway, 2011). Thus, the lack of *Igf2,* a

growth promoter for cardiomyocytes, could have contributed to the septal defects observed in *+/hIC1* and

ΔH19/hIC1 hearts. However, the thinned myocardium in *Igf2* knockout mouse embryos was resolved by

birth, resulting in normal cardiac morphology in neonates (Shen et al., 2020). In contrast, ventricular wall

thinning and septal defects in *+/hIC1* and *ΔH19/hIC1* hearts were aggravated towards the end of

gestation, indicating that these phenotypes are not exclusively attributable to the loss of *Igf2* expression.

Recovery of *Igf2* expression in *Δ3.8/hIC1* hearts rescued ventricular hypoplasia, consistent with

previous findings (Li et al., 2011; Shen et al., 2015). In contrast, septal defects persist in some *Δ3.8/+* and

Δ3.8/hIC1 embryos. Because *Igf2* expression varied substantially among *Δ3.8/+* and *Δ3.8/hIC1* hearts

(Figure 5D), we hypothesize that the varying penetrance of septal defects in these hearts reflects the range

of *H19/Igf2* levels. The only upregulated gene in *Δ3.8/hIC1* endothelial cells compared to wild-type was

Cadherin 18 (Cdh18) (Figure 6E), which was also upregulated in *Δ3.8/+* samples compared to wild-type

(Supplemental Figure 8C) and clinically reported to be mutated in CHDs including VSD (Chen et al.,

2018; Soemedi et al., 2012). Although we and others showed that restoring *Igf2* successfully rescues the

growth restriction in SRS-like mouse models (Han, Szabó, & Mann, 2010; Liao et al., 2021), septal

defects caused by *H19* and *Igf2* dysregulation are reported for the first time in this study. Our data provide

evidence that ventricular septation may be regulated separately from the ventricular wall thickening, and

that both events are extremely sensitive to the level of *H19* and *Igf2* expression. In humans, VSDs were

found in SRS patients with IC1 hypomethylation (Ghanim et al., 2013) and a patient carrying

chromosomal gain of chr11p15 (Serra et al., 2012). Thus, the VSD observed in our mouse models nicely

models the SRS-associated CHD in human patients. It should be noted, however, that mice are often more

susceptible to VSDs than humans. Mice with VSDs show a higher neonatal mortality rate, possibly due to

the higher heart rates and relatively larger size of VSD lesions (Snider & Conway, 2011). Additionally,

cardiac defects that are lethal in mice can lead to spontaneous miscarriages in humans due to longer

human gestation, making it difficult to observe human infants with similar defects.

Linked through fetoplacental blood circulation, the heart and placenta are closely connected

under the cardiac-placental axis, which is crucial for fetal growth and viability (Barak et al., 2019;

Maslen, 2018). Both organs are responsible for supplying nutrients and oxygen for developing fetuses,

and placental and cardiac defects are often coupled in mouse models (Perez-Garcia et al., 2018) and

humans (Matthiesen et al., 2016; Rychik et al., 2018). Placental maintenance of a normoxic fetal

environment is vital for cardiac morphogenesis, and hypoxia can lead to severe defects emerging

especially in cushions and septa (Dor et al., 2001). Epicardial *Igf2* expression in developing ventricles is

induced by a normoxic environment that is dependent on the placental function (Shen et al., 2015), while

H19 is upregulated under hypoxic conditions in mouse cardiomyocytes (Choong et al., 2019). This

suggests that *H19* and *Igf2* are important mediators of the interaction between heart and placenta.

Consistent with previous reports, gene sets of which expression is altered under hypoxic conditions

(HALLMARK_HYPOXIA, GROSS_HYPOXIA_VIA_HIF1A_DN) are differentially enriched in our

+/hIC1 endothelial cells compared to wild-type (data not shown). As the heart and placenta are

responsible for meeting the fetal demand for nutrients and oxygen, the malfunction of these two organs

would severely constrain embryonic growth. Therefore, the precise role of *H19* and *Igf2* in cardiac-

placental communication needs to be clarified to understand how the SRS-related growth restriction is

induced by IC1 hypomethylation.

The labyrinth, where *+/hIC1* placentas show abnormal vasculature morphology and thrombosis,

serves as a prime location for maternal-fetal blood exchange (Woods, Perez-Garcia, & Hemberger, 2018).

Placental thrombosis can be caused by defective labyrinth integrity, and diminished labyrinth function

could limit the nutritional and oxygen supply for a fetus, which can, in turn, lead to hypoxia and growth

restriction. Both *H19* and *Igf2* are highly expressed in fetoplacental endothelial cells in the labyrinth

(Aykroyd, Tunster, & Sferruzzi-Perri, 2022; Sandovici et al., 2022). Genetically depleting *Igf2* expression

in epiblast lineage led to decreased labyrinth size and caused the formation of thrombi in the labyrinth,

although the lesions were smaller in size than those observed in *+/hIC1* placentas (Sandovici et al., 2022).

Depletion of the placental-specific *Igf2* transcript in mice resulted in smaller labyrinth and fetal growth

restriction (Constância et al., 2002; Sibley et al., 2004), although these mice showed an increased fetal to

placental weight ratio, which was decreased in *+/hIC1* mice. This contrasting result could be explained by

the effect of the increased *H19* expression in *+/hIC1* mice. *H19* regulates vascular endothelial growth

factor (VEGF) in human endothelial cells *in vitro* (Conigliaro et al., 2015), which is involved in placental

angiogenesis, specifically for the branching of fetoplacental vessels beginning at mid-gestation (Woods

et al., 2018). Additionally, *H19* is highly expressed in trophoblasts (Marsh & Blelloch, 2020; Poirier et

al., 1991), and disrupted trophoblast development leads to defective vascular branching in the labyrinth

and restricted fetal growth (Ueno et al., 2013). Thus, it is possible that the morphological anomalies

observed in the *+/hIC1* placenta are exaggerated by abnormal *H19* expression in trophoblasts.

In summary, we provide evidence that the proper dosage of *H19* and *Igf2* is essential for normal

cardiac and placental development. Investigation of the role of *H19* and *Igf2* in the cardiac-placental axis

will enable a better understanding of how paternal *hIC1* transmission leads to the SRS-like growth

restriction and perinatal lethality. As many patients with SRS exhibit DNA methylation mosaicism, the

distribution of the epimutation, which reflects the severity of patient symptoms, varies. This work

provides insight into identifying organs that are most sensitive to *H19* and *Igf2* dysregulation, which

would allow us to develop early intervention methods for critical SRS pathologies with such variabilities.

Materials and Methods

Animal studies. All studies were approved by the Institutional Animal Care and Use Committee

(IACUC) at the University of Pennsylvania. *hIC1* (Hur et al., 2016) and *Δ3.8* (Thorvaldsen et al., 2006,

2002) mouse models were previously described. Generation of *ΔH19* line is described in Supplemental

Information. Timed breeding was performed as previously described (SanMiguel, Abramowitz, &

Bartolomei, 2018). Vaginal sperm plugs were checked to calculate the age, and the day of the plug was

marked as E0.5. Visual staging confirmed the embryonic days at the time of dissection. All mice were

maintained on C57BL/6 background for more than 10 generations if not noted otherwise. Genotyping was

performed as described in Supplemental Information.

Gene Expression Analysis. Mouse tissues were ground using pestles, syringes and needles in either

TRIzol (Thermo Fisher scientific, Waltham, MA) or RLP buffer included in RNeasy Mini kit (Qiagen,

Hilden, Germany). RNA was isolated according to the manufacturer's instructions. cDNA synthesis,

qRT-PCR, and allele-specific expression analysis was performed as previously described (Hur et al.,

2016). Primers and PCR conditions are listed in Supplemental Table 1. RNA Sequencing library

preparation and analysis are described in Supplemental Information.

Histological analysis. Mouse heart and placenta samples were collected in cold PBS, fixed overnight in

4% paraformaldehyde or 10% phosphate-buffered formalin and processed through ethanol dehydration.

Tissues were paraffin-embedded and sectioned for further staining analysis. Hematoxylin and eosin

staining was performed using a standard protocol. Immunohistochemistry for MF20 (Developmental

Studies Hybridoma Bank, Iowa City, IA) and Collagen I (cat# ab34710, Abcam, Cambridge, UK) was

performed with primary antibody incubations overnight at 4°C. Prior to antibody incubation, antigen

retrieval with citrate buffer was performed, followed by a 1 hour block in 10% normal serum. DAPI

(cat#32670-5MG-F, Sigma-Aldrich, St. Louis, MO) was used as a counter stain, and slides were mounted

with VECTASHILD (Vector, Burlingame, CA). Images were taken on a Leica DMi8S widefield

microscope. Placental CD34 staining was previously described (Vrooman et al., 2020). The thickness of

ventricular wall was measured on hematoxylin-eosin-stained heart sections using Adobe Photoshop. A

minimum of 3 distinct sections were quantified for each mouse of each genotype in a blinded manner.

Placental Jz/Lb ratio was measured using FIJI (ImageJ v2.0.0, Schindelin et al., 2012) in a blinded

manner. CD34 stained placental sections were digitally scanned using Aperio VERSA 200 platform in

Comparative Pathology Core (CPC) at School of Veterinary Medicine at the University of Pennsylvania.

Images were analyzed via Aperio Microvessel Analysis algorithm as previously described (Vrooman et

al., 2020).

Statistical Analysis. Differences between two groups were evaluated using Student's t-test. For three or

more groups, ordinary one-way analysis of variance (ANOVA), followed up with Tukey's multiple

comparisons test, was used. Two-sided Fisher's exact test was used to compare the occurrence of the

thrombi in the wild-type and *+/hIC1* placentas. All analyses were performed using GraphPad Prism

software. $*P < 0.05$, $**P < 0.01$, $***P < 0.001$, $****P < 0.0001$, n.s. = not significant.

Acknowledgments

Authors would like to express gratitude to Christopher Krapp, Lisa Vrooman, Joel Rurik, Olga Smirnova,

Jonathan Schug, Klaus Kaestner, and Colin Conine for their guidance on this study. This work was

supported by National Institutes of Health grant GM-051279-28 and R35HL140018-05, National Institute

of Arthritis and Musculoskeletal and Skin Diseases grant 5T32AR053461.

Author Contributions

S.C. and M.S.B. designed the study. S.C. and J.L.T. performed experiments and maintained mouse lines.

S.K.H. generated *ΔH19* allele. S.C., D.F., J.A.E. and L.L. evaluated cardiac histology. E.A.R. and X.C.

evaluated placental histology. S.C. and Y.L. analyzed RNA-seq data. S.C. and M.S.B. wrote the

manuscript with input from all authors.

Competing Interests

The authors declare no competing interests.

Data Availability

RNA sequencing data are deposited in GEO database under the accession number GSE199377.

References

Abi Habib, W., Brioude, F., Azzi, S., Salem, J., Das Neves, C., Personnier, C., … Netchine, I. (2017). 11p15 ICR1 Partial Deletions Associated with IGF2/H19 DMR Hypomethylation and Silver–Russell Syndrome. *Human Mutation*. https://doi.org/10.1002/humu.23131

Aykroyd, B. R. L., Tunster, S. J., & Sferruzzi-Perri, A. N. (2022). Loss of imprinting of the Igf2-H19 ICR1 enhances placental endocrine capacity via sex-specific alterations in signalling pathways in the mouse . *Development*, *149*(1). https://doi.org/10.1242/dev.199811

Barak, Y., Hemberger, M., & Sucov, H. M. (2019). Phases and Mechanisms of Embryonic Cardiomyocyte Proliferation and Ventricular Wall Morphogenesis. *Pediatric Cardiology*, *40*(7), 1359–1366. https://doi.org/10.1007/s00246-019-02164-6

Barlow, D. P., & Bartolomei, M. S. (2014). Genomic imprinting in mammals. *Cold Spring Harbor Perspectives in Biology*. https://doi.org/10.1101/cshperspect.a018382

Begemann, M., Zirn, B., Santen, G., Wirthgen, E., Soellner, L., Büttel, H.-M., … Eggermann, T. (2015). Paternally Inherited IGF2 Mutation and Growth Restriction . *New England Journal of Medicine*, *373*(4). https://doi.org/10.1056/nejmoa1415227

Borensztein, M., Viengchareun, S., Montarras, D., Journot, L., Binart, N., Lombès, M., & Dandolo, L. (2012). Double Myod and Igf2 inactivation promotes brown adipose tissue development by increasing Prdm16 expression. *FASEB Journal*. https://doi.org/10.1096/fj.12-208496

Chang, S., & Bartolomei, M. S. (2020). Modeling human epigenetic disorders in mice: Beckwith-

Wiedemann syndrome and Silver-Russell syndrome. *DMM Disease Models and Mechanisms*.

https://doi.org/10.1242/dmm.044123

Chen, C. P., Chang, S. Y., Lin, C. J., Chern, S. R., Wu, P. S., Chen, S. W., … Wang, W. (2018). Prenatal

diagnosis of a familial 5p14.3-p14.1 deletion encompassing CDH18, CDH12, PMCHL1, PRDM9

and CDH10 in a fetus with congenital heart disease on prenatal ultrasound. *Taiwanese Journal of*

Obstetrics and Gynecology, *57*(5). https://doi.org/10.1016/j.tjog.2018.08.023

Choong, O. K., Chen, C. Y., Zhang, J., Lin, J. H., Lin, P. J., Ruan, S. C., … Hsieh, P. C. H. (2019).

Hypoxia-induced H19/YB-1 cascade modulates cardiac remodeling after infarction. *Theranostics*,

9(22). https://doi.org/10.7150/thno.35218

Conigliaro, A., Costa, V., Lo Dico, A., Saieva, L., Buccheri, S., Dieli, F., … Alessandro, R. (2015).

CD90+ liver cancer cells modulate endothelial cell phenotype through the release of exosomes

containing H19 lncRNA. *Molecular Cancer*, *14*(1), 1–11. https://doi.org/10.1186/s12943-015-0426-

x

Constância, M., Hemberger, M., Hughes, J., Dean, W., Ferguson-Smith, A., Fundele, R., … Reik, W.

(2002). Placental-specific IGF-II is a major modulator of placental and fetal growth. *Nature*,

417(6892), 945–948. https://doi.org/10.1038/nature00819

DeChiara, T. M., Robertson, E. J., & Efstratiadis, A. (1991). Parental imprinting of the mouse insulin-like

growth factor II gene. *Cell*. https://doi.org/10.1016/0092-8674(91)90513-X

Dor, Y., Camenisch, T. D., Itin, A., Fishman, G. I., McDonald, J. A., Carmeliet, P., & Keshet, E. (2001).

A novel role for VEGF in endocardial cushion formation and its potential contribution to congenital

heart defects. *Development*, *128*(9). https://doi.org/10.1242/dev.128.9.1531

Drewell, R. A., Brenton, J. D., Ainscough, J. F. X., Barton, S. C., Hilton, K. J., Arney, K. L., … Surani,

M. A. (2000). Deletion of a silencer element disrupts H19 imprinting independently of a DNA

methylation epigenetic switch. *Development*.

Eggermann, T., Buiting, K., & Temple, I. K. (2011). Clinical utility gene card for: Silver-Russell

syndrome. *European Journal of Human Genetics*. https://doi.org/10.1038/ejhg.2010.202

Eisenberg, L. M., & Markwald, R. R. (1995). Molecular regulation of atrioventricular valvuloseptal

morphogenesis. *Circulation Research*. https://doi.org/10.1161/01.RES.77.1.1

Engel, N., West, A. G., Felsenfeld, G., & Bartolomei, M. S. (2004). Antagonism between DNA

hypermethylation and enhancer-blocking activity at the H19 DMD is uncovered by CpG mutations.

Nature Genetics. https://doi.org/10.1038/ng1399

Freschi, A., Del Prete, R., Pignata, L., Cecere, F., Manfrevola, F., Mattia, M., … Cerrato, F. (2021). The

number of the CTCF binding sites of the H19/IGF2:IG-DMR correlates with DNA methylation and

expression imprinting in a humanized mouse model. *Human Molecular Genetics*, *30*(16).

https://doi.org/10.1093/hmg/ddab132

Freschi, A., Hur, S. K., Valente, F. M., Ideraabdullah, F. Y., Sparago, A., Gentile, M. T., … Cerrato, F.

(2018). Tissue-specific and mosaic imprinting defects underlie opposite congenital growth disorders

in mice. *PLoS Genetics*. https://doi.org/10.1371/journal.pgen.1007243

Gabory, A., Jammes, H., & Dandolo, L. (2010). The H19 locus: Role of an imprinted non-coding RNA in

growth and development. *BioEssays*. https://doi.org/10.1002/bies.200900170

Gabory, A., Ripoche, M. A., Le Digarcher, A., Watrin, F., Ziyyat, A., Forné, T., … Dandolo, L. (2009).

H19 acts as a trans regulator of the imprinted gene network controlling growth in mice.

Development. https://doi.org/10.1242/dev.036061

García-Padilla, C., Domínguez, J. N., Aránega, A. E., & Franco, D. (2019). Differential chamber-specific

expression and regulation of long non-coding RNAs during cardiac development. *Biochimica et*

Biophysica Acta - Gene Regulatory Mechanisms, *1862*(10), 194435.

https://doi.org/10.1016/j.bbagrm.2019.194435

Ghanim, M., Rossignol, S., Delobel, B., Irving, M., Miller, O., Devisme, L., … Vincent-Delorme, C.

(2013). Possible association between complex congenital heart defects and 11p15 hypomethylation

in three patients with severe Silver-Russell syndrome. *American Journal of Medical Genetics, Part*

A, *161*(3), 572–577. https://doi.org/10.1002/ajmg.a.35691

Gicquel, C., Rossignol, S., Cabrol, S., Houang, M., Steunou, V., Barbu, V., … Le Bouc, Y. (2005).

Epimutation of the telomeric imprinting center region on chromosome 11p15 in Silver-Russell

syndrome. *Nature Genetics*, *37*(9), 1003–1007. https://doi.org/10.1038/ng1629

Greco, S., Zaccagnini, G., Perfetti, A., Fuschi, P., Valaperta, R., Voellenkle, C., … Martelli, F. (2016).

Long noncoding RNA dysregulation in ischemic heart failure. *Journal of Translational Medicine*,

14(1), 1–14. https://doi.org/10.1186/s12967-016-0926-5

Haley, V. L., Barnes, D. J., Sandovici, I., Constancia, M., Graham, C. F., Pezzella, F., … Hassan, A. B.

(2012). Igf2 pathway dependency of the Trp53 developmental and tumour phenotypes. *EMBO*

Molecular Medicine. https://doi.org/10.1002/emmm.201101105

Han, L., Szabó, P. E., & Mann, J. R. (2010). Postnatal survival of mice with maternal duplication of distal

chromosome 7 induced by a Igf2/H19 imprinting control region lacking insulator function. *PLoS*

Genetics. https://doi.org/10.1371/journal.pgen.1000803

Harris, L. K., & Westwood, M. (2012). Biology and significance of signalling pathways activated by

IGF-II. *Growth Factors*, *30*(1), 1–12. https://doi.org/10.3109/08977194.2011.640325

Hobuß, L., Foinquinos, A., Jung, M., Kenneweg, F., Xiao, K., Wang, Y., … Thum, T. (2020). Pleiotropic

cardiac functions controlled by ischemia-induced lncRNA H19. *Journal of Molecular and Cellular*

Cardiology, *146*(July 2019), 43–59. https://doi.org/10.1016/j.yjmcc.2020.07.001

Hubert, F., Payan, S. M., & Rochais, F. (2018). FGF10 Signaling in Heart Development, Homeostasis,

Disease and Repair. *Frontiers in Genetics*. https://doi.org/10.3389/fgene.2018.00599

Hur, S. K., Freschi, A., Ideraabdullah, F., Thorvaldsen, J. L., Luense, L. J., Weller, A. H., … Bartolomei,

M. S. (2016). Humanized H19/Igf2 locus reveals diverged imprinting mechanism between mouse

and human and reflects Silver-Russell syndrome phenotypes. *Proceedings of the National Academy*

of Sciences of the United States of America. https://doi.org/10.1073/pnas.1603066113

Keniry, A., Oxley, D., Monnier, P., Kyba, M., Dandolo, L., Smits, G., & Reik, W. (2012). The H19

lincRNA is a developmental reservoir of miR-675 that suppresses growth and Igf1r. *Nature Cell*

Biology. https://doi.org/10.1038/ncb2521

Kochilas, L. K., Li, J., Jin, F., Buck, C. A., & Epstein, J. A. (1999). p57Kip2expression is enhanced

during mid-cardiac murine development and is restricted to trabecular myocardium. *Pediatric*

Research, *45*(5). https://doi.org/10.1203/00006450-199905010-00004

Kruithof, B. P. T., Krawitz, S. A., & Gaussin, V. (2007). Atrioventricular valve development during late

embryonic and postnatal stages involves condensation and extracellular matrix remodeling.

Developmental Biology, *302*(1). https://doi.org/10.1016/j.ydbio.2006.09.024

Lee, J. H., Gao, C., Peng, G., Greer, C., Ren, S., Wang, Y., & Xiao, X. (2011). Analysis of transcriptome

complexity through RNA sequencing in normal and failing murine hearts. *Circulation Research*,

109(12), 1332–1341. https://doi.org/10.1161/CIRCRESAHA.111.249433

Li, P., Cavallero, S., Gu, Y., Chen, T. H. P., Hughes, J., Hassan, A. B., … Sucov, H. M. (2011). IGF

signaling directs ventricular cardiomyocyte proliferation during embryonic heart development.

Development, *138*(9), 1795–1805. https://doi.org/10.1242/dev.054338

Liao, J., Zeng, T. B., Pierce, N., Tran, D. A., Singh, P., Mann, J. R., & Szabó, P. E. (2021). Prenatal

correction of IGF2 to rescue the growth phenotypes in mouse models of Beckwith-Wiedemann and

Silver-Russell syndromes. *Cell Reports*, *34*(6). https://doi.org/10.1016/j.celrep.2021.108729

Marsh, B., & Blelloch, R. (2020). Single nuclei RNA-seq of mouse placental labyrinth development.

ELife, *9*. https://doi.org/10.7554/eLife.60266

Maslen, C. L. (2018). Recent advances in placenta-heart interactions. *Frontiers in Physiology*, *9*(JUN), 1–

9. https://doi.org/10.3389/fphys.2018.00735

Matthiesen, N. B., Henriksen, T. B., Agergaard, P., Gaynor, J. W., Bach, C. C., Hjortdal, V. E., &

Østergaard, J. R. (2016). Congenital Heart Defects and Indices of Placental and Fetal Growth in a

Nationwide Study of 924 422 Liveborn Infants. *Circulation*, *134*(20), 1546–1556.

https://doi.org/10.1161/CIRCULATIONAHA.116.021793

Mi, H., Muruganujan, A., Ebert, D., Huang, X., & Thomas, P. D. (2019). PANTHER version 14: More

genomes, a new PANTHER GO-slim and improvements in enrichment analysis tools. *Nucleic Acids*

Research, *47*(D1). https://doi.org/10.1093/nar/gky1038

Mootha, V. K., Lindgren, C. M., Eriksson, K. F., Subramanian, A., Sihag, S., Lehar, J., … Groop, L. C.

588 (2003). PGC-1α-responsive genes involved in oxidative phosphorylation are coordinately

589 downregulated in human diabetes. *Nature Genetics*, *34*(3). https://doi.org/10.1038/ng1180

590 Penny, D. J., & Vick, G. W. (2011). Ventricular septal defect. *The Lancet*, *377*(9771), 1103–1112.

591 https://doi.org/10.1016/S0140-6736(10)61339-6

592 Perez-Garcia, V., Fineberg, E., Wilson, R., Murray, A., Mazzeo, C. I., Tudor, C., … Hemberger, M.

593 (2018). Placentation defects are highly prevalent in embryonic lethal mouse mutants. *Nature*,

594 *555*(7697), 463–468. https://doi.org/10.1038/nature26002

595 Poirier, F., Chan, C. T. J., Timmons, P. M., Robertson, E. J., Evans, M. J., & Rigby, P. W. J. (1991). The

596 murine H19 gene is activated during embryonic stem cell differentiation in vitro and at the time of

597 implantation in the developing embryo. *Development*, *113*(4).

598 https://doi.org/10.1242/dev.113.4.1105

599 Rochais, F., Sturny, R., Chao, C. M., Mesbah, K., Bennett, M., Mohun, T. J., … Kelly, R. G. (2014).

600 FGF10 promotes regional foetal cardiomyocyte proliferation and adult cardiomyocyte cell-cycle re-

601 entry. *Cardiovascular Research*, *104*(3). https://doi.org/10.1093/cvr/cvu232

602 Rychik, J., Goff, D., McKay, E., Mott, A., Tian, Z., Licht, D. J., & Gaynor, J. W. (2018). Characterization

603 of the Placenta in the Newborn with Congenital Heart Disease: Distinctions Based on Type of

604 Cardiac Malformation. *Pediatric Cardiology*, *39*(6), 1165–1171. https://doi.org/10.1007/s00246-

605 018-1876-x

606 Sandovici, I., Georgopoulou, A., Pérez-García, V., Hufnagel, A., López-Tello, J., Lam, B. Y. H., …

607 Constância, M. (2022). The imprinted Igf2-Igf2r axis is critical for matching placental

608 microvasculature expansion to fetal growth. *Developmental Cell*, *57*(1), 63-79.e8.

609 https://doi.org/10.1016/j.devcel.2021.12.005

610 SanMiguel, J. M., Abramowitz, L. K., & Bartolomei, M. S. (2018). Imprinted gene dysregulation in a

611 Tet1 null mouse model is stochastic and variable in the germline and offspring. *Development*

612 *(Cambridge)*, *145*(7). https://doi.org/10.1242/dev.160622

613 Savolainen, S. M., Foley, J. F., & Elmore, S. A. (2009). Histology atlas of the developing mouse heart

with emphasis on E11.5 to E18.5. *Toxicologic Pathology*, *37*(4), 395–414.

    https://doi.org/10.1177/0192623309335060

Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T., … Cardona, A.

    (2012). Fiji: An open-source platform for biological-image analysis. *Nature Methods*.

    https://doi.org/10.1038/nmeth.2019

Serra, A., Eirich, K., Winkler, A. K., Mrasek, K., Ghring, G., Barbi, G., … Schindler, D. (2012). Shared

    copy number variation in simultaneous nephroblastoma and neuroblastoma due to Fanconi anemia.

    Molecular Syndromology, *3*(3), 120–130. https://doi.org/10.1159/000341935

Shen, H., Cavallero, S., Estrada, K. D., Sandovici, I., Kumar, S. R., Makita, T., … Sucov, H. M. (2015).

    Extracardiac control of embryonic cardiomyocyte proliferation and ventricular wall expansion.

    Cardiovascular Research, *105*(3), 271–278. https://doi.org/10.1093/cvr/cvu269

Shen, H., Gan, P., Wang, K., Darehzereshki, A., Wang, K., Ram Kumar, S., … Sucov, H. M. (2020).

    Mononuclear diploid cardiomyocytes support neonatal mouse heart regeneration in response to

    paracrine IGF2 signaling. *ELife*, *9*, 1–24. https://doi.org/10.7554/eLife.53071

Sibley, C. P., Coan, P. M., Dean, W., Hughes, J., Smith, P., Reik, W., & Burton, G. J. (2004). Regulates

    the Diffusional Exchange Characteristics of the Mouse Placenta, *2*.

Snider, P., & Conway, S. J. (2011). *Probing human cardiovascular congenital disease using transgenic*

    mouse models. *Progress in Molecular Biology and Translational Science* (1st ed., Vol. 100).

    Elsevier Inc. https://doi.org/10.1016/B978-0-12-384878-9.00003-0

Soellner, L., Kraft, F., Sauer, S., Begemann, M., Kurth, I., Elbracht, M., & Eggermann, T. (2019). Search

    for cis-acting factors and maternal effect variants in Silver-Russell patients with ICR1

    hypomethylation and their mothers. *European Journal of Human Genetics*, *27*(1).

    https://doi.org/10.1038/s41431-018-0269-1

Soemedi, R., Wilson, I. J., Bentham, J., Darlay, R., Töpf, A., Zelenika, D., … Keavney, B. D. (2012).

    Contribution of global rare copy-number variants to the risk of sporadic congenital heart disease.

    American Journal of Human Genetics, *91*(3). https://doi.org/10.1016/j.ajhg.2012.08.003

Spicer, D. E., Hsu, H. H., Co-Vu, J., Anderson, R. H., & Fricker, F. J. (2014). Ventricular septal defect.

Orphanet Journal of Rare Diseases, *9*, 144. https://doi.org/10.1186/s13023-014-0144-2

Subramanian, A., Tamayo, P., Mootha, V. K., Mukherjee, S., Ebert, B. L., Gillette, M. A., … Mesirov, J.

P. (2005). Gene set enrichment analysis: A knowledge-based approach for interpreting genome-wide

expression profiles. *Proceedings of the National Academy of Sciences of the United States of*

America, *102*(43). https://doi.org/10.1073/pnas.0506580102

Sullivan, K. E., & Black, L. D. (2013). The role of cardiac fibroblasts in extracellular matrix-mediated

signaling during normal and pathological cardiac development. *Journal of Biomechanical*

Engineering, *135*(7). https://doi.org/10.1115/1.4024349

Thorvaldsen, J. L., Fedoriw, A. M., Nguyen, S., & Bartolomei, M. S. (2006). Developmental Profile of

H19 Differentially Methylated Domain (DMD) Deletion Alleles Reveals Multiple Roles of the

DMD in Regulating Allelic Expression and DNA Methylation at the Imprinted H19/Igf2 Locus.

Molecular and Cellular Biology. https://doi.org/10.1128/mcb.26.4.1245-1258.2006

Thorvaldsen, J. L., Mann, M. R. W., Nwoko, O., Duran, K. L., & Bartolomei, M. S. (2002). Analysis of

Sequence Upstream of the Endogenous H19 Gene Reveals Elements Both Essential and Dispensable

for Imprinting. *Molecular and Cellular Biology*. https://doi.org/10.1128/mcb.22.8.2450-2462.2002

Ueno, M., Lee, L. K., Chhabra, A., Kim, Y. J., Sasidharan, R., VanHandel, B., … Mikkola, H. K. A.

(2013). C-Met-Dependent Multipotent Labyrinth Trophoblast Progenitors Establish Placental

Exchange Interface. *Developmental Cell*, *27*(4), 373–386.

https://doi.org/10.1016/j.devcel.2013.10.019

Vega-Hernández, M., Kovacs, A., de Langhe, S., & Ornitz, D. M. (2011). FGF10/FGFR2b signaling is

essential for cardiac fibroblast development and growth of the myocardium. *Development*, *138*(15).

https://doi.org/10.1242/dev.064410

Von Gise, A., & Pu, W. T. (2012). Endocardial and epicardial epithelial to mesenchymal transitions in

heart development and disease. *Circulation Research*, *110*(12), 1628–1645.

https://doi.org/10.1161/CIRCRESAHA.111.259960

Vrooman, L. A., Rhon-Calderon, E. A., Chao, O. Y., Nguyen, D. K., Narapareddy, L., Dahiya, A. K., …

Bartolomei, M. S. (2020). Assisted reproductive technologies induce temporally specific placental

defects and the preeclampsia risk marker sFLT1 in mouse. *Development (Cambridge, England)*,

147(11). https://doi.org/10.1242/dev.186551

Wakeling, E. L., Brioude, F., Lokulo-Sodipe, O., O'Connell, S. M., Salem, J., Bliek, J., … Netchine, I.

(2017). Diagnosis and management of Silver-Russell syndrome: First international consensus

statement. *Nature Reviews Endocrinology*, *13*(2), 105–124. https://doi.org/10.1038/nrendo.2016.138

Wang, Y., Sun, X., & Sun, X. (2021). The Functions of LncRNA H19 in the Heart. *Heart Lung and*

Circulation. https://doi.org/10.1016/j.hlc.2021.10.022

Woods, L., Perez-Garcia, V., & Hemberger, M. (2018). Regulation of Placental Development and Its

Impact on Fetal Growth—New Insights From Mouse Models. *Frontiers in Endocrinology*,

9(September), 1–18. https://doi.org/10.3389/fendo.2018.00570

Supplemental Information

Generation of *ΔH19* allele. Two pairs of gRNA targeting the *H19* gene are listed in Supplemental Table

1. gRNA was prepared following a protocol from Yang et al. (Yang, Wang, & Jaenisch, 2014) with

modifications. px335 plasmid (Addgene, Watertown, MA) was PCR amplified using Phusion high-

fidelity DNA polymerase (New England Biolabs, Ipswich, MA) and the primer set listed in Supplemental

Table 1. ~117bp PCR product was gel-purified using the Gel Extraction kit (Qiagen, Hilden, Germany)

according to the manufacturer's instruction. Using the gel-purified product as template, *in vitro*

transcription of gRNA was setup using T7 High Yield RNA Synthesis kit (New England Biolabs)

according to manufacturer's instructions. Transcribed gRNA was purified using the MEGAclear kit (Life

Technologies, Carlsbad, CA) according to manufacturer's instructions. 50ng/μl left and right sgRNA

together with 100ng/μl Cas9 mRNA was injected per zygote stage embryo by the Transgenic and

Chimeric Mouse Facility at the University of Pennsylvania. The targeted allele was validated using

Southern blot as previously described (Joanne L. Thorvaldsen, Duran, & Bartolomei, 1998) and PCR and

sequencing across junctions of *ΔH19* alleles. Obtained chimeras and germ line transmission animals were

PCR-genotyped for the *ΔH19* allele using primers (Supplemental Table 1).

Genotyping. Mouse genomic DNA for PCR genotyping was isolated from each animal as previously

described (SanMiguel, Abramowitz, & Bartolomei, 2018). Primers used for sex genotyping and

genotyping of *hIC1, ΔH19* and *Δ3.8* alleles are listed in Supplemental Table 1. For all genotypes, the

maternal allele is listed first and the paternal allele second.

RNA Sequencing library preparation and analysis. E12.5 hearts were lysed with Collagenase (Sigma-Aldrich), Dispase II (Sigma-Aldrich), and DNase I (Sigma-Aldrich). Cardiac endothelial cells were collected using MACS CD31 microbeads (Miltenyi Biotec, Bergisch Gladbach, Germany) and RNA was isolated using RNeasy Micro kit (Qiagen). After confirming RNA integrity using Bioanalyzer (Agilent Technologies, Santa Clara, CA), mRNA library was generated from 25ng RNA using NEBNext Poly(A) mRNA Magnetic Isolation Module and Ultra II RNA Library Prep Kit (New England Biolabs). Library quality was assessed by Bioanalyzer (Agilent Technologies) and TapeStation (Agilent Technologies). Sequencing was performed on NovaSeq 6000 (Illumina, San Diego, CA). Quality of raw fastq reads was assessed using FastQC version 0.11.5 (Andrews, Krueger, Seconds-Pichon, Biggins, & Wingett, 2015). Reads were aligned to the GRCm38/mm10 reference using STAR version 2.4.0i with default parameters and maximum fragment size of 2000 bp (Dobin et al., 2013). Properly paired primary alignments were retained for downstream analysis using Samtools version 1.9. Count matrices were generated using FeatureCounts version 1.6.2 against RefSeq gene annotation and read into DESeq2 (Love, Huber, & Anders, 2014) to perform normalization and statistical analysis.

Supplemental Figure 1. Serial cross-sections of E15.5 *+/hIC1* embryonic heart with BPV.
Sequential histological sections demonstrating the bicuspid pulmonary valve (BPV) phenotype.
Rudimentary anterior cusp can be seen fused with the right pulmonary cusp (raphe) in sections denoted
by * above the valve.

Supplemental Figure 2. Supplementary data for anomalies observed in *+/hIC1* placentas.
(A) Fetal to placental weight ratios of the wild-type (blue) and *+/hIC1* (red) samples at E11.5, E12.5,
E14.5 and E15.5. Each data point represents an average ratio of each genotype from one litter. (B)
Example of labyrinth zones that were used to quantify the microvessel density in CD34 immunostained
E17.5 wild-type and *+/hIC1* placental sections from Figure 3G. Thrombi in *+/hIC1* placentas were
excluded. Scale bars = 1mm. Statistics used are (A) multiple unpaired t-tests; ns = not significant.

Supplemental Figure 3. Characterization of *ΔH19* allele.
(A) Targeting strategy to generate the *ΔH19* allele. The endogenous *H19* locus is shown with restriction
sites, gRNA locations used to generate the deletion and binding sites for probes used in Southern blot
analysis (thick lines). (B) Southern blot analysis of *ΔH19* allele. Founder 4131 line was generated using

gRNA pair A, and founder 4133 line was generated using gRNA pair B (Supplemental Table 1). 3' probe

and 5' probe were hybridized to XbaI- and KpnI-digested DNA, respectively. The sizes of the DNA

fragments are shown on the right. (C) Allele-specific *Igf2* expression in wild-type and *ΔH19/+* neonatal

tongue, heart and liver analyzed by restriction fragment length polymorphism (RFLP). Ladder, genotypes

and c (*Mus castaneus*, paternal) and b (C57BL/6, maternal) allele controls are indicated above each gel.

Quantification of band densitometry is shown below each gel, with percent of maternal allele expression

relative to paternal allele indicated. No expression from the maternal *Igf2* allele was observed in liver. (D)

Neonatal body weight of *ΔH19/+* and *+/ΔH19* pups at postnatal day 0 (mean ± SD). Unpaired Student's t-

test; ****$P < 0.0001$, ns = not significant.

Supplemental Figure 4. Supplementary data for rescue upon maternal *ΔH19* transmission.

(A) Ratio of wild-type, *ΔH19/+, +/hIC1* and *ΔH19/hIC1* embryos observed in E11.5 and E17.5 litters (>5

pups). Each data point represents one litter. (B) (Left) Relative placental weights of E17.5 wild-type,

ΔH19/+, +/hIC1 and *ΔH19/hIC1* samples, normalized to the average placental weight of the wild-type

littermates (mean ± SD). (Right) Fetal to placental weight ratio in E17.5 wild-type, *ΔH19/+, +/hIC1* and

ΔH19/hIC1 samples (mean ± SD). (C) Relative total expression of *H19* and *Igf2* in E17.5 wild-type,

ΔH19/+, +/hIC1 and *ΔH19/hIC1* liver and tongue samples (mean ± SD). (B, C) Each data point

represents an individual conceptus from different litters. One-way ANOVA with Tukey's multiple

comparisons test; *$P < 0.05$, **$P < 0.01$, ***$P < 0.001$, ****$P < 0.0001$, ns = not significant.

Supplemental Figure 5. Supplementary data for rescue upon maternal *Δ3.8* transmission.

(A) (Left) Relative placental weights of E17.5 wild-type, *Δ3.8/+, +/hIC1* and *Δ3.8/hIC1* samples,

normalized to the average placental weight of the wild-type littermates (mean ± SD). (Right) Fetal to

placental weight ratio in E17.5 wild-type, *Δ3.8/+, +/hIC1* and *Δ3.8/hIC1* samples (mean ± SD). (B)

Relative total expression of *H19* and *Igf2* in E17.5 wild-type, *Δ3.8/+, +/hIC1* and *Δ3.8/hIC1* liver and

tongue samples (mean ± SD). (C) Junctional zone (Jz) to labyrinth (Lb) ratio in E17.5 wild-type, *Δ3.8/+,*

+/hIC1 and *Δ3.8/hIC1* placentas. (A, B, C) Each data point represents an individual conceptus from

different litters. One-way ANOVA with Tukey's multiple comparisons test; *$P < 0.05$, **$P < 0.01$, ***$P <$

0.001, ****$P < 0.0001$, ns = not significant.

Supplemental Figure 6. Supplementary data for differential gene expression of *+/hIC1* hearts.

(A) DAPI (blue) on E17.5 wild-type and *+/hIC1* hearts from Figure 6C. Images on the right are enlarged

from the boxed area of images on the left. Scale bars = 100 μm. (B) Immunofluorescence staining for

MF20 (red), collagen I (green), and DAPI (blue) on E17.5 *+/hIC1* hearts. Images on the right are enlarged

from the boxed area of images on the left. Scale Bars = 100 μm. (C) Same-sex comparisons between

+/hIC1 and the wild-type samples analyzed by RNA sequencing.

Supplemental Figure 7. (A) Volcano plot depicting comparison of *ΔH19/hIC1* and wild-type cardiac endothelial cells. (B) Volcano plot depicting comparison of *+/hIC1* and *ΔH19/hIC1* cardiac endothelial cells. (C) GO pathways that are enriched for 46 DEGs between *+/hIC1* and *ΔH19/hIC1* samples.

Supplemental Figure 8. (A) Volcano plot depicting comparison of *+/hIC1* and *Δ3.8/hIC1* cardiac endothelial cells. (B) GO pathways that are enriched for 116 DEGs that are commonly differentially expressed in *+/hIC1* compared to wild-type and *Δ3.8/hIC1* samples. (C) Volcano plot depicting comparison of *Δ3.8/+* and wild-type cardiac endothelial cells.

Supplemental Figure 9. Enrichment tests for BRIDEAU_IMPRINTED_GENES using GSEA.
Normalized enrichment score (NES), nominal p-value, enrichment plot and expression heatmap of the analyzed gene set comparing (A) *+/hIC1* and wild-type, (B) *ΔH19/+* and wild-type*,* (C) *ΔH19/hIC1* and wild-type cardiac endothelial cells.

Supplemental References

Andrews, S., Krueger, F., Seconds-Pichon, A., Biggins, F., & Wingett, S. (2015). FastQC. A quality control tool for high throughput sequence data. Babraham Bioinformatics.

de Waal, E., Vrooman, L. A., Fischer, E., Ord, T., Mainigi, M. A., Coutifaris, C., … Bartolomei, M. S. (2015). The cumulative effect of assisted reproduction procedures on placental development and epigenetic perturbations in a mouse model. *Human Molecular Genetics*. https://doi.org/10.1093/hmg/ddv400

Dobin, A., Davis, C. A., Schlesinger, F., Drenkow, J., Zaleski, C., Jha, S., … Gingeras, T. R. (2013). STAR: Ultrafast universal RNA-seq aligner. *Bioinformatics*, *29*(1). https://doi.org/10.1093/bioinformatics/bts635

Hur, S. K., Freschi, A., Ideraabdullah, F., Thorvaldsen, J. L., Luense, L. J., Weller, A. H., … Bartolomei, M. S. (2016). Humanized H19/Igf2 locus reveals diverged imprinting mechanism between mouse and human and reflects Silver-Russell syndrome phenotypes. *Proceedings of the National Academy of Sciences of the United States of America*. https://doi.org/10.1073/pnas.1603066113

795   Love, M. I., Huber, W., & Anders, S. (2014). Moderated estimation of fold change and dispersion for

796   RNA-seq data with DESeq2. *Genome Biology*, *15*(12). https://doi.org/10.1186/s13059-014-0550-8

797   Plasschaert, R. N., & Bartolomei, M. S. (2014). Genomic imprinting in development, growth, behavior

798   and stem cells. *Development (Cambridge)*. https://doi.org/10.1242/dev.101428

799   SanMiguel, J. M., Abramowitz, L. K., & Bartolomei, M. S. (2018). Imprinted gene dysregulation in a

800   Tet1 null mouse model is stochastic and variable in the germline and offspring. *Development*

801   *(Cambridge)*, *145*(7). https://doi.org/10.1242/dev.160622

802   Thorvaldsen, J. L., Mann, M. R. W., Nwoko, O., Duran, K. L., & Bartolomei, M. S. (2002). Analysis of

803   Sequence Upstream of the Endogenous H19 Gene Reveals Elements Both Essential and Dispensable

804   for Imprinting. *Molecular and Cellular Biology*. https://doi.org/10.1128/mcb.22.8.2450-2462.2002

805   Thorvaldsen, Joanne L., Duran, K. L., & Bartolomei, M. S. (1998). Deletion of the H19 differentially

806   methylated domain results in loss of imprinted expression of H19 and Igf2. *Genes and Development*.

807   https://doi.org/10.1101/gad.12.23.3693

808   Weaver, J. R., Susiarjo, M., & Bartolomei, M. S. (2009). Imprinting and epigenetic changes in the early

809   embryo. *Mammalian Genome*. https://doi.org/10.1007/s00335-009-9225-2

810   Yang, H., Wang, H., & Jaenisch, R. (2014). Generating genetically modified mice using CRISPR/Cas-

811   mediated genome engineering. *Nature Protocols*, *9*(8). https://doi.org/10.1038/nprot.2014.134