Health hazards posed by ionizing radiation to human space missions and newly developing methodologies to investigate them

Franco Ferrari and Ewa Szuszkiewicz

CASA* and Institute of Physics, University of Szczecin, Wielkopolska 15, 70-453 Szczecin, Poland

Abstract

Ionizing radiation is one of the main threats to human space exploration beyond low Earth orbit (BLEO). It is thus of primary importance to determine safe career dose limits for astronauts involved in BLEO missions. In this work the average equivalent doses received by a hypothetical human crew are established using the data of several robotic missions to the Moon and to Mars. The probabilities of the occurrence of deterministic effects due to radiation that could impair the success of a mission or lower the life expectancy of astronauts are evaluated using statistical methods. The limits of the obtained results concerning deterministic and stochastic effects basing on statistical methods are discussed. The conclusion is that new strategies are necessary in order to predict the insurgence of stochastic effects. We argue that the use of the so-called 3D models or organoids could be a good candidate of such a strategy that will be complementary to in vitro 2D models and animal models. Organoids are in fact able to recapitulate several relevant features of the real human organs and their in vivo surroundings. Yet, the main problem of detecting the onset of cancer in the human body remains outstanding. To this purpose, we propose here to combine organoid techniques with those of precision oncology and molecular medicine. Exactly these disciplines have greatly contributed in the recent time to the understanding of the patterns of cancer formation in several cases of tumors. Precision oncology and molecular medicine provide powerful markers to track down the events leading to tumorigenesis following exposure to radiation in organoids.

1   Introduction

The risks posed by energetic particle radiation to human crews implementing exploration missions in *Low Earth Orbit*/LEO and ***Beyond Low Earth Orbit/***BLEO are of a serious nature. For this reason, both the short term and long-term effects of irradiation in space should be taken into account when establishing safe career dose limits for astronauts, see McKenna-Lawlor et al. (2015) and Walsh et al. (2019).

In McKenna-Lawlor et al. (2015) it was pointed out that an important step toward the mitigation of human health risks in BLEO would be to gain a better understanding of, those flare related, 'hard' ***Solar Energetic Particle***/SEP) events which can significantly increase the severity of the doses incurred by astronauts. The latter hazard could be ameliorated were it to become possible to predict the occurrences of these events.

An obstacle in estimating for astronauts the maximum acceptable dose accrued in interplanetary space below which the risk of contracting cancer or other pathologies is not higher than the risk incurred by workers on Earth dealing with sources of ionizing radiation, is the lack of sufficient statistics concerning the effects of high LET radiation on human space travelers. Hazardous radiation (including energetic protons and heavy ions) is present outside the region protected by the geomagnetic field (i.e. outside the Earth's magnetosphere) and, hitherto, data concerning the effects of this ionizing radiation on living organisms in BLEO has not been available - apart from

measurements made during the Apollo lunar exploration missions of the 1960s and 1970s, see for example English et al., (1973). An experiment that is expected to quantify the level of biological damage incurred in space, as well as to validate existing models, is currently scheduled to be conducted aboard NASA's BioSentinel nanosatellite, utilizing a payload scheduled for launch in February 2022.

In the present work we provide a comprehensive view on the problem of determining the health hazards posed by ionizing radiation to human space missions to objects beyond the low Earth orbit. The main goal is to establish safe career dose limits for astronauts involved in such missions. The starting point is to estimate the equivalent doses that are expected to be received by an hypothetical crew during the travel to the Moon or to Mars and a short stay on these objects for exploration purposes. In doing that, it is of critical importance to evaluate not only doses and dose rates, but also the species and energy distributions of the particles composing the ionizing radiation.

In **Low Earth Orbits**/LEO these data have already been extensively measured aboard robotic spacecrafts or the International Space Station and are well known. In the case of BLEO missions, a satisfactory amount of relevant data is also available as a legacy of many robotic missions flown in the inner and outer Solar System. A review of doses, particle content and energy distributions of SEPs measured in space is contained in McKenna-Lawlor et al. (2015) and references therein. In the case of measured **Galactic Cosmic Radiation**/GCR, Ferrari & Szuszkiewicz, (2009) and Mrigakshi et al. (2012) contain useful accounts and references.

The analysis of data recorded by the RAD detector aboard the *Mars Science* Laboratory/MSL while on route to Mars, showed (Aghata et al. (2009)) that the inclusion in the calculations of ambient radiation due to pions and their decay products makes an important difference when estimating the dose present behind spacecraft shielding. Information on relevant pion transport codes is available in Sato et al. (2011) and Norman et al. (2012).

The second step of the present research consists in evaluating the risks related to the so-called deterministic effects that may occur in a typical mission to the Moon or Mars. To this aim we use the formulas derived in Edwards & Lloyd (1998), where sophisticated statistical methods have been applied to the analysis of events like the bombing of the Japanese cities of Nagasaki and Hiroshima or accidents involving radioactive substances. It is important to keep in mind that the parameters characterizing the exposure to radiation of the human population in such events are quite different from those describing the exposure in BLEO missions. For that reason, the coefficients computed for various types of deterministic effects in Edwards & Lloyd (1998) are not entirely suitable in the case of space explorations. Nonetheless, it turns out that during a BLEO mission the risks of the appearance of any of the deterministic effects considered in Edwards & Lloyd (1998) are so low, that they can be neglected even taking into account a large indetermination in the coefficients. This is not enough however to rule out almost all deterministic effects. There could be in fact exceptions, as for instance the light flashing that has been perceived by astronauts during the Apollo missions and is certainly related to ionizing radiation. Another effect that could interfere with the course of a mission is the risk of cardiac arrhythmia, see Stoupel et al. (2006).

The risks related to the insurgence of stochastic effects like cancer are much more difficult to be estimated. This problem is tackled in the literature using a two-fold strategy, epidemiologic studies and experiments, in vitro or in vivo. Here, we limit ourselves to a brief review of the basic statistical methods that help to determine the probability of cancer formation. In particular, basing ourselves on the excess relative risk (ERR) for the induction of fatal cancer computed from the data of the Hiroshima and Nagasaki bombings, we estrapolate the ERR for a typical mission on Mars. These rough estimations can be made more precise, but also the most refined evaluations of the risks related to ionizing radiation are still not enough to determine safe career dose limits for astronauts

participating in BLEO missions. Further studies are necessary in which alternative methods are exploited and new strategies should be found. Next, we briefly report here some of the conclusions coming from animal models and in vitro measurements. In doing that, we follow Ref. Shridharan et al. (2015), in which an extensive review is presented on the state-of-the-art of both in vitro and in vivo experiments attempting to assess the effects of ionizing radiation in space. Additionally, we suggest an approach based on the so-called 3D in vitro models. So far, these models, in which the human organs are simulated by chimera organoids, have been widely applied in medicine in order to investigate new methods for curing cancer. Moreover, they have also been used to track down the initial steps leading to tumorigenesis in individuals affected by particular mutations. We propose to adopt a similar strategy to study the onset of cancer following the exposure to ionizing radiation, especially in the case of radiation containing heavy nuclei accelerated at high energies. This kind of radiation is in fact one of the major challenges to the human exploration of space, see for instance Chancellor et al. (2014). The idea of investigating the effects of radiation with the help of 3D models is not entirely new, it has been mentioned for instance in the already cited work of Shridharan et al., (2015). Up to now, however, the follow up on this idea has been little, probably because its realization is challenging. Certainly, strategies based on miniorgans (organoids) present several advantages, like for example the possibility of studying directly human cells avoiding the limitations of animal models in this respect. Moreover, organoids are now able to reproduce many of the characteristics of the original organs and even of the surrounding epithelium and supporting mesenchymal stroma. The challenge comes from the fact that, even weakening organoids by mutations or by infecting them with bacteria in such a way that they will be more likely to develop cancer, the probability of tumor induction following irradiation could be still too low to be observed. Besides, cancer is a multistep process that in humans can last for decades. We are however convinced that these difficulties may be overcome by using the recent results of precision oncology and molecular medicine, in which the genetic and epigenetic changes leading to genomic instability and cancer formation have been pinpointed for several types of tumors. The revolution in understanding tumorigenesis in humans introduced by molecular medicine is illustrated here using the example of colorectal cancer, which has been extensively investigated in the recent past. We argue that some of the effects of radiation discussed in Shridharan et al., (2015) have a deeper interpretation in the context of precision oncology. At the end, we propose a possible strategy to establish safe career dose limits for astronauts based on the use of 3D models and the knowledge coming from molecular medicine as an useful complement of in vitro 2D studies and animal models.

2   Three case studies: Missions to the Moon, Mars and asteroids.

2.1   Doses and particle composition of the ionizing radiation in the case of missions to the Moon

Our knowledge on the effects of radiation in BLEO is good only in the case of the exploration of the Moon because of the extensive experience made during the Apollo missions, see Vernon Bailey (1975). The particle composition in this case included electrons, protons, neutrons, alpha particles and heavy ions. The radiation sources were provided by: particles trapped in the Van Allen belts (mainly energetic electrons and protons); Galactic Cosmic Rays (GCR); Solar Energetic Particles (SEPs) and secondary radiation created by the interaction of the primary particles with the spacecraft hull. According to Vernon Bailey (1975), the doses received by the crews of the Apollo missions were "significantly lower than the yearly average of 5 rem" (50 mSv) "set by the U.S. Atomic Energy Commission for workers" exposed to radioactive sources. The reason of these low levels is due to the fact that during Apollo missions 7 – 15 no significant solar event associated with the emission of SEPs took place. The next two Apollo missions (16 –17) flew when the solar

activity was at a minimum and the likelihood of large SEP events was accordingly low. It is recalled that Apollo 17 was the last human BLEO mission to be launched up to the year 2021. In all Apollo missions 7 – 17, the main source of ionizing radiation was Galactic Cosmic Radiation, in short GCR, the radiation levels of which are at their maximum during solar minimum. The estimated dose absorbed by each crewman on Apollo missions 7 – 15 averaged over the period of the travel from the Earth to the Moon and during the orbit around the Moon was of the order of 10μGy/hour. In the period of ten days typical of Apollo missions to the Moon the resulting absorbed dose was of the order of 2.4mGy (Vernon Bailey, 1975). On the lunar surface this value was less, i.e. ~ 6μGy/hour (Vernon Bailey, 1975). In this case, part of the absorbed radiation was due to neutrons produced in the collision of primary GCR with the lunar soil. The absorbed doses in Apollo missions 16 – 17 were higher, i.e. about 7.7mGy (Mazur et al., 2011), because these missions occurred at a time of solar minimum. More recent measurements made by the Cosmic Ray Telescope for the Effects of Radiation (CRaTER) instrument on the Lunar Reconnaissance Orbiter (LRO) in the absence of SEP events, recorded a total absorbed dose of about 3.8 mGy in a ten days period. It is noted that the shielding of CRaTER (32 mils of aluminum, 1mil=0.0254mm) is comparable with that installed on the Apollo spacecraft (39 mils). The reasons of the discrepancy in the data of the absorbed dose measured during solar minimum between CRaTER and the Apollo missions 16 – 17 are explained in Mazur et al. (2011).

Finally, the absorbed doses that will be taken into consideration in this study with regard to missions to the Moon are the following:

1. an absorbed dose of about 10mGy in the absence of SEP events

2. an absorbed dose of 4Gy, which corresponds to the maximum operational dose (MOD) limit for skin that was set for the Apollo missions. This dose is comparable with that of 3.6Gy that were estimated to be absorbed by the skin in the case of a large solar event (Vernon Bailey, 1975).

3. an absorbed dose of 0.5Gy for the blood forming organs set as a MOD limit in the Apollo missions, which is comparable to the estimated dose of 0.35Gy incurred during a large solar event (Vernon Bailey, 1975).

2.2 Doses and particle composition of the ionizing radiation in the case of missions to Mars

The doses to which a human crew will be exposed during a journey to Mars and back can be very roughly estimated thanks to the many unmanned missions to that planet. According to the data of the RAD detector on board of the Curiosity Rover, during the cruise phase to Mars of the Mars Science Laboratory mission, an average absorbed dose rate of:

$$\dot{D}_{\text{space}} = 0.48 \pm 0.08 \, \text{mGy/day} \tag{1}$$

composed of GCR (mainly protons and alpha particles), was observed (Zeitlin et al., 2013). The average quality factor of GCR radiation measured in space was taken to be:

$$Q_{\text{space}} = 3.82 \pm 0.30 \tag{2}$$

On the surface of Mars, the average absorbed dose rate amounted to

$$\dot{D}_{\text{Mars}} = 0.210 \pm 0.04 \, \text{mGy/day} \tag{3}$$

The average quality factor of the radiation on the surface (primary GCR plus neutrons and gamma rays coming from the secondary radiation) was:

$$Q_{\text{Mars}} = 3.05 \pm 0.30 \tag{4}$$

The SEP events registered during the Mars Science Laboratory mission consisted predominantly of protons for which $Q \sim 1.00$. The delivered absorbed dose ranged from 1.2mGy/event up to 19.5mGy/event. In total, five SEP events were recorded during the Cruise Phase with an absorbed dose peak of more than 2.5mGy/day. Of course, the above dose estimates could be strongly affected by the presence of large SEP events. The latter are usually rare but remain, thus far, impossible to predict, McKenna-Lawlor et al. (2015). The total equivalent dose estimated by the authors of Hassler et al. (2014) for a cruise to Mars and back of duration $180 \times 2 = 360$ days was $662 \pm 108 \, \text{mSv}$. The contribution due to a stay on the Martian surface of 500 days was $320 \pm 50 \, \text{mSv}$.

In total, this gives an equivalent dose of:

$$H_I = 982 \pm 158 \, \text{mSv} \tag{5}$$

The corresponding absorbed dose, computed taking into account the quality factors Q_{space} and Q_{Mars} of Eqs. (2) and (4) respectively is given by $D_I = 227.5 \pm 34.40 \, \text{mGy}$. Considering instead a typical Mars mission with a short surface stay (consisting of 400 days in deep space and 30 days on the surface – as described by McKenna-Lawlor et al. (2015), the equivalent dose extrapolated from the above data is:

$$H_{II} = 755 \pm 123 \, \text{mSv} \tag{6}$$

The related absorbed dose amounts to $D_{II} \sim 198.30 \pm 33.20 \, \text{mGy}$.

The total values of $982 \pm 158 \, \text{mSv}$ and of $755 \pm 123 \, \text{mSv}$ are both very high and comparable with the highest reference level recommended by ICRP for the rescue of workers on Earth in emergency situations.

3 Biological response of humans to energetic particle irradiation Beyond Low Earth Orbit/BLEO

3.1 Deterministic effects

Deterministic effects are medically diagnosable effects related to cell killing caused by the absorption of ionizing radiation. These effects appear when the absorbed dose is high – of the order of several Gy's – and affect single organs or tissues. The severity of the damage is dose dependent. In order to take into account the diversity in the response of individuals to ionizing radiation, it is possible to define lower and upper absorbed dose thresholds for the onset of a specific deterministic effect. The lower threshold D_{low} is such that the deterministic effect certainly does not appear for doses D such that $D < D_{low}$. The upper threshold represents the dose at which the cell killing is high enough that the effect becomes unavoidable when $D > D_{up}$. Of course, these thresholds depend on the way in which the dose is delivered such that their value is lower in the case of a single high dose delivered in a short time and higher when the dose is split into many smaller doses delivered over a long period. The risk R that a deterministic effect will appear in a sufficiently large sample consisting of individuals who have received a certain dose D, can be modeled using a sigmoid function. If R is defined as the percentage of individuals in which the effect becomes manifest, then the sigmoid function should approach zero when $D \sim D_{low}$ and be nearly equal to 1 if $D \sim D_{up}$. For practical purposes D_{low} can be defined as the absorbed dose for which the risk of a given effect is $R(D_{low}) = 0.01$ (Edwards & Lloyd, 1998). Analogously, D_{up} is defined as the dose for which $R(D_u) = 0.99$. A possible choice expressing the dependence of R on D is provided using a Weibull function, which is of the form (Edwards & Lloyd, 1998):

$$R(D) = 1 - \exp\left[\ln(2) \cdot \left(\frac{RBE \cdot D}{D_{50}}\right)^V\right] \tag{7}$$

The number $N(D)$ of affected individuals is given by the formula $N(D) = N_0 \cdot R(D)$ where N_0 is the total number of irradiated individuals.

The shape of the distribution in Eq. (7) is determined by three parameters:

- The absorbed dose D_{50} at which 50% of the exposed individuals are supposed to develop the given deterministic effect under study. This parameter depends on the dose rate according to the equation:

$$D_{50}(\dot{D}) = \theta_\infty + \frac{\theta_1}{RBE \cdot \dot{D}} \tag{8}$$

The first term in the right hand side of the above equation takes into account the case of acute dose rates delivered in a few minutes, while the second term which is proportional to θ_1 is the correction required by the fact that the effects of radiation vary as a function of the dose rate $\dot{D}$. It is pointed out that Eq. (8) is purely phenomenological and that there is no scientific evidence for it. The parameters θ_∞ and θ_1 are extrapolated applying sophisticated statistical methods to the existing data from nuclear accidents involving radioactive sources.

- The relative biological effectiveness factor *RBE*.

- A shape factor *V* determining the slope of the sigmoid. By adjusting *V,* it is possible to accommodate different values of the lower and upper thresholds D_{low} and D_{up}. Examples of risk-dose curves are given for various values of the parameter *V* in Figure 1. As in the case of the angles θ_{∞} and θ_1, the value of *V* is evaluated on the basis of statistical methods applied to the existing experimental data.

It is noted that the parameters $\theta_{\infty}, \theta_1$ and *V* are effect dependent.

Figure 1. Risk – dose curves computed using the sigmoid function defined in Eq. (7) for various values of the parameter *V*.

3.1.1 Deterministic effects in the case of Mars missions

Based on the above, we are now in a position to study the risk of the onset of deterministic effects during a space mission and discuss first the case of missions to Mars, with the goal to estimate the risk of occurrence of some deterministic effect using Eq. (7). Of course, it should be taken into account that this equation only provides indicative results. As pointed out in Cucinotta & Durante, (2006), there are not enough statistics available regarding the action of ionizing radiation of space

origin in producing deterministic and stochastic effects in the human body. Formulae that attempt to extrapolate the risk of mortal or morbidity effects based on data coming from accidents on the Earth in which radioactive sources were involved are subject to many uncertainties – for instance in determining parameters like the radiation quality factor Q or the dose and dose rate effectiveness factor (DDREF). The uncertainties in the DDREF, which is more or less related to the factor D_{50} defined in Eq. (8), are supposed to become particularly large when the dose rates are lower than 0.05Gy/h (Cucinotta & Durante, 2006). The dose rates expressed in Eqs. (1) and (3) which are relevant to Mars missions, are much lower than that value, even taking into account the occurrence of a large SEP.

Despite the many limitations encountered when attempting to predict the risk of stochastic and deterministic effects of radiation in space, it can be said that, at least in what concerns the radiation quality factors used below, namely the factors Q_{space} and Q_{Mars} shown in equations (2) and (4) respectively, these quantities have been estimated using data obtained during a 'real' Mars mission. Moreover, the formula for calculating risk in Eq. (7) is the result of several advances in the technique of data analysis that has contributed to greatly improve the statistics earlier utilized in Edwards & Lloyd, (1998).

On the basis of the parameters θ_1, θ_∞ and V used in Edwards & Lloyd, (1998) regarding various fatal and non-fatal deterministic effects, it is possible to also compute the coefficient D_{50}. In this regard, the values of the RBE are approximated using the radiation quality factor for space Q_{space} given by Eq. (2).

It is noted that the radiation quality factor Q_{space} was measured experimentally and so takes into account all kinds of radiation inside the spacecraft (i. e. coming both from primary and secondary particles). It is assumed that the stay on the surface of Mars is short, of the order of 30 days as it is typical of a short-term mission (McKenna-Lawlor, 2012). We choose as the effective dose for the whole mission $H = RBE \cdot D$ a value that is about four times larger than the values H_I and H_{II} coming from the data of the Curiosity mission given in Eqs. (5) and (6) respectively:

$$H \sim 3.82\,\mathrm{Sv} \tag{9}$$

This higher value allows us to estimate what happens in a limit situation in which there could be large SEP events. Moreover, in the calculation performed in this subsection the dose rate is the average absorbed dose rate in space given in Eq. (1). We rewrite it here using a slightly different set of units, that fits with the units in which the parameters θ_∞ and θ_1 will be defined afterwards:

$$\dot{D} = 2.00 \cdot 10^{-5}\,\mathrm{Gy/h} \tag{10}$$

We suppose that there is no attenuation of the radiation delivered to an organ due to its absorption by the surrounding tissues and by the hull of the space vessel. This hypothesis can be justified as follows. First of all, we expect that the overwhelming contribution to the absorbed dose comes from highly energetic protons and we consider a stay on the surface of Mars of the order of one terrestrial month. In this case, the total absorbed dose will be delivered mainly in space and will predominantly consist of GCR. As is well known, the main contribution to the particle content of GCR is made by protons (about 80%) whose energy spectrum peaks in the energy range between 300MeV and 1 GeV (Particle Data Group, 2014). Already at 300MeV, the range of protons in aluminum is about 6.57g/cm², i.e. 24.34cm, while at 1GeV the range grows to $4.10 \cdot 10^2\,\mathrm{g/cm^2}$,

corresponding to 152.60 cm. In consequence, the percentage of particles stopped by the hull of the spacecraft, even assuming a thickness of 24cm of aluminium, is limited. In this situation, in which there is a predominance of light, very energetic nuclei, the doses related to secondary particles, in particular neutrons, can be considered to be negligible, at least to a first approximation. A fortiori, it is possible to neglect the difference among the doses absorbed by different tissues in the human body. As a matter of fact, the average density of human tissues is lower than that of aluminium (by about a factor 2.70) and their stopping power is proportionally less. For example, approximating the density of a tissue with that of water, it turns out that the range of protons at 300MeV in water is about 5.14 g/cm^2 (~ 51.40 cm). At 1 GeV the range increases to 325.00 g/cm^2 (~ 300.00 cm). Clearly, in this situation there is no attenuation of the radiation delivered to an organ due to its absorption by surrounding tissues. On the other hand, apart from protons, another component of the cosmic rays that may have relevant health effects is composed of highly energetic heavy nuclei. The energy spectrum per nucleon of these particles is also peaked in the energy range between 300 MeV and 1 GeV (Particle Data Group, 2014). These particles have also a high penetrating power inside the human body. Their flux is many orders less than that of protons. Despite this fact, heavy nuclei can have disastrous consequences for the health of the crew of a manned mission (Cucinotta & Durante, 2006; Zaman et al., 2021), mainly because, when hitting the nucleus of a cell, they produce localized multiple damage in the DNA, which are much more difficult to repair than the damage caused by low-LET radiation. The effects of heavy nuclei cannot be taken into account by Eq. (7) for reasons that will be explained later. Also, some of these effects, for instance on the functionality of the brain and of the central nervous system, will be discussed separately.

On the basis of Eq. (7) it is possible to exclude the occurrence of several fatal deterministic effects, like the gastrointestinal syndrome and the pulmonary syndrome discussed in Edwards & Lloyd (1998). These pathologies are simply beyond the lower threshold of occurrence D_{low}. What cannot be entirely neglected is the risk R_{hs} of the hematopoietic syndrome, which amounts to:

$$R_{hs} \sim 1.20 \cdot 10^{-5} \qquad (11)$$

In this syndrome the killing of blood cell precursors in the bone marrow is so high that blood cells in the body cannot be substituted for them at the rate required. The main consequences are bleeding and a lower ability to fight infections for a long period after irradiation, see Edwards & Lloyd (1998) and references therein for more details.

In computing R_{hs}, the parameters $\theta_\infty, \theta_1, V$ are set following Edwards & Lloyd, (1998): $\theta_\infty = 3.00$ Gy, $\theta_1 = 0.07$ Gy^2/h and V=2. We recall that in the present subsection 3.1.1, the equivalent dose and the dose rate are given by Eqs. (9) and (10) respectively.

Morbidity effects seem not to be very likely. For instance, the risk R_{cat} of cataracts of the eye $(\theta_\infty = 3.00$ Gy, $\theta_1 = 0.01$ Gy^2/h, V=5$)$ amounts to:

$$R_{cat} \sim 1.31 \cdot 10^{-8}$$

The risk of vomiting R_v is instead given by $(\theta_\infty = 2.00$ Gy, $\theta_1 = 0.20$ Gy^2/h, V=3$)$:

$$R_v \sim 2.14 \cdot 10^{-9}$$

Relevantly higher is the risk of diarrhea $(\theta_\infty = 3.00$ Gy, $\theta_1 = 0.20$ Gy^2/h, V=2.50$)$:

$$R_d \sim 1.47 \cdot 10^{-6}$$

These differences are however not really significant. What is significant is the order of magnitude of the values of all the risks of deterministic effects computed above. They are each very low, implying that the occurrence of these effects is not very probable.

It is worth to stress that in the case of the central nervous system (CNS) syndrome, it is not possible to specify the parameters D_{50} and V with the statistical techniques used in Edwards & Lloyd, (1998). The reason is that to determine the symptoms of this syndrome is not an easy task. Even when very mild symptoms are presented, a patient affected by the CNS syndrome will die within 48 hours due to several possible causes. According to Edwards & Lloyd (1998), the CNS syndrome occurs at acute doses above 50 Gy. In the case of Mars missions. we have seen that the doses delivered are far below this threshold while also the dose rates are much lower than those typical of acute doses. Despite that, more subtle influences can be exerted on the CNS due to GCR, which do not lead directly to death, but constitute a serious danger of impairment for the crew members of a human BLEO mission. The reason is the presence in GCR of heavy nuclei accelerated to very high energies, the so-called HZE particles. Such particles are not efficiently stopped by the shielding of the space vessel and, when hitting a non-dividing cell such as a neuron, are very likely to compromise their functions in a relevant way. This problem has been known for many years. Already in 1973, an advisory panel to the Space Science Board of the US National Academy of Science urged a careful investigation of this aspect of GCR (Grahn, 1973). In Curtis et al. (1998) it was shown using numerical simulations that, during a three-year mission to Mars at solar minimum, HZE-particles may damage extended portions of a relevant percentage of hippocampal and thalamus cell nuclei. The effects of this radiation can be asymptomatic for a long time due to the exceptional abilities of the CNS to overcome damage. However, the authors of Curtis et al. (1998) did not exclude the related possibility of complete inactivation of the neurons and/or the onset of a series of events leading to cell killing or functional impairment. This subject has attracted considerable attention in recent times due to new results coming from experiments on mice. These results show that even relatively small doses of heavy nuclei (of the order of a few tens of cGy) lead to alterations in neurons and in synaptic connectivity, eventually causing a significant behavioral impairment in the irradiated mice (Limoli, 2015), (Kiffer et al., 2019).

In conclusion, the probability of deterministic effects studied here with the help of the sophisticated statistical methods of Edwards & Lloyd, (1998) is very low when doses typical of a mission to Mars are considered. However, this does not mean that there are not other deterministic effects that could become relevant during space travel. One example are the damages to the CNS induced by HZE radiation. Another example are the light flashes that some astronauts involved in lunar missions have perceived. This will be discussed in the next Subsection. Here we would still like to stress again the possibility of cardiac events like cardiac arrhythmias, sudden cardiac death, acute myocardial infarction, and cerebral stroke discussed in Steupel et al. (2006) (see also Hughson et al., 2018).

3.1.2 Deterministic effects in the case of Moon missions

1. In the case of short missions to the Moon, like the Apollo sequence, it is possible to rule out deterministic effects. In fact, absorbed doses of the order of 10mGy are far below the lower

threshold D_{low} of any lethal or non-lethal deterministic effect. However, a prolonged stay in the van Allen belts could be dangerous. Assuming an equivalent dose rate of about 13.33mSv/h (McKenna-Lawlor, 2012), it is possible to use Eq. (7) in order to evaluate the lower and upper threshold doses for the onset of a given deterministic effect. For example, in the case of the bone marrow syndrome described above, the lower threshold dose is D_{low}~3.30Gy, corresponding to a stay in the Van Allen belts of about 248h. The upper dose D_{up} amounts to 21.50Gy and the relative exposure time is about 1610 hours.

2. An important phenomenon is that of the light flashes which were perceived by individuals subjected to ionizing radiation mainly in dark environment conditions. This effect could be potentially dangerous if tasks should be performed which require a high level of visual efficiency. Further, light flashes could cause disturbances in the sleep cycle. According to data collected during the ALTEA mission (Narici, 2008), about 30 events were observed in the average by astronauts in the USLab module of the International Space Station (ISS). The rate of flashes varied depending on the shielding of the spacecraft and on individual physiological and psychological characteristics (Narici, 2008). The identity of particles causing light flashes is still a subject of debate. In space, candidates are light ions with charges ranging from $3 \leq Z \approx 8$, although heavier ions are also suspects (Narici, 2008). The latter, like for instance *Fe*, are more difficult to investigate because they are less abundant in LEO than are lighter ions. On the ground also light particles like pions, muons and alpha-particles have been shown to cause light flashes (Narici, 2008). More recently, cases involving high energy photons (6MV) have been reported (Blumenthal et al., 2015). The possibility that also protons could be associated with the phenomenon is under discussion (Narici, 2008). The mechanism by which individuals whose eyes and/or cortex are irradiated perceive light in the complete darkness is not yet fully understood. According to Narici, (2008), it is for instance not clear if there exist some mechanism to "see" ionizing radiation directly or if the observed light flashes are due to photons in the visible spectrum emitted after the high energetic particles hit an area near the retina.

3.2 Stochastic effects

In many cases cells survive after irradiation but their metabolism is impaired. After dividing, these cells produce other malfunctioning cells that lead to stochastic effects. Stochastic effects can be somatic – cancer induction (leukemia or solid tumors) – or heritable effects (also called hereditary effects). The latter are diseases appearing in the offspring of individuals exposed to ionizing radiation as a consequence of the exposure. In contrast to deterministic effects, it is believed that there is no threshold dose for the onset of the stochastic effects of ionizing radiation. More precisely, the probability of the onset of a stochastic effect, but not its severity, may be regarded as a function of the delivered dose (ICRP, 1977). According to this definition, every dose, no matter how small, is able to cause severe effects like cancer induction or mutations in individuals hit by ionizing radiation and in their next generations. Hereditary effects have proved to be very difficult to study in humans and, up to now, they have been predominantly tested on animals. According to UNSCEAR, (2012), there is an emerging consensus that transmissible chromosomal instabilities do not occur in response to low-LET radiation below 0.5 Gy, implying the existence of a threshold at least in this case. This conclusion does not however apply to the high-LET radiation, relevant to space missions.

3.2.1 Dose-response curves and low-high doses
Cells exposed to ionizing particles show a response to the dose of received radiation that is measurable. Possible effects can range from the molecular level, e. g. single and double strand

breakages (DSBs) up to more complex cytological and genetic lesions, such as the formation of micronuclei, chromosomal aberrations and mutations – including changes that lead to cancer. At fixed doses, the response of living cells depends on many factors, the most important of which is probably the dose rate. In mice, for instance, an acute dose of the order of 10Gy delivered in a short term causes severe illness and is mostly fatal, whereas the same dose delivered over a 450-day period is apparently harmless (Ina et al., 2005). However, it should be taken into account that high accumulated doses may result in the occurrence of tumors after many years, an outcome which makes the study of stochastic effects very difficult. In BLEO missions involving the exploration of Near-Earth Objects/NEOs, rather low dose rates turn out to be relevant, as is shown by the values of the average dose rates absorbed in space and on the surface of Mars given by Eqs. (1) and (3) respectively.

Other relevant factors that determine the response of cells or tissues include the type of radiation, as well as physiological and environmental conditions. The concepts of low dose and low dose rates have been extensively discussed in the literature, see e. g. UNSCEAR (2000) – Annex G, using as an effect solid tumor induction in humans and epidemiological studies on populations affected by radiation emitted during the Japan bombings or nuclear accidents. A low dose has been defined in UNCR, (1993) to be any exposure below $D_{max-low-dose} \sim 200 \, \text{mGy}$ [1].

At the moment, there is no satisfactory model which is able to predict the possible mutagenic, cancerogenic and teratogenic effects of radiation at low doses. The standard linear no-threshold model (LNT model), which assumes that the risk of such effects per unit dose is independent of the dose delivered, is currently a subject of debate. The linear no-threshold model will be presented in the next Subsection. Several alternatives to correct this model at low doses are discussed in ICRP, (2005). The results of many experiments aiming to determine cell damage at acute low doses can be found in UNSCEAR, (2006). See also Brenner et al., (2005) for a review of epidemiological studies. The investigation of low doses is not entirely relevant in the case of space missions, because the doses involved are much higher. Yet, some interesting related phenomena like the Bystander Effect (Nagasawa & Little, 1992) and the Adaptive Response may nevertheless be of some interest and thus will be briefly treated below. The peculiar situation of space missions, in which intermediate and high doses are delivered at relatively low dose rates, certainly deserves more attention. Studies of this kind, especially in the case of high-LET radiation deposited by heavy ions, are still scarce. In the following discussion much attention will be devoted to DSBs because they are strongly supposed to be related to carcinogenesis (Han & Yu, 2010), (Bonner, 2003). Moreover, a new technique based on the phosphorilation of the histone H2AX for the fast detection of DSBs during irradiation has been recently developed, see Firsanov et al. (2011) for a review on this topic.

3.2.2 An approximate calculation from statistical data assuming the linear no-threshold model

Assuming that the dose-response curve for radiation-induced cancer follows a linear model with no threshold, it is possible to predict the probability of the onset of stochastic effects under space conditions using the statistical data available from the Hiroshima and Nagasaki bombings and the extensive knowledge we have about low-LET radiation. Of course, these predictions should be considered as qualitative and rough. For example, they do not take into account several factors that

---

1   In Bonner, (2003) a low dose is defined as any dose from background to 150mSv. Other definitions of low doses based on different criteria   and methodologies can be found in UNSCEAR (2000) – Annex G.

are able to change the response of living organism to ionizing radiation, such as for instance the bystander effect, and the action of repair mechanisms in aerobic/hypoxic conditions. Some of these factors will be discussed later in more detail. It should also be stressed that nowadays very sophisticated models can be used in order to predict the risks of ionizing radiation. Examples of these models are for instance the BEIR VII risk models worked out by the BEIR VII Committee (BEIR VII, 2006). While their discussion is out of the scope of this work, the risk models in (IAEA, 1996) will be briefly mentioned below.

One of the main ingredients of our considerations is the excess relative risk per unit dose ERR_Z, a quantity that measures the percentage increase of risk of contracting fatal cancer in a group of N_0 individuals exposed to radiation of type Z with respect to a group of N_0 unexposed individuals. In the case of the atomic bombings in Japan, the statistical analysis of the data obtained from the survivors has shown that the excess relative risk $ERR_{HN} = \dfrac{N_{exp}}{N_{unexp}} - 1$ for the induction of fatal cancer per unit dose is

$$ERR_{HN} \sim 0.50 \, \text{Sv}^{-1} \tag{12}$$

where the subscript HN refers to the Japanese cities of Hiroshima and Nagasaki. Moreover, N_{exp} is the number of events occurring in the group of N_0 exposed individuals, while N_{unexp} denotes the number of events in the control group consisting of N_0 unexposed individuals. 'An event' means here a mutation leading to a fatal cancer. A crude estimate of the excess relative risk per unit of dose in space ERR_{space} during the journey to Mars can be obtained using the fact that the excess relative risk scales roughly according to the quality factor Q of the radiation type. Thus, denoting with the symbol Q_{HN} the quality factor of the radiation in the bombing of Japan, an approximate evaluation of ERR_{space} is:

$$ERR_{space} = \frac{Q_{space}}{Q_{HN}} ERR_{HN} \tag{13}$$

The quality factor Q_{space} is already known from Eq. (2). To determine Q_{HN}, we proceed as follows. The radiation due to the atomic bombs at Hiroshima and Nakasaki consisted not only of photons, which are low-LET radiation, but also of neutrons with a quality factor of about 10. Following Preston et al. (2004), we assume that the neutrons constituted 1% of the total radiation[2]. In consequence, the quality factor of the radiation in the bombings of Japan is approximately:

$$Q_{HN} \sim 1.09 \tag{14}$$

Substituting this value of Q_{HN} in Eq. (13), we obtain for ERR_{space}:

$$ERR_{space} = 1.75 \, \text{Sv}^{-1}$$

This value is quite high. It implies that the probability of fatal cancer during the entire life (about 20% in the unexposed population) raises to 55% after an exposure to a sievert of space radiation during the travel to Mars.

---

2 Actually, while the Hiroshima bomb was plutonium based, the Nagasaki one was uranium based and its percentage of neutrons was less by at leastby a factor 2 (Preston et al., 2004).

Based on the ERR_Z and other similar quantities, it is possible to establish models that attempt to assess the risks connected to irradiation. For instance, let $ERR_Z(D,e,s)$ be the excess relative risk of contracting fatal cancer after an exposure at the age e of an individual of sex s to a dose D of radiation of type Z. The so-called probability of causation, in short PC, estimating the chances that the absorbed dose of radiation has been the cause of fatal cancer after the exposure, may be approximated as follows (IAEA, 1996):

$$PC = \frac{ERR_Z(D,e,s)}{1 + ERR_Z(D,e,s)}$$

The formulas for evaluating $ERR_Z(D,e,s)$ may be found in (IAEA, 1996). Another relevant quantity is the Risk of Exposure Induced Death (REID). The REID accounts also for cancer deaths that would occur anyway, but are shifted to an earlier age due to radiation exposure. The NASA career dose limits in sieverts are established by requiring that the exposure to radiation during the career of a particular astronaut is not to exceed 3% Risk of Exposure Induced Death/REID from fatal cancers at the upper 95% confidence level, see for instance (McKenna-Lawlor et al, 2014). More information on the evaluation of the risks of stochastic effects due to ionizing radiation can be found in (L. Walsh et al., 2019).

3.2.3 The Bystander Effect

The Radiation Induced Bystander Effect (RIBE) is a term coined to describe the induction of biological effects in non-irradiated cells as a result of signals released by nearby cells that have been irradiated by some kind of ionizing radiation. Biological effects include cytologic abnormalities, i.e. they are visible at cell level, such as the formation of micronuclei or chromosomal aberrations, and genetic or epigenetic alterations. The relationship between the DSBs induced by the Bystander Effect and gene mutations have been studied by, for instance, Asur et al., (2010). RIBE refers also to effects occurring in non-irradiated cells when they are put in contact with a medium coming from irradiated cells. The Bystander Effect appears only at low doses imparted either by low-fluence radiation or when using microbeamer facilities. RIBE represents a departure from the linear no-threshold model explained in Subsection 3.2.2, because it implies that the damage due to radiation is not only proportional to the absorbed dose, but that an enhancement factor is related to the fact that non-irradiated cells are also affected. Incidentally, this means that there may be low radiation doses the detrimental effects of which could exceed that of higher doses, see for instance UNSCEAR (2000) – Annex G, paragraph 388. RIBE was first reported to occur in the ovary cells of a Chinese hamster (Nagasawa and Little, 1992) using α-particles. While 1% of the cells had been hit by the α-particles, it was observed that about 30% of the cells were affected after irradiation by sister chromatid exchanges (a cytologic effect). Later studies showed that RIBE could cause many other effects, including for instance: the appearance of micronuclei, chromosomal aberrations, double strand breaks and apoptosis - as well as gene mutations and genomic instability. Most of the evidence supporting RIBE comes either from in vitro studies in which the cultured cells belong to very established lineages, (including cancerous cell lines) or are obtained in vivo from mice models. There are also experiments supportive of the existence of RIBE in tissues. The non-irradiated cells affected by RIBE can be at different distances from the irradiated cells, ranging from less than a millimeter to more than a centimeter. As already mentioned, it is possible to induce Bystander

Effects by exposing the non-irradiated cells to a medium that has been in contact with irradiated cells (W. Han and K. N. Yu, 2010).

4   Need of a new strategy to ascertain the safety of BLEO missions

4.1   Statement of the problem and outline of its possible solution

Summarizing, the crew of an hypothetical BLEO mission will have to endure equivalent doses of ionizing radiation of about 1 Sv delivered during a period of approximately one year. To study the deterministic and stochastic effects on human crew members exposed to such doses, it is possible to apply the statistical methods discussed in the previous sections and based on epidemiological studies on the survivors of the bombings of Nagasaki and Hiroshima as well as on accident victims in which radioactive substances where involved. Deterministic effects have been the subject of Section 3.1. These obtained results hint unequivocally that the risk threshold $R(D_{low})$~0.01 for the appearance of medically diagnosable effects related to cell killing caused by the absorption of ionizing radiation is very far from being reached. However, it should be taken into account that the predictions of formulae like the Weibull function of Eq. (7) become inaccurate for dose rates that are not acute. Further knowledge could be gained from LEO mission exposures. It is known that during LEO space flights the crew is affected by disturbances related to the impairment of the immune system, see for instance (Sonnenfeld et al., 2003), (Crucian et al., 2008). Nevertheless, due to the limited statistics available thus far, it remains to be proved if such disturbances have long-term consequences on the health of astronauts or on their lifespan[3]. Besides, it is not easy to separate the short-term deterministic effects of ionizing radiation from those due to all the other factors that can impact the health state of a human crew in space. Examples of such factors are microgravity, isolation and emotional stress. Even more difficult to predict is the insurgence of stochastic and heritable effects. An additional complication for BLEO missions is provided by the fact that the radiation contains a relevant component of HZE particles whose effects on humans are still largely not understood (G. Horneck, 1992), (G. Horneck, 1994), (Cucinotta & Durante, 2006). A comprehensive review on the state of our understanding of the effects of HZE particles exposure can be found in Sridharan et al., (2015). In that publication, the phenomenon of tumorigenesis after HZE particle exposure is tackled by considering the relation between the risks of cancer induction and the oxidation stress following irradiation. An extensive series of interesting experiments are reported which show how the production of reactive oxygen species after a dose of HZE radiation may last for years, interfering with the cell repair system, affecting telomere stability and favoring chronic inflammation. It is also hinted that reactive oxygen species might affect the redox regulation of proteins and lipids in cells, changing in this way the epigenetic landscape, a fact that could lead to heritable effects. The recommendation of Sridharan et al., (2015) is that, in order to understand the phenomenon of tumorigenesis, it is critical to evaluate how the persistent production of reactive

---

3   We would like to cite on this point the interesting study based on epidemiological data of Cucinotta et al. (2001) on a late effect of ionizing radiation like the early appearance of cataracts. From the statistics obtained examining 295 astronauts involved in several different missions it turns out that radiation increases the risks for cataracts in a dose and LET dependent way.

oxygen species and their moieties depend on the charge, energy and LET of the imparted ionizing radiation. The work of Sridharan et al. (2015) presents also the status of the state of the art of in vivo experiments performed on mice models in order to assess the risks of cancer related to ionizing radiation in humans.

One of the conclusions coming from these studies on mice is that the risks after irradiation of the insurgence of a number of different types of carcinomas is largely dependent on LET. It is also possible that there exist LET thresholds for the induction of certain kinds of tumors. This seems to be for instance the case for the thymic lymphoma investigated by (Jacobus et al., 2013).

Despite these advances, which significantly improve our understanding of the risks of HZE radiation, several problems in determining safe career dose limits for astronauts still persist. Limitations in the prediction of the health risks incurred due to ionizing radiation in space are discussed in Chancellor et al., (2014) and Chancellor et al., (2018). The key point is that with experiments in vivo or in vitro[4] it is extremely difficult to determine the probability of stochastic events that, in humans, may require decades before they occur. To this purpose it is of uttermost importance to determine and apply markers that are able to signal in advance the early stages of carcinogenesis in humans. The human genomic features associated with a higher likelihood of developing given types of cancer should also be taken into account.

A new strategy to overcome at least in part these difficulties and to achieve the goal of establishing career limit for astronauts engaged in BLEO missions with a safety level comparable to that NASA criteria for LEO missions (McKenna-Lawlor et al, 2014) could come from some recent results of precision oncology and molecular medicine. In particular, thanks to an extensive effort in clinical testing supported in the laboratory also by the application of human organoids cultures and genome editing technology, it has been possible to characterize with a high precision the pathways of tumor formation in selected human organs. This knowledge, combined with the use of organoids could be helpful in solving the problems mentioned before by integrating the data coming from in vitro experiments and animal models. First of all, the possibility of detecting the early stages of cancer in an organ only by checking the activation of certain oncogenic pathways (such as the TGF-β signaling pathway), the loss of heterozygosity or other gene alterations, opens the possibility of predicting the risks due to irradiation to be made without the need of waiting for decades before being able to collect significant statistical data concerning the induction of cancer in a given organ. Secondly, organoids can now be formed using human cells, so that it is not necessary to rely on mouse models. If so required, it is always possible to transplant tissues from these organoids at any stage of the evolution of their health after irradiation and to check the level of tumorigenity in these tissues. Moreover, organoids are technically more challenging than the 2D models of cultured cells, but still retain the ease of maintainance and analysis typical of in vitro models. Organoids provide in the context of the study of the early steps of the onset of cancer a much more refined simulation of human organs than traditional cultured cells. It is sufficient here to mention the fact that cells in a 2D culture lose their differentiation in most cases (von der Mark et al., 1977), (Petersen et al., 1992) and their polarity. Both features are preserved when cells are put in a 3D environment (Debnath & Brugge, 2005), and sometimes can be restored when passing from 2D to 3D (Benya & Shaffer, 1982), (Yamada & Cukierman, 2007). Gene expression and reproduction are also very different than in vivo (Birgersdotter et al., 2005). For instance, the loss of polarity affects also the intracellular

---

4   In vitro experiments are easy to perform, but have strong limitations. The cells used in such experiments are very different from those of the human body: They duplicate in a different way, the cell polarity is lost, the cell differentiation that distinguishes the cells in an organ is lost just to mention a few disadvantages.

signalling pathways (Birgersdotter et al., 2005) that, as we will see later, play a crucial role in the development of cancer.

The present state-of-the-art of our knowledge of the early stages of cancer thanks to precision oncology will be illustrated in the next Section taking as an example the specific case of colorectal cancer. Next, the applications to space medicine outlined above will be explained in more detail.

4.2   Colorectal cancer as a study case

The events leading to colorectal cancer (CRC) in humans have been satisfactorily understood at the molecular level. This makes it possible to detect the early stages of colorectal carcinogenesis with the help of techniques that are already commonly applied like RNA-sequencing and Loss of Heterozygosity (LOH). Comparable knowledge already exists or is soon to be obtained for the case of solid tumors affecting other organs, like for instance pancreatic cancer (Du et al., 2017). All of these advances together with the use of organoids open the possibility to achieve a better understanding of the risks of stochastic effects related to exposure to ionizing radiation without the need of waiting decades for the appearance of such effects in vivo or reliance on complex epidemiological studies.

Coming back to colorectal cancer, its development follows a pathway of ordered events transforming normal epithelium cells into adenoma cells and finally leading to adenocarcinomas. "Driver" events are certain mutations – later discussed in details - activating oncogenes, tumor suppressor genes and genes responsible for activating certain signal transduction pathways that regulate the response of cells to stresses. Molecular studies of precision oncology have revealed that these driver events are not sufficient to initiate tumor progression. The presence of genomic instability is required in order to increase the rate of mutation necessary for cancer formation. Remarkably, the set of driver events and genomic instabilities that are relevant for colorectal cancer is limited, a fact that potentially allows verification in the laboratory if, at some stage after irradiation, the conditions for the formation of a tumor are present or not. Out of 80 genes that are involved, it has been found that the real drivers consist of mutations of a much smaller number of genes (Pino & Chung, 2010). The genomic instabilities involved are only three. With regard to driver events, relevant is the inactivation of the APC (Adenomatous Polyposis Coli) tumor suppressor gene followed in the majority of CRC's by the mutation of the KRAS gene, a proto-oncogene that regulates cell growth. More rare is the mutation of the BRAF proto-oncogene.

It is noted that the mutations of the KRAS and BRAF genes are mutually exclusive. The coexistence of both mutations has been observed in only 0.001% cases of CRC. Other important genes in KRAS mutated CRC's are the TP53 and SMAD4 tumor suppressor genes and the PIK3CA oncogene[5]. In BRAF the mutated CRC's affected are genes related to the DNA mismatch repair system like MLH1, MLH3, MSH2, MSH3, MSH6 and PMS2. These genes are usually not mutated, but rather epigenetically silenced, although their mutation is possible, but occurs less frequently. The signaling transduction pathways that are altered in CRC are those that regulate cell differentiation, migration, proliferation and apoptosis. In particular, the WNT-β-catenin and the transforming growth factor-β (TGF-β) pathways are affected. For instance, the APC gene

---

5   It is noted that some genes involved in signal transduction pathways play also a role as oncogenes or historically have been classified as oncogenes. In consequence, sometimes genes such as PIK3CA, which belong to the PI3K signaling pathway, are referred to in the literature as oncogenes.

synthesizes a large protein with multiple functions including apoptosis and cell differentiation and interacts with the β-catenin. The latter is a protein that regulates the adhesion of cells as well as gene transcription. It is noted that the activation of the WNT pathway is connected to an increased production of reactive oxygen species that cause further molecular damage (Yoon et al., 2010). The WNT pathway is activated in 93% of CRC's (Rodriguez-Salas et al., 2017). The role of reactive oxygen species is one of the key points for understanding cancer development after exposure to radiation, see Sridharan et al., (2015). As we see, the results of molecular medicine provide a more fundamental insight into the mechanisms that cause oxydation stresses in cells. The other crucial affected pathways are TP53, PI3K, the transforming growth factor beta (TGF-β) and MAPK (Rodriguez-Salas et al., 2017), Epithelial-to-mesenchymal transition (EMT) and MYC oncogene amplification. The gene mutations in the early development of CRC take place at a fast rate that cannot be explained by the random mutation rate occurring in normal cells. As already anticipated, for initiating cancer the presence of genomic instabilities favoring the accumulation of mutations is necessary. While it is not clear if the genomic instabilities start the mutations or vice-versa, the three types of genomic instabilities leading to CRC are known. In about 85% of CRC's, the driving events are accompanied by chromosomal instability (CIN), while the remaining 15% are characterized by microsatellite instability (MSI) (Dienstmann et al., 2017). In tumors with MSI, the occurrence of a high CpG island methylation phenotype (CIMP) is frequent (Dienstmann et al., 2017). Basically, CIN can be related to double strand breaks with subsequent defective repair leading to a fast rate of gain or loss of large portions of chromosomes, a phenomenon that causes somatic copy number alterations (SCNA) of genes. More precisely, CIN arises from defects of chromosomal segregation and of telomere stability (Dienstmann et al., 2017). The CIN condition is also favored by mutations in the TP53 pathway and other crucial genes (Dienstmann et al., 2017). The alterations related to CIN change the number and appearance of the chromosomes, in short, the karyotype of cells. While normal human cells are diploid, i. e. chromosomes are present in two copies, cells affected by CIN can have chromosomes with a different number of copies and some of their chromosomes may suffer the loss of heterozigosity (LOH). LOH occurs when one gene in one allele is lost. In cancer this becomes a crucial event when the gene is a tumor suppressor and the copy of that gene in the other allele is inactive due to a point mutation. The gain or loss of large portions of chromosomes usually ends up with the shortening of telomeres[6]. The CIN-pathway to tumorigenesis can be summarized as follows. With the activation of the WNT signaling pathway, whose proteins decrease the stability and levels of the proteins devoted to the degradation of β-catenin, the latter protein starts to accumulate in the cell cytoplasm. The damage of tumor suppressor genes like the APC gene, promoting the degradation of the β-catenin protein, contributes to this accumulation. The β-catenin proteins are responsible for cell growth and cell adhesion. When they translocate to the nucleus, one of their actions comprises interfering with the transcription of the MYC oncogene. The mutation of the KRAS gene further promotes cell proliferation and inhibits cell apoptosis. The SMAD2 and SMAD4 tumor suppressor genes also mutate inhibiting the TGF-β

---

6   Telomeres are structures consisting of a repetition of the DNA sequence TTAGGG. They are found at both ends of each chromosome and protect chromosomes from nucleolytic degradation, from inappropriate recombination, repair and fusion. The role played by the shortening of telomeres in making the genome unstable has been discussed in Shridharan et al., (2015). It is noted that exposure to radiation may also activate telomerase in human somatic cells stimulating the increase of the length of telomeres. The lengthening of telomeres has indeed been observed in irradiated limphoblasts (Neuhof et al., 2001) and, more recently, also in humans, see the NASA Twin Experiment reported in Segal, (2016).

pathway. This leads to uncontrolled cell proliferation. At a later stage the mutation of TP53 also occurs. As already said, this fast rate of mutations becomes possible due to CIN. The techniques used to detect CIN are described for instance in (Pino & Chung, 2010). They include cytometry, karyotyping, loss of heterozygosity, fluorescent in situ hybridization (FISH), and comparative genomic hybridization (CGH). The second pathway to genomic instability, MSI, is related to gene loci (the microsatellites) characterized by sequences of one to six nucleotides, for instance $(AC)_n$, that are repeated a number of times n ranging from five to one hundred. The human genome contains about $50000 - 100000$ microsatellites that are found mostly in the non-coding regions of DNA, but also in the coding part. The presence of repeated units creates motifs in the DNA that are used as markers for various enzymes, although most microsatellites look to be neutral markers. As markers, microsatellites play a role in DNA recombination and replication. Moreover, they influence gene expression. More details concerning their functions can be found in (Chistiakov et al., 2006). Microsatellites are areas intensively subjected to mutations because their regularity is easily altered during recombination and replication. For this reason, they are very sensitive to errors in the DNA mismatch repair system Methyl Mismatch Repair/MMR, which is responsible for the recognition and repair of mistakes occurring when DNA replicates or recombines. The proteins involved in MMR are coded by seven genes: MLH1, MLH3, MSH2, MSH3, MSH6, PMS1 and PMS2. The mutation or epigenetic silencing of some of these genes leads to MSI. The MSI pathway to cancer can be summarized as follows. Microsatellites are found in the powerful tumor suppressor gene TGF-β that regulates cell division. The malfunctioning of MMR leads to MSI and favors the further stages of carcinogenesis, accumulating mutations in genes responsible for cell differentiation, apoptosis and growth. The alteration of the TGF-β pathway with the subsequent coding of abnormal typeII TGF-β receptor and BAX protein leads to cell proliferation[7]. The mutation of the BRAF gene is an important event in MSI tumorigenesis too. Let us also note that the mutations of TP53 and of the ATM gene are mutually exclusive. The alteration of ATM gene is predominant in the MSI phenotype. In MSI driven CRC a relevant role is played by epigenetic silencing of the critical genes. For instance, MLH1 is epigenetically silenced by CIMP (CpG Island Methylator Phenotype). CIMP refers to the methylation of the CpG islands. These are regions of DNA in which there is a high frequency of CpG sites. Such sites consist of a cytosine nucleotide connected to a guanine nucleotide by only one phosphate. Formally, a CpG island is defined as a region of at least 200 base pairs in length in which the percentage of dinucleotides CpG is greater than 50%. The CpG islands in humans and, more in general, in vertebrates, are located in the promoters or near the promoters. In humans, CpG islands are present in about 70% of promoters. Promoters can be found in the initial part of genes and play an important role in transcription. As a matter of fact, to start the gene transcription, the enzyme that synthetizes RNA, known as RNA polymerase, must bind to the promoter. If the binding site is inactivated, the gene is no longer recognized by the RNA polymerase and its expression doesn't occur. In CRC tumor suppressor genes and other genes related to tumor development are epigenetically silenced by hypermethylation of the CpG islands. Methylation is a process in which the cytosine of the CpG sites has a methyl group attached. In CIMP, the CpG islands in the promoters of multiple genes are simultaneously hypermethylated. For the analysis of MSI several methods may be applied, like immunohistochemical (IHC) staining and PCR amplification. The IHC analysis is performed using antibodies to identify the loss of proteins that are involved in MMR. In the PCR analysis a set of

---

7   More information about the transforming growth factor TGF-β in CRC can be found for instance in (de Miranda et al., 2015).

suitable microsatellite markers is used, like for instance the BAT25, BAT26, D2S123, D5S346 and D17S2720 markers. For instance, D2S123 is a microsatellite linked to the MHS2 gene containing repeat of CA (cytosine – adenine) with chromosomal location 2p16. Concerning CIMP, up to now there is no a consensus methodology to check CIMP status (Powell et al., 2018). Different panels of markers used by independent groups can be found in the review (Powell et al., 2018).

To conclude, remarkable advances have been made in order to characterize at the molecular level the insurgence of CRC. To some extent, the gap in our knowledge about the processes behind the onset of cancer, starting from the physical and chemical effects visible immediately after exposure to radiation, like DSB, oxidation of nucleotides due to reactive oxygen species etc. and the later appearance of stochastic effects like solid tumors, has been filled. The initial steps of the development of cancer can be monitored looking at a limited set of genes that are mutated or epigenetically silenced together with the appearance of genomic instabilities like CIN or MSI. The progresses of molecular medicine are not only restricted to CRC. Similar results have been obtained also in the case of other cancers, like for instance hepatocellular cancer, biliary cancer and pancreatic cancer (Tanaka, 2018). Of course, it should not be forgotten that the research in this subject is still ongoing. In particular, a unique classification of the subtypes of CRC with respect to their response to given therapies still does not exist. Depending on the markers and methodology used, different groups have provided several classifications of CRC in which the number of subtypes varies from three to six. We report here the four consensus molecular subtypes (CMS) that have been distinguished after an extensive analysis of 4000 samples of CRC (Sadanandam et al., 2014), see also (Dienstmann et al., 2017) for a review:

CMS1: This subtype (14% of early-stage tumors) is characterized by MSI and hypermethylation of the CpG islands (CIMP). The level of CIN is low. Often the mutation of the BRAF gene is observed. CMS1 is prevalent in lesions appearing in the right colon (Rodriguez-Salas et al., 2017). In CMS1 cells are hypermutated[8].

CMS2: This group (37% of early-stage tumors) is microsatellite stable, while CIN is marked, but not so dominating as in CMS4. There is a strong upregulation of the WNT and MYC pathways[9]. Moreover, the expression of the oncogenes EGFR, ERBB2 is higher or amplification of these genes occurs. These events are accompanied by mutations of the gene TP53. CMS2 is preferably diagnosed in the left colon and rectum (Rodriguez-Salas et al., 2017).

CMS3: This subtype (13% of early-stage tumors) belongs to the CIN group like CMS2 and CMS4 but has distinctive characteristics. In particular, the level of CIN is low and the upregulation of the WNT and MYC pathways is moderate. With respect to CMS2, KRAS mutations are enriched. There are mutations in the genes belonging to the PI3K pathway. In this kind of CRC the most important feature is the ability of cancer cells to undergo metabolic reprogramming. About 30% of these tumors have MSI, CIMP and are hypermutated. This subtype appears in the large intestine with no preferred location.

---

8   Hypermutation means that cells have a heavy burden of mutations. In CRC this means a mutation number equal or greater than 12 mutations/Mb (Humphris et al., 2017), where Mb denotes mega base pairs. Usually tumors with MSI are hypermutated because of the errors in MMR, while chromosomally unstable tumors (CIN dominated) are not hypermutated.

9   Upregulation of a gene means that the expression of that gene is increased, leading to an increase of the amount of its products. In signal transducing pathways several genes are involved.

CMS4:   The CMS4 subtype (23% of early-stage tumors) is characterized by TGF-β
        activation and by upregulation of genes implicated in the epithelial-mesenchymal transition
        (EMT). With respect to epithelial cells, mesenchymal cells have migratory and invasive
        properties that worsen the prognosis for this kind of tumor. A better characterization of
        CMS4, in particular the knowledge of the driver events leading to CMS4 and the regulatory
        mechanisms underlying the expressions of the genes involved, can be found in Fessler et al.,
        (2016).

CMS2-4 follow the CIN phenotype described before in this Subsection, while CMS1 follows the
MSI phenotype. CMS1 is typically hypermutated, while CMS2-4 is not, with exceptions in the case
of CMS3. The mutation distributions of different genes within the consensus molecular subtypes
and other molecular data that help in distinguishing these groups can be found in Guinney et al.,
(2015). The consensus molecular subtypes have been obtained through combining the efforts of
different research teams and they provide the most robust classification of CRC available thus far.
Still, 13% of CRC do not fit in any of the four CMS groups presented above. Research is still
ongoing in this regard and the attainment of a more complete understanding of the molecular event,
markers and pathways is necessary before it will be possible to develop efficient drugs and therapies
to cure CRC. Deficiencies in the present knowledge of CRC certainly pose a limitation for clinical
purposes, but not for astrobiological purposes. In fact, the progress mentioned above provides
powerful criteria to support recognition of the first stages of cancer after irradiation as well as
concrete tests for checking the driving mutations of the critical genes, the presence of genomic
instability and of epigenetic alterations. In the next Subsection, the application of these ideas to the
case of organoid cells will be proposed.

4.3   Organoid technologies helping in establishing safe career dose limits for BLEO missions

Organoids, also called 3D in vivo cell cultures[10], are multicellular structures that are able to mimic
some of the characteristics of in vivo organs. They can be obtained starting both from adult cells or
pluripotent stem cells. Pluripotent stem cells are either embryonic stem cells extracted from
embryos or induced pluripotent stem cells obtained by reprogramming adult stem cells. Organoids
may develop both from normal cells and from tumor cells. In the latter case they are also called
tumoroids. In order to check the effects of radiation on a given organ, it is preferable to use normal
cells. In clinical applications, the mutations leading to cancer are introduced in organoids derived
from normal tissue through applying CRISP/CAS9 mediated genetic editing or induction with the
help of viruses (such as for instance the lentivirus). This allows reproduction of different stages in
the genesis of cancer. In the present context, the source of genetic alterations is ionizing radiation of
various types, especially HZE radiation the effects of which are still not very well understood. The
idea of using organoids to investigate the stochastic effects due to ionizing radiation is still in its
infancy, but it looks very promising. With organoids it is in fact possible to mimic many human
organs like the colon, pancreas, stomach, retina and many others. A more complete list of organs
that have been simulated thus far can be found in Dutta et al. (2017). It is worth noticing that the
majority of techniques applied in clinical tests for detecting the genetic mutations and genomic
instabilities mentioned in the previous Subsection can be applied to organoids also. Moreover,

---

10 Organoids are formed using stem cells. They are not to be confused with other 3D cell aggregates like for instance
    spheroids, which are not able to mimic the structure of in vivo organs (Dutta et al., 2017).

organoids have a long life (they can last for a few hundred days). This is very useful in verifying the effects of doses that are not acute as is the norm in BLEO. Of course, organoid technologies are still in development. As such, they have still some shortcomings that affect their application in medicine. The most important limitation in the present context is that organoids are not able to reproduce the full characteristics of in vivo organs. The original stem cells differentiate in organoids into different types of cells and many structures of the real organ are reproduced, although not all of them. Recently, the induction of myelinating oligodendrocytes in cortical spheroids recapitulating the human brain has been possible (Madhavan et al., 2018). In this situation it is not clear if it will be possible to study all consensus molecular subtypes CMS1-4 of CRC mentioned in the previous Subsection because a particular subtype could originate from a kind of cells that is lacking in organoids. In principle, it could be possible to stimulate the growth of organoids with that particular kind of cell. In practice, however, despite the dramatic improvements in our understanding of CRC thanks to molecular medicine, it is not yet recognized which kinds of cell tumors of a single molecular subtype are starting. On the positive side, organoids are based on human cells. For this reason, they are certainly suitable to study tumorigenesis following exposure to ionizing radiation in humans. Animals, even humanized mice, have not the same genetic characteristics as humans, so they also have a different oncogenic process.

5   Acknowledgements

The authors wish to thank heartily B. Górski and S. McKenna-Lawlor for participating in the early stages of this work and for several fruitful suggestions. The research of FF has been supported by the Polish National Science Centre grant 2020/37/B/ST3/01471. FF would also like to acknowledge the contribution of COST Action CA17139.

6   References

S.K. Aghara, S.R. Blattnig, J.W. Norbury and R.C. Singleterry, *Monte Carlo analysis of pion contribution to absorbed dose from galactic cosmic rays*, *Nucl. Inst. Methods* **B 267** (2009), 1115.

R. Asur, M. Balasubramaniam, B. Marples, R. A. Thomas, and J. D. Tucker, *Bystander effects induced by chemicals and ionizing radiation: Evaluation of changes in gene expression of downstream MAPK targets*, *Mutagenesis* **25** (3) (2010), 271.

NCR, *Health risks from exposures to low levels of ionizing radiation: BEIR VII Phase 2*. National Research Council, Washington DC: National Academy Press; 2006.

P. D. Benya and J. D. Shaffer, *Dedifferentiated chondrocytes reexpress the differentiated collagen phenotype when cultured in agarose gels*, *Cell* **30** (1) (1982), 215-224.

A. Birgersdotter, R. Sandberg and I. Ernberg, *Gene expression perturbation in vitro − A growing case for three-dimensional (3D) culture systems*, *Seminars in Cancer Biology* **15** (2005), 405 − 412.

D. T. Blumenthal, B. W. Corn and N. Shtraus, *Flashes of light-radiation therapy to the brain*, *Radioterapy and Oncology* **116** (2) (2015), 332.

W. M. Bonner, *Low-dose radiation: Thresholds, bystander effects, and adaptive* responses, *PNAS* **100** (9) (2003), 4973.

D. J. Brenner, R. Doll, D. T. Goodhead, E. J. Hall, C. E. Land, J. B. Little, J. H. Lubin, D. L. Preston, R. Julian Preston, J. S. Puskin, E. Ron, R. K. Sachs, J. M. Samet, R. B. Setlow and M. Zaider, *Cancer risks attributable to low doses of ionizing radiation: Assessing what we really know*, *Proc. Natl. Acad. Sci. U S A.* **100** (24) (2003), 13761.

J. C. Chancellor, G. B. I. Scott and J. P. Sutton, *Space Radiation: The Number One Risk to Astronaut Health beyond Low Earth Orbit, Life* **2014** (4) (2014), 491.

J. C. Chancellor, R. S. Blue, K. A. Cengel, S. M. Auñón-Chancellor, K. H. Rubins, H. G. Katzgraberand A. R. Kennedy, *Limitations in predicting the space radiation health risk for exploration astronauts*, Microgravity **4** (2018), 8.

D. A. Chistiakov, B. Hellemans, F. A. M. Volckaert, *Microsatellites and their genomic distribution, evolution, function, and applications: A review with special reference to fish genetics*, Aquaculture **255** (2006), 1.

B. E. Crucian, R. P. Stowe, D. L. Pierson, C. F. Sams, *Immune system dysregulation following short-vs long-duration spaceflight, Aviat. Space Environ. Med.* **79** (2008), 835.

F. A. Cucinotta and M. Durante, *Cancer risk from exposure to galactic cosmic rays: implications for space exploration by human beings*, *Lancet Oncol.* **7** (2006), 431.

F. A. Cucinotta, F. K. Manuel, J. Jones, G. Iszard, J. Murrey, B. Djojonegro and M. Wear, *Space Radiation and Cataracts in Astronauts, Radiation Research* **156** (2001), 460-466.

S. B. Curtis, M. E. Vazquez, J. W. Wilson, W. Atwell, M. Kim and J. Capala, *Cosmic ray hit frequencies in critical sites in the central nervous system, Adv. Space Res.* **22** (2) (1998), 197.

J. Debnath and Joan S. Brugge, *Modelling glandular epithelial cancers in three-dimensional cultures, Nature Reviews Cancer* **5** (9) (2005), 675.

N. F. C. C. de Miranda, M. van Dinther, B. E. W. M. van den Akker, T. van Wezel, P. ten Dijke, and H. Morreau, *Transforming Growth Factor b Signaling in Colorectal Cancer Cells With Microsatellite Instability Despite Biallelic Mutations in TGFBR2, Gastroenterology* **148** (2015), 1427.

R. Dienstmann, L. Vermeulen, J. Guinney, S. Kopetz, S. Tejpar,J. Tabernero, *Consensus molecular subtypes and the evolution of precision medicine in colorectal cancer, Nature Cancer Rev.* **17** (2) *(*2017), 79.

Y. Du, B. Zhao, Z. Liu, X. Ren, W. Zhao, Z. Li, L. You, Y. Zhao, *Molecular Subtyping of Pancreatic Cancer: Translating Genomics and Transcriptomics into the Clinic*, Jour. of Cancer, **8** (4) (2017), 513.

D. Dutta, I. Heo and H. Clevers, *Disease Modeling in Stemm Cell-Derived 3D Organoid Systems, Trends in Mol. Med.* **23** (5) (2017), 393.

A. A. Edwards and D. C. Lloyd, *Risks from ionising radiation: deterministic effects, J. Radiat. Prot.* **18** (3) (1998), 175.

R. A English, R. E . Benson, J. V. Bailey and C. M . Barnes, *Apollo Experience Report –Protection Against Radiation*, NASA Technical Note NASA TN D-7080 (1973).

F. Ferrari and E. Szuszkiewicz, *Cosmic Rays: A Review for Astrobiologists*, Astrobiology **9** (4) (2009), 413.

E. Fessler, M. Jansen, F.De Sousa E. Melo, L. Zhao, P.R. Prasetyanti, H. Rodermond, R. Kandimalla, J.F. Linnekamp, M. Franitza, S.R. van Hooff, J.H. de Jong, S.C. Oppeneer, C.J.M. van Noesel, E. Dekker, G. Stassi, X. Wang, J.P. Medema, L. Vermeulen, *A multidimensional network approach reveals microRNAs as determinants of the mesenchymal colorectal cancer subtype*, Oncogene **35** (2016), 6026.

D. V. Firsanov, L. V. Solovjeva, and M. P. Svetlova, *H2AX phosphorylation at the sites of DNA double-strand breaks in cultivated mammalian cells and tissues*, Clin. Epigenetics. **2** (2) (2011), 283.

D. T. Goodhead, *Initial events in the cellular effects of ionizing radiations: clustered damage in DNA. Int. J. Radiat. Biol.* **65** (1994), 7.

D. Grahn (Editor), *HZE-Particle effects in Manned Spaceflight*, National Academy of Sciences, Washington D. D., 1973.

J. Guinney et al., *The Consensus Molecular Subtypes of Colorectal Cancer*, Nat. Med. **21** (2015), 1350.

W. Han and K. N. Yu, I*onizing Radiation, DNA Double Strand Break and Mutation*, in: *Advances in Genetics Research*, Kevin V. Urbano (Ed.), Volume 4, Chapter 7, ISBN 978-1-61728-764-0 Editor: Kevin V. Urbano, Nova Science Publishers, Inc. 2010.

D. M. Hassler, C. Zeitlin, R. F. Wimmer-Schweingruber, B. Ehresmann, S. Rafkin, J. L. Eigenbrode, D. E. Brinza, G. Weigle, S. Böttcher, E. Böhm, S. Burmeister, J. Guo, J. Köhler, C. Martin, G. Reitz, F. A. Cucinotta, M.-H. Kim, D. Grinspoon, M. A. Bullock, A. Posner, J. Gòmez-Elvira, A. Vasavada, J. P. Grotzinger (The MSL Science Team), *Mars' Surface Radiation Environment Measured with the Mars Science Laboratory's Curiosity Rover*, Science **343** (2014), 6169.

G. Horneck, *Radiobiological experiments in space: A review*, Int. Jour. Rad. Appl. And Instr. Part D. Nuclear Tracks and Radiation Measurements **20** (1) (1992), 185-205.

G. Horneck, *HZE particle effects in space*, Acta astronautica, **32** (11) (1994), 749-55.

R. L. Hughson, A. Helm and M. Durante, *Heart in space: effect of the extraterrestrial environment on the cardiovascular system*, Nature Reviews Cardiology **15** (2018), 167–180.

J. L. Humphris et al., *Hypermutation in pancreatic cancer*, Gastroenterology, **152**(1), 68.

ICRP, *Recommendations of the International Commission on Radiological Protection*, ICRP Publication 26, *Ann. ICRP* **1** (3) (1977).

ICRP, *Low-dose extrapolation of radiation-related cancer* risk, Annals of the ICRP, **35** (4) (2005).

Y. Ina, H. Tanooka, T. Yamada and K. Sakai, *Suppression of thymic lymphoma induction by life-long low dose rate irradiation accompanied by immune activation of C57BL/6 mice*, Radiat. Res. **163** (2005), 153.

J.A. Jacobus, C. G. Duda, M. C. Coleman, S. M. Martin, K. Mapuskar, G. Mao et al., *Low-dose radiation induced enhancementof thymic lymphomagenesis in Lck-Bax miceis dependent on LET and gender, Radiat. Res.* **180** (2013), 156.

F. Kiffer, M. Boerma, A. Allen, *Behavioral effects of space radiation: A comprehensive review of animal studies, Life Sciences in Space Research* **21** (2019), 1–21

C. Limoli, *Your Brain on Mars, Rad. Res.* **184** (1) (2015), 1.

E. Mazur, W. R. Crain, M. D. Looper, D. J. Mabry and J. B. Blake, *New measurements of total ionizing dose in the lunar environment, Space Weather* **9** (2011), S07002.

Susan McKenna-Lawlor, P. Goncalves, A. Keating, G. Reitz and D. Matthiä, *Overview of energetic particle hazards during prospective manned missions to Mars, Planetary and Space Science* **63**–**64** (2012), 123.

S. McKenna-Lawlor, A. Bhardwaj, Franco Ferrari, Nikolay Kuznetsov, A. K. Lal, Yinghui Li, Aiko Nagamatsu, Rikho Nymmik, Michael Panasyuk, Vladislav Petrov, Guenther Reitz, Lawrence Pinsky, Sheikh Muszaphar Shukor, A.K. Singhvi, Ulrich Straube, Leena Tomi, Lawrence Townsend, *Feasibility study of astronaut standardized career dose limits in LEO and the outlook for BLEO, Acta Astronautica* **104** (2) (2014), Pages 565 – 573.

S. McKenna-Lawlor, A. Bhardwaj, F. Ferrari, N. Kuznetsov, A. K. Lal, Y. Li, A. Nagamatsu, R. Nymmik, M. Panasyuk, V. Petrov, G. Reitz, L. Pinsky, M. Shukor, A. K. Singhvi, U. Straube, L. Tomi, T. Lawrence, *Recommendations to mitigate against human health risks incurred due to energetic particle irradiation beyond low earth orbit/BLEO, Acta Astronautica* **109** (2015), 182.

Mayur Madhavan, Zachary S. Nevin, H. Elizabeth Shick, Eric Garrison, Cheryl Clarkson-Paredes, Molly Karl, Benjamin L. L. Clayton, Daniel C. Factor, Kevin C. Allan, Lilianne Barbar, Tanya Jain, Panagiotis Douvaras, Valentina Fossati, Robert H. Miller, Paul J. Tesar. *Induction of myelinating oligodendrocytes in human cortical spheroids*. *Nature Methods*, 2018; DOI: 10.138/s41592-018-0081-4.

H. Nagasawa and J. B. Little, *Induction of sister chromatid exchanges by extremely low doses of alpha-particles, Cancer Res.* **52** (1992), 6394.

L. Narici, *Heavy ions light flashes and brain functions: recent observations at accelerators and in spaceflight, New Jour. Phys.* **10** (2008), 075010.

D. Neuhof, A. Ruess, F. Wenz and K. J. Weber, *Induction of telomerase activity by irradiation in human limphoblasts, Radiat. Res.* **155** (2001), 693.

K.A. Olive et al. (Particle Data Group), Chin. Phys. C, 38, 090001 (2014).

O. W. Petersen, L. Rønnov-Jessen, A. R. Howlett and M. J. Bissell, *Interaction with basement membrane serves to rapidly distinguish growth and differentiation pattern of normal and malignant human breast epithelial cells, Proceedings of the National Academy of Sciences* **89** (19) (1992), 9064-9068.

M. S. Pino and D. C. Chung, *The chromosomal instability pathway in colon cancer, Gastroenterology,* **138** (6) (2010), 2059.

A. G. M. T. Powell, S. Soul, A. Christian and W. G. Lewis, *Meta-analysis of the prognostic value of CpG island methylator phenotype in gastric cancer*, *British Journal of Surgery* **105** (2018), e61.

D. L. Preston, D. A. Pierce, Y. Shimizu, H. M. Cullings, S. Fujita, S. Funamoto and K. Kodama, *Effect of recent changes in atomic bomb survivor dosimetry on cancer mortality risk estimates*, *Rad. Res.* **162** (2004), 377.

N. Rodriguez-Salas, G. Dominguez, R.Barderas, M. Mendiola, X. García-Albéniz, J. Maurel and J. Feliu Batlle, *Clinical relevance of colorectal cancer molecular subtypes*, *Critical Reviews in Oncology/Hematology* **109** (2017), 9.

N. L. Segal, *The 16^{th} International Twin Congress*, *Twin Research and Human Genetics*, **21** (1) (2018), 67.

J. D. Shadley, V. Afzal nad S. Wolff, *Characterization of the Adaptive Response to Ionizing Radiation Induced by Low Doses of X Rays to Human Lymphocytes*, Radiation Research 111 (1987), 511-517.

G. Sonnenfeld, J. Butel, W. Shearer, *Effects of the space flight environment on the immune system*, *Rev. Environ. Health* **18** (2003), 1.

D. M. Sridharan, A. Asaithamby, S. M. Bailey, S. V. Costes, P. W. Doetsch, W. S. Dynan, A. Kronenberg, K. N. Rithidech, I. Saha, A.M. Snijders, E. Werner, C. Wiese, F. A. Cucinotta and I. M. Piotr, *Understanding Cancer Development Processes after HZE-Particle Exposure: Roles of ROS, DNA Damage Repair and* Inflammation, Rad. *Res.* **183** (2015), 1.

E. Stoupel et Al., *Cardiac Arrhythmia and Geomagnetic Activity*, *Indian Pacing Electrophysical J.* **6** (1) (2006), 49.

S. Tanaka, *Precision medicine based on surgical oncology in the era of genome-scale analysis and genome editing technology, Ann Gastroenterol Surg.* **2** (2018), 106.
J. Vernon Bailey, Chapter 3: Radiation Protection and Instrumentation, in {\it SP-368 Biomedical Results of Apollo, (Richard S. Johnston, Lawrence F. Dietlein and Charles A. Berry Editors), Scientific and Technical Information Office, National Aeronautics and Space Administration Washington, D.C., 1975. http://history.nasa.gov/SP-368/sp368.htm

UNCR Report, *Sources and Effects of Ionizing Radiation*, United Nations Scientific Committee on the Effects of Atomic Radiation, 1993 Report to the General Assembly, with scientific annexes. United Nations sales publication E.94.IX.2. United Nations, New York, 1993.

United Nations Scientific Committee on the Effects of Atomic Radiation. "Sources and effects of ionizing radiation. UNSCEAR 2000 report to the General Assembly, with scientific annexes. Volume II: Effects." (2000).

UNSCEAR 2006 Report, *Effects of Ionizing Radiation*, Volume II, United Nations, New York, 2009.

UNSCEAR 2012 White Paper, *Biological Mechanisms of Radiation Actions at Low Doses*, A white paper to guide the future programme of work of the United Nations Scientific Committee on the Effects of Atomic Radiation, (United Nations, New York, 2012).

K. von der Mark, V. Gauss, H. von der Mark and P. Müller. *Relationship between cell shape and type of collagen synthesised as chondrocytes lose their cartilage phenotype in culture*, Nature **267** (5611) (1977), 531.

L. Walsh, U. Schneider, A. Fogtman, C. Kausch, S. McKenna-Lawlor, L. Narici, J. Ngo-Anh, G. Reitz, L. Sabatieer, G. Santin, L. Sihver, U. Straube, U. Weber and M. Durante, Research plans in Europe for radiation health hazerd assessment in exploratory space missions, Life Sciences in Space Research **21** (2019), 73-82.

K. M. Yamada and E. Cukierman, *Modeling tissue morphogenesis and cancer in 3D, Cell* **130** (4) (2007), 601-610.

J. C. Yoon, A. Ng, B. H. Kim, A. Bianco, R. J. Xavier and S. J. Elledge, *WNT signaling regulates mitochondrial physiology and insuling sensitivity*, *Genes & Development* **24** (14) (2010), 1507.

F. A. Zaman, L. W. Townsend, N. T. Burahmah, *Radiation Risks in a Mission to Mars for a Solar Particle Event Similar to the AD 993/4 Event, Aerospace* **8** (2021), 43.

C. Zeitlin, D. M. Hassler, F. A. Cucinotta, B. Ehresmann, R. F. Wimmer-Schweingruber, D. E. Brinza, S. Kang, G. Weigle,S. Böttcher, E. Böhm, S. Burmeister, J. Guo, J. Köhler, C. Martin, A. Posner, S. Rafkin and G. Reitz, *Measurements of energetic particle radiation in transit to Mars on the Mars Science Laboratory, Science* **340** (2013), 1080.