Xanthomonas infection transforms the apoplast into an accessible and habitable niche for

Salmonella enterica

Megan H. Dixon[a,b], Kimberly N. Cowles[a], Sonia C. Zaacks[a], Isabel N. Marciniak[a], and Jeri D.

Barak[a*]

[a]Department of Plant Pathology, University of Wisconsin, Madison, Wisconsin, USA;

[b]Microbiology Doctoral Training Program, University of Wisconsin, Madison, Wisconsin, USA

Running Head: Water-Soaking Protects *Salmonella enterica*

Address correspondence to Jeri D. Barak, jeri.barak@wisc.edu

Abstract word count: 215 (143 for *Importance* section)

Text word count: 4,863 (excluding Materials and Methods, References, Tables, and Figure

Legends)

ABSTRACT

The physiology of plant hosts can be dramatically altered by phytopathogens. *Xanthomonas hortorum* pv. *gardneri* is one such pathogen that creates an aqueous niche within the leaf apoplast by manipulating the plant via the transcription activator-like effector AvrHah1. Simultaneous immigration of *X. gardneri* and *Salmonella enterica* to healthy tomato leaves results in increased survival of *S. enterica* as *Xanthomonas* infection progresses. However, the fate of *S. enterica* following arrival on actively infected leaves has not been examined. We hypothesized that the water-soaking caused by *X. gardneri* could facilitate the ingression of *S. enterica* into the apoplast, and that this environment would be conducive for growth. We found that a water-soaked apoplast, abiotically or *Xanthomonas*-infected, enabled surface *S. enterica* to passively localize to the protective apoplast and facilitated migration of *S. enterica* to distal sites within the aqueous apoplast. *AvrHah1* contributed to the protection and migration of *S. enterica* early in *X. gardneri* infection. *Xanthomonas*-infected apoplasts facilitated prolonged survival and promoted *S. enterica* replication compared to healthy apoplasts. Access to an aqueous apoplast in general protects *S. enterica* from immediate exposure to irradiation whereas, the altered environment created by *Xanthomonas* infection provides growth-conducive conditions for *S. enterica*. Overall, we have characterized an ecological relationship in which host infection converts an unreachable niche to a habitable environment.

Importance: Bacterial spot disease caused by *Xanthomonas* species devastates tomato production worldwide. Salmonellosis outbreaks from consumption of raw produce have been linked to the arrival of *Salmonella enterica* on crop plants in the field via contaminated irrigation water. Considering that *Xanthomonas* is difficult to eradicate, it is highly likely that *S. enterica*

arrives on leaves pre-colonized by *Xanthomonas* with infection underway. Our study

demonstrates that infection and disease fundamentally alter the leaf, resulting in redistribution

and change in abundance of a phyllosphere bacterial member. These findings contribute to our

understanding of how *S. enterica* manages to persist on leaf tissue despite lacking the ability to

liberate nutrients from plant cells. More broadly, this study reveals a mechanism by which

physiochemical changes to a host environment imposed by a plant pathogen can convert an

uninhabitable leaf environment into a hospitable niche for select epiphytic microbes.

INTRODUCTION

Within an environment as dynamic and complex as the phyllosphere, water is one of the most critical factors for microorganism survival (1). The waxy cuticle limits the amount of moisture available to bacteria that colonize leaves. However, surface bacteria have been known to colonize particularly leaky areas of the cuticle such as leaf veins, cell junctions, and the base of glandular trichomes (2–4). Some phytopathogenic bacteria have adapted to the water scarcity issue by creating their own aqueous environment within the apoplast of leaf tissues (5–7). Infection relies on first entering the leaf through stomata and then colonizing the extracellular internal space, or apoplast, within the leaf. *Xanthomonas hortorum* pv. *gardneri* (hereafter, *X. gardneri*), a causal agent of bacterial spot disease of tomato and pepper, is one such plant pathogen that infects leaves. Once *X. gardneri* has actively migrated to the apoplast, it can deploy its type-III secretion system to manipulate the host into leaking cellular constituents into this space (6), creating an aqueous environment. These physiochemical modifications greatly benefit *X. gardneri* but devastate its host. Bacterial spot disease often results in necrotic lesions on leaves and fruit and defoliation (8) which are detrimental to the health of the plant.

Diseased plants are hosts to a diverse community of microbes, which can be influenced by the drastic changes to their environment caused by phytopathogen infection (9–11). Microbes that are non-pathogenic to plants can benefit from disease. For example, enteric human pathogens such as *Salmonella enterica* lack enzymes that would enable the liberation of nutrients directly from plant cells (12, 13) and are typically forced to scavenge scant resources on the leaf surface. On healthy plant tissues, *S. enterica* populations decline over time (14, 15). However, plant pathogens and even insect pests that exist in agricultural fields have been shown to enhance *S. enterica* populations on plant leaves, and such organisms that can promote *S. enterica*

persistence and/or replication have been coined "biomultipliers" (15–19). Bacterial pathogens

including *Xanthomonas* spp. and *Pectobacterium carotovorum* have been implicated as

biomultipliers for *S. enterica* on leaves (15–17, 20, 21). The mechanisms explaining how these

organisms benefit *S. enterica* are not well understood, posing ecological questions with

important agricultural and human disease epidemiological implications.

The prevalence of food-borne illness outbreaks linked to consumption of raw produce has

climbed in the past few decades (22). Such outbreaks have been linked to pre-harvest

contamination of crops in the field via irrigation water (23, 24). The presence of enteric

pathogens within internal plant tissues has been a concern of food safety researchers, especially

considering efforts to mitigate outbreaks through post-harvest sanitation of produce. Various

studies investigated *S. enterica* colonization of the leaf apoplast, and the extent to which *S.*

enterica can colonize this protective niche varies based on host and method. The flagella-

dependent localization of *S. enterica* to internal tissue in response to light has been demonstrated

via confocal microscopy of lettuce leaf pieces submerged in *S. enterica* inoculum (25). However,

tomato leaves have been shown to be impermeable to *S. enterica* internalization when compared

with lettuce leaves under comparable conditions (26, 27). Whether or not phytopathogen

infection could enhance the permeability of leaves to enteric pathogens was not addressed by

these studies.

Xanthomonas infection promotes *S. enterica* survival when both the plant and human

pathogen arrive on healthy tomato leaves simultaneously (15, 16, 21), however, the fate of *S.*

enterica on leaves actively infected with *Xanthomonas* spp. has not been described. *X. gardneri*

infection can facilitate the migration of surface *X. gardneri* cells to distal tissue via absorption of

X. gardneri suspensions into needle-damaged tissue (6), suggesting that water-soaking may

promote the ingression of bacteria into the apoplast of leaves when external water is present.

Water-soaking has been demonstrated to be enhanced by *X. gardneri* type-III effector AvrHah1

(6, 28). *AvrHah1* encodes a transcription activator-like effector that binds host DNA and

upregulates the activity of a host pectate lyase (6), forcing the host to degrade the pectin

"cement" that joins adjacent plant cells, resulting in an aqueous apoplast (29). Data from our lab

has shown that *avrHah1* expression by *X. gardneri* is necessary for enhancing the persistence of

S. enterica on tomato leaves when co-inoculated with *X. gardneri* (21). Based on these

observations, we hypothesized that *avrHah1*-dependent water-soaking and aqueous apoplasts in

general could serve as a mechanism for epiphytic *S. enterica* to enter protected sites of leaves.

The goal of our study was to examine the fate of epiphytic *S. enterica* following arrival

on the surface of infected leaves. We examined whether an abiotically water-soaked or

Xanthomonas-infected apoplast could facilitate the relocation of *S. enterica* to protected niches,

particularly in the absence of leaf wounding. *S. enterica* populations were considered protected if

they survived UV irradiation applied to leaves. We also identified the roles of *avrHah1*,

Xanthomonas infection progress, and *S. enterica* motility on *S. enterica* entry into the apoplast

and subsequent migration beyond the arrival site. Lastly, we investigated whether the aqueous

niche created by *Xanthomonas* species enhanced the prolonged survival of *S. enterica* within the

apoplast to determine whether factors aside from water availability contribute to *S. enterica*

survival in the apoplast. Identifying host and microbial factors that influence the redistribution

and survival of a model epiphyte will broaden our understanding of microbe-facilitated

adaptations by plant-associated bacteria.

METHODS

Bacterial Strains, Media, and Culture Conditions

Bacterial strains used in this study are described in Table 1. *Xanthomonas* cultures were prepared

by inoculating nutrient broth (NB) with frozen cells and incubating at 28°C with shaking at 200

rpm for two days. *S. enterica* cultures were prepared similarly but grown in lysogeny broth (LB)

and incubated overnight. Cultures were prepared to desired cell concentrations via dilution in

sterile water as described below.

Plant Infiltration

For all experiments, *Solanum lycopersicum* (tomato) cultivar Moneymaker seedlings were

cultivated in growth chambers with a 16 hr photoperiod at 24°C. For *Xanthomonas* infiltration

experiments, 5-week-old tomato plants were infiltrated with *Xanthomonas* and inoculated with

droplets of *S. enterica* as described previously (6) with modifications (Fig. 1). *Xanthomonas*

cultures were prepared to an optical density at 600 nm (OD_{600}) = 0.3, 0.25, or 0.2 (10^8 CFU/mL)

for *X. gardneri* WT, *X. gardneri avrHah1^{ADBD},* and *X. vesicatoria*, respectively, and infiltrated

into two leaflets per plant on the third true leaf via needleless syringe. Infiltration medium was

gently forced into the leaf tissue through the abaxial leaf surface until the infiltration zone was

approximately 2 cm long, and then the infiltration zone was outlined with permanent marker.

When applicable, a subset of plants was infiltrated with sterile water as a control. When *S.*

enterica infiltration was utilized as a control for one experiment, similar methods were applied,

although the *S. enterica* culture used for infiltration was prepared to a concentration of 10^7

CFU/mL and infiltrations were performed in a biosafety cabinet.

Following infiltration, plants were then incubated in bins with lids on at a relative

humidity (RH) > 90% for 1-2 days prior to subsequent *S. enterica* application. For some

experiments, infiltration with sterile water was delayed for a subset of plants until immediately

prior to *S. enterica* inoculation to simulate arrival of *S. enterica* on healthy water-soaked tissue.

These plants were incubated alongside the already infiltrated plants prior to their infiltration such

that environmental conditions would be consistent among all plants within an experiment.

Droplet Inoculation

Lids of plant bins were removed to inoculate leaflets with *S.* enterica, effectively reducing the

RH to ~70%. Infiltrated leaflets were marked again with permanent marker on the *S. enterica*

arrival day to designate the site of *S. enterica* droplet placement on the infiltration zone. *S.*

enterica cultures were prepared to an OD_{600}= 0.1 (10^8 CFU/mL) and 15 µL droplets of the

normalized *S. enterica* culture were applied to the adaxial surface within the part of the

infiltration zone closest to the leaf margin. For one experiment in which *S. enterica* was applied

at a lower inoculation level, the normalized culture was diluted 1000-fold prior to inoculation.

Note that droplets were applied such that they did not overlap with any minor tissue damage

caused by the needleless syringe infiltration from the abaxial surface. Plants inoculated with *S.*

enterica remained inside plastic bins, and the lids were kept off (RH ~70%) until droplets were

no longer visible (i.e., absorbed or evaporated). Visual observations of droplet absorption

(complete, partial, or none) were recorded at 2.5 hours after droplet application. When specified,

a micropipette was used for removing each remaining droplet and measuring their volume. Leaf

tissue was then sampled destructively at the indicated time post-*S. enterica* arrival to quantify

bacterial populations. For experiments requiring sampling beyond 2.5 hours post-*S. enterica*

arrival, bins lids were replaced to maintain plants at >90% RH until sampling.

Bacterial Population Sampling

Whole leaflets were removed at the petiolule and when specified, treated with 254 nm UV

radiation (Stratalinker® UV Crosslinker 1800) at 600,000 μJ/cm^2 with the adaxial leaf surface

facing upwards towards the bulb in order to inactivate surface *S. enterica* cells and enrich for

apoplastic *S. enterica*. The effect of 600,000 μJ/cm^2 of UV irradiation on surface *S. enterica*

populations was determined by removing surface *S. enterica* cells 1 day post-droplet inoculation

using a cotton swab dipped in 0.1% Tween-20 and gently wiping it across the adaxial surface of

the leaf for 5 seconds. The tip of the cotton swab was vortexed in 500 uL of water for 5 seconds,

and then the cell mixture was dilution-plated to enumerate surface *S. enterica* populations (Fig.

S1).

Leaflets irradiated with UV were chosen at random between the two leaflets sampled per

plant and the second leaflet was left non-irradiated as a control. Bacterial populations were

measured as described previously (18). Unless noted otherwise, all inoculated leaflets were

sampled regardless of observations made at the droplet absorption stage. Briefly, leaf tissue was

excised using a sterile 1 cm diameter tissue corer and homogenized in 500 μL of sterile water

using a hand Dremel. For *S. enterica* droplet-inoculated tissue, the site of *S. enterica* arrival was

sampled by centering the tissue corer between the marker guidelines. When indicated, tissue was

also sampled distal from the arrival site by excising a second tissue core 1-2 mm away from the

first leaf disk overlapping the infiltration zone. Leaf tissue homogenate was dilution-plated on

LB agar amended with 50 μg/mL kanamycin for enumeration of *S. enterica* populations. When

S. enterica was applied at a low dosage, the leaf tissue homogenate that remained after plating

was enriched with LB amended with kanamycin (50 ug/mL) to determine if *S. enterica* was

present, but below the limit of detection for dilution-plating. LB-enriched samples were plated

22-24 hours later. *S. enterica* colonies were enumerated after 1 day of incubation at 37°C.

Statistical Analyses

R Studio version 2021.09.1 (R Development Core Team, R Foundation for Statistical

Computing, Vienna, Austria [http://www.R-project.org]) was utilized for all statistical analyses

and plot generation. Each experiment was repeated three times, and each boxplot represents the

aggregation of three independent experiments found to be statistically similar ($P > 0.05$), unless

noted otherwise in the figure legend. Within each experiment, the number of biological replicates

$N = 4$-8 leaflets. For comparing *S. enterica* populations among treatment groups at the arrival

tissue site, the ANOVA and post-hoc Tukey tests were used. When two leaflets were inoculated

per plant, the data were fit to a linear mixed effects model using the lme4 package in which

individual plants were modeled as a random effect. *S. enterica* population data at distal tissue

sites were generally not found to be normally distributed due to zeroes resulting from the

absence of *S. enterica* migration beyond the initial site of arrival. Therefore, for comparing *S.*

enterica populations among treatment groups at the distal tissue site, the non-parametric

Kruskal-Wallis and post-hoc Wilcox tests were used. In addition, the proportion of samples with

detectable bacterial populations at the distal tissue site were compared using the Fisher's Exact

Test to compare the incidence of *S. enterica* colonization of distal tissue. For the low *S. enterica*

dosage experiment, data from the healthy/dry treatment was excluded from the mixed effects

model because the value of nearly every data point was zero. Preliminary analysis indicated that

for the remaining two treatments (healthy/wet and *X. gardneri*-infected), all means were greater

than zero based on one-way t-tests ($P < 0.05$). When applicable, the limit of detection minus one

(i.e., 49 CFU/leaf disk) was substituted for non-detects. The significance threshold for all

comparisons was $P < 0.05$.

RESULTS

Abiotically water-soaked apoplasts enable *S. enterica* to migrate via stomata to a

UV-protected niche. To characterize the role of water in the apoplast in the migration of *S.*

enterica to a protected niche and migration to sites distal to the arrival site, we developed a

method involving infiltration of intact tomato leaflets, application of droplets of *S. enterica* cell

suspensions, and measurement of UV-protected *S. enterica* populations (Fig. 1). Our methods

were inspired by work previously published by Schwartz et al. (6), with a few key differences

that included applying bacterial droplets to intact (not detached) leaflets onto the adaxial surface,

refraining from wounding the leaves at the droplet application site, and utilizing UV irradiation

to enrich for apoplastic *S. enterica* populations. We found that our UV irradiation methods

reduced the *S. enterica* populations on the leaf surface by 2.9 logs (Fig. S1). Additionally, we

manipulated the timing of *S. enterica* arrival on water-infiltrated leaves to observe how the

presence of water in the apoplast influences the relocation of *S. enterica*.

Infiltration of water into the apoplast simulated abiotic water-soaking that may occur

during periods of high precipitation (30). Under the tested conditions (~70% RH), water

infiltration created an obvious darkening or "wet" appearance to the tissue that lasted for a few

hours (data not shown). *S. enterica* droplets were applied to two treatments: leaves with a (i)

"healthy/dry" apoplast where the tissue had become macroscopically dry at time of *S. enterica*

application at 2 day post-infiltration (DPI) or (ii) "healthy/wet" apoplast where tissue was still

visibly dark and wet in appearance when *S. enterica* was applied within 10 minutes of water

infiltration (0 DPI). A third treatment was included in which we infiltrated *S. enterica* directly

into the apoplast to determine the relative amount of UV protection provided to apoplastic *S.*

enterica cells.

Within a few hours of *S. enterica* arrival on the leaf, we observed the disappearance of

the droplets specifically into healthy/wet tissue in the absence of any applied epidermal

wounding, suggesting that the *S. enterica* suspension had been absorbed through stomata.

Leaflets were observed for a duration of 2.5 hours post-application of *S. enterica,* and absorption

of the entire 15 µL suspension could be observed within a minimum of 30 minutes. The success

rate for complete absorption of the droplet for healthy/wet tissue varied between 50-83% among

three experimental replicates, although droplets that failed to be completely absorbed within the

2.5 hour timeframe had visibly reduced in size, suggesting that they were partially absorbed. By

contrast, healthy/dry leaflets had an absorption rate of 0%. Beyond the observation period, after

several hours had passed, all droplets disappeared either by absorption, evaporation, or both. *S.*

enterica populations were enumerated at 1 day post-arrival, and among the three treatments, UV

irradiation resulted in the lowest *S. enterica* populations when *S. enterica* was applied to leaflets

with a healthy/dry apoplast (Fig. 2). These surface-limited populations were reduced ~2.5 logs

by UV irradiation. Arrival on healthy/wet tissue and direct infiltration into the apoplast resulted

in equivalent UV-protected populations and relatively small population reductions of ~0.9 and

~0.7 logs, respectively, in response to UV irradiation (Fig. 2). The increased UV protection

conferred to *S. enterica* when applied to healthy water-soaked tissue suggests that an aqueous

apoplast provides a gateway for surface bacteria to move inside the leaf.

Next, we investigated whether movement of surface bacteria to the apoplast was limited

to the immediate substomatal chamber or extended beyond. We observed distinct differences in

the ability of *S. enterica* to migrate from the arrival site to tissue non-overlapping the droplet

application site depending on the physical state of the apoplast. A healthy/dry apoplast did not

readily facilitate lateral migration of *S. enterica* to areas within the apoplast distal from the

arrival site whereas, a healthy/wet apoplast enabled recovery of significantly higher *S. enterica*

populations at distal sites (Fig. 2). Distal sites within the apoplast provided *S. enterica* with a

protected niche if the apoplast was an aqueous environment at time of *S. enterica* arrival, as the

population reduction in response to UV irradiation was only ~0.5 logs (Fig. 2). These findings

support the conclusion that an aqueous apoplast permits surface bacteria to migrate beyond the

initial arrival site.

***Xanthomonas* water-soaking promotes *S. enterica* entry to the apoplast and**

facilitates migration to distal tissue. *X. gardneri* is known to cause water-soaking (6, 28), and

thus we examined whether *X. gardneri* infection could confer protection to *S. enterica* against

UV irradiation and facilitate its migration to infected sites distal from the initial site similarly to a

healthy/wet apoplast. We found that *X. gardneri*-infected tissue enabled complete absorption of

S. enterica droplets into the leaves at 1 DPI (Fig. S2) and 2 DPI (images not shown) within 0.5-

2.5 hours. To assess the protection of absorbed *S. enterica* cells against UV, we infiltrated an

excess number of plants with *X. gardneri*, and the first 50% of plants that completely absorbed

the droplets were specifically selected for *S. enterica* population sampling at 1 day post-*S.*

enterica arrival. *S. enterica* cells were found to be more protected against UV irradiation

following droplet absorption into *X. gardneri*-infected tissue at 2 DPI compared to *S. enterica*

cells that were limited to the surface of healthy/dry tissue (Fig. 3). Like a healthy/wet apoplast, a

X. gardneri-infected apoplast facilitated robust colonization of *S. enterica* cells to sites in the

apoplast beyond the initial arrival site, where *S. enterica* was protected from UV irradiation (Fig.

3b).

276   ***AvrHah1* facilitates *S. enterica* localization to the apoplast and lateral migration**

early in infection. Knowing that an aqueous apoplast facilitates the migration of *S. enterica* to a

protected niche within a day of arrival, we examined whether protection against UV irradiation

exists as early as 2.5 hours post-arrival, a time point at which *S. enterica* droplet suspensions

were often completely absorbed. Also, we compared the fate of *S. enterica* following arrival on

tissue infected by one of three *Xanthomonas* strains including *X. gardneri* WT, *X. gardneri*

avrHah1$^{\Delta DBD}$, and *X. vesicatoria*. *X. vesicatoria* is a pathogen within the bacterial spot disease

complex that lacks *avrHah1* and has been shown in co-inoculation experiments with *S. enterica*

to not promote *S. enterica* growth on tomato leaves, in contrast to *X. gardneri,* which does

facilitate *S. enterica* replication (15, 21).

286   The duration of infection with *Xanthomonas* spp. was found to affect visible water-

soaking, absorption of the *S. enterica* suspension, and *S. enterica* protection from UV irradiation

(Fig. 4). Tissue infected with *Xanthomonas* spp. was generally water-soaked at 1 and 2 DPI,

except for leaflets infected with the *X. gardneri avrHah1*$^{\Delta DBD}$ strain at 1 DPI (Fig. S3). Leaflets

infected with *X. vesicatoria* became water-soaked at a similar rate as *X. gardneri* WT-infected

leaflets despite lacking the *avrHah1* gene. As for effects on *S. enterica*, tissue that was

macroscopically dry at *S. enterica* arrival, which included the healthy/dry and *X. gardneri*

avrHah1$^{\Delta DBD}$ 1 DPI treatments, did not absorb the *S. enterica* suspension, resulting in the highest

droplet recovery volumes at 2.5 hours post *S. enterica* droplet application (Fig. 4a). In contrast to

dry tissue, the recovered droplet volumes were lower for water-soaked tissue with healthy/wet,

X. gardneri WT-infected, or *X. vesicatoria*-infected leaves, and these tissues often absorbed the

S. enterica suspension completely (Fig. 4a). Congruently, *S. enterica* arrival on water-soaked

tissues resulted in the greatest protection, as evident by smaller reductions in population in

response to UV (Fig. 4b). Overall, infection progress had a significant effect on *S. enterica*

protection in our model.

In addition to facilitating initial entry of *S. enterica* into the substomatal chamber,

avrHah1 was also found to contribute to the ability of *S. enterica* to migrate to distal tissue

beyond the site of arrival early in infection (Fig. 5). Leaves infiltrated with *X. gardneri*

avrHah1^{ΔDBD}, which generally had a dry apoplast at 1 DPI (Fig. S3) and did not result in visible

absorption of the *S. enterica* suspension (Fig. 4a), exhibited reduced migration of *S. enterica* to

distal tissue compared to leaves infiltrated with *X. gardneri* WT (Fig. 5). However, by 2 DPI,

when the tissue was visibly water-soaked (Fig. S3), *X. gardneri avrHah1^{ΔDBD}*-infected tissue did

facilitate colonization of distal tissue by *S. enterica* (Fig. 5). These findings demonstrate that the

role of *avrHah1* in altering the apoplast into an aqueous environment occurs early in the disease

progress, and these changes facilitate surface bacteria entry into the apoplast and subsequent

lateral migration to other water-soaked areas. We also note that additional bacterial factors

besides *avrHah1* contribute to water-soaking later in disease progression.

Active motility of *S. enterica* increases colonization of tissue beyond the arrival site

as *X. gardneri* infection progresses. To test the hypothesis that active *S. enterica* motility

contributes to migration away from the surface, bacterial populations for a *S. enterica* Δ*motA*

mutant with non-functional flagella were compared to a wild-type *S. enterica* strain. At 2.5 hours

post-*S. enterica* arrival, the Δ*motA* mutant colonized UV-protected sites of water-soaked leaves

equivalently to the wild-type *S. enterica* strain (Fig. S4), indicating that motility does not affect

S. enterica's ability to enter the aqueous apoplast. We also compared *S. enterica* WT and Δ*motA*

populations at tissue distal from the arrival site at 1 day post-*S. enterica* arrival. Motility did not

affect *S. enterica* colonization of distal tissue infected with *X. gardneri* when *S. enterica* arrived

at 1 DPI, however, at 2 DPI, *S. enterica* populations at distal sites were enhanced by motility

(Fig. 5). Deletion of *motA* did not eliminate *S. enterica* colonization of distal tissue nor affect the

proportion of distal sites with detectable *S. enterica* populations, indicating that motility was

useful but not required for lateral migration of *S. enterica*. Overall, motility was not required for

entry into the apoplast or migration beyond the initial arrival site, supporting a model of passive,

rather than active, *S. enterica* relocation within water-soaked tissue.

Prolonged survival of *S. enterica* in a protected niche is increased by *Xanthomonas*

infection. Having established that aqueous apoplasts result in similar migration of *S. enterica* to

UV-protected niches regardless of whether the apoplast is abiotically water-soaked or water-

soaked via *Xanthomonas* infection, we then tested the hypothesis that the apoplast of

Xanthomonas infected leaves would support higher *S. enterica* population growth than healthy

leaves following *S. enterica* application at 1 DPI. We found that *S. enterica* arrival on *X.*

gardneri WT or *X. vesicatoria*-infected tissue resulted in the highest UV-protected *S. enterica*

populations at three days post-arrival (Fig. 6). Although a healthy/wet apoplast facilitated equal

colonization of *S. enterica* within UV-protected areas compared to *X. gardneri* WT and *X.*

vesicatoria infected tissue at 2.5 hours post-*S. enterica* arrival, (Fig. 4b) the healthy/wet apoplast

supported lower *S. enterica* populations at the arrival and distal sites relative to infected/wet

apoplasts by 3 days post-*S. enterica* arrival (Fig. 6). *X. gardneri avrHah1^{ADBD}*-infected plants (1

DPI) did not permit droplet absorption (Table S1), yielded intermediate UV-protected *S. enterica*

populations at the arrival and distal sites (Fig. 6), and resulted in the most variable *S. enterica*

populations by 3 DPA (Fig. 6). Healthy/dry tissue also did not absorb the droplets (Table S1) and

supported the lowest *S. enterica* populations at the arrival and distal sites (Fig 6).

We also tested the hypothesis that *X. gardneri*-infected apoplasts support higher *S.*

enterica growth when *S. enterica* cells are applied to the leaves at a 1000 times lower inoculum

level (10^3 CFU/droplet) relative to the previous experiments. Reducing the *S. enterica* inoculum

dosage minimizes the possibility of overwhelming the plant immune response and more

realistically models foliar contamination of crops with *S. enterica*. Arrival of *S. enterica* on

healthy/dry leaves led to eradication of *S. enterica* populations in response to UV irradiation

(Fig. 7). Meanwhile, when *S. enterica* was applied to leaves with an aqueous apoplast (healthy or

X. gardneri WT-infected) it localized to UV-protected niches and migrated to distal tissue within

2.5 hours (Fig. 7). Additionally, the UV-protected populations increased between day 0 (2.5

HPA) and day 3 (3 DPA) (Fig. 7). *X. gardneri*-infected apoplasts supported the highest *S.*

enterica growth, resulting in *S. enterica* populations that increased at least 3.8 logs within 3 days

of arrival (Fig. 7). The higher apoplastic *S. enterica* populations observed in infected tissue

substantiates the conclusion that prolonged survival in the apoplast by surface bacteria is

supported by an aqueous environment resulting from phytobacterial infection, which is distinct

from a healthy, watery apoplast.

DISCUSSION

In this study, we determined the fate of *S. enterica* following arrival on *Xanthomonas*-

infected tissue, specifically regarding ingression into the apoplast and migration to distal sites.

We hypothesized that the modifications to the host caused by *Xanthomonas* infection could

facilitate ingression of *S. enterica* into the apoplast, where it could migrate to distal sites and

thrive. *S. enterica* was applied to leaves as a droplet to model splash dispersal of bacteria in the

field, which can occur during irrigation or precipitation events. Various host and microbial

factors including the presence of water in the apoplast, flagellar function by *S. enterica,* and

avrHah1 expression by *X. gardneri* were manipulated to understand how alterations to the

infection court, the apoplast, impact plant-associated surface bacteria.

Colonization of protected niches such as the apoplast is essential to the lifestyle of

various plant-associated bacteria. To determine whether *S. enterica* could localize to protected

niches of leaves, UV irradiation at 254 nm, which is an FDA-approved method for

decontamination of food surfaces (31) and has been applied in previous plant bacterial endophyte

studies (32, 33), was utilized to reduce surface *S. enterica* populations. The use of UV irradiation

additionally models the UV exposure that plant-associated microbes are routinely exposed to,

though it's worth noting UV wavelengths less than 300 nm are typically absorbed by our

atmosphere and do not reach Earth (34). Initially, we tested whether *S. enterica* cells that were

infiltrated directly into the apoplast of leaves would be protected from UV irradiation applied to

the adaxial surface, and we hypothesized that the palisade cells, upper epidermal cells, and

cuticle would occlude UV penetration to the apoplastic *S. enterica*. We determined that *S.*

enterica cells localized to the apoplast were more protected from UV compared to *S. enterica* on

the surface, although UV irradiation reduced the apoplastic *S. enterica* populations a small but

statistically significant amount (Fig. 2). Leaves must permit light to penetrate leaf tissue for

photosynthesis and act as "light traps (35)," which may explain why a subset of apoplastic *S.*

enterica cells were vulnerable to UV. The duration of UV exposure influenced the magnitude of

S. enterica population reduction (data not shown), and the exposure level of 0.6 J/cm^2 selected

for these experiments resulted in >2.0 log reduction of surface *S. enterica* populations with

minimal effects on the apoplastic populations. As for the surface-localized *S. enterica* cells, the

small subpopulations that survived UV irradiation (Fig. S1) likely resisted sterilization by being

canopied by leaf surface structures, such as trichomes, or wedged within the grooves between

adjacent cells (36–38).

S. enterica successfully colonized UV-protected niches of abiotically water-soaked or

Xanthomonas-infected tissue. We rationalized that *S. enterica* had accessed the leaf apoplast

based on a few lines of evidence. First, the extent to which *S. enterica* was protected from UV

irradiation following arrival on abiotically water-soaked or *X. gardneri*-infected apoplast was

similar to cells infiltrated directly into the apoplast (Fig. 2). Second, *S. enterica* droplets were

absorbed by abiotically water-soaked or infected tissue, as evident by rapid disappearance of the

droplets in as little as 30 minutes. Healthy tissue is typically impermeable to spontaneous

absorption of surface water because the surface tension of water is greater than the critical

surface tension of the leaf surface (39). However, the relatively rapid disappearance of droplets

into water-soaked leaves indicated that the wettability of the leaf surface had changed.

Considering that the leaf tissue was deliberately not wounded, open stomata likely served as

entry points for external water and suspended *S. enterica* cells to enter the apoplast. We theorize

that an aqueous apoplast enabled a watery film to extend upwards through the stomatal pores,

providing a continuous "watery highway" that could facilitate movement of *S. enterica* into the

substomatal space. Lastly, *S. enterica* was consistently isolated at high levels from abiotically

water-soaked (Fig. 2) and infected (Fig. 3) tissue ~1 cm distal from the arrival site when the

apoplast was water-soaked at *S. enterica* arrival, but not when it was dry. Schwartz et al. found

that *X. gardneri*-infected leaves facilitated migration of surface *X. gardneri* cells to the apoplast

(6), and here we have confirmed that other bacteria besides the phytopathogen *X. gardneri* could

be transported by water-soaking. To verify that *S. enterica* colonization of distal tissue was not

driven by migration across the surface but by migration through the apoplast, the adaxial surface

of leaf tissue disks were pressed to agar plates following UV exposure. *S. enterica* was rarely

isolated from the surface of leaves distal to the arrival site with a healthy/dry apoplasts (data not

shown). Overall, *S. enterica* was found to relocate from the surface of leaves into the substomatal

chamber and beyond. These observations posed the question of whether *S. enterica* motility is

necessary for the redistribution of cells following *S. enterica* arrival.

Flagellar motility is key to survival on the leaf surface for recent immigrants (40), but we

found that the relocation of *S. enterica* from the leaf surface to the aqueous apoplast can occur

independent of functional flagella (Fig. 5), thus favoring a model of passive *S. enterica*

relocation in the presence of a water-soaked apoplast. However, active motility was found to

provide *S. enterica* with an advantage in colonizing sites beyond the initial arrive site at 2 days

post-*X. gardneri* infection, but not earlier (Fig. 5). *In planta* microscopy of *Pseudomonas*

syringae colonization has revealed that distinct bacterial subpopulations exist within infected

tissue, suggesting that the apoplast is a heterogeneous environment (41). Later in *Xanthomonas*

infection when more nutrients may be available in the apoplast, motile *S. enterica* cells may

employ chemotaxis to colonize nutritional micro-oases within the apoplast. Therefore, as the

Xanthomonas infection court dynamically changes over time, the importance of specific traits

possessed by plant-associated microbes also changes.

In addition to determining how *S. enterica* redistribution is affected by one of its own

genetic traits, we also examined the role of *X. gardneri* type-III effector AvrHah1 to identify

mechanisms by which phytopathogen infection influences the fate of *S. enterica. AvrHah1*

contributed to *S. enterica's* ability to access the protective apoplast (Fig. 4) and migrate to distal

tissue in the early stage of an ongoing *X. gardneri* infection (Fig. 5). AvrHah1 is the only

Xanthomonas effector that has been directly linked to the water-soaking phenotype, despite

water-soaking being a characteristic symptom of bacterial spot disease. Other Xanthomonads

outside of the bacterial spot complex that infect diverse hosts, including leafy greens, cause

water-soaking (42, 43), yet no other water-soaking effectors have been identified within the

Xanthomonas genus. Among the various *Xanthomonas* species that cause bacterial spot disease,

which includes *X. vesicatoria, X. euvesicatoria, and X. perforans, avrHah1* was originally

specific to *X. gardneri* (44, 45). Recently, however, *X. perforans* strains carrying *avrHah1* have

been identified in the southeastern United States (46). *X. vesicatoria,* which lacks *avrHah1,*

induced visible water-soaking of tomato leaves as rapidly as *X. gardneri* and promoted

protection and survival of *S. enterica* equal to or better than *X. gardneri.* These results contrast

with previous studies documenting that *X. vesicatoria* infection does not promote *S. enterica*

persistence on tomato leaf tissue following co-arrival of *X. vesicatoria* and *S. enterica* on healthy

plants (15, 21). However, a key difference between those studies and the study presented here is

the physiological state of the leaf when *S. enterica* arrives: healthy leaves versus infected leaves.

Infected tissue can permit *S. enterica* to enter the apoplast, and therefore, the benefits conferred

to *S. enterica* by *X. vesicatoria* infection may be conditional on *S. enterica* entry. Overall, the

ability of *avrHah1*-lacking *Xanthomonas* strains to facilitate water-soaking mediated ingression

of *S. enterica* into the apoplast and the observed delay, not elimination, of water-soaking in *X.*

gardneri avrHah1$^{\Delta DBD}$ suggest that other factors within the *Xanthomonas* genome independent

of *avrHah1* contribute to creating an aqueous niche within the tomato leaf apoplast. Identifying

these bacterial factors in future studies would improve our understanding of phytopathogen-

mediated water-soaking.

Leaf surfaces are harsh environments that require plant-associated bacteria to flexibly adapt to

extreme and dynamic conditions, including desiccation, heavy rainfall, UV irradiation, and

sporadic nutrient availability (47–50). Many phytopathogenic bacteria have evolved to infect the

apoplast of leaves where they can replicate to high population levels whilst exploiting host

resources (51). The apoplast can be protective and provide bacteria with increased nutrient

availability, however, this environment is exclusive to microbes that can successfully migrate

there and tolerate the host immune response. Tomato leaves are relatively restrictive to *S.*

enterica ingression (26, 27), yet we found that *S. enterica* can access an apoplast that is water-

soaked either by *Xanthomonas* infection or abiotically. Furthermore, *Xanthomonas* infection

resulted in higher *S. enterica* populations within UV-protected niches (Fig. 6) and enhanced *S.*

enterica replication (Fig. 7) relative to healthy leaf controls. *S. enterica's* capacity to exhibit near

4 log population growth within three days of *S. enterica's* arrival on the surface of *X. gardneri*-

infected leaves (Fig. 7) emphasizes the extent that *Xanthomonas* pathogens modify the apoplast.

An infected tomato leaf apoplast relieves suppression of *S. enterica* replication, suggesting that

Xanthomonas infection aids in overcoming the host immune response and/or liberates nutrients

inaccessible to *S. enterica*. *S. enterica* is known to elicit the plant immune response (52) upon

recognition of microbe-associated molecular markers such as flg22 (53, 54), although some

evidence suggests that the host response provoked by *S. enterica* is relatively weak (55, 56). If

modulation of the immune response by *Xanthomonas* does enhance *S. enterica* fitness in the

apoplast, it is likely that this is not the only mechanism involved. The replication of *S. enterica* in

an infected apoplast also suggests that nutrients have been made bioavailable by *Xanthomonas*

infection. *Xanthomonas* spp. presumably liberate nutrients from leaf tissue, as supported by their

own replication to populations above 10^8 CFU/cm^2 (21) and increased electrolyte leakage of

tomato leaf tissue (15). One hypothesis is that an infected apoplast is enriched in carbohydrates,

and *S. enterica* metabolizes these nutrients that it can't access on its own. *S. enterica* can utilize

many different sugars as sole carbon sources (57), and although only 10% of *Salmonella* species

can metabolize sucrose (58), one of the most prevalent carbohydrates available in plant tissues

(51), other sugars such as glucose and fructose (59, 60) could be available to *S. enterica*. Future

studies will be informative in describing *avrHah1*-independent and -dependent mechanisms by

which *Xanthomonas* releases host nutrients and identifying resources scavenged by *S. enterica*

within infected leaves.

Overall, we have identified a common phytobacterial disease symptom of leaf pathogens,

water-soaking, as a mechanism by which *S. enterica* can be passively relocated to the apoplast of

leaves, resulting in access to protected niches including the substomatal chamber directly below

the initial arrival site and distal sites of the apoplast. Our results show that *avrHah1* expression

by *X. gardneri* influences the fate of *S. enterica,* and considering evidence that *avrHah1* is being

transferred between *Xanthomonas* species, gene flow within a plant pathogen complex may have

dire effects on the persistence of enteric human pathogens in crop plants that eventually will be

consumed raw. Altogether, our findings illustrate a model for how enteric bacterial human

pathogens are dispersed in agricultural fields via contaminated irrigation water and subsets of the

bacterial population can relocate from the hostile leaf surface to a habitable niche within the

apoplast (Fig. 7). As phytopathogen populations evolve and respond to global environmental

changes, understanding their effects on the redistribution and survival of other plant-associated

microbes will be critical.

REFERENCES

1. Beattie GA. 2011. Water Relations in the Interaction of Foliar Bacterial Pathogens with Plants. Annu Rev Phytopathol 49:533–555.

2. Schonherr J. 2006. Characterization of aqueous pores in plant cuticles and permeation of ionic solutes. J Exp Bot 57:2471–2491.

3. Monier J, Lindow SE. 2004. Frequency , Size , and Localization of Bacterial Aggregates on Bean Leaf Surfaces. Appl Environ Microbiol 70:346–355.

4. Barak JD, Kramer LC, Hao L-Y. 2011. Colonization of Tomato Plants by Salmonella enterica Is Cultivar Dependent, and Type 1 Trichomes Are Preferred Colonization Sites. Appl Environ Microbiol 77:498–504.

5. Xin XF, Nomura K, Aung K, Velásquez AC, Yao J, Boutrot F, Chang JH, Zipfel C, He SY. 2016. Bacteria establish an aqueous living space in plants crucial for virulence. Nature 539:524–529.

6. Schwartz AR, Morbitzer R, Lahaye T, Staskawicz BJ. 2017. TALE-induced bHLH transcription factors that activate a pectate lyase contribute to water soaking in bacterial spot of tomato. Proc Natl Acad Sci 114:E897–E903.

7. Roper MC, Burbank LP, Williams K, Viravathana P, Tien H-Y, von Bodman S. 2015. A Large Repetitive RTX-Like Protein Mediates Water-Soaked Lesion Development, Leakage of Plant Cell Content and Host Colonization in the Pantoea stewartii subsp. stewartii Pathosystem. Mol Plant-Microbe Interact 28:1374–1382.

8. Goode MJ, Sasser M. 1980. Prevention - the Key to Controlling Bacterial Spot and Bacterial Speck of Tomato. Plant Dis 64:831–834.

9. Ginnan NA, Dang T, Bodaghi S, Ruegger PM, McCollum G, England G, Vidalakis G,

Borneman J, Rolshausen PE, Caroline Roper M. 2020. Disease-induced microbial shifts in

citrus indicate microbiome-derived responses to huanglongbing across the disease severity

spectrum. Phytobiomes J 4:375–387.

10.   Gao M, Xiong C, Gao C, Tsui CKM, Wang MM, Zhou X, Zhang AM, Cai L. 2021.

Disease-induced changes in plant microbiome assembly and functional adaptation.

Microbiome 9:1–18.

11.   Yang F, Zhang J, Zhang H, Ji G, Zeng L, Li Y, Yu C, Fernando WGD, Chen W. 2020.

Bacterial Blight Induced Shifts in Endophytic Microbiome of Rice Leaves and the

Enrichment of Specific Bacterial Strains With Pathogen Antagonism. Front Plant Sci

11:1–14.

12.   Teplitski M, Barak JD, Schneider KR. 2009. Human enteric pathogens in produce : un-

answered ecological questions with direct implications for food safety. Curr Opin

Biotechnol 20:166–171.

13.   Abbott DW, Boraston AB. 2008. Structural Biology of Pectin Degradation by

Enterobacteriaceae. Microbiol Mol Biol Rev 72:301–316.

14.   Islam M, Morgan J, Doyle MP, Phatak SC, Millner P, Jiang X. 2004. Fate of Salmonella

enterica Serovar Typhimurium on Carrots and Radishes Grown in Fields Treated with

Contaminated Manure Composts or Irrigation Water. Appl Environ Microbiol 70:2497–

2502.

15.   Potnis N, Colee J, Jones JB, Barak JD. 2015. Plant Pathogen-Induced Water-Soaking

Promotes Salmonella enterica Growth on Tomato Leaves. Appl Environ Microbiol

81:8126–8134.

16.   Potnis N, Soto-Arias JP, Cowles KN, van Bruggen AHC, Jones JB, Barak JD. 2014.

Xanthomonas perforans Colonization Influences Salmonella enterica in the Tomato

Phyllosphere. Appl Environ Microbiol 80:3173–3180.

17.   Kwan G, Charkowski AO, Barak JD. 2013. Salmonella enterica Suppresses

Pectobacterium carotovorum subsp. carotovorum Population and Soft Rot Progression by

Acidifying the Microaerophilic Environment. MBio 4:1–9.

18.   Cowles KN, Groves RL, Barak JD. 2018. Leafhopper-Induced Activation of the Jasmonic

Acid Response Benefits Salmonella enterica in a Flagellum-Dependent Manner. Front

Microbiol 9:1–15.

19.   Harrod VL, Groves RL, Maurice MA, Barak JD. 2021. Frankliniella occidentalis facilitate

Salmonella enterica survival in the phyllosphere. PLoS One 16:1–16.

20.   Wells JM, Butterfield JE. 1997. Salmonella contamination associated with bacterial soft

rot of fresh fruits and vegetables in the marketplace. Plant Dis 81:867–872.

21.   Cowles KN, Block AK, Barak JD. Xanthomonas hortorum pv. gardneri TAL effector

AvrHah1 is neessary and sufficient for increased persistence of Salmonella enterica on

tomato leaves. Res Sq https://doi.org/10.21203/rs.3.rs-1229600/v1.

22.   Hoagland L, Ximenes E, Ku S, Ladisch M. 2018. Foodborne pathogens in horticultural

production systems : Ecology and mitigation. Sci Hortic (Amsterdam) 236:192–206.

23.   Centers for Disease Control and Prevention (CDC). 2008. Outbreak of Salmonella

Serotype Saintpaul Infections Associated with Multiple Raw Produce Items --- United

States, 2008. Morb Mortal Wkly Rep 57:929–934.

24.   Centers for Disease Control and Prevention (CDC). 2018. Multistate Outbreak of E. coli

O157:H7 Infections Linked to Romaine Lettuce (Final Update).

25.   Kroupitski Y, Golberg D, Belausov E, Pinto R, Swartzberg D, Granot D, Sela S. 2009.

Internalization of Salmonella enterica in Leaves Is Induced by Light and Involves

Chemotaxis and Penetration through Open Stomata   †. Appl Environ Microbiol

75:6076–6086.

26.   Golberg D, Kroupitski Y, Belausov E, Pinto R, Sela S. 2011. Salmonella Typhimurium

internalization is variable in leafy vegetables and fresh herbs. Int J Food Microbiol

145:250–257.

27.   Chahar M, Kroupitski Y, Gollop R, Belausov E, Melotto M, Sela-Saldinger S. 2021.

Determination of Salmonella enterica Leaf Internalization Varies Substantially According

to the Method and Conditions Used to Assess Bacterial Localization. Front Microbiol

12:1–12.

28.   Schornack S, Minsavage G V, Stall RE, Jones JB, Lahaye T, Jones JB. 2006.

Characterization of AvrHah1 , a novel AvrBs3-like effector from Xanthomonas gardneri

with virulence and avirulence activity. New Phytol 179:546–556.

29.   Collmer A, Keen NK. 1986. The Role of Pectic Enzymes in Plant Pathogenesis. Annu Rev

Phytopathol 24:383–409.

30.   Clayton E. 1936. Water-Soaking of leaves in relation to development of the wildfire

disease of tobacco. J Agric Res 52:239–269.

31.   Sommers CH, Sites JE, Musgrove M. 2010. Ultraviolet light (254 nm) inactivation of

pathogens on foods and stainless steel surfaces. J Food Saf 30:470–479.

32.   Wilson M, Hirano SS, Lindow SE. 1999. Location and survival of leaf-associated bacteria

in relation to pathogenicity and potential for growth within the leaf. Appl Environ

Microbiol 65:1435–1443.

33.   Kennedy BW, Ercolani GL. 1978. Soybean Primary Leaves as a Site for Epiphytic

Multiplication of Pseudomonas glycinea. Ecol Epidemiol 68:1196–1201.

34.   Döhring T. 2017. Critical discussion on the UV absorption properties of Earth's

atmosphere.Proceedings of SPIE 10453.

35.   Vogelmann TC, Bjorn LO. 1986. Plants as light traps. Physiol Plantarium 68:704–708.

36.   Doan HK, Ngassam VN, Gilmore SF, Tecon R, Parikh AN, Leveau JHJ. 2020.

Topography-driven shape, spread, and retention of leaf surface water impacts microbial

dispersion and activity in the phyllosphere. Phytobiomes J 4:268–280.

37.   Fino VR, Kniel KE. 2008. UV light inactivation of hepatitis A virus, aichi virus, and

feline calicivirus on strawberries, green onions, and lettuce. J Food Prot 71:908–913.

38.   Huang R, Chen H. 2019. Comparison of Water-Assisted Decontamination Systems of

Pulsed Light and Ultraviolet for Salmonella Inactivation on Blueberry, Tomato, and

Lettuce. J Food Sci 84:1145–1150.

39.   Schonherr J. 1972. Penetration of Stomata by Liquids 813–819.

40.   Haefele DM, Lindow SE. 1987. Flagellar Motility Confers Epiphytic Fitness Advantages

Pseudomonas syringae. Appl Environ Microbiol 53:2528–2533.

41.   Rufian J, Sanchez-Romero M, Lopez-Marquez D, Macho A, Mansfield J, Arnold D, Ruiz-

Albert J, Casadesus J, Beuzon C. 2016. Pseudomonas syringae differentiates into

phenotypically distinct subpopulations during colonization of a plant host. Environ

Microbiol 18:3593–3605.

42.   Sahin F, Miller SA. 1997. Identification of the bacterial leaf spot pathogen of lettuce,

Xanthomonas campestris pv. vitians, in Ohio, and assessment of cultivar resistance and

seed treatment. Plant Dis 81:1443–1446.

43.   Dia NC, Morinière L, Cottyn B, Bernal E, Jacobs JM, Koebnik R, Osdaghi E, Potnis N,

616   Pothier JF. 2022. Xanthomonas hortorum – beyond gardens: Current taxonomy,

617   genomics, and virulence repertoires . Mol Plant Pathol 1–25.

618 44.   Potnis N, Krasileva K, Chow V, Almeida NF, Patil PB, Ryan RP, Sharlach M, Behlau F,

619   Dow JM, Momol MT, White FF, Preston JF, Vinatzer BA, Koebnik R, Setubal JC,

620   Norman DJ, Staskawicz BJ, Jones JB. 2011. Comparative genomics reveals diversity

621   among xanthomonads infecting tomato and pepper. BMC Genomics 12:1–23.

622 45.   Schwartz AR, Potnis N, Timilsina S, Wilson M, Patane J, Martins J, Minsavage G V.,

623   Dahlbeck D, Akhunova A, Almeida N, Vallad GE, Barak JD, White FF, Miller SA,

624   Ritchie D, Goss E, Bart RS, Setubal JC, Jones JB, Staskawicz BJ. 2015. Phylogenomics of

625   Xanthomonas field strains infecting pepper and tomato reveals diversity in effector

626   repertoires and identifies determinants of host specificity. Front Microbiol 6:1–17.

627 46.   Newberry EA, Bhandari R, Minsavage G V., Timilsina S, Jibrin MO, Kemble J, Sikora

628   EJ, Jones JB, Potnis N. 2019. Independent Evolution with the Gene Flux Originating from

629   Multiple Xanthomonas Species Explains Genomic Heterogeneity in Xanthomonas

630   perforans. Appl Environ Microbiol 85:1–17.

631 47.   Beattie GA, Lindow SE. 1995. The secret life of foliar bacterial pathogens on leaves.

632   Annu Rev Phytopathol 33:145–172.

633 48.   Pfeilmeier S, Caly DL, Malone JG. 2016. Bacterial pathogenesis of plants: future

634   challenges from a microbial perspective. Mol Plant Pathol 7:1298–1313.

635 49.   Monier J, Lindow SE. 2003. Pseudomonas syringae Responds to the Environment on

636   Leaves by Cell Size Reduction 93:1209–1216.

637 50.   Leveau JHJ, Lindow SE. 2001. Appetite of an epiphyte: Quantitative monitoring of

638   bacterial sugar consumption in the phyllosphere. Proc Natl Acad Sci 98:3446–3453.

51.   Fatima U, Senthil-Kumar M. 2015. Plant and pathogen nutrient acquisition strategies.

Front Plant Sci 6:1–12.

52.   Ferelli AMC, Bolten S, Szczesny B, Micallef SA. 2020. Salmonella enterica Elicits and Is

Restricted by Nitric Oxide and Reactive Oxygen Species on Tomato. Front Microbiol

11:1–15.

53.   Meng F, Altier C, Martin GB. 2013. Salmonella colonization activates the plant immune

system and benefits from association with plant pathogenic bacteria. Environ Microbiol

15:2418–2430.

54.   Garcia AV, Charrier A, Schikora A, Bigeard J, Pateyron S, de Tauzia-moreau M-L,

Evrard A, Mithöfer A, Martin-magniette ML, Virlogeux-Payant I, Hirt H. 2014.

Salmonella enterica Flagellin Is Recognized via FLS2 and Activates PAMP-Triggered

Immunity in Arabidopsis thaliana. Mol Plant 7:657–674.

55.   Wahlig TA, Bixler BJ, Valdés-López O, Mysore KS, Wen J, Ané JM, Kaspar CW. 2019.

Salmonella enterica serovar Typhimurium ATCC 14028S is tolerant to plant defenses

triggered by the flagellin receptor FLS2. FEMS Microbiol Lett 366:1–7.

56.   Roy D, Panchal S, Rosa BA, Melotto M. 2013. Escherichia coli O157:H7 Induces

Stronger Plant Immunity than Salmonella enterica Typhimurium SL1344. Phytopathology

103:326–332.

57.   Tracy BS, Edwards KK, Eisenstark A. 2002. Carbon and nitrogen substrate utilization by

archival Salmonella typhimurium LT2 cells. BMC Evol Biol 2:1–6.

58.   Reid SJ, Abratt VR. 2005. Sucrose utilisation in bacteria: Genetic organisation and

regulation. Appl Microbiol Biotechnol 67:312–321.

59.   O'Leary BM, Neale HC, Geilfus CM, Jackson RW, Arnold DL, Preston GM. 2016. Early

changes in apoplast composition associated with defence and disease in interactions

between Phaseolus vulgaris and the halo blight pathogen Pseudomonas syringae Pv.

phaseolicola. Plant Cell Environ 39:2172–2184.

60.   Courbier S, Grevink S, Sluijs E, Bonhomme PO, Kajala K, Van Wees SCM, Pierik R.

2020. Far-red light promotes Botrytis cinerea disease development in tomato leaves via

jasmonate-dependent modulation of soluble sugars. Plant Cell Environ 43:2769–2781.

ACKNOWLEDGEMENTS

This material is based upon work supported by the National Science Foundation Graduate

Research Fellowship under Grant No. DGE-1747503 and by USDA-HATCH grant no.

WIS03022. Any opinions, findings, and conclusions or recommendations expressed in this

material are those of the author(s) and do not necessarily reflect the views of the National

Science Foundation. The funders had no role in study design, data collection, and interpretation,

or the decision to submit the work for publication.

We are grateful to J. Jones and G. Minsavage for providing the *Xanthomonas gardneri*

Δ*avrHah1* strain used in this work. Additionally, we thank C. Solís-Lemus and N. Keuler for

providing guidance on statistical analysis.

Author Contributions: MHD, KNC, and JDB developed the experiments. MHD, KNC, SCZ, and

INM performed the experiments. MHD and JDB wrote the manuscript, and all authors reviewed

the manuscript. MHD created the original illustrations.

TABLES

Table 1: Bacterial strains used in this study.

Strain Abbreviation	Description	Reference or Source
Xg WT	*Xanthomonas hortorum* pv. *gardneri* 444; Nal^R	(21)
Xg avr-	*Xanthomonas hortorum* pv. *gardneri* 444; *avrHah1^{ΔDBD}*	received from J. Jones; (21)
Xv	*Xanthomonas vesicatoria* 1111; Nal^R	ATCC
Se WT	*Salmonella enterica* sv. Typhimurium 14028s; Kan^R	(21)
Se ΔmotA	*Salmonella enterica* sv. Typhimurium 14028s; *ΔmotA*::Kan	(18)

Figure Legend

Figure 1: Infiltration and droplet inoculation assay. (A) Tomato leaflets were infiltrated with

water or bacteria through the abaxial surface; infiltration zones are indicated with a thick black

line and darker green shading. (B) Droplets of *S. enterica* (*Se*) cell suspension were then applied

to the adaxial surface of each infiltration zone at a site marked with permanent marker,

represented by black dash marks. The *Se* arrival day occurred 0-2 days post-infiltration (DPI).

(C) When the *Se* cells arrived on the plant via droplet application, several apoplast states were

possible depending on the infiltration treatment received. Circle symbols (healthy: open circle,

infected: bacteria-filled circle; dry: white circle, wet: turquoise circle) depict the various apoplast

states in subsequent data figures. (D) Designated leaflets were detached and irradiated with UV

at indicated timepoints, which ranged from 2.5 hours post-*Se* arrival (HPA) to 3 days post-Se

arrival (DPA) depending on the experiment. Tissue was sampled immediately after UV

irradiation at the site of *S. enterica* arrival (blue dotted circle) and non-overlapping distal tissue

(yellow dotted circles) for *Se* population enumeration.

Figure 2: Healthy leaf tissue with a water-filled apoplast facilitates localization to a protected

niche and migration to distal tissue. *S. enterica* (*Se*) was applied by droplet (10^6 CFU/leaflet) at 2

or 0 days post-infiltration (DPI) with water (blue boxplots). Half of leaflets were UV-irradiated

(triangle points; light-shaded boxplots) and the other half were left non-irradiated (circle points;

dark-shaded boxplots). *Se* populations were sampled at 1 day post *Se*-arrival (DPA) at the droplet

arrival site and distal from the arrival site. *Se* populations from droplet-inoculated leaflets were

compared to those from control leaflets infiltrated with *S. enterica* (gray boxplots). Uppercase

letters above the boxplots denote significance among treatment groups within the arrival site

based on an ANOVA and post-hoc Tukey test ($P < 0.05$). Lowercase letters denote significance

among treatment groups within the distal site based on a Kruskal-Wallis and post-hoc Wilcox

test ($P < 0.05$). Dotted line indicates limit of detection. Refer to Figure 1, panel C for circle

symbol key describing apoplast states.

Figure 3: *X. gardneri* infection facilitates *S. enterica* localization to protected niche and

migration beyond site of arrival. *S. enterica* (*Se*) was applied by droplet (10^6 CFU/leaflet) at 0

days post-infiltration (DPI) with water (blue boxplots) or 2 DPI with *X. gardneri* (*Xg* WT;

orange boxplots). Half of leaflets were UV-irradiated (triangle points; light-shaded boxplots) and

the other half were left non-irradiated (circle points; dark-shaded boxplots). *Se* populations were

sampled at 1 day post *Se*-arrival (DPA) at the droplet arrival site and distal from the arrival site

(distal site for *Xg*). Uppercase letters above the boxplots denote significance among treatment

groups within the arrival site based on an ANOVA and post-hoc Tukey test ($P < 0.05$).

Lowercase letters denote significance among treatment groups within the distal site based on a

Kruskal-Wallis and post-hoc Wilcox test ($P < 0.05$). Dotted line indicates limit of detection.

Refer to Figure 1, panel C for circle symbol key describing apoplast states.

Figure 4: *S. enterica* access to protected niche within 2.5 hours of arrival on water-soaked tissue

depends on *Xanthomonas* strain and duration of infection. *S. enterica* (*Se*) was applied by droplet

(10^6 CFU/leaflet) to leaflets infiltrated with water (blue boxplots), *X. gardneri* WT (*Xg* WT;

orange boxplots), *X. gardneri avrHah1$^{\Delta DBD}$* (*Xg avr;* pink boxplots), or *X. vesicatoria* (*Xv*; green

boxplots) at indicated days post-infiltration (DPI). (A) Unabsorbed *S. enterica* (*Se*) droplet

volumes measured 2.5 hours following droplet application. Differing letters above the plot

denote significantly different treatment groups based on a Kruskal-Wallis and post-hoc Wilcox

multiple comparison test ($P < 0.05$). (B) Reduction of *S. enterica* populations in response to UV

irradiation at the *Se* droplet arrival site at 2.5 hours-post *Se* arrival (HPA). Letters above the

boxplots denote significance among treatment groups based on an ANOVA and post-hoc Tukey

test ($P < 0.05$). Refer to Figure 1, panel C for circle symbol key describing apoplast states.

Figure 5: *X. gardneri avrHah1* enhances *S. enterica* migration during early infection, and

flagellar motility promotes *S. enterica* survival at distal tissue. *S. enterica* (*Se*) WT (circle points)

or Δ*motA* (asterisk points) was applied by droplet (10^6 CFU/leaflet) at 1 or 2 days post-

infiltration (DPI) with *X. gardneri* WT (*Xg* WT, orange boxplots) or *X. gardneri avrHah1$^{\Delta DBD}$*

(*Xg avr*; pink boxplots). *Se* populations were sampled at 1 day post *Se*-arrival (DPA) distal from

the *Se* droplet arrival site without UV irradiation treatment. Letters above the boxplots denote

significance among treatment groups based on a Kruskal-Wallis and post-hoc Wilcox test ($P <$

0.05). Dotted line indicates limit of detection. Refer to Figure 1, panel C for circle symbol key

describing apoplast states.

Figure 6: *Xanthomonas*-infected apoplast serves as a hospitable niche for *S. enterica* 3 days post-

arrival. *S. enterica* (*Se*) was applied by droplet (10^6 CFU/leaflet) to leaflets infiltrated with water

(blue boxplots), *X. gardneri* WT (*Xg* WT; orange boxplots), *X. gardneri avrHah1$^{\Delta DBD}$* (*Xg avr;*

pink boxplots), or *X. vesicatoria* (*Xv*; green boxplots) at indicated days post-infiltration (DPI). *Se*

populations were sampled at 3 days post *Se*-arrival (DPA) from the arrival site and distal from

the arrival site. All leaflets were UV irradiated prior to sampling. Uppercase letters above the

boxplots denote significance among treatment groups within the arrival site based on an

ANOVA and post-hoc Tukey test ($P < 0.05$). Lowercase letters denote significance among

treatment groups within the distal site based on a Kruskal-Wallis and post-hoc Wilcox test ($P <$

0.05). Dotted line indicates limit of detection. Refer to Figure 1, panel C for circle symbol key

describing apoplast states.

Figure 7: *X. gardneri*-infected apoplast enables replication of *S. enterica* within protected niches.

S. enterica (*Se*) was applied by droplet (10^3 CFU/leaflet) to leaflets infiltrated with water (blue

boxplots), *X. gardneri* WT (*Xg* WT; orange boxplots), *X. gardneri avrHah1^{\Delta DBD}* (*Xg avr;* pink

boxplots), or *X. vesicatoria* (*Xv*; green boxplots) at indicated days post-infiltration (DPI). *Se*

populations were destructively sampled at 2.5 hours post *Se*-arrival (HPA) and 3 days post *Se*-

arrival (DPA) at the droplet arrival site and distal from the arrival site. All leaflets were UV

irradiated prior to sampling. Letters above the boxplots denote significance between the

healthy/wet (H2O 0 DPI) and Xg treatment groups within the arrival (uppercase) and distal

(lowercase) sites based on an ANOVA and post-hoc Tukey test ($P < 0.05$). Dotted line indicates

limit of detection (LOD). Zero values were confirmed via LB enrichment. Points plotted at

LOD/2 indicate absence of colonies via direct plating and a positive enrichment sample. Refer to

Figure 1, panel C for circle symbol key describing apoplast states.

Figure 8: Model for *S. enterica* survival in a habitable apoplast. *S. enterica* can colonize crop

plants via spray-irrigation of contaminated water. Macroscopically dry tissue (a) does not enable

absorption of surface water, whereas tissue with an aqueous apoplast (b) facilitates the ingression

of *S. enterica* into the apoplast. Healthy tissue that is abiotically water-soaked eventually dries

out (c), creating an unfavorable niche for *S. enterica.* By contrast, an infected apoplast (d), which

maintains water-soaking via phytopathogen effectors and is likely enriched with nutrients and

other growth-promoting factors, facilitates *S. enterica* replication.

FIGURE 1

(A) Infiltration
H2O or bacteria

(B) *Se* Arrival Day
0-2 DPI

(D) Sampling & UV Irradiation
2.5 HPA-3 DPA

No UV   + UV

(C) Possible Apoplast States at *Se* Arrival

healthy dry   healthy wet   infeted dry   infected wet

FIGURE 2

FIGURE 3

FIGURE 4

FIGURE 5

FIGURE 6

FIGURE 7

FIGURE 8