Transcriptional dynamics of colorectal cancer risk associated variation at 11q23.1 are correlated with tuft cell abundance and marker expression *in silico*

Bradley T. Harris[1], Vidya Rajasekaran[1], James P. Blackmur[1,2], Alan O'Callaghan[2], Kevin Donnelly[1,2], Maria Timofeeva[1,3], Peter G. Vaughan-Shaw[1,2], Farhat V. N. Din[1], Malcolm G. Dunlop[1,2] and Susan M. Farrington[1,*]

[1] Cancer Research UK Edinburgh Centre, Institute of Genetics and Cancer, University of Edinburgh, United Kingdom

[2] MRC Human Genetics Unit, Institute of Genetics and Cancer, University of Edinburgh, Edinburgh, United Kingdom

[3] Department of Public Health, D-IAS, Danish Institute for Advanced Study, University of Southern Denmark, Odense, Denmark

* Corresponding author: susan.farrington@ed.ac.uk

Keywords: colorectal cancer; risk; eQTL (expression quantitative trait loci); scRNAseq (single cell RNA sequencing); WGCNA (weighted gene co-expression network analysis); linear modelling

Abstract

Colorectal cancer (CRC) is characterised by heritable risk that is not well understood. Heritable, genetic variation at 11q23.1 is associated with increased colorectal cancer (CRC) risk, demonstrating eQTL effects on 3 cis- and 23 trans-eQTL targets. We sought to determine the relationship between 11q23.1 cis- and trans-eQTL target expression and test for potential cell-specificity. scRNAseq from 32,361 healthy colonic epithelial cells was aggregated and subject to weighted gene co-expression network analysis (WGCNA). One module (blue) included 19 trans-eQTL targets and was correlated with *C11orf53* expression only. Following unsupervised clustering of single cells, the expression of 19 trans-eQTL targets was greatest and most variable in cluster number 11, which transcriptionally resembled tuft cells. 14 trans-eQTL targets were found to demarcate this cluster, 11 of which were corroborated in a second dataset. Intra-cluster WGCNA and module preservation analysis then identified twelve 11q23.1 trans-eQTL targets to comprise a network that was specific to cluster 11. Finally, linear modelling and differential abundance testing showed 11q23.1 trans-eQTL target

expression was predictive of cluster 11 abundance. Our findings suggest 11q23.1 trans-eQTL

targets comprise a *C11orf53*-related network that is likely tuft cell-specific and reduced

expression of these genes correlates with reduced tuft cell abundance *in silico.*

Introduction

Colorectal cancer (CRC) is the fourth most common cancer type UK and worldwide (Cancer

Research UK, 2017; Rawla et al., 2019). Approximately 40% of CRC risk is attributable to

heritable genetic variation (Graff et al., 2017), with rare, highly penetrant mutations

responsible for a small fraction of overall risk (Jasperson et al., 2010; Sud et al., 2017).

Genome wide association studies (GWAS) have identified 129 common genetic variants

associated with CRC risk (Huyghe et al., 2019; Law et al., 2019). Several common CRC

genetic risk variants are associated with heritable changes in the expression levels of genes

in colonic mucosa, known as expression quantitative trait loci (eQTLs) (Closa et al., 2014;

Hulur et al., 2015; Loo et al., 2017; Vaughan-Shaw et al., 2021).

Genetic variation at 11q23.1 is associated with increased CRC risk (Tenesa, Farrington et al.,

2008). However, the large number of variants in high linkage disequilibrium at 11q23.1 makes

identifying the causal variant, a key step to identifying the mechanism of gene dysregulation,

difficult. Studies have shown that CRC risk associated variation at several 11q23.1 variants is

correlated with the downregulation of three local genes; *C11orf53, COLCA1, COLCA2*

(Biancolella et al., 2014; Loo et al., 2017; Smillie, 2015), known as cis-eQTL targets. We have

recently shown that variation at rs3087967, a single nucleotide variant in the 3'UTR of

C11orf53, is significantly correlated with distal CRC risk and a myriad of distant, trans-eQTL

targets throughout the colon (Vaughan-Shaw *et al.,* 2021), Additional File 1. Of these, only

two have a common, well-described function; *IL17RB* and *TRPM5* are experimentally

determined markers of tuft cells - a rare epithelial cell-type (Haber et al., 2017; Kaske et al.,

2007). The function of several other trans-eQTL targets is currently unknown and their exact

relevance to both 11q23.1 cis-eQTL target expression and CRC risk is not characterised.

Current methods of eQTL detection, while widely used, exhibit some key limitations. eQTLs

are frequently identified by linear models of bulk RNA-seq/microarray transcriptome analysis

of healthy tissue (Michaelson et al., 2009; Richards et al., 2012; Stranger et al., 2007). These

methods often treat both gene expression and single-nucleotide polymorphisms as

independent, linear entities: assumptions which over-simplify the complex relationships

governing gene expression dynamics and limit results to additive gene-dosage related findings.

In addition, eQTL analyses require performance of an extremely large number of independent

tests to be conducted, limiting its sensitivity. Correlation-based methods of gene expression

analysis, such as weighted gene co-expression network analysis (WGCNA, Langfelder &

Horvath, 2008), circumvent this problem by agnostically identifying correlations between

individual genes, and entire, non-overlapping gene modules, with binarized categorical or

quantitative traits. Modules of correlated genes may themselves be correlated with sample

phenotypes. In addition, WGCNA also does not require a hard thresholding of correlations, a

major advantage over other correlation-based methods that rely on arbitrary cut offs. We have

recently shown that WGCNA can be efficacious at identifying genes driving transcriptional

dynamic changes in the colorectum of patients treated with Vitamin D (Blackmur et al., 2022),

even in the absence of statistically significant changes in differential expression analysis.

Moreover, as CRC risk-associated eQTL targets are identified by bulk expression methods,

findings are inherently limited in their potential to detect cell-specific changes, particularly in

rare cell-types; expression changes within which may be masked due to low relative

abundance.

We hypothesised that the expression of 11q23.1 trans-eQTL targets may be correlated with

only a single cis-eQTL target, a relationship which may in turn be specific to a single epithelial

cell-type in the colon. We utilised WGCNA across and within single-cell RNA sequencing

(scRNAseq) clusters to further characterise eQTL target expression relatedness in colonic

epithelial cell-types. An overview of the analysis performed in this study is shown in Figure 1.

Results

Understanding cis-eQTL specificity of 11q23.1 trans-eQTL effects

To test potential colonic epithelial cell-specificity of 11q23.1 eQTL effects, we sought to

analyse the expression of target genes within colonic epithelial cell-types. To this end, we

obtained scRNAseq from 32,361 healthy human colonic mucosa epithelial cells from eleven

individuals (Smillie et al., 2019). To first assess the validity of this dataset in studying 11q23.1

variation-related expression dynamics, we devised a method of pseudo-bulking the

expression from all cells in this dataset, aggregating the expression of every gene across all

cells from each sample (see Figure 1 and Methods). Pseudo-bulked expression from 11q23.1

nominally significant trans-eQTL targets, present in the scRNAseq dataset (p<0.01, n=273),

Figure 1. Overview of study design and analysis. This study makes use of two single cell RNA sequencing (scRNAseq) datasets: Smillie *et al.,* (2019) – n=32,261 and Elmentaite *et al.,* (2021) - n=11,651. The analysis performed on each dataset is outlined by arrow colour: blue=all cells from Smillie *et al.,* (2019), yellow=individual clusters from Smillie *et al.,* (2019), green=individual clusters from Elmentaite *et al.,* (2021). WGCNA=Weighted Gene Co-expression Network Analysis, GSEA=Gene Set Enrichment Analysis.

Figure 2A, was then subject to WGCNA (Langfelder & Horvath, 2008). Genes were
agnostically grouped into modules of correlated expression, and the correlation between the
eigenvector (first principal component) of each module with sample traits and pseudo-bulked
cis-eQTL target gene expression was calculated, Figure 2B. The blue module, comprising 77
genes, was found to correlate with *C11orf53* expression (cor=0.81, FDR=2e-04), but none of
the sample traits, *COLCA1* or *COLCA2* expression. The blue gene module includes 17 of the
20 significant 11q23.1 trans-eQTL targets that passed gene filtration quality control
(FDR<0.05, Vaughan-Shaw *et al.,* 2021); *ALOX5, SH2D6, TRPM5, BMX, PSTPIP2, GNG13,*
IL17RB, HTR3E, PTGS1, SH2D7, OGDHL, MATK, PLCG2, LRMP, PIK3CG, HTR3C and
CAMP, thus indicating the genes comprising this module are specifically related to the
expression of *C11orf53,* Supplementary Table 2.
116
117 To assess the contribution of 11q23.1 genes to the correlation of the blue module with
118 *C11orf53,* we performed gene set enrichment analysis of the 11q23.1 trans-eQTL target
119 genes (FDR<0.05) against the genes in the blue module, ranked by their module membership
120 - a measure of their relatedness to all other genes in the module (Langfelder & Horvath, 2008).
121 We found the genes in this module are highly enriched for 11q23.1 trans-eQTL targets,

Figure 2. 11q23.1 trans-eQTL targets are correlated with the expression of *C11orf53*, but not *COLCA1* or *COLCA2* in pseudo-bulked scRNASeq. (A) Hierachical clustering of complete distances between pairwise correlations of 11q23.1 nominal trans-eQTL targets pseudo-bulked expression (p<0.01, n=273) in 32,361 healthy colonic epithelial scRNAseq. (B) Weighted Gene Co-expression Network Analysis (WGCNA) identified module trait relationships. Pearson correlations shown above Benjamini-Hochberg corrected p-values in brackets. (C) Gene Set Enrichment Analysis (GSEA) of 11q23.1 trans-eQTL targets (FDR<0.05) in blue module genes, ranked by their module membership. (D) Kamadakawai network of Blue module relatedness (adjacency > 0.3). Red nodes indicate FDR<0.05 trans-eQTL target of 11q23.1.

normalised enrichment score (NES) = 2.04, p=7.78e-04, Figure 2C. In addition, 11 trans-eQTL

targets comprised the 12 genes with the greatest adjacency in the blue module (adjacency >

0.3), Figure 2D. We also find that module membership is highly correlated with gene

significance for *C11orf53* in the blue module, reinforcing the overall correlation of the network

with *C11orf53* expression (Supplementary Figure 1). Together, this replicates the trans-eQTL

target expression relatedness previously described (Vaughan-Shaw et al., 2021) and indicates

that the expression of the majority of significant 11q23.1 trans-eQTL target genes is correlated

with *C11orf53* specifically in this dataset.

Figure 3. 11q23.1 trans-eQTL expression demarcates tuft-like cell cluster. (A) UMAP of 32,361 epithelial scRNASeq, coloured by cell cluster - identified using Seurat (Hao et al., 2021). (B) Upper: GSEA of 11q23.1 trans-eQTL targets (Vaughan-Shaw *et al.,* 2021; FDR<0.05) in cluster 11 markers. Lower: GSEA of putative colonic tuft cell signature (Smillie *et al.,* 2019) in cluster 11 markers. P=P value, FDR=False Discovery Rate (C) Relative, pseudo-bulked expression (log transcripts per 10,000) of 11q23.1 trans-eQTL targets (FDR<0.05) across clusters.

Analysing cell-specificity of 11q23.1 eQTL target expression

To test the potential for cell-specific expression of 11q23.1 eQTL targets in this dataset, we

performed dimensionality reduction and clustering of the single cells, identifying a total of 12

transcriptionally distinct cell-clusters, named '0'-'11', Figure 3A. We then calculated the

markers of each cluster and found the markers of cluster 11, comprising 318 cells, includes

fourteen 11q23.1 trans-eQTL targets (FDR<0.05), Table 1 (full list of cluster 11 marker genes

available in Additional File 2). In addition, cluster 11 markers are significantly enriched for

11q23.1 trans-eQTL targets (NES=2.15, p=7.52e-06), Figure 3B. Cluster 11 was found to

transcriptionally resemble the Smillie *et al.,* tuft cell cluster, by enrichment of each cluster's

markers with the authors' putative markers (NES=2.37, FDR=1.23e-06, see Methods).

Together, this indicates cluster 11 is transcriptionally defined by the expression of 11q23.1

trans-eQTL targets, several of which are themselves putative tuft cell markers.

Because the expression of 11q23.1 trans-eQTL targets is reduced in individuals with the CRC

risk-associated genotype at 11q23.1 (Vaughan-Shaw et al., 2021), we wanted to assess the

variability in the expression of these genes within each cluster. We pseudo-bulked the

expression of all cells from each sample, within clusters, and analysed 11q23.1 trans-eQTL

target expression (Figure 3C). The relative expression level and variability of *C11orf53* and 18

of the 11q23.1 trans-eQTL targets is overwhelmingly greatest within cluster 11, indicating the

eQTL effect on these genes is exacerbated within this cluster. Notably, the relative variation

and expression of cis-eQTL targets *COLCA1* and *COLCA2* and trans-eQTL targets *ANKHD1*

and *GIN1,* is not greatest within this cluster, suggesting the eQTL effect on these genes may

not be driven by transcriptional dynamics within this cell-type. In addition, we analysed

11q23.1 eQTL target variability at the single cell level, Supplementary Figure 2. The variability

of *C11orf53* and the same 18 trans-eQTL targets was found to be greatest within cluster 11,

replicating the potential exacerbation of the eQTL effect within this cluster and supporting the

validity of our pseudo-bulk method. *C11orf53* expression, however, was not found to

demarcate cluster 11 by marker identification analysis (see Methods), data not shown.

To test the robustness of tuft cell-like mapping of 11q23.1 trans-eQTL target expression and

variability, we replicated this analysis in an independent dataset of 11,651 healthy adult

colonic epithelial cells from 3 individuals (Elmentaite et al., 2021), Supplementary Figure 4A.

In this instance, we identified 19 cell-clusters by dimensionality reduction and unsupervised

clustering, named '0'-'18'. The markers of cluster 18 were significantly enriched for the

Gene	p_val	avg_log2FC	pct.1	pct.2	p_val_adj
AZGP1	1.00E-279	2.5735445	0.72	0.029	1.00E-279
SH2D6	1.00E-279	2.43533298	0.736	0.007	1.00E-279
LRMP	1.00E-279	1.97284472	0.572	0.005	1.00E-279
PTGS1	1.18E-278	1.33766533	0.459	0.003	1.75E-274
IL17RB	1.00E-264	1.04046297	0.412	0.031	1.48E-260
BMX	7.18E-248	1.18575976	0.406	0.002	1.07E-243
ALOX5	8.25E-244	1.09872574	0.409	0.003	1.22E-239
MATK	2.88E-240	1.3710223	0.406	0.003	4.28E-236
SH2D7	2.50E-235	1.20590753	0.393	0.002	3.72E-231
GNG13	2.93E-220	0.98868335	0.362	0.001	4.34E-216
TRPM5	1.32E-214	0.94972732	0.377	0.004	1.96E-210
PLCG2	1.22E-209	0.98215989	0.365	0.002	1.82E-205
PSTPIP2	2.21E-197	0.87613246	0.333	0.004	3.28E-193
HTR3E	4.86E-184	0.8684383	0.308	0.001	7.21E-180

Table 1. 11q23.1 trans-eQTL targets identified as cluster 11 markers. Markers calculated using MAST (McDavid *et al.,* 2020). Avg_log2FC=average log2 fold change between cluster 11 and all other clusters. Pct1=Proportion of gene expression in cluster 11. Pct2=Proportion of expression in non-cluster 11. P_val_adj=FDR-corrected p-value.

expression of 11q23.1 trans-eQTL targets (NES=2.50, P=5.52e-09, Supplementary Figure

4B), 11 of which were identified as markers of this cluster, Supplementary Table 3. Cluster 18

was also enriched for the tuft cell signature (NES=2.41, P=7e-08), replicating our previous

finding. The relative variability and expression of the majority of 11q23.1 trans-eQTL targets

was also greatest within cluster 18, Supplementary Figure 4C. The expression of 13 trans-

eQTL targets was also most variable within cluster 18 when expression was analysed at the

single cell level, Supplementary Figure 4. Interestingly, *C11orf53* expression variability was

not found to vary greatest within cluster 18 when using expression from single cells.

Understanding 11q23.1 cis- and trans-eQTL relatedness within clusters

The demarcation and variability of 11q23.1 trans-eQTL target expression within a tuft-like cell

cluster strongly indicates the eQTL effect is derived from altered gene expression in this cell-

type specifically. However, it is possible that the gene-gene relatedness of 11q23.1 eQTL

targets, including those with *C11orf53*, is not specific, but rather exacerbated in this cluster.

To test this, we sought to divide samples by genotype at rs3087967, the variant associated

with expression changes of trans-eQTL targets (Vaughan-Shaw *et al.,* 2021)*,* and analyse the

congruence of trans-eQTL target expression relatedness within clusters. While genotype

information was unavailable for the Smillie *et al.,* (2019) dataset, raw sequencing reads were

available for the Elmentaite *et al.,* (2021) dataset and as rs3087967 lies within the 3' UTR of

C11orf53, we performed variant calling on these samples using orthogonal tools,

Supplementary Table 4. Using freebayes (Garrison & Marth, 2012), we found that all samples

were called as homozygous for the non-risk allele at rs3087967, except for one sample called

as heterozygous. However, as all 3 other samples from this individual were called as

homozygous non-risk, it is likely this is a technical error. Using bcftools (Danecek et al., 2021),

all samples were identified to be homozygous non-risk for rs3087967. The lack of genetic

variation at rs3087967 in these samples is consistent with the absence of heightened

C11orf53 variability in cluster 18 and indicates this dataset would be unlikely to be of use to

identify *C11orf53*-related expression dynamics. The relatively high variability of trans-eQTL

target expression within cluster 18 may be indicative of non-11q23.1-related dynamics, such

as changes during differentiation or cell cycle progression.

To test the potential efficacy of the Smillie *et al.,* (2019) dataset to further study the 11q23.1

eQTL target expression dynamics, we compared the standardized variability of 11q23.1 trans-

eQTL targets within the respective demarcated clusters at the single-cell level, Supplementary

Table 5. We found that the expression variability of 14 of the 15 11q23.1 trans-eQTL targets,

expressed in both Elmentaite *et al.,* (2021) cluster 18 and Smillie *et al.,* (2019) cluster 11, was

substantially increased in the latter (fold-change range 1.46-9.7, median=1.97, mean=2.83,

95% confidence interval=1.50-4.18). The only 11q23.1 trans-eQTL target not to exhibit

increased variability in Smillie *et al.,* (2019) cluster 11 was OGDHL (fold change=0.73).

Notably, the expression of *C11orf53,* was also greater in Smillie *et al.,* cluster 11 (fold

change=1.94). Subsequent analysis was therefore focussed on the Smillie *et al.,* (2019)

dataset.

To divide samples into high and low expressors of *C11orf53*-expression related genes, we

hierarchically clustered samples based on the relative expression of pseudo-bulk WGCNA-

identified blue module hub genes. Hub genes were defined by module membership > 0.7,

intramodular connectivity > 0.7 and network adjacency > 0.3 and include 10 genes: *TRPM5,*

PSTPIP2, SH2D6, ALOX5, BMX, GNG13, SH2D7, HCK, PLCG2, MATK, Figure 4A. Five

samples were separated at the first branch of clustering and exhibit a strong relative reduction

in the expression of these genes. This grouping of samples is henceforth referred to as the

'blue module hub gene grouping'. To assess the significance of this separation at representing

Figure 4. Several 11q23.1 trans-eQTL targets comprise *C11orf53*-correlated network in tuft-like cluster only. (A) Relative (z-score) expression of blue module hub genes from Fig.1D (MM>0.7, kIM>0.7, adj > 0.3). Robustness assessed by 10,000 permutations: p=0.055. (B) Module trait matrix of gene modules identified by WGCNA within pseudo-bulked expression from cluster 11. Only modules exhibiting a correlation with a covariate (FDR<0.1) are shown. Total number of modules=20. (C) Correlation of Gene Significance (GS) and Module Membership (MM) of genes in the black module from (B). 11q23.1 trans-eQTL targets are highlighted. 11q23.1 trans-eQTL targets with GS.C11orf53 > 0.5 and MM.black > 0.5 (red) were used as the auxiliary module. (D) Preservation of correlations between auxiliary module genes with the module eigengene (ME) and equivalent values in cluster 11. Dashed line indicates nominal significance threshold (p=0.05). (E) Average MM of genes within the auxiliary module across clusters. (F) Pairwise pearson correlations (p<0.05) between the pseudo-bulked expression of auxiliary module genes across clusters. Cluster number in upper right of each facet. Cluster 4 and 8 exhibited no significant correlations and so are not plotted.

underlying transcriptional differences, we performed 10,000 permutations, using 10 randomly

sampled genes, and generated a p-value of 0.055.

To first test gene-gene relatedness within cluster 11, we performed WGCNA on the relative,

pseudo-bulked expression of the top 5000 most variable genes within this cluster, identifying

a total of 20 modules, 7 of which exhibited correlations with sample covariates that approached

significance (FDR<0.1), Figure 4B. We found one module, 'cluster 11 black', which was highly

correlated with both the 'blue hub gene grouping' (cor=0.72, FDR=0.032) and *C11orf53*

expression (cor=0.68, FDR=0.048). 'Cluster 11 black' is comprised of 290 genes, including

fifteen 11q23.1 trans-eQTL targets, Additional File 3. As the 'blue module hub gene' grouping

is derived from analysis of genes correlated with *C11orf53* across all cells, the correlation of

'cluster 11 black' with both this grouping and *C11orf53* expression indicates that this

relationship is preserved within and potentially derived from this cell-cluster.

We then wanted to test whether the gene-gene relatedness of 11q23.1 trans-eQTL targets,

correlated with *C11orf53*, was specific to cluster 11. To this end, we defined an auxiliary

module, comprised of the twelve 11q23.1 trans-eQTL targets correlated with *C11orf53*

(cor>0.5, p<0.05) in cluster 11 black, which exhibited high module membership (MM.black >

0.5), Figure 4C. These genes include: *HTR3E, LRMP, GNG13, ALOX5, SH2D7, PTGS1,*

MATK, BMX, AZGP1, IL17RB, SH2D6 and *OGDHL.* We performed pairwise module

preservation of this auxiliary module, using identical parameters as used for intra-cluster 11

analysis, in all other clusters, see Methods. For two modules, cluster 8 and cluster 10, the

5000 most variable genes included only a single member of the auxiliary module and so were

excluded from this analysis. To analyse the preservation of the connectivity of these genes,

we assessed the similarity of each gene correlation with the module eigengene, and the

equivalent values in cluster 11 (cor.kME), Figure 4D. No module exhibited a significant

cor.kME with cluster 11 (p>0.05), indicating there is low overall preservation of the connectivity

between these genes in all other clusters. The average module membership of genes in this

module (average.MM) was also reduced in all clusters compared with cluster 11, Figure 4E.

Finally, we analysed the pairwise gene-gene correlations between all members of the auxiliary

module and *C11orf53* within each cluster, Figure 4F. While there are rare correlations between

these genes in other modules (cor>0.5, p<0.05), all comparisons reach this threshold in cluster

11. Taken together, this evidence indicates these 12 trans-eQTL targets comprise a

transcriptional network that is correlated with *C11orf53* expression and likely specific to cluster

11.

Identification of cluster 11 abundance associated genes

As many of the 11q23.1 eQTL targets, including those comprising the cluster 11-specific

network, demarcate this cluster, we wanted to examine the relationship between their

expression and cluster 11 abundance. First, we performed linear modelling of the relative

abundance of cluster 11 and the pseudo-bulked expression of *C11orf53* only. We found that

the expression of *C11orf53* is associated with the relative abundance of cluster 11

(coefficient=0.389, p=0.00431) indicating potential for the expression of *C11orf53* alone to be

moderately predictive of the abundance of this cell-type, Figure 5A.

To agnostically test the predictive power of 11q23.1 trans-eQTL target expression on cluster

11 abundance, we tested the association between the expression of all genes and the

proportion of cluster 11 in samples, Figure 5B. We found that all genes significantly associated

with the abundance of this cluster (FDR<0.05, log-fold change>1) were indeed 11q23.1 trans-

eQTL targets, including: *ALOX5, BMX, GNG13, MATK, SH2D7, PSTPIP2, TRPM5* and

PTGS1. In fact, the strength of gene association with cluster 11 abundance was also

significantly enriched for 11q23.1 trans-eQTL targets (NES=2.15, p=8.03e-10, Figure 5C).

While our linear modelling strongly supports a predictive role of 11q23.1 eQTL target

expression in the abundance of cluster 11, we wanted to agnostically test whether the

expression of the *C11orf53*-related trans-eQTL targets was correlated with abundance

changes to any cluster. To this end, we utilised miloR (Dann et al., 2021) to calculate cell-

neighbourhoods, Supplementary Figure 5, which were then used to perform differential

abundance testing across the 'blue module hub gene grouping', defined in Figure 4A. To

Figure 5. 11q23.1 trans-eQTL target expression is associated with the abundance of tuft cell-like cluster. (A) Linear modelling of pseudo-bulked *C11ORF53* expression and cluster 11 abundance. (B) Volcano plot of linear model result on pseudo-bulked expression of all 14,843 genes. Significantly associated 11q23.1 trans-eQTLs (logFC>1, FDR<0.05) highlighted. (C) GSEA of 11q23.1 trans-eQTLs (FDR<0.05) in (B), ranked by logFC. (D) Differential abundance of neighbourhoods in the 'Low' blue hub gene group compared to 'High'. Neighbourhoods were identified using miloR (Dann *et al.,* 2021) and grouped by cluster. Only significant neighbourhoods with major cluster proportion > 0.8 are plotted and only significant neighbourhoods (Spatial FDR < 0.01) are coloured by their logFC (red=down, blue=up). No neighbourhoods from cluster 10 were comprised of major proportion > 0.8.

generalise neighbourhoods to the cell clusters we have identified, we subsequently filtered for

neighbourhoods which represented a majority of a single cluster (majority proportion > 0.8)

Figure 5D. No neighbourhood from cluster 10 was comprised of a majority proportion of 0.8

and so this cluster was not included. In the low *C11orf53*-related gene expressing samples

there are significant reductions (Spatial FDR < 0.01) from cluster 0 and increases in

neighbourhoods in clusters 1, Figure 5D. The change in abundance of these neighbourhoods

represent a small proportion of the total number of neighbourhoods detected in these clusters

(2.1% and 1.3% respectively) and are therefore unlikely to represent a dramatic phenotype.

In contrast, all 7 of the neighbourhoods comprising a majority of cluster 11 cells were
significantly underrepresented in the low 'blue module hub gene group'. These results indicate
11q23.1 eQTL target expression is correlated with a significant and likely specific change in
abundance of cluster 11 cells.

Discussion

In this study, our pan-cluster WGCNA serves as validation of the expression relatedness
between trans-eQTL targets previously identified to be correlated with CRC associated
variation at 11q23.1 (Vaughan-Shaw *et al.,* 2021). 11q23.1 trans-eQTL target expression was
also found to be more correlated with *C11orf53* over other cis-eQTL targets. Following
clustering of single cells, many of the genes demarcating a single cluster, number 11, were
found to be 11q23.1 trans-eQTLs targets. Enrichment of these markers against putative gene
sets showed cluster 11 transcriptionally resembled tuft cells, replicated in an independent
dataset. WGCNA within cluster 11 identified several 11q23.1 trans-eQTL targets that exhibited
a high level of relatedness, subsequent analysis of the preservation of this relatedness
indicated this was likely to be specific to this cell cluster. Finally, samples which exhibited low
overall expression of 11q23.1 trans-eQTL targets most correlated with one another, were
found to exhibit a specific and dramatic reduction of the tuft cell-like cluster. Therefore, our
results potentiate CRC risk associated variation at 11q23.1 reduces the abundance of colonic
tuft cells.

To our knowledge, this is the first study to map CRC risk associated eQTL targets to specific
epithelial cell-types. Heritable inflammatory bowel disease risk loci have recently been
associated with shifts in transcriptional dynamics in individual colonic epithelial cells (Smillie
et al., 2019) and it is possible that other CRC risk variants, with robust eQTL effects, are
associated with cell-specific changes in transcriptional dynamics. Delineating the cell-specific
expression of CRC risk-associated eQTLs may provide valuable insights into the mechanism
of risk-associated pathophysiology and should be a focus of future work. The growing size
and availability of scRNAseq datasets will likely make cell-specificity mapping of heritable
disease associated eQTLs significantly easier and available, particularly if genotype data is
available. Indeed, our study is also not the first use of WGCNA to detect gene-gene
relatedness in scRNAseq data. WGCNA has been utilised to identify gene modules associated
with the activation of neuronal stem cells (Luo et al., 2015) and human induced pluripotent

stem cells (Wu et al., 2021), however, like our own study, these studies did not utilise

expression from individual cells as the input for WGCNA.

Surprisingly, the expression of the vast majority of 11q23.1 eQTL targets mapped to the cell-

type that transcriptionally resembles tuft cells, including; *LRMP, IL17RB, SH2D6, PLCG2,*

PSTPIP2, TRPM5, SH2D7, AXGP1, PTGS1, ALOX5, BMX. Many of the most significant

11q23.1 trans-eQTL targets, such as *LRMP, SH2D7* and *ALOX5*, have not previously been

associated with specific expression in this cell-type, potentiating their status as a marker in

the colonic epithelium. Additional markers of the tuft cell-like cluster include *HCK* and *HPGDS,*

for which there is some orthogonal evidence of expression within tuft cells (Gerbe et al., 2011;

Yamaga et al., 2018). This improves our confidence that cluster 11 represents this cell-type

and is not an artefact of the analysis.

It is noteworthy that many of the genes with expression found to be in association with

C11orf53 in the scRNAseq data were indeed identified as trans-eQTL targets in the bulk

analysis (Vaughan-Shaw et al., 2021). While the mapping of the expression of these genes to

a tuft-like cell cluster could only be achieved by using expression from individual cells, the

prior identification of these genes by bulk analysis is a testament to the power of such

approaches, and concordance of their findings with single cell-based methods.

Finally, the overall potentiation of tuft cell perturbation is of great interest regarding

characterisation of the mechanism governing CRC risk at 11q23.1. Tuft cells are associated

with stem-, neuro-transmitting and immune-related functions (Banerjee et al., 2020; Gerbe et

al., 2016; Middelhoff et al., 2017; Yi et al., 2019), however much of the evidence regarding

their function is derived from other organs and cannot necessarily be extrapolated to the colon.

Interestingly, genetic ablation of tuft cell abundance has been associated with exacerbated

tumour progression in mouse models of pancreatic cancer (DelGiorno et al., 2020; Hoffman

et al., 2021). Both studies show this is likely in association with perturbed immune cell function

and signalling. In line with this, tuft cell abundance has recently been shown to be reduced in

quiescent ulcerative colitis patients (Kjærgaard et al., 2021), suggesting tuft cells are involved

in immune regulation in the colon. Future work should aim to experimentally validate the

relationship between 11q23.1 variation and tuft cell abundance, examine how this impacts

tumourigenesis and identify the potential mechanism of CRC risk predisposition.

Materials and methods

Availability of data and materials

Raw gene-level expression counts of healthy colonic epithelial cells, published by Smillie *et al.,* (2019), were obtained from Single Cell Portal using accession code SCP259. Samples from one patient, N51, were removed as they were found to be outliers on the basis of cell-level mitochondrial and ribosomal protein gene expression, in addition to principal component analysis of pseudo-bulked expression. Fastq files for the second dataset, published by Elmentaite *et al.,* (2021) were obtained from https://www.ebi.ac.uk/arrayexpress, using accession code E-MTAB-9543. Fastq files were aligned to the hg19 transcriptome using the 10X Genomics Cell Ranger v3.02 pipeline (Zheng et al., 2017) to produce the raw gene-level counts. All code used for the analysis performed in this study is available at https://github.com/BradleyH017/Harris_et_al_WGCNAscRNA.

Pre-processing, dimensionality reduction and clustering of scRNAseq data

All subsequent expression analysis was completed in R version 4.0.2 (R Core Team, 2020). Once raw counts had been obtained for both datasets, bad quality droplets were filtered by a series of quality control steps: i) potential empty droplets were detected by thresholding at the inflection point of the barcode rank plot of cells, calculated using *DropletUtils* v1.8.0 (Griffiths et al., 2018), ii) genes expressed in fewer than 20 cells were removed, iii) cells with an expression sparsity > 0.99 were removed, iv) cells with proportion of mitochondrial gene expression greater than 2.5*(median absolute deviation) from the median proportion were removed.

Following filtration, counts were loaded into a Seurat object using Seurat v4.0.1 (Hao et al., 2021). Initial clustering was performed using Seurat's reciprocal PCA method of batch correction with *SCTransform* as per Hao *et al.,* (2021) guidelines (https://satijalab.org/seurat/articles/integration_rpca.html). The processed Seurat object was first split by sample and data integration was performed on the first 50 principal components. The principal components of the integrated data were then calculated, which was used to calculate the UMAP embedding for all cells, based on 50 principal components

To identify clusters, a nearest neighbour graph was constructed on the integrated data, using

50 principal components. Clusters were then identified via the *FindClusters* function using a

resolution of 0.6. For the Smillie *et al.,* (2019) data analysis, we utilised a k value of 250, as

this is concordant with the authors analysis and provided the greatest confidence in

identification of clusters than other k values. Concordance was tested by enrichment against

the authors cluster markers; data not shown. For the Elmentaite *et al.,* (2021) analysis, we

utilised a k value of 20. Any value greater than this resulted in failure to detect the tuft-cell

resembling cluster.

To identify potential doublets in the filtered dataset, we utilised *DoubletFinder* v2.0.3

(McGinnis et al., 2019) as per the authors guidelines (https://github.com/chris-mcginnis-

ucsf/DoubletFinder). The fully filtered dataset was then re-used as input for integration,

dimensionality reduction and clustering as described above.

To test the robustness of the clusters we identified, we performed pairwise differential gene

expression analysis by a receptor operator curve test using Seurat's *FindMarkers* function. In

order to only merge clusters which were extremely similar, without over-clustering, we defined

similar clusters as those with less than 30 differentially expressed genes with area under the

curve score of 0.6 differentiating them. We found no clusters that fall below this threshold and

so did not modify our initial clustering in either dataset.

Pan-cluster WGCNA

To analyse correlation of gene expression across all cells from the filtered Smillie *et al.,* (2019)

data, we first subset for the nominally significant 11q23.1 trans-eQTLs previously reported

(Vaughan-Shaw et al., 2021). The relative pseudo-bulked expression was then calculated by:

i) summing reads for each gene across all cells within samples, ii) recombining the summed

reads into a non-normalised bulk matrix, iii) log normalising using TMM normalised size factors,

calculated using *edgeR* v3.32.1 (McCarthy et al., 2012). Log-TMM normalised bulk expression

was then z-scored across genes before analysis.

To perform the network analysis, we used *WGCNA* v1.69 (Langfelder & Horvath, 2008). First,

C11orf53, *COLCA1* and *COLCA2* pseudo-bulk expression was extracted. A soft threshold of

14 was then selected after computation of mean connectivity and scale free topology, using

the recommended 'powerEstimate'. A signed adjacency matrix was then calculated, which

was subsequently used to calculate a topological overlap matrix (TOM). Modules were defined
by using a dynamic tree cut of hierarchically clustered gene expression by average distances.
We did not find any modules with a height of module separation below 0.25 and so did not
merge any module. Module eigengenes were then calculated and their correlation with
binarized sex, batch, site, and relative, pseudo-bulked expression of *C11orf53, COLCA1* and
COLCA2 was then assessed. Correlation p-values were multiple testing corrected by the
Benjamini-Hochberg method (Benjamini & Hochberg, 1995). To visualise the blue module
hub genes, a network object was generated from the TOM of blue module genes using
network v1.17.1 (Butts, 2008). Non-connected genes, along with adjacencies < 0.3, were then
removed and remaining genes were plotted using *ggplot2* v3.3.5 (Wickham, 2009).

Gene set enrichment analysis

All gene set enrichment analysis was carried out using R package fgsea v1.14.0 (Korotkevich
et al., 2021). Genes were ranked by their module membership, gene significance for *C11orf53*
expression, or log fold change of differential expression as stated. In the case where multiple
gene sets were being tested, i.e cluster 11 markers against all Smillie et al., putative markers,
p-values were multiple testing corrected by the false discovery rate method.

Calculation of cluster markers

To calculate the markers of each of our own and the Smillie *et al.,* (2019) clusters, we first
generated a log-transcripts per 10,000 expression matrix for the expression of genes within
each cell. This was done so that markers could be calculated using expression values that
were not affected by the relative expression of genes within this dataset and so were more
applicable to future use. Markers were identified by differential expression of genes within
each cluster and all other clusters combined, using MAST v1.160 (Finak et al., 2015).

Analysing the variability of trans-eQTL targets within clusters

To analyse the variability of the expression of 11q23.1 trans-eQTL targets within each cluster,
we applied the pseudo-bulking approach described in *Pseudo-bulk WGCNA* to each cluster
independently. To make expression variability comparable across samples and clusters, the
expression was z-scored across samples.

To analyse expression variability at the single level, we utilised Seurat's *FindVariableFeatures*
function and variance stabilising transformation. In concordance with the identification of many
trans-eQTL targets identified as markers, their mean expression was very low in several
clusters within each dataset. For within dataset comparison of variance, we therefore utilised
the raw variance, not normalising for mean expression. For the between dataset comparison
of variance, we utilised the normalised variance values.
466
467   **Genotyping of Elmentaite et al., (2021) samples**
468
To identify the rs3087967 genotype of Elmentaite *et al.,* (2021) samples, we utilised two
variant calling approaches. These were chosen on the basis of findings from a recent review
which identified these methods as the most sensitive for this purpose (Liu et al., 2019).
Freebayes (Garrison & Marth, 2012) was used with default settings, genotyping over a 10bp
region including rs3087967. Bcftools variant calling (Danecek et al., 2021) was performed on
chromosome 11 using a minimum base quality of 30, disabling read-pair overlap detection
and not discarding anomalous pairs.
476
477
478   **Sample group definition**
479
In the absence of genotype data for Smillie *et al.,* (2019) patients, we grouped samples into
high and low expressors of a *C11orf53*-associated signature by defining the hub genes of the
pseudo-bulk WGCNA blue module. These were defined by a module membership (MM) > 0.7,
intramodular connectivity (kIM) > 0.7 and network adjacency > 0.3. Samples were then
hierarchically clustered by their relative pseudo-bulked expression of these genes using
complete distance. We tested the robustness of our sample grouping by bootstrapping,
selecting 10 random genes 10,000 times, and counting the number of times this exact
separation was achieved – which was 550 times.
488
489   **Intra-cluster WGCNA**
490
491   To agnostically identify gene-gene relatedness within cluster 11, we performed WGCNA as
492   described (see *Pseudo-bulk WGCNA*)*,* selecting only the 5000 most variable genes using
493   Seurat's *FindVariableFeatures* and variance stabilising transformation. The scale free

494   topology threshold utilised was 6, as per the 'power estimate'. P-values were corrected for
495   multiple testing as before.
496

497   ***Module Preservation Analysis***
498

To analyse the preservation of the relatedness of *C11orf53*-correlated 11q23.1 trans-eQTL
targets, we defined an auxiliary module, comprised of the 12 trans-eQTL targets correlated
with *C11orf53* expression (cor>0.5, p<0.05) within the 'cluster 11 black' module. Pseudo-
bulked expression of each cluster was calculated and subset for the 5000 most variable genes
as before, see *Pseudo-bulk WGCNA* and *Intra-cluster WGCNA*. The expression from each
cluster was then raised to the same scale free topology threshold as cluster 11. The
preservation of all modules, including the auxiliary module, was calculated in a pairwise
manner with cluster 11, following the tutorial by Langfelder & Horvath (2008)
(https://horvath.genetics.ucla.edu/html/CoexpressionNetwork/ModulePreservation/Tutorials/).
To summarise module preservation results, we extracted cor.kME values – a measure of the
similarity of auxiliary module genes with the auxiliary module eigengene, and equivalent
results from cluster 11. We also extracted the nominal p-value for this correlation
(log.p.cor.kME), which was subsequently un-logged. The mean connectivity of auxiliary
module genes with the module eigengene was also calculated within each cluster. Pairwise
correlations (p<0.05) between the expression of the auxiliary module genes with one another
and *C11orf53* expression were plotted using corrplot (Wei & Simko, 2017).
515

516   ***Linear modelling of cluster abundance and pseudo-bulked gene expression***
517

518   Univariate linear modelling of cluster 11 abundance was performed on the TMM normalised,
519   non-z-scored pseudo-bulked expression matrix, so that results are more relevant to further
520   study. We fitted a linear model to all genes and performed empirical Bayes moderation using
521   *limma* v3.46.0 (Ritchie et al., 2015). P-values were adjusted by Benjamini-Hochberg multiple
522   testing correction (Benjamini & Hochberg, 1995).
523

524   ***Differential abundance testing***
525

Differential abundance testing was performed using miloR v0.99.8 (Dann et al., 2021). To
mitigate any potential for package-specific artefacts of the analysis, we first re-generated the
k nearest neighbour graph using 250 nearest neighbours from the integrated expression. The

graph was then built using 4 PCA components. Differential abundance testing was carried out

using a Quasi-Likelihood F-test on TMM normalised cell proportions. Differentially abundant

neighbourhood results were then annotated for their majority cluster proportion and those

comprised of fewer than 80% of the majority cluster were removed.

Author contributions

Conceptualisation: BTH, VR, MGD, SMF

Methodology: BTH, VR, SMF, MGD, AO, JPB, KD, MT, PGVS

Investigation: BTH

Original draft writing: BTH

Review & Editing: BTH, VR, JPB, KD, MT, AO, MGD, SMF

Funding Acquisition: SMF, MGD, FVND

Supervision: VR, MGD, SMF

    Supplementary Material

Subject ID	Status	Location	Gender
N8	Healthy	#N/A	Female
N10	Healthy	Right_Colon	Female
N11	Healthy	Right_Colon	Male
N13	Healthy	Right_Colon	Female
N15	Healthy	Right_Colon	Male
N16	Healthy	Right_Colon	Male
N17	Healthy	Transverse_Colon	Male
N18	Healthy	Transverse_Colon	Female
N20	Healthy	#N/A	Female
N21	Healthy	Left_Colon	Female
N46	Healthy	Right_Colon	Male

Supplementary Table 1. Smillie *et al.,* (2019) patient metadata

Supplementary Figure 1. Correlation of Module Membership (MM) and Gene Significance (GS) for *C11orf53* expression in the blue module. 11q23.1 trans-eQTL targets are highlighted, red if MM.blue and GS.C11orf53 > 0.5.

Gene_symbol	GS.C11orf53	p.GS.C11orf53
MFSD7	0.869516061	1.49E-07
P2RY2	0.852996421	4.57E-07
FANK1	0.849691779	5.63E-07
LINC00261	0.839485426	1.04E-06
KLK4	0.837697653	1.15E-06
PLEKHG4	0.771254176	2.65E-05
RNF32	0.754523148	4.97E-05
ALOX5	0.752415106	5.36E-05
PTPRD	0.740741398	8.05E-05
ASTE1	0.714297945	0.000188259
SH2D6	0.71110102	2.07E-04
HCK	0.698902984	2.96E-04
TRPM5	0.689672239	3.84E-04
BMX	0.689502913	3.85E-04
PSTPIP2	0.682088126	4.71E-04
MYOM1	0.670290717	6.42E-04
GNG13	0.657207652	8.90E-04
IL17RB	0.655879017	0.000918972
CYR61	0.655513165	0.000927169
HTR3E	0.652281	0.001002333
PTGS1	0.651790925	0.001014171
WWP2	0.643465569	0.001234253
FYB	0.638720231	0.001376965
SH2D7	0.63416094	0.001527043
SNRPA	0.632098175	0.00159937
AC025335.1	0.632010561	0.001602506
RAI1	0.631424527	0.001623611
MARCKSL1	0.627445473	0.001773235
OGDHL	0.626140466	0.00182478
CRYM	0.621805521	0.002005201
MATK	0.620641028	0.002056158
ALOX5AP	0.604524135	0.002881786
RP11-315O6.1	0.599692821	0.003177792
MSLN	0.587571695	0.004034646
PLCG2	0.573987877	0.005216381
SCRN1	0.571174034	0.005494069
CD96	0.555950868	0.007218247
L3MBTL1	0.547690051	0.008327128
RP11-1220K2.2	0.531364238	0.010932727
IKZF4	0.528361091	0.011477883
FAM65A	0.495417788	0.019052761
GFI1B	0.471411163	0.026778703
SGK223	0.467790056	0.028134237
RNF144A	0.466520671	0.028622119
CHSY1	0.460356935	0.031087422
DKK3	0.448035289	0.036517806
ZFP1	0.432558977	0.044363985
AFAP1L2	0.401466056	0.064037752
TBC1D24	0.397206607	0.067180992
LRMP	0.396518404	0.067699644
PODXL	0.394606685	0.069156315
SLC25A30	0.391914382	0.071247869
ARHGEF3	0.375600915	0.084958526
ARHGEF2	0.372293733	0.087962678
FILIP1L	0.370412873	0.089706055
C20orf194	0.362507448	0.097315042
NOTCH4	0.362506253	0.097316226
ARHGAP31	0.3494408	0.11091913
PGAM4	0.34556863	0.115204208
FGFRL1	0.32779629	0.136419162
PIK3CG	0.327080559	0.137328021
ATF3	0.305953301	0.16612758
PPP1R3B	0.301754652	0.172314946
CAPN9	0.290378683	0.189872811
APH1B	0.283453554	0.201136526
MT1H	0.283442955	0.201154103
GNB5	0.265793153	0.23186717
SNORA31	0.249996893	0.261829995
HTR3C	0.20872	0.351252864
TRIB1	0.203124291	0.364604113
COL1A1	0.197820332	0.377525123
PIP5KL1	0.174089375	0.438438491
CGNL1	0.149829973	0.505720737
CAMP	0.126754067	0.574047366
ISPD	0.099579334	0.659282892
CAPRIN2	0.053739047	0.812256115
DLG5-AS1	0.003234775	0.988601201

600 **Supplementary Table 2.** Blue module genes, corresponding Gene Significance for *C11orf53* (GS.C11orf53) and p-value of Gene Significance for *C11orf53* (p.GS.C11orf53). Red=11q23.1 trans-601 eQTL target (Vaughan-Shaw *et al.,* 2021)

11q23.1 trans-eQTL target variability across Smillie et al., clusters

Supplementary Figure 2. Investigating 11q23.1 eQTL target variance at the single cell level. Variance of 11q23.1 eQTL targets across clusters at the single cell level in the Smillie *et al.,* (2019) data. Variance is z-scored within genes.

Supplementary Figure 3. Replication of 11q23.1 eQTL expression mapping in independent dataset. (A) UMAP of 11,651 healthy colonic epithelial scRNASeq (Elmentaite *et al.,* 2021). (B) Upper: GSEA of 11q23.1 trans-eQTLs (Vaughan-Shaw *et al.,* 2021; FDR<0.05) in cluster 18. Lower: GSEA of putative colonic tuft cell signature (Smillie *et al.,* 2019) in cluster 18. (D) Relative expression of 11q23.1 trans-eQTLs (FDR<0.05) across clusters**.**

Gene	p_val	avg_log2FC	pct.1	pct.2	p_val_adj
LRMP	2.04E-73	2.614295875	0.903	0.004	3.19E-69
IL17RB	6.17E-69	1.454372429	0.516	0.033	9.64E-65
SH2D6	3.81E-64	2.521466548	0.806	0.002	5.95E-60
PLCG2	1.11E-50	2.182226523	0.71	0.011	1.73E-46
PSTPIP2	1.31E-45	1.656282707	0.581	0.006	2.04E-41
TRPM5	6.51E-44	1.403600676	0.581	0.002	1.02E-39
SH2D7	2.90E-39	1.090698798	0.484	0.001	4.53E-35
AZGP1	9.92E-39	1.061625996	0.452	0.005	1.55E-34
PTGS1	8.37E-30	1.282060245	0.516	0.01	1.31E-25
ALOX5	4.14E-29	1.32717779	0.452	0.004	6.46E-25
BMX	1.08E-22	1.007016149	0.419	0.012	1.68E-18

Supplementary Table 3. 11q23.1 trans-eQTL targets identified as cluster 18 markers from Elmentaite *et al.,* (2021). Markers calculated using MAST (McDavid *et al.,* 2020). Avg_log2FC=average log2 fold change between cluster 11 and all other clusters. Pct1=Proportion of cluster 18 cells expressing gene. Pct2=Proportion of non-cluster 18 cells expressing gene.

11q23.1 trans-eQTL target variability across Elmentaite et al., clusters

637 **Supplementary Figure 4.** Investigating 11q23.1 eQTL target variance at the single cell level. Variance of 11q23.1 eQTL targets across clusters at the single cell level in Elmentaite *et al.,* (2021) data. Variance is z-scored within genes.

Sample	Subject	freebayes_rs3087967	samtools_rs3087967
Human_colon_16S8123908	411C	CC	CC
Human_colon_16S8000473	417C	T/C	CC
Human_colon_16S8000477	417C	CC	CC
Human_colon_16S8000481	417C	CC	CC
Human_colon_16S8000479	417C	CC	CC
Human_colon_16S8000475	417C	CC	CC
Human_colon_16S8002627	432c	CC	CC
Human_colon_16S8002630	432c	CC	CC

Supplementary Table 4. Elmentaite *et al.,* (2019) samples are homozygous non-risk at rs3087967. rs3087967 genotype results of both freebayes (Garrison et al., 2012) and Bcftools (Danecek et al., 2021) variant calling methods. T=colorectal cancer risk allele, C=non-risk allele

Gene	Smillie_cluster11	Elmentaite_cluster18	FC
C11orf53	1.275873075	0.656341714	1.94391587
COLCA1	1.346177989	NA	NA
COLCA2	1.381191066	0.834903033	1.65431315
LRMP	1.695513426	0.173077239	9.79628192
SH2D6	0.518050076	0.196087378	2.64193484
PSTPIP2	0.961252255	0.332407278	2.89179064
HTR3C	1.850394057	NA	NA
ALOX5	0.855502479	0.467755885	1.82895075
OGDHL	0.83318323	1.141132077	0.73013742
MATK	0.815029157	NA	NA
PIK3CG	1.103714386	0.73167938	1.50846726
PLCG2	0.871942447	0.249476603	3.49508706
PTGS1	0.673534891	0.461933218	1.45807849
IL17RB	0.8969865	0.467309219	1.91947101
GNG13	1.024981369	NA	NA
CAMP	2.260972819	NA	NA
ANKHD1	0.897739169	NA	NA
GIN1	1	NA	NA
SH2D7	0.658046427	0.328744382	2.00169634
BMX	1.360003183	0.360535132	3.77217936
HTR3E	0.848772333	NA	NA
AZGP1	1.029538915	0.515296922	1.99795277
TRPM5	0.599018856	0.314108919	1.90704186

Supplementary Table 5. Variability of the majority of 11q23.1 trans-eQTL targets is greater in Smillie *et al.,* (2019) cluster 11 compared with Elmentaite *et al.,* (2021) cluster 18. Variability analysed within single cells using Seurat *FindVariableGenes* (Hao et al., 2021). FC = Fold change of variability in Smillie *et al.,* cluster 11 compared with Elmentaite *et al.,* (2021) cluster 18

Supplementary Figure 5. UMAP of cells from each Blue module hub gene group (left) and the differential abundance score of miloR-identified (Dann *et al.,* 2021) neighbourhoods (right).

References

Banerjee, A., Herring, C. A., Chen, B., Kim, H., Simmons, A. J., Southard-Smith, A. N., Allaman, M. M., White, J. R., Macedonia, M. C., Mckinley, E. T., Ramirez-Solano, M. A., Scoville, E. A., Liu, Q., Wilson, K. T., Coffey, R. J., Washington, M. K., Goettel, J. A., & Lau, K. S. (2020). Succinate Produced by Intestinal Microbes Promotes Specification of Tuft Cells to Suppress Ileal Inflammation. *Gastroenterology*, *159*(6). https://doi.org/10.1053/j.gastro.2020.08.029

Benjamini, Y., & Hochberg, Y. (1995). Controlling the false discovery rate: A practical and powerful approach to multiple testing. *Journal of the Royal Statistical Society*, *57*(1), 289–300.

Biancolella, M., Fortini, B. K., Tring, S., Plummer, S. J., Mendoza-Fandino, G. A., Hartiala, J., Hitchler, M. J., Yan, C., Schumacher, F. R., Conti, D. v., Edlund, C. K., Noushmehr, H., Coetzee, S. G., Bresalier, R. S., Ahnen, D. J., Barry, E. L., Berman, B. P., Rice, J. C., Coetzee, G. A., & Casey, G. (2014). Identification and characterization of functional risk variants for colorectal cancer mapping to chromosome 11q23.1. *Human Molecular Genetics*, *23*(8). https://doi.org/10.1093/hmg/ddt584

Blackmur, J. P., Vaughan-Shaw, P. G., Donnelly, K., Harris, B. T., Svinti, V., Ochocka-Fox, A.-M., Freile, P., Walker, M., Gurran, T., Reid, S., Semple, C. A., Din, F. V. N., Timofeeva, M., Dunlop, M. G., & Farrington, S. M. (2022). Gene Co-Expression Network Analysis Identifies Vitamin D-Associated Gene Modules in Adult Normal Rectal Epithelium Following Supplementation. *Frontiers in Genetics*, *12*. https://doi.org/10.3389/fgene.2021.783970

Butts, Carter. T. (2008). network: A package for managing relational data in R. *Journal of Statistical Software*, *24*(2).

Closa, A., Cordero, D., Sanz-Pamplona, R., Solé, X., Crous-Bou, M., Paré-Brunet, L., Berenguer, A., Guino, E., Lopez-Doriga, A., Guardiola, J., Biondo, S., Salazar, R., & Moreno, V. (2014). Identification of candidate susceptibility genes for colorectal cancer through eQTL analysis. *Carcinogenesis*, *35*(9). https://doi.org/10.1093/carcin/bgu092

Danecek, P., Bonfield, J. K., Liddle, J., Marshall, J., Ohan, V., Pollard, M. O., Whitwham, A., Keane, T., McCarthy, S. A., Davies, R. M., & Li, H. (2021). Twelve years of SAMtools and BCFtools. *GigaScience*, *10*(2). https://doi.org/10.1093/gigascience/giab008

Dann, E., Henderson, N. C., Teichmann, S. A., Morgan, M. D., & Marioni, J. C. (2021). Differential abundance testing on single-cell data using k-nearest neighbor graphs. *Nature Biotechnology*. https://doi.org/10.1038/s41587-021-01033-z

DelGiorno, K. E., Chung, C.-Y., Vavinskaya, V., Maurer, H. C., Novak, S. W., Lytle, N. K., Ma, Z., Giraddi, R. R., Wang, D., Fang, L., Naeem, R. F., Andrade, L. R., Ali, W. H., Tseng, H., Tsui, C., Gubbala, V. B., Ridinger-Saison, M., Ohmoto, M., Erikson, G. A., … Wahl, G. M. (2020). Tuft Cells Inhibit Pancreatic Tumorigenesis in Mice by Producing Prostaglandin D2. *Gastroenterology*, *159*(5). https://doi.org/10.1053/j.gastro.2020.07.037

Elmentaite, R., Kumasaka, N., Roberts, K., Fleming, A., Dann, E., King, H. W., Kleshchevnikov, V., Dabrowska, M., Pritchard, S., Bolt, L., Vieira, S. F., Mamanova, L., Huang, N., Perrone, F., Goh Kai'En, I., Lisgo, S. N., Katan, M., Leonard, S., Oliver, T. R. W., … Teichmann, S. A. (2021). Cells of the human intestinal tract mapped across space and time. *Nature*, *597*(7875). https://doi.org/10.1038/s41586-021-03852-1

Finak, G., McDavid, A., Yajima, M., Deng, J., Gersuk, V., Shalek, A. K., Slichter, C. K., Miller, H. W., McElrath, M. J., Prlic, M., Linsley, P. S., & Gottardo, R. (2015). MAST: a flexible statistical framework for assessing transcriptional changes and characterizing heterogeneity in single-cell RNA sequencing data. *Genome Biology*, *16*(1), 278. https://doi.org/10.1186/s13059-015-0844-5

Garrison, E., & Marth, G. (2012). *Haplotype-based variant detection from short-read sequencing*.

Gerbe, F., Sidot, E., Smyth, D. J., Ohmoto, M., Matsumoto, I., Dardalhon, V., Cesses, P., Garnier, L., Pouzolles, M., Brulin, B., Bruschi, M., Harcus, Y., Zimmermann, V. S., Taylor, N., Maizels, R. M., & Jay, P. (2016). Intestinal epithelial tuft cells initiate type 2 mucosal immunity to helminth parasites. *Nature*, *529*(7585). https://doi.org/10.1038/nature16527

Gerbe, F., van Es, J. H., Makrini, L., Brulin, B., Mellitzer, G., Robine, S., Romagnolo, B., Shroyer, N.
F., Bourgaux, J.-F., Pignodel, C., Clevers, H., & Jay, P. (2011). Distinct ATOH1 and Neurog3
requirements define tuft cells as a new secretory cell type in the intestinal epithelium. *Journal of*
Cell Biology, *192*(5). https://doi.org/10.1083/jcb.201010127

Graff, R. E., Möller, S., Passarelli, M. N., Witte, J. S., Skytthe, A., Christensen, K., Tan, Q., Adami, H.-
O., Czene, K., Harris, J. R., Pukkala, E., Kaprio, J., Giovannucci, E. L., Mucci, L. A., &
Hjelmborg, J. B. (2017). Familial Risk and Heritability of Colorectal Cancer in the Nordic Twin
Study of Cancer. *Clinical Gastroenterology and Hepatology*, *15*(8), 1256–1264.
https://doi.org/10.1016/j.cgh.2016.12.041

Griffiths, J. A., Richard, A. C., Bach, K., Lun, A. T. L., & Marioni, J. C. (2018). Detection and removal
of barcode swapping in single-cell RNA-seq data. *Nature Communications*, *9*(1), 2667.
https://doi.org/10.1038/s41467-018-05083-x

Haber, A. L., Biton, M., Rogel, N., Herbst, R. H., Shekhar, K., Smillie, C., Burgin, G., Delorey, T. M.,
Howitt, M. R., Katz, Y., Tirosh, I., Beyaz, S., Dionne, D., Zhang, M., Raychowdhury, R., Garrett,
W. S., Rozenblatt-Rosen, O., Shi, H. N., Yilmaz, O., … Regev, A. (2017). A single-cell survey of
the small intestinal epithelium. *Nature*, *551*(7680). https://doi.org/10.1038/nature24489

Hao, Y., Hao, S., Andersen-Nissen, E., Mauck, W. M., Zheng, S., Butler, A., Lee, M. J., Wilk, A. J.,
Darby, C., Zager, M., Hoffman, P., Stoeckius, M., Papalexi, E., Mimitou, E. P., Jain, J.,
Srivastava, A., Stuart, T., Fleming, L. M., Yeung, B., … Satija, R. (2021). Integrated analysis of
multimodal single-cell data. *Cell*, *184*(13). https://doi.org/10.1016/j.cell.2021.04.048

Hoffman, M. T., Kemp, S. B., Salas-Escabillas, D. J., Zhang, Y., Steele, N. G., The, S., Long, D.,
Benitz, S., Yan, W., Margolskee, R. F., Bednar, F., Pasca di Magliano, M., Wen, H.-J., &
Crawford, H. C. (2021). The Gustatory Sensory G-Protein GNAT3 Suppresses Pancreatic
Cancer Progression in Mice. *Cellular and Molecular Gastroenterology and Hepatology*, *11*(2).
https://doi.org/10.1016/j.jcmgh.2020.08.011

Hulur, I., Gamazon, E. R., Skol, A. D., Xicola, R. M., Llor, X., Onel, K., Ellis, N. A., & Kupfer, S. S.
(2015). Enrichment of inflammatory bowel disease and colorectal cancer risk variants in colon
expression quantitative trait loci. *BMC Genomics*, *16*(1). https://doi.org/10.1186/s12864-015-
1292-z

Huyghe, J. R., Bien, S. A., Harrison, T. A., Kang, H. M., Chen, S., Schmit, S. L., Conti, D. v., Qu, C.,
Jeon, J., Edlund, C. K., Greenside, P., Wainberg, M., Schumacher, F. R., Smith, J. D., Levine,
D. M., Nelson, S. C., Sinnott-Armstrong, N. A., Albanes, D., Alonso, M. H., … Peters, U. (2019).
Discovery of common and rare genetic risk variants for colorectal cancer. *Nature Genetics*,
51(1). https://doi.org/10.1038/s41588-018-0286-6

Jasperson, K. W., Tuohy, T. M., Neklason, D. W., & Burt, R. W. (2010). Hereditary and Familial Colon
Cancer. *Gastroenterology*, *138*(6). https://doi.org/10.1053/j.gastro.2010.01.054

Kaske, S., Krasteva, G., König, P., Kummer, W., Hofmann, T., Gudermann, T., & Chubanov, V.
(2007). TRPM5, a taste-signaling transient receptor potential ion-channel, is a ubiquitous
signaling component in chemosensory cells. *BMC Neuroscience*, *8*(1).
https://doi.org/10.1186/1471-2202-8-49

Kjærgaard, S., Jensen, T. S. R., Feddersen, U. R., Bindslev, N., Grunddal, K. v., Poulsen, S. S.,
Rasmussen, H. B., Budtz-Jørgensen, E., & Berner-Hansen, M. (2021). Decreased number of
colonic tuft cells in quiescent ulcerative colitis patients. *European Journal of Gastroenterology &*
Hepatology, *33*(6). https://doi.org/10.1097/MEG.0000000000001959

Korotkevich, G., Sukhov, V., Budin, N., Shpak, B., Artyomov, M. N., & Sergushichev, A. (2021). Fast
gene set enrichment analysis. *BioRxiv*.

Langfelder, P., & Horvath, S. (2008). WGCNA: an R package for weighted correlation network
analysis. *BMC Bioinformatics*, *9*(1). https://doi.org/10.1186/1471-2105-9-559

Law, P. J., Timofeeva, M., Fernandez-Rozadilla, C., Broderick, P., Studd, J., Fernandez-Tajes, J.,
Farrington, S., Svinti, V., Palles, C., Orlando, G., Sud, A., Holroyd, A., Penegar, S.,

Theodoratou, E., Vaughan-Shaw, P., Campbell, H., Zgaga, L., Hayward, C., Campbell, A., …
Dunlop, M. G. (2019). Association analyses identify 31 new risk loci for colorectal cancer
susceptibility. *Nature Communications*, *10*(1). https://doi.org/10.1038/s41467-019-09775-w

Liu, F., Zhang, Y., Zhang, L., Li, Z., Fang, Q., Gao, R., & Zhang, Z. (2019). Systematic comparative
analysis of single-nucleotide variant detection methods from single-cell RNA sequencing data.
Genome Biology, *20*(1), 242. https://doi.org/10.1186/s13059-019-1863-4

Loo, L. W. M., Lemire, M., & le Marchand, L. (2017). In silico pathway analysis and tissue specific cis-
eQTL for colorectal cancer GWAS risk variants. *BMC Genomics*, *18*(1).
https://doi.org/10.1186/s12864-017-3750-2

Luo, Y., Coskun, V., Liang, A., Yu, J., Cheng, L., Ge, W., Shi, Z., Zhang, K., Li, C., Cui, Y., Lin, H.,
Luo, D., Wang, J., Lin, C., Dai, Z., Zhu, H., Zhang, J., Liu, J., Liu, H., … Li, S. (2015). Single-Cell
Transcriptome Analyses Reveal Signals to Activate Dormant Neural Stem Cells. *Cell*, *161*(5).
https://doi.org/10.1016/j.cell.2015.04.001

McCarthy, D. J., Chen, Y., & Smyth, G. K. (2012). Differential expression analysis of multifactor RNA-
Seq experiments with respect to biological variation. *Nucleic Acids Research*, *40*(10).
https://doi.org/10.1093/nar/gks042

McGinnis, C. S., Murrow, L. M., & Gartner, Z. J. (2019). DoubletFinder: Doublet Detection in Single-
Cell RNA Sequencing Data Using Artificial Nearest Neighbors. *Cell Systems*, *8*(4).
https://doi.org/10.1016/j.cels.2019.03.003

Michaelson, J. J., Loguercio, S., & Beyer, A. (2009). Detection and interpretation of expression
quantitative trait loci (eQTL). *Methods*, *48*(3). https://doi.org/10.1016/j.ymeth.2009.03.004

Middelhoff, M., Westphalen, C. B., Hayakawa, Y., Yan, K. S., Gershon, M. D., Wang, T. C., & Quante,
M. (2017). Dclk1-expressing tuft cells: critical modulators of the intestinal niche? *American
Journal of Physiology-Gastrointestinal and Liver Physiology*, *313*(4).
https://doi.org/10.1152/ajpgi.00073.2017

Rawla, P., Sunkara, T., & Barsouk, A. (2019). Epidemiology of colorectal cancer: incidence, mortality,
survival, and risk factors. *Gastroenterology Review*, *14*(2).
https://doi.org/10.5114/pg.2018.81072

Richards, A. L., Jones, L., Moskvina, V., Kirov, G., Gejman, P. v, Levinson, D. F., Sanders, A. R.,
Purcell, S., Visscher, P. M., Craddock, N., Owen, M. J., Holmans, P., & O'Donovan, M. C.
(2012). Schizophrenia susceptibility alleles are enriched for alleles that affect gene expression in
adult human brain. *Molecular Psychiatry*, *17*(2). https://doi.org/10.1038/mp.2011.11

Ritchie, M. E., Phipson, B., Wu, D., Hu, Y., Law, C. W., Shi, W., & Smyth, G. K. (2015). limma powers
differential expression analyses for RNA-sequencing and microarray studies. *Nucleic Acids
Research*, *43*(7). https://doi.org/10.1093/nar/gkv007

Smillie, C. (2015). *Functional characterisation of the 11q23.1 colorectal cancer risk locus*, PhD thesis,
The University of Edinburgh, Edinburgh

Smillie, C. S., Biton, M., Ordovas-Montanes, J., Sullivan, K. M., Burgin, G., Graham, D. B., Herbst, R.
H., Rogel, N., Slyper, M., Waldman, J., Sud, M., Andrews, E., Velonias, G., Haber, A. L.,
Jagadeesh, K., Vickovic, S., Yao, J., Stevens, C., Dionne, D., … Regev, A. (2019). Intra- and
Inter-cellular Rewiring of the Human Colon during Ulcerative Colitis. *Cell*, *178*(3).
https://doi.org/10.1016/j.cell.2019.06.029

Stranger, B. E., Nica, A. C., Forrest, M. S., Dimas, A., Bird, C. P., Beazley, C., Ingle, C. E., Dunning,
M., Flicek, P., Koller, D., Montgomery, S., Tavaré, S., Deloukas, P., & Dermitzakis, E. T. (2007).
Population genomics of human gene expression. *Nature Genetics*, *39*(10).
https://doi.org/10.1038/ng2142

Sud, A., Kinnersley, B., & Houlston, R. S. (2017). Genome-wide association studies of cancer: current
insights and future perspectives. *Nature Reviews Cancer*, *17*(11).
https://doi.org/10.1038/nrc.2017.82

Tenesa, A., Farrington, S. M., Prendergast, J. G. D., Porteous, M. E., Walker, M., Haq, N., Barnetson, R. A., Theodoratou, E., Cetnarskyj, R., Cartwright, N., Semple, C., Clark, A. J., Reid, F. J. L., Smith, L. A., Kavoussanakis, K., Koessler, T., Pharoah, P. D. P., Buch, S., Schafmayer, C., … Dunlop, M. G. (2008). Genome-wide association scan identifies a colorectal cancer susceptibility locus on 11q23 and replicates risk loci at 8q24 and 18q21. *Nature Genetics*, *40*(5). https://doi.org/10.1038/ng.133

Vaughan-Shaw, Peter. G., Timofeeva, M., Ooi, L., Svinti, V., Grimes, G., Smillie, C., Blackmur, J. P., Donnelly, K., Theodoratou, E., Campbell, H., Zgaga, L., Din, F. V. N., Farrington, S. M., & Dunlop, M. G. (2021). Differential genetic influences over colorectal cancer risk and gene expression in large bowel mucosa. *International Journal of Cancer*, *149*(5). https://doi.org/10.1002/ijc.33616

Wei, T., & Simko, V. (2017). *"Corrplot": Visualization of a Correlation Matrix (Version 0.84)*.

Wickham, H. (2009). *ggplot2: Elegant graphics for data analysis*. Springer-Verlag.

Wu, C.-L., Dicks, A., Steward, N., Tang, R., Katz, D. B., Choi, Y.-R., & Guilak, F. (2021). Single cell transcriptomic analysis of human pluripotent stem cell chondrogenesis. *Nature Communications*, *12*(1). https://doi.org/10.1038/s41467-020-20598-y

Yamaga, Y., Fukuda, A., Nakanishi, Y., Goto, N., Matsumoto, Y., Yoshioka, T., Maruno, T., Chiba, T., & Seno, H. (2018). Gene expression profile of Dclk1+ cells in intestinal tumors. *Digestive and Liver Disease*, *50*(12). https://doi.org/10.1016/j.dld.2018.06.011

Yi, J., Bergstrom, K., Fu, J., Shan, X., McDaniel, J. M., McGee, S., Qu, D., Houchen, C. W., Liu, X., & Xia, L. (2019). Dclk1 in tuft cells promotes inflammation-driven epithelial restitution and mitigates chronic colitis. *Cell Death & Differentiation*, *26*(9). https://doi.org/10.1038/s41418-018-0237-x

Zheng, G. X. Y., Terry, J. M., Belgrader, P., Ryvkin, P., Bent, Z. W., Wilson, R., Ziraldo, S. B., Wheeler, T. D., McDermott, G. P., Zhu, J., Gregory, M. T., Shuga, J., Montesclaros, L., Underwood, J. G., Masquelier, D. A., Nishimura, S. Y., Schnall-Levin, M., Wyatt, P. W., Hindson, C. M., … Bielas, J. H. (2017). Massively parallel digital transcriptional profiling of single cells. *Nature Communications*, *8*(1). https://doi.org/10.1038/ncomms14049

Acknowledgements

Funding: This work was supported by Cancer Research UK (CRUK) PhD studentship at the Edinburgh CRUK Cancer Research Centre (Bradley T. Harris / Susan M. Farrington) and CRUK programme grant DRCPGM\100012 (Malcolm G. Dunlop / Susan M. Farrington). James P. Blackmur was supported by an ECAT-linked CRUK ECRC Clinical training award (C157/A23218). Peter Vaughan-Shaw was supported by a NES SCREDS clinical lectureship, MRC Clinical Research Training Fellowship (MR/M004007/1), a Research Fellowship from the Harold Bridges bequest and by the Melville Trust for the Care and Cure of Cancer. **Competing Interests:** The authors declare no competing interests. **Data availability:** All data used in this study has been referenced within the Materials and Methods. **Open Access:** For the purpose of open access, the author has applied a Creative Commons Attribution (CC BY) licence to any Author Accepted Manuscript version arising from this submission.