Development of a new 3D tracking system for multiple marmosets under free-moving conditions

NEW TRACKING SYSTEM FOR FREE-RUNNING CONDITIONS OF MARMOSETS

Terumi Yurimoto[1], Wakako Kumita[1], Kenya Sato[1], Rika Kikuchi[1], Yusuke Shibuki [1], Rino Hashimoto[1],

Michiko Kamioka[1], Yumi Hayasegawa[1], Eiko Yamazaki[1], Yoko Kurotaki[2], Norio Goda[3], Junichi

Kitakami[4], Tatsuya Fujita[5], Takashi Inoue[1], Erika Sasaki[1*]

[1]Department of Marmoset Biology and Medicine, Central Institute for Experimental Animals,

Kawasaki, 210-0821 JAPAN

[2]Center of Basic Technology in Marmoset, Central Institute for Experimental Animals, Kawasaki,

210-0821 JAPAN

[3]Public Digital Transformation Department, Hitachi, Ltd., Shinagawa, 140-8512 JAPAN

[4]Vision AI Solution Design Department Hitachi Solutions Technology, Ltd., Tachikawa, 190-0014

JAPAN

[5]Engineering Department Eastern Japan division, Totec Amenity Limited, Shinjuku, 163-0417 JAPAN

*Corresponding author: Erika Sasaki (esasaki@ciea.or.jp)

Abstract

Assessment of social interactions and behavioral changes in nonhuman primates is useful for

understanding brain function changes during life events and pathogenesis of neurological diseases.

The common marmoset (*Callithrix jacchus*), that lives in a nuclear family like humans, is a useful

model, but long-term automated behavioral observation of multiple animals has not been achieved.

Here, we developed a Full Monitoring and Animal Identification (FulMAI) system for long-term

detection of three-dimensional (3D) trajectories of each individual in multiple marmosets under free-

moving conditions by combining video tracking, Light Detection And Ranging, and deep learning.

Using this system, identification of each animal was more than 97% accurate. Location preferences

and inter-individual distance could be calculated, and deep learning could detect grooming behavior.

The FulMAI system allows us to analyze the natural behavior of individuals in a family over their

lifetime and understand how behavior changes due to life events together with other data.

Introduction

Behavioral analyses are important for understanding brain function changes during development and aging and for evaluating disease onset in model animals. In particular, nonhuman primates are often used to study brain function, and behavioral characteristics provide an indispensable source of data for hypothesis testing[1]. Nonhuman primates share anatomical and physiological features with humans, especially in specialized functions acquired through unique cerebral expansions, such as tool use, language (for example, vocal communication), and self-awareness[2, 3, 4, 5]. The common marmoset (*Callithrix jacchus*) is a small nonhuman primate with behavioral and social characteristics resembling those of humans. They are diurnal, form monogamous families, engage in altruistic behaviors such as food sharing, and cooperate with all family members to raise their youngest offspring[3, 6, 7]. As marmoset body size and social units are small, maintaining complete social units for research in a laboratory is easier than that for other primates. Therefore, marmosets are an ideal model for studying social behaviors in social units[3]. Recent developments in genetic modification techniques in marmosets have enabled the production of various disease-model marmosets, such as those for neurodegenerative diseases[8, 9, 10]. Using marmosets as model animals has advantages over rodents and may help elucidate basic gene functions in the brain, clarify molecular mechanisms underlying neurological disease onset using genetic modifications, and aid in implementing cognitive neuroscience techniques in primate research[5, 11]. For these purposes, national brain research projects

using marmosets have been conducted in the US and Japan[12, 13].

To evaluate long-term brain function changes due to development, aging, and disease onset

using marmosets as a model, it is important to capture changes in behavior over a lifespan, and an

analysis of multiple marmosets under free-moving conditions is necessary to capture changes in social

behavior. Therefore, a novel system is required to detect and quantify the individual behaviors of

multiple marmosets under free-moving conditions in real time, both automatically and accurately.

Automated behavior analysis in a home cage enable observations of more natural animal behavior[14,

15, 16]. This analysis has the advantages of being able to ignore environmental condition, monitor social

behavior in multiple animals, and analyze changes in social behavior over a relatively long period of

time[17].

The various systems used for analyzing animal behaviors under free-moving activities

include systems suitable for three dimensions (3D) tracking and behavior classification (DANNCE[18],

FreiPose [19], and MarmoDetecotor[20]), for generating pose estimation (DeepLabCut, SLEAP[21], MARS[22],

DANNCE, FreiPose, and OpenMonkeyStudio[1]), for analyzing behavior in home cages (DeepLabCut

23, 24, B-SoiD [17], SLEAP, and MARS), for analyzing multiple animals (DeepLabCut, MiceProfier[25],

SLEAP, and OpenMonkeyStudio), for finding novel behavioral abnormalities by unsupervised

clustering (MoSeq[26], B-SoiD, and FreiPose[19]), and for use across animal species (DeepLabCut,

DeepBhvTracking[27], B-SoiD, and DANNCE). These systems have been generally developed for mice;

only DeepLabCut, DeepBhvTracking, DANNCE, and MarmoDetector[20] have been adapted to

marmosets, which move relatively quickly in 3D. A previous study used the pose estimation system

to analyze the behavior of two marmosets for nine hours using individual identification markers[23].

These analysis systems have observed only a small proportion of an animal's life and longitudinal

observation over each marmoset's lifetime (15 years) or more than months of observing a family have

not been achieved. This is possibly because automatically detecting marmoset trajectories, behaviors,

and interactions between individuals under free-moving conditions in multiple animals is especially

challenging. For example, marmosets are often close to each other when grooming, mating, or

fighting[28]. This proximity makes it difficult to detect individual bodies, which complicates individual

tracking and behavior detection.

In this study, we developed a behavioral analysis system named FulMAI (Full Monitoring

and Animal Identification), that can record and analyze behavior of an animal living among multiple

animals over their lifetime. The concept of this system is to detect changes in behavior over an

individual's lifespan and understand the relation of changes in brain function. FulMAI is a 3D tracking

system that uses cameras, light detection and ranging (Lidar) devices, and deep learning to

simultaneously track a marmoset family of three individuals in one cage. Furthermore, to take

advantage of stress-free behavior analysis in the home cage, we also developed a system for individual

identification via facial recognition using deep learning. The FulMAI system can analyze the location

and the period each animal stayed in the cage and the distance between individuals. Furthermore, the

time spent in social behavior is an index for analyzing animal interactions. Therefore, as an example

of social behavior detection, we developed automatic grooming behavior detection system using deep

learning. The technology developed in this study would be applicable in detecting important

behavioral changes caused by development, aging, and various diseases in small, nonhuman model

primates.

Results

Marmoset facial recognition by deep learning

Five marmosets of the same generation (1–2 years old) were used to examine whether deep learning

for face identification could be adapted to marmosets. These marmosets were kept in different cages

in a conventional breeding colony and facial images were collected individually. To evaluate the

trained models, 500 new facial images (100 for each animal) were obtained from the videos not used

for training. The evaluation metrics for deep learning were as follows: specificity of 99.8%–100%,

accuracy of 99.4%–99.8%, recall of 98.0%–100%, precision of 99%–100%, and an F-measure of

98.5%–99.5% (Table 1). The overall identification accuracy (total number of correct images/total

number of images) of the five marmosets was 99.2%.

Subsequently, face identification accuracy was examined in a family of three, that was

raised in the set-up home cage. The trained model was evaluated using data that were not used for

training. For the evaluation, 5,600 facial images (1,400 for each class), three classes including each

marmoset, and an unknown class were used. The unknown class was used to exclude inappropriate

images, such as unclear images, images of the back of the head and faces in profile, and images not

showing faces. The evaluation metrics for deep learning for individual identification were as follows:

specificity of 98.7%–99.7%, accuracy of 98.6%–99.2%, recall of 96.1%–98.8%, precision of 98.4%–

99.1%, and an F-measure of 97.3%–98.3% (Table 2). The overall percentage of correct identifications

was 98.0%.

Multiple animals' 3D tracking

To optimize the Lidar for laser measurement and video recording, a part of the cage made of metal

mesh was replaced with acrylic panels, following which suitable measurement results were obtained

(Fig. 1a). The laser beam distance needed to be sufficiently narrow and cover the entire cage; thus, the

Lidars were placed 1000 mm from the front of the cage (Fig. 1b). All 3D tracking and grooming

behavior detection tests were conducted using data from one family consisting of three marmosets

(I5072M/father (I5072M/f), I5894F/mother (I5894F/m), and I940M/juvenile (I940M/j)) in a cage.

The 3D coordinate information of marmosets was obtained using video tracking and Lidar

(Fig. 2a-f, and Mov. S1). While the spatial resolution of Lidar was not enough to detect each marmoset

and its behavior, it was sufficient to detect the body and tail. Therefore, the centroids of the point

clouds were used as animal locations for 3D trajectory.

A raw image taken by the upper left video camera captured the entire cage and the three marmosets

(Fig. 2a), and the data was acquired by Lidar simultaneously (Fig. 2b). The centroid of cluster by Lidar

was indicated in white points (Fig. 2b), and each point was given a sequential tracking ID number

during background processing. A new number was assigned each time when the trajectory was broken.

The 2D pixel locations of the marmosets were then detected using an object detection algorithm

(Yolov3[29]) in the video image (Fig. 2c). The same process was applied to the four cameras to determine

the locations of the marmosets for each camera. The 3D location of each marmoset was calculated

using pre-calibrated camera positions and the pixel coordinates of the marmosets (Fig. 2d). The

extracted face images of marmosets by Yolo were identified using the trained VGG19 model, and

individual IDs were assigned to each marmoset by facial recognition (Fig. 2e, f). These series of

processed video data are shown in Mov. S1. In some cases, the marmosets could not be recognized by

video tracking when they moved too fast or behind an object. In such cases, the trajectory was

interpolated by Lidar (Mov. S2, S3). Fig. 3 shows 5 min of tracking trajectories without face

identifications (Fig. 3a, b) and with face recognition (Fig. 3c). When Lidar alone or Lidar and video

tracking were performed, the tracking serial numbers changed frequently, and it was impossible to

continuously track the individual (Fig. 3a, b). However, by combining face recognition with Lidar and

video tracking, the trajectory of each animal was continuously detected, and marmoset IDs were

assigned (Fig. 3c).

Trajectory analysis using the 3D tracking system

Using the results of 3D trajectories from this 3D tracking system, we analyzed where and for how long

each animal stayed in the cage during the one-hour period of the analyses (Fig. 4). This data indicated

that I5072M/f stayed on the lower floor of the cage for 88.8% (23642/26626 frames) of the one hour,

whereas both I5894F/m and I940M/j stayed on the lower floor for less than 0.5% (128/25597 and

12/24234 frames) of the one hour. In contrast, they stayed on middle floor for half of the one hour and

actively moved in various areas in the cage except the lower floor rest of the time (Fig. 4a-c, Table 3,

Supplementary Fig. 1). Furthermore, the individual distances were analyzed using the location data

(Fig. 4d). During this time, the distance between I940M/j and I5894F/m was less than 0.5 m 25.9%

(5044/19501 frames) of the total time. However, a distance of less than 0.5 m between I940M/j-

I5072M/f and I5072M/f-I5894F/m was detected 1.0% (203/20409 frames) and 2.2% (475/21599

frames) of the time, respectively. Conversely, the distance between I940M/j-I5894F/m was more than

1 m 2.0% (391/19501 frames) of the time, whereas I940M/j-I5072M/f and I5072M/f-I5894F/m were

separated by more than 1 m distance 20.0% (4081/20409 frames) and 11.0% (2364/21599 frames) of

the time, respectively. In this family, I940M/j and I5894F/m spent more time together than I5072M/f.

The same analyses were performed for long-term analysis on a different day during the

daytime, 7:00 am–7:00 pm. After 6:00 pm, all marmosets moved to the bed and clustered together and

their faces were not observed; therefore, the data were excluded from the analysis. The detailed

analyses indicated that, among the family members, I940M/j stayed on the lower floor of the cage the

longest, unlike the results of the 1-hour analysis (Supplementary Fig. 2). The time spent on the lower

floor of the cage between 7:00 am and 6:00 pm for each individual was 8.6% for I5072M/f, 13.0% for

I5894F/m, and 23.5% for I940M/j. Throughout the day, I940M/j spent the most time on the middle

floor (27.7%), I5072M/f on the upper floor (29.7%), and I5894F/m on the upper floor (29.9%).

Furthermore, locations of each animal were analyzed hourly for 12 hours (Supplementary Figs. 2, 3).

I5072M/f also spent most of his time in the upper area in the morning (9:00 am; 54%), but gradually

changed his activity area to the middle area in the afternoon (1:00 pm; 51%) and the lower area in the

evening (4:00 pm; 45%). I5894F/m rarely stayed in the middle area but moved to the upper section in

the morning (9:00 am; 48%) and to the lower section in the afternoon (3:00 pm; 55%). I940M/j moved

most evenly throughout the day among the three in the cage. At night, all individuals moved to the

upper bunk to their beds (Supplementary Fig. 2).

The distance between the marmoset pairs was also analyzed throughout the day. The

results showed that I5072M/f-I5894F/m were closer than 0.5 m for 21.1% of the daytime. However,

I5072M/f-I940M/j and I5894F/m-I940M/j were closer than 0.5 m for 19.6% and 16.7%, respectively.

Additionally, the closest and farthest pairs per hour switched frequently throughout the day

(Supplementary Fig. 5).

Detection of grooming behavior

In this system, behaviors could also be detected using video data from the two upper cameras in front

of the cage (Supplemental Fig. 6a, b). Grooming behavior, one of the representative social behaviors

of marmosets, excluding self-grooming behavior was detected. Training data on grooming behavior

were obtained through observations of video data by skilled animal technicians. To increase the

accuracy of grooming behavior detection, situations in which marmosets were close to each other but

not grooming were rigorously classified by skilled animal technicians as non-grooming images and

eliminated from the supervised images. The model was trained by annotated data of the grooming

positions in each video frame. Excluding annotation time, 880 hours were spent on training. Creation

of a csv file after processing of four movie files took 8 min. Face identification took 20 msec per

process, and matching face identification to trajectory took 1 min per 10-min video.

The coordinates of the grooming location and individuals were detected by separately

using the 3D tracking system, and determining which individual performed the grooming. Grooming

behaviors were observed more often in the early morning (7:30 am–8:00 am), midday (12:30 pm–1:30

pm), and evening (3:30 pm–4:00 pm, 5:00 pm), and all marmosets performed grooming behaviors

equally (Fig. 5a). All marmosets groomed at fixed locations (Fig. 5b), namely the two upper areas of

the cage. These two locations were different from the sleeping area (Supplemental Fig. 3).

Accuracy verification of the 3D tracking system

To determine the accuracy of the 3D tracking system, marmosets were fitted with colored belly band

to identify each animal, and individual tracking was conducted for 60 min. The Mov. S4 shows a

portion of this experiment and confirms that the system could track the marmosets accurately even

when they crossed. Although the marmoset cage had a swinging hammock and balls as enrichment

(Mov. S5), these movements did not affect the tracking accuracy of the marmosets when using a

combination of video tracking and Lidar. During this experiment, the marmoset IDs switched to

different IDs nine times for more than 5 sec, but seven of the nine ID switching events were resolved

within 15 sec. The maximum time length for accurate individual tracking was 19 min. Incorrect

marmoset IDs of longer than 5 sec were exhibited over 4.6% of the total 60 min.

The accuracy of the 3D trajectory data analysis in detecting the positions of animals in

cages was confirmed by comparing it with visual observation data (Table 3, Supplemental Fig. 2). For

example, in the visual observations of an hour, the position of the animals was on the upper wooden

bed 9.4% of the time, on the middle floor 63.8%, on the lower wooden bed 25.1%, on the lower floor

0%, and in the other areas 1.7% of the time. Similarly, the system-analyzed data showed that the

position of the animals on the upper wooden bed was 16.5%, middle floor was 52.6%, lower wooden

bed was 29.1%, lower floor was 0.1%, and other areas was 1.7%. A strong significant correlation was

found between the visual observation and system data ($R^2 = 0.988$, $P < 0.01$; Fig. 6a).

The accuracy of grooming behavior detection by the machine was compared with that of

visual observation for one hour (Fig. 6b). Each 30-sec video was considered as one unit. Twenty-four

grooming units were visually detected, and 21 units were detected by the machine. Among these, 19

units were detected by visual observation and machine (recall: 79.1%). In the system, 19 of 21 units

were confirmed as actual grooming behaviors (precision: 90.5%). The five missed units were difficult

to detect because of the placement of animals behind an object: four had marmosets grooming behind

the feeder, and one had marmosets grooming in the blind spot of the bed. Conversely, two units were

detected by the machine but not by direct observation. During these events, the marmoset nuzzled its

face into the other's tail or placed its hand on the other's shoulder. These actions seemed to be part of

grooming behavior; however, observers did not judge them as grooming behaviors because they were

not typical grooming actions. In one hour, there were 10 units during which the distances between

marmosets were close, and false positives were detected in only one unit, with a duration of 30 sec

(recall: 90%, Fig. 6b). The system also worked well in higher interaction situations, such as grooming.

A grooming situation is shown in Mov. S6. The two animals were very close to each other and the

animal ID switched frequently, but both were involved in grooming. After grooming, accurate

individual identification was resumed as soon as the face was visible, as shown in Mov. S4.

Discussion

In this study, we constructed the FulMAI system, a new automatic 3D tracking system for multiple

marmosets under free-moving conditions, using a novel approach that combines Lidar, video tracking,

and deep learning face recognition. These are all existing technologies, but this is the first study

combining them for 3D tracking of multiple unlabeled nonhuman primates with relatively rapid

movements in free-moving conditions. The FulMAI system allowed us to analyze the natural behavior

of individuals in a family for a month. Lidar and Yolo, the object detection algorithm, were fast

programs that could be run constantly for a month with little delay. In addition, Yolo's light memory

usage and easy integration with other systems will allow us to obtain more long-term and diverse data

through well-planned operations in the future. Based on these findings, FulMAI is expected to record

the lifetime behavior of marmosets with a lifespan of approximately 15 years and match it with MRI

and other data to determine what changes occurred in the brain during life events and transformed

their behavior.

In preliminary studies, we used metal mesh doors, but these reflected the laser, and the

marmosets' location could not be detected. The cage doors were changed to acrylic panels to make

them suitable for the analysis (Fig. 1). The cages can be divided into four compartments. As shown in

Supplemental Fig. 1, the system accurately tracked each animal even though the cage was divided.

Therefore, the cage size can be changed to the extent that it fits within the camera's view. In contrast,

thick poles should be avoided in front of the cage because they increase blind space for both the camera

and the Lidar.

Although either laser or video data can be used as the primary data for this system, we used

video tracking as a base for detecting marmoset positions and then completed the missing information

using Lidar, thereby compensating for the lost marmoset trajectory detections by video and false

recognitions by Lidar. Owing to its detection speed and resolution, Lidar is suitable for application to

fast-moving objects. However, it falsely recognized objects that were similar in size to a marmoset,

such as a marmoset tail, feeder boxes, or colored balls. On the other hand, a hammock in the family

cage, which was included as an environmental enrichment and was not firmly fixed to the cage, was

not mis-detected by either Lidar or video because of the differences in size and shape from those of

the marmoset, and it did not affect the tracking of the marmosets. Video tracking does not have this

issue because it uses color image data to identify objects (Mov. S5). However, video tracking lost

detection when marmoset motion was quicker than the camera's shutter speed or when the marmosets

were behind obstacles (Mov. S2, S3). By compensating for the respective shortcomings of Lidar and

video tracking and using each to complement the trajectories of the other, a tracking system for small,

high-speed objects in three dimensions was realized. In addition, because both Lidar and video

tracking can be processed at high speed without a time lag, data is not accumulated for subsequent

analysis, which is favorable for long-term analysis. Indeed, the 3D trajectories and grooming behaviors

of three marmosets could be continuously analyzed in their home cage for over six months without

data accumulation. The animals were healthy throughout the experiment, proving the safety of our

system. A similar 3D tracking system using Kinect senser or cameras has been reported for various

animal species, including marmosets[1, 20, 23, 24, 30, 31]. However, none of the existing 3D tracking systems

have achieved such a long-term analysis, whereas FulMAI is the only behavioral analysis system for

longitudinal observation over each marmoset's lifetime in the family. One of the advantages of

FulMAI is the flexibility in terms of cage size and structure as long as acrylic panels are located in

front of the camera and Lidar, and it can be applied to a wide variety of animal species. This system

would also be useful for other animal species that move as quickly as marmosets in 3D, such as

tamarins, free-swimming fish, and quick-flying birds and bats.

Our system successfully tracked multiple marker-less marmosets for the first time, which is

another advantage of our system over existing systems. The color marking causes animal stress

because it requires capture or anesthesia, and periodic hair dyeing requires time and effort, which is

unsuitable for analyzing marmosets' behaviors in the long-term. The face identification of marmosets,

which we were able to achieve for the first time, can also be undertaken independently and can be

applied to other motion analysis systems, such as DeepBhvTracking[27].

When marmosets are kept as a family, they repeat the behavior of huddling and then

suddenly dispersing several times a day. By this characteristic behavior, individual traceability is lost

from the trajectory. Therefore, we combined our system with facial recognition, using deep learning

to detect and track each animal in a family group, since the most distinctive feature of the marmoset's

body is its face. Usually marmoset individual identification is performed using microchips, hair

clippings, hair dyes, or necklaces[32]. Microchips and necklaces are unsuitable for MRI analysis, hair

clippings and hair dyes require periodic maintenance, and necklaces pose the risk of tightening around

the neck as they grow or of getting caught. In one part of this study, the marmosets were fitted with

colored belly band for visual tracking, but I5894F/m seemed uncomfortable with the band and was

constantly fidgeting with it. Thus, artificial markers have the risk of causing unexpected behavioral

changes. Therefore, one of the goals of this study was to achieve marker-less identification of

individuals.

The marmoset family used in this study had no blood relationships with other family

members since the juveniles (I940M) were obtained by an embryo from another embryo donor pair

and was transferred into the female's uterus (I5894F; foster mother)[10, 33]. Regarding facial

identification, the most accurate results within the family were those for I940M/j (Table 2). Although

I940M/j was beyond sexual maturity, younger animals have easily identifiable distinctive facial

features. In addition, when using five animals of the same age (1–2 years old), the animals could be

individually identified; therefore, highly accurate classification is possible by facial recognition even

when the animals are of similar age (Table 1). Incorporating facial recognition results into the

trajectory enabled long-term tracking, even for multiple animals in the same cage (Fig. 3). We

examined the accuracy of marmoset tracking using this system by color labeling the marmosets and

comparing this to visual tracking (Mov. S4, Table 3). This experiment confirmed that it is possible to

track animals using facial recognition with 95% accuracy, without labeling marmosets or relying on

color information.

In addition, using the 3D coordinates data, this system could calculate where each animal

stayed in the cage, for how long it stayed (Fig. 4a-c, Table 3, Supplemental Figs. 2, 3, 4), and the

distance between individuals (Fig. 4d, Supplemental Fig. 5), which suggests interactions between

individuals. Although switching of the marmoset IDs was observed several times during the hour-long

experiment, results of visual observations and system analysis for where each marmoset existed in

terms of 3D coordinates in the cage, and how long each marmoset spent in that area were highly

correlated ($R^2 = 0.988$, Fig. 6a, Supplemental Fig. 2). This indicated that the slight ID switching did

not affect the experimental results, and our tracking system was sufficiently accurate for automatically

analyzing marmosets' 3D coordinates in the cage.

Marmosets often change their preferred position and the individuals that they spend time

with during the day. For example, the male marmoset (I5072M/f) preferred the upper area in the

morning, but in the afternoon, it preferred the lower bed. This result was not apparent in the 1-hour

analysis compared with observations. Furthermore, one-hour and 12-hour analyses of position in the

cage and distance between marmoset individuals showed different results. For example, a one-hour

analysis showed that I5072M/f preferred the cage's lower floor and remained there 80% of the analysis

time (Table 3). However, a 12-hour analysis indicated that I940M/j spent the longest time (23.5%) on

the lower floor (Supplemental Fig. 3, 4). The analyses of the distance between individuals showed that

I5894F/m-I940M/j were within 0.5 m for 25.9% of the one-hour analysis, but the 12-hour analysis

indicated that the I5072M/f-I5894F/m remained closer than I5894F/m-I940M/j or I5072M/f-

I5894F/m. These results suggest that the results based on human observation for a limited time and

those based on machine observation throughout the day may differ. Indeed, when we directly observed

this family, I5072M/f and I5894F/m were usually far apart, but a 12-hour analysis suggests that the

I5072M/f and I5894F/m spent time close together. Therefore, it should also be considered that people's

presence alters marmoset behavior. To accurately understand marmoset behavior, it is necessary for

the behavior analysis system to constantly analyze a marmoset's life with the same evaluation criteria.

Our 3D tracking system can continuously analyze animal positions and distances between individuals.

In addition, some marmoset facilities, to provide more space or different environments

for marmosets, occasionally connect adjacent marmosets' cages[34]. In this study, an extra cage was

connected to the home cage. When a marmoset went off-screen due to movement to the connected

cage, the tracking of the marmoset was temporarily interrupted. However, other marmosets continued

to be tracked, and no other individual was tracked instead of the missing marmoset (Mov. S7).

Tracking was resumed when the marmoset returned to the cage.

Grooming, a typical social behavior, was detected through deep learning using video

analysis. The detector was Yolo, the same one used for the 3D tracking. Comparing detection of the

grooming behavior using visual observation by humans and the system showed that grooming

behaviors were detected with high precision (90.5%) and recall (79.1%; Fig. 6b). In some cases, the

system identified behaviors just before or after the grooming as grooming behaviors. The deep learning

system may have detected certain behaviors that might be unique to periods before and after grooming.

However, in other cases, grooming behavior could not be detected by the deep learning system because

the groomers were located in the camera's blind spot. Therefore, the training dataset must be further

updated to detect behavior behind obstacles. Combining the 3D tracking with the grooming behavior

detection by deep learning, this system could reveal when and where each marmoset displayed

grooming behavior using accurate quantification, and the social relationships among the animals (Fig.

5 a, b). The 12-hour grooming behavior analysis in this study indicated that grooming behavior was

found to occur at specific locations: upper beds and floor, and at specific times: early morning, midday,

and evening. This finding was obtained exclusively using automatic behavior analysis. While there

are existing methods for automatically detecting the behavior of a single animal[35], for the first time,

our system was able to directly detect the behavior of multiple animals in captivity and identify the

individuals and their locations. We aim to identify a variety of behaviors for analyzing lifelong

behavior changes. To find unpredictable unknown behavior, clustering by unsupervised learning

would be necessary. For unsupervised learning, it is necessary to analyze the videos of each individual

or to create a skeletal model of each marmoset and analyze the movement of each key point. Our study

is expected to further develop in the future because of its ability to detect accurate 3D location linked

to each individual. For more flexible behavior extraction, this system was made of Robot operating

system (ROS) and is extensible to incorporate other Python-based projects, such as DeepLabCut,

DeepEthogram[36], and Tweetynet[37]. Because all animal activities are recorded in the FulMAI, each

animal behavior in this home cage can be analyzed retrospectively in detail. By comparing with other

data, such as imaging, blood biochemistry, or biomarker data, it is also possible to analyze the behavior

of model animals from multiple perspectives without overlooking changes in behavior associated with

the life events of brain function changes or the onset of disease.

Moreover, Lidar can track the animal locations day and night; therefore, their nocturnal

behavior can be observed to obtain primary data. Our preliminary data showed that the combination

with a Lidar and starlight camera can capture details of nighttime behavior in a dark room

(Supplemental Mov. S8). Sleep disorders have recently been suggested as predictors of Alzheimer's

disease and Parkinson's disease[38, 39]. This system would be able to accurately quantify the behavior of

marmosets throughout their lives to discover behavioral changes before and after the onset of diseases,

even at an extremely early stage.

Recently Calapai et al. reported a new touch screen-based cognitive auditory analysis

system that allows marmosets to perform tasks in a free moving environment in a paired home cage[40].

By incorporating such a freely operable cognitive function task device into our system, it is possible

to provide a behavior analysis system that comprehensively and continuously analyzes the marmoset's

natural behavior, activity, and cognitive functions without inducing stress.

In conclusion, the combination of Lidar, video tracking, and facial recognition enabled

successful simultaneous tracking of each marmoset in a family without artificial labeling. Using data

from the 3D tracking system we developed, we could measure 3D coordinates, movement trajectories,

the quantity of activity, position in the cage, when, and for how long each animal stayed, distances

between individuals, and time spent in grooming behavior. These measurements will allow automated

analysis of activity and behavior under free-moving conditions throughout the marmoset's lifetime,

enabling quantitative analysis of "when, where, and what" of each animal and capturing changes in

brain function, including development, growth, aging, and disease.

Materials and Methods

Hardware Setup

The home cage was made of acrylic panels on the front and back, and all food dishes, water jugs, and

bedding inside the cage were fixed to a wire mesh on the side to prevent them from moving. This cage

was designed to conduct behavioral analysis in a marmoset's home cage (Fig. 1a). Four poles were set

up in front of the cage at 1000 mm, and four Lidars (VLP-16, Velodyne Lidar, San Jose, CA, USA)

and four cameras (C920, Logitech, Lausanne, Switzerland) were fixed to the poles (Fig. 1b). The Lidar

was controlled by motion measurement software, the path traced, and velocity of the marmosets

recorded (Hitachi, Ltd, Tokyo, Japan.). The lower Lidar was set at 0.9 m height and the upper Lidar

at 1.9 m height. The distance between the left and right Lidar devices was 0.3 m.

The laser beam of the Lidar was irradiated radially and adjusted to cover the entire interior

area of the cage. The recording parameters of the cameras were 1080p and 60 Hz, and all cameras

were adjusted to show the entire cage (Supplemental Fig. 6). The positions of the cage, Lidar, and

cameras were calibrated when the cage was moved, and the interiors were changed. Lidar calibration

was used to acquire background information, including the cage interior without marmosets.

Calibration of the camera was performed by detecting alco-markers at the four corners of the cage.

The specifications of the PC used to operate and record the camera and Lidar were corei7, RTX1060,

32GB. PC specifications used for analysis were corei7, RTX3090, 64 GB.

Ethical approval

This study was conducted at the animal facility of the Central Institute for Experimental Animals

(CIEA), Kawasaki, Japan. The animal experimental protocol was reviewed by the Institutional Animal

Care and Use Committee and approved (approval no. 17051) according to the Regulations for Animal

Experiments in CIEA, based on the Guidelines for the Proper Conduct of Animal Experiments by the

Science Council of Japan (2006).

Animal facilities

The marmosets were kept in a family cage with dimensions of 820 × 610 × 1578 mm. Although there

was only one marmoset family in the room, they were able to vocally communicate with the marmosets

in other rooms. All cages were equipped with a bunk, and the animal rooms were adjusted to 26–28°C

temperature, and 40–60% humidity with a 12:12 h light/dark cycle. The animals tested negative for

Salmonella spp., *Shigella* spp., and *Yersinia pseudotuberculosis* on annual fecal examination. The

study animals were inspected nine months before the study.

Animals

Each marmoset family was housed in a home cage. The observed family consisted of a male (I5072M/f,

ten years old), female (I5894F/m, six years old), and a juvenile (I940M/j, one year old). No artificial

or non-artificial physical features were used for individual identification, such as hair dye, necklaces,

piercings, or other visible signs, except during 3D tracking of marmosets during the color marker

experiments. In conventional animal care, a microchip is implanted subcutaneously for individual

identification. The individual I940M/j was born by embryo transfer to the mother's uterus[10], and there

was no blood relationship within the third degree of the family.

Five healthy marmosets approximately 1–2 years old, one male and four females, were

used in the preliminary study to develop deep learning facial recognition (I774 one year old, I6695

one year old, I6708 one year old, I6677 one year old, and I893 two years old). There were no artificial

or visible physical characteristics, such as hair dye, necklaces, or piercings, and each marmoset was

not related to any other marmoset within the third degree of kinship.

System workflow

The overview of this system is shown in Supplemental Fig. 7. The videos and Lidar data were

processed on the same PC, and acquisition times were perfectly synchronized for at least one month.

Yolo, an object recognition algorithm, detected the marmoset's face, body, actions, and environmental

enrichments (ball and hammock) in each video frame. Body coordinates were combined with

coordinates using Yolo and Lidar to calculate a short 3D trajectory. Furthermore, the cropped face

images were identified as individuals using a convolutional neural network, and individual IDs were

applied to the short-term 3D trajectories to achieve long-term 3D tracking of each individual.

Behaviors were detected simultaneously with the same model of Yolo, and each animal's behavior

was linked to the nearest marmoset ID.

Facial recognition

The cage (W820 × D610 × H1578 mm) was divided into four compartments to isolate each marmoset

for obtaining training images by video camera. Each animal was housed in a separate cage, and videos

were recorded from the cage front for 3–6 hours. The front door of the cage was then replaced with

clear acrylic panels. From these video frames, the face images were cropped only in the face area for

the training images, and these data were augmented by rotation. Using 1400 facial images in each

class for training, frontal images of the face were classified into four or five classes using the VGG19

model network. In the preliminary test, the five classes comprised five marmosets. For the family

group, the four classes included those members and an unknown class to exclude images inappropriate

for classification, and 1000 images were used for training.

Model training was performed by fine-tuning a pre-trained VGG19 model (learning rate

0.001, fixed number of layers 12, BATCH_SIZE 32)[41] (Supplementary Data S2). VGG19 is a network

model proposed by the Visual Geometry Group at the University of Oxford in 2014 that obtained

accurate classification performance on the ImageNet dataset. The pre-trained models used in this study

were trained using the ImageNet dataset.

3D tracking

Lidar and video tracking performed 3D tracking. A Lidar measures the distance to an object by

irradiating a laser and analyzing the reflected laser. Marmoset detection using Lidar involves the

following steps: background information removal, clustering, and selection of marmoset clusters[42].

Background removal was performed during calibration. The observed points were moving points

belonging to objects whose spatial positions changed and static points from the background

environment. Marmosets were then selected from among all the moving clusters. The Euclidean

cluster extraction algorithm was used for clustering, with a minimum cluster size of S1 and maximum

cluster size of S2 (S1: 100 mm, S2: 500 mm)[43]. The centroid of the cluster is regarded as the position

of the marmoset for tracking. Three-dimensional video tracking was performed using four time-

synchronized video cameras. The positions of the whole body and head of the marmoset were detected

in each video frame using Yolo, an object detection algorithm, and the 3D position was calculated

from each camera information[29]. The Yolo model was trained using 1164 images in which the whole

body and head area were annotated (learning rate: 0.0003, momentum: 0.9, batch: 64). For tracking,

the primary information used was the 3D video tracking coordinates, and the information was

supplemented with Lidar coordinates when video coordinates were missing. When the primary video

information was lost, the system searched for available Lidar coordinates close to the coordinates just

before the location was lost and switched to Lidar tracking. When Lidar coordinates were available,

continued tracking was undertaken using only Lidar. When the camera re-detected the coordinates,

the tracking was switched back to the camera again. The switching was set to occur when the

coordinate relationship between the camera and Lidar coordinates was fixed.

The program for linkage between video tracking and Lidar for individual information was

produced by Hitachi, Ltd.   When the marmoset coordinates were detected outside the cage, they were

considered as missing values.

Detecting grooming behavior

The software Yolo was used to detect grooming behavior. A total of 252 annotated image files in which

grooming marmosets were surrounded by a minimum rectangle were prepared as training images, and

the model was trained using the dataset. Grooming images were strictly categorized by skilled animal

technicians to exclude situations in which marmosets did not groom. The trained model analyzed the

top two camera view images and detected grooming behavior in each video frame. To improve the

accuracy by removing noise, a Gaussian-weighted moving average of 50 sec window and 5 sec slide

was applied to the frequency of grooming occurrence each time to create the probability distribution

of grooming occurrence. The same process was performed for the left and right cameras and the

calculated probability distributions were averaged. When the average value exceeded a certain

threshold, grooming was considered to have occurred (Supplementary Data S2). To test the accuracy

of this detection system, skilled breeding staff visually detected marmoset grooming in one hour of

video that was not used for training data. This visual result was used as the true label, and the accuracy

and repeatability of grooming detection were calculated in 30 sec increments of time. Because

grooming was analyzed with two cameras, it was possible to link location and time. Grooming

locations were calculated using the same method as used for video tracking. The grooming time was

based on the video file.

Statistical analysis

Pearson's correlation coefficient was calculated using GraphPad Prism version 9 (GraphPad Software,

La Jolla, CA, USA).

Acknowledgments

This research was partially supported by the Brain Mapping by Integrated Neurotechnologies for

Disease Studies (Brain/MINDS), "Study of developing neurodegenerative model marmosets and

establishment of novel reproductive methodology (JP19dm0207065)" from the Japan Agency for

Medical Research and Development (AMED) and Grant-in-Aid for Scientific Research (A)

"Development of an Automated Behavioral Analysis System for Evaluation of Genetically Modified

Disease Model Marmosets" JSPS KAKENHI Grant Numbers JP21H04756 to ES and Grant-in-Aid

for Early-Career Scientists "Effects of transient alcohol exposure to the fetus in early pregnancy on

brain development." JSPS KAKENHI Grant Number JP20K16908 to TY.

Author contributions

T.Y. and E.S. conceived and designed the study. T.Y., N.G., J.K., and T.F. performed the setup of the

measurement system. R.K., R.H., M.K., Y.H., and E.Y. performed the annotation of the training data.

T.Y., Y.S., R.H., J.K., and T.F. processed the data, trained the models, and analyzed the results. T.Y.

and R.K. performed the experiment. T.Y., W.K., K.S., N.G., T.I., and E.S. discussed and interpreted

the results. T.Y., T.I., and E.S. wrote the manuscript. E.S. supervised the project. All authors

commented on the manuscript.

Competing interests

NG is an employee of Hitachi, Ltd. JK is an employee of Hitachi Solutions Technology Ltd. TF is an

employee of Totec Amenity Limited. The remaining authors declare no competing interests.

Resources Availability

Information and requests for the program for linkage between video tracking and Lidar should be

directed to and will be fulfilled by Norio Goda (norio.goda.ws@hitachi.com). Requests for other

resources and information should be addressed to Erika Sasaki (esasaki@cieea.or.jp).

Data and code availability

The readily tested code and data has been presented in the Supplemental information (Supplementary

Data S1, S2).

Figure legends

Figure 1. Installation of hardware for the behavior analysis system

(a) A home cage with acrylic panels on the front and back and metal mesh on the sides. (b) Schematic

diagram of light and ranging (Lidar) systems and cameras installed in front of the cage. Lidars and

cameras were installed 1000 mm in front of the cage.

Figure 2. 3D tracking of marmosets using Lidar and video tracking

(a) Raw video frame. (b) Detected 3D coordinates of marmosets by Lidar. White point: centroid of

marmosets; white box: virtual space of the cage. (c) Detected location of marmosets in video frame

by Yolo. Rectangle: detected marmosets. (d) Calculated 3D coordinates of marmosets by video

tracking, colored point: coordinate of marmoset. (e, f) Individual information added to c and d using

face identification, green: I5072M/f, red: I5894F/m, yellow: I940M/j.

Figure 3. 3D position trajectory of a marmoset over a period of five minutes

(a) Marmoset trajectories analyzed by Lidar. White points indicate centroid of marmosets. (b) Lidar

and video tracking. Different colors were assigned to each tracking serial number that numbered the

background. (c) Lidar and video trajectory tracking combined with face recognition. Green: I5072M/f;

red: I5894F/m; yellow: I940M/j.

Figure 4. Place preference of marmosets over a period of 1 hour

Color map of the time spent by marmosets in the cage for one hour (a-c). Locations where marmosets

spent more than 1% of one hour are shown in red, and others are shown as color bars. (a) I5072M/f,

(b) I5894F/m, (c) I940M/j. (d) Histogram of the distance between each family member. Blue bars

indicate the individual distances between I940M/j and I5072M/f, orange bars indicate the individual

distances between I940M/j and I5894F/m, and green bars indicate the individual distances between

I5072M/f and I5894F/m.

Figure 5. Detection of grooming behavior for a day

(a) Number of grooming behaviors detected per day. Bars show the number of frames detected each 5

min. (b) The 3D coordinates where the grooming behavior of each individual was detected are shown.

Red indicates I940M/j, green indicates I5072M/f, and blue indicates I5894F/m.

Figure 6. Comparison of human visual observations and the system

(a) The time the marmosets stayed in a particular location was compared between visual observations

and those by the system. The marmosets stayed in four locations: upper bunk, middle floor, lower

bunk, and lower floor. Dotted lines indicate regression lines. (b) Each cell represented 30 s. Blue cells:

Visual observation detected grooming at least once in 30 s. Yellow cells: Visual observation detected

the close distance between two marmosets. Red cells: The system detected grooming by the intensity

exceeding a threshold value.

References

1.	Bala, P. C. *et al.* Automated markerless pose estimation in freely moving macaques with

OpenMonkeyStudio. *Nat. Commun.* **11**, 4560 (2020) (DOI 10.1038/s41467-020-18441-5)

2.	Yamazaki, Y. *et al.* Tool-use learning by common marmosets (Callithrix jacchus). *Exp. Brain*

Res. **213**, 63–71 (2011) (DOI 10.1007/s00221-011-2778-9)

3.	Miller, C. T. *et al.* Marmosets: A neuroscientific model of human social behavior. *Neuron*

90, 219–233 (2016) (DOI 10.1016/j.neuron.2016.03.018)

4.	Mitchell, J. F. & Leopold, D. A. The marmoset monkey as a model for visual neuroscience.

Neurosci. Res. **93**, 20–46 (2015) (DOI 10.1016/j.neures.2015.01.008)

5.	Roelfsema, P. R. & Treue, S. Basic neuroscience research with nonhuman primates: a small

but indispensable component of biomedical research. *Neuron* **82**, 1200–1204 (2014) (DOI

10.1016/j.neuron.2014.06.003)

6.   Burkart, J. M., Fehr, E., Efferson, C. & van Schaik, C. P. Other-regarding preferences in a

non-human primate: common marmosets provision food altruistically. *Proc. Natl Acad. Sci. U. S. A.*

104, 19762–19766 (2007) (DOI 10.1073/pnas.0710310104)

7.   Abbott, D. H., Barnett, D. K., Colman, R. J., Yamamoto, M. E. & Schultz-Darken, N. J.

Aspects of common marmoset basic biology and life history important for biomedical research. *Comp.*

Med. **53**, 339–350 (2003)

8.   Sasaki, E., *et al*. Generation of transgenic non-human primates with germline transmission.

Nature **459**, 523–527 (2009) (DOI 10.1038/nature08090)

9.   Sato, K., *et al*. Generation of a nonhuman primate model of severe combined

immunodeficiency using highly efficient genome editing. *Cell Stem Cell* **19**, 127–138 (2016) (DOI

10.1016/j.stem.2016.06.003)

10.   Sato, K., *et al*. A nonhuman primate model of familial Alzheimer's disease. *bioRxiv*,

2020.2008.2024.264259 (2020)

11.   Kishi, N., Sato, K., Sasaki, E. & Okano, H. Common marmoset as a new model animal for

neuroscience research and genome editing technology. *Dev. Growth Differ.* **56**, 53–62 (2014) (DOI

10.1111/dgd.12109)

12.   Okano, H., Miyawaki, A. & Kasai, Kasai K.. Brain/MINDS: brain-mapping project in Japan.

Philos. Trans. R. Soc. Lond. B Biol. Sci. **370** (2015) (DOI 10.1098/rstb.2014.0310)

13. network, B. I. C. C. A multimodal cell census and atlas of the mammalian primary motor

cortex. *Nature* **598**, 86–102 (2021) (DOI 10.1038/s41586-021-03950-0)

14. Hong, W., *et al*. Automated measurement of mouse social behaviors using depth sensing,

video tracking, and machine learning. *Proc. Natl Acad. Sci. U. S. A.* **112**, E5351–E5360 (2015) (DOI

10.1073/pnas.1515982112)

15. Redfern, W. S., *et al*. Automated recording of home cage activity and temperature of

individual rats housed in social groups: the Rodent Big Brother project. *PLOS ONE* **12**, e0181068

(2017) (DOI 10.1371/journal.pone.0181068)

16. Sutoko, S., *et al.* Early identification of Alzheimer's disease in mouse models: application

of deep neural network algorithm to cognitive behavioral parameters. *iScience* **24**, 102198 (2021)

(DOI 10.1016/j.isci.2021.102198)

17. Hsu, A. I. & Yttri, E. A. B-SOiD, an open-source unsupervised algorithm for identification

and fast prediction of behaviors. *Nat. Commun.* **12**, 5188 (2021) (DOI 10.1038/s41467-021-25420-x)

18. Karashchuk, P., Tuthill, J. C. & Brunton, B. W. The DANNCE of the rats: a new toolkit for

3D tracking of animal behavior. *Nat. Methods* **18**, 460–462 (2021) (DOI 10.1038/s41592-021-01110-

w)

19. Zimmermann, C., Schneider, A., Alyahyay, M., Brox, T. & Diester, I. FreiPose: A deep

learning framework for precise animal motion capture in 3D spaces. *bioRxiv*, 2020.2002.2027.967620

(2020)

20. Yabumoto, T., *et al.* MarmoDetector: A novel 3D automated system for the quantitative

assessment of marmoset behavior. *J. Neurosci. Methods* **322**, 23–33 (2019) (DOI

10.1016/j.jneumeth.2019.03.016)

21. Pereira, T. D., *et al.* SLEAP: A deep learning system for multi-animal pose tracking. *Nat.*

Methods **19**, 486–495 (2022) (DOI 10.1038/s41592-022-01426-1)

22. Segalin, C., *et al*. The Mouse Action Recognition System (MARS) software pipeline for

automated analysis of social behaviors in mice. *eLife* **10** (2021) (DOI 10.7554/eLife.63720)

23. Lauer, J., *et al*. Multi-animal pose estimation, identification and tracking with DeepLabCut.

Nat. Methods **19**, 496–504 (2022) (DOI 10.1038/s41592-022-01443-0)

24. Mathis, A., *et al.* DeepLabCut: markerless pose estimation of user-defined body parts with

deep learning. *Nat. Neurosci.* **21**, 1281–1289 (2018) (DOI 10.1038/s41593-018-0209-y)

25. de Chaumont, F., *et al*. Computerized video analysis of social interactions in mice. *Nat.*

Methods **9**, 410–417 (2012) (DOI 10.1038/nmeth.1924)

26. Wiltschko, A. B., *et al*. Revealing the structure of pharmacobehavioral space through motion

sequencing. *Nat. Neurosci.* **23**, 1433–1443 (2020) (DOI 10.1038/s41593-020-00706-3)

27. Sun, G., *et al.* DeepBhvTracking: A novel behavior tracking method for laboratory animals

based on deep learning. *Front. Behav. Neurosci.* **15**, 750894 (2021) (DOI 10.3389/fnbeh.2021.750894)

28.   Schultz-Darken, N., Ace, L. & Chapter, A. H. 7. Behavior and behavioral management in

The Common Marmoset in Captivity and Biomedical Research (eds Marini, R., Wachtman, L., Tardif,

S., Mansfield, K. & Fox, J.) (Academic Press, 2019)

29.   Redmon, J. & Yolov, F. A. An incremental improvement. *arXiv Preprint ArXiv:180402767*,

(2018)

30.   Matsumoto, J., *et al.* A 3D-video-based computerized analysis of social and sexual

interactions in rats. *PLOS ONE* **8**, e78460 (2013) (DOI 10.1371/journal.pone.0078460)

31.   Lyons, D. M., MacDonall, J. S. & Cunningham, K. M. A Kinect-based system for automatic

recording of some pigeon behaviors. *Behav. Res. Methods* **47**, 1044–1054 (2015) (DOI

10.3758/s13428-014-0531-6)

32.   Susanne, R. & Ann-Kathrin, O. Husbandry and management of New World species:

marmosets and tamarins. *The Lab. Primate*, 145–162 (2005)

33.   Kurotaki, Y. & Sasaki, E. Practical reproductive techniques for the common marmoset. *J.*

Mamm. Ova Res. **34**, 3–12 (2017) (DOI 10.1274/032.034.0103)

34.   Pines, M. K., Kaplan, G. & Rogers, L. J. A note on indoor and outdoor housing preferences

of common marmosets (Callithrix jacchus). *Appl. Anim. Behav. Sci.* **108**, 348–353 (2007) (DOI

10.1016/j.applanim.2006.12.001)

35.   Kobayashi, K. *et al.* Automated detection of mouse scratching behaviour using

convolutional recurrent neural network. *Sci. Rep.* **11**, 658 (2021) (DOI 10.1038/s41598-020-79965-

w) (Pubmed:33436724)

36.   Bohnslav, J. P., *et al.* DeepEthogram, a machine learning pipeline for supervised behavior

classification from raw pixels. *eLife* **10** (2021) (DOI 10.7554/eLife.63377)

37.   Cohen, Y. *et al.* Automated annotation of birdsong with a neural network that segments

spectrograms. *eLife* **11** (2022) (DOI 10.7554/eLife.63853)

38.   Borges, C. R., Poyares, D., Piovezan, R., Nitrini, R. & Brucki, S. Alzheimer's disease and

sleep disturbances: a review. *Arq. Neuro Psiquiatr.* **77**, 815–824 (2019) (DOI 10.1590/0004-

282X20190149)

39.   dos Santos, A. B., Kohlmeier, K. A. & Barreto, G. E. Are sleep disturbances preclinical

markers of Parkinson's disease? *Neurochem. Res.* **40**, 421–427 (2015) (DOI 10.1007/s11064-014-

1488-7)

40.   Calapai, A., Cabrera-Moreno, J., Moser, T. & Jeschke, M. Flexible auditory training,

psychophysics, and enrichment of common marmosets with an automated, touchscreen-based system.

Nat. Commun. **13**, 1648 (2022) (DOI 10.1038/s41467-022-29185-9)

41.   Zhou, J. *et al.* with Transposed Convolution for Rotating Machinery Fault Diagnosis Based

on Deep Transfer Learning. *Shock Vib.* **2020**, 1–12 (2020)

42. Zhang, J., Xiao, W., Coifman, B. & Mills, J. P. Vehicle tracking and speed estimation from

roadside lidar. *IEEE J. Sel. Top. Appl. Earth Obs. Remote Sensing* **13**, 5597–5608 (2020)

43. Rusu, R. B. Semantic 3D object maps for everyday manipulation in human living

environments. *KI Künstliche Intell.* **24**, 345–348 (2010) (DOI 10.1007/s13218-010-0059-6)

Fig. 1

a

b

Fig. 2

a

c

e

b

d

f

Fig. 3

Fig. 4

a

(m)

Male (I5072M)

b

(m)

Female (I5894F)

c

(m)

Juvenile (I940M)

d

I940M-I5072M
I940M-I5894F
I5072M-I5894F

(m)

Fig. 5

a

b

Fig. 6

a

Human vs. System

R² = 0.9883
P < 0.01

b

Table 1. Classification accuracy of face identification in five marmosets aged 1 to 2 years.

	Animal ID				
	I774	I6695	I6708	I6677	I893
Specificity	100.0%	99.8%	99.8%	99.8%	99.8%
Accuracy	99.8%	99.6%	99.4%	99.8%	99.8%
Recall	99.0%	99.0%	98.0%	100.0%	100.0%
Precision	100.0%	99.0%	99.0%	99.0%	99.0%
F-Measure	99.5%	99.0%	98.5%	99.5%	99.5%

Table 2. Classification accuracy of face identification in a marmoset family.

	Male (I5072M)	Female (I5894F)	Juvenile (I940M)	Unknown
Specificity	98.7%	99.5%	99.7%	99.5%
Accuracy	98.7%	98.6%	99.2%	99.5%
Recall	98.8%	96.1%	97.6%	99.6%
Precision	96.1%	98.4%	99.1%	98.5%
F-Measure	97.4%	97.3%	98.3%	99.0%

Table 3. Comparison of visual observations and the system analysis of time spent in specific locations in the cage for an hour

	Male (I5072M)		Female (I5894F)		Juvenile (I940M)	
	Human	System	Human	System	Human	System
Upper bed	0.4%	0.6%	10.7%	16.5%	9.4%	16.5%
Middle floor	0.3%	0.6%	38.8%	52.6%	63.8%	52.6%
Lower bed	8.8%	8.9%	41.8%	29.1%	25.1%	29.1%
Lower floor	90.3%	88.8%	0.5%	0.5%	0.0%	0.1%
Other	0.2%	1.11%	8.2%	8.2%	1.7%	1.7%