1
2
3
2-deoxyglucose transiently inhibits yeast AMPK signaling and triggers glucose transporter endocytosis, potentiating the drug toxicity

Clotilde Laussel[#,1], Véronique Albanèse[#,1], Francisco Javier García-Rodríguez[&,1], Alberto Ballin[&,1], Quentin Defenouillère[1], Sébastien Léon[1,2]

Affiliations:

[#]: Equal contributions
[&]: Equal contributions
[1]: Université Paris Cité, CNRS, Institut Jacques Monod, F-75013, Paris, France.

[2]: Author for correspondence:

    Sébastien Léon, PhD
    Institut Jacques Monod
    15 Rue Hélène Brion
    75205 Paris cedex 13, France.
    email: sebastien.leon@ijm.fr ; tel. +33 (0)1 57 27 80 57.

Running title:

Effects of 2-deoxyglucose on yeast AMPK signaling and endocytosis, and contribution to 2DG tolerance

Abstract

2-deoxyglucose is a glucose analog that impacts many aspects of cellular physiology. After its uptake and its phosphorylation into 2-deoxyglucose-6-phosphate (2DG6P), it interferes with several metabolic pathways including glycolysis and protein N-glycosylation. Despite this systemic effect, resistance can arise through strategies that are only partially understood. In yeast, 2DG resistance is often associated with mutations causing increased activity of the yeast 5'-AMP activated protein kinase (AMPK), Snf1. Here we focus on the contribution of a Snf1 substrate in 2DG resistance, namely the alpha-arrestin Rod1 involved in nutrient transporter endocytosis. We report that 2DG triggers the endocytosis of many plasma membrane proteins, mostly in a Rod1-dependent manner. Rod1 participates in 2DG-induced endocytosis because 2DG, following its phosphorylation by hexokinase Hxk2, triggers changes in Rod1 post-translational modifications and promotes its function in endocytosis. Mechanistically, this is explained by a transient, 2DG-induced inactivation of Snf1/AMPK by protein phosphatase 1 (PP1). We show that 2DG-induced endocytosis is detrimental to cells, and the lack of Rod1 counteracts this process by stabilizing glucose transporters at the plasma membrane. This facilitates glucose uptake, which may help override the metabolic blockade caused by 2DG, and 2DG export - thus terminating the process of 2DG detoxification. Altogether, these results shed a new light on the regulation of AMPK signaling in yeast and highlight a remarkable strategy to bypass 2DG toxicity involving glucose transporter regulation.

Author summary

In this article, we study the resistance to a drug named 2-deoxyglucose (2DG). 2DG is an efficient inhibitor of several metabolic pathways, particularly glycolysis which is of prime importance for tumor cell proliferation. Cancer cells but also cells from other organisms can develop various resistance strategies that only being unraveled. Here, using baker's yeast as a model organism, we document the resistance mechanism of a mutant lacking the arrestin gene *ROD1*, known to regulate nutrient transporter endocytosis. First, we show that 2DG triggers the endocytosis of many nutrient transporters. Mechanistically, 2DG leads to a transient inhibition of the AMPK signaling pathway, which in turn promotes the activation of Rod1, a pre-requisite for 2DG-induced endocytosis. Consequently, cells lacking Rod1 maintain various proteins at the plasma membrane including hexose transporters, the latter being essential for 2DG resistance. Hexose transporter stabilization at the cell surface may sustain glucose import and help cells cope with the 2DG-induced metabolic blockade, while also allowing 2DG detoxication.

Introduction

2DG is a glucose analog which lacks a hydroxyl group in position 2, and mainly acts as a competitive

inhibitor of glucose metabolism [1]. As such, it competes with glucose for entry into cells and for its

subsequent phosphorylation by hexokinase, but is almost not further metabolized. This leads to the

intracellular accumulation of 2DG-6-phosphate, a toxic metabolite that inhibits glycolysis and thus leads to a

rapid depletion of ATP stores, and the activation of nutrient starvation-activated pathways such as 5'-AMP

activated protein kinase, AMPK [2]. Moreover, since mannose is the C2 epimer of glucose, 2-deoxyglucose is

also 2-deoxymannose and thus also interferes with mannose metabolism, notably inhibiting protein N-

glycosylation [reviewed in 3]. This causes a series of cellular effects, and triggers the onset of the Unfolded

Protein Response pathway to face the stress encountered by a defective glycosylation of proteins at the

endoplasmic reticulum, which is reportedly the main toxic effect of 2DG in some conditions [4].

The realization that many cancer cells display an aberrantly high glucose uptake and metabolism has led

to the idea that glycolysis inhibitors could be used to preferentially target and kill tumor cells [5]. This also

allowed the use of radiolabeled glucose derivatives, such as 18-fluoro-2-deoxyglucose, to preferentially label

tumors for cancer imaging (PET-scans) [6]. So far, trials using 2DG as the sole chemotherapeutic agent have

failed because of the relative toxicity of this molecule, however strategies using 2DG or other 2-deoxy-

substituted versions of glucose in combination with other drugs are under study and are promising [6].

2DG has also been extensively used to understand glucose signaling in micro-organisms such as the

baker's yeast *Saccharomyces cerevisiae*, where it has been instrumental to delineate the molecular

mechanisms by which glucose regulates gene expression [reviewed in 3]. The metabolism of *S. cerevisiae* is

biased towards the preferential use of glucose as a carbon source. At the molecular level, glucose represses

the expression of many genes involved in respiratory metabolism or in the use of other carbon sources,

thereby favoring the preferential use of glucose by fermentation [7]. Among the actors in charge of adapting

the transcriptional program with respect to glucose availability is the yeast orthologue of AMPK, Snf1 [7, 8].

Snf1 is the catalytic subunit of a heterotrimeric complex which is active in the absence of glucose in the

medium. A canonical example of a Snf1-regulated transcription is that of the invertase *SUC2*, required for

sucrose hydrolysis and metabolism. Transferring yeast cells to sucrose medium activates Snf1, causing the

phosphorylation of the transcriptional repressor Mig1, its translocation out of the nucleus and the

derepression of *SUC2* [9, 10].

Snf1, like other kinases of the AMPK family, is activated by phosphorylation of a residue in its activation

loop [11]. This involves one of 3 upstream kinases whose activity is, surprisingly, not regulated by glucose.

Rather, evidence suggests that Snf1 activity is regulated by its dephosphorylation by the Protein Phosphatase

1 (PP1) complex made of the catalytic subunit, Glc7, and the glucose-specific regulatory subunit Reg1 [12].

How PP1 activity is linked to glucose availability is not yet clear, but deletion of *REG1* leads to constitutive

Snf1 activity and to the expression of glucose-repressed genes even in the presence of glucose [13].

Another phenotype displayed by the *reg1Δ* strain is its ability to grow robustly in presence of 2DG [3, 14,

15]. This is attributed to the constitutive activation of Snf1 in this mutant, as mutations in *SNF1* cause

hypersensitivity to 2DG and yeast lacking both Reg1 and Snf1 are also hypersensitive [16].

Despite the multifaceted cellular effects of 2DG, cells can indeed overcome 2DG toxicity by mechanisms

that are currently being unraveled [reviewed in 3, 15]. A prolonged exposure of HeLa cells in 2DG-containing

medium gave rise to resistant clones [17]. These cells displayed a high phosphatase activity towards 2DG-6-

phosphate (2DG-6-P), suggesting the presence of an enzyme that could detoxify this metabolite back into

2DG and thus prevent the metabolic blockade. Studies in yeast also identified phosphatases, named Dog1

and Dog2, with a similar activity *in vitro* and whose overexpression allows growth in 2DG-containing medium

[18, 19]. Recently, we found that the most abundant isoform, Dog2, is itself a glucose-repressed gene and

thus is regulated by Snf1/AMPK function [20]. This partially explains why strains with high Snf1/AMPK activity

are resistant to 2DG, such as *reg1Δ* [14, 16], other *reg1* mutant alleles [20, 21], a viable *glc7* mutation (Q48K)

[21] or gain-of-function mutations in components of the AMPK complex [16, 21, 22]. However, additional

resistance mechanisms independent of *DOG2* expression are at stake in these strains [20, 21]. This is

suggested by the observation that *DOG2* deletion does not completely re-sensitize *reg1Δ* mutants to 2DG,

and a point mutation in *REG1* (*reg1-P231L*) confers 2DG resistance but does not impact on the glucose-

mediated repression of genes [21]. Thus, Snf1 likely controls additional factors participating in 2DG

resistance.

One such candidate is the arrestin-related protein, Rod1. Arrestin-related proteins (ARTs) are important

regulators of plasma membrane protein endocytosis in response to extracellular cues [23, 24]. ARTs recruit

the ubiquitin ligase Rsp5 to nutrient transporters at the plasma membrane, promoting their subsequent

ubiquitylation and endocytosis [25]. ART activity is regulated by nutrient signaling pathways, allowing to

remodel the landscape of transporters at the plasma membrane to meet the physiological needs of cells

facing a nutrient challenge [26, 27]. Particularly, Rod1 regulates endocytosis in response to glucose

availability, and its activity is oppositely regulated by Snf1 and PP1 [28-30], being a direct target of Snf1 [31].

A genome-wide screen revealed that *rod1Δ* is indeed resistant to 2DG [14], thus the Snf1-mediated inhibition

of Rod1 might contribute to 2DG resistance.

Point mutations aimed at abolishing Rod1 interaction with the ubiquitin ligase Rsp5, thereby annihilating

their function as adaptor proteins in endocytosis, increased resistance to 2DG suggesting that it mediates

2DG toxicity through its function in endocytosis [32]. 2DG resistance of the *rod1Δ* strain is further increased

by the additional deletion of the *ROD1* paralogue *ROG3* [32]. Rod1 (and Rog3, to a lesser extent) controls the

endocytosis of the glucose transporters Hxt1 and Hxt3 triggered in response to 2DG exposure, and

overexpression of the same transporters in a *snf1Δ* mutant restored partial resistance to 2DG [32]. Deletion

of *ROD1* and *ROG3* stabilized hexose transporters at the plasma membrane in *snf1Δ* and restored its 2DG-

sensitivity to WT levels [32]. However, whether the effects of 2DG on endocytosis extend beyond glucose

transporters is unknown, and there is currently no molecular understanding of how transporter

stability/localization contributes to 2DG resistance.

In this study, we examined the effects of 2DG on endocytosis and found that 2DG triggers the endocytosis

of many plasma membrane proteins, most of which depend on Rod1. Mechanistically, we show that opposite

to the situation in mammalian cells, 2DG treatment leads to the dephosphorylation of Snf1/AMPK and

several targets including Rod1 in a-PP1 dependent manner. This favors Rod1 function and thus provides a

rationale as to why this arrestin is central to 2DG-induced endocytosis. Finally, we demonstrate that 2DG

resistance of the *rod1Δ* strain relies on the maintenance of active glucose transporters at the membrane,

where they can increase glucose uptake and contribute to detoxifying 2DG out of the cells, thus terminating

the detoxification process.

Results

2DG triggers the endocytosis of many plasma membrane proteins

We previously described the proteomic changes occurring in response to 2DG and focused on proteins upregulated in this condition [20]. In addition, a few transmembrane proteins of the plasma membrane displayed a significantly decreased abundance after 2h30 2DG treatment, including the amino acid permease Tat1 and the hexose transporter Hxt2 [20]. This could either be caused by a decreased synthesis or an active degradation by endocytosis. We verified this by looking at the regulation of the endogenously GFP-tagged proteins. Treatment of glucose-grown cells with 0.2% 2DG led to the targeting of plasma membrane-localized Tat1 and Hxt2 (**Fig 1A**) to the vacuole, which was accompanied by their degradation (**Fig 1B**). Of note, we observed that the vacuole was strongly fragmented in response to 2DG (**Fig S1**), confirming recent findings [33]. This may be due to the fact that 2DG causes ER stress in yeast [20], which itself induces vacuolar fragmentation [34].

Tat2 and Hxt2 endocytosis in response to 2DG is reminiscent of the behavior of the low affinity hexose transporters Hxt1 and Hxt3 [32]. To document the extent of plasma membrane remodeling caused by 2DG exposure, we studied the regulation of a representative subset of plasma membrane proteins tagged with GFP at their endogenous loci before and after 2DG treatment. This included transporters of various substrates or structural classes, as well as receptor/sensor proteins whose expression could be detected in glucose-grown cells. This showed that many of these membrane proteins are endocytosed in response to 2DG (**Fig 1D, Fig S2**).

In yeast, nutrient transporter endocytosis is driven by their ubiquitylation at the plasma membrane, which relies on the action of the ubiquitin ligase Rsp5 and adaptor proteins of the arrestin-like family (ARTs) [24, 27]. Previous work highlighted the importance of the arrestin-related protein Rod1 in the 2DG-induced endocytosis of Hxt1 and Hxt3 [32]. Indeed, we found that most of the cargoes studied here depended on Rod1 for their 2DG-induced endocytosis (**Fig 1D, Fig S2**). Although the Rod1 paralogue Rog3 displays some level of functional redundancy with Rod1 [32], Rog3 was not responsible for the endocytosis of the Rod1-independent cargo Lyp1 (**Fig S3**), although this may be the case for other cargoes. Altogether, these results indicate a prominent role for the ART protein Rod1 in 2DG-induced endocytosis.

2DG triggers endocytosis by inducing Rod1 dephosphorylation and ubiquitylation

We then focused on understanding why Rod1 function is so central for 2-deoxyglucose endocytosis. In the case of glucose-induced endocytosis, Rod1 activity is regulated through changes in its post-translational modifications [28, 29]. In a glucose-deprived medium, AMPK/Snf1 is active and phosphorylates Rod1,

inhibiting its function [28, 29, 31, 35]. Conversely, glucose addition in the medium leads to Rod1

dephosphorylation in a PP1-dependent manner, followed by its ubiquitylation by Rsp5 [28, 29]. These

modifications promote its activity as an Rsp5 adaptor, leading to transporter ubiquitylation, endocytosis and

vacuolar sorting [28, 29, 35]. Similarly, 2DG addition to glucose-grown cells triggers Rod1 ubiquitylation in an

Rsp5- and PP1-dependent manner, but paradoxically 2DG was also reported to increase Snf1 activity [30, 32].

Thus, how 2DG regulates the relative activities of Snf1/PP1 and how this impacts on Rod1 activity are not

fully understood.

Previously, the effect of 2DG on Rod1 post-translational modifications was evaluated using an 3HA-

tagged construct [32]. Although Rod1-3HA could support the glucose-induced endocytosis of Jen1 [28], we

realized that it was unable to restore the 2DG-induced endocytosis of Hxt1-GFP and Hxt3-GFP in a *rod1Δ*

context , contrary to Flag-tagged Rod1 (Flag) which functionally behaved like the wild-type protein (**Fig S4A**

and **B**). We then re-evaluated Rod1 post-translational modifications using this functional construct. As for

Rod1-3HA [32], we observed drastic changes in Rod1 mobility on gel after 2DG treatment (**Fig 2A**). In

glucose-grown cells, Rod1-Flag migrated as a diffuse band resulting from a mild phosphorylation, as

determined by phosphatase treatment, and similarly to previously published results using Rod1-3HA [28, 32]

(**Fig 2A**). This pattern gave rise to two discrete bands upon 2DG exposure, with the upper band

corresponding to ubiquitylated Rod1, as shown by its disappearance when mutating the known ubiquitylated

lysine(s) (K235, K245, K264, K267) on Rod1 as described previously [28, 32]. The lower band corresponded to

a dephosphorylated species of Rod1, as demonstrated by the fact that it migrated similarly as Rod1-Flag from

phosphatase-treated extracts (**Fig 2B**). This fits with previous results obtained in the context of glucose-

induced endocytosis, in which Rod1 ubiquitylation was preceded by its dephosphorylation [28, 29]. As

expected, Rod1 was constitutively phosphorylated in the PP1 phosphatase mutant *reg1Δ* and this

phosphorylation was no longer affected by 2DG treatment (**Fig 2B**). In contrast, mutations in the described

Rod1 phosphatase, calcineurin (*cnb1Δ*) [36, 37] or in various signaling pathways whose activation is triggered

by 2DG treatment [20] (MAPK= *slt2Δ*, *hog1Δ*; UPR: *hac1Δ*) had no impact (**Fig 2C**). This suggested that PP1

activity is required for the 2DG-induced dephosphorylation of Rod1. Moreover, deletion of *SNF1* or mutation

of 12 serine residues within potential Snf1 consensus sequences (see Material and Methods) [36] led to a

profile comparable to that obtained in 2DG-treated cells, suggesting that lack of Rod1 phosphorylation

mimics 2DG-treatment (**Fig 2D**).

Having confirmed how Rod1-Flag is post-translationally modified in response to 2DG, we assessed the

functional consequences abrogating these modifications on the function of Rod1 in endocytosis. Instead of

focusing on Hxt1 or Hxt3, which are glucose-regulated [32], we used Ina1-GFP as a model cargo (see **Fig 1D**

and **Fig S5**). Ina1 is a protein of the SUR7/Pall family, with carries 3 predicted transmembrane domains and

localizes to the plasma membrane [38]. Mutation of the ubiquitylation site(s) within Rod1 resulted in a

slowdown of Ina1 endocytosis, whereas mutation of the PPxY motifs preventing the interaction with Rsp5

[28] completely abolished it (**Fig 2E**). This was confirmed by data obtained on the endocytosis of Hxt1-GFP

and Hxt3-GFP (**Fig S6**). Growth assays demonstrate that Rod1 activity in endocytosis correlated with 2DG

sensitivity (**Fig 2F**). Interestingly, despite the fact that Rod1-S12A-Flag appeared constitutively

dephosphorylated/ubiquitylated (see **Fig 2D**), Ina1-GFP endocytosis still relied on 2DG addition, suggesting

yet another layer of regulation. Accordingly, Ina1 endocytosis was still regulated by 2DG in the *snf1Δ* mutant

(**Fig 2G**), and so were other Rod1 cargoes (**Fig S7**). Thus, Rod1 dephosphorylation (or lack of phosphorylation)

is not sufficient to trigger endocytosis, suggesting yet an additional level of regulation.

2DG promotes the PP1-dependent dephosphorylation of Snf1 and several of its targets

The fact that 2DG triggers a PP1-dependent dephosphorylation of Rod1 contrasted with previous results

showing that 2DG induces a mild phosphorylation (activation) of the PP1 substrate Snf1 within 2h of

treatment [16], similar to observations in mammalian cells [3, 39]. To explain this discrepancy, we

reexamined the effects of 2DG on Snf1 signaling. For this, we followed Snf1 phosphorylation after 2DG

treatment using antibodies directed against the activated (phosphorylated) form of human AMPKα, which

cross-reacts with yeast Snf1 [40]. As a control for total Snf1, anti-polyhistidine antibodies can be used

because Snf1 contains a stretch of 13 histidine residues that can be used for its detection [40]. However, the

signals were sometimes very weak and so we completed our results with an endogenously GFP-tagged

version of Snf1. We found this version to be fully functional (**Fig S8**), as follows. Briefly, (i) the Snf1-GFP strain

was able to use sucrose as a C source (**Fig S8A**), (ii) Mig1-Flag phosphorylation was similar in the Snf1-GFP

strain as it was in the WT upon varying glucose concentrations, (iii) Snf1-GFP phosphorylation increased at

low glucose concentrations, which was counteracted by glucose addition, and finally (iv) Snf1-GFP

phosphorylation increased in the *reg1Δ* mutant (**Fig S8B**). These data validate Snf1-GFP as a suitable epitope-

tagged version of Snf1.

After 10 min 2DG treatment, we never observed Snf1 activation, even at higher 2DG concentration (**Fig**

3A). On the contrary, 2DG consistently decreased Snf1 phosphorylation within minutes after treatment (**Fig**

3B-C). A kinetic analysis of phospho-Snf1 response to 2DG confirmed a drop in Snf1 phosphorylation that was

followed by an increase at later time points (**Fig 3C**). Snf1 dephosphorylation by 2DG required the PP1

subunit Reg1, thus mimicking the effect that glucose has on glucose-starved cells. Similar observations were

made for the Snf1 substrate Rod1-Flag (**Fig 3C**). Altogether, we conclude that 2DG triggers an immediate and

transient PP1-dependent dephosphorylation of Snf1, consistent with our (**Fig 2**) and others' [32] observations

on Rod1.

This was further confirmed by studying another Snf1 substrate, the transcriptional repressor Mig1 [41,

42] which was also dephosphorylated after 2DG treatment in a PP1-dependent manner (**Fig 3D**). Again, this

mimicked the situation described when glucose-starved cells are treated with glucose [43]. Thus, 2DG

triggers both an immediate and transitory PP1-dependent inhibition of Snf1, and the dephosphorylation of

Snf1 substrates such as Mig1 or Rod1.

Snf1 inactivation by 2DG requires 2DG phosphorylation by Hxk2

We then investigated the origin of the signal triggering the PP1-dependent dephosphorylation of

substrates. 2DG is taken up by hexose transporters and is then phosphorylated into 2DG6P [44], which

interferes with several metabolic pathways. Hxk2 is the main hexokinase isoform expressed in glucose-grown

yeast cells [45] and we studied its contribution in 2DG phosphorylation by measuring 2DG6P appearance

after 2DG treatment using an enzyme-based assay [46]. After 15 min exposure to 0.2% 2DG, 2DG6P was

readily detectable in WT cells (**Fig 4A**). In *hxk2Δ*, 2DG6P only reached 25% of the WT value (**Fig 4A**),

suggesting that Hxk2 is the main 2DG-phosphorylating enzyme.

2DG phosphorylation by hexokinase into the dead-end metabolite 2DG6P requires ATP and causes a

strong ATP depletion [47, 48]. This can be visualized within minutes after 2DG addition in WT cells using an

optimized, pH-insensitive version of the AT1.03 ATP FRET biosensor [49] (**Fig 4B**). The decrease in FRET signal

occurred only upon 2DG treatment and revealed changes in ATP content as demonstrated using a mutant

version of the FRET sensor which does not bind ATP [49] (**Fig 4B**). We found that comparatively to the WT,

the drop in FRET was lower in *hxk2Δ* cells, again suggesting that Hxk2 is key for 2DG phosphorylation (**Fig 4C**).

In line with these findings, we found that neither Snf1 nor Rod1 were dephosphorylated in response to

2DG in the *hxk2Δ* mutant (**Fig 4D**). Similarly, 2DG no longer triggered Ina1-GFP endocytosis in this strain (**Fig**

4E), nor that of Hxt1 and Hxt3 (**Fig S9A**). Overall, our data suggest that either Hxk2 itself or 2DG

phosphorylation is required for the 2DG-mediated activation of PP1. Noteworthy, overexpression of the

2DG6P phosphatase Dog2, but not that of its catalytic-dead mutant (Dog2-DDAA), also abolished Ina1-GFP

endocytosis (**Fig 4F**) and that of Hxt1 and Hxt3 (**Fig S9B**), suggesting that 2DG-6-P levels directly or indirectly

control PP1 activity. Altogether, we propose that 2DG phosphorylation by Hxk2 triggers a PP1-dependent

Snf1 dephosphorylation, Rod1 dephosphorylation and endocytosis.

Stabilization of glucose transporters at the plasma membrane confers 2DG resistance to the *rod1Δ* mutant

So far, our data provide mechanistic insights as to how 2DG triggers endocytosis through AMPK inhibition and Rod1 activation. *ROD1* deletion leads to 2DG resistance [14], but the underlying mechanism remains unclear. We previously described that a frequent strategy leading to 2DG resistance involves an increased expression of the Dog2 phosphatase at the transcriptional level [20]. However, using a *pDOG2*:LacZ reporter, we found that this was not the case in the *rod1Δ* mutant, suggesting a distinct mechanism (**Fig 5A**).

The 2DG resistance of the *rod1Δ* strain is exacerbated upon the further deletion of its paralogue *ROG3* [32], and other data indicate that 2DG resistance correlates with Rod1 activity in endocytosis [32] (see also **Fig 2F**). Since Rod1 has a central role in plasma membrane protein remodeling by 2DG (**Fig 1D** and **Fig S2**), we first hypothesized that the 2DG-mediated activation of Rod1 might trigger excessive endocytosis that would be deleterious for the cell, and could be counteracted by *ROD1* deletion. We postulated that 2DG resistance originates from the stabilization of one or several Rod1-regulated cargoes at the plasma membrane. We initially focused on the low affinity hexose transporters, Hxt1 and Hxt3, for the following reasons. First, the hypersensitivity of the *snf1Δ* mutant to 2DG is rescued by overexpression of Hxt1 and Hxt3 through an unknown mechanism [32]. Second, duplication of a chromosomal region containing hexose transporter genes (*HXT3*, *HXT6* and *HXT7*) confers 2DG-resistance in a dominant manner [22]. Since these transporters are not endocytosed in the *rod1Δ* mutant (**Fig 1C** and **S2**) [32], we thus considered that their stabilization could contribute to 2DG resistance. Indeed, deletion of either *HXT3* or *HXT1* abolished the 2DG resistance of the *rod1Δ* mutant strain (**Fig 5B**), showing that both transporters are required for this process. In contrast, deletion of the high-affinity hexose transporter gene *HXT6* did not cause the same phenotype (**Fig 5B**). Although, it seems that only low-affinity hexose transporters promote 2DG resistance, this conclusion may be biased by the fact that low- and high-affinity hexose transporters are expressed to a lower level, the former contributing more to glucose uptake than the latter in glucose-rich conditions [50].

The growth defect of the obtained *rod1Δ hxt3Δ* strain on 2DG was restored by expressing a GFP-tagged version of Hxt3 driven by its own promoter on a low-copy (centromeric) plasmid (**Fig 5C**). This mild overexpression of Hxt3 also further increased 2DG resistance of the already resistant *rod1Δ* strain but had no effect on WT cells. Only a stronger expression of Hxt3 from a multicopy (2μ) plasmid conferred resistance to WT cells (**Fig 5D**), whereas that of Hxt1 had little or no effect as found previously [32]. Hxt1 and Hxt3 may have different functions regarding 2DG resistance, since Hxt1 overexpression did not compensate for the loss of Hxt3 in the *rod1Δ hxt3Δ* background (**Fig 5D**). Overall, we conclude that a high expression of glucose transporters generally correlates with 2DG resistance.

Structure/function studies of transporters [reviewed in 51] identified residues critical for hexose binding

and translocation. In particular, residue N317 of the human glucose transporter GLUT1 binds glucose [52, 53]

and mutation of the corresponding residue in yeast transporters alters substrate specificity and transport

[54-56]. Particularly, mutation of this residue in Hxt1 (N370A) [57] and other yeast hexose transporters [54,

58] abolished glucose uptake. We mutated the corresponding residue in Hxt3 (N367A) and found that its

overexpression no longer rescued growth on 2DG, further linking glucose transporter function with 2DG

resistance (Fig 5C).

Stabilization of glucose transporters at the plasma membrane facilitates glucose uptake and 2DG

export

The fact that glucose transport correlates with 2DG resistance was counterintuitive because 2DG enters

into cells through glucose transporters [44], thus their stabilization at the cell surface may increase 2DG

uptake. Indeed, deleting both *HXT1* and *HXT3* causes partial resistance to 2DG (Fig 6A), indicating that 2DG is

mostly transported into cells through these transporters.

We considered two hypotheses to explain how glucose transporter accumulation at the plasma

membrane results in 2DG resistance. First, by depleting glucose transporters from the plasma membrane,

2DG-induced endocytosis could lower glucose intake at a time where glycolysis is already impacted, resulting

in energy shortage as previously proposed [32]. Preventing endocytosis may therefore attenuate the

glycolysis blockade by maintaining glucose uptake. Alternatively, transporters present at the cell surface may

instead favor 2DG detoxification and release into the medium, as previously observed [44] since the

directionality of transport (import or export) through hexose transporters is dictated by the concentration

gradient.

We first tested whether *rod1Δ* cells could detoxify 2DG6P more efficiently than WT cells. Cells were

treated for 2h with 2DG, and 2DG6P was generated at comparable levels in both strains (Fig 6B).

Interestingly, in WT cells, the concentration of 2DG6P after 2h 2DG treatment was comparable to that

observed after 15 min (Fig 6C), showing that saturation is reached very early after treatment. Cells were then

transferred to a 2DG-free medium for 30 min and 2DG6P was assayed again. In both strains, 2DG6P

concentration decreased during these 30 min, revealing they could detoxify 2DG6P (Fig 6B). However, the

2DG6P content was much lower in *rod1Δ* than in WT cells, suggesting that 2DG6P is better detoxified in this

strain. Therefore, the endocytosis blockade caused by the deletion of *ROD1* allows a better detoxification of

2DG. To address whether this involved Hxt3, we repeated this experiment in a *rod1Δ hxt3Δ* context (Fig 6D).

We found that this restored 2DG6P content to comparable levels as those of the WT strain, whereas the

deletion of *HXT3* alone had no significant effect on 2DG6P accumulation or disappearance. Therefore, Hxt3 is

required for the increased detoxification of 2DG6P observed in the *rod1Δ* cells.

We also tested whether hexose transporters maintenance at the membrane impacts on glucose uptake,

by using an intracellular glucose FRET sensor based on the bacterial glucose/galactose-binding transport

protein MglB [59]. Previous work established that 2DG does not bind significantly to this probe, allowing its

use to measure glucose content in our setting [60]. We assessed intracellular glucose following glucose

exposure after a 4h-treatment with 2DG, and found that glucose accumulated slightly more in *rod1Δ* than in

WT cells (**Fig 6E-F**). This depended on the presence of the glucose transporter Hxt3, since the additional

deletion of *HXT3* in the context of *rod1Δ* restored glucose uptake to WT levels (**Fig 6E-F**) and 2DG sensitivity

(**Fig 5B-D**). Finally, deletion of both *HXT1* and *HXT3* strongly impaired glucose uptake, as expected. Thus,

ROD1 deletion could facilitate cell survival independently of 2DG detoxification by allowing a sustained

glucose uptake. This was further illustrated by the fact that deletion of *ROD1* conferred 2DG resistance to a

strain lacking both *DOG1* and *DOG2*, that are central to 2DG detoxification (**Fig 6G**). Therefore, the lack of

endocytosis of hexose transporters protects cells from 2DG toxicity by several mechanisms.

Discussion

Several strategies are known to modulate 2DG sensitivity/resistance in yeast [3]. The most common

mechanism is through the aberrant activation of the AMPK orthologue, Snf1, either by gain-of-function

mutation in constituents of the AMPK heterotrimeric complex, or by loss-of-function mutations of its

inhibitory phosphatase (PP1: Glc7/Reg1) [3, 21]. How AMPK hyperactivation contributes to 2DG resistance is

not fully understood, but we previously found that this partially occurs through the expression of the

detoxification enzyme Dog2 [20]. Here, we provide important insights into the cellular effects that 2DG has

on AMPK signaling, how this impacts Rod1 activity and subsequently, on endocytosis, and how endocytosis

interference leads to 2DG resistance through the control of glucose transport activity.

We first report that 2DG acts as a strong endocytosis signal, leading to the endocytosis of many plasma

membrane proteins. These include a GPCR (Ste2), various classes of transporters, such as permeases, solute

carriers such as multidrug transporters of the major facilitator superfamily (MFS) or ATP-binding cassette

(ABC) types. This massive 2DG-induced endocytosis of membrane proteins is reminiscent of that observed in

other situations of metabolic stresses such as acute glucose or nitrogen starvation [61-63], or upon

treatment with other drugs, stressors or heat shock [26, 64-69]. Because 2DG is an artificial molecule, the

observed endocytosis is unlikely to originate from an adaptation strategy but rather informs us that 2DG-

induced metabolic blockade triggers endocytosis through mechanisms that we further explored.

First, our work highlights the major role played by the arrestin Rod1 in 2DG-induced endocytosis. Rod1

participates in the endocytosis of the glucose transporters Hxt1 and Hxt3 upon 2DG treatment [32], and we

extend these findings to many other membrane proteins. Using a tagged but fully functional protein, we

confirm previous findings [32] that 2DG treatment strongly affects Rod1 post-translational modifications

(phosphorylation and ubiquitylation), reminiscent of past studies focusing on the regulation of Rod1 by

glucose availability [28, 29, 31, 36]. Phosphorylation is a frequent mechanism for arrestin inhibition [28, 65,

70, 71], perhaps by introducing negative charges that may hinder transporter interactions [63] [reviewed in

27]. Accordingly, Rod1 dephosphorylation and its subsequent ubiquitylation are required for its activity in

endocytosis [28, 29]. Arrestin ubiquitylation is often required for their function [28, 70, 72], and this may

favor interaction with Rsp5 and thus the Rsp5-mediated ubiquitylation of transporters [73]. Overall, we

propose that 2DG induces an aberrant activation of Rod1 which explains the increased endocytosis observed

in response to 2DG.

Second, we gained insights into how 2DG triggers these changes on Rod1, given that the precise effect

of 2DG on Rod1 regulatory kinase(s) or phosphatase(s) was unclear until now. 2DG is well known to trigger

AMPK activation in mammalian cells even at short timepoints [3, 39, 74], and a similar but milder effect was

reported for Snf1 in yeast after 2h treatment [16]. Yet, this apparent activation of Snf1 did not translate into

an increased phosphorylation of its substrate Mig1 [16], and the reported dephosphorylation of Rod1 in

these conditions was also not in line with an increased Snf1 activity. Rather, by focusing on shorter

timepoints, we observed a sharp decrease in Snf1 phosphorylation shortly after 2DG addition. Thus, 2DG

treatment rapidly inhibits yeast Snf1 activity, contrary to observations on mammalian AMPK. We also found

that after this initial drop in phosphorylation, Snf1 activity slowly increases over time to reach back its initial

activity, or even higher levels [16]. However, we expect that hours of 2DG treatment strongly perturb cellular

physiology and it may be delicate to specifically pinpoint the mechanisms leading to Snf1 activation in this

context.

Beyond the fact that it involves PP1, the mechanism by which Snf1 dephosphorylation is stimulated by

2DG shortly after treatment remains to be established. However, several possibilities can be considered.

First, through a derived metabolite or a metabolic imbalance caused by the glycolytic inhibition, 2DG could

activate the Glc7/Reg1 PP1 subcomplex, for example by favoring its assembly or its recruitment onto

substrates. A second and more likely possibility is derived from previous work on Snf1 dephosphorylation in

response to glucose. Using analog-sensitive alleles of a Snf1-activating kinase or of Snf1 itself, it was shown

that PP1 activity towards Snf1 is stimulated by glucose whereas PP1 activity towards Mig1 remains constant

regardless of the glucose status [12]. Whether this is due to a better accessibility of Snf1, an increased

recruitment of PP1, or a stimulation of PP1 activity, the same mechanism could be at stake in response to

2DG. The maintenance of PP1 activity towards Snf1 substrates (eg. Mig1) in a context where Snf1 is

inactivated would be sufficient to explain why phosphorylation of Mig1 decreased in response to 2DG within

a few minutes. This may also explain observations on Rod1 which also requires PP1 for its dephosphorylation

in response to glucose [28, 29] or 2DG [this study and 32]. The increased dephosphorylation of Snf1 in

response to 2DG is reminiscent to that observed when glucose is added to cells deprived of glucose [75].

Thus, shortly after treatment and even in glucose-grown cells, 2DG treatment mimics glucose-replete

conditions rather than starvation, and part of the early 2DG response involves a drop in Snf1 activity.

This inactivation of Snf1 by 2DG involves Hxk2, as 2DG was no longer able to trigger Snf1/Rod1

dephosphorylation or the endocytosis of Ina1, Hxt1 and Hxt3 in the *hxk2Δ* strain. Hxk2 may either have a

unique function in 2DG toxicity beyond its enzymatic function, but we favor the hypothesis that Hxk2

mediates 2DG-induced signaling through its unique ability to generate 2DG6P. Despite the fact that 2DG6P

can be generated by other sugar-phosphorylating enzymes (Hxk1 and glucokinase Glk1) *in vitro* [21], only

HXK2 deletion causes 2DG resistance [14, 16, 20, 21]. Moreover, our *in vivo* data suggest that Hxk2 is the

most prominent 2DG-phosphorylating enzyme, since *hxk2Δ* cells have a lower 2DG6P content and 2DG has a

lower impact on the ATP status in these cells. The lack of effect of Hxk1 or Glk1 on 2DG toxicity could be

explained by a lower expression level or other *in vivo* regulatory mechanisms that would prevent an efficient

2DG phosphorylation. Interestingly, both *HXK2* deletion and overexpression of the 2DG-6-P phosphatase

Dog2, which converts 2DG-6-P back into 2DG, protect *snf1Δ* against 2DG hypersensitivity [16, 20], so both

may act through a decreased 2DG6P content. More generally, 2DG has virtually no effect in strains

overexpressing Dog2, further suggesting that 2DG6P (or a derived metabolite) contributes to the

dephosphorylation of Snf1 and its substrates. Further work should be aimed at understanding how the

PP1/Snf1 balance is controlled by 2DG at the molecular level, which may help decipher the long-sought

mechanism by which glucose-starved cells react to glucose addition [12, 75-77].

After delving into the mechanisms by which 2DG triggers endocytosis through the modulation of glucose

signaling pathways and Rod1 activity, we also obtained insights into how *ROD1* deletion results in 2DG

resistance. The study of Rod1 point mutants indicate a correlation between endocytosis and 2DG resistance

[this work, and 32] and Hxt1 and Hxt3 stabilization at the plasma membrane was necessary for 2DG

resistance, in line with the fact that their overexpression allows 2DG resistance [32]. Although Hxt1 and Hxt3

are the main transporters involved in 2DG uptake, their maintenance at the plasma membrane is required

for long-term resistance through two possible pathways that possibly synergize (**Fig 7**). The first pathway

involves 2DG detoxification. Indeed, 2DG triggers the induction of the Dog1 and Dog2 phosphatases which

dephosphorylate 2DG6P into 2DG [20], the latter of which could presumably be exported once its cytosolic

concentration is higher than that of the medium and provided that hexose transporters be present at the

plasma membrane. This is in line with observations showing that (i) transport across the membrane is

required to confer resistance, (ii) the same transport systems are used both for 2DG and glucose [44], and

(iii) proteins of the GLUT/HXT family support bidirectional fluxes across the membrane [eg. GLUT2 in

hepatocytes, reviewed in 78]. Accordingly, we found that *ROD1* deletion increases the ability to detoxify

2DG6P, and this depended on the presence of Hxt3 . Thus, stabilizing glucose transporters at the plasma

membrane might allow the release of 2DG back to the medium, but this possibility should be further

explored in the future. The second pathway posits that transporter stabilization at the plasma membrane

allows a sustained glucose import, and thus may help bypass the 2DG-induced glycolytic blockade. Over time,

2DG6P synthesis and accumulation should inhibit hexokinase and thus blocks glucose phosphorylation, which

itself drives glucose import [79] [reviewed in 80], but given the existence of detoxification mechanisms, the

glycolytic blockade may be only temporary. Increasing transporter availability should in principle boost

glucose import and help glycolysis recover, and accordingly, *rod1Δ* cells contain more glucose than WT cells

when exposed to glucose after 2DG treatment.

In mammalian cells, 2DG causes a fast energy depletion and AMPK activation [39]. This is exemplified by

the opposite effect that 2DG has on the endocytosis of glucose transporters. In hepatocytes, the 2DG-

induced activation of AMPK triggers the phosphorylation of the arrestin-like protein TXNIP, resulting in its

degradation and stabilization of the glucose transporter GLUT1 [74], a response which is consistent with

energy depletion and starvation response. In contrast, 2DG transiently inhibits yeast Snf1, thus promoting

Rod1-mediated glucose transporter endocytosis [this work and 32]. This somewhat paradoxical response

likely reveals that in yeast, 2DG causes energy deprivation but is also likely sensed as excess glucose (causing

a decreased Snf1/AMPK activity). The differences in 2DG signaling between these two model systems may

originate from evolutionary differences regarding their ability to detect glucose as a signaling molecule, in

addition to being a source of energy. Glucose is the preferred carbon source for *Saccharomyces cerevisiae*

and is mostly used by fermentation, implying a high glycolytic flux to sustain proliferation. Yeast has evolved

various sensing systems at the plasma membrane to detect extracellular glucose, that are wired into various

signaling pathways [81]. It is likely that Hxk2 and/or glucose phosphorylation itself acts as a glucose sensor

within the Snf1/AMPK pathway [82-84], and 2DG phosphorylation by Hxk2 may produce a glucose-replete

signal, leading to Snf1 inactivation despite changes in the energy load. This is probably only true shortly after

2DG treatment, before 2DG inhibits other aspects of cellular physiology (such as protein glycosylation or

structural sugar metabolism) leading to a complex response through stress-signaling pathway (such as UPR

or MAPK-based signaling) [3, 20]. In conclusion, the characterization of 2DG resistance strategies led once

more to a better understanding of the cellular effects of 2DG and the diversity of mechanisms that can be

deployed to bypass 2DG-induced metabolic inhibition.

Materials and Methods

Yeast strain construction and growth conditions

All yeast strains used in this study derive from the *Saccharomyces cerevisiae* BY4741 or BY4742 background and are listed in <u>table S1</u>. Apart from the mutant strains obtained from the yeast deletion collection (Euroscarf) and the fluorescent GFP-tagged strains originating from the yeast GFP clone collection {Huh, 2003 #123}, all yeast strains were constructed by transformation with the standard lithium acetate–polyethylene glycol protocol using homologous recombination and verified by polymerase chain reaction (PCR) on genomic DNA prepared with a lithium acetate (200 mM)/SDS (0.1%) method (104). Yeast cells were grown in YPD medium (2%) or in SC medium [containing yeast nitrogen base (1.7 g/liter; MP Biomedicals), ammonium sulfate (5 g/liter; Sigma-Aldrich), the appropriate drop-out amino acid preparations (MP Biomedicals), and 2% (w/v) glucose, unless otherwise indicated]. Precultures were incubated at 30°C for 8 hours and diluted in the evening to 20-ml cultures to reach mid-log phase the next morning. 2DG (Sigma) was added to mid-log phase yeast cultures grown overnight to final concentrations of 0.2% (w/v) and incubated for the indicated times.

Plasmid construction

All plasmids used in this study are listed in <u>table S2.</u>

Plasmid pSL553 (pRS313, p_{ROD1}:*ROD1*-Ncol-Flag) was generated by site-directed mutagenesis on pSL234 (pRS313-based, p_{ROD1}:*ROD1*-Flag, a kind gift of O. Vincent, CSIC, Madrid, Spain) to integrate a Ncol site between *ROD1* and the Flag-tag, allowing to substitute the WT ORF of *ROD1* by mutated versions at Sall-Ncol sites as follows. The inserts p_{ROD1}:*ROD1*-WT-Flag (pSL559), p_{ROD1}:*ROD1*-PYm-Flag (pSL560) p_{ROD1}:*ROD1*-S12A-Flag (pSL561) and p_{ROD1}:*ROD1*-KR-Flag (pSL563) were PCR amplified (oSL1662/oSL1663) using as templates pSL94 (p_{ROD1}:*ROD1*-3HA) [28], pSL119 (p_{ROD1}:*ROD1*-PYm-3HA) [28], pSL152 (p_{ROD1}:*ROD1*-S12A-3HA) (see below), and pSL147 (p_{ROD1}:*ROD1*-KR-3HA, carrying mutations K235R, K245R, K264R, K267R) [28] respectively. They were digested with Sall/Ncol and cloned at these sites into pSL553 (p_{ROD1}:*ROD1*-Ncol-Flag). Plasmid pSL152 (pRS415, p_{ROD1}-ROD1(S12A)-3HA) was made from a synthetic gene designed to mutate S125, S138, S315, S358, S447, S623, S641, S706, S720, S734, S781, and S821 residues into A (Eurofins), digested Pacl/Xmal and cloned Pacl/Xmal into pSL94 at the place of WT *ROD1* (pRS415, p_{ROD1}:*ROD1*-3HA) [28].

To construct pSL589 (pUG35-based, p_{HXT3}-Hxt3-GFP), the *HXT3* promoter (1kb) was first cloned into pUG35 after amplification by PCR on WT gDNA (oSL1739/oSL1740) at BamHI/Spel sites to give pSL592. The *HXT3-GFP* ORF was PCR-amplified (oSL1770/oSL1771) from gDNA of the Hxt3-GFP strain (ySL1027) using primers overlapping with the pSL592 sequence and cloned into pSL592 by gap-repair, and later rescued in bacteria to obtain pSL589. The N367A mutation was introduced using 2 overlapping PCR products amplified from gDNA of the HXT3-GFP strain (ySL1027) (oSL1730/oSL682 and oSL1731/oSL1563) and cloned by gap-repair into pSL592 to give pSL591. Plasmid pSL599 (pRS426, p_{HXT1}:*HXT1*) was obtained by (1) amplifying the *HXT1* gene (-515/+225) from WT genomic DNA (oSL435/oSL932) into a pCRII-blunt topo vector (pSL597), (2) cloning a BamHI/Xhol fragment of pSL597 into pRS425 at BamHI/Xhol sites. For plasmid pSL602 (pRS426, p_{HXT1}:*HXT1*), A PCR amplifying the *HXT3* gene (and 1kb of promoter) was obtained from WT genomic DNA (oSL1739/oSL1734), digested BamhI/Spel and cloned BamHI/Spel in pRS426.

All constructs were verified by sequencing.

Total protein extracts and immunoblotting

Yeast were always grown in SC medium. For each protein sample, 1.4 ml of culture was incubated with 140μl of 100% TCA for 10 min on ice to precipitate proteins, centrifuged at 16,000g at 4°C for 10 min, and broken for 10 min with glass beads. Lysates were transferred to another 1.5-ml tube to remove glass beads and centrifuged for 5 min at 16,000g at 4°C, supernatants were discarded, and protein pellets were resuspended in sample buffer [50 mM tris-HCl (pH 6.8), 100 mM dithiothreitol, 2% SDS, 0.1% bromophenol blue, and 10%

glycerol, complemented with 50 mM tris-base (pH 8.8)] (50μl/initial OD). Protein samples were heated at
37°C for 5 min and 10 μl was loaded on SDS–polyacrylamide gel electrophoresis (PAGE) gels (4-20% Mini-
PROTEAN TGX Stain-Free, Bio-Rad). After electrophoresis for 30 min at 200V, total proteins were visualized
by in-gel fluorescence using a trihalo compound incorporated in SDS–PAGE gels (stain-free TGX gels, 4–20%;
Bio-Rad) after 45 sec UV-induced photoactivation using a ChemiDoc MP imager (BioRad), serving as a loading
control. Gels were transferred on nitrocellulose membranes for 60 min (100V) in a liquid transfer system
(Bio-Rad). Membranes were blocked in Tris-buffered saline solution containing 0.5% Tween-20 (TBS-T) and
2% milk for 30 min and incubated for at least 2 hours with the corresponding primary antibodies.
Membranes were washed 3x10 min in TBS-T and incubated for at least an hour with the corresponding
secondary antibody (coupled with horseradish peroxidase). Membranes were then washed again 3x10 min in
TBS-T and incubated with SuperSignal West Femto reagent (Thermo). Luminescence signals were acquired
using a ChemiDoc MP (BioRad). Primary and secondary antibodies used in this study as well as their dilutions
are listed in table S3.
533
Treatment of protein extracts with calf intestinal phosphatase (CIP) or Endoglycosidase H (EndoH)
Calf Intestine Phosphatase (CIP, Roche ref 11097075001) was used to dephosphorylate proteins prior to
electrophoresis. Samples were prepared using 20 μL of TCA protein extract (see "Protein extracts and
immunoblotting", above), to which 2 μL of 10X CIP buffer supplied with the enzyme, as well as 0.6 μL of 1.5M
Tris pH 8.8, 0.2 μL 0.1M $MgSO_4$, and 17.2 μL of H_2O were added. The extract was split in 2 and 0.6 U CIP was
added to one of the samples. Samples were incubated at 37°C for 3 h, and loaded on SDS-PAGE.
Endoglycosidase H (EndoH) (Promega V4871, 500 U/μL) was used to deglycosylate Ina1-GFP-containing
samples. Samples were prepared using 10 μL of TCA protein extract (see "Protein extracts and
immunoblotting", above) to which 2 μL of the supplied buffer, 1 μL of EndoH and 7 μL of water were added.
A control sample without EndoH was also prepared. Samples were incubated for 2 h at 37°C and loaded on
SDS-PAGE.
545
Beta-galactosidase assays
Beta-galactosidase assays were performed using 1 ml of mid-log phase yeast cultures carrying the *pDOG1-*
LacZ (pSL409) or *pDOG2-LacZ* plasmids (pSL410) [20], grown overnight to the mid-log phase in SC medium
without uracil with 2% glucose and switched to the specified conditions. The OD(600 nm) of the culture was
measured, and samples were taken and centrifuged at 16,000g at 4°C for 10 min. Cell pellets were snap-
frozen in liquid nitrogen and resuspended in 800 μl of buffer Z (pH 7, 50 mM NaH_2PO_4, 45 mM Na_2HPO_4, 10
mM $MgSO_4$, 10 mM KCl, and 38 mM beta-mercaptoethanol). After addition of 160μl of ONPG (ortho-
nitrophenyl-d-galactopyranoside, 4 mg/ml; Sigma-Aldrich), samples were incubated at 37°C. Enzymatic
reactions were stopped in the linear phase (60-min incubation for *pDOG2-LacZ* and 120-min incubation for
the *pDOG1-LacZ* plasmid, as per initial tests) [20] by addition of 400 μl of Na_2CO_3, and cell debris were
discarded by centrifugation at 16,000g. The absorbance of clarified samples was measured with a
spectrophotometer set at 420 nm.
Galactosidase activities (arbitrary units) were calculated using the formula 1000*[A420/(A600*t)], where
A420 refers to the measured enzyme activity, A600 is the turbidity of the culture, and t is the incubation
time. Each enzymatic assay was repeated independently at least three times.
561
2DG-6-phosphate assays
Cells were grown overnight in SC media until mid-log phase. 2DG was added into the culture to a final
concentration of 0.2% for the indicated time. For the chase experiments, the cultures were centrifuged after
2 hours, washed with water and re-suspended in the same volume of SC medium without 2DG. For each time
points, 1 OD equivalent of cells were taken, immediately placed on ice, centrifuged at 4°C for 4' at 16,000g,
washed once with cold PBS and lysed using Y-PER Yeast Protein Extraction Reagent (ThermoFisher Scientific)

as previously described [85]. Briefly, 1 OD equivalent of cells was resuspended in 100µL of Y-PER buffer,
incubated for 20' at 30°C, and centrifuged for 4' at 16100g at room temperature. The supernatant (between
0.0125 and 0.0625 OD equivalents) was used for the assay and diluted in Y-PER when needed. The assay was
performed using the 2DG uptake measurement kit (Cosmo Bio USA, Carlsbad, CA, USA;, Ref. CSR-OKP-PMG-
K01H) following the manufacturer's protocol adapted for a total reaction volume of 100 µL. Absorbance at
420 nm was measured every minute for 1 hour on a SpectraMax M2 Microplate Reader (Molecular Devices).
Measures taken between 20 and 35 minutes (linear range) were taken into account to compare the slopes.
Each sample was assayed at least twice in each experiment and at least three independent experiments were
performed. Mean values were calculated and were plotted with error bars representing SEM. Statistical
significance was determined using a t-test for paired variables assuming a normal distribution of the values
(unless otherwise stated) using GraphPad Prism7.
579
Glucose FRET sensor experiments
Yeast transformed with the plasmid FLII12Pglu-700µδ6 (pSL590, [a gift from Wolf Frommer: Addgene
#28002, 59] were grown overnight in SC-URA medium to reach an OD600nm of 0.3-0.7 in the morning. Cells
were centrifuged at 3000g for 5 minutes and the cell pellets were resuspended in 20mM MES pH=6 to a final
OD of 0.6. 180µl of the cell suspension were distributed in 12 wells of a 96-well black PS microplate with flat
bottom (Greiner). Fluorescence was measured with a plate reader (Spark, TECAN) set at 30°C before the
experiment was started.
A first measurement for 3 cycles was performed to establish a baseline (each cycle is around 90 seconds with
10 seconds shaking before each measurement of the fluorescence). Cells were excited at CFP wavelength
excitation (428 nm) and the emission intensities at 485 nm (CFP) and 528 nm (Citrine) were collected. The
cycle was then paused to inject 20µl of the glucose solutions using a multichannel pipette, and the plate
rapidly reloaded in the reader to restart measurements for another 17 cycles. Data were collected and
analyzed in an Excel spreadsheet and in Prism 7.0 (GraphPad). Emission intensity values of yeast transformed
with an empty plasmid were subtracted from emissions values at 485 nm and 528 nm for each measurement
at the different glucose concentrations. The emission intensity ratio for Citrine (528 nm) over CFP (485 nm)
was calculated. All analyses were repeated (with three technical replicates) at least three times
independently. Emission ratios were normalized to the average of the three initial ratio values before glucose
addition.
598
ATP FRET sensor experiments
Yeast transformed with the plasmid pDRF1-GW yAT1.03 (pSL608) [a gift from Bas Teusink: Addgene #132781,
49] were grown overnight in SC-URA medium to reach an OD600nm of 0.3-0.7 in the morning. FRET
measurements were performed as above except cells were excited at 438 nm and the emission intensities at
483 nm (ymTq2Δ11) and 593nm (tdTomato) were collected. A plasmid encoding a mutated sensor which
does not bind ATP (pSL609) [a gift from Bas Teusink: Addgene #132782, 49] was used as a negative control.
605
606 **Drop tests**
607 Yeast cells grown in liquid-rich or SC medium for at least 6 hours at 30°C were adjusted to an optical density
608 (600 nm) of 1. Serial 10-fold dilutions were prepared in 96-well plates and spotted on plates containing rich
609 or SC medium containing 2% (w/v) agar and, when indicated, 2DG [0.05 or 0.2% (w/v)]. Plates were
610 incubated at 30°C for 3 to 4 days before scanning on a desktop scanner (Epson).
611
Microscopy
Cells were mounted in synthetic complete medium with the appropriate carbon source and observed with a
motorized Olympus BX-61 fluorescence microscope equipped with an Olympus PlanApo 100× oil-immersion
objective (1.40 NA), a Spot 4.05 charge-coupled device camera and the MetaVue acquisition software

(Molecular Devices; Sunnyvale, CA) and imaged at room temperature. GFP-tagged proteins were visualized
using a Chroma GFP II filter (excitation 440–470 nm). mCherry-tagged proteins were visualized using an
HcRed I filter (excitation 525–575 nm). Images were processed in ImageJ (NIH). Vacuolar staining was
obtained by incubating cells grown in the desired condition with 100 μM CMAC (Life Technologies) for 10 min
under agitation at 30°C. Cells are then washed twice with water before observations with a fluorescence
microscope equipped with a DAPI filter.
622
623 **Quantification of vacuolar fragmentation**
624 Quantification of vacuolar fragmentation (Fig. S1B) was performed with ImageJ. For each experiment and
625 each strain, at least 30 cells were observed and the quantification was done blindly. Cells with more than 3
626 vacuoles were considered as cells with highly fragmented vacuoles.
627
Quantification of Western-blots
Western blot quantifications were performed with ImageJ. The integrated density of proteins of interest was
divided by the integrated density of total proteins. Results were normalized to the Glc ON condition of each
protein. The ratio of undegraded protein is shown. The experiments were repeated three times. Datas were
analyzed and processed with Prism7 software (GraphPad). The statistical test performed was a one-way
analysis of variance (ANOVA) test.
634
635 **Statistical analysis**
636 Mean values were calculated using a minimum of three independent measurements from three biological
637 replicates and were plotted with error bars representing SEM. Statistical significance was determined using a
638 t-test for paired variables assuming a normal distribution of the values using GraphPad Prism7.
639

References

1. Xi H, Kurtoglu M, Lampidis TJ. The wonders of 2-deoxy-D-glucose. IUBMB Life. 2014;66(2):110-21. Epub 2014/03/01. doi: 10.1002/iub.1251. PubMed PMID: 24578297.
2. Hardie DG. Regulation of AMP-activated protein kinase by natural and synthetic activators. Acta Pharm Sin B. 2016;6(1):1-19. doi: 10.1016/j.apsb.2015.06.002. PubMed PMID: 26904394; PubMed Central PMCID: PMCPMC4724661.
3. Laussel C, Leon S. Cellular toxicity of the metabolic inhibitor 2-deoxyglucose and associated resistance mechanisms. Biochem Pharmacol. 2020;182:114213. doi: 10.1016/j.bcp.2020.114213. PubMed PMID: 32890467.
4. Kurtoglu M, Gao N, Shang J, Maher JC, Lehrman MA, Wangpaichitr M, et al. Under normoxia, 2-deoxy-D-glucose elicits cell death in select tumor types not by inhibition of glycolysis but by interfering with N-linked glycosylation. Molecular cancer therapeutics. 2007;6(11):3049-58. Epub 2007/11/21. doi: 10.1158/1535-7163.MCT-07-0310. PubMed PMID: 18025288.
5. Luengo A, Gui DY, Vander Heiden MG. Targeting Metabolism for Cancer Therapy. Cell Chem Biol. 2017;24(9):1161-80. doi: 10.1016/j.chembiol.2017.08.028. PubMed PMID: 28938091; PubMed Central PMCID: PMCPMC5744685.
6. Pajak B, Siwiak E, Soltyka M, Priebe A, Zielinski R, Fokt I, et al. 2-Deoxy-d-Glucose and Its Analogs: From Diagnostic to Therapeutic Agents. Int J Mol Sci. 2019;21(1). doi: 10.3390/ijms21010234. PubMed PMID: 31905745; PubMed Central PMCID: PMCPMC6982256.
7. Kayikci O, Nielsen J. Glucose repression in Saccharomyces cerevisiae. FEMS Yeast Res. 2015;15(6):fov068. doi: 10.1093/femsyr/fov068. PubMed PMID: 26205245; PubMed Central PMCID: PMCPMC4629793.
8. Coccetti P, Nicastro R, Tripodi F. Conventional and emerging roles of the energy sensor Snf1/AMPK in Saccharomyces cerevisiae. Microb Cell. 2018;5(11):482-94. doi: 10.15698/mic2018.11.655. PubMed PMID: 30483520; PubMed Central PMCID: PMCPMC6244292.
9. De Vit MJ, Waddle JA, Johnston M. Regulated nuclear translocation of the Mig1 glucose repressor. Molecular Biology of the Cell. 1997;8(8):1603-18. PubMed PMID: 9285828; PubMed Central PMCID: PMCPMC276179.
10. Papamichos-Chronakis M, Gligoris T, Tzamarias D. The Snf1 kinase controls glucose repression in yeast by modulating interactions between the Mig1 repressor and the Cyc8-Tup1 co-repressor. EMBO Rep. 2004;5(4):368-72. doi: 10.1038/sj.embor.7400120. PubMed PMID: 15031717; PubMed Central PMCID: PMCPMC1299031.
11. McCartney RR, Schmidt MC. Regulation of Snf1 kinase. Activation requires phosphorylation of threonine 210 by an upstream kinase as well as a distinct step mediated by the Snf4 subunit. The Journal of biological chemistry. 2001;276(39):36460-6. doi: 10.1074/jbc.M104418200. PubMed PMID: 11486005.
12. Rubenstein EM, McCartney RR, Zhang C, Shokat KM, Shirra MK, Arndt KM, et al. Access denied: Snf1 activation loop phosphorylation is controlled by availability of the phosphorylated threonine 210 to the PP1 phosphatase. J Biol Chem. 2008;283(1):222-30. Epub 2007/11/10. doi: 10.1074/jbc.M707957200. PubMed PMID: 17991748.
13. Tu J, Carlson M. REG1 binds to protein phosphatase type 1 and regulates glucose repression in Saccharomyces cerevisiae. EMBO J. 1995;14(23):5939-46. Epub 1995/12/01. PubMed PMID: 8846786; PubMed Central PMCID: PMCPMC394713.
14. Ralser M, Wamelink MM, Struys EA, Joppich C, Krobitsch S, Jakobs C, et al. A catabolic block does not sufficiently explain how 2-deoxy-D-glucose inhibits cell growth. Proc Natl Acad Sci U S A. 2008;105(46):17807-11. doi: 10.1073/pnas.0803090105. PubMed PMID: 19004802; PubMed Central PMCID: PMC2584745.

15. Schmidt MC, O'Donnell AF. 'Sugarcoating' 2-deoxyglucose: mechanisms that suppress its toxic effects.
Current genetics. 2021;67(1):107-14. doi: 10.1007/s00294-020-01122-7. PubMed PMID: 33136227.

16. McCartney RR, Chandrashekarappa DG, Zhang BB, Schmidt MC. Genetic analysis of resistance and
sensitivity to 2-deoxyglucose in Saccharomyces cerevisiae. Genetics. 2014;198(2):635-46. doi:
10.1534/genetics.114.169060. PubMed PMID: 25116136; PubMed Central PMCID: PMC4196618.

17. Barban S. Studies on the mechanism of resistance to 2-deoxy-D-glucose in mammalian cell cultures. J Biol
Chem. 1962;237:291-5. Epub 1962/02/01. PubMed PMID: 13864809.

18. Sanz P, Randez-Gil F, Prieto JA. Molecular characterization of a gene that confers 2-deoxyglucose
resistance in yeast. Yeast. 1994;10(9):1195-202. Epub 1994/09/01. doi: 10.1002/yea.320100907. PubMed
PMID: 7754708.

19. Randez-Gil F, Blasco A, Prieto JA, Sanz P. DOGR1 and DOGR2: two genes from Saccharomyces cerevisiae
that confer 2-deoxyglucose resistance when overexpressed. Yeast. 1995;11(13):1233-40. Epub
1995/10/01. doi: 10.1002/yea.320111303. PubMed PMID: 8553694.

20. Defenouillère Q, Verraes A, Laussel C, Friedrich A, Schacherer J, Leon S. The induction of HAD-like
phosphatases by multiple signaling pathways confers resistance to the metabolic inhibitor 2-
deoxyglucose. Sci Signal. 2019;12(597):aaw8000. doi: 10.1126/scisignal.aaw8000. PubMed PMID:
31481524.

21. Soncini SR, Chandrashekarappa DG, Augustine DA, Callahan KP, O'Donnell AF, Schmidt MC. Spontaneous
mutations that confer resistance to 2-deoxyglucose act through Hxk2 and Snf1 pathways to regulate gene
expression and HXT endocytosis. PLoS Genet. 2020;16(7):e1008484. doi: 10.1371/journal.pgen.1008484.
PubMed PMID: 32673313; PubMed Central PMCID: PMCPMC7386655.

22. Barney JB, Chandrashekarappa DG, Soncini SR, Schmidt MC. Drug resistance in diploid yeast is acquired
through dominant alleles, haploinsufficiency, gene duplication and aneuploidy. PLoS Genet.
2021;17(9):e1009800. doi: 10.1371/journal.pgen.1009800. PubMed PMID: 34555030; PubMed Central
PMCID: PMCPMC8460028.

23. Piper RC, Dikic I, Lukacs GL. Ubiquitin-dependent sorting in endocytosis. Cold Spring Harb Perspect Biol.
2014;6(1). Epub 2014/01/05. doi: 10.1101/cshperspect.a016808. PubMed PMID: 24384571; PubMed
Central PMCID: PMCPMC3941215.

24. MacGurn JA, Hsu PC, Emr SD. Ubiquitin and membrane protein turnover: from cradle to grave. Annu Rev
Biochem. 2012;81:231-59. Epub 2012/03/13. doi: 10.1146/annurev-biochem-060210-093619. PubMed
PMID: 22404628.

25. Sardana R, Emr SD. Membrane Protein Quality Control Mechanisms in the Endo-Lysosome System. Trends
Cell Biol. 2021;31(4):269-83. doi: 10.1016/j.tcb.2020.11.011. PubMed PMID: 33414051.

26. Babst M. Regulation of nutrient transporters by metabolic and environmental stresses. Curr Opin Cell Biol.
2020;65:35-41. doi: 10.1016/j.ceb.2020.02.009. PubMed PMID: 32200208; PubMed Central PMCID:
PMCPMC7501145.

27. Kahlhofer J, Leon S, Teis D, Schmidt O. The alpha-arrestin family of ubiquitin ligase adaptors links
metabolism with selective endocytosis. Biology of the cell / under the auspices of the European Cell
Biology Organization. 2021;113(4):183-219. Epub 2020/12/15. doi: 10.1111/boc.202000137. PubMed
PMID: 33314196.

28. Becuwe M, Vieira N, Lara D, Gomes-Rezende J, Soares-Cunha C, Casal M, et al. A molecular switch on an
arrestin-like protein relays glucose signaling to transporter endocytosis. J Cell Biol. 2012;196(2):247-59.
Epub 2012/01/18. doi: 10.1083/jcb.201109113. PubMed PMID: 22249293.

29. Llopis-Torregrosa V, Ferri-Blazquez A, Adam-Artigues A, Deffontaines E, van Heusden GP, Yenush L.
Regulation of the Yeast Hxt6 Hexose Transporter by the Rod1 alpha-Arrestin, the Snf1 Protein Kinase, and
the Bmh2 14-3-3 Protein. J Biol Chem. 2016;291(29):14973-85. doi: 10.1074/jbc.M116.733923. PubMed
PMID: 27261460; PubMed Central PMCID: PMC4946916.

30. O'Donnell AF, Schmidt MC. AMPK-Mediated Regulation of Alpha-Arrestins and Protein Trafficking. Int J
Mol Sci. 2019;20(3). Epub 2019/01/30. doi: 10.3390/ijms20030515. PubMed PMID: 30691068; PubMed
Central PMCID: PMCPMC6387238.
31. Shinoda J, Kikuchi Y. Rod1, an arrestin-related protein, is phosphorylated by Snf1-kinase in *Saccharomyces*
cerevisiae. Biochem Biophys Res Commun. 2007;364(2):258-63. Epub 2007/10/24. doi:
10.1016/j.bbrc.2007.09.134. PubMed PMID: 17949685.
32. O'Donnell AF, McCartney RR, Chandrashekarappa DG, Zhang BB, Thorner J, Schmidt MC. 2-Deoxyglucose
impairs *Saccharomyces cerevisiae* growth by stimulating Snf1-regulated and alpha-arrestin-mediated
trafficking of hexose transporters 1 and 3. Mol Cell Biol. 2015;35(6):939-55. doi: 10.1128/MCB.01183-14.
PubMed PMID: 25547292; PubMed Central PMCID: PMCPMC4333089.
33. Yang X, Reist L, Chomchai DA, Chen L, Arines FM, Li M. ESCRT, not intralumenal fragments, sorts
ubiquitinated vacuole membrane proteins for degradation. J Cell Biol. 2021;220(8). doi:
10.1083/jcb.202012104. PubMed PMID: 34047770; PubMed Central PMCID: PMCPMC8167898.
34. Stauffer B, Powers T. Target of rapamycin signaling mediates vacuolar fission caused by endoplasmic
reticulum stress in Saccharomyces cerevisiae. Mol Biol Cell. 2015;26(25):4618-30. Epub 2015/10/16. doi:
10.1091/mbc.E15-06-0344. PubMed PMID: 26466677; PubMed Central PMCID: PMCPMC4678019.
35. Becuwe M, Leon S. Integrated control of transporter endocytosis and recycling by the arrestin-related
protein Rod1 and the ubiquitin ligase Rsp5. eLife. 2014;3:03307. doi: 10.7554/eLife.03307. PubMed PMID:
25380227; PubMed Central PMCID: PMC4244573.
36. Alvaro CG, Aindow A, Thorner J. Differential Phosphorylation Provides a Switch to Control How alpha-
Arrestin Rod1 Down-regulates Mating Pheromone Response in Saccharomyces cerevisiae. Genetics.
2016;203(1):299-317. Epub 2016/02/28. doi: 10.1534/genetics.115.186122. PubMed PMID: 26920760;
PubMed Central PMCID: PMCPMC4858781.
37. Alvaro CG, O'Donnell AF, Prosser DC, Augustine AA, Goldman A, Brodsky JL, et al. Specific alpha-arrestins
negatively regulate *Saccharomyces cerevisiae* pheromone response by down-modulating the G-protein-
coupled receptor Ste2. Mol Cell Biol. 2014;34(14):2660-81. doi: 10.1128/MCB.00230-14. PubMed PMID:
24820415; PubMed Central PMCID: PMCPMC4097657.
38. Germann SM, Schramke V, Pedersen RT, Gallina I, Eckert-Boulet N, Oestergaard VH, et al. TopBP1/Dpb11
binds DNA anaphase bridges to prevent genome instability. J Cell Biol. 2014;204(1):45-59. doi:
10.1083/jcb.201305157. PubMed PMID: 24379413; PubMed Central PMCID: PMCPMC3882784.
39. Vincent EE, Coelho PP, Blagih J, Griss T, Viollet B, Jones RG. Differential effects of AMPK agonists on cell
growth and metabolism. Oncogene. 2015;34(28):3627-39. doi: 10.1038/onc.2014.301. PubMed PMID:
25241895; PubMed Central PMCID: PMCPMC4980123.
40. Orlova M, Barrett L, Kuchin S. Detection of endogenous Snf1 and its activation state: application to
Saccharomyces and Candida species. Yeast. 2008;25(10):745-54. doi: 10.1002/yea.1628. PubMed PMID:
18949820.
41. Smith FC, Davies SP, Wilson WA, Carling D, Hardie DG. The SNF1 kinase complex from Saccharomyces
cerevisiae phosphorylates the transcriptional repressor protein Mig1p in vitro at four sites within or near
regulatory domain 1. FEBS Lett. 1999;453(1-2):219-23. doi: 10.1016/s0014-5793(99)00725-5. PubMed
PMID: 10403407.
42. Treitel MA, Kuchin S, Carlson M. Snf1 protein kinase regulates phosphorylation of the Mig1 repressor in
Saccharomyces cerevisiae. Mol Cell Biol. 1998;18(11):6273-80. Epub 1998/10/17. PubMed PMID:
9774644; PubMed Central PMCID: PMCPMC109214.
43. Shashkova S, Wollman AJM, Leake MC, Hohmann S. The yeast Mig1 transcriptional repressor is
dephosphorylated by glucose-dependent and -independent mechanisms. FEMS Microbiol Lett.
2017;364(14). Epub 2017/09/01. doi: 10.1093/femsle/fnx133. PubMed PMID: 28854669.

44. van Steveninck J. Transport-associated phosphorylation of 2-deoxy-D-glucose in yeast. Biochim Biophys
Acta. 1968;163(3):386-94. Epub 1968/11/05. doi: 10.1016/0005-2736(68)90123-5. PubMed PMID:
5721900.
45. Herrero P, Galindez J, Ruiz N, Martinez-Campa C, Moreno F. Transcriptional regulation of the
Saccharomyces cerevisiae HXK1, HXK2 and GLK1 genes. Yeast. 1995;11(2):137-44. doi:
10.1002/yea.320110205. PubMed PMID: 7732723.
46. Saito K, Lee S, Shiuchi T, Toda C, Kamijo M, Inagaki-Ohara K, et al. An enzymatic photometric assay for 2-
deoxyglucose uptake in insulin-responsive tissues and 3T3-L1 adipocytes. Anal Biochem. 2011;412(1):9-
17. doi: 10.1016/j.ab.2011.01.022. PubMed PMID: 21262191.
47. Karczmar GS, Arbeit JM, Toy BJ, Speder A, Weiner MW. Selective depletion of tumor ATP by 2-
deoxyglucose and insulin, detected by 31P magnetic resonance spectroscopy. Cancer Res. 1992;52(1):71-
6. Epub 1992/01/01. PubMed PMID: 1727388.
48. Ibanez AJ, Fagerer SR, Schmidt AM, Urban PL, Jefimovs K, Geiger P, et al. Mass spectrometry-based
metabolomics of single yeast cells. Proc Natl Acad Sci U S A. 2013;110(22):8790-4. doi:
10.1073/pnas.1209302110. PubMed PMID: 23671112; PubMed Central PMCID: PMCPMC3670324.
49. Botman D, van Heerden JH, Teusink B. An Improved ATP FRET Sensor For Yeast Shows Heterogeneity
During Nutrient Transitions. ACS Sens. 2020;5(3):814-22. doi: 10.1021/acssensors.9b02475. PubMed
PMID: 32077276; PubMed Central PMCID: PMCPMC7106129.
50. Ozcan S, Johnston M. Function and regulation of yeast hexose transporters. Microbiol Mol Biol Rev.
1999;63(3):554-69. doi: 10.1042/bj3630737. PubMed PMID: 10477308; PubMed Central PMCID:
PMCPMC103746.
51. Yan N. A Glimpse of Membrane Transport through Structures-Advances in the Structural Biology of the
GLUT Glucose Transporters. J Mol Biol. 2017;429(17):2710-25. doi: 10.1016/j.jmb.2017.07.009. PubMed
PMID: 28756087.
52. Mueckler M, Makepeace C. Model of the exofacial substrate-binding site and helical folding of the human
Glut1 glucose transporter based on scanning mutagenesis. Biochemistry. 2009;48(25):5934-42. doi:
10.1021/bi900521n. PubMed PMID: 19449892; PubMed Central PMCID: PMCPMC2776625.
53. Deng D, Xu C, Sun P, Wu J, Yan C, Hu M, et al. Crystal structure of the human glucose transporter GLUT1.
Nature. 2014;510(7503):121-5. doi: 10.1038/nature13306. PubMed PMID: 24847886.
54. Farwick A, Bruder S, Schadeweg V, Oreb M, Boles E. Engineering of yeast hexose transporters to transport
D-xylose without inhibition by D-glucose. Proc Natl Acad Sci U S A. 2014;111(14):5159-64. doi:
10.1073/pnas.1323464111. PubMed PMID: 24706835; PubMed Central PMCID: PMCPMC3986176.
55. Nijland JG, Shin HY, de Jong RM, de Waal PP, Klaassen P, Driessen AJ. Engineering of an endogenous
hexose transporter into a specific D-xylose transporter facilitates glucose-xylose co-consumption in
Saccharomyces cerevisiae. Biotechnol Biofuels. 2014;7(1):168. doi: 10.1186/s13068-014-0168-9. PubMed
PMID: 25505932; PubMed Central PMCID: PMCPMC4263072.
56. Shin HY, Nijland JG, de Waal PP, de Jong RM, Klaassen P, Driessen AJ. An engineered cryptic Hxt11 sugar
transporter facilitates glucose-xylose co-consumption in Saccharomyces cerevisiae. Biotechnol Biofuels.
2015;8:176. doi: 10.1186/s13068-015-0360-6. PubMed PMID: 26535057; PubMed Central PMCID:
PMCPMC4630928.
57. Roy A, Dement AD, Cho KH, Kim JH. Assessing glucose uptake through the yeast hexose transporter 1
(Hxt1). PLoS One. 2015;10(3):e0121985. doi: 10.1371/journal.pone.0121985. PubMed PMID: 25816250;
PubMed Central PMCID: PMC4376911.
58. Nijland JG, Li X, Shin HY, de Waal PP, Driessen AJM. Efficient, D-glucose insensitive, growth on D-xylose by
an evolutionary engineered Saccharomyces cerevisiae strain. FEMS Yeast Res. 2019;19(8). doi:
10.1093/femsyr/foz083. PubMed PMID: 31782779.

59. Bermejo C, Haerizadeh F, Takanaga H, Chermak D, Frommer WB. Dynamic analysis of cytosolic glucose
and ATP levels in yeast using optical sensors. Biochem J. 2010;432(2):399-406. doi: 10.1042/BJ20100946.
PubMed PMID: 20854260; PubMed Central PMCID: PMC2992555.
60. Fehr M, Lalonde S, Lager I, Wolff MW, Frommer WB. In vivo imaging of the dynamics of glucose uptake in
the cytosol of COS-7 cells by fluorescent nanosensors. J Biol Chem. 2003;278(21):19127-33. Epub
20030320. doi: 10.1074/jbc.M301333200. PubMed PMID: 12649277.
61. Lang MJ, Martinez-Marquez JY, Prosser DC, Ganser LR, Buelto D, Wendland B, et al. Glucose starvation
inhibits autophagy via vacuolar hydrolysis and induces plasma membrane internalization by down-
regulating recycling. J Biol Chem. 2014;289(24):16736-47. doi: 10.1074/jbc.M113.525782. PubMed PMID:
24753258; PubMed Central PMCID: PMC4059118.
62. Müller M, Schmidt O, Angelova M, Faserl K, Weys S, Kremser L, et al. The coordinated action of the MVB
pathway and autophagy ensures cell survival during starvation. eLife. 2015;4:e07736. doi:
10.7554/eLife.07736. PubMed PMID: 25902403; PubMed Central PMCID: PMCPMC4424281.
63. Ivashov V, Zimmer J, Schwabl S, Kahlhofer J, Weys S, Gstir R, et al. Complementary alpha-arrestin-
ubiquitin ligase complexes control nutrient transporter endocytosis in response to amino acids. eLife.
2020;9. doi: 10.7554/eLife.58246. PubMed PMID: 32744498.
64. Volland C, Urban-Grimal D, Geraud G, Haguenauer-Tsapis R. Endocytosis and degradation of the yeast
uracil permease under adverse conditions. J Biol Chem. 1994;269(13):9833-41. Epub 1994/04/01.
PubMed PMID: 8144575.
65. MacGurn JA, Hsu PC, Smolka MB, Emr SD. TORC1 regulates endocytosis via Npr1-mediated
phosphoinhibition of a ubiquitin ligase adaptor. Cell. 2011;147(5):1104-17. Epub 2011/11/29. doi:
10.1016/j.cell.2011.09.054. PubMed PMID: 22118465.
66. Nikko E, Pelham HR. Arrestin-mediated endocytosis of yeast plasma membrane transporters. Traffic.
2009;10(12):1856-67. Epub 2009/11/17. doi: 10.1111/j.1600-0854.2009.00990.x. PubMed PMID:
19912579.
67. Megarioti AH, Primo C, Kapetanakis GC, Athanasopoulos A, Sophianopoulou V, Andre B, et al. The Bul1/2
Alpha-Arrestins Promote Ubiquitylation and Endocytosis of the Can1 Permease upon Cycloheximide-
Induced TORC1-Hyperactivation. Int J Mol Sci. 2021;22(19). doi: 10.3390/ijms221910208. PubMed PMID:
34638549; PubMed Central PMCID: PMCPMC8508209.
68. Crapeau M, Merhi A, Andre B. Stress Conditions Promote Yeast Gap1 Permease Ubiquitylation and Down-
regulation via the Arrestin-like Bul and Aly Proteins. J Biol Chem. 2014;289(32):22103-16. doi:
10.1074/jbc.M114.582320. PubMed PMID: 24942738; PubMed Central PMCID: PMC4139224.
69. Zhao Y, Macgurn JA, Liu M, Emr S. The ART-Rsp5 ubiquitin ligase network comprises a plasma membrane
quality control system that protects yeast cells from proteotoxic stress. eLife. 2013;2:e00459. doi:
10.7554/eLife.00459. PubMed PMID: 23599894; PubMed Central PMCID: PMC3628405.
70. Merhi A, André B. Internal amino acids promote Gap1 permease ubiquitylation via TORC1/Npr1/14-3-3-
dependent control of the Bul arrestin-like adaptors. Mol Cell Biol. 2012;32(22):4510-22. Epub 2012/09/12.
doi: 10.1128/mcb.00463-12. PubMed PMID: 22966204; PubMed Central PMCID: PMC3486192.
71. Lee S, Ho HC, Tumolo JM, Hsu PC, MacGurn JA. Methionine triggers Ppz-mediated dephosphorylation of
Art1 to promote cargo-specific endocytosis. J Cell Biol. 2019;218(3):977-92. doi: 10.1083/jcb.201712144.
PubMed PMID: 30610170; PubMed Central PMCID: PMCPMC6400557.
72. Lin CH, MacGurn JA, Chu T, Stefan CJ, Emr SD. Arrestin-related ubiquitin-ligase adaptors regulate
endocytosis and protein turnover at the cell surface. Cell. 2008;135(4):714-25. Epub 2008/11/04. doi:
10.1016/j.cell.2008.09.025. PubMed PMID: 18976803.
73. MacDonald C, Shields SB, Williams CA, Winistorfer S, Piper RC. A Cycle of Ubiquitination Regulates
Adaptor Function of the Nedd4-Family Ubiquitin Ligase Rsp5. Curr Biol. 2020;30(3):465-79 e5. Epub
2020/01/21. doi: 10.1016/j.cub.2019.11.086. PubMed PMID: 31956026; PubMed Central PMCID:
PMCPMC7197006.

74.Wu N, Zheng B, Shaywitz A, Dagon Y, Tower C, Bellinger G, et al. AMPK-dependent degradation of TXNIP
upon energy stress leads to enhanced glucose uptake via GLUT1. Mol Cell. 2013;49(6):1167-75. doi:
10.1016/j.molcel.2013.01.035. PubMed PMID: 23453806; PubMed Central PMCID: PMC3615143.
75.Castermans D, Somers I, Kriel J, Louwet W, Wera S, Versele M, et al. Glucose-induced posttranslational
activation of protein phosphatases PP2A and PP1 in yeast. Cell Res. 2012;22(6):1058-77. doi:
10.1038/cr.2012.20. PubMed PMID: 22290422; PubMed Central PMCID: PMCPMC3367521.
76.Randez-Gil F, Prieto JA, Sanz P. The expression of a specific 2-deoxyglucose-6P phosphatase prevents
catabolite repression mediated by 2-deoxyglucose in yeast. Current genetics. 1995;28(2):101-7. Epub
1995/07/01. PubMed PMID: 8590459.
77.Momcilovic M, Carlson M. Alterations at dispersed sites cause phosphorylation and activation of SNF1
protein kinase during growth on high glucose. J Biol Chem. 2011;286(26):23544-51. Epub 2011/05/13. doi:
10.1074/jbc.M111.244111. PubMed PMID: 21561858.
78.Mueckler M, Thorens B. The SLC2 (GLUT) family of membrane transporters. Molecular aspects of
medicine. 2013;34(2-3):121-38. doi: 10.1016/j.mam.2012.07.001. PubMed PMID: 23506862; PubMed
Central PMCID: PMCPMC4104978.
79.Bisson LF, Fraenkel DG. Involvement of kinases in glucose and fructose uptake by Saccharomyces
cerevisiae. Proc Natl Acad Sci U S A. 1983;80(6):1730-4. Epub 1983/03/01. doi: 10.1073/pnas.80.6.1730.
PubMed PMID: 6300872; PubMed Central PMCID: PMCPMC393677.
80.Lagunas R. Sugar transport in Saccharomyces cerevisiae. FEMS Microbiol Rev. 1993;10(3-4):229-42. Epub
1993/04/01. doi: 10.1016/0378-1097(93)90598-v. PubMed PMID: 8318258.
81.Santangelo GM. Glucose signaling in Saccharomyces cerevisiae. Microbiol Mol Biol Rev. 2006;70(1):253-
82. doi: 10.1128/MMBR.70.1.253-282.2006. PubMed PMID: 16524925; PubMed Central PMCID:
PMCPMC1393250.
82.Vega M, Riera A, Fernandez-Cid A, Herrero P, Moreno F. Hexokinase 2 Is an Intracellular Glucose Sensor of
Yeast Cells That Maintains the Structure and Activity of Mig1 Protein Repressor Complex. J Biol Chem.
2016;291(14):7267-85. doi: 10.1074/jbc.M115.711408. PubMed PMID: 26865637; PubMed Central
PMCID: PMCPMC4817161.
83.Moreno F, Herrero P. The hexokinase 2-dependent glucose signal transduction pathway of Saccharomyces
cerevisiae. FEMS Microbiol Rev. 2002;26(1):83-90. doi: 10.1111/j.1574-6976.2002.tb00600.x. PubMed
PMID: 12007644.
84.Ma H, Bloom LM, Walsh CT, Botstein D. The residual enzymatic phosphorylation activity of hexokinase II
mutants is correlated with glucose repression in Saccharomyces cerevisiae. Mol Cell Biol.
1989;9(12):5643-9. doi: 10.1128/mcb.9.12.5643-5649.1989. PubMed PMID: 2685572; PubMed Central
PMCID: PMCPMC363735.
85.Jayakody LN, Kadowaki M, Tsuge K, Horie K, Suzuki A, Hayashi N, et al. SUMO expression shortens the lag
phase of Saccharomyces cerevisiae yeast growth caused by complex interactive effects of major mixed
fermentation inhibitors found in hot-compressed water-treated lignocellulosic hydrolysate. Appl
Microbiol Biotechnol. 2015;99(1):501-15. doi: 10.1007/s00253-014-6174-9. PubMed PMID: 25359478.
86.Brachmann CB, Davies A, Cost GJ, Caputo E, Li J, Hieter P, et al. Designer deletion strains derived from
Saccharomyces cerevisiae S288C: a useful set of strains and plasmids for PCR-mediated gene disruption
and other applications. Yeast. 1998;14(2):115-32. Epub 1998/03/04. doi: 10.1002/(SICI)1097-
0061(19980130)14:2<115::AID-YEA204>3.0.CO;2-2. PubMed PMID: 9483801.
87.Hovsepian J, Defenouillere Q, Albanese V, Vachova L, Garcia C, Palkova Z, et al. Multilevel regulation of an
alpha-arrestin by glucose depletion controls hexose transporter endocytosis. J Cell Biol.
2017;216(6):1811-31. Epub 2017/05/05. doi: 10.1083/jcb.201610094. PubMed PMID: 28468835; PubMed
Central PMCID: PMCPMC5461024.
88.Huh WK, Falvo JV, Gerke LC, Carroll AS, Howson RW, Weissman JS, et al. Global analysis of protein
localization in budding yeast. Nature. 2003;425(6959):686-91. PubMed PMID: 14562095.

89. Dohmen RJ, Stappen R, McGrath JP, Forrova H, Kolarov J, Goffeau A, et al. An essential yeast gene
921   encoding a homolog of ubiquitin-activating enzyme. J Biol Chem. 1995;270(30):18099-109. doi:
922   10.1074/jbc.270.30.18099. PubMed PMID: 7629121.

Acknowledgments

We thank Agathe Verraes for technical assistance, Olivier Vincent (CSIC, Madrid, Spain) and Michel Becuwe (Harvard University, Cambridge, MA, USA) for sharing reagents and Alexandre Soulard (Université Claude Bernard Lyon 1, Lyon, France) for helpful discussions and critical reading of the manuscript. We also acknowledge the IJM ImagoSeine facility, member of IBiSA and the France-BioImaging infrastructure (https://anr.fr/ProjetIA-10-INBS-0004; ANR-10-INBS-04). This work was supported by grants from the Fondation ARC pour la recherche sur le cancer (https://www.fondation-arc.org, ARCPJA32020060002096 to SL), Université de Paris IDEx (https://u-paris.fr; ANR-18-IDEX-0001, "Emergence en Recherche" RS30J20IDXA1_EMERLEON) and individual fellowships from the Ligue contre le cancer (https://www.ligue-cancer.net; TAZK20115 to CL), the Fondation pour la Recherche Médicale (https://www.frm.org; SPF20150934065 to QD) and the French Ministry for Research and Education (https://www.enseignementsup-recherche.gouv.fr; to AB). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing interests: The authors have declared that no competing interests exist.

                                       FIGURE LEGENDS

Figure 1 - 2DG treatment induces the endocytosis of many plasma membrane proteins, that is mostly
Rod1-dependent. (**A**) Cells expressing Hxt2-GFP and Tat1-GFP cells were grown overnight in a glucose-
containing medium (exponential phase) and treated with 2DG for 2h30 and observed by fluorescence
microscopy. PM: plasma membrane ; V: vacuole. Scale bar: 5 μm. (**B**) Western blot on total protein extracts
from Hxt2-GFP- and Tat1-GFP-expressing cells before and after 2DG addition for 2h30, using anti-GFP
antibodies. (**C**) Plasma membrane proteins whose localization was examined by fluorescence microscopy
before and after 2h30 2DG. When endocytosed, the same proteins were also observed in a *rod1Δ* context to
check for Rod1 dependence (NT, not tested). (**D**) Fluorescence microscopy of WT and *rod1Δ* cells expressing
the indicated transporters tagged with GFP before and after 2DG treatment for 4h. Scale bar: 5 μm.

Figure 2 - Following 2DG treatment, Rod1 is dephosphorylated in a PP1-dependent manner. (**A**) Total
protein extracts of WT cells expressing either Rod1-Flag or Rod1-KR-Flag were prepared before and after 2DG
addition for 10', and immunoblotted using an anti-Flag antibody. Ub: ubiquitylated Rod1, Nat: native Rod1.
(**B**) Total protein extracts of WT or *reg1Δ* cells expressing Rod1-Flag were prepared before and after 2DG
addition for 10', and immunoblotted using an anti-Flag antibody. Samples were dephosphorylated by CIP
where indicated. Ub: ubiquitylated Rod1, Nat: native Rod1, P: phosphorylated Rod1. (**C**) Total protein
extracts of WT, *cnb1Δ, slt2Δ, hac1Δ,* or *hog1Δ* expressing Rod1-Flag were prepared before and after 2DG
addition for 10', and immunoblotted using an anti-Flag antibody. (**D**) Total protein extract of WT or *snf1Δ*
cells expressing Rod1-Flag or Rod1-S12A-Flag were prepared before and after 2DG addition for 10' and
immunoblotted using an anti-Flag antibody. (**E**) WT or *rod1Δ* cells expressing the various *ROD1*-Flag
constructs were grown in a glucose-containing medium and observed by fluorescence microscopy before and
after 2DG treatment for 4h. Scale bar, 5 μm. (**F**) Serial dilutions of cultures of the indicated strains were
spotted on SC medium or SC + 0.2% 2DG medium and grown for 4 days at 30°C. (**G**) WT or *snf1Δ* cells
expressing Ina1-GFP were grown in a glucose-containing medium and observed by fluorescence microscopy
before and after 2DG treatment for 4h. Scale bar, 5 μm.

Figure 3 - 2DG causes a fast and temporary dephosphorylation of Snf1 and its substrates through PP1
activation (**A**) Total protein extracts of WT, *reg1Δ* and *snf1Δ* cells were prepared before and after 2DG
addition at the indicated concentrations for 10 min and immunoblotted using anti-phospho-AMPK and anti-
polyHistidine antibodies. (**B**) *Left*, Total protein extracts of WT, *reg1Δ* and *snf1Δ* cells were prepared before
and after 2DG addition for the indicated time and immunoblotted using anti-phospho-AMPK and anti-
polyHistidine antibodies. *Right,* quantification of the signals (adjusted *p*-value is indicated, one-way ANOVA).
(**C**) Total protein extracts of WT and *reg1Δ* cells expressing Snf1-GFP and Rod1-Flag were prepared before
and after 2DG addition for the indicated time and immunoblotted using anti-Flag, anti-phospho-AMPK and
anti-GFP antibodies. (**D**) Total protein extracts of WT, *reg1Δ* and *snf1Δ* cells expressing Mig1-Flag were
prepared before or after 2DG addition for 10' and 2h and immunoblotted using anti-Flag antibodies.

Figure 4 - Hexokinase 2 (Hxk2) is the main 2DG-phosphorylating enzyme *in vivo* (**A**) Intracellular 2DG6P
was assayed enzymatically (see Methods) in WT and *hxk2Δ* cells grown overnight in a glucose-containing
medium and treated or not for 15 min with 0.2% 2DG. Values are normalized to the value of the WT / 15 min
(n=3 independent experiments ± SEM, *p*value indicated, paired *t*-test) (**B**) 2DG causes a decrease in ATP
content as visualized using a FRET ATP biosensor. WT cells expressing the ATP biosensor (WT or mutated
version) were grown overnight in a glucose-containing medium and treated with 0.2% 2DG (arrow). The FRET
ratio (535/570 nm) was measured over time in a plate reader (see Methods) and is represented as
normalized to the t0 value. (*n*=3 independent experiments ± SEM). (**C**) ATP levels were measured as in (B) in

WT and *hxk2Δ* cells in response to 2DG. The FRET ratio (535/570 nm) is represented as normalized to the t0
value (n=4 independent experiments ± SEM). A paired *t*-test was used to compare WT + 2DG vs *hxk2Δ* + 2DG
(*p*-value indicated). (**D**) Total protein extracts of WT and *hxk2Δ* expressing Snf1-GFP and Rod1-Flag were
prepared before and after 10' 2DG treatment and immunoblotted using anti-Flag, anti-phospho-AMPK and
anti-GFP antibodies. (**E**) WT or *hxk2Δ* cells expressing Ina1-GFP were grown in a glucose-containing medium
and observed by fluorescence microscopy before and after 2DG treatment for 4h. Scale bar, 5 μm. (**F**) WT
cells expressing Ina1-GFP and transformed with an empty plasmid, or with plasmids allowing the
overexpression of *DOG2* or its catalytic mutant *DOG2-DDAA* were grown in a glucose-containing medium and
observed by fluorescence microscopy before and after 2DG treatment for 4h. Scale bar, 5 μm.
995
996 **Figure 5- Low affinity glucose transporters are required for the resistance of the *rod1Δ* mutant to 2DG**
997 (**A**) Beta-galactosidase assays of WT and *rod1Δ* cells expressing *LacZ* under the control of the p*DOG1* or
998 p*DOG2* promoters, before and after 2DG treatments for 4h (+- SEM, n=3 independent experiments, *t*-test).
999 A.U., arbitrary units. (**B-D**) Serial dilutions of cultures of the indicated strains were spotted on SC medium or
1000 SC + 0.2% 2DG medium and grown for 4 days at 30°C.
1001

1002

Figure 6- The maintenance of glucose transporters in the *rod1Δ* mutant promotes 2DG resistance by
increasing glucose uptake and limiting 2DG toxicity (**A**) Serial dilutions of cultures of the indicated strains
were spotted on SC medium or SC + 0.2% 2DG medium and grown for 4 days at 30°C. (**B**) Intracellular 2DG6P
was assayed enzymatically in WT and *rod1Δ* cells grown overnight in a glucose-containing medium, treated
for 2h with 0.2% 2DG, and then transferred into a 2DG-free glucose-containing medium ("chase"). Values are
normalized to the value of the WT / 2h (n=4 independent experiments ± SEM, paired *t*-test, *p*-value
indicated). (**C**) 2DG6P content was assayed from WT cells as in (**B**) after 15 min and 2h treatment with 0.2%
2DG n=4 independent experiments ± SEM, paired *t*-test, *p*-value indicated). (**D**) Intracellular 2DG6P was
assayed enzymatically in WT, *hxt3Δ*, *rod1Δ* and *rod1Δ hxt3Δ* cells as in (**B**). Values are normalized to the value
of the WT / 2h (n=3 independent experiments ± SEM, paired *t*-test, *p*-value indicated).(**E**) Intracellular
glucose measurement using a FRET-based glucose biosensor (representative experiment). Cells were grown
overnight in glucose medium (exponential phase), treated for 4h with 0.2% 2DG, and transferred in a
glucose-free buffer. Fluorescence at 485 and 528 nm was measured every 90 sec in a plate reader before and
after glucose addition (2%) (see Methods) and the FRET ratio (485/528 nm) is indicated over time. (**F**)
Intracellular glucose was evaluated as in (**D**) at 2000 sec across *n*=5 independent experiments (± SEM, paired
t-test) and is represented as normalized to the "before glucose" value. (**G**) Serial dilutions of cultures of the
indicated strains were spotted on SC medium or SC + 0,1% 2DG medium and grown for 4 days at 30°C.
1020

Figure 7 - Hypothetical model for a role of hexose transporters in 2DG detoxification. Treatment of cells
with 2DG triggers the accumulation of intracellular 2DG6P. This activates stress-signaling pathways such as
cell-wall integrity signaling (CWI) or the unfolded protein response (UPR), converging onto *DOG2* expression
(1) [20]. On the other hand, 2DG6P elicits a fast PP1 activation (2) which causes AMPK inhibition and, either
directly or indirectly, Rod1 dephosphorylation. By the time Dog2 is induced and begins to dephosphorylate
2DG6P (3), Rod1 has already triggered the endocytosis of many plasma membrane proteins (4), including
that of low affinity glucose transporters. This may prevent 2DG exit and detoxification but also affect cell
survival by reducing glucose import (5).

A.

Hxt2-GFP Tat1-GFP

Glc
PM

2DG
2h30
V

B.

Hxt2-GFP Tat1-GFP

MW (kDa) Glc | 2DG 2h30 | Glc | 2DG 2h30

100
70
55

35
25

Anti-GFP

Total proteins

C.

Name	Function	Endocy-tosis ?	Rod1-depdt?
Hxt1	Glucose transporter (low aff)	Yes	Yes
Hxt3	Glucose transporter (low aff)	Yes	Yes
Hxt2	Glucose transporter (med/high aff)	Yes	Yes
Hxt4	Glucose transporter (med/high aff)	Yes	Yes
Qdr3	Multidrug transporter (MFS type)	Yes	Yes
Pdr12	Multidrug transporter (ABC type)	Yes	Yes
Vht1	Biotin transporter	Yes	No
Fui1	Uridine permease	Yes	Yes
Tat1	Tryptophan permease	Yes	Yes
Lyp1	Lysine permease	Yes	No
Can1	Arginine permease	Yes	No
Itr1	Inositol transporter	Yes	Yes
Bap2	Leucine permease	Yes	NT
Ste2	GPCR (α-factor pheromone)	Yes	No
Ina1	Unknown	Yes	Yes
Slg1/Wsc1	Cell wall stress sensor	No	-
Wsc2	Cell wall stress sensor	No	-
Wsc3	Cell wall stress sensor	No	-
Mid2	Cell wall stress sensor	No	-
Fps1	Aquaglyceroporin	No	-

D.

Rod1-dependent endocytosis   **Rod1-independent endocytosis**

Ina1-GFP   Fui1-GFP   Itr1-GFP   Can1-GFP
WT | rod1Δ   WT | rod1Δ   WT | rod1Δ   WT | rod1Δ

Glc

2DG 4h

Laussel et al. Fig 1

Laussel et al. Fig 2

Laussel et al. Fig 3

Laussel et al. Fig 4

Laussel et al. Fig 5

Laussel et al. Fig 6

Laussel et al. Figure 7.

A.

Vph1-
mCherry

| Glc | 2DG 15' | 2DG 30' | 2DG 1h | 2DG 2h | 2DG 4h |

B.

Laussel et al. Fig S1

Endocytosis in response to 2DG

Rod1-dependent

Hxt1-GFP		Hxt3-GFP		Itr1-GFP		Hxt4-GFP	
WT	*rod1Δ*	WT	*rod1Δ*	WT	*rod1Δ*	WT	*rod1Δ*

Glc

2DG 4h

Qdr3-GFP		Pdr12-GFP		Hxt2-GFP	
WT	*rod1Δ*	WT	*rod1Δ*	WT	*rod1Δ*

Glc

2DG 4h

Rod1-independent

Ste2-GFP		Lyp1-GFP		Vht1-GFP	
WT	*rod1Δ*	WT	*rod1Δ*	WT	*rod1Δ*

Glc

2DG 4h

Not tested

Bap2-GFP
WT

Glc

2DG 4h

No endocytosis in response to 2DG

Slg1-GFP	Wsc2-GFP	Wsc3-GFP	Mid2-GFP	Fps1-GFP

Glc

2DG 4h

Laussel et al. Figure S2

Laussel et al. Figure S3

A.

B.

Strain	*LEU2* plasmid	*URA3* plasmid	SC-LEU-URA	SC-LEU-URA + 0.2% 2DG
WT	Ø	Ø		
reg1Δ	Ø	Ø		
rod1Δ	Ø	Ø		
rod1Δ	Rod1-3HA	Ø		
rod1Δ	Ø	Rod1-GFP		
rod1Δ	Ø	Rod1-Flag		

Laussel et al. Figure S4

A.

B.

Laussel et al. Fig S5

Laussel et al. Figure S6

Laussel et al. Figure S7

A.

B.

Laussel et al. Figure S8

Laussel et al. Figure S9