Fast Evolution of SOS-Independent Multi-Drug Resistance in Bacteria

Le Zhang[1,†], Yunpeng Guan[1†],Yuen Yee Cheng[1], Nural N. Cokcetin[2], Amy L. Bottomley[2], Andrew Robinson[3], Elizabeth J. Harry[2], Antoine van Oijen[3], Qian Peter Su[1#], Dayong Jin[1#]

[1]Institute for Biomedical Materials and Devices (IBMD), University of Technology Sydney, Ultimo, Australia.

[2]Australian Institute for Microbiology & Infection (AIMI), University of Technology Sydney, Ultimo, Australia.

[3]School of Chemistry and Molecular Bioscience, University of Wollongong, Wollongong, Australia.

†These authors contributed equally to this work

#Corresponding author. Email: qian.su@uts.edu.au; dayong.jin@uts.edu.au

Abstract

The killing mechanism of many antibiotics involves the induction of DNA damage, either directly or indirectly, which activates the SOS response. RecA, the master regulator of the SOS response, has shown to play a central role in the evolution of resistance to fluoroquinolones, even after short-term exposure. While this paradigm is well established for DNA-damaging antibiotics, it remains unclear whether β-lactams elicit similar resistance dynamics or depend on RecA and SOS-mediated mechanisms. In this study, we observed a rapid and stable evolution of β-lactam resistance (20-fold MIC increase within 8 hours) in *Escherichia coli* lacking RecA after a single exposure to ampicillin. Contrary to expectation, this resistance emerged through an SOS-independent mechanism involving two distinct evolutionary forces: increased mutational supply and antibiotic-driven selection. Specifically, we found that RecA deletion impaired DNA repair and downregulated base excision repair pathways, while concurrently repressing the transcription of antioxidative defence genes. This dual impairment led to excessive accumulation of reactive oxygen species (ROS), which in turn promoted the emergence of resistance-conferring mutations. While ampicillin treatment did not alter survival, it selectively enriched for rare mutants arising in the RecA-deficient and ROS-elevated background. Collectively, our findings demonstrate that this oxidative environment, together with compromised DNA repair capacity, increases genetic instability and creates a selective landscape favouring the expansion of resistant clones. These results highlight the repair-redox axis as a key determinant of bacterial evolvability under antimicrobial stress.

Introduction

Addressing bacterial infections caused by emerging and drug-resistant pathogens represents a major global health priority. Bactericidal antibiotics can exert their effects on cells by directly or indirectly causing DNA damage or triggering the production of highly destructive hydroxyl radicals (1-3). This, in turn, initiates a protective mechanism known as the SOS response, which enables bacterial survival against the lethal impacts of antibiotics by activating intrinsic pathways for DNA repair (4-7). The activation of DNA repair processes relies on specific genes, such as *recA*, which encodes a recombinase involved in DNA repair, and *lexA*, a repressor of the SOS response that can be inactivated by RecA (8).

Studies have demonstrated that a single exposure to fluoroquinolones, a type of antibiotic that induces DNA breaks and triggers the SOS response, leads to the development of bacterial resistance in *Escherichia coli* (*E. coli*) through a RecA and SOS response-dependent mechanism (9). Given the crucial role of RecA in the SOS response, inhibiting RecA activity to deactivate the SOS response presents an appealing strategy for preventing the evolution of bacterial resistance to antibiotics (10). Similarly, exposure to fluoroquinolones induces the SOS response and mutagenesis in *Pseudomonas aeruginosa*, and the deletion of *recA* in this pathogen results in a significant reduction in resistance to fluoroquinolones (11).

Unlike fluoroquinolones, β-lactam antibiotics induce a RecA-dependent SOS response in *E. coli* through impaired cell wall synthesis, mediated by the DpiBA two-component signal system (12). The development of antibiotic resistance, triggered by exposure to β-lactams, has been extensively investigated using the cyclic adaptive laboratory evolution (ALE) method. Mutations that arise during cyclic ALE experiments are attributed to errors occurring during continued growth, necessitating multiple rounds of β-lactam exposure to drive the evolution of resistance in *E. coli* cells (13,14). However, the precise roles of RecA and SOS responses in the development of resistance under short-term β-lactam antibiotics exposure remain unclear.

Recently, there has been a growing interest in understanding the impact of the stress-induced accumulation of reactive oxygen species (ROS) on bacterial cells (15,16). While exploring methods to harness ROS-mediated killing has the potential to enhance the effectiveness of various antibiotics (17-19), the role of ROS in antimicrobial activity has become a topic of controversy following challenges to the initial observations (20,21). The generation of ROS

has been found to contribute to the development of multidrug resistance, as prolonged exposure

to antibiotics in cyclic ALE experiments is known to generate ROS, leading to DNA damage

and increased mutagenesis (22,23). Nevertheless, there is still limited knowledge regarding the

consequences of ROS accumulation in bacteria when the activity of RecA or the SOS response

is suppressed.

Here, we report that a single exposure to β-lactam antibiotics can rapidly drive the evolution

of multidrug resistance in *E. coli* lacking RecA. This process reflects a two-step evolutionary

mechanism: RecA deficiency increases mutational supply by impairing DNA repair, repressing

antioxidant gene expression, and promoting ROS accumulation; subsequently, antibiotic

pressure selectively enriches resistant variants from this hypermutable population.

Results

Single β-lactam exposure accelerates resistance evolution in the *recA* mutant strain

through a selection-driven mechanism

To investigate the impact of the SOS response on bacterial evolution towards β-lactam

resistance, we generated a *recA* mutant strain (*ΔrecA*) from the *E. coli* MG1655 strain. Initially,

we conducted an ALE experiment using a slightly modified treatment protocol (24) on the wild

type and *ΔrecA* strains. During a period of three weeks, the cells were subjected to cycles of

ampicillin exposure for 4.5 hours at a concentration of 50 µg/mL (10 times the MIC) each day

(Fig. S1A) (25). As anticipated based on previous studies, the intermittent ampicillin treatments

over the course of three weeks resulted in the evolution of antibiotic resistance in the wild type

strain (Fig. S1B). However, a significantly accelerated development of resistance was found in

the *ΔrecA* strain, with the average time to resistance reduced to 2 days (Fig. S1B). More

importantly, we observed that resistance even emerged in the *ΔrecA* strain after a single

exposure for 8 hours to ampicillin (Fig. 1A-C). Meanwhile, after 8 hours of treatment with 50

µg/mL ampicillin, the survival rates of both wild type and *ΔrecA* strain were consistent (Fig.

S2). To ensure that the emergence of resistance we observed was not illusory due to technical

issues during the *recA* knockout process, we employed another *ΔrecA* strain (JW2669-1)

provided by the Coli Genetic Stock Centre (CGSC) with the same killing procedure. The results

from both bacterial strains were consistent (Fig. S3A and B). To further investigate, we treated

both the wild type and *ΔrecA* cells with other β-lactams, including penicillin G and carbenicillin,

at concentrations equivalent to 10 times the MIC (1 mg/mL and 200 µg/mL, respectively) for

8 hours (26,27). Consistently, these treatments also led to a fast evolution of antibiotic
resistance in the *ΔrecA* strain (Fig. S4A and B).

To assess the stability of this accelerated antibiotic resistance acquired by the *ΔrecA* strain, we
conducted a study wherein the *ΔrecA* resistant isolates, originating from the initial 8-hour
treatment with ampicillin, were continuously cultivated in a medium devoid of antibiotics for
a period of seven days. Our findings revealed that once resistance was established, resistance
remained stable and was able to be passed on to subsequent generations even in the absence of
ampicillin (Fig. 1D). Moreover, we performed a complementation experiment by introducing
a plasmid containing *recA* under its native promoter into the *ΔrecA* strain prior to Step i in Fig.
1A, that is, before exposing the cells to ampicillin. Interestingly, this complemented strain
displayed a comparable MIC to the isogenic wild type strain and maintained its sensitivity even
after ampicillin treatment for up to 8 hours (Fig. 1E).

Antibiotic resistance evolves through the combined effects of mutational events and selection
imposed by antimicrobial pressure (28). To clarify which mechanism underlies the rapid
emergence of resistance in the *ΔrecA* strain, we systematically analysed the mutation frequency
and distribution patterns in response to ampicillin treatment. Fig. 1F shows the distribution of
rifampicin-resistant colony-forming units (CFUs) across 96 independent cultures of the wild
type and *ΔrecA* strains, with and without exposure to ampicillin (50 μg/mL for 8 hours).
Although the treatment of ampicillin in the *ΔrecA* strain displayed a highly skewed distribution
with apparent jackpot cultures, non-parametric statistical comparison (Kruskal-Wallis with
Dunn's test) did not detect a significant difference in overall mutation frequency compared to
untreated *ΔrecA*. This suggests that while the median mutation burden remained stable, rare
resistant outliers were significantly enriched, which was a pattern consistent with selection
rather than broad mutagenesis (29). Further, using the maximum likelihood estimation (MLE)
(30), we calculated the mutation rates (mutations per culture) for each group (Fig. 1G). While
the *ΔrecA* group exhibited a modest increase in baseline mutation rate compared to the wild
type strain, the addition of ampicillin led to a significant increase in estimated mutation rate.

To further assess the statistical properties of these distributions, we fitted the observed data to
a Poisson model (Fig. S5A). The mutation frequency distributions in the wild type strain with
or without the treatment of ampicillin conformed well to Poisson expectations, consistent with
random spontaneous mutation events. In contrast, the single exposure to ampicillin

significantly deviated from the Poisson model, indicating a non-random clonal enrichment
process in the $\Delta recA$ strain (31). Finally, we applied a fluctuation test-based inference
framework grounded in the Luria-Delbrück model to distinguish between mutation induction
and clonal selection (32). As shown in Fig. S5B, only the $\Delta recA$ strain treated with ampicillin
exhibited a markedly non-Poisson distribution of mutant counts, characterized by a long right-
skewed tail and the emergence of jackpot cultures. This distribution is inconsistent with
uniform mutagenesis and instead supports a model in which antibiotic treatment selectively
enriches rare early-arising resistant subpopulations. Together, these findings demonstrate that
a single exposure to β-lactam antibiotics promotes a rapid and heritable evolution of antibiotic
resistance in the $\Delta recA$ strain, predominantly through selection rather than *de novo* induction
of mutations.
147

Figure 1. Fast evolution of antibiotic resistance in *E. coli recA* mutant strain. (A)
Experimental flow for the single exposures to antibiotics in *E. coli* strains. Step i: an overnight
culture (1 x 10^9 CFU/mL cells) was diluted 1:50 into 30 mL LB medium supplemented with
50 μg/mL ampicillin and incubated at 37ºC with shaking at 250 rpm for 0, 2, 4, 6 and 8 hours;
Step ii: after each treatment, the ampicillin containing medium was removed by washing twice
in a fresh LB medium; Step iii: the surviving isolates were resuspended in 30 mL fresh LB
medium and regrown overnight at 37ºC with shaking at 250 rpm; Step iv: cell cultures were
plated onto LB agar and incubated for 16 hours at 37ºC; Step v: single colonies were inoculated
in 30 mL fresh LB medium and cultured at 37ºC with shaking at 250 rpm for 4 to 6 hours. **(B)**
MICs of ampicillin were measured against the wild type *E coli* strain after single exposures to

ampicillin. **(C)** MICs of ampicillin were measured against the *ΔrecA* strain after single

exposures to ampicillin. **(D)** After the treatment of Step v, cells were continuously cultured in

an antibiotic-free medium for seven days. MICs of ampicillin were measured each day. **(E)**

MICs of ampicillin were measured against the *ΔrecA* strain treated with ampicillin, where the

expression of RecA was restored using plasmid-based constitutive expression of *recA* before

the treatment of Step i. **(F)** Distribution of rifampicin-resistant mutant counts following single

β-lactam exposure (96 replicate cultures in each group). Statistical comparison of mutation

frequency (median values) used the Kruskal-Wallis test followed by Dunn's multiple

comparisons. **(G)** Mutation rate (mutations per culture) estimates derived by maximum

likelihood analysis. Each experiment was independently repeated at least six times using

parallel replicates, and the data are shown as mean ± SEM. Significant differences among

different treatment groups are analysed by independent t-test, $*P < 0.05$, $**P < 0.01$, $***P <$

0.001, *ns*, no significance.

Detection of drug resistance associated DNA mutations in *recA* mutant resistant isolates

In bacteria, resistance to most antibiotics requires the accumulation of drug resistance

associated DNA mutations that can arise stochastically and, under stress conditions, become

enriched through selection over time to confer high levels of resistance (33). Having observed

a non-random and right-skewed distribution of mutation frequencies in *ΔrecA* isolates

following ampicillin exposure, we next sought to determine whether specific resistance-

conferring mutations were enriched in *ΔrecA* isolates following antibiotic exposure. We

thereby randomly selected 15 colonies on non-selected LB agar plates from the wild type

surviving isolates, and antibiotic screening plates containing 50 μg/mL ampicillin from the

ΔrecA resistant isolates, respectively, and performed whole-genome sequencing. We found that

drug resistance associated mutations were present in all resistant isolates, including the

mutations in the promoter of the β-lactamase *ampC* (P_{ampC}) in 8 isolates, the ampicillin-binding

target PBP3 (*ftsI*) in 1 isolate, and the AcrAB-TolC subunit AcrB (*acrB*) mutations in 6 isolates

(Fig. 2A). A mutation in gene *nudG* was detected in wild type surviving isolates after the single

exposure to ampicillin (Fig. 2A), which is involved in pyrimidine (d)NTP hydrolysis to avoid

DNA damage (34). Other mutations were listed in the Table S1.

The presence of P_{ampC} mutations was accompanied by a significant increase in the production

of β-lactamase in bacteria (Fig. S6). This leads to specific resistance to β-lactam antibiotics.

The gene *acrB* codes for a sub-component of the AcrAB-TolC multi-drug efflux pump, which

is central in Gram-negative bacteria (35,36). Mutations in AcrAB-TolC enhance the efflux of

antibiotics and confer resistance to multiple drugs (35). Consequently, after short-term

exposure to ampicillin, the *ΔrecA* isolates carrying the *acrB* mutations exhibited resistance to

other types of antibiotics, such as chloramphenicol and kanamycin (Fig. 2B and C). Treatment

with high concentrations of 1-(1-Naphthylmethyl) piperazine (NMP), an efflux pump inhibitor

(EPI) that competitively blocks TolC-composed efflux pumps, successfully restored the

sensitivity of *ΔrecA* resistant isolates to ampicillin, bringing them back to the equivalent

concentrations found in the wild type (Fig. 2D). Collectively, these results suggest that β-

lactam treatment rapidly selects for resistance-conferring mutations, which were enriched in

ΔrecA isolates following short-term exposure.

Figure 2. Rapid induction of drug resistance associated DNA mutations in the *recA*

mutant strain. (A) Detection of drug resistance associated DNA mutations in the wild type

and *ΔrecA* strain after the single exposures to ampicillin at 50 μg/mL for 8 hours. **(B)** MICs of

chloramphenicol against the wild type and *ΔrecA* strain after the single treatment with

ampicillin were tested. **(C)** MICs of kanamycin against the wild type and *ΔrecA* strain after the

single treatment with ampicillin were tested. **(D)** The wild type, *ΔrecA*, and *ΔrecA*[acrB mutant]

resistant isolates were incubated with NMP at different concentrations for 12 hours.

Subsequently, MICs were tested in these strains for resistance to ampicillin. Each experiment

was independently repeated at least six times, and the data is shown as mean $\pm$ SEM.

Significant differences among different treatment groups are analyzed by independent t-test,

*$P < 0.05$, **$P < 0.01$, ***$P < 0.001$, *ns*, no significance.

Hindrance of SOS-independent DNA repair in *recA* mutant resistant isolates

Impairment of DNA damage repair can accelerate the accumulation of mutations and influence

bacterial adaptability under antibiotic stress. While RecA is best known for regulating the SOS

response, we asked whether its absence also impacts resistance evolution through SOS-

dependent or -independent repair mechanisms. To investigate it, we first tested the ability of

various mutants involved in different pathways of the SOS response to evolving antibiotic

resistance following a single treatment with ampicillin for 8 hours at 50 µg/mL. A mutant form

of the SOS master regulator LexA (*lexA3*), which is incapable of being cleaved and thus

defective in SOS induction, did not exhibit antibiotic resistance evolution (Fig. 3A).

Additionally, the deletion of either DpiB or DpiA (encoded by *citB* or *citA*, respectively),

inhibiting the DpiBA two-component signal transduction system, did not result in the

development of antibiotic resistance after ampicillin exposure (Fig. 3A). Moreover, the

deletion of several downstream effectors of the SOS response, including those involved in cell

division inhibition (SulA and YmfM encoded by *sulA* and *ymfM*) (37) (Fig. 3A), and DNA

repair (DNA Pol II, DNA Pol IV, and DNA Pol V encoded by *polB*, *dinB* and *umuDC*) (38)

also did not lead to the evolution of antibiotic resistance (Fig. 3A). These results indicate that

the rapid resistance evolution observed in the *ΔrecA* strain is not mediated by SOS pathway

mutants, suggesting that RecA governs resistance evolution through SOS-independent

mechanisms.

In addition to the DNA repair components associated with the SOS response, DNA Pol I plays

a role in processing RNA primers during lagging-strand synthesis and filling small gaps during

DNA repair reactions (39). Since DNA Pol I (encoded by *polA*) has been demonstrated as an

essential gene required for the growth of *E. coli* in rich medium, including the LB medium

(40,41), we next utilized Single Molecule Localization Microscopy (SMLM) to precisely locate

the chromosome and DNA Pol I in a dynamic manner. During an 8 hour exposure to ampicillin,

we observed the formation of multinucleated filaments in both the wild type and *ΔrecA* strain, indicating a pause in cell division and suggesting a time period for bacterial DNA repair to take place (Fig. 3B and C) (42). However, the expression level of DNA Pol I was significantly suppressed in the *ΔrecA* strain compared to the wild type strain after 4 hours of ampicillin exposure (Fig. 3D). More notably, the super-resolution colocalization analysis revealed a significantly lower ratio of co-localization between the chromosome and DNA Pol I in the *ΔrecA* strain (Fig. 3E). Together, these findings demonstrate that the RecA loss impairs DNA repair capacity beyond SOS regulon. This repair deficiency contributes to genetic instability and is able to facilitate the rapid evolution of antibiotic resistance through SOS-independent pathways.

Figure 3. SOS-independent impairment of DNA repair in *ΔrecA* resistant isolates. (A) MICs of ampicillin were tested against the wild type strain, *ΔrecA* strain, and mutants lacking specific genes from the SOS response after single exposures to ampicillin at 50 µg/mL for 8 hours. **(B)** Filament cell lengths in the wild type (*n=253*) and the *ΔrecA* strain (*n=216*) after single treatments with ampicillin at 50 µg/mL. **(C)** Multinucleated filaments were observed in the wild type (i) and the *ΔrecA* (ii) strain after single exposures to ampicillin at 50 µg/mL. Purple: *E. coli* chromosome; green: DNA Pol I. **(D)** Relative fold changes of DNA Pol I in the wild type and *ΔrecA* strain after single treatments with ampicillin at 50 µg/mL. **(E)** Co-localization between the *E. coli* chromosome and DNA Pol I in the wild type and *ΔrecA* strain after the single exposures to ampicillin at 50 µg/mL. Data is shown as mean ± SEM. Significant differences among different treatment groups are analyzed by independent t-test, *$P < 0.05$, **$P < 0.01$, ***$P < 0.001$, *ns*, no significance.

Repression of antioxidative gene expression promotes the evolution of antibiotic resistance in the *recA* mutant strain

To further comprehend the fast evolution of β-lactam resistance observed in this study, we investigated the gene expression changes induced by ampicillin using a comprehensive transcriptome sequencing approach (total RNA-seq). Our analysis revealed significant transcriptomic alterations in both the wild type and *ΔrecA* strain isolates following a single treatment with ampicillin, compared to untreated controls (Fig. 4A). Specifically, we identified changes in the expression of 161 and 248 coding sequences (with $\log_2 FC > 2$ and P value < 0.05) in the wild type and *ΔrecA* strains, respectively. Principal component analysis (PCA) demonstrated a notable disparity in the effects of ampicillin on the *ΔrecA* strain compared to the wild type strain (Fig. 4B). Additionally, Venn diagram analysis confirmed that 138 and 225 genes were uniquely regulated by ampicillin exposure in the wild type and *ΔrecA* strains, respectively (Fig. 4C).

To elucidate the differential expression of genes associated with specific biological functions, we conducted Gene Ontology (GO) enrichment analyses (Fig. S7A and B) and Kyoto Encyclopedia of Genes and Genomes (KEGG) pathway analyses (Fig. S7C and D). Our findings indicate that ampicillin profoundly impacted persistence pathways in the wild type strain, specifically affecting pathways related to quorum sensing, flagellar assembly, biofilm formation, and bacterial chemotaxis (43,44). Conversely, in the *ΔrecA* strain, a distinct

functional category associated with the oxidative stress response exhibited significant and

unique down-regulation. This category included activities such as sulfate transporter activity,

iron-sulfur cluster assembly, oxidoreductase activity, and carboxylate reductase activity.

To identify specific genes showing significant fold changes ($\log_2$FC > 2 and P value < 0.05),

we used volcano plots to visualize the comprehensive changes in gene expression across the

genome (Fig. 4D). We examined the transcription levels associated with the SOS response

system and found that the transcription of several proteins in the wild type strain can be

significantly induced by the single exposure to ampicillin, including *citB* and *dinB* (Fig. 4E).

However, in the *recA* mutant strain, antibiotic exposure does not affect the transcription levels

of any SOS system-related proteins, suggesting that antibiotic exposure induced the SOS

response in the wild type strain but not in the *ΔrecA* strain. More importantly, we discovered

that the induction of the transcription level of the DNA Pol I was significantly suppressed after

the single treatment of ampicillin in the *ΔrecA* strain compared with that in the wild type strain

(Fig. 4E). This is consistent with our imaging results and further supports the notion that an

SOS-independent evolutionary mechanism dominates the development of antibiotic resistance

in *recA* mutant *E. coli*.

Further, significant downregulation in the transcription of antioxidative-related genes in the

ΔrecA strain was detected, including *cysJ*, *cysI*, *cysH*, *soda*, and *sufD* (Fig. 4F). This

downregulation suggested an excessive accumulation of reactive oxygen species (ROS) due to

compromised cell antioxidative defences. It has been previously reported that the induction of

mutagenesis can be stimulated by the overproduction of ROS during antibiotic administration,

leading to the evolution of antibiotic resistance both *in vivo* and *in vitro* (22,23). Therefore, we

hypothesized that elevated ROS levels facilitate the rapid evolution of antibiotic resistance in

the *ΔrecA* strain by increasing genetic instability and enabling the selection of resistant variants.

To test it, we first examined the level of ROS generation in the wild type and *ΔrecA* strains

treated with ampicillin for 8 hours by using the fluorescent probe DCFDA/H2DCFDA. We

found that ROS levels significantly increased in both the wild type and *ΔrecA* strain after 8

hours of ampicillin treatment. However, ROS levels in the *ΔrecA* strain showed a significant

further increase compared to the wild type strain (Fig. 4G). Additionally, with the addition of

50 mM glutathione (GSH), a natural antioxidative compound, no significant change in ROS

levels was observed in either the wild type or *ΔrecA* strain before and after ampicillin treatment

(Fig. 4H). Further, we supplemented the wild type and *ΔrecA* strains with 50 mM GSH and

treated them with ampicillin at a concentration of 50 µg/ml for 8 hours. Remarkably, the
addition of GSH prevented the development of resistance to ampicillin in the *ΔrecA* strain (Fig.
4I), without impairing the bactericidal effectiveness of ampicillin (Fig. 4J).

Apart from the SOS response, bacterial cells coordinate other DNA repair activities through a
network of regulatory pathways, including base excision repair (BER) (45-47). The excessive
generation of ROS results in elevated levels of deoxy-8-oxo-guanosine triphosphate (8-oxo-
dGTP), an oxidized form of dGTP that becomes both highly toxic and mutagenic upon
integration into DNA. The presence of 8-oxo-dG can induce SNP mutations, especially those
occurring in guanine, which can be actively rectified by the BER repair pathway (47). Notably,
BER glycosylases MutH and MutY can identify and repair these 8-oxo-dG-dependent
mutations; however, when MutY and MutH are inactivated, unrepaired 8-oxo-dG can lead to
the accumulation of SNP mutations within cells (47). As a result, we conducted a further
assessment of the transcription levels of the BER repair pathway in both the wild type and
ΔrecA strain before and after a single exposure to ampicillin. We discovered that after an 8-
hour treatment of ampicillin, three DNA repair-associated proteins, including MutH, MutY,
and MutM, were notably suppressed in the *ΔrecA* strain (Fig. 4K).

Finally, we sequenced the surviving *ΔrecA* isolates and found that the addition of GSH
inhibited drug resistance associated mutations in the *ΔrecA* strain, which was detected in the
ΔrecA resistant isolates including genes of the promoter of *ampC* and *acrB* (Fig. 4L). Given
the DNA repair impairment resulting in the generation of ROS, it would have been expected
for genes involved in the oxidative stress response to be induced in RecA-deficient cells.
However, the repression of antioxidative-related genes indicated the involvement of
transcriptional repressors that might be regulated by RecA. Consequently, we examined the
transcription levels of all transcriptional repressors in both the wild type and *ΔrecA* strain.
Remarkably, we observed a significant upregulation of H-NS, a crucial transcriptional
repressor (Fig. 4M). This finding suggests that the upregulation of H-NS could potentially
contribute to the suppression of antioxidant gene expression in the *ΔrecA* strain, thereby
promoting ROS accumulation and subsequent resistance evolution. Together, these findings
demonstrate that RecA deficiency not only impairs DNA repair but also suppresses the
oxidative stress response, leading to elevated ROS and increased mutational load. This
oxidative-genetic imbalance forms the basis for enhanced mutational supply in RecA-deficient
cells

Figure 4. Overaccumulation of ROS drives the fast evolution of multi-drug resistance in the *ΔrecA* strain. (A) Clustered heatmap of relative expression of coding sequences in the wild type and *ΔrecA* strain with significant fold changes ($\log_2FC > 2$ and P value < 0.05). **(B)** Principal-component analysis (PCA) of normalised read counts for all strains. **(C)** Venn diagram of differentially expressed genes ($\log_2FC > 2$) after treatment with ampicillin at 50 µg/ml for 8 hours in the wild type and *ΔrecA* strain. **(D)** The top 10 most differentially expressed genes in the *ΔrecA* strain after the single treatment with ampicillin are labelled in each plot. Blue dots indicate genes with a significant downregulation compared to the untreated control ($\log_2FC > 2$ and P value < 0.05), and yellow dots indicate genes with a significant upregulation compared to the untreated control ($\log_2FC > 2$ and P value < 0.05). **(E)** Levels of transcription of SOS response system-associated genes and gene *polA* in the wild type and *ΔrecA* strain after single exposures to ampicillin for 8 hours. **(F)** Levels of transcription of different antioxidative associated genes in the wild type and *ΔrecA* strain after single exposures to ampicillin for 8 hours. **(G)** ROS levels were measured by flow cytometry before and after 8 hours of ampicillin treatment (50 µg/mL) in the wild type and *ΔrecA* strains. **(H)** ROS levels were measured by flow cytometry in the wild type and *ΔrecA* strains before and after 8 hours of ampicillin treatment at 50 µg/mL with the addition of GSH (50 mM). **(I)** The addition of 50 mM antioxidative compound GSH prevented the evolution of antibiotic resistance to ampicillin in the *ΔrecA* strain treated with ampicillin at 50 µg/ml for 8 hours. **(J)** Survival fraction after a single exposure to ampicillin at 50 µg/ml for 8 hours in the wild type and the *ΔrecA* strain with or without the addition of GSH at 50 mM. **(K)** Levels of transcription of proteins involved in the BER DNA repair system in the wild type and *ΔrecA* strain after single exposures to ampicillin for 8 hours. **(L)** Whole genome sequencing confirms undetectable DNA mutations in the wild type and *ΔrecA* strain treated with single exposures to ampicillin with the addition of GSH at 50 mM for 8 hours. **(M)** Transcription levels of all transcriptional repressors in the wild type and *ΔrecA* strain after single treatments with ampicillin for 8 hours. Total RNA-seq was performed with three repeats in each group. Each experiment was independently repeated at least six times, and the data are shown as mean ± SEM. Significant differences among different treatment groups are analyzed by independent t-test, $*P < 0.05$, $**P < 0.01$, $***P < 0.001$, $\#P < 0.05$.

Discussion

In this study, we challenge the prevailing view that disabling RecA and thereby inhibiting the
SOS response can prevent bacteria from developing antibiotic resistance (10). While the SOS
response and RecA have been extensively studied for their roles in antibiotic resistance
evolution (48), we observed a remarkably fast and stable evolution of multidrug resistance in
the *E. coli ΔrecA* strains following a single exposure to β-lactam antibiotics. This phenomenon
cannot be explained by canonical SOS-driven mutagenesis (49), but instead reflects the
interplay of two distinct evolution forces: RecA loss increases mutational supply through DNA
repair suppression and ROS accumulation, while antibiotic-induced lethal stress provides a
selective environment that promotes the expansion of rare and resistance-conferring variants
(Fig. 5).

Figure 5. Mechanism of rapid development of multidrug resistance in the *recA* mutant *E.*
***coli* strain.** Deletion of *recA* impairs DNA damage repair and upregulates the transcription of
the global repressor H-NS, though the mechanism of this regulation remains unclear. Elevated
H-NS levels repress the expression of multiple antioxidant-related genes (*cysI*, *cysJ*, *cysH*,
sodA, *sufD*), leading to excessive accumulation of ROS. The resulting oxidative stress increases
the overall mutational burden. Upon single exposure to β-lactam antibiotics, drug-resistant
subpopulations carrying specific mutations, such as in *acrB* or P_{ampC}, are selectively enriched,
ultimately driving the rapid emergence of multidrug resistance. This process reflects a two-step
mechanism involving enhanced mutational supply and subsequent selection under antibiotic
pressure.

To determine whether antibiotic exposure in *ΔrecA* cells directly induces mutations or
selectively enriches resistant variants, we combined statistical modelling, fluctuation analysis,

and whole-genome sequencing. Although mutation rates estimated by maximum likelihood

were moderately elevated in *ΔrecA* cells after ampicillin treatment, the mutation frequency

distributions were highly right-skewed and deviated from Poisson expectations, a hallmark of

clonal selection rather than uniform and population-wide mutagenesis. These findings align

with the classical Luria-Delbrück framework and indicate that resistance evolution in *ΔrecA* is

primarily driven by selection (29). However, our data also demonstrate that the mutational

supply is enhanced in this background due to antibiotic-induced oxidative stress and impaired

DNA repair, which together increase the likelihood that resistance-conferring mutations arise

and persist. Thus, this process reflects a selection-dominated evolutionary trajectory facilitated

by stress-enhanced mutagenesis, rather than classical induced mutation or selection on pre-

existing variants.

Mechanistically, our data reveal that RecA plays broader roles in genome stability beyond its

function in SOS activation. Disruption of individual SOS components, including *lexA*, *citA/B*,

or translesion polymerases, did not recapitulate the *ΔrecA*-specific resistance phenotype.

Instead, *ΔrecA* strains exhibited suppression of DNA Pol I. and key BER genes, such as *mutH*,

mutY, and *mutM*, confirming that RecA is essential for the maintenance of repair fidelity under

stress conditions. This defect likely enables ROS-induced DNA lesions to persist and become

fixed as mutations.

In parallel, we identified a striking repression of antioxidative stress response genes in *ΔrecA*

strains exposed to ampicillin, including *cysJIH*, *sodA*, and *sufD*. This transcriptional

suppression was associated with markedly elevated ROS levels. Crucially, supplementation

with the antioxidant GSH reversed both ROS accumulation and the emergence of resistance,

without impairing the bactericidal activity of ampicillin. This decouples the mutagenic and

lethal effects of ROS, and highlights ROS as a driver of mutational supply rather than survival.

To explore the transcriptional basis for oxidative dysregulation, we examined global repressors

and discovered that *hns* was significantly upregulated in *ΔrecA* strains after ampicillin exposure.

H-NS is a known transcriptional silencer of stress response genes, including those involved in

redox regulation (50,51). We propose that RecA deficiency may lead to H-NS derepression,

thereby silencing antioxidant defences, exacerbating ROS accumulation, and enabling

mutation accumulation under stress. Although previous findings report that H-NS can down-

regulate the transcriptional expression of *cysIJH* through the mediation of *cysB* (52,53), the

mechanistic relationship between RecA and H-NS regulation remains to be experimentally validated.

More broadly, our findings highlight the repair-redox axis as a central regulator of bacterial evolvability. Rather than solely targeting bacterial growth or survival, future antimicrobial strategies might focus on constraining mutational potential. For example, co-administration of antioxidants or repair stabilizers could buffer stress-induced mutagenesis without compromising antibiotic efficacy, which has been a concept already under exploration in cancer therapeutics (54,55).

Clinically, the emergence of *acrB* mutations and enhanced activity of the AcrAB-TolC efflux pump in *ΔrecA* strains recapitulates known multidrug resistance pathways. These observations raise concerns for therapies combining DNA repair inhibitors with ROS-inducing antibiotics or anticancer drugs (56). Our data suggest that such combinations may inadvertently promote resistance evolution, particularly in immunocompromised patients or during chemotherapy.

Finally, ROS-driven mutagenesis and repair suppression are not unique to β-lactams. Antimicrobial technologies such as antimicrobial photodynamic therapy (aPDT) and cold atmospheric plasma (CAP) (57,58) also rely on oxidative stress. In RecA-deficient or stress-sensitized bacteria, these approaches may risk accelerating resistance evolution unless accompanied by safeguards that preserve genomic integrity. Thus, maintaining the transcription of genes involved in the oxidative stress defence or combining antibiotics represents a promising strategy to prevent ROS-driven mutagenesis and thereby limit the evolutionary emergence of resistance during antimicrobial therapy.

Acknowledgements
This work was supported by the Australian Research Council (ARC grant no: APP1165135), Science and Technology Innovation Commission of Shenzhen (KQTD20170810110913065), Australia China Science and Research Fund Joint Research Centre for Point-of-Care Testing (ACSRF658277, SQ2017YFGH001190).

Author contributions

482 L.Z., G.P., Q.S., A.R., E.H., A.O. and D.J. designed the experiments. L.Z., N.C., A.B., E.H.,
483 and D.J. wrote the manuscript with input from all co-authors. L.Z., G.P., Q.S., and Y.Y.C.
484 conducted experiments and analysed data.
485

486 **Competing interests**
487 The authors declare that they have no competing interests.
488

489 **Data and materials availability**
490 All data are available in the main text or the supplementary materials.
491

Reference

1. Rebecca S. Antimicrobial-Induced DNA Damage and Genomic Instability in Microbial
Pathogens. *PLOS Pathogens*. **11**, e1004678 (2015).
2. Kohanski MA, Dwyer DJ, Hayete B, Lawrence CA, Collins JJ. A common mechanism of
cellular death induced by bactericidal antibiotics. *Cell*. **7**, 130 (5): 797-810 (2007).
3. Baquero F, Levin BR. Proximate and ultimate causes of the bactericidal action of antibiotics.
Nat Rev Microbiol. **19** (2): 123-132 (2021).
4. Žgur-Bertok D. DNA Damage Repair and Bacterial Pathogens. *PLOS Pathogens*. **9** *PLOS*
Pathogens, e1003711 (2013).
5. Dale B. Wigley. Bacterial DNA Repair: Recent Insights into The Mechanism of RecBCD,
AddAB and AdnAB. *Nature Reviews Microbiology*. **11**, 9-13 (2013).
6. Harms A, Maisonneuve E, Gerdes K. Mechanisms of Bacterial Persistence during Stress and
Antibiotic Exposure. *Science*. **354**, aaf4268 (2016).
7. Katarzyna H. Maslowska, Karolina Makiela-Dzbenska, Iwona J. Fijalkowska. The SOS
System: A Complex and Tightly Regulated Response to DNA Damage. *Environ Mol*
Mutagen. **60**, 368-384 (2019).
8. Michael M. Cox. Motoring Along with the Bacterial RecA Protein. *Nature Reviews*
Molecular Cell Biology. **8**, 127-138 (2007).
9. Barrett, T.C., Mok, W.W.K., Murawski, A.M. et al. Enhanced antibiotic resistance
development from fluoroquinolone persisters after a single exposure to antibiotic. *Nat*
Commun. **10**, 1177 (2019).
10. Athanasia Pavlopoulou. RecA: A Universal Drug Target in Pathogenic Bacteria. *Frontiers*
in Bioscience. **23**, 36-42 (2018).

11. Mercolino J, Lo Sciuto A, Spinnato MC, Rampioni G, Imperi F. RecA and Specialized Error-Prone DNA Polymerases Are Not Required for Mutagenesis and Antibiotic Resistance Induced by Fluoroquinolones in Pseudomonas aeruginosa. *Antibiotics (Basel)*. **11**, 3, 325 (2022).

12. Miller C, Thomsen LE, Gaggero C, Mosseri R, Ingmer H, Cohen SN. SOS response induction by beta-lactams and bacterial defense against antibiotic lethality. *Science*. **305** (5690), 1629-31 (2004).

13. Jahn LJ, Munck C, Ellabaan MMH, Sommer MOA. Adaptive Laboratory Evolution of Antibiotic Resistance Using Different Selection Regimes Lead to Similar Phenotypes and Genotypes. *Front Microbiol*. **8**:816 (2017).

14. Levin-Reisman I, Ronin I, Gefen O, Braniss I, Shoresh N, Balaban NQ. Antibiotic Tolerance Facilitates the Evolution of Resistance. *Science*. **355**, 826-830 (2017).

15. Dwyer DJ, Kohanski MA, Hayete B, Collins JJ. Gyrase inhibitors induce an oxidative damage cellular death pathway in *Escherichia coli*. *Mol Syst Biol*. **3**: 91 (2007).

16. Kohanski MA, Dwyer DJ, Hayete B, Lawrence CA, Collins JJ. A common mechanism of cellular death induced by bactericidal antibiotics. *Cell*. **7**, 130(5): 797-810 (2007).

17. Luan, G, Hong, Y, Drlica, K, and Zhao, X. Suppression of reactive oxygen species accumulation accounts for paradoxical bacterial survival at high quinolone concentration. *Antimicrob. Agents Chemother*. 62: e01617-e01622 (2018).

18. Zhao X, Drlica K. Reactive oxygen species and the bacterial response to lethal stress. *Curr Opin Microbiol*. **21**, 1-6 (2014).

19. Zhao X, Hong Y, Drlica K. Moving forward with reactive oxygen species involvement in antimicrobial lethality. *J Antimicrob Chemother*. **70** (3): 639-642 (2015).

20. Keren I, Wu Y, Inocencio J, Mulcahy LR, Lewis K. Killing by bactericidal antibiotics does not depend on reactive oxygen species. *Science*. **339** (6124):1213-1216 (2013).

21. Imlay JA. Diagnosing oxidative stress in bacteria: not as easy as you might think. *Curr Opin Microbiol*. **24**, 124-131 (2015).

22. Kohanski MA, DePristo MA, Collins JJ. Sublethal antibiotic treatment leads to multidrug resistance via radical-induced mutagenesis. *Mol Cell*. **37**(3): 311-320 (2010).

23. Takahashi N, Gruber CC, Yang JH, Liu X, Braff D, Yashaswini CN, Bhubhanil S, Furuta Y, Andreescu S, Collins JJ, Walker GC. Lethality of MalE-LacZ hybrid protein shares mechanistic attributes with oxidative component of antibiotic lethality. *Proc Natl Acad Sci USA*. **114** (34): 9164-9169 (2017).

24. Andersson DI, Hughes D. Microbiological effects of sublethal levels of antibiotics. *Nat.*
Rev. Microbiol. **12**, 465-478 (2014).

25. Fridman, O., Goldberg, A., Ronin, I. et al. Optimization of lag time underlies antibiotic
tolerance in evolved bacterial populations. *Nature,* **513**, 418-421 (2014).

26. Reimer LG, Stratton CW, Reller LB. Minimum inhibitory and bactericidal concentrations
of 44 antimicrobial agents against three standard control strains in broth with and without
human serum. *Antimicrob Agents Chemother*., **19**, 1050-1055 (1981).

27. Jones RN, Fuchs PC, Gerlach EH, Gavan TL. Piperacillin and carbenicillin; a collaborative
in vitro comparison against 10,838 clinical bacterial isolates. *Cleveland Clinic Journal of*
Medicine, **46**, 49-55 (1979).

28. Baquero F, Coque TM, Galán JC, Martinez JL, Cantón R. Evolutionary pathways and
trajectories in antibiotic resistance. *Clinical Microbiology Reviews*, **34**, e00050-19 (2021).

29. Luria SE, Delbrück M. Mutations of bacteria from virus sensitivity to virus resistance.
Genetics, **28**, 491-511 (1943).

30. Zheng Q. Progress of a half century in the study of the Luria–Delbrück distribution.
Mathematical Biosciences, **162**, 1-32 (1999).

31. Sarkar S, Ma WT, Sandri GVH. On fluctuation analysis: a new, simple and efficient method
for computing the expected number of mutants. *Genetica*, **85**, 173-179 (1992).

32. Foster PL. Stress-induced mutagenesis in bacteria. *Critical Reviews in Biochemistry and*
Molecular Biology, **42**, 373–397 (2007).

33. Toprak E, Veres A, Michel JB, Chait R, Hartl DL, Kishony R. Evolutionary paths to
antibiotic resistance under dynamically sustained drug selection. *Nat Genet.* **44** (1), 101-105
(2011).

34. Kamiya H, Murata-Kamiya N, Iida E, Harashima H. Hydrolysis of oxidized nucleotides by
the Escherichia coli Orf135 protein. *Biochem Biophys Res Commun.* **288** (3): 499-502
(2001).

35. El Meouche I, Dunlop MJ.. Heterogeneity in Efflux Pump Expression Predisposes
Antibiotic-Resistant Cells to Mutation. *Science.* **362**, 686-690 (2018).

36. Jessica M. A. Blair et al. Molecular Mechanisms of Antibiotic Resistance. *Nature Reviews*
Microbiology. **13**, 42-51 (2015).

37. Simmons LA, Foti JJ, Cohen SE, Walker GC. The SOS regulatory network. *EcoSal Plus*.
5.4.3 (2008).

38. Ansari S, Walsh JC, Bottomley AL, Duggin IG, Burke C, Harry EJ. A newly identified
prophage-encoded gene, *ymfM*, causes SOS-inducible filamentation in *Escherichia coli*. *J*
Bacteriol. **15**, 203(11): e00646-20 (2015).

39. Monk M, Peacey M, Gross JD. Gross. Repair of Damage Induced By Ultraviolet Light in
DNA Polymerase-Defective *Escherichia Coli* Cells. *Journal of Molecular Biology*. **58**, 623-
624 (1971).

40. Joyce CM, Grindley ND. Method for determining whether a gene of *Escherichia coli* is
essential: application to the *polA* gene. *J Bacteriol*. 158(2): 636-643 (1984).

41. Konrad EB, Lehman IR. A conditional lethal mutant of *Escherichia coli* K12 defective in
the 5' leads to 3' exonuclease associated with DNA polymerase I. *Proc Natl Acad Sci*. 71(5):
2048-2051 (1974).

42. Justice SS, Hunstad DA, Cegelski L, Hultgren SJ. Morphological Plasticity as a Bacterial
Survival Strategy. *Nature Review Microbiology*. **6**, 162-168 (2008).

43. Helaine S, Kugelberg E. Bacterial Persisters: Formation, Eradication, and Experimental
Systems. *Trends in Microbiology*. **22**, 417-424 (2014).

44. Amy L. Spoering, Kim Lewis. Bio¢Lms and Planktonic Cells of *Pseudomonas Aeruginosa*
Have Similar Resistance to Killing by Antimicrobials. *J. Bacteriol*. **183**, 6746-6751 (2001).

45. Foti JJ, Devadoss B, Winkler JA, Collins JJ, Walker GC. Oxidation of the guanine
nucleotide pool underlies cell death by bactericidal antibiotics. *Science*. 336(6079): 315-319
(2012).

46. DNA repair and mutagenesis (2nd ed.). *ASM Press* (2005).

47. Hahm JY, Park J, Jang ES, Chi SW. 8-Oxoguanine: from oxidative damage to epigenetic
and epitranscriptional modification. *Exp Mol Med*. 54(10): 1626-1642 (2022).

48. Pavlopoulou A. RecA: A Universal Drug Target in Pathogenic Bacteria. *Frontiers in*
Bioscience. **23**, 36-42 (2018).

49. Janion C. Inducible SOS response system of DNA repair and mutagenesis in Escherichia
coli. *International Journal of Biological Sciences*, **4**, 338-344 (2008).

50. Dorman CJ. H-NS: a universal regulator for a dynamic genome. *Nat Rev Microbiol*. **2**
(5):391-400 (2004).

51. Bracco L, Kotlarz D, Kolb A, Diekmann S, Buc H. Synthetic curved DNA sequences can
act as transcriptional activators in Escherichia coli. *EMBO J*. **8** (13):4289-96 (1989).

52. Álvarez R, Neumann G, Frávega J, Díaz F, Tejías C, Collao B, Fuentes JA, Paredes-Sabja
D, Calderón IL, Gil F. CysB-dependent upregulation of the Salmonella Typhimurium

cysJIH operon in response to antimicrobial compounds that induce oxidative stress. *Biochem Biophys Res Commun*. **458** (1):46-51 (2015).

53. Ishihama A, Shimada T. Hierarchy of transcription factor network in *Escherichia coli* K-12: H-NS-mediated silencing and Anti-silencing by global regulators. *FEMS Microbiol Rev*. **45** (6):fuab032 (2021).

54. Aguilera A, Gómez-González B. Genome instability: a mechanistic view of its causes and consequences. *Nature Reviews Genetics*, **9**, 204-217 (2008).

55. Klinakis A, Karagiannis D, Rampias T. Targeting DNA repair in cancer: current state and novel approaches. *Cellular and Molecular Life Sciences*, **77**, 677-703 (2020).

56. Wang M, Chen S, Ao D. Targeting DNA repair pathway in cancer: mechanisms and clinical application. *MedComm*, **2**, 654-691 (2021).

57. Wilson, BC, and Patterson, MS. The physics, biophysics and technology of photodynamic therapy. *Phys. Med. Biol*. **53**, R61-R109 (2008).

58. Vatansever F, de Melo WC, Avci P, Vecchio D, Sadasivam M, Gupta A, Chandran R, Karimi M, Parizotto NA, Yin R, Tegos GP, Hamblin MR. Antimicrobial strategies centered around reactive oxygen species-bactericidal antibiotics, photodynamic therapy, and beyond. *FEMS Microbiol Rev*. **37**(6): 955-989 (2013).

59. Ghodke H, Paudel BP, Lewis JS, Jergic S, Gopal K, Romero ZJ, Wood EA, Woodgate R, Cox MM, van Oijen AM. Spatial and temporal organization of RecA in the *Escherichia coli* DNA-damage response. *Elife*, **8**: e42761 (2019).

60. Eric M. Scholar, William B. Pratt. *The Antimicrobial Drugs*. Oxford University Press.

61. A. Gutierrez, L. Laureti, S. Crussard, H. Abida, A. Rodríguez-Rojas, J. Blázquez, Z. Baharoglu, D. Mazel, F. Darfeuille, J. Vogel, I. Matic, β-Lactam antibiotics promote bacterial mutagenesis via a RpoS-mediated reduction in replication fidelity. *Nat. Commun*. 4, 1610 (2013).

62. Kirill A. Datsenko, Barry L. Wanner. One-Step Inactivation of Chromosomal Genes in *Escherichia Coli* K-12 Using PCR Products. *Proc Natl Acad Sci*. **97**, 6640-6645 (2000).

63. Baba T, Ara T, Hasegawa M, Takai Y, Okumura Y, Baba M, Datsenko KA, Tomita M, Wanner BL, Mori H. Construction of *Escherichia Coli* K-12 In-Frame, Single-Gene Knockout Mutants: The Keio Collection. *Mol Syst Biol*, **2**, 2006.0008 (2006).

64. Van der Auwera GA & O'Connor BD. Genomics in the Cloud: Using Docker, GATK, and WDL in Terra (1st Edition). *O'Reilly Media*. (2020).

65. Cingolani P, Platts A, Wang le L, Coon M, Nguyen T, Wang L, Land SJ, Lu X, Ruden DM. A program for annotating and predicting the effects of single nucleotide polymorphisms,

SnpEff: SNPs in the genome of Drosophila melanogaster strain w1118; iso-2; iso-3. *Fly*
(Austin). **6**, 2, 80-92 (2012)

66. Liao Y, Smyth GK, Shi W. The R package Rsubread is easier, faster, cheaper and better
for alignment and quantification of RNA sequencing reads. *Nucleic Acids Res.* **30**, 923
(2019).

67. Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for
RNA-seq data with DESeq2. *Genome Biol.* **15**, 550 (2014).

68. Risso D, Ngai J, Speed TP, Dudoit S. Normalization of RNA-seq data using factor analysis
of control genes or samples. *Nat Biotechnol* **32**, 896-902 (2014).

69. Su QP, Zhao ZW, Meng L, Ding M, Zhang W, Li Y, Liu M, Li R, Gao YQ, Xie XS, Sun
Y. Superresolution Imaging Reveals Spatiotemporal Propagation of Human Replication
Foci Mediated By CTCF-Organized Chromatin Structures. *Proceedings of the National*
Academy of Sciences. **117**, 15036-15046 (2020).

70. Wang G, Li Y, Wang P, Liang H, Cui M, Zhu M, Guo L, Su Q, Sun Y, McNutt MA, Yin
Y. PTEN Regulates RPA1 and Protects DNA Replication Forks. *Cell Res*. **25**, 1189-1204
(2015).

71. Huang B, Wang W, Bates M, Zhuang X. Three-Dimensional Super-Resolution Imaging by
Stochastic Optical Reconstruction Microscopy. *Science*. **319**, 810-813 (2008).

72. Levet F, Hosy E, Kechkar A, Butler C, Beghin A, Choquet D, Sibarita JB. SR-tesseler: A
Method to Segment and Quantify Localization-Based Super-Resolution Microscopy Data.
Nat. Methods. **12**, 1065–1071 (2015).

Materials and Methods

Bacterial strains, medium and antibiotics

Bacterial strains and plasmids used in this work are described in Table S2 and Table S3. Luria-
Bertani (LB) was used as broth or in agar plates. *E. coli* cells were grown in LB liquid medium
or on LB agar (1.5% w/v) plates at 37ºC, unless stated otherwise, antibiotics were
supplemented, where appropriate. Antibiotic stock solutions were prepared by dissolving
antibiotics in MilliQ filter sterilising, including ampicillin (50 mg/mL), penicillin G (100
mg/mL), carbenicillin (20 mg/mL), kanamycin (50 mg/mL) and tetracycline (10 mg/mL).
Chloramphenicol stock solution was prepared in 95% EtOH (25 mg/mL). Antibiotic solutions
were stored at -20ºC (long-term) or 4ºC (short-term).

Treatment with antibiotics to induce evolutionary resistance

For the single exposure to antibiotic experiment, an overnight culture (0.6 mL; 1 x 10^9 CFU/mL

cells) was diluted 1:50 into 30 mL LB medium supplemented with antibiotics (50 μg/mL

ampicillin, 1 mg/mL penicillin G, or 200 μg/mL carbenicillin) and incubated at 37ºC with

shaking at 250 rpm for 0, 2, 4, 6 and 8 hours, respectively. After each treatment, the antibiotic-

containing medium was removed by washing twice (20 min centrifugation at 1500 g) in fresh

LB medium (See Fig. 1A for method overview).

To test resistance, the surviving isolates were first resuspended in 30 mL LB medium and

grown overnight at 37ºC with shaking at 250 rpm. The regrown culture was then plated onto

LB agar and incubated overnight at 37ºC. Single colonies were isolated and grown in LB

medium for 4-6 hours at 37ºC with shaking at 250 rpm, which were then used to test the

resistance or stored at -80ºC for future use.

For the ALE antibiotic treatment experiments, an overnight culture (0.6 mL; 1 x 10^9 CFU/mL

cells) was diluted 1:50 into 30 mL LB medium supplemented with 50 μg/mL ampicillin and

incubated at 37ºC with shaking at 250 rpm for 4.5 hours. After treatment, the antibiotic-

containing medium was removed by washing twice (20 min centrifugation at 1500 g) in a fresh

LB medium. The remaining pellet was resuspended in 30 mL LB medium and grown overnight

at 37ºC with shaking at 250 rpm. Ampicillin treatment was applied to the regrown culture and

repeated until resistance was established, as confirmed by MIC measurement.

Antibiotic susceptibility testing

The susceptibility of *E. coli* cells to antibiotics was measured using minimum inhibitory

concentration (MIC) testing (60). In brief, overnight cultures were diluted and incubated at

37ºC for 4-6 hours with shanking at 250 rpm. Cells were then diluted 1:100 and incubated with

increasing concentrations of antibiotics in the Synergy HT BioTek plate reader (BioTek

Instruments Inc., USA) at 37ºC for 16 hours. It was programmed to measure the OD hourly at

595 nm (Gen5 software, BioTek Instruments Inc., USA). The minimum inhibitory

concentration was determined as the concentration of antibiotic where no visible growth was

observed.

Survival assays

Overnight cultures of *E. coli* were prepared from single colonies in LB medium and incubated

at 37°C with shaking (250 rpm). The overnight cultures were diluted 1:50 in fresh LB medium

containing 50 μg/mL ampicillin to initiate antibiotic treatment. Cultures were incubated at
37°C for the indicated times under shaking conditions. Following treatment, cells were
collected by centrifugation (1500 g, 20 minutes), and the antibiotic-containing medium was
removed by washing twice with fresh LB medium. Serial dilutions of the washed cultures were
prepared, and 25 μL of each dilution was plated onto LB agar plates. Plates were incubated
overnight at 37°C, and CFUs were counted the following day to evaluate bacterial survival
rates.

Mutation frequency and fluctuation analysis

723
Overnight cultures inoculated from single colonies in LB medium were diluted 1:1,000,000
and incubated at 37°C with shaking until the OD_{600} reached to 1~1.3. This extreme dilution
minimizes the presence of pre-existing stationary phase mutants and allows *de novo* mutation
events to occur during exponential growth. For each biological condition, 96 independent
parallel cultures were prepared to perform a fluctuation analysis. The total number of colony-
forming units per ml (CFU/ml) was determined by plating on LB agar. To identify rifampicin-
resistant mutants, the remaining culture volume was centrifuged and plated on LB agar
containing rifampicin (100 μg/mL). LB plates were incubated for 24 hours at 37°C and
selective plates were incubated for 48-72 hours at 37°C (61).

The mutation frequency was calculated as the ratio of CFU/ml on rifampicin-containing plates
to CFU/ml on non-selective LB plates for each culture. Distributions of mutation frequencies
across replicate cultures were plotted, and deviations from Poisson expectations were assessed
by model fitting.

To estimate the mutation rate (mutations per culture), maximum likelihood estimation (MLE)
was applied using the Ma-Sandri-Sarkar algorithm implemented in the FALCOR toolset
(https://www.keshavsingh.org/protocols/FALCOR.html). The inferred mutation rate
distributions were compared across treatment conditions, and non-Poisson distributions
indicative of jackpot cultures were further analysed using fluctuation test-based inference
following the Luria–Delbrück framework (29).

Construction of *recA* deletion mutant

746
Lambda Red recombination was used to generate the gene *recA* deletion in the *E. coli* K-12
strain, followed by previously reported methods with modifications (62,63). Primers (*recA-*

FWD and *recA-REV,* Table S4) were designed approximately 50 bp upstream and downstream

to the gene *recA* on the chromosome to amplify the tetracycline cassette as well as the flanking

DNA sequence needed for homologous recombination. Phusion polymerase (NEB) was used

to amplify the DNA sequence (Table S4), and the reaction was cleaned up using a PureLink™

PCR purification kit (ThermoFisher Scientific) as per the manufacturer's instructions. Electro-

competent *E. coli* MG1655 containing the recombinase plasmid pKD46 was transformed with

50 ng of amplified DNA a 30°C. The transformation was plated onto LB agar plates containing

10 μg/mL tetracycline and incubated overnight at 37°C. PCR was used to confirm the insertion

of the tetracycline resistance cassette at the correct site on the chromosome using primers

upstream and downstream to the gene *recA*. The newly constructed mutant strains were cured

of plasmid pKD46 by incubating LB streak plates at 42°C overnight. Loss of the plasmid was

confirmed by lack of ampicillin sensitivity on LB agar plates. Mutant strains were made

electro-competent, and 50 μL of cells were transformed with plasmid pCP20 and incubated on

100 μg/mL ampicillin plates at 30°C overnight. A few colonies were then restreaked onto LB

plates and incubated overnight at 42°C. PCR products confirmed the loss of cassette and

plasmid.

β-lactamase assay

The amount of β-lactamase was measured using a β-lactamase Activity Assay Kit (Sigma-

Aldrich, US). Briefly, cells were collected by centrifugation at 10,000 g for 10 min, and the

pellet was resuspended with 5 μL of assay buffer per mg of sample. Then, 48 μL of the sample

was mixed with 2 μL of nitrocefin. The β-Lactamase activity was monitored by measuring the

absorbance at 490 nm for 30 min at 28°C. The level of β-lactamase was determined by the

absorbance at OD_{390}.

Whole genome sequencing

Resistant clones were isolated by selection using LB agar plates with the supplementation of

ampicillin at 50 μg/mL, and non-resistant clones were isolated from the LB agar plates without

the supplementation of antibiotics. Chromosomal DNA was extracted and purified using the

PureLink™ Genomic DNA mini kit following the manufacturer's instructions (ThermoFisher

Scientific). Whole genome sequencing (WGS) was conducted following the Nextera Flex

library preparation kit process (Illumina). Briefly, genomic DNA was quantitatively assessed

using Quant-iT picogreen dsDNA assay kit (Invitrogen, USA). The sample was normalised to

the concentration of 1 ng/μL. 10 ng of DNA was used for library preparation. After

tagmentation, the tagmented DNA was amplified using the facility's custom-designed i7 or i5
barcodes, with 12 cycles of PCR. The quality control for the samples was done by sequencing
a pool of samples using MiSeq V2 nano kit - 300 cycles. After library amplification, 3 μL of
each library was pooled into a library pool. The pool was then cleaned up using SPRI beads
following the Nextera Flex clean-up and size selection protocol. The pool was then sequenced
using a MiSeq V2 nano kit (Illumina, USA). Based on the sequencing data generated, the read
count for each sample was used to identify the failed libraries (i.e., libraries with less than 100
reads).
791
Moreover, libraries were pooled at different amounts based on the read count to ensure equal
representation in the final pool. The final pool was sequenced on Illumina NovaSeq 6000 Xp
S4 lane, 2 × 150 bp. WGS read quality was assessed using FASTQC (version 0.11.5) and
trimmed using Trimmomatic (version 0.36) with default parameters and trimmed of adaptor
sequences (TruSeq3 paired-ended). Reads were aligned to the *E. coli* MG1655 genome
(http://bacteria.ensembl.org/Escherichia_coli_str_k_12_substr_mg1655_gca_000005845/Info
/Index/, assembly ASM584v2) and then analysed variants following GATK Best Practices for
Variant Discovery (HaplotypeCaller) (64). Further genome variant annotation was conducted
using the software SnpEff (65).
801
Global transcriptome sequencing
After ampicillin treatment for 0 and 8, surviving isolates were immediately washed and
harvested for global transcriptome sequencing. Total RNA was extracted from the cell pellets
using a PureLink RNA mini kit (Invitrogen) as per the manufacturer's instructions. The global
transcriptome sequencing was processed and analysed by Genewiz, Jiangsu, China. Primers
used in this work are listed in Table S4. RNA-Seq read quality was assessed using FASTQC
and trimmed using Trimmomatic with default parameters. Reads were aligned to the *E. coli*
MG1655 genome
(http://bacteria.ensembl.org/Escherichia_coli_str_k_12_substr_mg1655_gca_000005845/Info
/Index/, assembly ASM584v2) and then counted using the RSubread aligner with default
parameters (66). After mapping, differential expression analysis was carried out using strand-
specific gene-wise quantification using the DESeq2 package (67). Further normalisation was
conducted using RUVSeq and the RUV correction method, with $k = 1$ to correct for batch
effects, using replicate samples to estimate the factors of unwanted variation (68). Absolute
counts were transformed into standard z-scores for each gene over all treatments, that is,

absolute read for a gene minus mean read count for that gene over all samples and then divided

by the standard deviation for all counts over all samples. Genes with an adjusted P value

(P_{adj}) of ≤0.05 were considered differentially expressed. PseudoCAP analysis was conducted

by calculating the percentage of genes in each classification that were differentially expressed

($\log_2$FC ≥ ±2, P_{adj} ≤ 0.05).

ROS measurement

Intracellular ROS accumulation was measured using the Cellular ROS Assay Kit (Abcam,

ab113851) according to the manufacturer's protocol. Overnight cultures of both wild type and

Δ*recA* strains were grown in LB medium. Cells were treated with 50 µg/mL ampicillin for 8

hours at 37°C with shaking (250 rpm). After the treatment, cells were washed twice with 1x

buffer to remove residual antibiotics and debris. The collected cells were resuspended in 1x

buffer and incubated with the fluorescent ROS probe DCFDA (2',7'–dichlorofluorescin

diacetate)/H2DCFDA at a final concentration of 10 µM for 30 minutes at 37°C in the dark to

prevent probe degradation.

After incubation, 500 µL of the stained cells were immediately analysed using a CytoFLEX

LX flow cytometer (Beckman Coulter). Fluorescence was detected in the FITC channel

(excitation at 488 nm, emission at 525 nm). Data were analysed using FlowJo software

(Version X, Tree Star Inc.), with fluorescence intensity serving as an indicator of intracellular

ROS levels. Appropriate controls, including unstained cells and cells without ampicillin

treatment, were included to ensure accurate ROS measurement.

Single-molecule localisation imaging and data analysis

Single-molecule localisation imaging was performed on a custom-built Stochastic Optical

Reconstruction Microscope (STORM) with an Olympus IX81 microscope frame, a 100x

magnification NA 1.45 objective (Olympus) and an EMCCD camera (DU-897, Andor) as

described previously (69-71). In summary, 35 mm cell culture dishes (0.17 mm No.1

coverglass) were cleaned with 1 M KOH for 30 minutes in an ultrasonic cleaning machine,

followed by three washes with MilliQ water. The dishes were air-dried with high-purity

nitrogen blowing and sterilised by UV exposure for 30 minutes. *E. coli* cells were fixed with

NaPO4 (30 nM), formaldehyde (2.4%), and glutaraldehyde (0.04%) at room temperature for

15 minutes, followed by 45 minutes on ice. Samples were then centrifuged to collect the pellet

cells, and the supernatant was discarded. Cell pellets were washed twice with phosphate-buffered saline (PBS), pH 7.4. Cells were resuspended in 200 μL of GTE buffer and kept on ice until 200 μL was placed onto the coverslip bottom of the cleaned 35 mm culture dish. To label the bacterial chromosome, a Click-iT EdU kit was used prior to fixation following the manufacturer's instruction (ThermoFisher) and as described before. To label DNA polymerase I, fixed cells were blocked and permeabilised with blocking buffer (5% wt/vol bovine serum albumin (Sigma-Aldrich) and 0.5% vol/vol Triton X-100 in PBS) for 30 min and then incubated with 1 μg/mL primary antibody against DNA polymerase I (ab188424, Abcam) in blocking buffer for 60 min at room temperature. After washing with PBS three times, the cells were incubated with 2 μg/mL fluorescently labelled secondary antibody (Alexa 647, A20006, ThermoFisher) against the primary antibody in the blocking buffer for 40 min at room temperature. After washing with PBS three times, the cells were postfixed with 4% (wt/vol) paraformaldehyde in PBS for 10 min and stored in PBS before imaging. STORM image analysis, drift correction, image rendering, protein cluster identification and images presentation were performed using Insight342, custom-written Matlab (2012a, MathWorks) codes, SR-Tesseler (IINS, Interdisciplinary Institute for Neuroscience) (72), and Image J (National Institutes of Health).

Statistical analysis

Statistical analysis was performed using GraphPad Prism v.9.0.0. All data are presented as individual values and mean or mean ± SEM. Unless otherwise specified, statistical comparisons were conducted using one-way ANOVA (for multiple groups) or unpaired two-tailed Student's t-tests (for two-group comparisons), assuming a 95% confidence interval.. A probability value of $P < 0.05$ was considered significant. Statistical significance is indicated in each figure. All remaining experiments were repeated independently, at least six with similar results.

Data availability

Sequence data supporting this study's findings have been deposited in the GEO repository with the GEO accession number GSE179434.

881 **Table S1. Other mutations detected in the *ΔrecA* resistant isolates**

Gene	Mutation	Genomic Position	Animo Acid Changes
stfE	A>AGGTTTTCGAGAGC	1209618	p.Val13fs
puuC	T > G	1363695	p.Phe318Cys
cpxA	T > C	4104711	p.Thr89Ala

883 **Table S2. Strains used in this study**

Strain	Relevant Genotype	Parent strain	Source
DH5α	-	-	Lab stock
E. coli MG1655	$recA^+ lexA^+$	-	Lab stock
RW1570	$recA^+ lexA3$	MG1655	Lab stock
JW0612-1	$recA^+ lexA^+ \Delta citB$	MG1655	Keio Collection
JW0611-1	$recA^+ lexA^+ \Delta citA$	MG1655	Keio Collection
JW1134-1	$recA^+ lexA^+ \Delta ymfm$	MG1655	Keio Collection
JW0059-1	$recA^+ lexA^+ \Delta polB$	MG1655	Keio Collection
JW0221-1	$recA^+ lexA^+ \Delta dinB$	MG1655	Keio Collection
RW120	$recA^+ lexA^+ \Delta umuDC$	MG1655	Lab stock
EAW13	$recA^+ lexA^+ \Delta sulA$	MG1655	Lab stock
JW2669-1	$lexA^+ \Delta recA$	MG1655	Keio Collection

885 **Table S3. Plasmids used in this study**

Plasmid	Features	Source
pJM1071-*recA*	Low copy vector with constitutive *recA* promoter and native *recA* RBS; *recA* cloned in pJM1071 between *NdeI/XbaI*	Lab stock (59)
pKD46	Temperature sensitive replication (*repA101ts*); encodes lambda Red genes	Lab stock
pCP20	Temperature sensitive origin of replication; encodes the FLP recombinase.	Lab stock

887 **Table S4. Primers used in this study**

Name	Sequence	Function

recA-FWD	AAAAAAGCAAAAGGGCCGCAGATGCGACCCTTGT GTATCAAACAAGACGAGAAACGAGAGAGGATGCT CAC	Construction of recA deletion mutant
recA-REV	CAACAGAACATATTGACTATCCGGTATTACCCGGC ATGACAGGAGTAAAAGACGTCTAAGAAACCATTA TTATCATGAC	

Figure S1. Long-term exposures to ampicillin induced the evolution of resistance in the

wild type and *ΔrecA* E. coli strain. (A) The experimental flow of ALE antibiotic treatment

experiment. An overnight culture (0.6 mL; 1 x 10⁹ CFU/mL cells) was diluted 1:50 into 30 mL

LB medium supplemented with 50 μg/mL ampicillin and incubated at 37ºC with shaking at

250 rpm for 4.5 hours. After treatment, the antibiotic-containing medium was removed by

washing twice (20 min centrifugation at 1500 g) in fresh LB medium. The surviving isolates

were resuspended in 30 mL LB medium and grown overnight at 37ºC with shaking at 250 rpm.

Ampicillin treatment was applied to the regrown culture and repeated until resistance was

established. **(B)** Changes in the MICs of ampicillin in the wild type and *ΔrecA* strain after 21

days of treatment with ampicillin at 50 μg/ml for 4.5 hours each day. MICs were measured

after each daily treatment. Each experiment was independently repeated at least twice, and the

data are shown as mean ± SEM.

Figure S2. The survival rate after a single exposure to ampicillin at 50 μg/ml for 0,2,4,6, and

8 hours in the wild type and *ΔrecA* strain.

907
Figure S3. Single exposures to ampicillin induced the evolution of resistance in the
ΔrecA[CGSC] **strain (JW2669-1). (A)** MICs of ampicillin were measured against the wild
type[CGSC] *E coli* strain after single exposures to ampicillin. **(B)** MICs of ampicillin were
measured against the *ΔrecA*[CGSC] strain after single exposures to ampicillin. Each experiment
was independently repeated at least six times using parallel replicates, and the data are shown
as mean ± SEM. Significant differences among different treatment groups are analyzed by
independent t-test, $*P < 0.05$, $**P < 0.01$, $***P < 0.001$.
915

Figure S4. Single exposures to other β-lactam antibiotics induced the evolution of
resistance in the *ΔrecA* strain. (A) MICs of penicillin G in the wild type, *ΔrecA*, and
complemented *ΔrecA* strain, where the expression of RecA was restored, after single exposures
to penicillin G at 1 mg/mL for 8 hours. **(B)** MICs of carbenicillin in the wild type, *ΔrecA*, and
complemented *ΔrecA* strain, where the expression of RecA was restored, after single exposures
to carbenicillin at 200 µg/mL for 8 hours. Each experiment was independently repeated at least
six, and the data are shown as mean ± SEM.

Fig. S5. Distribution fitting and fluctuation analysis support a selection-driven resistance mechanism in the *ΔrecA* strain. (A) Rifampicin-resistant mutant counts from replicate cultures were fitted to theoretical Poisson distributions based on observed means. **(B)** Fluctuation analysis using the Luria-Delbrück framework revealed long-tailed, highly skewed mutation count distributions in the *ΔrecA* strain under the treatment of ampicillin, characteristic of jackpot cultures arising from early-arising resistant mutants.

934
Figure S6. The activity of β-lactamase was increased in the *ΔrecA* culture supernatants.
The *ΔrecA* strain was treated with ampicillin at 50 μg/ml for 8 hours, and surviving isolates
harbouring the *ampC* mutations were selected. The level of β-lactamase in cell culture
supernatants was determined by the absorbance at OD_{490}. The levels of β-lactamase in the wild
type or *ΔrecA* strain culture supernatants without exposure to ampicillin were tested as a control.
Each experiment was independently repeated six times. Each experiment was independently
repeated at least six, and the data are shown as mean ± SEM.

Figure S7. Transcriptional responses of the wild type and *ΔrecA* strain after single treatments with ampicillin for 8 hours. GO analysis was performed following the GOseq approach, and different genes in the wild type **(A, left)** and the *ΔrecA* strain **(B, left)** were plotted. KEGG pathway enrichment was assigned according to the KEGG database, and different genes in the wild type **(C, left)** and the *ΔrecA* strain **(D, left)** were plotted. The top 20 enrichment pathways are listed in the GO and KEGG enrichment analysis **(A-D, right)**. Total RNA-seq was performed with three repeats in each group.