1   **PinR as a Stress Inducible Inhibitory Transcription Factor in Stopping Stress-induced**
2   **Mutagenesis Bacterial Evolution of Antibiotic Resistance**
3
4

5   Le Zhang[1,†], Yunpeng Guan[1,2,†], Qian Peter Su[1], Yan Liao[3], Iain Duggin[3], Dayong Jin[1,2]*
6
[1]Institute for biomedical materials and devices, University of Technology Sydney, Ultimo,
Australia.
[2]Department of biomedical engineering, Southern University of Science and Technology,
Shenzhen, China.
[3]iThree institute, University of Technology Sydney, Ultimo, Australia.
†These authors contributed equally to this work
*Corresponding author. Email: dayong.jin@uts.edu.au

PinR is a lambdoid prophage Rac existing in many different bacteria, but its function is
yet to be discovered. Here, we report PinR as an effective "anti-evolution" target towards
combating antibiotic resistant bacteria. We show the function of PinR as a stress
inducible inhibitory transcription factor in the bacterial evolution of antibiotic resistance,
which represses the transcriptions of many oxidative stress response genes and
consequently orchestrate the ROS burst-induced evolution of antibiotic resistance. We
show that deletion of *pinR* gene can avoid the accelerated evolution of antibiotic resistance
induced by the deficiency of DNA repair, while the strategies to target RecA, e.g., the
deletion of *recA* gene, can be safely implemented to disarm the bacterial resistances to a
variety of antibiotics.

Since the first antibiotic penicillin was discovered in 1940s (1), antibiotics are used to prevent
and treat the bacterial infections. Antibiotics have been designed to target the cell wall or
membranes, the bacterial DNA or RNA, and the machinery that produces proteins (2).
However, the antibiotic resistance can occur naturally along with the administration of
antibiotics, and the misuse of antibiotics is accelerating this process which poses a potentially
catastrophic threat to the public health globally (3, 4).
32

Bacterial antibiotic resistance can arise by either *de novo* mutations or bacterial recombination,
in which the DNA recombinases catalyse site-specific recombination events within DNA and

promote the spread of antibiotic resistant genes (3, 4). The putative site-specific recombinase

PinR is a lambdoid prophage Rac encoded by the gene *pinR* (or named *ynaD*) and exists in

many bacterium types, including the *Escherichia coli* (*E. coli), Tetragenococcus halophilus,*

Lactococcus lactis, Salmonella enterica and Streptococcus (5-8). Although the function of

PinR is yet to be clear, it has been shown that PinR shares a similar structure with the DNA

invertase (Fig. S1) and can catalyse an inversion of a 177-bp DNA fragment in the

Streptococcus acting as a bacterial recombinase (6). Meanwhile, using MobileElementFinder,

the *pinR* is predicted as a mobile genetic element carrying the antibiotic resistant genes (7),

which has been experimentally evidenced in *L. pentosus* plasmid (8).

The SOS response pathway, involving its master regulator RecA, induces the DNA repair,

which consequently helps to drive the development and spreading of antibiotic resistance

determinants (9-11). Therefore, deactivating the RecA may disarm the bacterial resistance to

antibiotics (12). However, as a double-edged sword, the deficiency of DNA repair, as a result

of *recA* deletion, may also increase the rate of drug resistant mutagenesis induced by the

antibiotic exposure towards the antibiotic resistance (13, 14). Notably, our recent work reports

that the deletion of RecA can cause an unexpected superfast establishment of antibiotic

resistance in *E. coli ΔrecA* strain (15), but its mechanism is unknown.

Here, we find that, in the *ΔrecA* strain, further deletion of *pinR* not only prevented the

emergence of resistance after a single exposure to ampicillin (Fig. 1A), but also up to 3-week

intermittent treatment of ampicillin was unable to induce the evolution of resistance (Fig. 1B).

This was in a clear contrast to the commonly occurred resistance in the wild type strain treated

with an experimental evolution model that mimics the administration of antibiotics in clinic.

This suggests that targeting PinR offers a way out of the dilemma whether or not to target RecA

to suppress the SOS response, towards preventing the antibiotic resistance. As the deletion of

recA alone causes the DNA repair deficiency and consequently increases the rate of antibiotic

exposure-induced resistant mutagenesis that facilitates the antibiotic resistance, we further

sequenced the surviving isolates in *ΔrecA*/*ΔpinR* strain and confirmed that the drug resistant-

associated gene mutations were not detected in the *ΔrecA*/*ΔpinR* strain after neither the single

exposure to ampicillin for 8 hours nor the 3-week intermittent treatment with ampicillin (Fig.

1C). This further suggests the strategies to target both RecA and PinR may stop the bacteria

from developing the long-term antibiotic resistance.

To further understand the role of *pinR* and how the deletion of *pinR* can prevent the *ΔrecA*
strain from antibiotic resistance, we first compared the global transcriptome analysis in the
wild type and the *ΔrecA* surviving isolates after the single exposure to ampicillin. The treatment
of ampicillin affected the transcriptomic profile of both in the wild type and the *ΔrecA* strain,
compared with that of untreated control cells, with changes to the expression of 4373 and 4286
coding sequences, respectively (Fig. 1D). Venn diagram show the differentially expressed
genes with a $\log_2$ fold change ($\log_2$FC) of $\geq \pm 2$, and principal component analysis (PCA)
confirmed that the effect of ampicillin exposure on the *ΔrecA* strain remarkably differed from
that of the wild type cells (Fig. S2). Genome-wide expression changes were visualized as
volcano plots (Fig. S3) to identify specific genes with large fold changes and statistical
significance. With Gene Ontology (GO) enrichment analysis and Kyoto Encyclopedia of Genes
and Genomes (KEGG) pathway analysis shown in Fig. S4. We compared the selected
functional groups corresponding to biological processes, including DNA recombination,
quorum-sensing system, SOS response and ABC transport system (Fig. 1E and Fig. S5), and
found it is of much significance that a series of genes associated with the DNA damage and
repair were slightly changed, but the transcription of the gene *pinR* was considerably
upregulated (10-fold) in the *ΔrecA* resistant populations compared with that of the ampicillin-
treated wild type strain (Fig. 1E). These results collectively reveal that the PinR is a stress
inducible protein, and therefore as a new "evolution potentiator" of the antibiotic resistance

We further studied the oxidative stress response, which was uniquely regulated in the *ΔrecA*
strain and may cause the evolution of antibiotic resistance, as the induction of mutagenesis can
be stimulated by the production of reactive oxygen species (ROS) throughout the
administration of antibiotics in a few weeks (16). The transcriptome analysis revealed that the
single treatment of the *ΔrecA* strain with the ampicillin caused a significant downregulation of
transcription of many oxidative stress response genes, including the genes *cysJ, cysI, cysH,*
soda and *sufD* (Fig. 2A), indicating the upregulation of ROS after the single exposure to
ampicillin in the *ΔrecA* strain. By adding 50 mM glutathione (GSH), a natural antioxidant, into
the *ΔrecA* culture medium and exposing them to 50μg/ml ampicillin for 8 hours, we observed
that the antioxidant prevented the emergence of resistance to ampicillin in the *ΔrecA* strain (Fig.
2B) but did not affect the survival rate (Fig. 2C), which suggested that the overproduction of
ROS was not a lethal effect induced by the single exposure to ampicillin in the *ΔrecA* strain.
We sequenced the survived isolates and found that the addition of GSH prevented the DNA
mutations associated with the drug resistance in the *ΔrecA* strain (Fig. 2D). These data

demonstrated that the ROS burst happened as the result of the suppression on the oxidative stress response system, and consequently triggered the resistant mutations, which caused the superfast emergence of resistance in the *ΔrecA* strain. More importantly, by further comparing the transcription of oxidoreductase genes in the *ΔrecA/ΔpinR* strain, we did not find the changes of the transcription of any oxidative stress response genes in the *ΔrecA/ΔpinR* strain after the single exposure to ampicillin (Fig. 2A). These results, taking together, indicated that the PinR services as an inhibitory transcription factor repressing on the transcription of oxidoreductase genes, and the absence of PinR stops the evolution of antibiotic resistance caused by the overaccumulation of ROS.

In short, we conclude PinR as an inhibitory transcription factor in the pathway indicated in Fig. 2E, as the transcription of *pinR* can be upregulated in the *ΔrecA* strain under the treatment of ampicillin. Although the involvement of ROS in the evolution of antibiotic resistance has been identified, we first find that the upregulation of PinR can orchestrate the ROS burst induced mutagenesis through the repression on the oxidative stress response system, which consequently facilitates the bacterial evolution of antibiotic resistance. From a drug development perspective, strategies to directly suppress the transcription of 'potentiator' genes or to inhibit their encoding proteins activities may combat the antibiotic resistance (8). Our result shows the deletion of *pinR* can dramatically delay the evolution of resistance to ampicillin in both the short-term and long-term models. With the role of PinR being clear, our evolutionary model can be generic, as the overproduction of ROS is a common action upon the treatment of many antibiotics (16), maintaining the transcription of oxidative stress response genes by inhibiting the transcription of 'potentiator' genes will become a direction to stop the stress-induced mutagenesis that is otherwise unavoidably induced by the treatments of a variety of antibiotic drugs. Therefore, our work suggests future studies on the 'potentiator' genes towards the safe and more effective strategies to be developed for infectious disease treatments.

C

Antibiotic exposure	Gene	Mutations	Genomic Position	Phenotype
8 hours	P_{ampC}	G > A	4379014	$\Delta recA^{Resistance}$
8 hours	P_{ampC}	A > T	4379035	$\Delta recA^{Resistance}$
8 hours	acrB	G > A	482581	$\Delta recA^{Resistance}$
8 hours	acrB	C > T	482387	$\Delta recA^{Resistance}$
3 weeks	P_{ampC}	G > A	4379014	Wild type^{Resistance}
3 weeks	P_{ampC}	+ TA	4379017	Wild type^{Resistance}
3 weeks	P_{ampC}	G > A	4379014	$\Delta recA^{Resistance}$
-	-	-	-	$\Delta recA/\Delta pinR^{Sensitive}$

Figure 1. PinR as a new "evolution potentiator" of the antibiotic resistance. (A) Deletion of *pinR* delayed the evolution of antibiotic resistance after an intermittent treatment with ampicillin (50 μg/ml) for 21 days in the *ΔrecA/ΔpinR* strain. **(B)** Deletion of *pinR* prevented the evolution of antibiotic resistance in the *ΔrecA/ΔpinR* strain after a single exposure to ampicillin at 50 μg/ml for 8 hours. **(C)** DNA mutations detected in the wild type, *ΔrecA*, and *ΔrecA/ΔpinR* strains after a single exposure to ampicillin (50 μg/ml) for 8 hours or an intermittent treatment of ampicillin at 50 μg/ml for 21 days. **(D)** The clustered heatmap of relative expression of coding sequences in the wild type and the *ΔrecA* strain. **(E)** After the exposure to ampicillin at 50 μg/ml for 8 hours, the regulation of transcription of DNA recombination and repair genes. The global transcriptome sequencing was performed twice.

Other experiments were independently repeated at least 4 times. $*\,p < 0.05$; $**\,p < 0.01$; $***$

$p < 0.001$.

Figure 2. PinR orchestrates the ROS burst induced mutagenesis facilitating the bacterial

evolution of antibiotic resistance. (A) After the exposure to ampicillin at 50 μg/ml for 8 hours,

the regulation of the transcription of oxidative stress response genes in the wild type, *ΔrecA*,

and the *ΔrecA/ΔpinR* strain. **(B)** Addition of 50 mM antioxidative compound GSH prevented

the evolution of antibiotic resistance in the *ΔrecA* strain. **(C)** Survival fraction after the

exposure to ampicillin in the wild type and the *ΔrecA* strain with or without the addition of

GSH. **(D)** DNA mutation detection after the single treatment of ampicillin (50 μg/ml) in the

ΔrecA strain with or without the addition of GSH. **(E)** The mechanism by PinR mediating the

bacterial evolution of antibiotic resistance. The global transcriptome sequencing was

performed twice. Other experiments were independently repeated at least 4 times. *$p < 0.05$;

** $p < 0.01$; *** $p < 0.001$.*

Materials and Methods

Bacterial strains, medium and antibiotics

Bacterial strains and plasmids used in this work are described in Table S1 and Table S2. Luria-Bertani (LB) was used as broth or in agar plates. *E. coli* cells were grown on LB agar (1.5% w/v) plates at 37ºC, unless stated otherwise, antibiotics were supplemented, where appropriate. Whenever possible, antibiotic stock solutions were prepared fresh before the use.

Treatment with antibiotics to induce evolutionary resistance

For the single exposure to antibiotic experiment, an overnight culture (0.6 ml; 1 x 10^9 CFU/ml cells) was diluted 1:50 into 30 ml LB medium supplemented with antibiotics (50 μg/ml ampicillin, 1 mg/ml penicillin G, or 200 μg/ml carbenicillin) and incubated at 37ºC with shaking at 250 rpm for 0 or 8 hours, respectively. After each treatment, the antibiotic-containing medium was removed by washing twice (20 min centrifugation at 1500 g) in fresh LB medium.

To test resistance, the surviving isolates were first resuspended in 30 ml LB medium and grown overnight at 37ºC with shaking at 250 rpm. The regrown culture was then plated onto LB agar supplemented with the appropriate selective antibiotics and incubated 16 hours at 37ºC. Single colonies were isolated and used to test the resistance or stored at -80ºC for future use.

For the intermittent antibiotic treatment experiments, an overnight culture (0.6 ml; 1 x 10^9 CFU/ml cells) was diluted 1:50 into 30 ml LB medium supplemented with 50 μg/ml ampicillin and incubated at 37ºC with shaking at 250 rpm for 4 hours. After treatment, the antibiotic-containing medium was removed by washing twice (20 min centrifugation at 1500 g) in fresh LB medium. The surviving isolates were resuspended in 30 ml LB medium and grown overnight at 37ºC with shaking at 250 rpm. The killing treatment was applied as above to the regrown culture and repeated until resistance was established.

Antibiotic susceptibility testing

The susceptibility of *E. coli* cells to antibiotics was measured by using minimum inhibitory concentration (MIC) testing (17).

Construction of deletion mutants

190   Lambda Red recombination was used to generate various gene deletions in *E. coli* strains
191   followed by previous reported methods with modifications (18, 19). Primers used in this work
192   are listed in the Table S3.
193
194   **DNA extraction**
195   Chromosomal DNA was extracted and purified using the PureLink™ Genomic DNA mini kit
196   (ThermoFisher Scientific). Plasmid DNA was extracted and purified using the PureLink™
197   Quick Plasmid Miniprep kit (ThermoFisher Scientific).
198
199   **Whole genome sequencing**
200   The genomic sequencing was conducted following the Nextera Flex library preparation kit
201   process (Illumina), and processed by Sangon Biotech, Shanghai, China.
202
203   **Global transcriptome sequencing**
204   The global transcriptome sequencing was processed and analysed by Genewiz, Jiangsu, China.
205   Primers used in this work are listed in the Table S3.
206
Statistical analysis
Statistical analysis was performed using GraphPad Prism v.9.0.0. All data are presented as
individual values and mean or mean ± s.e.m. A two-tailed unpaired Student's t-test using a 95%
confidence interval was used to evaluate the difference between two groups. For more than two
groups, a one-way ANOVA was used. A probability value of $P < 0.05$ was considered
significant. Statistical significance is indicated in each figure. All remaining experiments were
repeated independently at least fourth with similar results.
214

Figure S1. The proposed structure of PinR compared with the site-specific DNA recombinase and DNA invertase.

Figure S2. Transcriptional response of individual strain treated with ampicillin for 8 hours. (A) Venn diagram of differentially expressed genes ($\log_2$FC $\geq \pm 2$). (B) Bi-plot of the principal-component analysis of normalized read counts for all strains.

Figure S3. Summary of genome-wide expression changes in the wild type strain (A) and the
ΔrecA strain (B) after the single exposure to ampicillin. The top 10 most differentially
expressed genes are labelled in each plot. Gray dots indicate genes with no significant
difference compared to the untreated control (P > 0.05), blue dots indicate genes with a
significant difference compared to the untreated control (P < 0.05), and red dots indicate genes
with both a significant difference (P < 0.05) and $\log_2$FC of > 2 compared to the untreated
control, with numerical annotations to indicate the number of differentially expressed genes.

Figure S4. Transcriptional response of the wild type and *ΔrecA* strains after the exposure

to ampicillin at 50 μg/ml for 8 hours. GO enrichment analysis of different genes including in

biological process (red), cellular component (green), and molecular function (white) in the wild

type strain (A) and the *ΔrecA* strain (B). KEGG pathway enrichment scatter plot of different

genes in (C) the wild type strain and the *ΔrecA* strain (D). Top 20 enrichment pathways were

listed in the GO and KEGG enrichment analysis.

Figure S5. Heatmaps show log₂FC data for each strain treated with single exposure to
$$\text{log}_2\text{FC}$$

Figure S5. Heatmaps show log$_2$FC data for each strain treated with single exposure to ampicillin at 50 μg/ml for 8 hours. (A) Quorum-sensing system genes. (B) SOS response genes. (C) ABC transport system genes.

Table S1. Strains used in this study.

Strain	Relevant Genotype	Parent strain	Source/technique
DH5α	-	-	Lab stock
E. coli K-12	*recA⁺ pinR⁺*	-	Lab stock
LZ101	*ΔrecA::Tet*	*K-12*	Lab stock
LZ102	*ΔrecA::Tet, ΔpinR::Kan*	LZ101	This study

Table S2. Plasmids used in this study.

Plasmid	Description	Source
pKD46	Helper plasmid for Lambda RED recombination	Lab stock

Table S3. Primers used in this study.

Name	Sequence
recA-FWD	AAAAAAGCAAAAGGGCCGCAGATGCGACCCTTGTGTATCAAAC AAGACGAGAAACGAGAGAGGATGCTCAC
recA-REV	CAACAGAACATATTGACTATCCGGTATTACCCGGCATGACAGGA GTAAAAGACGTCTAAGAAACCATTATTATCATGAC
pinR-FWD	CAAGGAAGGGGGCTTGGAAGACGTAAAGCATCTCACACCGAGA TTATTTTCATATGAATATCCTCCTTAGTTCCTATTC
pinR-REV	TTTCTGAGATGCATTATGATATGAACACCAATTTCGTATAGAGTC TCACTGAGCTGCTTCGAAGTTCCTA
P5 adapter	AGATCGGAAGAGCGTCGTGTAGGGAAAGAGTGT
P7 adapter	AGATCGGAAGAGCACACGTCTGAACTCCAGTCA

Reference

1. E. P. Abraham, and E. Chain. An Enzyme from Bacteria able to Destroy Penicillin. *Nature*, **146**, 837 (1940).

2. Michael A. Kohanski, Daniel J. Dwyer, and James J. Collins. How antibiotics kill bacteria: from targets to networks. *Nature Review Microbiology*. **8**, 423-435 (2010).

3. Blair J. M, Webber M. A, Baylay A. J, Ogbolu D. O, Piddock L. J. Molecular mechanisms of antibiotic resistance. *Nature Review Microbiology*. **13,** 1:42-51 (2015).

4. Walsh C. Molecular mechanisms that confer antibacterial drug resistance. *Nature*. **17**, 406:775-81 (2000).

5. Butland G, Peregrín-Alvarez JM, Li J, Yang W, et al. Interaction network containing conserved and essential protein complexes in *Escherichia coli*. *Nature*. **433**, 7025:531-537 (2005).

6. Steward KF, Harrison T, Robinson C, Slater J, et al. PinR mediates the generation of reversible population diversity in *Streptococcus zooepidemicus*. *Microbiology*. **161**, 5:1105-1112 (2015).

7. Johansson MHK, Bortolaia V, Tansirichaiya S, et al. Detection of mobile genetic elements associated with antibiotic resistance in *Salmonella enterica* using a newly developed web tool: MobileElementFinder. *J Antimicrob Chemother*. **76**, 1:101-109 (2021).

8. Abriouel H, Pérez Montoro B, de la Fuente Ordoñez JJ, et al. New insights into the role of
plasmids from probiotic *Lactobacillus pentosus* MP-10 in Aloreña table olive brine
fermentation. *Sci Rep*. **9**, 1:10938 (2019).

9. Darja Žgur-Bertok. DNA Damage Repair and Bacterial Pathogens. *PLOS Pathogens.* **9**
PLOS Pathogens, e1003711 (2013).

10. Dale B. Wigley. Bacterial DNA Repair: Recent Insights into The Mechanism of
RecBCD, AddAB and AdnAB. *Nature Reviews Microbiology*. **11**, 9-13 (2013).

11. Jason C. Bell, Stephen C. Kowalczykowski. RecA: Regulation and Mechanism of a
Molecular Search Engine. *Trends in Biochemical Sciences*. **41**, 491-507 (2016).

12. Athanasia Pavlopoulou. RecA: A Universal Drug Target in Pathogenic Bacteria.
Frontiers in Bioscience. **23**, 36-42 (2018).

13. Jiafeng Liu, Orit Gefen, Irine Ronin, Maskit Bar-Meir, Nathalie Q Balaban. Effect of
tolerance on the evolution of antibiotic resistance under drug combinations. *Science*. **367**,
200-204 (2020).

14. Irit Levin-Reisman et al. Antibiotic Tolerance Facilitates the Evolution of Resistance.
Science. **355**, 826-830 (2017).

15. Le Zhang et al. Superfast Evolution of Antibioitc Resistance in *Escherichia coli*. (2021).
in submission.

16. Michael A Kohanski, Mark A DePristo, James J Collins. Sublethal antibiotic treatment
leads to multidrug resistance via radical-induced mutagenesis. *Mol. Cell*. **37**, 311-320
(2010).

17. Eric M. Scholar, William B. Pratt. *The Antimicrobial Drugs*. Oxford University Press.

18. Kirill A. Datsenko, Barry L. Wanner. One-Step Inactivation of Chromosomal Genes in
Escherichia Coli K-12 Using PCR Products. *Proc Natl Acad Sci*. **97**, 6640-6645 (2000).

19. Tomoya Baba et al. Construction of *Escherichia Coli* K-12 In-Frame, Single-Gene
Knockout Mutants: The Keio Collection. *Mol Syst Biol*, **2**, 2006.0008 (2006).