**Phylogenomics Illuminates the Evolutionary History of Wild Silkmoths in Space and Time (Lepidoptera: Saturniidae)**

Rodolphe Rougerie[1*£], Astrid Cruaud[2£], Pierre Arnal[1,2£], Liliana Ballesteros-Mejia[1,3], Fabien L. Condamine[4], Thibaud Decaëns[5], Marianne Elias[1], Delphine Gey[6,7], Paul D. N. Hebert[8], Ian J. Kitching[9], Sébastien Lavergne[10], Carlos Lopez-Vaamonde[11,12], Jérôme Murienne[13], Yves Cuenot[13], Sabine Nidelet[2], Jean-Yves Rasplus[2]

[1] *Institut de Systématique, Évolution, Biodiversité (ISYEB), Muséum national d'Histoire naturelle, CNRS, EPHE, Sorbonne Université, Université des Antilles, Paris, France (rodolphe.rougerie@mnhn.fr)*

[2] *Centre de Biologie pour la Gestion des Populations (CBGP), INRAE, CIRAD, IRD, Montpellier SupAgro, Université de Montpellier, Montpellier, France*

[3] *CESAB, Centre de Synthèse et d'Analyse sur la Biodiversité, Montpellier, France*

[4] *Institut des Sciences de l'Évolution de Montpellier (ISEM), CNRS, Université de Montpellier, Montpellier, France*

[5] *Centre d'Écologie Fonctionnelle et Évolutive (CEFE), Université de Montpellier, CNRS, EPHE, IRD, Montpellier, France*

[6] *Service de Systématique Moléculaire, UAR 2700 2AD, Muséum national d'Histoire naturelle, Paris, France*

[7] *Molécules de Communication et Adaptation des Microorganismes (MCAM), Muséum national d'Histoire naturelle, CNRS, Paris, France*

[8] *Center for Biodiversity Genomics, University of Guelph, Guelph, Canada*

[9] *Department of Life Sciences, Natural History Museum, London, United Kingdom*

[10] *Université Grenoble Alpes, CNRS, Université Savoie Mont Blanc, Laboratoire d'Écologie Alpine*

*(LECA), Grenoble, France*

[11] *Institut de Recherche sur la Biologie de l'Insecte, UMR7261 CNRS - Université de Tours, France*

[12] *INRAE, UR633, Zoologie Forestière, F-45075 Orléans, France*

[13] *Laboratoire Évolution et Diversité Biologique (EDB UMR5174), Université Toulouse 3 Paul*

*Sabatier, CNRS, IRD, Toulouse, France*

[*]corresponding author

[£]equal contributors

**ABSTRACT**

Wild silkmoths (Saturniidae) are one of the most emblematic and most studied families

of moths. Yet, the absence of a robust phylogenetic framework based on a comprehensive

taxonomic sampling impedes our understanding of their evolutionary history. We analyzed

1,024 ultraconserved elements (UCEs) and their flanking regions to infer the relationships

among 338 species of Saturniidae representing all described subfamilies, tribes, and genera.

We investigated systematic biases in genomic data and performed dating and historical

biogeographic analyses to reconstruct the evolutionary history of wild silkmoths in space and

time. Using Gene Genealogy Interrogation, we showed that saturation of nucleotide sequence

data blurred our understanding of early divergences and first biogeographic events. Our

analyses support a Neotropical origin of saturniids, shortly after the Cretaceous-Paleogene

extinction event (*ca* 64.0 [stem] - 52.0 [crown] Ma), and two independent colonization events

of the Old World during the Eocene, presumably through the Bering Land Bridge. Early

divergences strongly shaped the distribution of extant subfamilies as they showed very limited

mobility across biogeographical regions, except for Saturniinae, a subfamily now present on

all continents but Antarctica. Overall, our results provide a framework for in-depth

investigations into the spatial and temporal dynamics of all saturniid lineages and for the

integration of their evolutionary history into further global studies of biodiversity and

conservation. Rather unexpectedly for a taxonomically well-known family such as

Saturniidae, the proper alignment of taxonomic divisions and ranks with our phylogenetic

results leads us to propose substantial rearrangements of the family classification, including

the description of one new subfamily and two new tribes.

**KEYWORDS**: Historical Biogeography; Moths; Phylogenomics; Saturation; Taxonomy; Ultra-

Conserved Elements

Thanks to generations of naturalists, moths in the family Saturniidae – also known as

wild silkmoths – are now among the best documented insects, particularly because of their

spectacular variation in size and form in both adult and immature stages. The family

encompasses some of the largest (e.g., *Coscinocera hercules* and *Attacus* species that reach

30cm wingspan) and most emblematic moths (e.g., the moon moths, *Actias* species and the

comet moth, *Argema mittrei*). Wild silkmoths also bear importance to humans for silk

production (Kundu et al. 2012, Dong et al. 2015), as a food resource (Fogang Mba et al.

2019), and for health concern (Carrijo-Carvalho and Chudzinski-Tavassi 2007, Battisti et al.

2011, Ciminera et al. 2018). However, despite being one of the best-known families of

Lepidoptera, a robust phylogenetic hypothesis of interrelationships among its genera based on

extensive sampling is lacking. Saturniids are considered to be poor-dispersers, because of

their large and short-lived non-feeding adults that represent an extreme case of capital-

breeding organisms (Davis et al. 2016). These moths nonetheless managed to colonize and

diversify on almost all continents, but the spatial and temporal dynamics of their evolutionary

history remain largely unknown, for neither formal biogeographical analyses nor inferences

from time trees have been drawn for the family.

Saturniidae comprise 3,454 species and 180 genera worldwide, classified into eight

subfamilies and 11 tribes (Kitching et al. 2018). Five of these subfamilies (Arsenurinae,

Ceratocampinae, Cercophaninae, Hemileucinae and Oxyteninae) are restricted to the New

World and comprise 60% of the diversity of the family (>2,100 species). The subfamily

Agliinae (5 species) occurs throughout the Palearctic region while most species of Salassinae

(32 species) are in the Sino-Himalayan region with a few species in the East Palearctic and

Oriental regions. Finally, the highly diversified Saturniinae (>1,000 species) are present in all

biogeographical regions, except for polar and desertic regions. The family is considered

monophyletic based on both morphological (Minet 1994) and molecular evidence (Regier et

al. 2008, Breinholt et al. 2018, Hamilton et al. 2019). A few studies have addressed higher-

level relationships within the Saturniidae although with sparse taxon and often reduced gene

sampling (Regier et al. 2008: 8 species, 7 subfamilies, 5 loci, Barber et al. 2015: 80 species, 8

subfamilies, 6 loci, Hamilton et al. 2019: 16 species, 8 subfamilies, 650 loci). Nevertheless, a

consensus has emerged that the Neotropical Oxyteninae is sister to all other Saturniidae while

the Cercophaninae is the second earliest-diverging subfamily. The remaining lineages are

divided into two large clades, a Neotropical clade that includes three subfamilies (Arsenurinae

+ Ceratocampinae + Hemileucinae), and an old-world clade composed of Salassinae as sister

to Saturniinae. However, the position of the eighth subfamily – the palearctic Agliinae –

remains contentious. In recent phylogenomic analyses based on anchored hybrid enrichment

(Rubin et al. 2018, Hamilton et al. 2019), Agliinae is recovered as sister to the old-world

clade, but with low support, rather than as sister to the new-world clade as found in studies

based on Sanger sequencing of a few loci and with low support as well (Regier et al. 2008,

Barber et al. 2015, Rubinoff and Doorenweerd 2020). Beside this uncertainty, previous

studies have also raised questions regarding the placement of certain taxa, such as the genus

*Rhodinia* which has been placed as a member of the tribes Saturniini (Regier et al. 2002) or

Attacini (Rubin et al 2018; Chen et al. 2021), or poorly known genera such as the African

*Eosia* and *Eochroa* – doubtfully placed, respectively, in Micragonini and Bunaeini tribes

within Saturniinae (Oberprieler and Nässig 1994), or *Hirpida* and *Catharisa* of unclear

affinities within Hemileucinae (Michener 1952, Lemaire 2002). The latter is the most

speciose subfamily with more than 1,600 species in 55 genera, of which just eleven have been

included in a formal phylogenetic analysis (Barber et al 2015). Furthermore, although both

morphology and molecules have revealed the paraphyly of Urotini with respect to

Afrotropical Bunaeini and Micragonini (Regier et al 2008, Rubin et al 2018), incomplete

taxon sampling led taxonomists to refrain from revising a classification recognized as

unnatural for African Saturniidae (Oberprieler, 1997).

While genome-scale data can often help resolving phylogenetic relationships, they have

proven frustrating when addressing difficult nodes within the Saturniidae, presumably

because of the lack of signal associated with an old, rapid radiation (Hamilton et al. 2021) and

of the increasing probability of observing conflicting signals between markers (Kumar et al.

2012). In particular, although these properties are still rarely thoroughly explored,

heterogeneity in base composition and evolutionary rates of taxa and markers (e.g., Boussau

et al. 2014, Romiguier et al. 2016, Bossert et al. 2017, Borowiec et al. 2019, Cruaud et al.

2021, Rasplus et al. 2021) as well as saturation (Borowiec 2019, Duchêne et al. 2021) were

identified as major causes of analytical bias in phylogenomic analyses. These factors, in

concert with limited taxon sampling, may, for instance, explain the poorly resolved placement

of Agliinae within the Saturniidae (Hamilton et al. 2019) and constrain our understanding of

the early evolutionary history of this family.

In this study, we present a new time-calibrated phylogeny of Saturniidae reconstructed

with 1,024 ultraconserved elements (UCEs) and their flanking regions obtained from 338

species representing all described subfamilies, tribes, and genera of Saturniidae. We analyzed

potential systematic bias in genomic data and used the best-supported phylogenomic

hypothesis to (1) propose a revised classification of the family that reflects these inferred

relationships, and (2) infer a timeline for the origin and dispersal patterns of the family

through the estimation of its biogeographical history.

**MATERIALS AND METHODS**

*Taxon Sampling*

Our taxonomic sampling included 9.8% of all species of Saturniidae (338/3454) and

representatives of all recognized subfamilies (8), tribes (11), and genera (180) (Kitching et al.

2018). Our samples also included representatives of all subgenera of *Psilopygida*, *Meroleuca*,

*Gonimbrasia,* and *Antheraea*, and five of the six subgenera of *Hylesia* (Table S1). Sixteen

outgroup species were used, including (1) ten species of Sphingidae, the sister family of

Saturniidae (Hamilton et al. 2019), with representatives from all four of its subfamilies, and

(2) six species from four other families of Bombycoidea (Brahmaeidae, Endromidae,

Bombycidae and Eupterotidae).

*Library Preparations and Sequencing*

Samples were obtained from tissue collections at CBGP and MNHN, and from

historical specimen collections at MNHN when necessary. DNA was extracted from thorax

muscles or legs using the Qiagen DNeasy Blood and Tissue kit and following manufacturer's

protocol with modifications detailed in Cruaud et al. (2019). Alternatively, we also used DNA

extracts prepared for DNA barcoding studies at the Centre for Biodiversity Genomics

(University of Guelph, Ontario, Canada) following the protocol in Ivanova et al. (2006).

Library preparations followed Cruaud et al. (2019). Pools of 16 specimens made at equimolar

ratio were enriched in 1,381 UCEs using the 14,363 baits designed by Faircloth (2017)

(ordered as a myBaits UCE kit; Arbor Biosciences). Batches of 96 samples were sequenced

on different flow cells of an Illumina MiSeq platform with a 2x250 pb paired-end sequencing.

*Assembly of Data Sets into Loci, Data Set Cleaning*

Filtering of raw data and assembly into loci followed Cruaud et al. (2019). Only UCEs

that had a sequence from at least 50% of the samples were retained for analysis (n=1,024).

Loci were aligned with MAFFT (*-linsi* option; Katoh and Standley 2013). Sites with more

than 50% gaps and sequences with more than 25% gaps were removed from each UCE using

SEQTOOLS (package *PASTA*; Mirarab et al. 2015)). For all loci, two successive rounds of

TreeShrink (Mai and Mirarab 2018) were performed to detect and remove abnormally long

branches in individual gene trees, using the *per-species* mode and *b* parameter set to 20. Loci

were realigned with MAFFT between and after the two rounds of TreeShrink; gene trees were

inferred with IQ-TREE 2.0.6 (Minh et al. 2020) with the best fit model selected by

ModelFinder (Kalyaanamoorthy et al. 2017).

*Phylogenetic Analyses*

UCEs were analyzed with concatenation (IQ-TREE 2.0.6) and tree reconciliation

(ASTRAL-III, Zhang et al. 2018) approaches. For the concatenation approach, aligned UCEs

were merged and the resulting data set was analyzed without (hereafter referred to as

IQTREE-unpartitioned) and with partitions within the alignment. For the partitioned analysis,

each UCE was first divided into one core and two flanking regions using the Sliding-Window

Site Characteristics (SWSC) method (Tagliacollo et al. 2018). We then tested two partitioning

schemes: (1) a partition that grouped all core regions and another that grouped all flanking

regions (hereafter referred to as IQTREE-CoreVsFlanking); and (2) the best partitioning

scheme inferred by PartitionFinder2 (Lanfear et al. 2017) that joins core and flanking regions

under a similar evolutionary model (model selection = AICc; algorithm = rclusterf; branch

lengths linked; hereafter referred to as IQTREE-PartitionFinder). Best-fit models for the

unpartitioned data set and for all partitions were selected with the Bayesian Information

Criterion (BIC) as implemented in ModelFinder (Kalyaanamoorthy et al. 2017). FreeRate

models with up to ten categories of rates were included in tests for the unpartitioned data set

and for partitions that grouped all core regions or all flanking regions, but only common

substitution models were tested for the subsets identified by PartitionFinder. The candidate

tree set for all tree searches was composed of 98 parsimony trees + 1 BIONJ tree and only the

20 best initial trees were retained for NNI search. Node supports were assessed with ultrafast

bootstrap (UFBoot) (Minh et al. 2013) with a minimum correlation coefficient set to 0.99 and

1,000 replicates of SH-aLRT tests (Guindon et al. 2010). UFBoot values ≥ 95 and SH-aLRT ≥

80 were considered as strong support for a node. In addition, we computed gene (gCF) and

site (sCF) concordance factors (Minh et al. 2020) with IQ-TREE.

For ASTRAL analyses, nodes in gene trees with UFBoot support lower than 50, 70, or

90 were collapsed following the approach of Zhu (2014, perl script AfterPhylo.pl) before

reconciliation (trees are hereafter referred to as ASTRAL50, ASTRAL70 and ASTRAL90,

respectively). Node supports for the ASTRAL trees were evaluated with local posterior

probabilities (local PP). RF distances (Robinson and Foulds 1981) among recovered trees

were calculated with RAxML-NG_v0.9.0 (Kozlov et al. 2019) and Approximately Unbiased

(AU) tests (Shimodaira 2002) were performed with IQ-TREE (number of bootstrap replicates

set to 20,000).

*Analysis of Conflicting Topologies and Bias Exploration*

Systematic biases were investigated with respect to influences from both loci and taxa.

For analyses based on loci, Gene Genealogy Interrogation (GGI) (Arcila et al. 2017, Zhong

and Betancur-R 2017, Betancur-R et al. 2019) was used to identify UCEs that supported each

of the competing topologies. Each UCE was analyzed with IQ-TREE to infer trees using each

of the competing topologies as multifurcating constraint trees (the structure of the backbone

was fixed, but taxa within identified clades were free to move around). Per-site log-likelihood

scores for all constrained trees were then calculated with RAxML-8.2.11 (Stamatakis 2014)

and used to perform AU tests in CONSEL (Shimodaira and Hasegawa 2001). The program

*makermt* in CONSEL was used to generate K = 10 sets of bootstrap replicates with each set

consisting of 100,000 replicates of the row sums. Properties of the UCEs that supported each

topology were compared to detect whether significant differences could be identified between

groups of UCEs and whether any topology could result from analytical bias. Two properties

that are known to bias phylogenetic analyses, GC content (Romiguier and Roux 2017) and

saturation (Philippe et al. 2001, Duchêne et al. 2021), were analyzed. GC content was

calculated with AMAS (Borowiec 2016) and saturation (R squared of the linear regression of

uncorrected p-distances against inferred distances in individual gene trees) was calculated

following Borowiec et al. (2015). Results were analyzed in R (R Core Team 2018) using the

packages *ggpubr* (Kassambara 2020) and *rich* (Rossi 2011).

For analyses based on taxa, GC content and long branch (LB) score heterogeneity were

calculated for each taxon in each UCE and in the concatenated data set. GC content was

calculated with AMAS and LB heterogeneity scores were calculated with TreSpEx (Struck

2014). In each tree, the per sample LB score measures how the patristic distance (PD) of a

sample to all others deviates (as a percentage) from the average PD across all samples

(Struck, 2014). The calculation of the heterogeneity in LB scores can therefore help to reveal

possible Long Branch Attraction (LBA) artefacts (Bergsten 2005). Hierarchical clustering of

taxa based on GC content and LB scores was performed with the R package *cluster* (Maechler

et al. 2018).

*Divergence Time Estimates*

We estimated divergence times for Saturniidae with MCMCtree (Yang and Rannala

2006), using uniform distributions as calibration priors. A sphingid fossil (†*Mioclanis*

*shanwangiana*) provided a minimum age for the crown of Smerinthinii set to 15.0 Ma (Zhang

et al. 1994, Yang et al. 2007), while a maximum of 44.9 Ma was set following the maximum

age estimate for Sphingidae in Kawahara et al. (2019). For secondary calibration, the

minimum and maximum bounds of four selected nodes were respectively set to the mean of

minimum ages and the mean of maximum ages obtained by Kawahara et al. (2019) with

different calibration schemes and relaxed-clock models: (1) root (crown Bombycoidea): 59.6-

81.1 Ma; (2) crown Saturniidae + Sphingidae: 48.8-69.0 Ma; (3) crown Saturniidae: 35.1-52.5

Ma; and (4) crown Sphingidae: 29.6-44.9 Ma. Analyses were run with uncorrelated and

correlated relaxed clock models. The IQ-TREE topology obtained with the 50% least

saturated UCEs loci was used as input tree. Five data sets composed of 30,000 randomly

selected sites (custom script) were used as input data to make computation tractable. Two

chains were run for each data set, 20,000 generations were discarded as burnin, and chains

were run for at least 2 million generations with sampling every 200 generations and until

effective sample sizes of parameters reached at least 200 as reported by Tracer (Rambaut et

al. 2018). Possible conflicts between priors and data were controlled by sampling directly

from the prior of times and rates. Prior and posterior distributions were then compared in

Tracer. Marginal densities of parameters obtained with different data sets were compared with

Tracer. Posterior estimates obtained with the different chains were combined with

LogCombiner 2.6.0 (Bouckaert et al. 2019).

*Historical Biogeography*

Species distributions were assigned as presence/absence to the nine following

biogeographical areas: Neotropical, West Nearctic, East Nearctic, Afrotropical, Australasian,

Madagascar, Oriental, West Palearctic, and East Palearctic (Table S2). Ancestral area

estimations were performed using the R package *BioGeoBEARS 1.1.1* (Matzke 2014). The

chronogram built with MCMCtree was used as input but only one specimen per

genus/subgenus was included in the analysis and outgroups were removed to avoid artefacts

(taxa were pruned from the chronogram with the R package *ape* (Paradis and Schliep 2018)).

Dispersal-Extinction-Cladogenesis (DEC; Ree and Smith 2008), BAYAREALIKE (Landis et

al. 2013) and DIVALIKE (Ronquist 1997) models were used with and without considering

the jump parameter for founder events (+$J$; Matzke 2014). Model selection was performed by

considering statistical (AICc; Klaus and Matzke 2020, Matzke 2021) and non-statistical (i.e.,

biological and geographical) information (Ree and Sanmartin 2018). The maximum number

of areas that a species could occupy was set to 5. To consider the main geological events that

occurred during the diversification of Saturniidae, we defined five time periods with different

dispersal rate scalers: (1) from the mean crown age of Saturniidae (ca. 52 Ma) to 34 Ma; (2)

34 to 23 Ma (Oligocene); (3) 23 to 14 Ma (early to mid-Miocene); (4) 14 to 5 Ma (mid to late

Miocene); (5) 5 Ma to present. Dispersal rate matrices between areas and adjacency matrices

are provided in Table S2.

**RESULTS**

*Initial Data Set and Trees*

The initial phylogenomic data set included sequence information for 354 taxa (338

species from 185 genera of Saturniidae + 16 outgroup species) and 1,024 UCEs representing

579,775 aligned bp (215,449 bp for core regions and 364,326 bp for flanking regions) of

which 66.7% were parsimony informative (Table S3). PartitionFinder grouped the 3,072

core/flanking regions into 1,943 subsets. The six trees (IQTREE-unpartitioned, IQTREE-

CoreVsFlanking, IQTREE-PartitionFinder, ASTRAL50, ASTRAL70, ASTRAL90) are

shown in Figure S1. Relationships within Saturniidae inferred by IQTREE were well

resolved with only 6-7% of the nodes receiving SH-aLRT/UFBoot scores below 100/100. IQ-

TREE trees were considered as equally good explanations of the data set (AU test *P*-value >

0.05; Table S3), regardless of the partitioning scheme.

The differences between IQ-TREE topologies (RF pairwise distances 4-8; Table S3)

were located in a few poorly supported regions of the topology (within Bunaeini and

Arsenurinae). Except for 8-10% of the nodes, local PP were above 0.99 in ASTRAL trees.

The differences between ASTRAL topologies (RF pairwise distances 6-14; Table S3) were

restricted to parts that were poorly supported. Depending on the pairs of topologies compared,

from 19 to 24 clades were observed in IQ-TREE trees but not in ASTRAL trees and vice

versa. All conflicts (but one, see next section) were either unsupported in IQ-TREE trees,

ASTRAL trees, or in both types of trees.

Saturniidae was recovered monophyletic with maximal node support in all analyses. All

subfamilies were recovered as monophyletic with strong support except Hemileucinae which

was polyphyletic with *Hirpida + Hirpsinjaevia* well separated from other species. The

Oxyteninae and Cercophaninae formed a grade sister to all other saturniids. Except for

Agliinae and *Hirpida + Hirpsinjaevia* (Fig. 1; see next section), relationships among the main

lineages were strongly supported with Arsenurinae recovered as sister to Ceratocampinae +

Hemileucinae (excluding *Hirpida + Hirpsinjaevia*) and Salassinae recovered as sister to

Saturniinae. All but five tribes (Hemileucini, Urotini, Micragonini, Saturniini, Bunaeini) were

recovered as monophyletic with strong support. Finally, 95 (92%) of the 103 genera

represented by multiple species were recovered as monophyletic. The six genera *Automeris*,

*Eacles, Ludia*, *Neodiphthera*, and *Pseudodirphia* were problematic as they were rendered

paraphyletic, respectively, by *Eubergioides, Catharisa, Vegetia, Pararhodia,* and

*Kentroleuca*. Two genera (*Gonimbrasia*, *Molippa*) appeared polyphyletic while *Syssphinx*

was recovered as a grade that also included the North American *Dryocampa* and *Anisota*.

**Figure 1. Conflicting topologies observed after analysis of the initial data set (a) and results of the gene genealogy interrogation (GGI) approach (b)**. In panel (a), topology *A* is supported by IQ-TREE analyses; *Abis* is supported by ASTRAL90; *B* is supported by ASTRAL70 and *C* is supported by ASTRAL50. All corresponding trees are available in Figure S1. In simplified topologies, Hemileucinae is used for Hemileucinae minus *Hirpida* and *Hirpsinjaevia* and Hirpi/Hirpsi stands for *Hirpida + Hirpsinjaevia.* For the GGI approach, Hemileucinae (minus *Hirpida* and *Hirpsinjaevia*) + Ceratocampinae + Arsenurinae; Saturniinae + Salassinae; *Hirpida + Hirpsinjaevia* and Agliinae were considered as different clades and trees were inferred using each of the competing topologies as multifurcating constraint trees. The structure of the backbone was fixed, but taxa within clades were

free to move around. Topology *A* was considered equivalent to *Abis* as we focused on sister taxa

relationships of Agliinae / *Hirpida* + *Hirpsinjaevia* with either Hemileucinae + Ceratocampinae +

Arsenurinae or Saturniinae + Salassinae. In panel (b), average values of GC content and saturation (R-

squared of the regression between pairwise distances calculated from aligned sequences and branch

length) for loci that significantly supported topologies *A/Abis* (i.e., Pvalue of the AU test $\leqq 0.05$; N

loci = 189; Table S3) or not (Nloci=796) were compared using a randomization test (N randomization

= 9999; c2m function of the R package *rich*) and P values are reported on graph.

*Analysis of Conflicting Topologies*

A single important conflict was observed among the trees with respect to the position of

Agliinae and of species in the *Hirpida* + *Hirpsinjaevia* clade (Figs. 1 and S1): (1) Agliinae

and *Hirpida* + *Hirpsinjaevia* either formed a clade sister to Arsenurinae + Ceratocampinae +

other Hemileucinae (ACH) (IQ-TREE; topology A), or branched sequentially with Agliinae

sister to a clade grouping *Hirpida* + *Hirpsinjaevia* and ACH (ASTRAL90; topology Abis); or

(2) *Hirpida* + *Hirpsinjaevia* grouped with ACH while Agliinae grouped with Salassinae +

Saturniinae (ASTRAL70; topology B); or (3) Agliinae and *Hirpida* + *Hirpsinjaevia* formed a

grade basal to Salassinae + Saturniinae (ASTRAL50; topology C). The contrasting results

obtained with ASTRAL and the gCFs of the Agliinae + *Hirpida* + *Hirpsinjaevia* clade in the

IQ-TREE trees (8.6%; Fig. S1) highlight conflict among gene trees. GGI analyses show that

of the 985 UCEs available for both Agliinae and *Hirpida* + *Hirpsinjaevia* species, 540

(54.8%) favored topologies A/Abis; 228 (23.2%) favored topology B; and 217 (22.0%)

favored topology C (Table S3). In addition, topologies A/Abis were a significantly better

explanation for 189 (19.2%) of the gene trees while topology B was a significantly better

explanation for only 3 (0.3%) of the gene trees, and topology C was a significantly better

explanation for 9 (0.9%) of the gene trees. Statistical analyses of UCE properties show that

topologies A/Abis were supported by UCEs with lower saturation (Fig. 1(2)). Further

analyses of data subsets comprising the 20% least saturated (*n*=204) or 50% most saturated

(*n*=512) loci confirmed that the placement of Agliinae as sister to Salassinae + Saturniinae

was likely an artefact caused by saturation of phylogenetic signal (Fig. S2). Both IQ-TREE

and ASTRAL90 analyses of the least saturated UCEs recovered Agliinae close to ACH

(topologies A/Abis; Fig. 1a), while Agliinae always jumped into the Saturniinae + Salassinae

part of the tree when analysis focused on the most saturated loci (Fig. S2). In addition, taxon-

based analysis of systematic bias did not reveal evidence that the sister taxa relationship

between Agliinae and *Hirpida + Hirpsinjaevia* (topologies A/Abis; Fig. 1a) resulted from

compositional (GC) bias or LBA (Fig. S3).

Finally, regardless of the conflicting placement of the subfamily Agliinae, IQ-TREE

and ASTRAL analyses based on the least saturated half of the loci produced the most similar

topologies (RF distances; Table S3). Consequently, subsequent analyses focused on the

phylogram derived from the IQ-TREE analyses of the 50% least saturated UCEs (*n*=512;

Figs. 2 and S2I).

Saturniinae, *Rothschildia*

Bunaeinae, *Gonimbrasia*

Salassinae, *Salassa*

Hemileucinae, *Leucanella*

Ceratocampinae, *Eacles*

Arsenurinae, *Caio*

Hirpidinae, *Hirpida*

Agliinae, *Aglia*

Cercophaninae, *Cercophana*

Oxyteninae, *Oxytenis*

**Figure 2. Saturniidae tree of life.** The classification proposed in this study is displayed on the

chronogram (MCMCtree, uncorrelated relaxed clock model) inferred from the topology selected after

exploration of systematic bias (i.e., the IQ-TREE tree inferred from the 50% least saturated loci). Red

dots at nodes identify poorly supported nodes (SHaLRT<80 or UFBoot<95). Abbreviations used: A =

Antistathmopterini; Agl = Agliinae; Al = Almeidaiini; Bathy = Bathyphlebiini; C = Cercophanini; Cer

= Cercophaninae; Cith = Citheroniini; E = Eudaemoniini; Eo = Eochroini; Hir = Hirpidinae; J =

Janiodini; Lono = Lonomiini; Oxyteninae; P = Pseudapheliini; Poly = Polythysanini; Sal = Salassinae;

So = Solini; Urot = Urotini.

*Divergence Time Estimates and Historical Biogeography*

366            Analyses based on both autocorrelated and uncorrelated relaxed clock models provided

similar results (median of differences between mean or median ages of all nodes = 1.65 Myr;

Table S4), therefore we only represent (Figs. 2 and 3) and cite estimates based on the

uncorrelated relaxed clock model in main text for brevity. The divergence times and

confidence intervals for correlated and uncorrelated clock models are provided in Table S4 for

all nodes (see also Figure S4 for the chronograms), while those of clades discussed below are

provided in Table 1.

| | Stem (in Myr) | | Crown (in Myr) | |
|---|---|---|---|---|
| **Clades** | **Uncorrelated** | **Correlated** | **Uncorrelated** | **Correlated** |
| ACH | 45.11 [42.93-47.18] | 42.94 [39.69-45.31] | 44.08 [41.85-46.19] | 42.05 [38.85-44.51] |
| *Agapema* | 11.90 [9.76-14] | 14.02 [11.89-16.60] | 4.85 [2.81-7.06] | 5.78 [3.7-7.94] |
| Agliinae | 42.95 [39.88-45.81] | 41.44 [38.10-44.26] | 3.09 [2.20-4.06] | 13.01 [8.94-16.40] |
| Agliinae + Hirpidinae + ACH | 45.98 [43.88-47.97] | 43.65 [40.50-45.94] | 45.11 [42.93-47.18] | 42.94 [39.69-45.31] |
| American *Antheraea* (*Telea*) | 8.36 [5.95-10.91] | 9.51 [7.10-11.77] | 3.93 [2.34-5.67] | 4.67 [3.16-6.30] |
| *Antherina* + *Ceranchia* | 27.20 [24.78-29.65] | 28.34 [25.43-31.24] | 7.61 [4.42-11.23] | 11.05 [7.21-14.94] |
| *Argema* | 14.53 [12.12-17.01] | 13.30 [10.63-15.68] | 10.63 [8.03-13.22] | 9.81 [7.54-12.05] |
| Arsenurinae | 44.08 [41.85-46.19] | 42.05 [38.85-44.51] | 40.04 [36.06-43.62] | 39.52 [36.06-42.25] |
| BSS | 45.98 [43.88-47.97] | 43.65 [40.50-45.94] | 43.36 [40.97-45.65] | 41.46 [38.34-43.87] |
| *Bunaea* (Madagascar colonization) | [-] | [-] | 2.00 [1.21-2.87] | 1.91 [1.32-2.53] |
| Bunaeinae | 40.17 [37.61-42.67] | 38.27 [35.48-40.87] | 37.22 [34.66-39.78] | 35.24 [32.48-37.74] |
| *Calosaturnia* | 14.16 [12.01-16.12] | 16.58 [14.42-19.38] | [-] | [-] |

| | | | | |
|---|---|---|---|---|
| Ceratocampinae | 41.84 [39.51-44.07] | 39.33 [36.32-41.92] | 35.97 [33.04-38.83] | 32.02 [28.43-35.05] |
| Ceratocampinae + Hemileucinae | 44.08 [41.85-46.19] | 42.05 [38.85-44.51] | 41.84 [39.51-44.07] | 39.33 [36.32-41.92] |
| Cercophaninae | 49.61 [47.96-51.04] | 48.92 [46.36-50.67] | 38.08 [31.29-43.79] | 39.99 [34.91-44.22] |
| *Copaxa* | 21.11 [18.71-23.48] | 22.37 [19.87-25.02] | 10.89 [7.96-14.03] | 9.92 [6.94-13.03] |
| *Coscinocera* | 11.21 [8.61-13.93] | 11.28 [8.75-14.14] | 2.04 [1.15-3.06] | 1.87 [1.03-2.81] |
| *Eosia* | 17.02 [13.86-20.19] | 15.98 [13.05-19.48] | 4.18 [2.49-6.05] | 2.76 [1.50-4.19] |
| *Epiphora* | 14.51 [12.18-17.06] | 14.60 [11.88-17.77] | 6.62 [4.64-8.73] | 5.91 [3.86-8.13] |
| *Eupackardia + Rothschildia* | 21.72 [18.92-24.58] | 22.20 [19.05-25.43] | 10.83 [8.12-13.71] | 11.65 [8.48-14.97] |
| Hemileucinae | 41.84 [39.51-44.07] | 39.33 [36.32-41.92] | 36.48 [33.72-39.27] | 32.17 [29.08-35.06] |
| Hirpidinae | 42.95 [39.88-45.81] | 41.44 [38.10-44.26] | 19.03 [13.31-25.18] | 23.87 [17.69-29.91] |
| *Maltagorea* | 11.56 [9.09-14.00] | 8.78 [6.46-11.13] | 8.34 [5.94-10.82] | 6.63 [4.69-8.58] |
| *Opodiphthera* clade | 20.92 [18.65-23.28] | 21.77 [19.27-24.38] | 11.80 [9.46-14.25] | 11.60 [8.80-14.47] |
| Oxyteninae | 51.90 [50.66-52.55] | 51.52 [49.25-52.55] | 37.10 [31.08-43.13] | 31.63 [24.19-38.56] |
| Saturniidae | 63.27 [58.69-68.51] | 63.96 [59.74-68.52] | 51.90 [50.66-52.55] | 51.52 [49.25-52.55] |
| *Sinobirma* | 14.02 [11.46-16.53] | 10.56 [7.91-13.16] | [-] | [-] |

**Table 1. Summary of mean divergence time estimates and 95% highest posterior density (HPD)**

**for groups or colonization events discussed.** Mean and median ages with 95% HPD for all nodes are

provided in Table S4. Units are in Myr. Our new classification is used for taxon names. ACH =

Arsenurinae + Ceratocampinae + Hemileucinae; BSS = Bunaeinae + Salassinae + Saturniinae.

The stem age of Saturniidae was estimated in the early Cenozoic at about 63 Ma. The

Oxyteninae and Cercophaninae originated about 52 Ma and 50 Ma, respectively. All other

subfamilies originated during a brief interval (4 Myr) in the middle Eocene (46-42 Ma) and

diversified during the Oligocene and Miocene. While only 32 (18%) of the 180 extant genera

had appeared before the end of the Oligocene (including all 9 genera of Arsenurini), the

remaining bulk of extant saturniid genera originated during the Miocene (Fig. 3).

**Figure 3. Global historical biogeography of Saturniidae.** Ancestral ranges were inferred with the

BAYAREALIKE+J model (best-fit and most plausible model according to empirical considerations;

see text) as implemented in *BioGeoBEARS*. Alternative inferences of ancestral ranges are provided in

Figure S5. The chronogram is derived from the species-level chronogram obtained with the

uncorrelated relaxed clock model implemented in MCMCtree (see Fig. 2) that was pruned to keep

only one specimen per genus/subgenus. The classification proposed in this study is used to annotate

the tree. Colored boxes following each terminal name represent biogeographical regions A to I as

illustrated in the upper-left map; colored arrows refer to inferred dispersal/colonization events.

Abbreviations used: A = Agliinae; Ars = Arsenurinae; C=Cercophaninae; H = Hirpidinae; O =

Oxyteninae; S = Salassinae. PPH=Pliocene, Pleistocene and Holocene. Illustrative photograph =

*Antherina suraka* (Saturniinae); image courtesy of Armin Dett.

With respect to historical biogeography, inclusion of the parameter *J* (jump dispersal

events during cladogenesis) in models of geographic range evolution remains controversial, as

is the validity of statistical comparisons of models with and without *J* (Ree and Sanmartín

2018; Matzke 2021). Empirical considerations have been advocated as an alternative way to

select among biogeographical scenarios (Ree and Sanmartin 2018). Here, both AICc and the

observation that most extant subfamilies occupy a single biogeographical area favor

biogeographical models that include the *J* parameter (Table S2, Fig. S5). Without it, the DEC

model had the best fit, but inferred a widespread ancestor for the family with multiple

subsequent local extinctions. When the *J* parameter was included, BAYAREALIKE+J was

inferred as the best-fit model (Table S2) and all ancestral range estimations unambiguously

supported a Neotropical origin for saturniids.

Given the most likely ancestral areas estimation, saturniid biology, and

palaeogeological knowledge, following their origin in the Neotropics, the Saturniidae likely

colonized the Old World twice during the mid-Eocene, probably through Beringia. These two

dispersal events involved an ancestor of Salassinae + Saturniinae between 46 and 43 Ma, and

an ancestor of Agliinae at 43 Ma (red arrows in Figure 3). Shortly later, a lineage (represented

by three tribes - Bunaeini, Micragonini and Urotini - in the current classification) colonized

Africa between 40 and 37 Ma and subsequently diversified extensively (with 55 extant

genera). This lineage remained restricted to this continent except for three lineages which

dispersed to new areas: *Sinobirma* invaded the Oriental region (*ca*. 14 Ma), while *Maltagorea*

(*ca*. 12 Ma) and *Bunaea* (*ca*. 2 Ma) colonized Madagascar. By contrast, the Attacini and

Saturniini tribes in Saturniinae not only diversified in the Oriental and East Palearctic regions,

but also dispersed multiple times from there to adjacent areas (Fig. 3). They colonized Africa

and Madagascar on four occasions during the mid-Miocene Climatic Optimum (*Epiphora* 14

Ma, *Argema* 14 Ma, *Eosia* 17 Ma, and the clade *Antherina* + *Ceranchia* 27 Ma). They also

dispersed through Sunda and Wallacea on multiple times, crossing Lydekker's line on three

occasions (*Attacus*, *Coscinocera*, *Austrocaligula* clade). Finally, they returned seven times to

the New World with two relatively ancient dispersal events (early Miocene; *Copaxa* 21 Ma

and the clade *Eupackardia* + *Rothschildia* 21 Ma) and five others during the second half of

Miocene (*Actias* 14 Ma, *Antheraea* (*Telea*) 13 Ma, *Calosaturnia* 13 Ma, *Agapema* 12 Ma, and

the clade *Hyalophora* + *Callosamia* 7 Ma). The diverse Neotropical fauna of saturniids

colonized the Nearctic region only seven times (Fig. 3, right-hand section of tree, green

arrows) and all these dispersal events postdated the final formation of the Isthmus of Panama

(*ca*. 15 Ma).

**DISCUSSION**

*A Comprehensive Genus-level Phylogeny of Saturniidae*

This study expands upon earlier investigations (Regier et al. 2008, Barber et al. 2015,

Hamilton et al. 2019) to yield a comprehensive and robust phylogenetic framework for higher

relationships within Saturniidae. The results confirm that Oxyteninae and Cercophaninae

were the first two subfamilies to diverge and that Salassinae is sister to Saturniinae. The

present results agree with those of Barber et al. (2015) and Hamilton et al. (2019) as they

place Arsenurinae as sister to Ceratocampinae + Hemileucinae (but without *Hirpida* +

*Hirpsinjaevia*; see below). However, in-depth analysis of conflict among gene trees shows

that subfamily Agliinae, whose position remained unelucidated, belongs to a lineage that also

includes the new-world subfamilies Arsenurinae, Ceratocampinae, and Hemileucinae. This

suggests that its alternative placement as sister to Salassinae and Saturniinae as obtained by

Hamilton et al. (2019) was also likely an artefact of tree reconstruction, possibly caused by

the saturation of nucleotide sequences. Indeed, when these authors analyzed the AHE loci

using protein sequence data which saturate less rapidly owing to a larger state space (20

possible amino acids; see their additional file 8), they recovered Agliinae as more closely

related to new-world subfamilies. These relationships are corroborated by larval characters

shared with Arsenurinae (Michener 1952; e.g., D2 setae on abdominal segment A9 of first

instar larva form a median scolus or chalaza; first larval stages bear two or more pairs of large

thoracic horns forked at their apices). Interestingly, our analyses are the first to examine the

genera *Hirpida* and *Hirpsinjaevia* and they reveal that these two closely related Neotropical

genera are sister to Agliinae. Unfortunately, the host plant(s) and immature stages of *Hirpida*

and *Hirpsinjaevia* are unknown but, based on their phylogenetic placement, we predict that

their first instar larvae bear large thoracic horns and may feed on Fagaceae and/or Betulaceae

like the Agliinae.

In the arsenurine tribe Arsenurini, Hamilton et al. (2021) found from a thorough

phylogenomic analysis of hundreds of AHE genes that phylogenetic signal remained elusive,

likely reflecting a burst of diversification. Our analysis also found that relationships among its

component genera are poorly supported (Figs. 2 and S2I) and are discordant from the

hypothesis preferred by Hamilton et al. (2021). This casts doubt on current understanding of

their evolutionary history, an uncertainty that can only be resolved by further investigation,

perhaps by combining AHE and UCE loci and expanding taxon sampling. Within the

Ceratocampinae and Hemileucinae, our results provide the first robust, comprehensive

hypothesis of relationships among its genera. They fully agree with conclusions based on a 6-

gene data set and limited taxon sampling (Barber et al 2015), but largely contradict groupings

proposed for Ceratocampinae by Balcazar-Lara & Wolfe (1997) based on their cladistic

analysis of both adult and immature morphology.

Within Saturniinae, our results unequivocally support the position of *Rhodinia* as sister

to members of the Attacini, in agreement with AHE-based conclusions (Rubin et al. 2018)

and recent findings from the analysis of mitogenomes (Chen et al. 2021). In addition, as our

sampling examined all genera in the tribe Urotini – considered to represent an artificial

grouping of several unrelated lineages (Oberprieler 1997; Rougerie 2005; Barber et al 2015;

Rubin et al 2018) – the outcome of our analyses offers the first robust framework of the

evolutionary history of African saturniids to lay down a new classification.

*Taxonomic Implications*

Our phylogenetic hypothesis has important and unexpected implications for the

classification of Saturniidae (Fig. 2). We propose several taxonomic changes that are detailed

below and summarized as a complete genus-level checklist (SI Appendix S1; see also Table

S1). The unexpected placement of *Hirpida + Hirpsinjaevia* as sister to Agliinae as well as the

age of the lineage justify the erection of a new subfamily: Hirpidinae Rougerie **subfam. nov**.

As all three previously recognized Afrotropical tribes of Saturniinae (Bunaeini, Urotini,

Micragonini) form a monophyletic group sister to all other Saturniinae (Attacini, Saturniini

and Solus), we adopt here the classification proposed by Nässig et al. (2015) that recognizes

this clade as subfamily Bunaeinae Bouvier, **1927 stat. rev.**. Within this subfamily, we

consider Bunaeini, Micragonini, and Urotini as monophyletic tribes but with new

circumscriptions ensuring their monophyly. To achieve this, we describe a new tribe and

resurrect three others involving the following seven genera: *Eochroa* (previously in

Bunaeini), *Parusta* and *Usta* are assigned to Eochroini Cooper, 2002 **stat. rev.**; *Pseudaphelia*

and *Pselaphelia* to Pseudapheliini Packard, 1914 **stat. rev.**; *Eudaemonia* to Eudaemoniini

Packard, 1902 **stat. rev.**; and *Antistathmoptera* to Antistathmopterini Rougerie **trib. nov.**.

The rare, enigmatic east African genus *Eosia* Le Cerf,1911 (Bouyer 2002), previously placed

within Micragonini (Bunaeinae), is recovered as sister to *Cricula* Walker, 1855 of the

Saturniini (Saturniinae) to which it is transferred, a position which agrees with evidence from

the morphology of first instar larva (Rougerie 2005). The genus *Solus*, whose phylogenetic

placement is clarified here for the first time, is sister to Attacini + Saturniini, a position that

requires the erection of a new tribe: Solini Rougerie **trib. nov.**. We also propose to follow

Nässig (1991) in considering *Graellsia* Grote, 1896 to be a junior synonym of *Actias* Leach,

1815. Since our results propose the first robust, comprehensive phylogeny for the two most

diverse subfamilies of new-world Saturniidae, we propose a new tribal arrangement for their

genera (Fig. 2; see Table S1 and Appendix S1 for details): (1) in Ceratocampinae, we

resurrect the following three tribes: Bathyphlebiini Travassos & Noronha, 1967 **stat. rev.**,

Ceratocampini Harris, 1841 **stat. rev.**, and Dryocampini Grote & Robinson, 1866 **stat. rev.**;

and (2) in Hemileucinae, we expand the previously monotypic tribe Polythysanini Michener

1952 by adding the genera *Adetomeris*, *Callodirphia*, and *Cinommata*; resurrect tribe

Lonomiini Bouvier, 1930 **stat. rev.** for genera *Lonomia* and *Periga*; and exclude *Catharisa*

*cerina* Jordan 1911, considered until now as an hemileucine (Lemaire 2002, Smith et al.

2013), while we recovered it nested within the genus *Eacles* of the Ceratocampinae. This

unexpected placement agrees with recent observations of the morphology of its first instar

caterpillars (P. Smith, pers. comm.) and leads us to treat *Catharisa* Jordan, 1911 **syn. nov.** as

a junior synonym of *Eacles* Hübner, [1819], with *Eacles cerina* **comb. nov.** transferred within

Bathyphlebiini of the Ceratocampinae. Within the Hemileucinae, the monotypic genus

*Eubergioides* Michener, 1949 **syn. nov.** appears nested within *Automeris* Hübner, [1819], an

affinity already suggested by Michener (1952) based on the comparison of male genitalia; it is

considered a junior synonym of the latter, with *Automeris bertha* (Schaus, 1896) **comb. nov.**

transferred to genus *Automeris*. Among the African bunaeine saturniids, we resurrect the

genus *Pinheyella* Cooper, 2002 **stat. rev.** for *P. anna* Maassen, 1985 **comb. nov.** whose

current placement in genus *Gonimbrasia* Butler, 1878 makes that genus polyphyletic. Finally,

we note that our results question the monophyly of six genera, but we refrain from proposing

nomenclatural change as limited sampling within each genus impedes proper circumscription

of natural groups: *Neodiphthera* Fletcher, 1982; *Ludia* Wallengren, 1865; *Molippa* Walker,

1855; *Pseudodirphia* Bouvier, 1928; *Kentroleuca* Draudt, 1929; and *Syssphinx* Hübner,

[1819]. From here on, subfamily, tribe and genus names refer to the new classification

proposed in this study (Fig. 2; Appendix S1).

*Historical Biogeography*

This work represents a first investigation of the historical biogeography of saturniids

(Fig. 3) based on a robust dated phylogeny that includes all known extant genera. It reveals

that the family originated in the Neotropics shortly after the Cretaceous-Paleogene boundary.

Its diversification was marked by a key split about 46 Ma between lineages AACHH (formed

by subfamilies Agliinae, Arsenurinae, Ceratocampinae, Hemileucinae and Hirpidinae; >2,000

extant species in 96 genera) and BSS (Bunaeinae, Salassinae and Saturniinae; >1,350 species

in 86 genera). The former remained confined to the New World except for the Agliinae (see

below), while the later diversified in all biogeographical regions, before recolonizing the New

World on several occasions. This split had been preceded by two earlier branching lineages

(first Oxyteninae 52 Ma, then Cercophaninae 50 Ma) that have remained restricted to South

America where they comprise 70 and 122 extant species respectively (Kitching et al. 2018,

Brechlin 2020).

Dispersal out of the Neotropics and into the Old World of the ancestors of lineage BSS

occurred during the middle Eocene (46-43 Ma). As dispersal across the oceanic barriers

separating the New and Old Worlds seems unlikely, it is more likely that dispersal occurred

via the Nearctic during the mid-Eocene Climatic Optimum, a transient period of global

warming (Westerhold et al. 2020). This would have required over-water dispersal through an

archipelago in the Panamanian gap (Pindell et al. 2005, Iturralde-Vinent 2006) or via the

Caribbean Arc that subsided toward the end of the Eocene (Pindell et al. 2005, Philippon et al.

2020, Cornée et al. 2021, Garrocq et al. 2021) before their northward dispersal. At this time

period, further dispersal of the BSS ancestors from the Nearctic into the Eastern Palearctic,

and from there southwards into the Oriental region would have been possible through the

Boreotropical forest belt (Morley 2003, 2011), which extended as far as the Arctic Circle,

facilitating faunal exchanges (Jahren 2007). Our results also suggest that ancestors of the

Agliinae, within the AACHH lineage, likely used the same route through Beringia about 43

Ma as an independent long-distance dispersal from the New to Old World. Interestingly, the

food plants (Ballesteros-Mejia et al 2020) of both Agliinae and early lineages of the BSS

lineage (e.g., beech, oak, maple, plum trees, sumac) were well represented in the forests of the

northern hemisphere during middle Eocene (Jahren, 2007), thus lending support to this

dispersal route.

During this same period (Fig. 3; 40-37 Ma), ancestors of Bunaeinae dispersed from the

Oriental to African region. This dispersal was facilitated by the mixed deciduous and

evergreen forests occurring in Central Asia and by the presence of archipelagoes and land

connections.  Between 41 and 35 Ma, Balkanotolia, an archipelago that spanned the

Neotethyan margin (Licht et al., 2022) and the Alboran/Apulian platforms

emerged (Vandenberghe et al. 2012), possibly facilitating the colonization of Africa. This

dispersal may have been driven by the climatic deterioration (or "Grande Coupure")

(Hartenberger 1998, Licht et al. 2022) that occurred during the Eocene-Oligocene transition

period, effectively forcing the Bunaeinae ancestor to seek warmer conditions southward.

Interestingly, several groups of mammals (e.g., rodents, primates) with excellent fossil

records also colonized Africa near the Bartonian-Priabonian boundary (*ca* 37.8 Ma) (Huchon

et al. 2007, Chaimanee et al. 2012, Marivaux et al. 2014, Coster et al. 2015, Licht et al. 2022).

After colonizing Africa, the ancestral Bunaeinae experienced a spectacular diversification as

all extant tribes had appeared by the early Oligocene. Further investigation of the importance

of different key innovations (ecological or morphological) that have triggered this

diversification burst constitute an interesting perspective.

As cooling during the Eocene–Oligocene disrupted the belt of Boreotropical forests, the

accretion of ice sheets broke the Beringian connection, leading to the divergence of ancestral

Saturniinae in these forests. Our analyses (Figs. 3 and S5) show that the Saturniinae and most

of its lineages originated in the Oriental + Eastern Palearctic region, but as the Boreotropical

forests contracted drastically during the "Grande Coupure", this likely forced southward

displacements of their ranges into refuges in the Sino-Himalayan mountains of the Eastern

Palearctic region (LePage et al. 2005, Xing and Ree 2017, Jia and Bartish 2018). The

observation that the diversity of non-African members of the BSS clade peaks in the Sino-

Himalayan mountains may reflect the role of this transition zone as a hub for further

diversification of these moths. A number of Saturniinae lineages subsequently dispersed and

diversified extensively in the Oriental and Palearctic regions, colonizing the Western

Palearctic four times: *Neoris* (stem age 20 Ma), *Perisomena* (13 Ma), *Saturnia* (12Ma), and

*Actias* (7 Ma). They also colonized Australasia (the *Opodiphthera* group (21 Ma) and

*Coscinocera* (11 Ma) within the Attacini), the Afrotropics (*Epiphora* (14 Ma), *Eosia* (17 Ma)

and *Argema* (14 Ma)), and the Nearctic and Neotropical regions (*Eupackardia* + *Rothschildia*

(22 Ma), *Copaxa* (21 Ma), *Calosaturnia* (13 Ma), and *Agapema* (12 Ma)). Interestingly, a

single dispersal event by a member of the genus *Sinobirma* (subfamily Bunaeinae) occurred

from Africa to the Eastern Palearctic region, nearly synchronous (14 Ma) with the

colonization of the Afrotropics by *Epiphora*, *Eosia,* and *Argema.* These range expansions

may have been facilitated by the collision of the Arabian plate with Eurasia *ca.* 15 Ma (Rogl

1999, Harzhauser et al. 2007).

*Final Remarks*

Our knowledge of insect evolution remains strongly impeded by both the scale of their

diversity and the difficulty to generate accurate and robust spatial and evolutionary

frameworks based on comprehensive coverage. Currently, only a few densely sampled and

time-calibrated phylogenies have been generated in diverse groups of insects, bringing

insights into the origins of observed diversity patterns such as the latitudinal diversity gradient

(e.g., Condamine et al. 2012, Chazot et al. 2021 for butterflies; Economo et al. 2018 for ants),

or the influence of life-history traits on diversification dynamics (Sota et al. 2022). We

believe that the Saturniidae family, as one of the best-documented family of insects with

global distribution and high species-richness, represents a new promising model to further

investigate the mechanisms driving diversification in insects. Our results have revealed that

the spatial dynamism of Saturniinae (33 genera, 741 species) is in striking contrast (Fig. 3) to

the situation in other subfamilies that also diversified extensively (Bunaeinae: 50 genera, 580

species; Ceratocampinae: 30 genera, ca. 300 species; Hemileucinae: 55 genera, ca. 1600

species; see Kitching et al. (2018)), but remained restricted to the biogeographical region

where they originated: Bunaeinae in the Afrotropical region (except for *Sinobirma*, and

including two independent colonization of Madagascar); Ceratocampinae and Hemileucinae

in the Neotropics with limited dispersal to the Nearctic region. The reasons for these

contrasting dispersal patterns remain unclear, but differing life-history traits, such as larval

diet and pupation modes (cocoons in Saturniinae are among the strongest within the family)

may have facilitated long-distance dispersal and range expansion. In general, the new spatial

and temporal framework produced here for the evolution of wild silkmoths opens new

avenues of research to understand the intriguing success of these capital-breeding moths that

managed to occupy all biogeographical regions today and to represent one of the most diverse

and abundant group of large-sized insects.

**FUNDING**

This work was supported by French National Research Agency (SPHINX: ANR-16-

CE02-0011-01, GAARAnti: ANR-17-CE31-0009, LABEX CEBA: ANR-10-LABX-25-01

and LABEX TULIP: ANR-10-LABX-41); French Foundation for Research on Biodiversity

(FRB) and Centre for the Synthesis and Analysis of Biodiversity (CESAB: ACTIAS grant).

**ACKNOWLEDGEMENTS**

We are thankful to the Canadian Centre for DNA Barcoding and Centre for Biodiversity

Genomics at University of Guelph (Ontario, Canada) as well as to all the participants to the

Saturniidae DNA barcoding campaign for their contributions to the assembly of samples. We

thank Corinne Poitout, Hélène Vignes and Audrey Weber (INRAE and CIRAD, AGAP,

France) for their assistance to sequencing, and the Genotoul bioinformatics platform

(Toulouse, Occitanie, France) for providing computing resources.

**TAXONOMIC APPENDIX – NEW HIGHER TAXA**

Hirpidinae Rougerie, **subfamilia nova**

http://zoobank.org/LSID######

Type genus: *Hirpida* Draudt, 1930

This subfamily comprises two Andean genera, *Hirpida* and *Hirpsinjaevia*. A suggested

combination of diagnostic features includes the following characters of the adults proposed by

Michener (1952) as distinctive for genus *Hirpida*: head with the sides of the frons slightly

convex; tarsomeres with spines on their ventral surface; well-developed arolia and pulvilli;

absence or reduction of vein 3A in hindwings; tegulae long, reaching the anterior median

angle of mesothoracic scutellum; anepisternal suture largely horizontal but directed slightly

downward posteriorly; male genitalia with free juxta and articulated valvae. The status of

these characters remains unknown in the recently described, rare genus *Hirpsinjaevia* as it

could not be examined for this study. Further comparative morphology and a formal

phylogenetic analysis of characters are needed to identify unique characters or more likely a

unique combination of characters that can accurately diagnose members of the subfamily

Hirpidinae.

Antistathmopterini Rougerie, **tribus nova**

http://zoobank.org/LSID######

Type genus: *Antistathmoptera* Tams, 1935

This tribe only includes species in the genus *Antistathmoptera*, all of which appear

restricted to Eastern Africa. Because of their rarity in collections, little is known of their

morphology, but they can be easily distinguished by a unique combination of adult characters:

long tailed hindwings; pointed forewing apex; both pairs of wings with a discal spot formed

of two contiguous hyaline areas.

Solini Rougerie, **tribus nova**

http://zoobank.org/LSID######

Type genus: *Solus* Watson, 1913

This tribe only includes the genus *Solus* which is restricted to the continental part of the

Oriental region. The following combination of characters in adults are diagnostic: vein d3 of

forewing discal cell missing, resulting in an open discal cell; lower branch of fork at the start

of forewing vein 1A+2A much thinner than the upper branch; extremity of axillary sclerite 1

of forewings rounded; anapleural suture of mesothorax reduced; galeae present, short and

broad.

**LITERATURE CITED**

Arcila D., Orti G., Vari R., Armbruster J.W., Stiassny M.L.J., Ko K.D., Sabaj M.H.,

Lundberg J., Revell L.J., Betancur-R R. 2017. Genome-wide interrogation advances

resolution of recalcitrant groups in the tree of life. Nat. Ecol. Evol. 1: 0020.

Balcazar-Lara M., Wolfe K.L. 1997. Cladistics of the Ceratocampinae (Lepidoptera:

Saturniidae). Trop. Lepid. Res. 8: 1-53.

Barber J.R., Leavell B.C., Keener A.L., Breinholt J.W., Chadwell B.A., McClure C.J.W., Hill

G.M., Kawahara A.Y. 2015. Moth tails divert bat attack: evolution of acoustic

deflection. PNAS. 112: 2812-2816.

Battisti A., Holm G., Fagrell B., Larsson S. 2011. Urticating hairs in arthropods: their nature

and medical significance. Annu. Rev. Entomol. 56: 203-220.

Bergsten J. 2005. A review of long-branch attraction. Cladistics. 21:163-193.

Betancur-R R., Arcila D., Vari R.P., Hughes L.C., Oliveira C., Sabaj M.H., Orti G. 2019.

Phylogenomic incongruence, hypothesis testing, and taxonomic sampling: the

monophyly of characiform fishes. Evolution. 73: 329-345.

Borowiec M. 2019. Convergent evolution of the army ant syndrome and congruence in big-

data phylogenetics. Syst. Biol. 68: 642-656.

Borowiec M., Rabeling C., Brady S., Fisher B., Schultz T., Ward P. 2019. Compositional

heterogeneity and outgroup choice influence the internal phylogeny of the ants. Mol.

Phylogenet. Evol. 134: 111-121.

Borowiec M.L. 2016. AMAS: a fast tool for alignment manipulation and computing of

summary statistics. PeerJ. 4: e1660.

Borowiec M.L., Lee E.K., Chiu J.C., Plachetzki D.C. 2015. Extracting phylogenetic signal

and accounting for bias in whole-genome data sets supports the Ctenophora as sister to

remaining Metazoa. BMC genomics. 16:987.

Bossert S., Murray E.A., Blaimer B.B., Danforth B.N. 2017. The impact of GC bias on

phylogenetic accuracy using targeted enrichment phylogenomic data. Mol. Phylogenet.

Evol. 111: 149-157.

Bouckaert R., Vaughan T., Barido-Sottani J., Duchêne S., Fourment M., Gavryushkina A.,

Heled J., Jones G., Kühnert D., De Maio N., Matschiner M., Mendes F.K., Müller N.F.,

Ogilvie H.A., du Plessis L., Popinga A., Rambaut A., Rasmussen D., Siveroni I., Suchar

M.A., Wu C.-H., Xie D., Zhang C., Stadler T., Drummond A.J. 2019. BEAST 2.5: An

advanced software platform for Bayesian evolutionary analysis. PLoS Comput. Biol.

15: e1006650.

Boussau B., Walton Z., Delgado J.A., Collantes F., Beani L., Stewart I.J., Cameron S.A.,

Whitfield J.B., Johnston J.S., Holland P.W.H., Bachtrog D., Kathirithamby J.,

Huelsenbeck J.P. 2014. Strepsiptera, phylogenomics and the Long Branch Attraction

problem. Plos One. 9: e107709.

Bouyer T. 2002. Description du mâle supposé d'*Eosia insignis* Le Cerf, 1911 (Lepidoptera

Saturniidae). Entomologia Africana. 7: 3-14.

Brechlin R 2020. A revision of the genus *Janiodes* Jordan, 1924 (Lepidoptera: Saturniidae;

Cercophaninae). Entomo-Satsphingia 13: 4-78.

Breinholt J.W., Earl C., Lemmon A.R., Lemmon E.M., Xiao L., Kawahara A.Y. 2018.

Resolving relationships among the megadiverse butterflies and moths with a novel

pipeline for anchored phylogenomics. Syst. Biol. 67: 78-93.

Carrijo-Carvalho L.C., Chudzinski-Tavassi A.M. 2007. The venom of the *Lonomia*

caterpillar: an overview. Toxicon. 49: 741-757.

Chaimanee Y., Chavasseau O., Beard K.C., Kyaw A.A., Soe A.N., Sein C., Lazzari V.,

Marivaux L., Marandat B., Swe M., Rugbumrung M., Lwin T., Valentin X., MAung-

Thein Z.-M., Jaeger J.-J. 2012. Late Middle Eocene primate from Myanmar and the

initial anthropoid colonization of Africa. PNAS. 109: 10293-10297.

Chazot N., Condamine F.L., Dudas G., Pena C., Kodandaramaiah U., Matos-Maravi P.,

Aduse-Poku K., Elias M., Warren A.D., Lohmman D.J., Penz C.M, DeVries P., Zdenek

F.F., Nylin S., Müller C., Kawahara A.Y., Silva-Brandao K.L., Lamas G., Kleckova I.,

Zubek A., Ortiz-Acevedo E., Vila R., Vane-Wright R.I., Mullen S.P., Jiggins C.D.,

Wheat C.W., Freitas A.V.L., Wahlberg N. 2021. Conserved ancestral tropical niche but

different continental histories explain the latitudinal diversity gradient in brush-footed

butterflies. Nat. Commun. 12: 5717.

Chen D.B., Zhang R.S., Jin X.D., Yang J., Li P., Liu Y.Q. 2021. First complete mitochondrial

genome of *Rhodinia* species (Lepidoptera: Saturniidae): genome description and

phylogenetic implication. Bull. Entomol. Res.: 1-10.

Ciminera M., Auger-Rozenberg M.A., Caron H., Herrera M., Scotti-Saintagne C., Scotti I., N.

T., Roques A. 2018. Genetic variation and differentiation of *Hylesia metabus*

(Lepidoptera: Saturniidae): Moths of public health importance in French Guiana and in

Venezuela. J. Med. Entomol. 56: 137-148.

Condamine F.L., Sperling F.A., Wahlberg N., Rasplus J.-Y., Kergoat G.J. 2012. What causes

latitudinal gradients in species diversity? Evolutionary processes and ecological

constraints on swallowtail biodiversity. Ecol. Lett. 15: 267-277.

Cornee J.J., Munch P., Philippon M., BouDagher-Fadel M., Quillevere F., Melinte-

Dobrinescu M., Lebrun J.F., Gay A., Meyer S., Montheil L., Lallemand S., Marcaillou

B., Laurencin M., Legendre L., Garrocq C., Boucard M., Beslier M.-O., Laigle M.,

Schenini L., Fabre P.-H., Antoine P.-O., Marivaux L. 2021. Lost islands in the northern

Lesser Antilles: possible milestones in the Cenozoic dispersal of terrestrial organisms

between South America and the Greater Antilles. Earth-Sci. Rev. 217: 103617.

Coster P.M.C., Beard K.C., Salem M.J., Chaimanee Y., Jaeger J.-J. 2015. New fossils from

the Paleogene of central Libya illuminate the evolutionary history of endemic African

anomaluroid rodents. Front. Earth Sci. 3: 00056.

Cruaud A., Delvare G., Nidelet S., Sauné L., Ratnasingham S., Chartois M., Blaimer B.B.,

Gates M., Brady S.G., Faure S., van Noort S., Rossi J.-P., Rasplus J.-Y. 2021. Ultra-

Conserved Elements and morphology reciprocally illuminate conflicting phylogenetic

hypotheses in Chalcididae (Hymenoptera, Chalcidoidea). Cladistics. 37: 1-35.

Cruaud A., Nidelet S., Arnal P., Weber A., Fusu L., Gumovsky A., Huber J., Polaszek A.,

Rasplus J.-Y. 2019. Optimised DNA extraction and library preparation for small

arthropods: application to target enrichment in chalcid wasps used for biocontrol. Mol.

Ecol. Resour. 19: 702-710.

Davis R. B., Javois J., Kaasik A., Ounap E., Tammaru T. 2016. An ordination of life histories

using morphological proxies: capital vs. income breeding in insects. Ecology. 97: 2112-

2124.

Dong Y., Dai F.Y., Ren Y.D., Liu H., Chen L., Yang P.C., Liu Y.Q., Li X., Wang W., Xiang

H. 2015. Comparative transcriptome analyses on silk glands of six silkmoths imply the

genetic basis of silk structure and coloration. BMC Genomics. 16: 203.

Duchêne D., Mather N., Van Der Wal C., Ho S. 2021. Excluding loci with substitution

saturation improves inferences from phylogenomic data. Syst. Biol. Syab075.

Economo E.P., Narula N., Friedman N.R., Weiser M.D., Guénard B. 2018. Macroecology and

macroevolution of the latitudinal diversity gradient in ants. Nat. Comm. 9: 1778.

Faircloth B.C. 2017. Identifying conserved genomic elements and designing universal bait

sets to enrich them. Methods Ecol. Evol. 8: 1103-1112.

Fogang Mba A.R., Kansci G., Viau M., Rougerie R., Genot C. 2019. Edible caterpillars of

*Imbrasia truncata* and *Imbrasia epimethea* contain lipids and proteins of high potential

for nutrition. J Food Compost Anal. 79: 70-79.

Garrocq C., Lallemand S., Marcaillou B., Lebrun J.F., Padron C., Klingelhoefer F., Laigle M.,

Munch P., Gay A., Schenini L., Beslier M.-O., Cornee J.-J., Mercier de Lépinay B.,

Quillévéré F., Boudagher-fadel M. 2021. Genetic relations between the Aves Ridge and

the Grenada back-arc basin, East Caribbean sea. J. Geophys. Res. Solid Earth. 126:

e2020JB020466.

Guindon S., Dufayard J.F., Lefort V., Anisimova M., Hordijk W., Gascuel O. 2010. New

algorithms and methods to estimate maximum-likelihood phylogenies: assessing the

performance of PhyML 3.0. Syst. Biol. 59: 307-321.

Hamilton C., St Laurent R., Dexter K., Kitching I., Breinholt J., Zwick A., Timmermans M.,

Barber J., Kawahara A. 2019. Phylogenomics resolves major relationships and reveals

significant diversification rate shifts in the evolution of silk moths and relatives. BMC

Evol. Biol. 19: 182.

Hamilton C.A., Winiger N., Rubin J.J., Breinholt J., Rougerie R., Kitching I.J., Barber J.R.,

Kawahara A.Y. 2021. Hidden phylogenomic signal helps elucidate arsenurine silkmoth

phylogeny and the evolution of body size and wing shape trade-offs. Syst. Biol.:

syab090.

Hartenberger J. 1998. An Asian Grande Coupure. Nature. 394: 321.

Harzhauser M., Kroh A., Mandic O., Piller W.E., Göhlich U., Reuter M., Berning B. 2007.

Biogeographic responses to geodynamics: A key study all around the Oligo-Miocene

Tethyan Seaway. Zool. Anz. 246: 241-256.

Huchon D., Chevret P., Jordan U., Kilpatrick C.W., Ranwez V., Jenkins P.D., Brosius J.,

Schmitz J. 2007. Multiple molecular evidences for a living mammalian fossil. PNAS.

104: 7495-7499.

Iturralde-Vinent M.A. 2006. Meso-Cenozoic Caribbean paleogeography: Implications for the

historical biogeography of the region. Int. Geol. Rev. 48: 791-827.

Ivanova N.V., Dewaard J.R., Hebert P.D.N. 2006. An inexpensive, automation-friendly

protocol for recovering high-quality DNA. Mol. Ecol. Notes. 6: 998-1002.

Jahren A.H. 2007. The Arctic Forest of the Middle Eocene. Annu. Rev. Earth Planet. Sci. 35:

509-540.

Jia D.-R., Bartish I.V. 2018. Climatic changes and orogeneses in the late Miocene of Eurasia:

The main triggers of an expansion at a continental scale? Front. Plant Sci. 9: 1400.

Kalyaanamoorthy S., Minh B.Q., Wong T.K.F., von Haeseler A., Jermiin L.S. 2017.

ModelFinder: fast model selection for accurate phylogenetic estimates. Nat. Methods.

14: 587-589.

Kassambara A. 2020. ggpubr: ggplot2 Based Publication Ready Plots. R package version

0.3.0 https://CRAN.R-project.org/package=ggpubr.

Katoh K., Standley D.M. 2013. MAFFT multiple sequence alignment software version 7:

improvements in performance and usability. Mol. Biol. Evol. 30: 772-780.

Kawahara A.Y., Plotkin D., Espeland M., Meusemann K., Toussaint E.F.A., Donath A.,

Gimnich F., Frandsen P.B., Zwick A., Reis dos M., Barber J.R., Peters R.S., Liu S.,

Zhou X., Mayer C., Podsiadlowski L., Storer C., Yack J.E., Misof B., Breinholt J.W.

2019. Phylogenomics reveals the evolutionary timing and pattern of butterflies and

moths. PNAS. 116: 22657–22663.

Kitching I.J., Rougerie R., Zwick A., Hamilton C.A., St Laurent R.A., Naumann S., Mejia

L.B., Kawahara A.Y. 2018. A global checklist of the Bombycoidea (Insecta:

Lepidoptera). BDJ. 6: e22236.

Klaus K.V., Matzke N.J. 2020. Statistical comparison of trait-dependent biogeographical

models indicates that Podocarpaceae dispersal is influenced by both seed cone traits and

geographical distance. Syst. Biol. 69: 61–75.

Kozlov A.M., Darriba D., Flouri T., Morel B., Stamatakis A. 2019. RAxML-NG: a fast,

scalable and user-friendly tool for maximum likelihood phylogenetic inference.

Bioinformatics, 35: 4453-4455.

Kumar S., Filipski A.J., Battistuzzi F.U., Pond S.L.K., Tamura K. 2012. Statistics and truth in

phylogenomics. Mol. Biol. Evol. 29: 457-472.

Kundu S.C., Kundu B., Talukdar S., Bano S., Nayak S., Kundu J., Mandal B.B., Bhardwaj N.,

Botlagunta M., Dash B.C., et al. 2012. Invited review nonmulberry silk biopolymers.

Biopolymers. 97: 455-467.

Landis M., Matzke N., Moore B., Huelsenbeck J. 2013. Bayesian analysis of biogeography

when the number of areas is large. Syst. Biol. 62: 789-804.

Lanfear R., Frandsen P.B., Wright A.M., Senfeld T., Calcott B. 2017. PartitionFinder 2: New

methods for selecting partitioned models of evolution for molecular and morphological

phylogenetic analyses. Mol. Biol. Evol. 34: 772-773.

Lemaire C. 2002. The Saturniidae of America. Vol. 4. Hemileucinae. 1388 pp. Goecke &

Evers Eds., Keltern, Germany.

LePage B., Yang H., Matsumoto M. 2005. The Evolution and biogeographic history of

Metasequoia. In: LePage B., Williams C.J., Yang H. eds. The Geobiology and Ecology

of Metasequoia. Dordrecht, The Netherlands, Springer: 3-114.

Licht A., Métais G., Coster P., İbilioğlu D., Ocakoğlu F., Westerweel J., Mueller M.,

Campbell C., Mattingly S., Wood M.C., Beard K.C. 2022. Balkanatolia: The insular

mammalian biogeographic province that partly paved the way to the Grande Coupure.

Earth Sci. Rev. 226: 103929.

Maechler M., Rousseeuw P., Struyf A., Hubert M., Hornik K. 2018. cluster: Cluster Analysis

Basics and Extensions. R package version 2.0.7-1.

Mai U., Mirarab S. 2018. TreeShrink: fast and accurate detection of outlier long branches in

collections of phylogenetic trees. BMC Genomics. 19: 272.

Marivaux L., Essid E.M., Marzougui W., Ammar H.K., Adnet S., Marandat B., Merzeraud

G., Tabuce R., Vianey-Liaud M. 2014. A new and primitive species of *Protophiomys*

(Rodentia, Hystricognathi) from the late middle Eocene of Djebel el Kébar, central

Tunisia. Palaeovertebrata. 38: e2.

Matzke N. 2014. Model selection in historical biogeography reveals that founder-event

speciation is a crucial process in island clades. Syst. Biol. 63: 951-970.

Matzke N.J. 2021. Statistical comparison of DEC and DEC+J is identical to comparison of

two ClaSSE submodels, and is therefore valid. OSF Preprints.

Michener C.D. 1952. The Saturniidae (Lepidoptera) of the western hemisphere: morphology,

phylogeny and classification. Bull. Am. Mus. Nat. Hist., 98: 341–501.

Minet J. 1994. The Bombycoidea phylogeny and higher classification (Lepidoptera,

Glossata). Insect. Syst. Evol. 25: 63-88.

Minh B., Hahn M., Lanfear R. 2020 New methods to calculate concordance factors for

phylogenomic datasets. Mol. Biol. Evol. 37: 2727-2733.

Minh B.Q., Nguyen M.A.T., von Haeseler A. 2013. Ultrafast approximation for phylogenetic

bootstrap. Mol. Biol. Evol. 30: 1188-1195.

Minh B.Q., Schmidt H.A., Chernomor O., Schrempf D., Woodhams M.D., von Haeseler A.,

Lanfear R. 2020. IQ-TREE 2: New models and efficient methods for phylogenetic

inference in the genomic era. Mol. Biol. Evol. 37: 1530-1534.

Mirarab S., Nguyen N., Guo S., Wang L.S., Kim J., Warnow T. 2015. PASTA: Ultra-Large

Multiple Sequence Alignment for Nucleotide and Amino-Acid Sequences. J. Comput.

Biol. 22: 377-386.

Morley R.J. 2003. Interplate dispersal paths for megathermal angiosperms. Perspect. Plant

Ecol. Evol. Syst. 6: 5-20.

Morley R.J. 2011. Cretaceous and Tertiary climate change and the past distribution of

megathermal rainforests. In Tropical Rainforest Responses to Climatic Change. Eds.

Bush M., Flenley J., Gosling W. Springer-Verlag, Berlin Heidelberg, Germany: 1-34.

Nässig W.A. 1991. Biological observations and taxonomic notes on *Actias isabellae* (Graells)

(Lepidoptera, Saturniidae). Nota lep. 14: 131-143.

Nässig W.A., Naumann S., Oberprieler R.G. 2015. Notes on the Saturniidae of the Arabian

Peninsula, with description of a new species (Lepidoptera: Saturniidae). Nachr.

Entomol. Ver. Apollo, N.F. 36: 31-38.

Nguyen L.T., Schmidt H.A., von Haeseler A., Minh B.Q. 2015. IQ-TREE: A fast and

effective stochastic algorithm for estimating maximum likelihood phylogenies. Mol.

Biol. Evol. 32: 268-274.

Oberprieler R.G., Nässig W.A. 1994. Tarn- oder Warntrachten - ein Vergleich larvaler und

imaginaler Strategien bei Saturniinen (Lepidoptera: Saturniinen). Nachr. Entomol. Ver.

Apollo, N.F. 15: 267-303.

Oberprieler R.G. 1997. Classification of the African Saturniidae (Lepidoptera) - The quest for

natural groups and relationships. Metamorphosis. Suppl. 3: 142-155.

Paradis E., Schliep K. 2018. ape 5.0: an environment for modern phylogenetics and

evolutionary analyses in R. Bioinformatics 35: 526-528.

Philippe H., Brinkmann H., Lavrov D., Littlewood D., Manuel M., Wörheide G., Baurain D.

2001. Resolving difficult phylogenetic questions: why more sequences are not enough.

PLoS Biol. 9: e1000602.

Philippon M., Cornee J.J., Munch P., van Hinsbergen D.J.J., BouDagher-Fadel M., Gailler L.,

Boschman L.M., Quillevere F., Montheil L., Gay A., Lebrun J.F., Lallemand S.,

Marivaux L., Antoine P.-O. 2020. Eocene intra-plate shortening responsible for the rise

of a faunal pathway in the northeastern Caribbean realm. PLoS One. 15: e0241000.

Pindell J., Kennan L., Maresch W.V., Stanek K., Draper G., Higgs R. 2005. Plate-kinematics

and crustal dynamics of circum-Caribbean arc-continent interactions: tectonic controls

on basin development in Proto-Caribbean margins. Geological Society of America

Special Paper, 394: 7–52.

R Core Team. 2018. R version 3.5.1 (Feather Spray): A language and environment for

statistical computing. R Foundation for Statistical Computing, Vienna, Austria. URL

https://www.R-project.org/.

Rambaut A., Drummond A., Xie D., Baele G., MA S. 2018. Posterior summarization in

Bayesian phylogenetics using Tracer 1.7. Syst. Biol. 67: 901-904.

Rasplus J.-Y., Rodriguez L.J., Sauné L., Peng Y.-Q., Bain A., Kjellberg F., Harrison R.D.,

Pereira R.A.S., Ubaidillah R., Tollon-Cordet C., Gautier M., Rossi J.-P., Cruaud A.

2021. Exploring systematic biases, rooting methods and morphological evidence to

unravel the evolutionary history of the genus *Ficus* (Moraceae). Cladistics. 37: 402-422.

Ree R., Sanmartin I. 2018. Conceptual and statistical problems with the DEC+J model of

founder☐event speciation and its comparison with DEC via model selection. J.

Biogeogr. 45: 741-749.

Ree R.H., Smith S.A. 2008. Maximum-likelihood inference of geographic range evolution by

dispersal, local extinction, and cladogenesis. Syst. Biol. 57: 4-14.

Regier J.C., Cook C.P., Mitter C., Hussey A. 2008. A phylogenetic study of the 'bombycoid

complex' (Lepidoptera) using five protein-coding nuclear genes, with comments on the

problem of macrolepidopteran phylogeny. Syst. Ent. 33: 175-189.

Regier J.C., Mitter C., Peigler R.S., Friedlander T.P. 2002. Monophyly, composition, and

relationships within Saturniinae (Lepidoptera : Saturniidae): Evidence from two nuclear

genes. Insect. Syst. Evol. 33: 9-21.

Robinson D., Foulds L. 1981. Comparison of phylogenetic trees. Math. Biosci. 53: 131-147.

Rogl F. 1999. Mediterranean and Paratethys. Facts and hypotheses of an Oligocene to

Miocene paleogeography (short overview). Geol. Carpath. 50: 339-349.

Romiguier J., Cameron S.A., Woodard S.H., Fischman B.J., Keller L., Praz C.J. 2016.

Phylogenomics controlling for base compositional bias reveals a single origin of

eusociality in corbiculate bees. Mol. Biol. Evol. 33: 670-678.

Romiguier J., Roux C. 2017. Analytical biases associated with GC-content in molecular

evolution. Front. Genet. 8: 00016.

Ronquist F. 1997. Dispersal-vicariance analysis: a new approach to the quantification of

historical biogeography. Syst. Biol. 46: 195-203.

Rossi J.-P. 2011. rich: an R package to analyse species richness. Diversity. 3: 112-120.

Rougerie R. 2005. Phylogeny and biogeography of the Saturniinae (Lepidoptera:

Bombycoidea, Saturniidae). PhD Thesis. Muséum National d'Histoire Naturelle, Paris,

France: 530 pp + 45 pl.

Rubin J.J., Hamilton C.A., McClure C.J.W., Chadwell B.A., Kawahara A.Y., Barber J.R.

2018. The evolution of anti-bat sensory illusions in moths. Sci. Adv. 4: eaar7428.

Rubinoff D., Doorenweerd C. 2020. In and out of America: Ecological and species diversity
in Holarctic giant silkmoths suggests unusual dispersal, defying the dogma of an Asian
origin. J. Biogeogr. 47: 903-914.

Shimodaira H. 2002. An approximately unbiased test of phylogenetic tree selection. Syst.
Biol. 51: 492-508.

Shimodaira H., Hasegawa M. 2001. CONSEL: for assessing the confidence of phylogenetic
tree selection. Bioinformatics 17: 1246-1247.

Smith P., Rios S., Atkinson K., Mielke O., Casagrande M. 2013. New Paraguayan specimens
and first confirmed phenological data for *Catharisa cerina* Jordan 1911 (Lepidoptera:
Saturniidae). Rev. Bras. Biocienc. 11: 349-351.

Sohn J.C., Labandeira C., Davis D., Mitter C. 2012. An annotated catalog of fossil and
subfossil Lepidoptera (Insecta: Holometabola) of the world. Zootaxa 3286:1-116.

Sota T., Takami Y., Ikeda H., Liang H., Karagyan G., Scholtz C., Hori M. 2022. Global
dispersal and diversification in ground beetles of the subfamily Carabinae. Mol.
Phylogenet. Evol. 167: 107355.

Stamatakis A. 2014. RAxML version 8: a tool for phylogenetic analysis and post-analysis of
large phylogenies. Bioinformatics 30: 1312-1313.

Struck T.H. 2014. TreSpEx - Detection of misleading signal in phylogenetic reconstructions
based on tree information. Evol. Bioinform. Online. 10:51-67.

Tagliacollo V.A., Lanfear R., Townsend J. 2018. Estimating Improved Partitioning Schemes
for Ultraconserved Elements. Mol. Biol. Evol. 35: 1798-1811.

Vandenberghe N., Hilgen F.J., Speijer R.P. 2012. Chapter 28. The Paleogene Period. In:
Gradstein F.M., Ogg J.G., Schmitz M.D., Ogg G.M. eds. The Geologic Time Scale.
Elsevier: 855-921.

Westerhold T., Marwan N., Drury A.J., Liebrand D., Agnini C., Anagnostou E., Barnet

J.S.K., Bohaty S.M., De Vleeschouwer D., Florindo F., Frederichs T., Hodell D.A.,

Holbourn A.E., Kroon D., Lauretano V., Littler K., Lourens L.J., Lyle M., Palike H.,

Rohl U., Tian J. Wilkens R.H., Wilson P.A., Zachos J.C. 2020. An astronomically dated

record of Earth's climate and its predictability over the last 66 million years. Science.

369: 1383-1387.

Xing Y., Ree R.H. 2017. Uplift-driven diversification in the Hengduan Mountains, a

temperate biodiversity hotspot. PNAS. 114: E3444-E3451.

Yang Z., Rannala B. 2006. Bayesian estimation of species divergence times under a

molecular clock using multiple fossil calibrations with soft bounds. Mol. Biol. Evol.,

23: 212-226.

Yang J., Wang Y.-F., Spicer R.A., Mosbrugger V., Li C.-S., Sun Q.-G. 2007. Climatic

Reconstruction at the Miocene Shanwang Basin, China, Using Leaf Margin Analysis,

CLAMP, Coexistence Approach, and Overlapping Distribution Analysis. Am. J. Bot.

94: 599-608.

Zhang C., Rabiee M., Sayyari E., Mirarab S. 2018. ASTRAL-III: polynomial time species

tree reconstruction from partially resolved gene trees. BMC Bioinformatics. 19: 153.

Zhang J., Sun B., Zhang X.J. 1994. Miocene Insects and Spiders from Shanwang, Shandong.

Beijing, China. Beijing Science Press. v + 298 pp.

Zhong B., Betancur-R R. 2017. Expanded taxonomic sampling coupled with gene genealogy

interrogation provides unambiguous resolution for the evolutionary root of

angiosperms. Genome Biol. Evol. 9: 3154-3161.

Zhu Q. 2014. AfterPhylo. A Perl script for manipulating trees after phylogenetic

reconstruction. Available from https://github.com/qiyunzhu/AfterPhylo/.

**AUTHOR CONTRIBUTIONS**

Designed the study: AC, JYR, PA, RR

Obtained funding: AC, CLV, FLC, JM, JYR, ME, RR, SL, TD

Contributed samples: CLV, JYR, PDNH, RR, TD

Identified samples: IJK, JYR, RR, TD

Performed laboratory work: AC, DG, JYR, PA, RR, SN, YC

Analyzed data: AC, JYR, PA, RR

Discussed results: AC, CLV, FLC, IJK, JM, JYR, LBM, ME, PA, RR, SL, TD

Drafted the manuscript: AC, JYR, PA, RR

All authors commented on the manuscript

**DATA AVAILABILITY**

Fastq paired reads for analyzed samples are available as a NCBI Sequence Read

Archive (ID#PRJNA XXXXX). Custom script for random sampling of sites within data sets

is available from https://github.com/acruaud/saturniidae_phylogenomics_2022. Data sets

(concatenated UCEs) and trees have been uploaded on Dryad

(http://dx.doi.org/10.5061/dryad.[NNNN]).

1004

**TABLE AND FIGURE LEGENDS**

**Table 1. Summary of mean divergence time estimates and 95% highest posterior density**

**(HPD) for groups or colonization events discussed.** Mean and median ages with 95% HPD

for all nodes are provided in Table S4. Units are in Myr. Our new classification is used for

taxon names. ACH = Arsenurinae + Ceratocampinae + Hemileucinae; BSS = Bunaeinae +

Salassinae + Saturniinae.

1011

**Figure 1. Conflicting topologies observed after analysis of the initial data set (a) and**

**results of the gene genealogy interrogation (GGI) approach (b)**. In panel (a), topology *A* is

supported by IQ-TREE analyses; *Abis* is supported by ASTRAL90; *B* is supported by

ASTRAL70 and *C* is supported by ASTRAL50. All corresponding trees are available in

Figure S1. In simplified topologies, Hemileucinae is used for Hemileucinae minus *Hirpida*

and *Hirpsinjaevia* and Hirpi/Hirpsi stands for *Hirpida + Hirpsinjaevia.* For the GGI approach,

Hemileucinae (minus *Hirpida* and *Hirpsinjaevia*) + Ceratocampinae + Arsenurinae;

Saturniinae + Salassinae; *Hirpida + Hirpsinjaevia* and Agliinae were considered as different

clades and trees were inferred using each of the competing topologies as multifurcating

constraint trees. The structure of the backbone was fixed, but taxa within clades were free to

move around. Topology *A* was considered equivalent to *Abis* as we focused on sister taxa

relationships of Agliinae / *Hirpida + Hirpsinjaevia* with either Hemileucinae +

Ceratocampinae + Arsenurinae or Saturniinae + Salassinae. In panel (b), average values of

GC content and saturation (R-squared of the regression between pairwise distances calculated

from aligned sequences and branch length) for loci that significantly supported topologies

*A/Abis* (i.e., Pvalue of the AU test $\leqq$ 0.05; N loci = 189; Table S3) or not (Nloci=796) were

compared using a randomization test (N randomization = 9999; c2m function of the R

package *rich*) and P values are reported on graph.

**Figure 2. Saturniidae tree of life.** The classification proposed in this study is mapped on the

chronogram (MCMCtree, uncorrelated relaxed clock model) inferred from the topology

selected after exploration of systematic bias (i.e., the IQ-TREE tree inferred from the 50%

least saturated loci). Red dots at nodes identify poorly supported nodes (SHaLRT<80 or

UFBoot<95). Abbreviations used: A = Antistathmopterini; Agl = Agliinae; Al = Almeidaiini;

Bathy = Bathyphlebiini; C = Cercophanini; Cer = Cercophaninae; Cith = Citheroniini; E =

Eudaemoniini; Eo = Eochroini; Hir = Hirpidinae; J = Janiodini; Lono = Lonomiini;

Oxyteninae; P = Pseudapheliini; Poly = Polythysanini; Sal = Salassinae; So = Solini; Urot =

Urotini.

**Figure 3. Global historical biogeography of Saturniidae.** Ancestral ranges were inferred

with the BAYAREALIKE+J model (best-fit and most plausible model according to empirical

considerations; see text) as implemented in *BioGeoBEARS*. Alternative inferences of ancestral

ranges are provided in Figure S5. The chronogram is derived from the species-level

chronogram obtained with the uncorrelated relaxed clock model implemented in MCMCtree

(see Fig. 2) that was pruned to keep only one specimen per genus/subgenus. The classification

proposed in this study is used to annotate the tree. Colored boxes following each terminal

name represent biogeographical regions A to I as illustrated in the upper-left map; colored

arrows refer to inferred dispersal/colonization events. Abbreviations used: A = Agliinae; Ars

= Arsenurinae; C=Cercophaninae; H = Hirpidinae; O = Oxyteninae; S = Salassinae.

PPH=Pliocene, Pleistocene and Holocene. Illustrative photograph = *Antherina suraka*

(Saturniinae); image courtesy of Armin Dett.

Saturniinae, *Rothschildia*

Bunaeinae, *Gonimbrasia*

Salassinae, *Salassa*

Hemileucinae, *Leucanella*

Ceratocampinae, *Eacles*

Arsenurinae, *Caio*

Hirpidinae, *Hirpida*

Agliinae, *Aglia*

Cercophaninae, *Cercophana*

Oxyteninae, *Oxytenis*

63.3 Ma
(58.7-68.5)

51.9 Ma
(50.7-52.6)