Generation of raptor diversity in Europe: linking speciation with climate

changes and the ability to migrate

Running title: climate changes and raptor diversification

Juan J. Negro[1]

Eduardo Rodríguez-Rodríguez[2]

Airam Rodríguez[3,4,5]

Keith Bildstein[6]

[1]Estación Biológica de Doñana-CSIC, Avda. Americo Vespucio 26. 41092. Sevilla. Spain.

[2]Department of Integrated Sciences, Faculty of Experimental Sciences, University of Huelva,
Huelva, Spain.·

[3]Grupo de Ornitología e Historia Natural de las islas Canarias (GOHNIC), C/La Malecita s/n,
Buenavista del Norte, Canary Islands, Spain.

[4]Terrestrial Ecology Group (TEG-UAM), Department of Ecology, Universidad Autónoma de
Madrid, Madrid, Spain.

[5]Centro de Investigación en Biodiversidad y Cambio Global (CIBC-UAM), Universidad Autónoma
de Madrid, Madrid, Spain.

[6]116 Village Drive, Blandon, PA 19510, USA

Corresponding author: Juan J. Negro. E-mail: negro@ebd.csic.es

Acknowledgements

We thank M.A. Rojas, M.Cayuela, J.M. Irastorza, J.M. Sayago and A. Kovacs for sharing pictures
included in Figures 1 and 2, and Figure S2. We also thank all the photographers who have
shared pictures under Creative Commons licenses. This study received no funding.

Key Words

Glacial cycles, migration, interspecific competition, species richness, birds of prey, speciation

Abstract

Aim: Europe should be considered as a diversity hotspot for diurnal raptors, but just during the
breeding season, as it holds the higher proportion of transcontinental migratory species of any
landmass, and the area becomes depleted during the winter period. This study will test the
hypothesis that the high diversification of the raptor assemblage in Europe is a recent event,
occurring mainly during the Quaternary, and that closely related species sharing the same
trophic niches can only coexist in sympatry during the breeding period, when food availability is
higher.

Location: Continental Europe*.*

Taxon: Diurnal birds of prey (Accipitriformes and Falconiformes).

Methods: A consensus molecular phylogeny for the 38 regular breeding species of raptors in
Europe was obtained from BirdTree. For the same species, a trophic niche cluster dendrogram
was constructed. Size and migratory strategy were introduced in the resulting phylogeny, where
trophic groups were also identified.

Results: A total of 16 trophic groups were identified. Multispecific trophic groups tended to be
composed of reciprocal sister species, while monospecific groups (only three) were composed
of highly specialized species. According to time calibrated phylogenies, the speciation events
took place during the glacial cycles of the Quaternary period in a majority of cases. During the
non-breeding season, the smaller species in every trophic group migrate to sub-Saharan Africa,
whereas larger species are either non-migratory or perform shorter migrations within Europe
and/or northern Africa.

Main conclusions: This investigation illustrates how the rich assemblage of diurnal birds of prey
in Europe, more diverse and more migratory than the North American assemblage at equivalent
latitudes, has emerged recently due to the multiplication of look-alike species with similar
trophic ecologies, mainly in climate refugia during cold periods. In the non-breeding season,
when shared food resources are limited, smaller species migrate to Africa and alleviate
competition.

Introduction

Raptor migration is a conspicuous phenomenon in many locations globally (Bildstein,
2018). Numerous raptors, of 183 out of 313 (58%) species worldwide, abandon their breeding
grounds and, often gathering in large flocks, move away at predictable times and places to reach
distant areas located sometimes in different continents (Bildstein et al., 2007; Ferguson-Lees &
Christie, 2001). Raptor migration has been extensively monitored to assess long-term
population trends and to determine raptor flight behavior, migration corridors, as well as
breeding and non-breeding grounds (Bildstein, 2006). However, much remains to be
investigated concerning the causes and function of raptor migration, or why a majority of
species are migratory but still others remain sedentary. Recent molecular advances have shown
that there are in fact two distantly related groups of diurnal raptors, convergent in their
morphology and general ecology (Jarvis et al., 2015). On one side, are the true falcons and allies,
in the Order Falconiformes. On the other side, eagles, hawks, kites, harriers, and vultures
constitute the Order Accipitriformes, with the New World vultures (Family Cathartidae or Order
Cathartiformes, depending on the source, Del Hoyo et al., 1994; Del Hoyo, 2020) forming a sister
group of all remaining Accipitriformes (Jarvis et al., 2015). According to Ericsson (2011), the early
radiation of Falconiformes took place in South America, whereas Accipitriformes emerged in
Africa. Today, these two orders have gained a cosmopolitan distribution and species of the two
orders tend to co-occur in most environments. There are many migratory species or populations
within both Falconiformes and Accipitriformes (Bildstein, 2006) in both the Nearctic and the
Palearctic , implying that long-distance migratory behavior has evolved multiple times and that
it is a rather plastic trait (Nagy et al., 2017).

Europe is the land mass with the highest proportion of migratory species of birds of prey
in the World (Bildstein, 2006), and there is also a large absolute number of species considering
its relatively small size compared to the remaining of the Palearctic and also the Nearctic (see
results). Why this is so has never been explained nor explored so far. Also overlooked, even
though it should be obvious glancing at any bird guide (e.g., Svensson et al., 1999) or raptor
biology book (Ferguson-Lees & Christie, 2001), is the fact that there are numerous groups
(mainly duets and trios) of morphologically and ecologically similar raptorial species in the
European portion of the Western Palearctic. The species conforming these groups may look so
similar in appearance that their field identification at a distance may be challenging (Negro,
1991; Forsman, 2016). We refer to pairs of species such as the Eurasian (*Falco tinnunculus*) and
the lesser kestrels (*F. naumanni*), the lesser and greater spotted eagles (*Clanga pomarina* and *C.
clanga*), or trios including, for instance, Montagu's harrier (*Circus pygargus*), hen harrier (*C.
cyaneus*) and pallid harrier (*C. macrourus*).

Here we aim to link raptor migration, interspecific competition and finally diversity and
speciation in the European diurnal raptor assemblage, including both Falconiformes and
Accipitriformes. Controlling for phylogeny and considering some regularities in size differences
and migratory behaviour, we will test the hypothesis that the raptor community in Europe has
recently experienced a diversification of breeding species, mainly during the Quaternary.
Additionally, the coexistence of multiple look-alike species in the same trophic niches is only
possible during the breeding season (i.e., spring and summer), when food resources are less
limiting. In support of our hypothesis, that implies recent speciation events, we have gathered
information on the timing of species splitting derived from time-calibrated molecular
phylogenies (Fuchs et al., 2015; Mindell et al., 2018).

Methods

Study area and selected species

This study focuses on the 38 diurnal raptor species breeding regularly and historically in
continental Europe (see Table 1). We use the term Europe as defined in the seven-continents
model (i.e., Europe separated from Asia as a conglomerate of peninsulas delineated by the
Mediterranean sea to the South, the Atlantic ocean to the West, the Northern sea in high
latitudes, and the Ural mountains, the Caucasus and the Black and Caspian seas to the East)
because it suits well the differences in migrating behaviour of birds including raptors in northern
and temperate areas (e.g., Herrera, 1978; Newton & Dale, 1996; Newton & Dale, 1998, Bildstein,
2006). Using continental Europe as a biogeographical area allows us to cleanly separate
sedentary, migrants within continental Europe, and sub-Saharan bird species migrating through
the Mediterranean flyways (i.e., Straits of Gibraltar, Messina and Bosphorus). The Western
Palearctic, a zoogeographical region widely used in bird studies (e.g., Cramp & Simmons, 1979;
Perrins & Snow, 1998), and that fully encompasses Europe, introduces, however, African and
Asian raptorial species that would add noise to our analyses. These species, either at the
southern limit of the Western Palearctic in the African continent or at the eastern border with
Asia, are all rare breeders or vagrants in the area that have never established breeding
populations in continental Europe. Raptors of the Western Palearctic which are clearly African
taxa include at least (Svensson et al., 1999), the lappet-faced vulture (*Torgos tracheliotos*), the
dark chanting goshawk (*Melierax metabates*), the barbary falcon (*Falco pelegrinoides*), the sooty
falcon (*Falco concolor*), the tawny eagle (*Aquila rapax*), and Verreaux's eagle (*A. verreauxi*).

Phylogeny and trophic niche

As a first step to determine the relationships among the different species under study, a consensus phylogeny was constructed using FigTree v 1.4.4 (Rambaut & Drummond, 2018) and 100 permutations obtained from BirdTree (Jetz et al., 2012; Jetz et al., 2014). Addressing migratory behaviour in birds requires not only the consideration of biogeographic aspects, but also a knowledge of diet specializations and prey availability (Nagy & Tokolyi, 2014). A trophic niche cluster dendrogram, based on Euclidean distances, was constructed using a dataset with data of trophic niche variables of each species, including hunting or scavenging behaviour, prey type and size (i.e., small birds, large birds, small mammals, large mammals, arthropods, herps or fishes), scavenging strategy (small pieces or big pieces) and foraging method (perch hunting, foraging in open areas, forest ambusher). These data can be consulted in Supplementary Material Appendix 1. The software utilised to construct the dendrogram was in R statistics (R Core Team, 2013). Presentation of the Figures was improved with Inkscape 1.0.1 (Harrington et al., 2020).

Relating body mass and trans-Saharan migration

137   We compiled the following information of the raptor species: the minimum and maximum value of both total length and wingspan (2) from del Hoyo et al. (1994), the minimum and maximum values of length, tail, and tarsus, as well as ranges of male and female body masses (4 mass values in total) from Ferguson-Lees & Christie (2001). To estimate body mass, we averaged the minimum and maximum values for males and females. To estimate bird size, we ran a principal component analysis (PCA) on the above mentioned fourteen centered and scaled morphometric variables. The first principal component was used as a body size index. The first principal component retained 90.1% of variation (see PCA details in Table S1). The fourteen morphometric variables showed the same sign for their factor loadings and highly significant correlations to the first principal component (Figure S1). Factor loadings of the fourteen morphometric variables and importance of the components are given in Table S1. Pairwise correlations of the first principal component (body size index) and seven morphometric variables (i.e. those representing the lower ranges of each morphometric) are shown in Figure S1.

151   Based on results from trophic niche clustering, groupings of species sharing a similar niche were established. These groups were identified on the previously built phylogeny. Using body mass data and migratory strategy (Table 1), we established two categories: smaller raptors inside their groups and larger raptors inside their groups, and estimated the proportion of trans-continental migratory species among large and small species. Significance of the differences in frequency of migration for both groups of large and small species were tested using a contingency Table. In addition, we plotted the mass of paired migratory and resident species in the foraging and phylogenetic groups that we had previously identified (Figure 3a). By adding a diagonal line (y = x), the resulting plot graphically shows in which cases the migratory species was either larger (above the diagonal) or smaller (below the diagonal) than its resident or less-migratory counterpart. We additionally plotted the previously described body size index, because the anatomical design of migratory species is known to be slender, i.e., much longer wings, or different wing loads, compared to non-migratory species (Figure 3b).

164   To build the maps displayed in Figure 4, distributions ranges of raptor species were downloaded from the IUCN Red List Data (IUCN, 2021) and imported in *Qgis* (v3.20) (Qgis Development Team, 2021). We deleted passage areas for migrating species and mapped pairs and trios of species according to the analysis of trophic niche.

168   **Results**

169   **Foraging groups**

170   Based on Euclidean distances of behavioural and dietary variables, we obtained 16
171   trophic or ecological niche groups, including three monospecific niches (i.e., the reed-bank
172   specialist *Circus aeruginosus*, the snake eating *Circaetus gallicus* and the himenopteran-eating
173   *Pernis apivorus*), 7 groups of two species, 4 groups of three species, one group of 4 species,
174   and one group of five species conformed by the bird-eating falcons (Table 1, Figure 1, Figure
175   S2).

176   **Phylogenetic relationships of European raptors**

177   The resulting phylogeny for the European raptor species matches well with the
178   obtained trophic groups in a majority of cases, except for the *Pandion haliaetus/Haliaeetus
179   albicilla* and *Aquila fasciata/Hieraaetus pennatus* species pairs, which are polyphyletic (Figure
180   2).

181   **Implications of size and migration strategy**

182   A majority of the smaller species inside trophic groups reach Africa during migration
(81% of cases, n = 16, and only 19% do not. On the contrary, only 20% (4 out of the 20) of the
larger species inside trophic groups perform a trans-Saharan migration (i.e., *Clanga clanga*, *Falco
eleonorae, Circus macrourus* and *Buteo rufinus*). However, these "large" species, except F.
eleonore, which overwinters in Madagascar, perform a much shorter migration than their
smaller counterparts in their respective groups (see Figure 4). The difference in migratory
behaviour among larger and smaller species is highly significant (χ^2=11,2754, P=0.0057, see
Figure 3).

190   When plotting the body masses for paired species derived from the foraging groups than
were established above (Figure 3a), the values fall below the diagonal of the graph (x axis, mass
of less migratory species, y axis, mass of more migratory species in the pair). This demonstrates
that the fully or more migratory species are systematically smaller than the less migratory or
resident species. Essentially the same results are obtained when instead of plotting body
masses, body size indices were plotted (Figure 3b). Larger species are the less migratory ones in
the species pairs. The distribution ranges for a selection of the species pairs that we considered
in the above plots have been mapped in Figure 4. The less-migratory species remain within
Europe (although it has to be noted that some species may have resident populations in Africa
such as the peregrine falcon and the Eurasian kestrel), whereas the fully migratory species have
non-overlapping breeding (in Europe) and non-breeding ranges (in Africa).

201   **Timing of the splits from the common ancestor**

202   According to the time-calibrated phylogenies of falcons (Falconiformes) on the one hand
(Fuchs et al. 2015) and of all other birds of prey (eagles, hawks, vultures and kites in
Accipitriformes) on the other hand (Mindell et al., 2018), six of our species groupings (both in
the phylogeny and in the trophic cladogram) originated in the Quaternary (i.e., the last 2.5
million years) as sister species. These groupings are the following: 1. The red and the black kite.
2. The two spotted eagles, lesser and greater. 3. The two imperial eagles (*A. heliaca* and *A.
adalberti*), that we have matched ecologically to the most distantly related *A. chrysaetos*. 4. The
three European buzzards: common, rough legged and long-legged. 5. The two late-breeding
falcons, dependent on the return migration of passerines for reproduction: hobby and

Eleonora's falcon. 6. The large European falcons: Lanner, Saker and Gyrfalcon. Apart from this,
we may add that the two colonial and tree-nesting falcons of temperate Eurasia, i.e., the red-
footed falcon in the Western Palearctic and Amur falcon in the Eastern Palearctic, which are also
sister species splitting from a common ancestor whose range may have spanned the temperate
forests of the whole of Eurasia.

The remaining groupings may have formed earlier in time, again according to Fuchs et
al (2015) and Mindell et al. (2018): 1. the Eurasian kestrel and the lesser kestrel emerged as
species in the kestrel radiation of the Pliocene, about 4-5 million years (Negro et al. 2020). 2.
The harriers (hen, Montagu´s and pallid) may have radiated about 5 million years ago. 3. The
hawks of the genus *Accipiter* radiated much earlier, and the common ancestor of the northern
goshawk and both the European sparrowhawk and levant sparrowhawk possibly evolved 10-15
million years ago in the Miocene. 4. The two small and ventrally white eagles, Bonelli's and
booted, emerged from a common ancestor living in the Pliocene 4 million years ago. 5. The two
large European vultures, griffon and cinereous, shared a common ancestor about 9 million years
ago. 6. Coalescence time for Bearded and Egyptian vultures, which are sister groups to each
other, is about 12 million years.

Discussion

The diversity of species of any given taxa in a land mass typically depends on its own
evolutionary history, the interaction with other species in the area, and the past and present
physical patterns of the environment (e.g., Hanssen & Urban, 1992; Mönkkönen, 1994). This
work was prompted by the fact that the number of breeding species of diurnal raptors in
continental Europe (38 species in a land-mass of 10,180,000 km^2) is higher than the number of
breeding species of raptors in Canada and continental US combined (30 species in about
19,000,000 km^2, including the 3 endemic New World vultures), and also by the disproportionate
number of transcontinental migratory species in Europe (Bildstein 2006, see below). The
latitudinal range is much wider in the American side and includes large areas below 35° North
latitude in the southern states of the US that are not represented in continental Europe.
However, this does not bring over a significant number of tropical species that do inhabit Mexico
and Central America (the total number of raptorial species in the latter area rises to 69 species,
Clark & Schmitt, 2017).

Seven species are shared between Europe and North America: five Holarctic raptor
species, the gyrfalcon, the merlin, the golden eagle, the rough-legged hawk, and the northern
goshawk, plus two cosmopolitan species, the peregrine falcon and the osprey, which hold
breeding populations in all continents except Antarctica (Ferguson-Lees & Christie, 2001).
Leaving aside these few shared species, mainly from northern or mountainous regions, there is
a marked difference among Europe and Canada-US combined in the number and proportion of
regular transcontinental migratory species, which is much higher in Europe (22 species crossing
the Mediterranean Sea to Africa in the non-breeding season) than in the above mentioned area
of North America, where only five species make the bulk of the migrants moving down to South
America (i.e., *Ictinia mississippiensis*, *Buteo swainsoni*, *B. platypterus*, *Cathartes aura* and
Pandion haliaetus) (Bildstein, 2006).

Our results show that there is a duplicity, or even multiplicity of species in the same
trophic and ecological niches in Europe. In many cases the species involved are so similar that
identification at a distance may be difficult (Forsman, 2016). This similarity may be due to shared
ancestry in congeneric species, such as in the duet conformed by *C. pomarina* and *C. clanga*, or

in the case of *Milvus milvus* and *M. migrans* (Mindell et al., 2018), but it may be due to convergence in more distantly related species such as *A. fasciata* and *H. pennatus*. Duplicated species are not apparent in North America, where the two more diverse genera (*Buteo* and *Falco*), have rather distinct species in terms of size and ecology. That Europe has been a recent hotspot for raptor speciation is also supported by the pattern of endemism (Weir and Schluter, 2004) in the group: six species are only or mainly distributed in Europe (i.e., red kite, honey buzzard, lesser spotted eagle, Spanish imperial eagle, Montagu's harrier and Eleonora's falcon). Five other species breed in Europe and more into the East, but may have a Western Palearctic origin: lesser kestrel (with the older fossil found in the Iberian Peninsula, Negro et al., 2020), red-footed falcon (sister species of the Amur falcon breeding in eastern Asia, as commented above), common buzzard (the nominal subspecies is sedentary across Europe, and only the Eastern subspecies *B. b. vulpinus* migrates to Africa), and finally the levant sparrowhawk, and the western marsh harrier.

Although our work does not focus on the evolution of raptor migration, our results shed light on the biogeographical origin of at least some of the migratory species. A majority of the duplicated species would adhere to the "northern origin" hypothesis (Bell, 2000; Bruderer and Selewski, 2008, Nagy et al., 2014), according to which harsh winter conditions would impel northern species to move to Africa in the non-breeding season. Some species in the system, such as the short-toed eagle, the Egyptian vulture and the bearded vulture, and two small eagles, Bonelli's and booted eagles, may have an afro-tropical origin, as they belong to phylogenetic groups that are highly diverse in Africa (Mindell et al., 2018). None-the-less, the migratory species among them may have acquired the migratory behaviour after colonizing Europe and competing with other species there.

Taken together, our results support the notion that the current raptor assemblage in Europe experienced a recent enrichment, and may be considered a hotspot accounting for its area and geographic location, and particularly by comparison with the smaller raptor community in much larger North America. This does not mirror the situation for the whole of the avifaunas, as there are about 556 bird species in Europe (Keller et al. 2021), compared to 740 in North America to the north of Mexico (Cornell lab of Ornithology, https://www.notesfromtheroad.com/roam/how-many-birds-north-america.html). The North American avifauna contains a plethora of insectivorous passerines with a probable tropical or subtropical origin (i.e., Parulidae, Turannidae, Vireonidae, Icteridae) which exploit the insect-rich summer of North America and migrate to the south in the winter (Snow 1978). The European migratory songbirds are fewer and belong to families (i.e., Sylviidae, Turdidae, Muscicapidae and Motacillidae) whose evolutionary centres are Palearctic but not African (Snow 1978). As a result, the assemblages of forest passerine birds of the Nearctic region are more diverse than those of the Western Palearctic, thus including Europe (Mönkkönen 1994). Other taxa follow the same pattern: Western North America contains, for instance, 41% more mammal species than Europe (Mönkkönen & Viro, 1997).

One mechanism that may explain the generation of raptor diversity in Europe is the existence of climate-driven vicariant events during the glacial cycles of the Pleistocene (Voelker 2010), and before during the Pliocene or earlier in the Miocene. It has been suggested, for instance, that the genus *Falco* radiated during the Miocene several million years ago, coincidental with a period of increased aridity favouring grasslands (Zachos et al., 2001, Fuchs et al. 2015). Glacial cycles have been invoked as drivers of speciation for numerous bird taxa in both the Old and the New World: fragmentation of populations, isolation in refugia during glacial

times and subsequent reunion without admixture is a common model for explaining the origin
of closely-related bird species (Bermingham et al., 1992 Weir and Schluter, 2004, Lovette, 2005).
Such a model was in fact invoked to explain the emergence as separate species of the Eastern
Imperial Eagle and the Spanish imperial eagle (Ferrer and Negro, 2004). The latter is the species
in the genus *Aquila* with the smallest distribution range, as it is endemic to the Iberian peninsula.

Lacking enough dated fossils, the recent avian speciation during the Pleistocene glacial
cycles has remained controversial for some groups (Lovette, 2005; Zink et al., 2004; Wang et al.,
2018). However, recent advances in molecular techniques, such as whole-genome sequencing,
clearly support that many bird species suffered dramatic expansions and contractions during the
Quaternary coincidental with climate changes (Nadachowska-Bryzska et al., 2015). These
population changes may have prompted speciation.

The generation of sister species of birds of prey may be explained by vicariance events.
The difference among Europe and North America possibly has to do with the different effects of
the glaciation in the two land masses, and the alignment of the major mountain barriers that
may have acted as geographical barriers, with a north-south component in the Nearctic and
east-west in the Palearctic (Albach et al., 2006). In North America, numerous pairs of sister bird
species tend to have east-west distributions (e.g., Lovette, 2005). In Europe, the existence of
several southern peninsulas and islands offered different refugia and thus more possibilities for
population fragmentation and isolation (Covas & Blondel, 1998). After ice cover retreated during
the interglacials, isolated populations might have expanded and established contact but, in
some cases, without genetic admixture. However, closely related species of predators coexisting
in sympatry would compete dearly in the harsh winter season. Competition would be resolved
by the less competitive species moving away and thus migrating to a different continent. This
may also be seen as several concurrent cases of character displacement, with competing species
diverging in size precisely to diminish potential competition (Dayan & Simberloff, 1998). Also of
interest may the fact that the assemblage of European carnivores (Mammalia) is much smaller,
with 22 species than the North American assemblage, with 45 species (Burgin et al., 2020).
Lessened potential competition with mammalian carnivores may have facilitated the
diversification of European breeding raptors. None-the-less, competition with other predators
alone is not the only possible explanation, as Bergmann's rule (Bergmann 1847), for instance,
would also predict larger bird species having a thermoregulatory advantage compared to smaller
ones during the cooler non-breeding season (Salewski & Watt, 2017). As additional support to
our hypothesis, the three New World vultures of North America conform to the size and
migration "rules", with the largest species, the California Condor, being sedentary, and the much
smaller species turkey and black vultures being migratory (Ferguson-Lees & Christie 2001).

An extraordinary case of rapid increase of size in the absence of competitors, therefore
linking body size to competition in the raptor group, is provided by two extinct birds of prey in
New Zealand (Knapp et al., 2019): the largest known eagle in the world, Haast's eagle *Hieraaetus*
moorei, and the largest known harrier in the world, Eyles' harrier *Circus teauteensis*. Both
derived from Australian vagrants, the much smaller *Hieraaetus morphnoides* and *Circus*
assimilis, respectively. These two concurrent cases of island gigantism occurred in the early
Pleistocene when the clearing of closed forests due to climate changes allowed for successful
colonization of Australian raptor immigrants hunting in open woodlands and grasslands (Knapp
et al., 2019). The fact that body size is in turn related to migration propensity and ability is also
supported at the intraspecific level. Raptors tend to present reversed sexual dimorphism, as
females are larger than males (Newton, 1979). Numerous raptor species show different

migration timings and non-breeding distributions, although it is true that there is variation in which sex migrates first (both out-bound and in-bound) or overwinters in more northerly locations (see Bildstein, 2006).

Finally, we call attention to the fact that most raptorial species in Europe have vulnerable populations (Burfield, 2008) and some, as with the Spanish imperial eagle and the bearded vulture, were on the brink of extinction in the late 20th century (Bustamante, 1998; Martínez-Cruz et al., 2004). The system we have described here seems to be unique to Europe and has resulted in a hotspot for raptors, many of which are migratory and face conservation problems both inside and outside Europe. The current scenario of climate change adds uncertainty to population persistence (Rodríguez-Rodríguez et al., 2020), while some trans-Saharan migratory species are already changing their behaviour and becoming non-migratory in southern Europe. This includes at least the Egyptian vulture (Di Vittorio et al. 2016), lesser kestrel (Negro et al. 1991), booted eagle (Mellone et al., 2013) and short-toed eagle (Martínez & Sánchez-Zapata 1999). In addition, African species are entering Europe from the south, as it possibly happened with the black-shouldered kite in historical times (Balbontín et al., 2008) and it is just starting with the Ruppell's vulture (Ramírez et al., 2011). If interspecific competition was central in explaining diversity, distribution and migratory behaviour in the European raptor community, as our results suggest, the rapid contemporary changes in climate and bird's ecology will favour some species at the expense of others, generating new conservation challenges.

References

Albach, D.C, Schönswetter, P., Tribsch, A. (2006). Comparative phylogeography of the *Veronica alpina* complex in Europe and North America. *Molecular Ecology, 15*, 3269–3286

Balbontín, J., Negro, J. J., Sarasola, J. H., Ferrero, J. J., & Rivera, D. (2008). Land-use changes may explain the recent range expansion of the Black-shouldered Kite *Elanus caeruleus* in southern Europe. *Ibis*, *150*, 707-716.

Bell, C. P. (2000). Process in the evolution of bird migration and pattern in avian ecogeography. *Journal of Avian Biology, 31*, 258–265. DOI: 10.1034/j.160 0048X.2000.310218.x.

Bergmann, C. (1847). Ueber die Verhaltnisse der Warmeokonomie der Thiere zu ihrer Grosse. *Gottinger Studien, 1*, 595–708.

Bermingham, E., Rohwer, S., Freeman, S., & Wood, C. (1992). Vicariance biogeography in the Pleistocene and speciation in North American wood warblers: a test of Mengel's model. *Proceedings of the National Academy of Sciences*, *89*, 6624-6628.

Bildstein, K. L. (2006). *Migrating raptors of the World: their ecology & conservation*. Cornell University Press.

Bildstein, K. L. (2018). Raptor Migration. In *Birds of Prey*: *Biology and Conservation in the XXI Century*. Edited by J.H. Sarasola, J.M. Grande & J.J. Negro (pp. 123-138). Springer, Cham.

Bildstein, K. L., Smith J.P. & Yosef, R. (2007). Migration counts and monitoring. In Raptor Research and Management Techniques. Eds. D. Bird and K.L. Bildstein. Hancock House.

Bruderer, B., Salewski, V. (2008). Evolution of bird migration in a biogeographical
context. *Journal of Biogeography, 35*, 1951–1959. DOI: 10.1111/j.1365-2699.2008.01992.x

Burfield, I. J. (2008). The conservation status and trends of raptors and owls in Europe.
AMBIO: A Journal of the Human Environment,*37*, 401-407.

Burgin, C. J., Wilson, D. E., Mittermeier, R. A., Rylands, A. B., Lacher, T. E. & Sechrest,
W. (Eds). (2020). 'Illustrated Checklist of the Mammals of the World.' Lynx Edicions, Barcelona.

Bustamante, J. (1998): Use of simulation models to plan species reintroductions: the
case of the bearded vulture in southern Spain. In *Animal Conservation forum* (Vol. 1, No. 4, pp.
229-238). Cambridge University Press.

Clark, W. S., & Schmitt, N. J. (2017). *Raptors of Mexico and Central America*. Princeton
University Press.

Covas, R., & Blondel, J. (1998). Biogeography and history of the Mediterranean bird
fauna. *Ibis*, *140*, 395-407.

Cramp, S., & Simmons, K. E. L. (1979). The birds of the Western Palearctic. Vol. II:
Hawks to Bustards. 695 pp.

Dayan, T. & Simberloff, D. (1998). Size patterns among competitors: ecological
character displacement and character release in mammals, with special reference to island
populations. *Mammal Review*, *28*, 99–124.
408
409　　　　　Del Hoyo, J. (Ed.). (2020). All the Birds of the World. Lynx Edicions, Barcelona.
410
Del Hoyo, J., Elliott, A. & Sargatal, J. (Eds). (1994). Handbook of the Birds of the World.
Vol. 2. New World Vultures to Guineafowl. Lynx Edicions, Barcelona.

Di Vittorio, M., Barbera, A., Di Trapani, E., Faraone, F. P., Ciaccio, A., Sciagura, N.,
D'Amico, D., Giacalone, G., Zafarana, M., Grenci, S. & Sarto, A. (2016). Wintering of Egyptian
vultures (*Neophron percnopterus*) in Sicily: new data. *Arxius de Miscellània Zoològica, 14*, 114–
116.

Dunning Jr, J. B. (2007). *CRC handbook of avian body masses*. CRC press.

Ericson, P.G.P. (2012) Evolution of terrestrial birds in three continents: biogeography
and parallel radiations. *Journal of Biogeography*,*39*, 813–824.
420
Ferguson-Lees, J., & Christie, D. A. (2001). *Raptors of the World*. Houghton Mifflin
Harcourt.

Ferrer, M. & Negro, J.J. 2004. The near extinction of two large European predators: super
specialists pay a price. *Conservation Biology, 18*, 344–349. htps://doi.org/10.1111/j.1523-
1739.2004.00096.x

Forsman, D. (2016). *Flight identification of raptors of Europe, North Africa and the
Middle East*. Bloomsbury Publishing.

Fuchs, J., Johnson, J. A., & Mindell, D. P. (2015). Rapid diversification of falcons (Aves:
Falconidae) due to expansion of open habitats in the Late Miocene. *Molecular Phylogenetics and
Evolution*, *82*, 166-182.

Harrington, B. et al. (2020). Inkscape. http://www.inkscape.org/.

Herrera, C. M. (1978). On the breeding distribution pattern of European migrant birds:
MacArthur's theme reexamined. *The Auk*, *95*, 496-509.

IUCN (2021). The IUCN Red List of Threatened Species. Version 2021-2.
https://www.iucnredlist.org.

Jarvis, E.D. et al. (2014) Whole-genome analyses resolve early branches in the tree of
life of modern birds. *Science 346* (6215), 1320-1331. DOI: 10.1126/science.1253451.
438

Jetz, W., Thomas, G. H., Joy, J. B., Hartmann, K., Redding, D., Mooers, A. O. (2014).
Distribution and conservation of global evolutionary distinctness in birds. *Current Biology, 24*, 1-
12.

Jetz, W., Thomas, G.H, Joy, J. B., Hartmann, K., Mooers A. O. (2012). The global diversity
of birds in space and time. *Nature*, *491*, 444-448.

Keller V, Herrando S, Voříšek P, Franch M, Kipson M, Milanesi P, Martí D, Anton M,
Klvaňová A, Kalyakin MV, Bauer H-G & Foppen RPB (red). (2020). European Breeding Bird Atlas
2: Distribution, Abundance and Change. European Bird Census Council & Lynx Edicions,
Barcelona. 967 pages. ISBN 978-84-16728-38-1.

Knapp M., Thomas J.E., Haile J., Prost S., Ho S.Y.W., Dussex N. *et al.* (2019).
Mitogenomic evidence of close relationships between New Zealand's extinct giant raptors and
small-sized Australian sister-taxa. *Molecular Phylogenetics and Evolution, 134*, 122-128.

Lovette, I. J. (2005). Glacial cycles and the tempo of avian speciation. *Trends in Ecology
& Evolution*, *20*, 57-59.

Martínez, J.E. & Sánchez-Zapata, J.A. (1999). Wintering of booted eagles (*Hieraaetus
pennatus*) and short-toed eagles (*Circaetus gallicus*) in Spain. *Ardeola, 46*, 93-96.

Martínez-Cruz, B., Godoy, J. A., & Negro, J.J. (2004). Population genetics after
fragmentation: the case of the endangered Spanish imperial eagle (*Aquila adalberti*). *Molecular
Ecology*, *13*, 2243-2255.

Massemin, S., Korpimäki, E., & Wiehn, J. (2000). Reversed sexual size dimorphism in
raptors: evaluation of the hypotheses in kestrels breeding in a temporally changing
environment. *Oecologia*, *124*, 26-32.

Mellone, U., De La Puente, J., López-López, P., Limiñana, R., Bermejo, A., & Urios, V.
(2013). Migration routes and wintering areas of Booted Eagles Aquila pennata breeding in Spain.
Bird Study, *60*, 409-413.

Mindell D.P., Fuchs, J., & Johnson, J.A. (2018). Phylogeny, Taxonomy, and Geographic
Diversity of Diurnal Raptors: Falconiformes, Accipitriformes, and Cathartiformes. In: Birds of
Prey: Biology and Conservation in the XXI Century. Eds J.H. Sarasola, J.M. Grande & J.J. Negro.
Springer.

Mönkkönen. M. (1994) Diversity patterns in Palaearctic and Nearctic forest bird
assemblages. *Journal of Biogeography, 21*, 183-95.

Mönkkönen, M. & Viro, P. (1997) Taxonomic diversity of the terrestrial bird and mammal
fauna in temperate and boreal biomes of the northern hemisphere. *Journal of Biogeography*,
24, 603– 612

Nadachowska-Brzyska, K., Li, C., Smeds, L., Zhang, G., & Ellegren, H. (2015). Temporal
dynamics of avian populations during Pleistocene revealed by whole-genome sequences.
Current Biology,*25*(, 1375-1380.

Nagy, J., & Tökölyi, J. (2014). Phylogeny, historical biogeography and the evolution of
migration in accipitrid birds of prey (Aves: Accipitriformes). *Ornis Hungarica*, *22*, 15-35.

Nagy, J., Végvári, Z., & Varga, Z. (2017). Life history traits, bioclimate, and migratory
systems of accipitrid birds of prey (Aves: Accipitriformes). *Biological Journal of the Linnean*
Society, *121*, 63-71.

Negro, J.J. (1991). Problemas de identificación. El Cernícalo Vulgar y el Primilla. *La*
Garcilla, *82*, 5-7.

Negro, J.J. (2018). Raptors and people: An ancient relationship persisting today. In: Birds
of Prey: Biology and Conservation in the XXI Century. Eds J.H. Sarasola, J.M. Grande & J.J. Negro.
Springer.

Negro, J.J., De la Riva, M., & Bustamante, J. (1991). Patterns of winter distribution and
abundance of sedentary lesser Kestrel (*Falco naumanni*) in Spain. *Journal of Raptor Research*,
25, 30-35.

Negro, J.J., Prenda, J., Ferrero, J.J., Rodríguez, A. & Reig-Ferrer, A. (2020). A timeline for
the urbanization of wild birds: the case of the lesser kestrel. *Quaternary Science Reviews*, *249*,
106638.

Newton, I. (1979). Population ecology of raptors. Buteo Books, Vermillion, South Dakota.

Newton, I., & Dale, L. (1996). Relationship between migration and latitude among west
European birds. *Journal of Animal Ecology*, *65*, 137-146.

Newton, I., & Dale, L. (1997). Effects of seasonal migration on the latitudinal
distribution ofWest Palaearctic bird species. *Journal of Biogeography, 24*, 781–789.

Perrins, C. M., & Snow, D. W. (1998). The Birds of the Western Palearctic. Concise
Edition. *New York*.

QGIS Development Team (2021). QGIS Geographic Information System. Open Source
Geospatial Foundation. URL http://qgis.org.

R Core Team (2013). R: A language and environment for statistical computing. R
Foundation for Statistical Computing, Vienna, Austria. URL http://www.R-project.org/.inburgh.

Rambaut, A., & Drummond, A. J. (2018). FigTree v1. 4.4. Institute of Evolutionary
Biology. University of Edinburgh. UK.

Ramírez, J., Muñoz, A. R., Onrubia, A., De la Cruz, A., Cuenca, D., González, J. M., &
Arroyo, G. M. (2011). Spring movements of Rüppell's Vulture Gyps rueppellii across the Strait of
Gibraltar. *Ostrich*, *82*, 71-73.

509     Rodríguez-Rodríguez, E. J., Beltrán, J. F., El Mouden, E. H., Slimani, T., Márquez, R., &
510 Donaire-Barroso, D. (2020). Climate change challenges IUCN conservation priorities: A test with
511 western Mediterranean amphibians. *SN Applied Sciences*, *2*, 1-11.

512     Salewski, V. & Watt, C. (2017). Bergmann's rule: a biophysiological rule examined in
513 birds. *Oikos, 126*, 161–172.

514     Snow, D. W. (1978). Relationships between the European and African avifaunas. *Bird*
515 *Study*, *25*, 134-148.

516     Svensson, L., Mullarney, K., Zetterström, D. &Grant P. J. (1999). Collins Bird Guide: the
517 most complete guide to the birds of Britain and Europe, 2nd ed. Harper Collins.

518     Voelker, G. (2010) Repeated vicariance of Eurasian songbird lineages since the Late
519 Miocene. *Journal of Biogeography*,*37,* 1251–1261.
520
521     Wang P, Yao H, Gilbert KJ, Lu Q, Hao Y. et al. (2018). Glaciation-based isolation
522 contributed to speciation in a palearctic alpine biodiversity hotspot: evidence from endemic
523 species. *Molecular Phylogenetics and Evolution, 129*,315–324.
524
525     Weir, J. T., & Schluter, D. (2004). Ice sheets promote speciation in boreal birds.
526 *Proceedings of the Royal Society of London. Series B: Biological Sciences*, *271*(1551), 1881-
527 1887.

528     Zachos, J., Pagani, M. Sloan, L., Thomas, E., Billups, K. (2001). Trends, Rhythms, and
529 Aberrations in Global Climate 65 Ma to Present. *Science, 292*, 686-693.
530
531     Zink, R.M., Klicka, J. & Barber, B.R. (2004). The tempo of avian diversification during
532 the Quaternary. *Philosophical Transactions of the Royal Society of London* B., *359*, 215-220.
533 doi: 10.1098/rstb.2003.1392, 215-220359 2004.
534

Biosketch

Juan José Negro has been investigating raptor ecology and behavior for more than three
decades. He was Director of Doñana Biological Station-CSIC in Spain, and currently holds a
position as Research Professor in the Spanish National Research Council.

Authors' contributions: JJN conceived the hypothesis, wrote the first draft of the manuscript
and revised the last versions. EJJRR analyzed the trophic niche data and generated trophic
niche and phylogeny of European raptors, contributing also to the method and results
sections. AR performed the body mass analyses in relation with migration, prepared
distribution maps and contributed with manuscript writing and editing. KB provided insights,
and numerous references. All authors helped to draft the paper and edited the text.

Group	Species	Body mass male(g)	Body mass female (g)	Transaharian migration	IUCN
1	*Pandion haliaetus*	1120-1740	1210-2050	yes	LC
1	*Haliaeetus albicilla*	4900-6000	6800-9000	no	LC
2	*Circaetus gallicus*	1200-2000	1300-2300	yes	LC
3	*Pernis apivorus*	440-943	450-1050	yes	LC
4	*Circus aeruginosus*	405-730	540-960	yes	LC
5	*Circus macrourus*	235-416	402-550	yes	NT
5	*Circus pygargus*	227-305	254-445	yes	LC
5	*Circus cyaneus*	300-400	370-708	no	LC
6	*Accipiter gentilis*	517-1110	820-2200	no	LC
6	*Accipiter brevipes*	135-223	232-275	yes	LC
6	*Accipiter nisus*	105-196	185-350	yes	LC
7	*Milvus migrans*	630-928	750-1080	yes	LC
7	*Milvus milvus*	757-1220	960-1600	no	LC
8	*Buteo buteo*	427-1180	486-1360	yes	LC
8	*Buteo lagopus*	600-1130	783-1660	no	LC
8	*Buteo rufinus*	590-1281	945-1760	yes	LC
9	*Hieraaetus pennatus*	510-770	840-1250	yes	LC
9	*Aquila fasciata*	1560-1560	2000-2500	no	LC
10	*Aquila chrysaetos*	2800-4600	3800-6700	no	LC
10	*Aquila adalberti*	2500-3500	2500-3500	no	VU
10	*Aquila heliaca*	2450-2720	3160-4530	yes	VU
11	*Clanga clanga*	1700-1900	1800-2500	no/ yes	VU
11	*Clanga pomarina*	1000-1400	1300-2200	yes	LC
12	*Aegypius monachus*	7000-11500	7500-12500	no	NT
12	*Gyps fulvus*	6200-10500	6500-1100	yes	LC
13	*Gypaetus barbatus*	4500-7000	5600-6700	no	NT
13	*Neophron percnopterus*	1600-2400	1600-2400	yes	EN
14	*Falco eleonorae*	350-390	340-460	yes	LC
14	*Falco subbuteo*	131-232	141-40	yes	LC
15	*Falco peregrinus*	550-750	740-1300	no	LC
15	*Falco pelegrinoides*	549-610	657-924	no	LC
15	*Falco biarmicus*	500-600	700-900	no	LC
15	*Falco rusticolus*	800-1320	1130-2100	no	LC
15	*Falco cherrug*	730-950	970-100	yes	EN
16	*Falco naumanni*	90-172	128-216	yes	LC
16	*Falco vespertinus*	115-190	130-197	yes	NT
16	*Falco tinnunculus*	136-252	154-314	no	LC
16	*Elanus caeruleus*	197-277	219-343	no	LC

Table 1. Assemblage of diurnal raptors breeding in continental Europe, ordered according to
trophic niche (group, first column). Some of the groups (2 to 4) are monospecific, as they are
constituted for highly niche-specific species. Range of body masses for males and females were
taken from Ferguson-Lees & Christies (2001), as well as their migratory status (i.e., trans-
Saharan migrant or not). IUCN conservation status is also given (IUCN 2021).

Figure 1: Trophic niche cluster dendrogram based on Euclidean distances for the
European species of raptors. Picture credits: *Gypaetus barbatus*, *Accipitier nisus* and *Falco*
eleonorae by Manuel Cayuela López. *Circus pygargus* by Miguel Ángel Rojas. Remaining
photographs by the authors.

Figure 2: Phylogeny of included European raptors. Colour areas indicate groups of species with direct niche competition. Colour stars are used to join species not phylogenetically directly related but engage in niche competition (colour areas). Red asterisk indicated long distance migration. Number are species not include in the niche groups (1. Circus aeruginosus; 2. Circaetus gallicus; 3. Pernis apivorus). Picture credits: Clanga clanga modified from J.M. Garg (CC BY-SA 3.0). Buteo lagopus cropped from Walter Siegmund (CC BY 2.5). Haliaeetus albicilla cropped from Christoph Müller (CC BY 4.0). Falco vespertinus cropped from Martin Mecnarowski (CC BY 3.0). Falco biarmicus cropped from Derek Keats (CC BY 2.0). Falco cherrug cropped from DickDaniels (http://carolinabirds.org/) (CC BY-SA 3.0). Falco pelegrinoides cropped from Peter Wächtershäuser/ naturlichter.de (CC BY-SA 3.0). Falco eleonorae and Accipiter gentilis with permission of Manuel Cayuela López. Remaining pictures taken by the authors.

License links: https://creativecommons.org/licenses/by-sa/3.0/legalcode;
https://creativecommons.org/licenses/by/2.5/legalcode;
https://creativecommons.org/licenses/by/4.0/legalcode;
https://creativecommons.org/licenses/by/2.0/legalcode

Figure 3: Pairwise comparisons of body mass (a) and body size (b) of raptor species derived of the foraging groups in Figure 1. Y-axis represents the migratory species in the pair, while the X-axis the non-migratory species (resident). Dashed lines indicate the diagonal (y = x). Letters refers to species pairs: a) *Gypaetus barbatus - Neophron percnopterus*, b) *Aegypius monachus - Gyps fulvus*, c) *Clanga clanga - Clanga pomarina*, d) *Aquila chrysaetos - Aquila heliacal*, e) *Aquila fasciata - Aquila pennatus*, f) *Circus cyaneus - Circus macrourus*, g) *Circus cyaneus - Circus pygargus*, h) *Accipiter gentilis - Accipiter brevipes*, i) *Accipiter gentilis - Accipiter nisus*, j) *Buteo rufinus - Buteo buteo*, k) *Buteo rufinus - Buteo lagopus*, l) *Milvus milvus - Milvus migrans*, m) *Falco tinnunculus - Falco naumanni*, n) *Falco tinnunculus - Falco vespertinus*, o) *Falco peregrinus - Falco eleonorae*, p) *Falco peregrinus - Falco subbuteo*, q) *Falco rusticolus - Falco columbarius*, r) *Falco rusticolus - Falco cherrug*.

598 **Figure 4**: Year-round distribution maps of the raptor species with pairs or trios resulting from
599 the trophic-niche cluster analysis. Ranges of larger in size species are displayed in blue, while
600 ranges of the smaller and typically intercontinental migratory species are displayed in red (and
601 green in case of trios). Distribution ranges were taken from the IUCN Red List Data (IUCN
602 2021).