Generation of raptor diversity in Europe: linking speciation to climate
changes and the ability to migrate

Running title:

Juan J. Negro[1]

Eduardo Rodríguez-Rodríguez[3]

Airam Rodríguez[1,2]

Keith Bildstein

[1]Estación Biológica de Doñana-CSIC, Avda. Americo Vespucio 26. 41092. Sevilla. Spain.

[2]Grupo de Ornitología e Historia Natural de las islas Canarias (GOHNIC), C/La Malecita s/n,
Buenavista del Norte, Canary Islands, Spain.

[3]Integrative Sciences Department. University of Huelva (Huelva). Spain.

41012. Sevilla. Spain.

Corresponding author: Juan J. Negro. E-mail: negro@ebd.csic.es

Abstract

The majority (58%) of raptors abandons their breeding ground and migrates to distant
areas, sometimes in other continents. In the case of European breeding raptors, we can
observe that in groups of species that share similar trophic niches (sometimes these species
are taxonomically closely related), only some of them (one in the case of groups formed by
only a pair) do a large distance migration. In this work we analyze the trophic niche of
European raptors, with the objective of obtaining trophic niche groups. Once we have these
niche groups, we related them with the migration behavior and taxonomy of species. Our
results show that the smallest species inside these niche groups are those that migrates at long
distance, suggesting that competence by resources in the less favorable seasons promote the
migratory behavior in the "weakest" species.

Introduction

Raptor migration is a conspicuous phenomenon (Bildstein, 2018). Numerous raptor
species, 183 out of 313 (58%), abandon their breeding grounds and, often gathering in large
flocks, move away at predictable times and places to reach distant areas located sometimes in
different continents (Bildstein et al., 2007, Fergusson-Lees and Christie, 2001). Raptor
migration has been extensively studied and monitored to assess long-term population trends
and to determine raptor flight behavior, migration corridors, as well as breeding and non-
breeding grounds (Bildstein, 2006). However, much remains to be investigated concerning the
causes and function of raptor migration, or why a majority of species are migratory but still
others remain sedentary. Recent molecular advances have shown that there are in fact two
distantly related groups of diurnal raptors, convergent in their morphology and general
ecology (Jarvis et al., 2015). On one side, we have the true falcons and allies, in the Order
Falconiformes. On the other side, eagles, hawks, kites, harriers, and vultures constitute the

Order Accipitriformes. According to Ericsson (2011), the early radiation of Falconiformes took
place in South America, whereas Accipitriformes emerged as a group in Africa. Today, both
orders have gained a cosmopolitan distribution and species of the two orders tend to co-occur
in most environments. There are many migratory species or populations within both
Falconiformes and Accipitriformes (Bildstein, 2006) in both the Nearctic, the Palearctic and
inside Africa, implying that long-distance migratory behavior has evolved multiple times and
that it is a rather plastic trait (Nagy et al., 2017).

Europe is the land mass with the highest proportion of migratory species of birds of
prey in the World (Bildstein, 2006), and there is also a large absolute number of species
considering its relatively small size compared to the remaining of the Palearctic and also the
Nearctic (see results). Why this is so has never been explained nor explored so far. Also
overlooked, even though it should be obvious glancing at any bird guide (e.g., Svensson et al.,
1999) or raptor biology book (Ferguson-Lees and Christie, 2001), is the fact that there are
numerous groups (duets and trios, in fact) of morphologically and ecologically similar raptorial
species in the European portion of the Western Palearctic. The species conforming these
groups may look so similar in appearance that their field identification at a distance may be
challenging (Negro, 1991; Forsman, 2016). We refer to pairs of species such as the Eurasian
(*Falco tinnunculus*) and the Lesser kestrels (*F. naumanni*), the lesser and greater spotted eagles
(*Aquila pomarina* and *A. clanga*), or trios including, for instance, Montagu's harrier (*Circus
pygargus*), hen harrier (*C. cyaneus*) and pallid harrier (*C. macrourus*).

Here we aim to link raptor migration, interspecific competition and finally diversity and
speciation in the European diurnal raptor assemblage, including both Falconiformes and
Accipitriformes. Controlling for phylogeny and considering some regularities in size differences
and migratory behaviour, we will test the hypothesis that the raptor community in Europe has
recently experienced a diversification of breeding species, mainly during the Quaternary. To
get further support for our hypothesis, we have gathered information on the timing of species
splitting derived from time-calibrated molecular phylogenies (Fuchs et al., 2015; Mindell et al.,
2018).

Methods

Study area and selected species

This study focuses on the 38 diurnal raptor species breeding regularly and historically
in continental Europe (therefore excluding the Macaronesian archipelagos of Azores, Madeira
and the Canary Islands). We use the term Europe as defined in the seven-continents model
(i.e., Europe separated from Asia as a conglomerate of peninsulas delineated by the
Mediterranean sea to the South, the Atlantic ocean to the West, the Northern sea in high
latitudes, and the Ural mountains, the Caucasus and the Black and Caspian seas to the East)
because it suits well the differences in migrating behaviour of birds including raptors in
northern and temperate areas (see, e.g., Herrera, 1978; Newton and Dale, 1996, Newton and
Dale, 1998, Bildstein, 2006). Using continental Europe as a biogeographical area permits to
cleanly separate sedentary, migrants within continental Europe, and sub-Saharan bird species
migrating through the Mediterranean flyways (i.e., Straits of Gibraltar, Messina and
Bosphorus). The Western Palearctic, a zoogeographical region widely used in bird studies
(Cramp & Simmons, 1979; Perrins & Snow, 1998), and that fully encompasses Europe,
introduces, however, African and Asian raptorial species that would add noise to our analyses.
These species, either at the southern limit of the Western Palearctic in the African continent or

at the eastern border with Asia, are all rare breeders or vagrants in the area that have never
established breeding populations in continental Europe. Raptors of the Western Palearctic
which are clearly African taxa include at least (Svensson et al., 1999), the lappet-faced vulture
(*Torgos tracheliotos*), the Dark chanting goshawk (*Melierax metabates*), the Barbary falcon
(*Falco pelegrinoides*), the Sooty falcon (*Falco concolor*), the tawny eagle (*Aquila rapax*), and
Verreaux's eagle (*A. verreauxi*).

Phylogeny and trophic niche

As a first step to determine the relationships among the different species under study,
a consensus phylogeny was constructed using FigTree v 1.4.4 (Rambaut & Drummond, 2018)
and 100 permutations obtained from BirdTree (Jetz et al., 2012; Jetz et al., 2014). Addressing
migratory behaviour in birds requires not only the consideration of biogeographic aspects, but
also a knowledge of diet specializations and prey availability (Nagy and Tokolyi, 2014). A
trophic niche cluster dendrogram, based on Euclidean distances, was constructed using a
dataset with information of trophic niche variables of each species, including hunting or
scavenging behaviour, prey type and size (i.e., small birds, large birds, small mammals, large
mammals, arthropods, herps or fishes), scavenging strategy (small pieces or big pieces) and
foraging method (perch hunting, foraging in open areas, forest ambusher). These data can be
consulted in Supplementary Material appendix 1. The software utilised to construct the
dendrogram was in R statistics (R Core Team, 2013). Presentation of the figures was improved
with Inkscape 1.0.1 (Harrington et al., 2020).

Relating body mass and trans-Saharan migration

We compiled the following information of the raptor species: the minimum and
maximum value of both total length (2 values) and wingspan (2) from del Hoyo et al. (1994),
the minimum and maximum values (2 values each) of length, tail, and tarsus, as well as male
body mass and female body mass (4 mass values) from Fergusson-Lees & Christie (2001). To
estimate body mass, we averaged the minimum and maximum values for males and females.
To estimate bird size, we ran a principal component analysis (PCA) on the above mentioned
fourteen centered and scaled morphometric variables. The first principal component was used
as a body size index. The first principal component retained 90.1% of variation (see PCA details
in Table S1). The fourteen morphometric variables showed the same sign for their factor
loadings and highly significant correlations to the first principal component (Figure S1). Factor
loadings of the fourteen morphometric variables and importance of the components are given
in Table S1. Pairwise correlations of the first principal component (body size index) and seven
morphometric variables (i.e. those representing the lower ranges of each morphometric) are
shown in Fig. S1.

Based on results from trophic niche clustering, groupings of species sharing a similar
niche were established. These groups were identified on the previously built phylogeny also
indicating body mass for every species, and we also indicated if the species performed a trans-
Saharan migration (Bildstein 2006). With these data, we established two categories: smaller
raptors inside their groups and larger raptors inside their groups, and estimated the proportion
of trans-continental migratory species among large and small species. Significance of the
differences in frequency of migration for both groups of large and small species were tested
using a contingency table. In addition, we plotted the mass of paired migratory and resident
species in the foraging and phylogenetic groups that we had previously identified (Figure 3a).
By adding a diagonal line (y = x), the resulting plot would graphically show in which cases the

migratory species was either larger (above the diagonal) or smaller (below the diagonal) than
its resident counterpart. An additionally plotted the previously described body size index
instead of just body mass, because the anatomical design of migratory species is known to be
slender, i.e., much longer wings, or different wing loads, compared to non-migratory species
(Figure 3b).

To build the maps displayed in Figure 4, distributions ranges of raptor species were
downloaded from the IUCN Red List Data (IUCN, 2021) and imported in *Qgis* (v3.20) (Qgis
Development Team, 2021). We deleted passage areas for migrating species and mapped pairs
and trios of species depending on results of trophic niche clustering of the previous analyses.

Results

Foraging groups

Based on Euclidean distances of behavioural and dietary variables, we obtained 16
trophic/ecological niche groups, including four monospecific niches (i.e., the crepuscular and
small-mammal specialist *Elanus caeruleus*; the reed-bank specialist *Circus aeruginosus*, the
snake eating *Circaetus gallicus* and the Himenopteran-eater *Pernis apivorus*), ten groups of
two species, two groups of three species –the harriers and the accipiter hawks-, and one group
of five species conformed by the bird-eating falcons (Figure 1).

Phylogenetic relationships of European raptors

148   The resulting phylogeny for the European raptor species matches well with the
149 obtained trophic groups in a majority of cases, except for the *Pandion haliaetus/Haliaeetus*
150 *albicilla* and *Aquila fasciatus/Hieraetus pennatus* species pairs, that are polyphyletic.
151 Phylogeny, niche groups and average body mass are provided in Figure 2. We also indicate
152 whether the species performs a trans-Saharan migration or not.

Implications of size and migration strategy

A majority of the smaller species inside trophic groups reach Africa during migration
(81% of cases, n = 16, and only 19% do not. On the contrary, only 20% (4 out of the 20) of the
larger species inside trophic groups perform a trans-Saharan migration (i.e, *Clanga clanga* or
Falco eleonorae, Circus macrourus and *Buteo rufinus*). However, these "large" species perform
a much shorter migration than their smaller counterparts in their respective groups (see Fig.
4). The difference in migratory behaviour among larger and smaller species is highly significant
(x^2=11,6298, P=0.00649, see figure 3).

When plotting the body masses for paired species derived from the foraging groups
than were stablished above (Fig. 3a), the values fall below the diagonal of the graph (x axis,
mass of less migratory species, y axis, mass of more migratory species in the pair). This
demonstrates that the fully or more migratory species are systematically smaller than the less
migratory or resident counterparts. Essentially the same results are obtained when instead of
plotting body masses, body size indices were plotted (Fig. 3b). Larger species are the less
migratory ones in the species pairs. The distribution ranges for a selection of the species pairs
that we considered in the above plots have been mapped in Fig. 4. The less-migratory species
remain within Europe (although it has to be noted some may have resident populations in
North-Africa), whereas the fully migratory species have non-overlapping breeding (in Europe)
and non-breeding ranges (in Africa).

Timing of the splits from the common ancestor

Using the time-calibrated phylogenies of falcons (Falconiformes) on the one hand (Fuchs et al. 2015) and of all other birds of prey (eagles, hawks, vultures and kites in Accipitriformes) on the other hand (Mindell et al., 2018), we propose that six of our species groupings (both in the phylogeny and in the trophic cladogram) originated in the Quaternary (i.e., the last 2.5 million years) as sister species. These groupings are the following: 1. The red and the black kite. 2. The two spotted eagles, lesser and greater. 3. The two imperial eagles (*A. heliaca* and *A. adalberti*), that we have matched ecologically to the most distantly related *A. chrysaetos*. 4. The three European buzzards: common, rough legged and long-legged. 5. The two late-breeding falcons, dependent on the return migration of passerines for reproduction: hobby and Eleonora's falcon. 6. The large European falcons: Lanner, Saker and Gyrfalcon. Apart from this, we may add that the two colonial and tree-nesting falcons of temperate Eurasia, i.e., the red-footed falcon in the Western Palearctic and Amur falcon in the Eastern Palearctic, are also sister species that descend from a common ancestor in the Quaternary.

The remaining groupings may have formed earlier in time, again according to Fuchs et al (2015) and Mindell et al. (2018): 1. the Eurasian kestrel and the lesser kestrel emerged as species in the Pliocene kestrel radiation about 4-5 million years (see also Negro et al., 2020). 2. The harriers (hen, Montagu's and pallid) radiated about 5 million years ago. 3. The hawks of the genus *Accipiter* radiated much earlier, and the common ancestor of the northern goshawk and both the European sparrowhawk and levant sparrowhawk possibly evolved 10-15 million years ago in the Miocene. 4. The two white eagles, Bonelli's and booted, emerged from a common ancestor living in the Pliocene 4 million years ago. 5. The two large European vultures, griffon and cinereous, shared a common ancestor about 9 million years ago. 6. Coalescence time for Bearded and Egyptian vultures, which are sister groups to each other, is about 12 million years.

Discussion

The diversity of species of any given taxa in a land mass typically depends on its own evolutionary history, the interaction with other species in the area, and the past and present physical patterns of the environment (e.g., Hanssen & Urban, 1992; Mönkkönen, 1994). This work was prompted by the fact that the number of breeding species of diurnal raptors in continental Europe (38 species in a land-mass of 10,180,000 km^2) is higher than the number of breeding species of raptors in Canada and continental US combined (30 species in about 19,000,000 km^2), and also by the disproportionate number of transcontinental migratory species in Europe (Bildstein 2006, see below). The latitudinal range is much wider in the American side and includes large areas below 35° North latitude in the southern states of the US that are not represented in continental Europe. However, this does not bring over a significant number of tropical species that do inhabit Mexico and Central America (the total number of raptorial species in the latter area rises to 69 species, Clark and Schmitt, 2017).

Seven raptorial species are shared among Europe and North America: five Holarctic raptor species, the Gyrfalcon, the merlin, the golden eagle, the rough-legged hawk, and the northern goshawk, plus two cosmopolitan species, the peregrine falcon and the osprey, which hold breeding populations in all continents except Antarctica (Fergusson-Lees and Christie,

2001). Leaving aside these few shared species, mainly from northern or mountainous regions,
there is a marked difference among Europe and Canada-US combined in the number and
proportion of regular transcontinental migratory species, which is much higher in Europe (22
species crossing the Mediterranean Sea to Africa in the non-breeding season) than in the
above mentioned area of North America, where only five species make the bulk of the
migrants moving down to South America (i.e., *Ictinia mississippiensis*, *Buteo swainsoni*, *Buteo*
platypterus, *Cathartes aura* and *Pandion haliaetus*) (Bildstein, 2006).

Our results show that there is a duplicity, or even multiplicity of species in the same
trophic /ecological niches in Europe. In many cases the species involved are so similar that
identification at a distance may be difficult (Forsman, 2016). This similarity may be due to
shared ancestry in congeneric species, such as in the duet conformed by *Aquila pomarina* and
A. clanga, or in the case of *M. milvus* and *M. migrans* (Mindell et al., 2018), but it may be due
to convergence in more distantly related species such as *A. fasciatus* and *Hieraaetus pennatus*.
Duplicated species are not apparent in North America, where the two more diverse genera
(*Buteo* and *Falco*, with 6 species each in Canada and the USA), have rather distinct species in
terms of size and ecology. That Europe has been a recent hotspot for raptor speciation is also
supported by the pattern of endemism (Weir and Schluter, 2004) in the group: six species are
only or mainly distributed in Europe (i.e., red kite, honey buzzard, lesser spotted eagle, Spanish
imperial eagle, Montagu's harrier and Eleonora's falcon). Five other species breed in Europe
and more into the East, but may have a Western Palearctic origin: lesser kestrel (with the older
fossil found in the Iberian peninsula, Negro et al., 2020), red-footed falcon (sister species of
the Amur falcon breeding in eastern Asia, as commented above), Common Buzzard (the
nominal subspecies is sedentary across Europe, and only the Eastern subspecies *B. b. vulpinus*
migrates to Africa), and finally the Levant sparrowhawk, and the Western marsh harrier.

Our work is not focused on the evolution of raptor migration, but our results may shed
light on the biogeographical origin of at least some of the migratory species. A majority of the
duplicated species would adhere to the "northern origin" hypothesis (Bell, 2000; Bruderer and
Selewski, 2008, Nagy et al., 2014), according to which harsh winter conditions would impel
northern species to move to Africa in the non-breeding season. Some species in the system,
such as the short-toed eagle, the Egyptian vulture and the bearded vulture, and the two white
eagles, *A. fasciata* and *H. pennatus* may have an afro-tropical origin, as they belong to
phylogenetic groups which are highly diverse in Africa (Mindell et al., 2018), even if the
migratory species among them may have acquired the migratory behaviour after colonizing
Europe and competing with other species there.

Taken together, our results support the notion that the raptor community in Europe
experienced a recent enrichment, and may be considered a hotspot accounting for its area and
geographic location, and particularly by comparison with the smaller raptor community in
much larger North America. This does not mirror the situation for the whole of the avifaunas,
as there are about 556 bird species in Europe (Keller et al. 2021), compared to 740 in North
America to the north of Mexico (Cornell lab of Ornithology,
https://www.notesfromtheroad.com/roam/how-many-birds-north-america.html). The North
American avifauna contains a plethora of insectivorous passerines with a probable tropical or
subtropical origin (i.e., Parulidae, Turannidae, Vireonidae, Icteridae) which exploit the insect-
rich summer of North America and migrate to the south in the winter (Snow 1978). The
European migratory songbirds are fewer and belong to families (i.e, Sylviidae, Turdidae,
Muscicapidae and Motacillidae) whose evolutionary centres are Palearctic but not African

(Snow 1978). As a result, the assemblages of forest passerine birds of the Nearctic region are
more diverse than those of the Western Palearctic, thus including Europe (Mönkkönen 1994).
Other taxa follow the same pattern: Western North America contains, for instance, 41% more
mammal species than Europe (Mönkkönen and Viro, 1997).

The mechanism that may explain the generation of raptor diversity in Europe is the
existence of climate-driven vicariant events during the glacial cycles of the Pleistocene
(Voelker 2010), and before during the Pliocene or earlier in the Miocene. It has been
suggested, for instance, that the genus *Falco* radiated during the Miocene several million years
ago, coincidental with a period of increased aridity favouring grasslands (Zachos et al., 2001,
Fuchs et al. 2015). Glacial cycles have been invoked as drivers of speciation for numerous bird
taxa in both the Old and the New World: fragmentation of populations, isolation in refugia
during glacial times and subsequent reunion without admixture is a common model for
explaining the origin of closely-related bird species (Bermingham et al., 1992 Weir and
Schluter, 2004, Lovette, 2005). Such a model was in fact used to explain the emergence as a
separate species of the Spanish imperial eagle (Ferrer and Negro, 2004), the species in the
genus *Aquila* with the smallest distribution range, as it is endemic to the Iberian Peninsula.

Lacking enough dated fossils, the recent avian speciation during the Pleistocene glacial
cycles has remained controversial for some groups (Lovette, 2005 ; Zink et al., 2004, Wang et
al., 2018). However, recent advances in molecular techniques, such as whole-genome
sequencing, clearly support that many bird species suffered dramatic expansions and
contractions during the Quaternary coincidental with climate changes (Nadachowska-Bryzska
et al., 2015). These population changes may have prompted speciation.

The generation of sister species of birds of prey may be explained by vicariance events
as explained above. The difference among Europe and North America possibly has to do with
the different effects of the glaciation in the two land masses, and the alignment of the major
mountain barriers that may have acted as geographical barriers, with a north-south
component in the Nearctic and east-west in the Palearctic (Albach et al., 2006). In North
America, numerous pairs of sister bird species tend to have east-west distributions (e.g.,
Lovette, 2005). In Europe, the existence of several southern peninsulas and islands offered
different refugia and thus more possibilities for population fragmentation and isolation (Covas
& Blondel, 1998). After ice retreated during the interglacials, isolated populations might have
expanded and established contact but, in some cases, without genetic admixture. However,
closely related species of predators coexisting in sympatry would compete dearly in the harsh
winter season. Competition would be resolved by the less competitive species moving away
and thus migrating to a different continent. This may also be seen as several concurrent cases
of character displacement., with competing species diverging in size precisely to diminish
potential competition (Dayan & Simberloff, 1998). Also of interest may the fact that the
assemblage of European carnivores (Mammalia) is much smaller, with 22 species than the
North American assemblage, with 45 species (Burgin et al., 2020). Lessened potential
competition with mammalian carnivores may have facilitated the diversification of European
breeding raptors. None-the-less, competition with other predators alone is not the only
possible explanation, as Bergmann's rule (Bergmann 1847), for instance, would also predict
larger bird species having a thermoregulatory advantage compared to smaller ones during the
cooler non-breeding season (Salewski & Watt, 2017).

An extraordinary case of rapid increase of size in the absence of competitors, therefore
linking body size to competition in the raptor group, is provided by two extinct birds of prey in

New Zealand (Knapp et al., 2019): the largest known eagle in the world, Haast's eagle
Hieraaetus moorei, and the largest known harrier in the world, Eyles' harrier *Circus*
teauteensis. Both derived from Australian vagrants, the much smaller *Hieraaetus morphnoides*
and *Circus assimilis*, respectively. These two concurrent cases of island gigantism occurred in
the early Pleistocene when the clearing of closed forests due to climate changes allowed for
successful colonization of Australian raptor immigrants hunting in open woodlands and
grasslands (Knapp et al., 2019). The fact that body size is in turn related to migration
propensity and ability is also supported at the intraspecific level. Raptors tend to present
reversed sexual dimorphism, as females are larger than males (Newton, 1979). Numerous
raptor species show different migration timings and non-breeding distributions, although it is
true that there is variation in which sex migrates first (both out-bound and in-bound) or
overwinters in more northerly locations (see Bildstein, 2006).

Last, we wish to call attention to the fact that most raptorial species in Europe have
vulnerable populations (Burfield, 2008) and some, as with the Spanish Imperial eagle and the
Bearded vulture, were on the brink of extinction in the late 20th century (Bustamante, 1998;
Martínez-Cruz et al., 2004). The system we have described here seems to be unique to Europe
and has resulted in a hotspot for raptors, many of which are migratory and face conservation
problems both inside and outside Europe. The current scenario of climate change adds
uncertainty to population persistence (Rodríguez-Rodríguez et al., 2020), while some trans-
Saharan migratory species are already changing their behaviour and becoming sedentary in
the southern countries of Europe. This includes at least the Egyptian vulture (Di Vittorio et al.
2016), lesser kestrel (Negro et al. 1991), booted eagle (Mellone et al., 2013) and short-toed
eagle (Martínez & Sánchez-Zapata 1999). In addition, African species are entering Europe from
the south, as it possibly happened with the black-shouldered kite in historical times (Balbontín
et al., 2008) and it is just starting with Ruppell's vulture (Ramírez et al., 2011). If interspecific
competition was central in explaining diversity, distribution and migratory behaviour in the
European raptor community, as our results suggest, the rapid contemporary changes in
climate and bird's ecology will favour some species at the expense of others, generating new
conservation challenges.

Acknowledgements

References

Albach, D.C, Schönswetter, P., Tribsch, A. (2006). Comparative phylogeography of the
Veronica alpina complex in Europe and North America. *Molecular Ecology, 15*, 3269–3286

Balbontín, J., Negro, J. J., Sarasola, J. H., Ferrero, J. J., & Rivera, D. (2008). Land-use
changes may explain the recent range expansion of the Black-shouldered Kite *Elanus caeruleus*
in southern Europe. *Ibis, 150*(4), 707-716.

Bell, C. P. (2000). Process in the evolution of bird migration and pattern in avian
ecogeography. *Journal of Avian Biology,* 31, 258–265. DOI: 10.1034/j.160
0048X.2000.310218.x.

Bergmann, C. (1847). Ueber die Verhaltnisse der Warmeokonomie der Thiere zu ihrer
Grosse. *Gottinger Studien,* 1, 595–708.

Bermingham, E., Rohwer, S., Freeman, S., & Wood, C. (1992). Vicariance biogeography
in the Pleistocene and speciation in North American wood warblers: a test of Mengel's model.
Proceedings of the National Academy of Sciences, *89*(14), 6624-6628.

Bildstein, K. L. (2006). *Migrating raptors of the World: their ecology & conservation.*
Cornell University Press.

Bildstein, K. L. (2018). Raptor Migration. In *Birds of Prey*. Biology and Conservation in
the XXI Century. Edited by J.H. Sarasola, J.M. Grande & J.J. Negro (pp. 123-138). Springer,
Cham.

Bildstein, K. L., Smith J.P., Yosef, R. (2007). Migration counts and monitoring. In Raptor
Research and Management Techniques. Eds. D. Bird and K.L. Bildstein. Hancock House.

Bruderer, B., Salewski, V. (2008). Evolution of bird migration in a biogeographical
context. *Journal of Biogeography,* 35, 1951–1959. DOI: 10.1111/j.1365-2699.2008.01992.x

Burfield, I. J. (2008). The conservation status and trends of raptors and owls in
Europe. *AMBIO: A Journal of the Human Environment*, *37*(6), 401-407.

Burgin, C. J., Wilson, D. E., Mittermeier, R. A., Rylands, A. B., Lacher, T. E. and Sechrest,
W. (Eds). (2020). 'Illustrated Checklist of the Mammals of the World.' (Lynx Edicions:
Barcelona.)

Bustamante, J. (1998): Use of simulation models to plan species reintroductions: the
case of the bearded vulture in southern Spain. In *Animal Conservation forum* (Vol. 1, No. 4, pp.
229-238). Cambridge University Press.

Clark, W. S., & Schmitt, N. J. (2017). *Raptors of Mexico and Central America.* Princeton
University Press.

Covas, R., & Blondel, J. (1998). Biogeography and history of the Mediterranean bird
fauna. *Ibis*, *140*(3), 395-407.

Cramp, S., & Simmons, K. E. L. (1979). The birds of the Western Palearctic. Vol. II:
Hawks to Bustards. 695 pp.

Dayan, T. & Simberloff, D. (1998). Size patterns among competitors: ecological
character displacement and character release in mammals, with special reference to island
populations. *Mammal Review*, 28 (3),, 99–124.
381
Del Hoyo, J., Elliott, A., Sargatal, J. (Eds). (1994). Handbook of the Birds of the World.
Vol. 2. New World Vultures to Guineafowl. Lynx Edicions, Barcelona.

Di Vittorio, M., Barbera, A., Di Trapani, E., Faraone, F. P., Ciaccio, A., Sciagura, N.,
D'Amico, D., Giacalone, G., Zafarana, M., Grenci, S. & Sarto, A.. (2016). Wintering of Egyptian
vultures (Neophron percnopterus) in Sicily: new data. *Arxius de Miscellània Zoològica,* 14, 114–
116.

Dunning Jr, J. B. (2007). *CRC handbook of avian body masses.* CRC press.

Ericson, P.G.P. (2012) Evolution of terrestrial birds in three continents: biogeography
and parallel radiations. *Journal of Biogeography,*39, 813–824.
391

Ferguson-Lees, J., & Christie, D. A. (2001). *Raptors of the World*. Houghton Mifflin
Harcourt.

Ferrer, M. & Negro, J.J. 2004. The near extinction of two large European predators:
super specialists pay a price. *Conservation Biology* 18(2), 344–349.
https://doi.org/10.1111/j.1523-1739.2004.00096.x

Forsman, D. (2016). *Flight identification of raptors of Europe, North Africa and the
Middle East*. Bloomsbury Publishing.

Fuchs, J., Johnson, J. A., & Mindell, D. P. (2015). Rapid diversification of falcons (Aves:
Falconidae) due to expansion of open habitats in the Late Miocene. *Molecular Phylogenetics
and Evolution*, *82*, 166-182.

Harrington, B. et al. (2020). Inkscape. http://www.inkscape.org/.

Herrera, C. M. (1978). On the breeding distribution pattern of European migrant birds:
MacArthur's theme reexamined. *The Auk*, *95*(3), 496-509.

IUCN (2021). The IUCN Red List of Threatened Species. Version 2021-2.
https://www.iucnredlist.org.

Jarvis, E.D. et al. (2014) Whole-genome analyses resolve early branches in the tree of
life of modern birds. Science 346 (6215), 1320-1331. DOI: 10.1126/science.1253451.
409
Jetz, W., Thomas, G. H., Joy, J. B., Hartmann, K., Redding, D., Mooers, A. O. (2014).
Distribution and conservation of global evolutionary distinctness in birds. *Current Biology,* 24,
1-12.

Jetz, W., Thomas, G.H, Joy, J. B., Hartmann, K., Mooers A. O. (2012). The global
diversity of birds in space and time. *Nature*, 491, 444-448.

Keller V, Herrando S, Voříšek P, Franch M, Kipson M, Milanesi P, Martí D, Anton M,
Klvaňová A, Kalyakin MV, Bauer H-G & Foppen RPB (red). 2020. European Breeding Bird Atlas
2: Distribution, Abundance and Change. European Bird Census Council & Lynx Edicions,
Barcelona. 967 pages. ISBN 978-84-16728-38-1.

Knapp M., Thomas J.E., Haile J., Prost S., Ho S.Y.W., Dussex N. *et al.* (2019).
Mitogenomic evidence of close relationships between New Zealand's extinct giant raptors and
small-sized Australian sister-taxa. *Molecular Phylogenetics and Evolution*, 134, 122-128.

Lovette, I. J. (2005). Glacial cycles and the tempo of avian speciation. *Trends in Ecology
& Evolution*, *20*(2), 57-59.

Martínez, J.E. & Sanchez-Zapata, J.A. (1999). Wintering of booted eagles (*Hieraaetus
pennatus*) and short-toed eagles (*Circaetus gallicus*) in Spain. *Ardeola* 46, 93-96.

Martínez-Cruz, B., Godoy, J. A., & Negro, J. J. (2004). Population genetics after
fragmentation: the case of the endangered Spanish imperial eagle (*Aquila adalberti*).
Molecular Ecology, *13*(8), 2243-2255.

Massemin, S., Korpimäki, E., & Wiehn, J. (2000). Reversed sexual size dimorphism in
raptors: evaluation of the hypotheses in kestrels breeding in a temporally changing
environment. *Oecologia*, *124*(1), 26-32.

Mellone, U., De La Puente, J., López-López, P., Limiñana, R., Bermejo, A., & Urios, V.
(2013). Migration routes and wintering areas of Booted Eagles Aquila pennata breeding in
Spain. *Bird Study*, 60(3), 409-413.

Mindell D.P., Fuchs, J.., Johnson, J.A. (2018). Phylogeny, Taxonomy, and Geographic
Diversity of Diurnal Raptors: Falconiformes, Accipitriformes, and Cathartiformes. In: Birds of
Prey: Biology and Conservation in the XXI Century. Eds J.H. Sarasola, J.M. Grande & J.J. Negro.
Springer.

Mönkkönen, M. & Viro, P. (1997) Taxonomic diversity of the terrestrial bird and
mammal fauna in temperate and boreal biomes of the northern hemisphere. *Journal of*
Biogeogrphy,24, 603– 612

Mönkkönen. M. (1994) Diversity patterns in Palaearctic and Nearctic forest bird
assemblages. *Journal of Biogeography,* 21, 183-95.

Nadachowska-Brzyska, K., Li, C., Smeds, L., Zhang, G., & Ellegren, H. (2015). Temporal
dynamics of avian populations during Pleistocene revealed by whole-genome sequences.
Current Biology,25(10), 1375-1380.

Nagy, J., & Tökölyi, J. (2014). Phylogeny, historical biogeography and the evolution of
migration in accipitrid birds of prey (Aves: Accipitriformes). *Ornis Hungarica*, 22(1), 15-35.

Nagy, J., Végvári, Z., & Varga, Z. (2017). Life history traits, bioclimate, and migratory
systems of accipitrid birds of prey (Aves: Accipitriformes). *Biological Journal of the Linnean*
Society, 121(1), 63-71.

Negro, J. J. (1991). Problemas de identificación. El Cernícalo Vulgar y el Primilla. *La*
Garcilla, 82, 5-7.

Negro, J. J. (2018). Raptors and people: An ancient relationship persisting today. In:
Birds of Prey: Biology and Conservation in the XXI Century. Eds J.H. Sarasola, J.M. Grande & J.J.
Negro. Springer.

Negro, J. J., De la Riva, M., & Bustamante, J. (1991). Patterns of winter distribution and
abundance of sedentary lesser Kestrel (*Falco naumanni*) in Spain. *Journal of Raptor*
Research,25, 30-35.

Negro, J.J., Prenda, J., Ferrero, J.J., Rodríguez, A. & Reig-Ferrer, A. (2020). A timeline for
the urbanization of wild birds: the case of the lesser kestrel. *Quaternary Science Reviews*, 249,
106638.

Newton, I. (1979). Population ecology of raptors. Buteo Books, Vermillion, South
Dakota.

Newton, I., & Dale, L. (1996). Relationship between migration and latitude among west
European birds. *Journal of Animal Ecology*, 137-146.

Newton, I., & Dale, L. (1997). Effects of seasonal migration on the latitudinal
distribution ofWest Palaearctic bird species. *Journal of Biogeography.* 24, 781–789.
469
Perrins, C. M., & Snow, D. W. (1998). The Birds of the Western Palearctic. Concise
Edition. *New York.*

QGIS Development Team, (2021). QGIS Geographic Information System. Open Source
Geospatial Foundation. URL http://qgis.org

R Core Team (2013). R: A language and environment for statistical computing. R
Foundation for Statistical Computing, Vienna, Austria. URL http://www.R-project.org/.inburgh.

Rambaut, A., & Drummond, A. J. (2018). FigTree v1. 4.4. Institute of Evolutionary
Biology. University of Edinburgh. UK.

Ramírez, J., Muñoz, A. R., Onrubia, A., De la Cruz, A., Cuenca, D., González, J. M., &
Arroyo, G. M. (2011). Spring movements of Rüppell's Vulture Gyps rueppellii across the Strait
of Gibraltar. *Ostrich*, 82(1), 71-73.

Rodríguez-Rodríguez, E. J., Beltrán, J. F., El Mouden, E. H., Slimani, T., Márquez, R., &
Donaire-Barroso, D. (2020). Climate change challenges IUCN conservation priorities: A test
with western Mediterranean amphibians. *SN Applied Sciences*, 2(2), 1-11.

Salewski, V. and Watt, C. (2017). Bergmann's rule: a biophysiological rule examined in
birds. *Oikos* 126, 161–172.

Snow, D. W. (1978). Relationships between the European and African avifaunas. *Bird
Study*, 25(3), 134-148.

Svensson, L., Mullarney, K., Zetterström, D. and Grant P. J. (1999). Collins Bird Guide:
the most complete guide to the birds of Britain and Europe, 2nd ed. Harper Collins.

Voelker, G. (2010) Repeated vicariance of Eurasian songbird lineages since the Late
Miocene. *Journal of Biogeography*. 37, 1251–1261.
492

493   Wang P, Yao H, Gilbert KJ, Lu Q, Hao Y et al. (2018). Glaciation-based isolation
494 contributed to speciation in a palearctic alpine biodiversity hotspot: evidence from endemic
495 species. *Molecular Phylogenetics and Evolution,* 129,315–324.
496

497   Weir, J. T., & Schluter, D. (2004). Ice sheets promote speciation in boreal birds.
498 *Proceedings of the Royal Society of London. Series B: Biological Sciences*, 271(1551), 1881-
499 1887.

500   Zachos, J., Pagani, M. Sloan, L., Thomas, E., Billups, K. (2001). Trends, Rhythms, and
501 Aberrations in Global Climate 65 Ma to Present. *Science* 292, 686-693.
502

503   Zink, R.M., Klicka, J. & Barber, B.R. (2004). The tempo of avian diversification during
504 the Quaternary. *Philosophical Transactions of the Royal Society of London* B. 359, 215-220. doi:
505 10.1098/rstb.2003.1392, 215-220359 2004.
506

Group	Species	Body mass male(g)	Body mass female (g)	Transaharian migration	IUCN
1	*Pandion haliaetus*	1120-1740	1210-2050	yes	LC
1	*Haliaetus albicilla*	4900-6000	6800-9000	no	LC
2	*Elanus caeruleus*	197-277	219-343	no	LC
3	*Circaetus gallicus*	1200-2000	1300-2300	yes	LC
4	*Pernis aviporus*	440-943	450-1050	yes	LC
5	*Circus aeruginosus*	405-730	540-960	yes	LC
6	*Circus macrorus*	235-416	402-550	yes	NT
6	*Circus pygargus*	227-305	254-445	yes	LC
6	*Circus cyaneus*	300-400	370-708	no	LC
7	*Accipiter gentilis*	517-1110	820-2200	no	LC
7	*Accipiter brevipeps*	135-223	232-275	yes	LC
7	*Accipiter nisus*	105-196	185-350	yes	LC
8	*Milvus migrans*	630-928	750-1080	yes	LC
8	*Milvus milvus*	757-1220	960-1600	no	LC
9	*Buteo buteo*	427-1180	486-1360	yes	LC
9	*Buteo lagopus*	600-1130	783-1660	no	LC
9	*Buteo rufinus*	590-1281	945-1760	yes	LC
10	*Hieraaetus pennatus*	510-770	840-1250	yes	LC
10	*Aquila fasciata*	1560-1560	2000-2500	no	LC
11	*Aquila chrysaetos*	2800-4600	3800-6700	no	LC
11	*Aquila adalberti*	2500-3500	2500-3500	no	VU
11	*Aquila heliaca*	2450-2720	3160-4530	yes	VU
12	*Clanga clanga*	1700-1900	1800-2500	no/ yes	VU
12	*Clanga pomarina*	1000-1400	1300-2200	yes	LC
13	*Aegypius monachus*	7000-11500	7500-12500	no	NT
13	*gyps fulvus*	6200-10500	6500-1100	yes	LC
14	*Gypaetus barbatus*	4500-7000	5600-6700	no	NT
14	*Neophron pernocterus*	1600-2400	1600-2400	yes	EN
15	*Falco eleonorae*	350-390	340-460	yes	LC
15	*Falco subbuteo*	131-232	141-40	yes	LC
16	*Falco peregrinus*	550-750	740-1300	no	LC
16	*Falco pelegrionoides*	610	NA	no	LC
16	*Falco biarmicus*	500-600	700-900	no	LC
16	*Falco rusticolus*	800-1320	1130-2100	no	LC
16	*Falco cherrug*	730-950	970-100	yes	EN
17	*Falco naumanni*	90-172	128-216	yes	LC
17	*Falco vespertinus*	115-190	130-197	yes	NT
17	*Falco tinnunculus*	136-252	154-314	no	LC

509 Table 1. Assemblage of diurnal raptors breeding in continental Europe, ordered according to
510 trophic niche (group, first column). Some of the groups (2 to 5) are monospecific, as they are
511 constituted for highly niche-specific species. Range of body masses for males and females are
512 included, as well as their migratory status (ie., trans-Saharan migrant or not) and IUCN
513 conservation status.

514
515

519　　　　**Figure 1**: Trophic niche cluster dendrogram based on Euclidean distances for the
520　　studied species. * Indicates this species is in its own niche group and not included in others.
521　　Picture credits: *Gypaetus barbatus*, *Accipitier nisus* and *Falco eleonorae* by Manuel Cayuela
522　　López. *Circus pygargus* by Miguel Ángel Rojas. Remaining photographs by the authors.

Figure 2: Phylogeny of included European raptors. Colour areas indicate groups of species with direct niche competition. Colour stars are used to join species not phylogenetically directly related but engage in niche competition (colour areas). Red asterisk indicated long distance migration. Number are species not include in the niche groups (1. *Circus aeruginosus*; 2. *Circaetus gallicus*; 3. *Pernis apivorus*). Picture credits: *Clanga clanga* modified from J.M. Garg (CC BY-SA 3.0). *Buteo lagopus* cropped from Walter Siegmund (CC BY 2.5). *Haliaeetus albicilla* cropped from Christoph Müller (CC BY 4.0). *Falco vespertinus* cropped from Martin Mecnarowski (CC BY 3.0). *Falco biarmicus* cropped from Derek Keats (CC BY 2.0). *Falco cherrug* cropped from DickDaniels (http://carolinabirds.org/) (CC BY-SA 3.0). *Falco pelegrinoides* cropped from Peter Wächtershäuser/ naturlichter.de (CC BY-SA 3.0). *Falco eleonorae* and *Accipiter gentilis* with permission of Manuel Cayuela López. Remaining pictures taken by the authors.

545   **Figure 3:** Pairwise comparisons of body mass (a) and body size (b) of raptor species
546   derived of the foraging groups in Fig. 1. Y-axis represents the migratory species in the pair,
547   while the X-axis the non-migratory species (resident). Dashed lines indicate the diagonal (y =
548   x). Letters refers to species pairs displayed in Table x.

Figure 4: Distribution maps of the raptor species with pairs or trios in our trophic niche
clustering. Ranges of resident species are displayed in blue, while ranges of migrant species are
displayed in red (and green in case of trios). Distribution ranges are taken from the IUCN Red
List Data (IUCN 2021).