1 **Running title: The ptDNAs of nonphotosynthetic Chlamydomonadalean algae**
2
3
4
5 **The plastomes of *Hyalomonas oviformis* and *Hyalogonium fusiforme* evolved**
6 **dissimilar architecture after the loss of photosynthesis**
7
8

9 Alexandra E. DeShaw[1], Francisco Figueroa-Martinez[2], Thomas Pröschold[3], Maike Lorenz[4],
10 Aurora M. Nedelcu[1], David R. Smith[5] and Adrián Reyes-Prieto[1*]
11
12
13 [1]*Department of Biology, University of New Brunswick, Canada*
14 [2]*CONACyT and Universidad Autónoma Metropolitana, México*
15 [3]*Research Department for Limnology Mondsee, University of Innsbruck, Austria*
16 [4]*Experimental Phycology and Culture Collection of Algae, University of Göttingen, Germany*
17 [5]*Western University, Canada*
18
19
*Author for Correspondence:
Adrian Reyes-Prieto
Department of Biology
University of New Brunswick
10 Bailey Drive Fredericton, New Brunswick E3B 5A3, Canada
Telephone: (506) 453 4578
Email: areyes@unb.ca
27
28

29 **Author Contributions**
30
AED performed all laboratory work and computational analyses, in collaboration with FFM.
ARP, AED, DRS, AMN and FFM conceived the original research plan; ML and TP provided
cell cultures and purified DNA, performed microscopy documentation, and prepared the
taxonomic description; ARP supervised experimental and data analysis; AED, DRS and ARP led
the writing of the article with contributions from all the authors; ARP agrees to serve as the
author responsible for contact and communication.
37
38
39 **One sentence summary**
40
41 The plastid genomes of two free-living chlamydomonadalean algae, *Hyalomonas oviformis* and
42 *Hyalogonium fusiforme,* reveal different evolutionary stages following the loss of
43 photosynthesis.
44
45
46

Abstract

The loss of photosynthesis in land plants and algae is typically associated with parasitism but can

also occur in free-living species, including chlamydomonadalean green algae. The plastid

genomes (ptDNAs) of colorless chlamydomonadalean species are surprisingly diverse in

architecture, including highly expanded forms (*Polytoma uvella, Leontynka pallida*) as well as

outright genome loss (*Polytomella* species). Here, we explore the ptDNAs of *Hyalomonas (Hm.)*

oviformis (SAG 62-27; formerly known as *Polytoma oviforme*) and *Hyalogonium (Hg.)*

fusiforme (SAG 62-1c), each representing independent losses of photosynthesis within the

Chlamydomonadales. The *Hm. oviformis* ptDNA is moderately sized (132 kb), smaller than that

of its photosynthetic relative *Hyalomonas chlamydogama* SAG 11-48b (198.3 kb), with a

reduced gene complement but still encoding the ATPase subunits. The *Hg. fusiforme* plastome,

however, is the largest yet observed in colorless plants or algae (~463 kb) and has a coding

repertoire that is almost identical to that of its photosynthetic relatives in the genus

Chlorogonium. Furthermore, the ptDNA of *Hg. fusiforme* shows no clear evidence of

pseudogenization, which is consistent with our analyses showing that *Hg. fusiforme* is the non-

photosynthetic lineage of most recent origin among the known colorless Chlamydomonadales.

Together, these new ptDNAs clearly show that, in contrast to parasitic algae, plastid genome

compaction is not an obligatory route following the loss of photosynthesis in free-living algae,

and that certain chlamydomonadalean algae have a remarkable propensity for genomic

expansion, which can persist regardless of the trophic strategy.

Keywords

Hyalomonas, Polytoma, *Hyalogonium*, *Polytomella,* loss of photosynthesis, plastid genome,

nonphotosynthetic algae, genome reduction, gene losses

Introduction

The loss of photosynthesis has occurred many times throughout eukaryotic evolution (Figueroa-Martinez et al., 2015) and is typically associated with a switch to a parasitic or pathogenic lifestyle in both land plants and algae (Wicke et al., 2013). Nevertheless, the transition to a nonphotosynthetic lifestyle has also occurred in free-living species, such as in various members of the green algal order Chlamydomonadales (Chlorophyceae). There are at least six distinct, free-living chlamydomonadalean lineages that have independently lost photosynthetic capabilities, including species in the genera *Polytomella*, *Polytoma*, *Leontynka*, the isolate *Volvocales sp.* NrCl902, *Hyalogonium fusiforme* and *Hylaomonas oviformis* gen. et sp. nov (Nedelcu, 2001; Vernon et al., 2001; Figueroa-Martinez et al., 2017; Kayama et al., 2020, Pánek et al., 2022; Figure 1).

Previous studies of colorless Chlamydomonadales have identified disparate and intriguing patterns of plastid genome (ptDNA) evolution. *Polytomella* species, for instance, retain a colorless plastid but no longer have ptDNA (Smith & Lee, 2014). *Polytoma uvella* on the other hand, has a highly expanded plastome (~240 kb), resulting from an abundance of noncoding DNA (Figueroa-Martinez et al., 2017). In fact, despite having lost all genes related to photosynthesis, this ptDNA is larger than those of its close photosynthetic relative, *Chlamydomonas leiostraca* (Figueroa-Martinez et al., 2017). However, *Polytomella* and *P. uvella* belong to just two of the six known independently evolved non-photosynthetic chlamydomondalean algae (Figure 1). To improve our understanding of the evolutionary mechanisms underlying the changes in ptDNA organization associated with the loss of

Figure 1

Figure 1. Phylogenetic trees estimated from **A)** nuclear 18S rRNA and **B)** plastid 16S rRNA gene sequences. Both maximum likelihood (ML) trees were estimated with IQ-TREE considering the best fitting nucleotide substitution model in each case (see text for details). The 18S rRNA ML tree was rooted using the Trebouxiophyceae clade as outgroup, in the case of the 16S rRNA tree the outgroup reference were the three members of the OCC (Oedogoniales, Chaetophorales and Chaetopeltidales) clade. Branch lengths are proportional to the number of substitutions per site as indicated by the

photosynthesis in free-living species, we explored ptDNA evolution in *Polytoma* sp. SAG 62-27

(hereafter *Hyalomonas oviformis* gen. et sp. nov.; Figure 2A) and *Hyalogonium fusiforme* SAG

62-1c (Figure 2B), which represent two distinct and poorly studied nonphotosynthetic lineages

within the Chlamydomonadales. These colorless algae are closely related to the photosynthetic

species *Chlamydomonas chlamydogama* (hereafter transferred to *Hyalomonas*) and

Chlorogonium spp., respectively. We found that the *Hm. oviformis* ptDNA is moderately sized

with a reduced gene content typical of other nonphotosynthetic plastomes. The *Hg. fusiforme*

ptDNA, however, retains a near-complete collection of genes involved in photosynthesis with no

obvious examples of pseudogenization. Even more surprising, it represents one of the largest

plastomes yet observed in a nonphotosynthetic plant or algae.

Figure 2. Differential interference contrast micrographs of A) *Hyalomonas oviformis* (SAG 62-27) in low-melting point agarose and B) *Hyalogonium fusiforme* (SAG 62-1c), with the carotene-rich eyespot (e) indicated in the inset image.

Results and Discussion

Hyalomonas, a new non-photosynthetic genus

The strain SAG 62-27 was originally assigned to the genus *Polytoma* with the species epithet

oviforme Pringsheim (nom. nud., without valid description; Koch 1964). Phylogenetic analyses

revealed that this taxon is not closely related to the type species of *Polytoma*, *P. uvella*, and

represents therefore a new genus (Nedelcu, 2001; Figueroa-Martinez et al., 2017). The other

species that belongs to this new genus is the photosynthetic species *Chlamydomonas*

chlamydogama originally described by Bold 1949.

Hyalomonas gen. nov.

Description: Cells biflagellated, with chloroplast of *Agloe*-type containing a single pyrenoid or

reduced parietal leucoplast. Cells surrounded with glycoprotein cell wall. Plastids with

prominent eyespot. Asexual reproduction by 4-8 zoospores. Sexual reproduction by isogamy.

Diagnosis: Differs from other genera of biflagellated chlorophytes by SSU-ITS and plastome

sequence.

Type species (designated here): *Hyalomonas oviformis* sp. nov.

Hyalomonas oviformis sp. nov.

Description: Cells oviform, 11-15 μm long, 8-10 μm wide, two flagella with 1-1½ body length,

with parietal leucoplast containing spot-like anterior eyespot. Nucleus in posterior position. Two

contractile vacuoles (Figure 2A).

Diagnosis: Differs from other species of non-photosynthetic biflagellated chlorophytes by SSU-

ITS (GenBank accession OM985702) and plastome (GenBank accession XXXXXX) sequences.

Holotype (designated here): The strain SAG 62-27 cryopreserved in metabolically inactive state

at the Culture Collection of Algae (SAG), University Göttingen, Germany.

Hyalomonas chlamydogama (Bold) comb. nov.

Basionym: *Chlamydomonas chlamydogama* Bold 1949, *Bull. Torrey Bot. Cl.* **76**, 101, fig. 1

(lectotype designated here).

Epitype (designated here): The strain SAG 11-48b cryopreserved in metabolically inactive state

at the Culture Collection of Algae (SAG), University Göttingen, Germany.

The plastomes of two non-photosynthetic chlamydomonadalean lineages of distinct age

Our phylogenetic analyses show that *Hm. oviformis* and *Hg. fusiforme* lost photosynthesis

independently from one another as well as from other colorless lineages within the

Chlamydomonadales (Figure 2). Among the closest known photosynthetic relatives of *Hm.*

oviformis and *Hg. fusiforme* are *Hm. chlamydogama* and *Cg. euchlorum*, respectively (Figure 2).

RelTime analyses with protein and rDNA sequence data suggest that *Hm. oviformis* and *P. uvella*

likely diverged from their respective photosynthetic siblings during close, likely overlapping

periods (~90-120 MYA) (Figures S1, S4, S5; Table S1). However, our analyses consistently

recovered *Hg. fusiforme* as the colorless lineage of most recent origin among the

chlamydomonadalean taxa investigated here (~45-75 MYA) (Figure S1).

To understand the consequences of the loss of photosynthesis in these two non-photosynthetic

lineages of distinct evolutionary age, we sequenced the ptDNAs of *Hm. oviformis* and *Hg.*

fusiforme as well as those of *Hm. chlamydogama* and *Cg. euchlorum*. The plastomes of *Hm.*

oviformis (132 kb) and *Hm. chlamydogama* (198.3 kb) were assembled into circular-mapping

molecules (Figure 3; Table S2). Unlike *Hm. chlamydogama*, the *Hm. oviformis* ptDNA lacks

inverted repeats, which is a recurring theme among plastomes of nonphotosynthetic plants and

algae (Figueroa-Martinez et al., 2015, 2017; Wicke et al., 2013). However, the presence of a

large number of repeats prevented us from generating intact assemblies of the *Hg. fusiforme* and

Cg. euchlorum ptDNAs, despite employing long-read PacBio sequencing. The *Hg. fusiforme* and

Cg. euchlorum plastome sequences are distributed across 43 and 4 scaffolds with accumulative

lengths of 463 kb and 314 kb, respectively (Figure S2; Table S2).

Figure 3. Circular gene maps of the plastid genomes of **A)** *Hyalomonas oviformis* (SAG 62-27) and **B)** *Hm. chlamydogama* (SAG 11-48b). Large single copy (LSC), small single copy (SSC) and inverted repeat (IRN) regions are indicated. The graphic representations were prepared with the OrganellarGenomeDRAW (OGDRAW) v1.3.1 tool (Greiner et al., 2019) implemented in the CHLORO-BOX server (https://chlorobox.mpimp-golm.mpg.de/index.html).

Chlamydomonadalean ptDNAs are renowned for being extremely hard to assemble (Gaouda et

al., 2018), with many reported assemblies existing in highly fragmented forms (Figueroa-

Martinez et al., 2017; Gaouda et al., 2018). Our read coverage analyses indicate that all the *Hg.*

fusiforme and *Cg. euchlorum* scaffolds are genuine components of the plastid genome and not

ptDNA-like sequences from the mitochondrial or nuclear genomes. The read coverage survey

also supports the existence of an inverted repeat in *Hg. fusiforme* (Table S3).

Comparative plastid genomics among photosynthetic and nonphotosynthetic species

To gain a better understanding of the evolutionary mechanisms that shaped the genetic

architecture of non-photosynthetic plastomes in Chlamydomonadales, we compared the ptDNAs

Figure 4. Comparison of nonsynonymous (dN) nucleotide substitution rates estimated in the five plastid genes encoding subunits of the F-ATPase. Box plots in green colour correspond to nucleotide substitution rates estimated between the species *Hm. chlamydogama, Chlamydomonas leiostraca, Chlamydomonas reinhardtii, Chlorogonium capillatum, Chlorogonium euchlorum* and *Hyalogonium fusiforme*. Box plots in gray color summarize the nucleotide substitution rates between the former six chlamydomonadalean taxa and Hyalomonas oviformis. The Z and U scores and the p values estimated

of the two non-photosynthetic species with those of their closest known photosynthetic relatives

as well as other colorless species. The *Hm. oviformis* plastome is circa 65 kb smaller than that of

the photosynthetic *Hm. chlamydogama* but still large when compared to the ptDNAs from other

colorless plants and algae, which are typically under 75 kb. One of the reasons for this is that

Hm. oviformis encodes all subunits of the cF_0-CF_1 ATPase (*atpA, atpB, atpE, atpF, atpH*, and

atpI). These genes are similar in length to their orthologues in *Hm. chlamydogama,* but some of

them (4/6) show elevated rates of nucleotide substitution (Figure 4). The genes of the *atp* series

are not generally retained in colorless plastids, but notable exceptions to varying degrees come

from the chlamydomonadalean *Leontynka pallida* (Pánek et al., 2022), other diverse colorless

algae (Tanifuji et al., 2020) and angiosperms (Wicke et al., 2013). Nevertheless, like the

plastomes of other colorless plants and algae, the *Hm. oviformis* ptDNA has lost all genes

encoding subunits of the two photosystems, the b_6f complex, and the enzymes for chlorophyll

biosynthesis (Table S4). However, it is still possible to distinguish conserved collinear plastid

gene arrangements between *Hm. oviformis* and the photosynthetic *Hm. chlamydogama* (Figure

S3).

The presence of the cF_0-CF_1 ATPase in a nonphotosynthetic context is intriguing (Dorrell et al.,

2019; Tanifuji et al., 2020). Some have suggested that its hydrolytic activity sustains a proton

electrochemical gradient essential for the internalization of proteins into the thylakoid via the

twin arginine translocator (TAT) system (Dorrell et al., 2019). Support for this view in *Hm.*

oviformis comes from the fact that, in addition to plastid-encoded *atp* genes, we uncovered

nuclear genes for *atpD* and *atpG*, which respectively encode the delta and gamma (γ) subunits of

the plastid cF_0-CF_1 ATPase. The deduced amino acid sequences of *atpD* and *atpG* both contain

standard N-terminus plastid transit peptides, and the latter contains the two cysteine residues

involved in the thioredoxin-mediated modulation of the enzyme hydrolytic activity (Hisabori et

al., 2012) (Figure S7). We also searched the *Hm. oviformis* nuclear contigs for *tat*A, *tat*B, and

*tat*C, which encode the TAT protein import system, but only recovered evidence of *tat*C. Similar

findings were recently reported in the gene repertoire of *L. pallida*, but in that case *tat*A and *tat*B

were identified in transcriptomic data (Pánek et al., 2022).

At circa 463 kb, the ptDNA of *Hg. fusiforme* is not only larger than that of its close

photosynthetic relative *Cg. euchlorum* (314 kb; intron-rich genome), but also the largest

plastome ever observed in a colorless plant or alga, and larger than the vast majority of

sequenced ptDNAs from photosynthetic species. This expanded architecture is primarily the

consequence of inflated intergenic regions as well as apparent accumulation of intron-resembling

sequences (Tables 1 and S6). Close inspection of the intergenic ptDNA of *Hg. fusiforme* revealed

dozens of TrRnEn-like ORFs belonging to different families of homing endonucleases, including

members of the LAGLIDADG (2), GIY-YIG (4) and H-N-H (14) families as well as ORFs

matching to reverse transcriptases and maturases (see Table S6). These ORFs were not evidently

associated with introns and were one of the key reasons why we were unable to bridge the

scaffolds during ptDNA assembly. Other chlamydomonadalean algae, including *Dunaliella*

salina, also have a preponderance of free-standing intronic ORFs within their intergenic regions

(Smith et al., 2010).

The large size of *Hg. fusiforme* ptDNA is not only associated with increased intergenic and

intronic regions. In contrast to the ptDNAs of *Hm. oviformis* and other nonphotosynthetic

chlamydomonadalean species, the *Hg. fusiforme* plastome shows virtually no gene loss as

compared to its photosynthetic close relatives. Indeed, it encodes most subunits of both

photosystems (*psa* and *psb*), the b_6f complex, and the proteins for chlorophyll biosynthesis

present in *Chlorogonium* species (Table S4). What is more intriguing is that the vast majority of

the *Hg. fusiforme* plastid genes typically involved in photosynthesis display no obvious

reductions in size or classic signs of pseudogenization (Table S5), with the exception *chl*B (1,068

bp), which is around 130 amino acids shorter than its *Cg. elongatum* counterpart. Still, the *Hg.*

*fusiforme chl*B encodes all the amino acid residues important for the activity of the

protochlorophyllide oxidoreductase (Bröcker et al., 2010) (Figure S8). It is notable as well that

*psb*E and *psb*Z from *Hg. fusiforme* are split into two pieces by an intergenic spacer (no intron

was detected), but their conceptual translations reveal protein products of similar lengths to those

of photosynthetic relatives. Evolutionary rate analyses indicate that some *Hg. fusiforme* plastid-

encoded proteins involved in light reactions and carbon fixation have higher rates of substitution

than their homologs in photosynthetic taxa (Figure S6). But, altogether, there is no clear

evidence of rampant plastid pseudogenization in *Hg. fusiforme*, despite it having forfeited

photosynthetic capabilities (i.e., no chlorophyll presence; Figure 2A; Pringsheim 1963).

Plastome evolution following the loss of photosynthesis

The loss of photosynthesis has traditionally been associated with plastid gene loss and genomic

streamlining. In parallel, plastome gene repertoires of nonphotosynthetic plants and algae,

regardless of their trophic strategy, tend to converge to relatively homogeneous small gene sets

involved in transcription and translation (Figure S9). Only in rare cases, the plastome remodeling

process after the loss of photosynthesis resulted in the complete loss of the organelle genome

(Smith & Lee, 2014). But as more and more ptDNA sequences from colorless species become

available, it is increasingly obvious that there is not a single narrative for plastome evolution

following the loss of photosynthetic capabilities. This is certainly true for colorless members of

the Chlamydomonadales, where plastid genome information exists for at least six

nonphotosynthetic lineages, each representing an independent loss of photosynthesis within the

order. A key theme emerging from these data is that the ptDNAs of colorless plastids, in

particular those of free-living Chlamydomonadales, are not always characterized by small size,

reduced gene content and compactness. In fact, diverse combinations of genome size, gene

content, and gene density exist in this green algal group. The notable cases include the large

gene-rich plastome of *Hg. fusiforme* as well as the large but gene-reduced ptDNAs of *Leontynka*

pallida (Pánek et al., 2022) and *Polytoma uvella* (Figueroa-Martinez et al., 2017); nevertheless,

all three of these plastid genomes are significantly larger than those of their close photosynthetic

relatives. But this group of colorless algae also includes the relatively small and compact

plastomes of *Hm. oviformis* and *Volvocales* sp. NrCl902 and the complete loss of ptDNA in

Polytomella species.

The observed diversity among ptDNAs of colorless chlamydomonadalean algae highlights the

fact that various aspects of genome organization – size, compaction, and gene content – evolve

independently, even under seemingly similar selective pressures associated with the switch to a

strictly heterotrophic lifestyle. This is because they are likely driven by different evolutionary

mechanisms and forces. For instance, gene loss is primarily a consequence of (1) reduced

functional constraints on photosynthesis-related genes and (2) evolutionary time, whereby the

longer the time since the loss of photosynthesis the greater the potential for gene loss.

Conversely, changes in genome compaction are largely driven by mutational forces and DNA

maintenance machineries acting upon intergenic and intronic regions (Figueroa-Martinez et al.,

2015; Wicke et al., 2013). Thus, it is possible for a plastid genome to be large and gene poor, and

vice versa.

Our data suggest that *Hg. fusiforme* may have lost photosynthetic capabilities in recent

evolutionary time, thus, explaining why it has a near-complete cohort of photosynthetic genes. In

the future, it is possible/expected that some of these genes will be lost, but its propensity for

genomic expansion may (or may not) persist. It should be noted that chlamydomonadalean

ptDNAs, including those of photosynthetic species, are particularly prone to genomic inflation, a

topic that has been discussed in detail elsewhere (Smith et al., 2010; Figueroa-Martinez et al.,

2017; Gaouda et al., 2018). In fact, the largest plastid genome on record belongs to

Haematococcus pluvialis (>1 Mb) (Gaouda et al., 2018), a relatively close relative of *Hg.*

fusiforme. As it stands, the data presented here add to the astonishing amount of plastome

architectural diversity observed across the Chlamydomonadales (in both photosynthetic and

nonphotosynthetic species). How much more diversity remains to be uncovered? Our guess is a

lot.

Materials and Methods

Growth conditions

Hyalomonas oviformis (SAG 62-27) and *Hyalogonium fusiforme* (SAG 62-1c) were grown in

ErbsMS media (Pringsheim, 1946) under dark conditions and no shaking at 16 °C. Cultures of

Hm. chlamydogama (SAG 11-48b) were maintained in standard *Volvox* medium (Kirk et al.,

1999) under constant shaking (200 rpm) at 18°C and a 16-hour-light/8-hour-dark cycle. *Cg.*

euchlorum (SAG 12-2a) cultures were grown in MiEg 1:1 and Eg media (Schlösser, 1994).

DNA extraction and sequencing

Total DNA of *Hm. chlamydogama* was extracted from a 250-mL culture in exponential growth.

The cell pellet was washed three times with saline-EDTA (50 mM Tris-HCl, pH 8, 50 mM NaCl,

and 5 mM EDTA) and then incubated in the presence of proteinase K (33 mg/mL) and 0.5%

SDS for 1 h at 50°C. DNA was extracted by standard phenol-chloroform procedures. Total

genomic DNA from *Hm. oviformis*, *Hg. fusiforme* and *Cg. euchlorum* was extracted using the

DNeasy PlantMini Kit (Qiagen, Hilden, Germany) following manufacturer instructions. Paired-

end libraries were sequenced using Illumina technology (150 bp length reads; 450-bp insert

length; Genome Quebec, McGill University). We also produced low-coverage PacBio reads

(PacBio Sequel, v3 chemistry; IMR Dalhousie University) of *Hg. fusiforme* (413 219 reads; circa

604 Mb) and *Cg. euchlorum* (555 390 reads; circa 680 Mb) from selected DNA fragments of ~

350 Kbp. The 18S rRNA and ITS rRNA sequences of the used strains are available in GenBank

(SAG 62-27: OM985702; SAG 11-48b: OM985703; SAG 62-1c: OM985704; SAG 12-2a:

OM985705).

Genome assembly and sequence annotation

After quality control filtering and trimming, we assembled two high-quality paired-end read sets

of each alga alternatively with Ray v 2.2.0 [k-mer lengths of 21, 31, and 77] (Boisvert et al.,

2010). PacBio SMRT reads of *Hg. fusiforme* and *Cg. euchlorum* were corrected and assembled

with Canu v 2.1 [predicted genome size of 0.4 megabases, using raw PacBio reads for correction

and then corrected reads for assembly] (Koren et al., 2017).

We identified contigs containing typical plastid genes using the Automatic Annotation tool of

Geneious Prime v2019.1.3 (Kearse et al., 2012) with published complete ptDNAs from diverse

Chlamydomonadales as reference. Manual sequence refinement and read coverage evaluations

were carried out with Geneious Prime. Genes encoding tRNAs were predicted with the

tRNAscan-SE Search Server (Lowe & Chan, 2016).

Phylogenetics, relative divergence time and nucleotide substitution rates analyses

Sequence multiple alignments for all analyses were prepared alternatively with the MAFFT

(Katoh & Standley, 2013) tools implemented in Geneious Prime. Resulting raw sequence

matrices were visually inspected and manually edited to discard poorly aligned regions. To test

the independent origin of the colourless lineages *Hm. oviformis* and *Hg. fusiforme* analyzed here,

we estimated maximum likelihood (ML) trees from multiple alignments of nuclear 18S rRNA

(1,521 bp long; 121 taxa) and plastid 16S rRNA (1,257 bp long and 35 taxa) sequences of

diverse chlorophyte green algae. Both ML reconstruction were estimated independently with IQ-

TREE v1.6.11 (Nguyen et al., 2015) considering the best fitting nucleotide substitution model

(TIM3+R3 and TVMe+I+G4, selected respectively by the ModelFinder version implemented in

IQ-TREE v1.6.11) and 10,000 ultra-fast bootstrap replicates in each case. Then, to estimate the

relative divergence times (rDT) of the Chlamydomonadalean colorless lineages, we prepared

additional alignments of 16S rRNA+18S rRNA (2,763 nucleotides long; 28 taxa) sequences and

of conceptual translations of the protein-coding genes *tufa*, *rps*4, *rps*7, *rp*l2 and *rp*l5 (1,163

amino acid residues long; 28 taxa) considering a reduced sample of chlorophyte algae (see

supplementary Table S7). The five protein-coding sequences were selected based of their

presence in ptDNAs of the colorless algae analyzed here, the length of the translated protein

(>150 aa) and reliability (i.e., limited ambiguously aligned or gap-rich regions) of the multiple

alignments. Maximum likelihood trees of the 16S rRNA, 18S rRNA, the concatenated 16S rRNA

and 18S rRNA sets and concatenated protein alignments were independently estimated with IQ-

TREE considering the best fitting model for each sequence matrix (-m TEST option). Then, we

used those maximum likelihood phylogenetic trees as references to estimate rDT with RelTime

approximation implemented in MEGA X (Mello, 2018). We considered the *Proterocladus*

fossils (780 -1000 MY) as the minimum divergence time between Ulvophyceae-Briopsidales and

Chlorophyceae as calibration reference point (Tang et al., 2020).

Finally, we also prepared multiple alignments at nucleotide sequence level of selected protein-

coding regions from diverse photosynthetic and free-living non-photosynthetic

chlamydomonadalean algae to estimate rates of synonymous (*dS*) and nonsynonymous (*dN*)

substitutions with the CODEML option of PAMLX (Xu & Yang, 2013).

Accession numbers

Data deposition: sequences have been deposited in GenBank under accession numbers XXXXX

Acknowledgments

This work was supported by Discovery Grant to A.R.P. from the Natural Sciences and

Engineering Research Council (NSERC) of Canada. A.R.P. was also supported by the Canada

Foundation for Innovation (project 28276) and the New Brunswick Innovation Foundation

(project RIF2012-006).

Tables

Table 1. Plastid genome features of diverse chlorophytes

Taxa	Genbank Accession	Size (kb)	Inverted repeat length (kb)	Coding DNA (%)	GC content (%)	Mean intergenic spacer (bp)	Protein-coding genes	rRNA genes	tRNA genes	Introns
Chlorophyceae										
Hyalomonas oviformis[1]	pending	132	NA	40.6	27.3	617	33	5	27	3
Hyalomonas chlamydogama[1]	pending	198	12.1	40.9	27.6	886	66	10	29	6
Hyalogonium fusiforme[1]	pending	~442	NA	18.4	44	2454	66	6	21	4
Chlorogonium euchlorum[1]	pending	~314	27.9	27.9	36.9	1692	67	6	31	14
Chlorogonium capillatum	KT625085.1-KT620591.1	~271	NA	34.7	38.8	1616	65	3	27	10
Polytoma uvella	KX828177.1	~230	NA	26.8	27.3	2967	24	2	27	1
Chlamydomonas leiostraca	NC_032109	167	13.1	52.4	42.5	600	68	6	28	8
Volvocales sp. NrCl902	LC516060.1	176	NA	39.2	40.1	1974	24	5	30	5
Chlamydomonas reinhardtii	NC_005353.1	204	22.2	37.8	34.5	700	68	10	29	6
Volvox carteri	GU084820.1	>420	16.6	<20	42.8	5103	60	4	27	8
Dunaliella salina	GQ250045.1	269	14	45.5	32.1	935	66	3	28	43
Haematococcus pluvialis	MG677935.1	1352	NA	~90	~50	NA	~71-125	6	31	NA
Leontynka pallida	OM479425.1	362	NA	18	36.1	4726	32	3	26	4
Trebouxiophyceae										
Helicosporidium sp.	NC_008100.1	37	NA	95.2	26.9	34	21	3	25	1
Prototheca wickerhamii	KJ000176.1	56	NA	83.17	31.2	135	34	3	26	0

375   Gray boxes indicate nonphotosynthetic taxa
376   [1]This work
377   NA, no detected or no present.

Figure legends

Figure 1. Phylogenetic trees estimated from **A)** nuclear 18S rRNA and **B)** plastid 16S rRNA

gene sequences. Both maximum likelihood (ML) trees were estimated with IQ-TREE

considering the best fitting nucleotide substitution model in each case (see text for details). The

18S rRNA ML tree was rooted using the Trebouxiophyceae clade as outgroup, in the case of the

16S rRNA tree the outgroup reference were the three members of the OCC (Oedogoniales,

Chaetophorales and Chaetopeltidales) clade. Branch lengths are proportional to the number of

substitutions per site as indicated by the corresponding scale bars. Numbers near nodes indicate

values of bootstrap proportion support (10,000 ultra-fast replicates) and thick branches denote

100% bootstrap. Nonphotosynthetic chlamydomonadalean species are indicated in orange; green

branches represent photosynthetic/mixotrophic lineages.

Figure 2. Differential interference contrast micrographs of **A)** *Hyalomonas oviformis* (SAG 62-

27) in low-melting point agarose and **B)** *Hyalogonium fusiforme* (SAG 62-1c), with the carotene-

rich eyespot (e) indicated in the inset image.

Figure 3. Circular gene maps of the plastid genomes of **A)** *Hyalomonas oviformis* (SAG 62-27)

and **B)** *Hm. chlamydogama* (SAG 11-48b). Large single copy (LSC), small single copy (SSC)

and inverted repeat (IR*N*) regions are indicated. The graphic representations were prepared with

the OrganellarGenomeDRAW (OGDRAW) v1.3.1 tool (Greiner et al., 2019) implemented in the

CHLOROBOX server (https://chlorobox.mpimp-golm.mpg.de/index.html).

Figure 4. Comparison of nonsynonymous (dN) nucleotide substitution rates estimated in the five

plastid genes encoding subunits of the F-ATPase. Box plots in green colour correspond to

nucleotide substitution rates estimated between the species *Hm. chlamydogama*,

Chlamydomonas leiostraca, *Chlamydomonas reinhardtii*, *Chlorogonium capillatum*,

Chlorogonium euchlorum and *Hyalogonium fusiforme.* Box plots in gray color summarize the

nucleotide substitution rates between the former six chlamydomonadalean taxa and *Hyalomonas*

oviformis. The *Z* and *U* scores and the *p* values estimated with the Wilcoxon-Mann-Whitney U

test are indicated in each case.

Supplemental data

Supplemental tables

Supplemental Table S1. RelTime estimated divergence times of non-photosynthetic

chlamydomonadalean algae

Supplemental Table S2. General information of the Illumina and PacBio assemblies

Supplemental Table S3. Average read coverage estimated per genomic compartment

Supplemental Table S4. Comparative gene content of plastid genomes from diverse

chlamydomonadalean algae

Supplemental Table S5. Size of predicted plastid protein-coding regions in *Hyalogonium*

Supplemental Table S6. Functional classification of the plastid *Hyalogonium fusiforme*

TrRnEn-like ORFs

Supplemental Table S7. Taxa considered in the RelTime analyses

Supplemental Figures

Supplemental Figure S1. Divergence time estimates of nonphotosynthetic lineages among the

Order Chlamydomonadales based on the *RelTime* approximation (Tamura et al., 2012) applied to

A) an alignment of 5 plastid encoded proteins (*tuf*A, *rps*4, *rps*7, *rpl*2 and *rpl*5) and **B**) a

concatenated alignment of the 16SrRNA (plastid) and 18S rRNA (nuclear) sequences. For

illustration purposes, the presented trees correspond to subsections of the highest-likelihood trees

estimated with IQ-TREE in each case (LG+R4 for the protein set and GTR+R3 for the

concatenated rRNA data). The complete rooted trees including the outgroup (*Chlorella* clade)

and indicating the fossil calibration point (*Proteolcadus* fossils; 780 -1000 MY) are available in

the supplementary information (Figs. S4 and S5). Black numbers near nodes indicate values of

bootstrap support (10,000 ultra-fast replicates) and red numbers are the *RelTime* estimated

divergence times. Yellow rectangles delimit 95% credibility intervals on node ages. The branch

lengths are proportional to the absolute ages of the nodes.

Supplemental Figure S2. Linear maps of the four assembled scaffolds of the *Chlorogonium*

euchlorum ptDNA. The graphic representations were created with OGDraw (Greiner et al.,

2019).

Supplemental Figure S3. Alignment of the complete plastid genomes of *Hyalomonas oviformis* and *Hm. chlamydogama.* Colored diagonal lines indicate conserved gene blocks. Rectangles in red highlight protein-coding genes associated to the photosynthetic function. Grey rectangles indicate genes encoding rRNAs. The genome alignment was elaborated with the MAUVE v 2.4.0 tool (progressive algorithm default options) implemented in Geneious Prime.

Supplemental Figure S4. Divergence time estimates of nonphotosynthetic lineages among the Order Chlamydomonadales based on a RelTime approximation (Tamura et al., 2012) applied to an alignment of 5 plastid encoded proteins (*tuf*A, *rps*4*, rps*7*, rp*l2 and *rp*l5). The tree topology corresponds to a the highest-likelihood tree (Log Likelihood = -16540.505) estimated with IQ-TREE considering the best fitting amino acid substitution model (LG+R4) identified by the ModelFinder version implemented in IQ-TREE v1.6.11. Nonphotosynthetic taxa are highlighted with orange circles to the right. The outgroup (*Chlorella* clade) and the fossil calibration point (*Proteolcadus* fossils; 780 -1000 MY; red dot) are indicated. Black numbers near nodes indicate values of bootstrap support (10,000 ultra-fast replicates) and red numbers are the *RelTime*-estimated divergence times. Grey rectangles delimit 95% credibility intervals on node ages. The branch lengths proportional to the absolute ages of the nodes.

Supplemental Figure S5. Divergence time estimates of nonphotosynthetic lineages among the Order Chlamydomonadales based on a RelTime approximation (Tamura et al., 2012) applied to a concatenated alignment of the 16SrRNA and 18S rRNA sequences. The tree topology corresponds to a the highest-likelihood tree (Log Likelihood = -14261.073) estimated with IQ-

TREE considering the best fitting model (GTR+R3) identified by identified by the ModelFinder

version implemented in IQ-TREE v1.6.11. Nonphotosynthetic taxa are highlighted with orange

circles to the right. The outgroup (*Chlorella* clade) and the fossil calibration point (*Proteolcadus*

fossils; 780 -1000 MY; red dot) are indicated. Black numbers near nodes indicate values of

bootstrap support (10,000 ultra-fast replicates), red numbers are the *RelTime*-estimated

divergence times. Grey rectangles delimit 95% credibility intervals on node ages. The branch

lengths proportional to the absolute ages of the nodes.

Supplemental Figure S6. Comparison of nonsynonymous (dN) and synonymous (dS)

nucleotide substitution rates estimated in photosynthesis-related plastid genes of *Hyalogonium*

fusiforme. Box plots in green color correspond to nucleotide substitution rates estimated between

the photosynthetic species *Hyalomonas chlamydogama*, *Chlamydomonas leiostraca*,

Chlamydomonas reinhardtii, *Chlorogonium capillatum* and *Chlorogonium euchlorum.* Box plots

colored in gray summarize the nucleotide substitution rates between the five mentioned

photosynthetic taxa and *Hyalogonium fusiforme*.

Supplemental Figure S7. Multiple protein sequence alignments of the **A**) γ and **B**) δ subunits of

the cF_0-CF_1 ATPase from diverse viridiplants. Amino acid residues highlighted in yellow colour

represent the transit peptides for import into the plastid predicted by TargetP 2.0

[http://www.cbs.dtu.dk/services/TargetP/] (Armenteros et al., 2019). In the case of the γ subunit

multiple alignment (panel **A**), the columns emphasized in black background indicate the two

cysteine residues typically involved in the thiol-modulation of the cF_0-CF_1 ATPase activity.

Supplemental Figure S8. Multiple protein sequence alignments of the ChlB from

Chlorogonium species and *Hyalogonium fusiforme*. Residues highlighted in magenta and cyan

represent amino acids involved in the binding and stabilization of the [4Fe-4S] cluster of the

protochlorophyllide oxidoreductase. The residues with green background indicate amino acids

participating in the interaction of ChlB (binding) with ChlL (Bröcker et al., 2010).

Supplemental Figure S9. Comparative content of protein-coding genes in plastid genomes of

diverse non photosynthetic green algae and *Euglena (Astasia) longa,* which contains a secondary

plastid of green algal origin. Gene names with asterisks refer to *gene series* encoding subunit of

the CF-ATPase (*atp* series), the b_6f complex (*pet* series), photosystem I (*psa* series) and

photosystem II (*psb* series). Taxa codes: As, *Euglena longa*; Pw, *Prototheca wickerhamii*; He,

Helicosporidium sp.; Pu, *Polytoma uvella*; Vs, *Volvocales* NrCl902; Ho, *Hyalomonas oviformis*;

Hf, *Hyalogonium fusiforme*.

Parsed Citations

Armenteros JJA, Salvatore, M, Emanuelsson, O, Winther, O, Heijne, G von, Elofsson A, Nielsen H (2019). Detecting sequence signals in targeting peptides using deep learning. Life Science Alliance, 2: e201900429 . https://doi.org/10.26508/LSA.201900429
  Google Scholar: Author Only Title Only Author and Title

Boisvert, S, Laviolette, F, Corbeil J (2010) Ray: Simultaneous assembly of reads from a mix of high-throughput sequencing technologies. J Comput Biol 17: 1519–1533. https://doi.org/10.1089/cmb.2009.0238
  Google Scholar: Author Only Title Only Author and Title

Bold HC (1949). The morphology of Chlamydomonas chlamydogama, sp. nov. Bull Torrey Bot Cl., 76: 101-108. https://doi.org/10.2307/2482218
  Google Scholar: Author Only Title Only Author and Title

Bröcker MJ, Schomburg, S, Heinz, DW, Jahn, D, Schubert, WD, Moser J (2010). Crystal structure of the nitrogenase-like dark operative protochlorophyllide pxidoreductase catalytic complex (ChlN/ChlB)2. J Biol Chem 285: 27336–27345. https://doi.org/10.1074/jbc.M110.126698
  Google Scholar: Author Only Title Only Author and Title

Dorrell, R.G., Azuma T, Nomura M, de Kerdrel GA, Paoli L, Yang S, Bowler C, Ishii K, Miyashita H, Gile GH, Kamikawa R (2019) Principles of plastid reductive evolution illuminated by nonphotosynthetic chrysophytes. Proc Natl Acad Sc USA 116: 6914–6923. https://doi.org/10.1073/pnas.1819976116
  Google Scholar: Author Only Title Only Author and Title

Figueroa-Martinez F, Nedelcu AM, Smith DR, Reyes-Prieto A (2015). When the lights go out: the evolutionary fate of free-living colorless green algae. New Phytologist 206: 972–982. https://doi.org/10.1111/nph.13279
  Google Scholar: Author Only Title Only Author and Title

Figueroa-Martinez F, Nedelcu AM, Smith DR, Reyes-Prieto A (2017) The plastid genome of Polytoma uvella is the largest known among colorless algae and plants and reflects contrasting evolutionary paths to nonphotosynthetic lifestyles. Plant Physiol 173: 932–943. https://doi.org/10.1104/pp.16.01628
  Google Scholar: Author Only Title Only Author and Title

Gaouda H, Hamaji T, Yamamoto, K, Kawai-Toyooka H, Suzuki M, Noguchi H, Minakuchi Y, Toyoda A, Fujiyama A, Nozaki H, Smith D. R. (2018). Exploring the limits and causes of plastid genome expansion in Volvocine green algae. Genome Biol Evol 10: 2248–2254. https://doi.org/10.1093/GBE/EVY175
  Google Scholar: Author Only Title Only Author and Title

Greiner S, Lehwark P, Bock R (2019) OrganellarGenomeDRAW (OGDRAW) version 1.3.1: expanded toolkit for the graphical visualization of organellar genomes. Nucleic Acids Res 47: W59–W64. https://doi.org/10.1093/nar/gkz238
  Google Scholar: Author Only Title Only Author and Title

Hisabori T, Sunamura EI, Kim, Y, Konno H (2012). The chloroplast ATP synthase features the characteristic redox regulation machinery. Antioxid Redox Signal 19: 1846–1854. https://doi.org/10.1089/ars.2012.5044
  Google Scholar: Author Only Title Only Author and Title

Katoh K, Standley DM (2013) MAFFT Multiple Sequence Alignment Software Version 7: Improvements in performance and psability. Mol Biol Evol 30: 772–780. https://doi.org/10.1093/MOLBEV/MST010
  Google Scholar: Author Only Title Only Author and Title

Kearse M, Moir R, Wilson A, Stones-Havas S, Cheung M, Sturrock S, Buxton S, Cooper A, Markowitz S, Duran C, Thierer T, Ashton B, Meintjes P, Drummond A (2012). Geneious Basic: An integrated and extendable desktop software platform for the organization and analysis of sequence data. Bioinformatics 28: 1647–1649. https://doi.org/10.1093/bioinformatics/bts199
  Google Scholar: Author Only Title Only Author and Title

Kirk MM, Stark K, Miller SM, Muller W, Taillon BE, Gruber H, Schmitt R, Kirk DL (1999) regA, a Volvox gene that plays a central role in germ-soma differentiation, encodes a novel regulatory protein. Development 126: 639–647. https://dev.biologists.org/content/126/4/639
  Google Scholar: Author Only Title Only Author and Title

Kayama M, Chen JF, Nakada T, Nishimura Y, Shikanai T, Azuma T, Miyashita H, Takaichi S, Kashiyama Y, Kamikawa R (2020) A non-photosynthetic green alga illuminates the reductive evolution of plastid electron transport systems. BMC Biology 18: 126. https://doi.org/10.1186/s12915-020-00853-w
  Google Scholar: Author Only Title Only Author and Title

Koren S, Walenz BP, Berlin K, Miller JR, Bergman NH, Phillippy AM (2017) Canu: scalable and accurate long-read assembly via adaptive k-mer weighting and repeat separation. Genome Res 27: 722-736 https://doi.org/10.1101/gr.215087.116
  Google Scholar: Author Only Title Only Author and Title

Lowe TM, Chan PP (2016) tRNAscan-SE On-line: integrating search and context for analysis of transfer RNA genes. Nucleic Acids

Res 44: W54-57. https://doi.org/10.1093/nar/gkw413

Google Scholar: Author Only Title Only Author and Title

Mello B (2018). Estimating TimeTrees with MEGA and the TimeTree Resource. Mol Biol Evol 35: 2334–2342. https://doi.org/10.1093/molbev/msy133

Google Scholar: Author Only Title Only Author and Title

Nedelcu AM (2001) Complex patterns of plastid 16S rRNA gene evolution in nonphotosynthetic green algae. J Mol Evol 53: 670–679. https://doi.org/10.1007/s002390010254

Google Scholar: Author Only Title Only Author and Title

Nguyen LT, Schmidt HA, von Haeseler A, Minh BQ (2015) IQ-TREE: A fast and effective stochastic algorithm for estimating maximum-likelihood phylogenies. Mol Biology Evol 32: 268–274. https://doi.org/10.1093/molbev/msu300

Google Scholar: Author Only Title Only Author and Title

Pánek T, Barcytė D, Treitli SC, Záhonová K, Sokol M, Ševčíková T, Zadrobílková E, Jaške K, Yubuki N, Čepička I, Eliáš M (2022). A new lineage of non-photosynthetic green algae with extreme organellar genomes. BMC Biology 20: 66 https://doi.org/10.1186/s12915-022-01263-w

Google Scholar: Author Only Title Only Author and Title

Pringsheim EG (1946). Pure Cultures of Algae: their preparation and maintenance. Cambridge University Press, UK.

Google Scholar: Author Only Title Only Author and Title

Pringsheim EG (1963). Farblose Algen. Ein Beitrag zur Evolutionsforschung. Gustav Fischer Verlag, Stuttgart, Germany.

Google Scholar: Author Only Title Only Author and Title

Schlösser UG (1994) SAG-Sammlung von Algenkulturen at the University of Göttingen. Catalogue of strains. Bot Acta 107: 113-186. https://doi.org/10.1111/j.1438-8677.1994.tb00784.x

Google Scholar: Author Only Title Only Author and Title

Smith DR, Lee RW (2014) A Plastid without a Genome: Evidence from the Nonphotosynthetic Green Algal Genus Polytomella. Plant Physiol 164: 1812–1819. https://doi.org/10.1104/pp.113.233718

Google Scholar: Author Only Title Only Author and Title

Smith DR, Lee RW, Cushman JC, Magnuson JK, Tran D, Polle JEW (2010). The Dunaliella salina organelle genomes: large sequences, inflated with intronic and intergenic DNA. BMC Plant Biol 10: 83. https://doi.org/10.1186/1471-2229-10-83

Google Scholar: Author Only Title Only Author and Title

Tamura K, Battistuzzi, FU, Billing-Ross P, Murillo O, Filipski A, Kumar S (2012) Estimating divergence times in large molecular phylogenies. Proc Natl Acad Sc USA 109: 19333–19338. https://doi.org/10.1073/pnas.1213199109

Google Scholar: Author Only Title Only Author and Title

Tang Q, Pang K, Yuan, X, Xiao S (2020) A one-billion-year-old multicellular chlorophyte. Nat. Ecol Evol 4: 543–549. https://doi.org/10.1038/s41559-020-1122-9

Google Scholar: Author Only Title Only Author and Title

Tanifuji G, Kamikawa R, Moore CE, Mills T, Onodera NT, Kashiyama, Archibald JM, Inagaki Y, Hashimoto T (2020) Comparative plastid genomics of Cryptomonas species reveals fine-scale genomic responses to loss of photosynthesis. Genome Biol Evol 12: 3926–3937. https://doi.org/10.1093/gbe/evaa001

Google Scholar: Author Only Title Only Author and Title

Vernon D, Gutell RR, Cannone JJ, Rumpf RW, Birky J (2001) Accelerated evolution of functional plastid rRNA and elongation factor genes due to reduced protein synthetic load after the loss of photosynthesis in the chlorophyte alga Polytoma. Mol Biol Evol 18: 1810–1822. https://doi.org/10.1093/oxfordjournals.molbev.a003968

Google Scholar: Author Only Title Only Author and Title

Wicke S, Müller KF, de Pamphilis CW, Quandt D, Wickett NJ, Zhang Y, Renner SS, Schneeweiss GM (2013) Mechanisms of functional and physical penome peduction in photosynthetic and nonphotosynthetic parasitic plants of the broomrape family. Plant Cell 25: 3711–3725. https://doi.org/10.1105/tpc.113.113373

Google Scholar: Author Only Title Only Author and Title

Xu B, Yang Z (2013) PamlX: A graphical user interface for PAML. Mol Biol Evol 30: 2723–2724. https://doi.org/10.1093/molbev/mst179

Google Scholar: Author Only Title Only Author and Title