Accounting for *cis*-regulatory constraint prioritizes genes likely to affect species-specific traits

Alexander L. Starr[1], David Gokhman[1,2], Hunter B. Fraser[1,3]

1. Department of Biology, Stanford University, Stanford, CA, USA

2. Current address: Department of Molecular Genetics, Weizmann Institute of Science, Rehovot, Israel

3. Correspondence: hbfraser@stanford.edu

Abstract

Measuring allele-specific expression in interspecies hybrids is a powerful way to detect cis-regulatory changes underlying adaptation. However, it remains difficult to identify genes most likely to explain species-specific traits. Here, we outline a simple strategy that leverages population-scale allele-specific RNA-seq data to identify genes that have constrained *cis*-regulation within species yet show divergence between species. Applying this strategy to data from human-chimpanzee hybrid cortical spheroids, we identify signatures of lineage-specific selection on genes related to cellular proliferation, speech, and glucose metabolism. We also highlight cis-regulatory divergence in *CUX1* and *EDNRB* that may shape the unique trajectory of human brain development.

Key words: Hybrid, gene expression, evolution, human, chimpanzee, primate, brain, development

Background

Changes in gene expression are thought to play a major role in the evolution of complex traits [1–4]. As a result, comparing gene expression between species can enable the identification of molecular changes underlying phenotypic divergence. However, obtaining accurate comparisons of gene expression between species is challenging due to confounding factors like age, environmental effects, differential cell type abundances, differences in developmental timing, and batch effects [5–7]. The use of interspecies hybrids overcomes these issues through the measurement of allele-specific expression (ASE) [8,9]. In hybrids, the genomes of both species share the same nucleus and are exposed to identical environments, so there are no confounding environmental, batch, compositional, or developmental effects. This approach has been successfully applied in many species and advanced our understanding of the evolution of gene regulation and its role in establishing phenotypic variation [10–13]. Furthermore, the recent development of human-chimpanzee allotetraploid "hybrid" cells and organoids enables detailed, accurate quantification of differences in gene expression between humans and our closest living relatives [8,9,14].

Hybrids also enable the separation of *cis* and *trans* components of interspecies differences in gene expression [8,9]. The *cis*-component is caused by differences in regulatory elements such as promoters or enhancers that only affect the expression of a nearby gene or genes on the same DNA molecule. The *trans*-component stems from changes in diffusible molecules such as transcription factors that can regulate gene expression throughout the genome. In hybrids the genomes of the two species are exposed to the same *trans* factors. As a result, allele-specific differences in gene expression can only be explained by *cis*-regulatory differences. In addition, using ASE to identify differentially expressed genes (referred to as AS-DEGs) enables

elimination of many confounding factors (including the environmental, batch, compositional, and

developmental timing effects mentioned above). This not only increases the signal-to-noise

ratio, but also disentangles important, potentially evolutionarily significant gene-specific *cis*-

regulation from broad *trans*-acting changes [4,15,16].

While the resulting list of AS-DEGs from hybrids is likely more accurate and isolates the *cis*-

regulatory component, there are often thousands of AS-DEGs which makes it difficult to

prioritize candidate genes and pathways that may have played a major role in evolution.

Differential expression p-values and fold changes are commonly used to rank genes in

comparative RNA-seq studies. However, large and significant fold changes may often result

from low evolutionary constraint on their expression levels, as opposed to being under positive

directional selection. These large fold changes in unconstrained genes (e.g. pseudogenes)

could result in no or very limited phenotypic changes, since a lack of constraint implies a lack of

phenotypic consequence of changes in expression. Therefore, a large and significant fold

change alone is not sufficient to determine the importance of the gene in the evolution of the

parental species. For example, consider a gene whose expression varies by two-fold between

species. If this gene also varies by two-fold within each species individual members of the same

species, it is unlikely to account for any species-specific phenotypes. In contrast, a gene that is

under strong stabilizing selection—with little variation in expression within species but with a

two-fold change between species–is more likely to have contributed to phenotypic divergence

between species. Most studies of expression divergence between species do not include any

comparison to within-species variation; the few exceptions have been limited by small sample

sizes and the confounding factors inherent to any between-species comparison [17–20].

Here, we describe a method that leverages population-scale ASE data to approximate

constraint on *cis*-regulation of gene expression. This method ranks AS-DEGs identified from

interspecies hybrids in a way that is likely more able to prioritize adaptive, functionally significant

changes. We apply this method to ASE data from human-chimpanzee hybrid cortical spheroids

that recapitulate the gene expression patterns of the developing cerebral cortex. Using this

dataset, we identify lineage-specific selection on the expression of genes related to speech,

glucose metabolism, cellular proliferation, and glycan degradation [8,9]. In addition, we highlight

divergence in the expression of *CUX1* and *EDNRB* that may have played an important role in

human brain evolution.

Results

In a typical ASE pipeline, AS-DEGs are identified by comparing the RNA-seq read counts from

the allele from species 1 and the allele from species 2 and ranked using a p-value for differential

expression (Fig. 1A). Various enrichment tests and knowledge from the literature can then be

used to identify interesting trends and prioritize candidate genes. However, these previous

methods do not consider within-species variation in gene expression levels (Fig. 1B). If some

genes have highly variable expression even within a single species, then differences of a similar

(or smaller) magnitude between species are unlikely to explain species-specific traits (e.g.

PDPR in Fig. 1B). Conversely, differential expression of genes with highly constrained

expression in at least one species are more likely to be responsible for differences in organismal

phenotypes between species. For example, ZNF331 and RPS16 have similar fold change

magnitudes in human-chimpanzee hybrid cortical spheroids (Fig. 1C). However, the fold change

for ZNF331 lies well within the distribution of fold changes between alleles in the human

population whereas the fold changes for RPS16 are nearly outside the human population

distribution (Fig. 1C). This indicates that the expression level of RPS16 is much more

constrained and that its differential *cis*-regulation between human and chimpanzee is more likely

to have phenotypic consequences.

Figure 1 Outline of methodology: A) Outline of a typical ASE pipeline. Hybrids are generated

and RNA-seq is used to determine the relative expression of each allele. The ASE ratio is

computed as the ratio of species-specific read counts between the two alleles. **B)** The

distribution of the variance in ASE ratio for each gene in the GTEx data. Insets in orange show

two genes at the extreme ends (EEF2 with low variance suggesting strong stabilizing selection,

and PDPR with high variance suggesting less constraint on gene expression). **C)** Schematic of

incorporation of the interspecies ASE and population level ASE. ZNF331 has a wide range of

ASE values, and the human-chimpanzee ASE is well within the population distribution whereas

human-chimp ASE in RPS16 is on the edge of the population distribution indicating greater

potential for functional relevance. For both **B** and **C**, only GTEx brain samples were used as this

provided clearer illustrative examples, though results are similar using all GTEx samples (Fig

2A).

To systematically apply this concept, we compute the distribution of ASE in one population and

use the Mann-Whitney U test to compare the population-level ASE distribution to the

interspecies ASE distribution for that gene (see Methods) [21]. The p-value reflects how

divergent the ASE of the gene is between species compared to its divergence within species.

We then use those p-values to separately rank genes with DESeq2 false discovery rate (FDR)

less than or equal to 0.1 and greater than 0.1 such that genes with FDR less than or equal to

0.1 are always ranked higher than genes with FDR greater than 0.1. This method prioritizes

genes based on whether the interspecies difference in ASE is of greater magnitude than would

be expected from the intra-species ASE distribution while minimizing false positives. Notably,

ranking solely by the Mann-Whitney U Test p-value would potentially rank lowly expressed

genes that are not significantly differentially expressed between species very highly, increasing

the number of genes falsely identified as differentially expressed. This problem is eliminated by

incorporating the DESeq2 FDR. The ranked list is then used in enrichment analyses and to

identify of top candidates [22,23].

We use a large publicly available data set, GTEx, to estimate ASE variance for every gene. The

GTEx v8 data includes RNA-seq from 838 individuals and 54 tissues, a total of 15,253 samples

in which ASE values have been previously estimated [21]. To minimize any allelic bias in the

population distribution, we normalize the median to 1. The resulting rankings are robust to

choosing either the reference or alternative allele as the numerator for the population

distribution (spearman's rho = 0.999 p < 10^{-300}). Notably, we minimize any batch effects

between GTEx and the human-chimpanzee hybrid dataset by focusing on ASE since direct

comparisons of expression levels between data sets are not involved. For example, if some

technical factor (e.g. sequencing platform or RNA isolation method) caused a gene to show a

spurious 2-fold higher expression in GTEx samples compared to the hybrid data, this would

cancel out in the GTEx ASE calculation.

First, we tested whether the variance in ASE in the human population is a reasonable proxy for

constraint on gene expression. We compared the variance of the GTEx ASE distribution for

each gene to its probability of haploinsufficiency score (pHI), a measure of sensitivity to a 50%

reduction in gene dosage [24]. We observe a significant negative correlation (Spearman's rho =

-0.28, $p < 10^{-170}$) between the two measures (Supp. Fig. 1) indicating that the variance of the

population-scale ASE distribution provides a reasonable proxy for constraint on expression

levels. This correlation remains significant when only five (as opposed to ten) reads from each

allele for a gene in a sample are required for inclusion (Spearman's rho = -0.27, $p < 10^{-163}$

). Furthermore, as pHI is computed from copy number variation, there is no circularity in

reasoning when comparing pHI to the variance of the GTEx ASE distribution [24]. While this

indicates that variation in ASE is sufficient to approximate evolutionary constraint on gene

expression, the strength of this relationship should not be interpreted as a quantitative estimate

of how well ASE represents constraint. Overall, the correlation with pHI indicates that within-

species ASE variance contains useful information about evolutionary constraint on gene

expression.

In addition, it is important to determine the extent to which sample heterogeneity may affect

ASE variance and thereby impact our results. For example, GTEx contains data from a wide

range of adult tissues and donors have variable sex and ancestry. To determine to what extent

these factors affect our results, we divided the GTEx dataset along lines of sex (male/female),

ancestry (African descent/European descent), and brain/non-brain tissues. In all cases, the

results obtained were very highly correlated (spearman's rho > 0.97, $p < 10^{-170}$, Figure 2 A-C).

This indicates that sample heterogeneity is unlikely to have major effects on our results and that

aggregating across all the GTEx samples provides a reasonable measure of constraint on gene

expression within the human population.

In addition, the wide range of adult tissues in GTEx may not accurately reflect gene expression

constraint in cortical spheroids, which mimic fetal development [25,26]. To test this, we used an

ASE dataset generated from neural progenitors, neurons, and fetal cortical wall which

resembles the developmental stage of cortical spheroids more closely than GTEx samples [27].

There is a strong correlation between the rankings generated by comparing to the fetal cortex

dataset and GTEx (Spearman's rho = 0.92, p < 10^{-170} for each of day 50, day 100, and day 150

after initial differentiation of cortical spheroids; Fig 2D). While it is likely that the ASE variance

for some genes changes over development, these genes appear to be somewhat rare. Overall,

these results (Fig 2) indicate that even though gene expression levels vary considerably across

samples, ASE variance is robust to sample heterogeneity. This suggests that within- and

between-species ASE values can be meaningfully compared even when they are not based on

the same tissues or developmental timepoints. We therefore focused our analysis on the full

GTEx data in order to maximize statistical power but verified results for specific genes in the

fetal cortex data when appropriate.

Figure 2 Influence of population differences on gene rankings: A) Comparison of ranks

generated by our method derived from brain vs. non-brain samples. Spearman's rho = 0.98 p <

10^{-170}. **B)** Comparison of ranks using only GTEx subjects of African descent vs. subjects of

European descent. Spearman's rho = 0.98, p < 10^{-170}. **C)** Comparison of ranks using only male

vs. female GTEx subjects. Spearman's rho = 0.99, p < 10^{-170}. **D)** Comparison of ranks derived

from GTEx (across all tissues) vs. data from fetal cortex and primary fetal neurons/neural

progenitors. Spearman's rho = 0.92, p < 10^{-170}.

To highlight differences between our method and the traditional method for ranking genes (Fig

1A vs. Fig 1C), we computed the difference in ranks between the two methods and used

GSEAPY to identify gene sets enriched near the top or bottom of the resulting sorted list

(referred to as the difference in ranks list, Fig 3A) [28]. A top-ranking gene would be one that is

lowly ranked by the traditional method but highly ranked by the population comparison method.

Many gene ontology (GO) categories related to cortical development were enriched near the top

of the difference in ranks list including calcium and potassium channel activity, various

transcription factor (TF) related terms, and cytokine activity (Fig 3B, C and Additional file 1).

This is consistent with previous observations that TFs, ion channel subunits, and important

players in canonical signaling pathways tend to have strongly constrained expression compared

to other genes [24]. Many genes that drive the enrichment of the term "RNA polymerase II

regulatory region sequence-specific DNA binding" are haploinsufficient and play key roles in

neurodevelopment including *MEF2C*, *NEUROD2*, and *CUX1* (Supp. Fig. 2A) [29–31].

Differences in the expression of haploinsufficient genes are more likely to have phenotypic

consequences than differences in the expression of genes for which loss of one copy has no

clear phenotypic effect. In particular, there is strong evidence that both increases and decreases

in CUX1 expression alter neurodevelopment in humans, which implies that the CUX1

expression divergence between humans and chimpanzees is likely to have phenotypic

consequences (Box 1) [29,32].

Figure 3 Enrichment summary for difference in ranks: A) Pipeline for gene set enrichment

analysis. Gene set databases tested include Gene Ontology (GO) Cellular Component, GO

Molecular Function, Kyoto Encyclopedia of Genes and Genomes (KEGG) human gene sets,

Human Phenotype Ontology (HPO), and REACTOME [33–37]. Enrichment analysis is

performed for each time point that cortical spheroids frozen for RNAseq (day 50, day 100, and

day 150) separately and significant terms were aggregated with only one copy of redundant

terms that were significant in multiple timepoints included. Only genes with sufficient reads in

the human-chimpanzee cortical spheroid dataset and a sufficient number of samples in GTEx

were included (see Methods), leaving approximately 10,000 genes in the list used for

enrichment analysis. **B)** Summary of GO Cellular Component enrichments across all time points

with false discovery rate (FDR) < 0.05. REVIGO in conjunction with a custom python script was

used to generate the plot. The axes are derived from multidimensional scaling and measure

semantic similarity, enabling removal of redundant GO terms and visualization of the similarity

between GO categories. Each circle represents a GO term and circles near each other contain

similar genes in the corresponding gene set. Labeled gene sets are generally those with the

lowest FDR in a cluster of terms on the plot. The size of the circles indicates the number of

genes that are driving the enrichment for that category. **C)** Summary of GO Molecular Function

enrichments across all time points with FDR < 0.05.

Box 1: Recent work has linked a mutation in a human accelerated region (HAR) that likely

increases *CUX1* expression to autism spectrum disorder, suggesting that human-derived

changes in *CUX1* expression alter human behavior [32,38]. *CUX1* is expressed at a lower level

in humans than chimpanzees across time points in both hybrid and parental cortical spheroids

(e.g. $\log_2$ fold-change of -0.93, FDR < 0.005 in parentals, mean $\log_2$ fold-change of -0.74, FDR <

0.023 in hybrids at day 150 of differentiation) and the per-sample ASE ratios are well outside the

human fetal cortex population ASE distribution (Supp. Fig. 2 B-E). This reduced expression in

humans is surprising considering that a recent massively parallel reporter assay (MPRA) found

that the HAR linked to *CUX1* should increase expression in humans [38]. As expression is lower

in humans, haploinsufficiency of *CUX1* might provide a reasonable model of the phenotypic

consequences of this change. *CUX1* haploinsufficiency in humans leads to delayed

development of speech and motor skills [29]. One aspect of this condition is that individuals with

one functional *CUX1* allele often close the developmental gap over time (i.e. cognitive

impairments and delays disappear with age) [29]. This "catch up" phenotype is very rare and

may even be specific to *CUX1* haploinsufficiency [29]. Interestingly, humans develop more

slowly than other great apes (known as neoteny) but eventually "catch up," reminiscent of the

CUX1 phenotype. While changes in *CUX1* expression may have played a role in causing

human neoteny, investigation of the development of layer II-III cortical neurons and behavior of

CUX1 haploinsufficient mice will be required to explore this further.

Applying the Sign Test to Constrained Differentially Expressed Genes

While our method is designed to prioritize genes that are relevant to phenotypic differences

between species, it does not, on its own, imply selection on changes in gene expression. On the

other hand, the sign test is designed to detect lineage-specific selection based on systematic

up- or down-regulation of genes that deviates from neutral expectation [39,40]. One important

requirement of the test is that the expression divergence of every gene being tested should be

driven by independent genetic differences. For example, a single mutation in a *trans*-acting

factor could cause a whole pathway to be down-regulated, but this would only count as one

genetic difference. For this reason, hybrids are ideal for applying the sign test to gene

expression, since *cis*-regulation of genes is typically independent (except in the case of some

neighboring genes that share *cis*-elements, which can be accounted for by using a minimum

distance threshold between genes). For example, in the human-chimpanzee hybrid, if a

particular pathway contains significantly more genes with higher expression from the human

allele than the chimpanzee allele (significance determined by a binomial test), that would

provide evidence of selection on that pathway in the human and/or chimpanzee lineage. Similar

to all tests of selection, the sign test cannot discern what the cause of the selective pressure is.

For example, many *cis*-regulatory changes could be compensating for a change in a single

trans-acting factor. Alternatively, changes in gene expression might cause a phenotypic change

that is being selected for. In either case, the sign test provides evidence of lineage-specific

selection on cis-regulation.

While our method (Fig 1C) does not require comparison to the traditional method (Fig 1A), it is

still of interest to determine if any of the categories that contain genes with strong differences in

ranking between our method and the traditional method show signatures of lineage-specific

selection. We therefore applied the sign test to identify gene sets in which highly ranked genes

(i.e. those with a difference in ranks between the population comparison method and the

traditional method above the cutoff found by GSEAPY) show a systematic bias in directionality.

Three gene sets are significant by the binomial test at an FDR cutoff of 0.1: *Neuronal Cell Body*,

Cytokine Activity and *Cytoskeleton* (Additional file 2, FDR = 0.085 for all three terms). The

Cytokine Activity category is dominated by genes that promote stem cell proliferation which may

have consequences for the proliferation of human and chimpanzee radial glia (although we

caution against overinterpretation of this result given the low fold changes of most genes in this

category; Box 2).

Figure 4 Changes in expression of cytokine activity genes: A) Summary of Cytokine Activity

enrichment. Each blue line represents a gene in the gene set and the green curve is the

cumulative enrichment score. Top ranking genes could be human or chimpanzee-biased in

expression. B) Volcano plot showing expression levels of genes driving Cytokine Activity

enrichment. The $\log_2$ fold-change ($\log_2$ FC, which refers to the value computed by DESeq2) is

chimpanzee-biased for 15 out of 16 genes. Mean $\log_2$ FC between chimpanzee-referenced and

human-referenced $\log_2$ FC values is shown. Negative $\log_2$ FC indicates chimpanzee-biased

expression. EDNRB does not appear because it is a cytokine receptor rather than a cytokine. C)

Comparison of human-chimpanzee EDNRB ASE to within-human ASE from GTEx and fetal

cortical samples. Raw ASE ratios (as opposed to the value derived from DESeq2) are indicated

by "ASE Ratio". The human-chimpanzee ASE is significantly outside of the human ASE

distribution. P-values are from the Mann-Whitney U Test comparing the distribution of human

population ASE to human-chimpanzee ASE. D) Comparison of human-chimpanzee EDN1 ASE

to within-human GTEx ASE. Fetal cortical ASE is not shown due to insufficient data. P-value is

the same as in C. E) ASE of EDNRB across timepoints in cortical spheroids. Expression from

the human allele is consistently higher than expression from the chimpanzee allele. F) ASE of

EDN1 across timepoints in cortical spheroids. Expression from the chimpanzee allele is

consistently higher than expression from the human allele except at D200.

Box 2: The *Cytokine Activity* category is enriched near the top of the difference in ranks list with

15 out of 16 genes driving the enrichment displaying chimpanzee-biased expression (Fig. 4A,

B). It is dominated by genes that generally promote neural stem cell proliferation, which is

unexpected considering the directionality bias and the higher proliferative capacity of human

neural stem cells. *EDN1* had the strongest chimpanzee bias (mean $\log_2$ fold-change = -0.92,

FDR < 0.025 at Day 150, Fig 4B). As *EDN1* primarily signals through *EDNRB* in the brain [41],

we also investigated the expression of *EDNRB*. Surprisingly, *EDNRB* is one of the most strongly

human-biased genes across all timepoints in both hybrid and parental cortical spheroids (mean

$\log_2$ fold-change = 4.46, FDR < 0.005 at Day 150 in hybrids, $\log_2$ fold-change = 2.85, FDR <

0.0005 in parental samples). Human-chimpanzee ASE generally exceeds ASE found in human

populations for *EDNRB* (Fig 4C; Mann Whitney U Test comparing EDNRB distribution to human

population distribution p = 0.00027 for *EDNRB*), although this is not the case for *EDN1* (Fig 4D).

The human-biased *EDNRB* expression and chimpanzee-biased *EDN1* expression is generally

consistent across timepoints in hybrid cortical spheroids (Fig 4E, F).

Changes in *EDNRB* expression appear to be human-derived with respect to gorillas and

macaques (although orangutans may have independently acquired similar expression to

humans in early-stage brain organoids) (Supp. Fig. 3 A, B). We were unable to confidently

determine whether changes in *EDN1* expression were human- or chimpanzee-derived using

currently available data (data not shown). Next, we analyzed single-cell RNA-seq data

generated from 1 month, 2 month, and 4 month-old human and chimpanzee brain organoids to

identify the cell types driving increased EDNRB expression. We found that a previously

identified radial glial cell (RGC) cluster was characterized by high *EDNRB* expression with non-

zero expression in over 50% of cells [42]. Furthermore, this cluster had higher expression than

any chimpanzee cluster (Mann-Whitney U test, $p < 10^{-16}$, Supp. Fig. 4-5). Overall, our results

suggest that a subpopulation of human radial glia have much higher *EDNRB* expression than

chimpanzee radial glia.

EDNRB haploinsufficiency reduces proliferation of cerebellar granule precursor cells and

chemical inhibition of *EDNRB* signaling reduces proliferation of mouse radial glia [43,44]. Based

on this, the change in *EDNRB* expression may have promoted human brain expansion by

increasing the proliferation of the subpopulation of radial glia that express *EDNRB*. In addition, a

recent study identified a population of caudal late interneuron progenitor (CLIP) cells marked by

expression of *EDNRB* and *PTGDS* along with caudal ganglionic eminence markers [45]. As

both *EDNRB* and *PTGDS* have strongly human-biased ASE it would be interesting to

investigate if this population of cells exists in chimpanzee brain organoids and if it may have

expanded in humans. The phenotypic implications of the higher *ENDRB* and *PTGDS* but lower

EDN1 expression in humans will be an exciting area for further research.

Thus far, we have primarily focused on the differences between our new method and the

traditional method for ranking genes. Having established their differences, we now turn to

analysis of results from our new approach (Fig. 1 and Methods). To examine human- and

chimpanzee-biased genes separately, we sorted the list so that highly ranked genes with

human-biased expression are at the top of the list and highly ranked genes with chimpanzee-

biased expression are at the bottom of the list. In effect, this results in a test for directionally-

biased cis-regulatory divergence that exceeds the cis-regulatory variation among most human

alleles present in the GTEx population. Enrichment testing with GSEAPY identified several

enriched gene sets at an FDR cutoff of 0.25 (the cutoff suggested by the GSEA authors)

including *Other glycan degradation* (human-biased), *Spastic Dysarthria* (human-biased), and

Gluconeogenesis (chimpanzee-biased) (Fig 5 A-F, Supp. Table 3). In all three cases, there

were zero genes showing expression bias in the other direction at an identical rank cutoff (p =

0.00012 for *gluconeogenesis*, p = 0.031 for *Other glycan degradation*, and p = 0.00195 for

Spastic Dysarthria by binomial test), suggesting lineage-specific selection on genes with

constrained expression in these gene sets [40]. Notably, the bias in gene expression found in

the hybrids for these genes generally matched the bias in expression found in the parental

cortical spheroids (Supp. Fig. 6A-C).

All three of these enriched categories may influence human-specific phenotypes.

Gluconeogenesis siphons oxaloacetate from the TCA cycle and eventually produces glucose

[46]. Many of the gene expression changes driving the gluconeogenesis enrichment appear to

be human-derived compared to other great apes (Supp. Fig. 7). Decreased gluconeogenesis in

the human lineage would likely enable increased flux through other anabolic pathways and the

TCA cycle possibly increasing the availability of oxaloacetate for anabolic pathways that

promote proliferation. Spastic dysarthria is a condition in which patients speak in a characteristic

slow, regular, monotone manner [47]. Loss of function of genes in this category are associated

with spastic dysarthria and show systematic human bias, which may be connected to the human

capacity for speech (Fig 5D). Finally, six genes that were ranked very highly by our method and

all have DESeq2 FDR < 0.1 drive the "Other glycan degradation" enrichment. Interestingly, loss

of function of three of the six human-biased glycan degradation genes (MANBA, MAN2B2, and

MAN2B1) is associated with intellectual disability [48–50].

Figure 5 Evidence of lineage-specific selection: A) Summary of *Spastic dysarthria*

enrichment (from Human Phenotype Ontology) [33,37]. In A, B, and C, each blue line

represents a gene in the gene set and the green curve is the cumulative enrichment score. **B)**

Summary of *Other glycan degradation* enrichment (from KEGG). **C)** Summary of

Gluconeogenesis enrichment (from REACTOME). **D)** Volcano plot summarizing of Log2 fold-

changes for genes driving the enrichments for *Spastic Dysarthria*, *Other glycan degradation*,

and *Gluconeogenesis*. All genes are human-biased for *Other glycan degradation* and *Spastic*

Dysarthria whereas all genes are chimpanzee-biased for *Gluconeogenesis*.

Discussion

Here we presented a method incorporating population-scale ASE data as a proxy for constraint

on expression. This ranking method helps reveal candidate genes and signatures of selection

that may explain phenotypic differences between humans and chimpanzees. The test is based

on the logic of comparing within- to between-species variation, similar in spirit to the Hudson-

Kreitman-Aguade test although it is based on variation in ASE rather than protein-coding

sequences [51]. Although the phenotypic consequences of these differences remain to be

determined, our finding of polygenic lineage-specific selection on several gene sets suggests

that these changes must have some phenotypic effects in order to be under natural selection.

Collectively, these findings more than double the number of known cases of lineage-specific

polygenic selection on gene expression between humans and chimpanzees (the two previous

examples being Hedgehog signaling and astrocyte-related genes) [8,9].

Importantly, the strong correlation between the GTEx brain vs. non-brain rankings (Fig. 2A) and

GTEx vs. fetal cortex rankings (Fig. 2D) suggests that comparison to the GTEx ASE distribution

can be meaningfully compared with between-species ASE measured in diverse cell types and

organoids and that comparison to the GTEx population distribution will be useful for other cell

types and organoids. The method can also be applied to any species with sufficient gene

expression data, e.g. comparing ASE in *Arabidopsis* interspecies hybrids to ASE within *A.*

thaliana [52].

Conclusion:

We outlined a strategy that uses allele-specific expression data from interspecies hybrids and

population-scale studies to prioritize genes that are more likely to impact species-specific traits

and applied this method to data from human-chimpanzee cortical spheroids. Our findings

provide opportunities for targeted follow-up experiments and increase our understanding of how

polygenic selection has shaped human and chimpanzee evolution. Overall, we anticipate that

our method will become a useful tool for identifying functionally significant gene expression

changes between species, and will contribute to our understanding of how gene expression

drives phenotypic diversification.

Methods:

Read Alignment and RNA-seq Data Processing

Data from hybrid cortical spheroids was mapped as previously described [9]. Briefly, Hornet, a

rewritten version of WASP, was used in conjunction with a curated list of human-chimpanzee

SNPs and indels to correct for mapping bias. Reads for every sample were aligned to both the

human and chimpanzee genomes and the $\log_2$ fold-change from both alignments was

compared. Any genes with $\log_2$ fold-change that differed by greater than 1 were removed. We

used the ASE $\log_2$ fold-change ($\log_2$ FC) values available in the supplemental tables of Agoglia

et al. Although this dataset is restricted to hybrids from two humans and two chimpanzees,

previous work has shown that interspecies differences dominate over differences between

populations within a species so we expect that our results generalize well. It also contains

multiple independently derived hybrid lines and independent differentiations, reducing

confounding by technical differences. Throughout, for hybrids the mean $\log_2$ FC between human

genome mapped and chimpanzee genome mapped reads is stated as well as the highest p-

value. Additional data was downloaded from GSE127898, GSE106245, GSE153076, and

phs000755.v2.p1 and mapped separately for each dataset to the respective species' genome

(PanTro6 for chimpanzee, hg38 for human, mmul10 for rhesus macaque, Gorgor6 for gorilla,

and PonAbe3 for orangutan) [53–56]. We used STAR v2.5.4 with arguments: -outSAMattributes

MD NH -outFilterMultimapNmax 1 -sjdbGTFfile -sjdbOverhang N where N is 1 less than the

read length used for each respective dataset [57]. For paired end reads, we used Picard to

remove duplicates with argument: DUPLICATE_SCORING_STRATEGY = RANDOM [58]. We

used HT-Seq with the following arguments: -t exon -i gene_name -m intersection-strict -r pos to

count reads overlapping gene bodies [59]. Transcripts per million (TPM) was computed as

previously described [60]. We used the likelihood ratio test in DESeq2 to test for differential

expression in the downloaded datasets with RIN and sex included as covariates for the

Khrameeva et al. dataset [23,61]. We binarized RIN values as high if greater than or equal to

7.5 and low otherwise as we do not generally expect the expression level of genes to scale

linearly with RIN.

Comparison of Population and Interspecies ASE Distributions

GTEx data was downloaded from https://www.gtexportal.org/home/datasets and the fetal cortex

ASE data was kindly provided by the Stein laboratory [21,27]. GTEx contains data from 838

individuals and the data from the Stein laboratory was generated from approximately 235

individuals. To preprocess the GTEx data, we split the file into two files each containing read

counts from 1 of the alleles with a custom R script. As cortical spheroids are a mixture of

different cell types including neural progenitors and immature neurons, we pooled fetal cortical

wall, neural progenitor, and neuron counts per individual in the fetal cortex dataset. For each

gene in each sample, we added one count to each gene (to prevent division by zero) and

computed the ASE ratio as the ratio of counts from allele 1 (the reference) to counts from allele

2. The distribution for each gene was then normalized so that the median was 1. This

normalization ensures that the Mann-Whitney U-test p-values only take into account the

variance in allelic expression in the human population and are not confounded by consistently

higher/lower expression from a particular allele. Notably, flipping the sign of each value in the

GTEx ASE distribution had minimal effect on the rankings (Spearman's rho = 0.999, p < 10^{-300}

for day 50, day 100, and day 150 after the beginning of cortical spheroid differentiation)

supporting the efficacy of the correcting the median to 1 in isolating the variance of the human

population expression distribution. All samples with at least 10 counts (not including the single

added count) from each allele for a sample were included in the ASE population distribution.

Notably the rankings and our results are robust to requiring at least 5 counts from each allele

instead of 10 (Spearman's rho = 0.996, p < 10^{-300}). To filter out genes that are lowly expressed

in cortical spheroids, we removed genes with an average number of counts from the

chimpanzee and human alleles less than 25 (i.e. mean of human and chimpanzee read count

less than 25 and mean of chimpanzee read count less than 25). In addition, we filtered out any

genes showing mapping bias (listed in the supplemental tables of Agoglia et al.) as well as

genes on chromosomes 18 and 20 as parts of these chromosomes were duplicated in some

cortical spheroid samples. Previous work has shown that these structural changes have minimal

effect on the computation of ASE values for genes outside the duplicated region [9]. After

filtering, we computed the interspecies ASE distribution in similar manner to the population ASE

distribution (i.e. by taking the ratio of the counts from the human allele to the ratio of the counts

of the chimpanzee allele). However, we did not require 10 counts from each allele and did not

normalize the medians. We did not require 10 counts from each allele because we expect

extreme differences in expression to be relatively common in between species comparisons.

We compared the log2(ASE Ratio) interspecies distribution to the population distribution using

the Mann-Whitney U Test (a nonparametric test robust to the distribution of data) and used the

resulting p-values to rank genes as described below.

Generation of Gene Rankings and Enrichment Analysis

First, we ranked genes by the Mann-Whitney U Test p-value with the lowest p-value receiving

the highest rank. To reduce false positives at the top of the list, we separately ranked genes

with DESeq2 FDR less than or equal to 0.1 and greater than 0.1. We then concatenated the list

so that genes with FDR less than or equal to 0.1 were always ranked higher than genes with

FDR greater than 0.1 (referred to as the MWU ranking). This consensus gene ranking was then

used in GSEAPY preranked with the rankings used as the score that GSEAPY uses to sort the

list. All results highlighted in the text replicated when using 0.05 as a cutoff instead of 0.1. We

used REVIGO in conjunction with a custom python script to generate the plots shown in figure 2

[62].

To compare to the traditional method, we also ranked genes by the DESeq2 derived FDR and

used that in GSEAPY preranked (referred to as the DESeq2 ranking). To highlight differences

between the two methods, we computed the difference in ranks between the two methods by

subtracting the DESeq2 ranking from the MWU ranking and sorting the list on those rankings for

use in GSEAPY. In this context, highly ranked genes are likely those that show relatively mild

gene expression changes but have more constrained expression.

We next performed the expression sign test. First, we generated a list of all gene sets across all

tested ontologies that were nominally enriched at an FDR of 0.25 (using the FDR from GSEAPY

preranked) in at least one time point and that had greater than 10 genes driving the enrichment.

To avoid testing the same gene set multiple times, we only tested each gene set at the

timepoint that had the lowest GSEAPY FDR. We used the average of the DESeq2 $\log_2$ FC from

mapping to the human allele and from mapping to the chimpanzee allele as input for the

binomial test to identify gene sets with significantly more human-biased or chimpanzee-biased

changes than expected by chance. This $\log_2$ FC was generated by comparing the reads from

each species' allele in the cortical spheroid data. For example, if a gene set had 8 human-

biased and 3 chimpanzee-biased genes, then the binomial test was used with $k = 8$, $n = 11$, and

$p = 0.5$. We considered any gene set with Benjamini-Hochberg corrected FDR < 0.1 to be

significant.

Finally, we ranked the genes using a signed version of the MWU ranking. More specifically,

genes were effectively ranked by the $\log_{10}$(MWU p-value) multiplied by the sign of mean

DESeq2 $\log_2$ fold-change so that top ranked genes with negative L2FC are at the bottom of the

list and top ranked genes with positive L2FC are at the top of the list. This ranking was then

used in GSEAPY preranked. All statistical tests (Mann Whitney U Test, Binomial Test,

correlations) were performed in python using the implementation in scipy.

For enrichment testing, we tested gene sets from the Gene Ontology Cellular Component and

Molecular Function categories, the Human Phenotype Ontology, KEGG, and REACTOME using

the same version as in Gokhman et. al [8,33–37]. Regardless of which ranking was used, we

used GSEAPY preranked with the following arguments: processes=4, permutation_num=1000,

seed=6, min_size = 10, max_size = 300 to test for enrichment [28]. Following the authors

suggestion, we considered any category with an FDR below 0.25 to be nominally enriched [28].

We required that ASE data be available from at least 50 individuals in GTEx for a gene to be

included in the ranking. We conducted gene set enrichment analysis with the four different

rankings described above.

Single Cell RNA-seq Data Processing and Analysis

Single cell data from human and chimpanzee organoids and associated metadata were

downloaded from E-MTAB-7552 [42]. We used SCANPY to read in the counts matrix and filter

the data so that only data from 1 month, 2 month, and 4 month old organoids remained [63]. We

used a two-sided Mann Whitney U Test to compare *EDNRB* log2(counts per million) between

the "RGC early 2" cluster and all chimpanzee clusters with and without cells with 0 *EDNRB*

counts included. Mean counts by cell type and tissue for fetal human expression were

downloaded from GSE156793 [64]. We compared cerebrum counts to all other organs except

eye and cerebellum due to their similarity to the cerebrum and used a binomial test to determine

if more tissues exhibited higher expression than brain than expected by chance.

Declarations:

Ethics approval and consent to participate: Access to restricted data was approved by

dbGaP.

Consent for publication: Not relevant to our study.

Availability of data and materials: No new data nor materials were generated during this

study. The datasets supporting the conclusions of this article are available in GSE, dbGaP, or

the supplemental material of Agoglia et al. 2021 (https://www.nature.com/articles/s41586-021-

03343-3#Sec35), GSE127898

(https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE127898), GSE106245

(https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE106245), GSE153076

(https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE153076), and phs000755.v2.p1

(https://www.ncbi.nlm.nih.gov/projects/gap/cgi-bin/study.cgi?study_id=phs000755.v2.p1),

phs002493.v1.p1 (link unavailable, data was received directly from the Stein lab). The GTEx

data was downloaded from https://www.gtexportal.org/home/datasets (although the specific file

that was downloaded is no longer available). The software used to perform the analysis in this

paper is available at: https://github.com/astarr97/Z_Score. There is no restriction on the use of

this code.

Competing interests: The authors have no competing interests to declare.

Funding: Funding for this work came from NIH R01GM097171. A.L.S was supported by an

NDSEG fellowship.

Author contributions: HBF, DG, and ALS conceptualized the study. ALS performed all

analysis, validation, visualization, and writing of software. ALS wrote the original manuscript and

HBF and DG reviewed and edited the manuscript. All authors approved the publication of the

manuscript.

Acknowledgements: We thank Jason Stein and Nil Aygun for providing processed fetal cortex,

neuron, and neural progenitor ASE calls. We thank members of the Fraser lab for helpful

discussion. Illustrations were created with BioRender.

Description of Additional file 1: File name is Additional file 1.csv. Enrichment testing for

difference between traditional and population-based method. This file contains the output from

GSEAPY for enrichment testing as well as the dataset that was used for the enrichment

analysis.

Description of Additional file 2: File name is Additional file 2.csv. Sign test on enriched terms

for difference between traditional and population-based method. This file contains the results for

the sign test applied to a subset (see Methods) of the significant terms from Additional file 1.

Supplementary Figure 1 Relationship between the variance of the GTEx ASE distribution

and the probability of Haploinsufficiency score: A) Shows the relationship between the

variance of the GTEx ASE distribution and the probability of Haploinsufficiency score (a

measure of constraint on gene expression) for each gene tested in this manuscript. As the

probability of Haploinsufficiency increases, the variance of the GTEx distribution decreases. The

spearman correlation is -0.28 with $p < 10^{-170}$.

Supplementary Figure 2 Exploration of changes in CUX1 expression: A) Summary of RNA

polymerase II regulatory region sequence-specific DNA binding. Each blue line represents a

gene in the gene set and the green curve is the cumulative enrichment score. Genes in the

gene set are enriched at the top of the list. **B)** Expression of CUX1 in human-chimpanzee hybrid

cortical spheroids (CS). Expression is shown in transcripts per million (TPM). Expression from

the chimpanzee allele is consistently higher than expression from the human allele. **C)**

Expression of CUX1 in human and chimpanzee parental cortical spheroids (CS). Expression is

shown in transcripts per million (TPM). **D)** Comparison of the GTEx ASE distribution for CUX1 to

the human-chimpanzee hybrid ASE distribution. **E)** Comparison of the fetal cortex ASE

distribution for CUX1 to the human-chimpanzee ASE distribution. The human-chimpanzee

hybrid ASE values lie well outside the human fetal cortex distribution.

Supplementary Figure 3 Changes in EDNRB expression are human derived: A)

Comparison of EDNRB expression between early-stage human and gorilla cerebral organoids

[55]. Human expression is considerably higher than gorilla expression across timepoints

indicating that EDRNB is upregulated in the human lineage (as opposed to downregulated in the

chimpanzee lineage). **B)** Comparison of EDNRB expression between early-stage human,

chimpanzee, orangutan, and rhesus macaque cerebral organoids [53]. Human expression is

considerably higher than rhesus macaque expression across timepoints. However, orangutan

EDNRB expression is high as well, indicating an independent increase in expression in the

orangutan lineage. **C)** Phylogenetic tree of old-world primates. Red text indicates a high *EDNRB*

expression and blue text indicates low expression. Notably, as gorillas are more closely related

to humans than organgutans this implies that the most parsimonious explanation for the data is

that the last common ancestor of gorillas, chimpanzees, and humans had low *EDNRB*

expression and that there was an increase in the human lineage.

Comparison of mean *EDNRB* expression

Human • Human
• Chimp

Supplementary Figure 4 A population of human radial glia expresses EDNRB: A) Plot

showing mean expression of EDNRB across chimpanzee and human cell clusters [42].

Expression in the "RGCs early 2" cluster is significantly higher than in all chimpanzee clusters

by Mann-Whitney U Test (p < 10^{-16} for all comparisons).

Supplementary Figure 5 Expression of EDNRB is higher in human radial glia: The distribution of EDNRB expression in cells with non-zero counts [42]. As there were many more chimpanzee cells in the Chimp NSC/radial glia category, we down-sampled the number of cells so that there were an equal number of EDNRB-expressing cells. The distribution in chimpanzee NSC/radial glia is much more shifted left than that of the human "RGCs Early 2" cluster indicating higher expression in the human cluster.

Supplementary Figure 6 Many gene expression changes have the same direction in

hybrid and parental organoids: A) Barplot showing the $\log_2$ fold-changes of genes driving the

Spastic dysarthria enrichment comparing parental human and chimpanzee cortical spheroids at

day 100 of differentiation. Asterisks indicate a significant difference for that gene at and FDR

cutoff of 0.1. **B)** Same as in A but for genes driving the *Other glycan degradation* enrichment. **C)**

Same as in A, but for genes driving the *Gluconeogenesis* enrichment.

Supplementary Figure 7 Changes in many gluconeogenesis genes are human-derived: A)
Barplot showing that many changes in the expression of gluconeogenesis genes are human-
derived and occur in parental organoids. Data are from Week 5 cerebral organoids [53].
Asterisks indicate genes for which the human-orangutan difference is significant at an FDR
cutoff of 0.1 and with lower expression in human. Genes whose down-regulation is not human-
derived generally show insignificant differences in expression between humans and
chimpanzees potentially indicating compensatory trans-acting genetic changes.

Supplementary Figure 9: A) Explanation of computation of difference in ranks. Population ASE

Ranking was subtracted from the traditional ranking. Due to this, genes that are highly ranked in

both the traditional and population ASE rankings (likely those with low p-values and high

difference in ASE) are near the middle of the list in the difference of ranks. On the other hand,

genes with moderately high p-values and high constraint on expression are ranked lowly in the

traditional ranking, but near the middle in the population ASE ranking, and so are very high in

the difference in ranks.

References

1. Reilly SK, Noonan JP. Evolution of Gene Regulation in Humans. Annual Review of Genomics

and Human Genetics. 2016;17:45–67.

2. Romero IG, Ruvinsky I, Gilad Y. Comparative studies of gene expression and the evolution of

gene regulation. Nature Reviews Genetics. 2012;13:505–16.

3. King M-C, Wilson AC. Evolution at Two Levels in Humans and Chimpanzees. Science

(1979). 1975;188:107–16.

4. Fraser HB. Gene expression drives local adaptation in humans. Genome Research.

2013;23:1089–96.

5. Kelley JL, Gilad Y. Effective study design for comparative functional genomics. Nature

Reviews Genetics. 2020;21:385–6.

6. Housman G, Gilad Y. Prime time for primate functional genomics. Current Opinion in

Genetics & Development. 2020;62:1–7.

7. Zhu Y, Sousa AMM, Gao T, Skarica M, Li M, Santpere G, et al. Spatiotemporal transcriptomic

divergence across human and macaque brain development. Science (1979). 2018;362.

8. Gokhman D, Agoglia RM, Kinnebrew M, Gordon W, Sun D, Bajpai VK, et al. Human–

chimpanzee fused cells reveal cis-regulatory divergence underlying skeletal evolution. Nature

Genetics. 2021;53:467–76.

9. Agoglia RM, Sun D, Birey F, Yoon S-J, Miura Y, Sabatini K, et al. Primate cell fusion

disentangles gene regulatory divergence in neurodevelopment. Nature. 2021;592:421–7.

10. Hu CK, York RA, Metz HC, Bedford NL, Fraser HB, Hoekstra HE. cis-Regulatory changes in

locomotor genes are associated with the evolution of burrowing behavior. Cell Reports.

2022;38:110360.

11. Mack KL, Campbell P, Nachman MW. Gene regulation and speciation in house mice.

Genome Research. 2016;26:451–61.

12. Combs PA, Krupp JJ, Khosla NM, Bua D, Petrov DA, Levine JD, et al. Tissue-Specific cis-

Regulatory Divergence Implicates eloF in Inhibiting Interspecies Mating in Drosophila. Current

Biology. 2018;28:3969-3975.e3.

13. Zhang X, Borevitz JO. Global Analysis of Allele-Specific Expression in *Arabidopsis thaliana*.

Genetics. 2009;182:943–54.

14. Song JHT, Grant RL, Behrens VC, Kučka M, Roberts Kingman GA, Soltys V, et al. Genetic

studies of human–chimpanzee divergence using stem cell fusions. Proceedings of the National

Academy of Sciences. 2021;118.

15. Prud'homme B, Gompel N, Carroll SB. Emerging principles of regulatory evolution.

Proceedings of the National Academy of Sciences. 2007;104:8605–12.

16. Wittkopp PJ, Kalay G. Cis-regulatory elements: molecular mechanisms and evolutionary

processes underlying divergence. Nature Reviews Genetics. 2012;13:59–69.

17. Blekhman R, Oshlack A, Chabot AE, Smyth GK, Gilad Y. Gene Regulation in Primates

Evolves under Tissue-Specific Selection Pressures. PLoS Genetics. 2008;4:e1000271.

18. Gilad Y, Oshlack A, Smyth GK, Speed TP, White KP. Expression profiling in primates

reveals a rapid evolution of human transcription factors. Nature. 2006;440:242–5.

19. Roy S, Wapinski I, Pfiffner J, French C, Socha A, Konieczka J, et al. Arboretum:

Reconstruction and analysis of the evolutionary history of condition-specific transcriptional

modules. Genome Research. 2013;23:1039–50.

20. Rohlfs R v., Nielsen R. Phylogenetic ANOVA: The Expression Variance and Evolution

Model for Quantitative Trait Evolution. Systematic Biology. 2015;64:695–708.

21. Castel SE, Aguet F, Mohammadi P, Aguet F, Anand S, Ardlie KG, et al. A vast resource of

allelic expression data spanning human tissues. Genome Biology. 2020;21:234.

22. Benjamini Y, Hochberg Y. Controlling the False Discovery Rate: A Practical and Powerful

Approach to Multiple Testing. Journal of the Royal Statistical Society: Series B (Methodological).

1995;57:289–300.

23. Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for RNA-

seq data with DESeq2. Genome Biology. 2014;15:550.

24. Collins RL, Glessner JT, Porcu E, Niestroj L-M, Ulirsch J, Kellaris G, et al. A cross-disorder

dosage sensitivity map of the human genome. Genomic Medicine Institute [Internet]. Lerner

Research Institute; 15:23–5. Available from: https://doi.org/10.1101/2021.01.26.21250098

25. Paşca AM, Sloan SA, Clarke LE, Tian Y, Makinson CD, Huber N, et al. Functional cortical

neurons and astrocytes from human pluripotent stem cells in 3D culture. Nature Methods.

2015;12:671–8.

26. Ferraro NM, Strober BJ, Einson J, Abell NS, Aguet F, Barbeira AN, et al. Transcriptomic

signatures across human tissues identify functional rare genetic variation. Science (1979).

2020;369:eaaz5900.

27. Aygün N, Elwell AL, Liang D, Lafferty MJ, Cheek KE, Courtney KP, et al. Brain-trait-

associated variants impact cell-type-specific gene regulation during neurogenesis. The

American Journal of Human Genetics. 2021;108:1647–68.

28. Subramanian A, Tamayo P, Mootha VK, Mukherjee S, Ebert BL, Gillette MA, et al. Gene set

enrichment analysis: A knowledge-based approach for interpreting genome-wide expression

profiles. Proceedings of the National Academy of Sciences. 2005;102:15545–50.

29. Platzer K, Cogné B, Hague J, Marcelis CL, Mitter D, Oberndorff K, et al. Haploinsufficiency

of *CUX1* Causes Nonsyndromic Global Developmental Delay With Possible Catch-up

Development. Annals of Neurology. 2018;84:200–7.

30. Paciorkowski AR, Traylor RN, Rosenfeld JA, Hoover JM, Harris CJ, Winter S, et al. MEF2C

Haploinsufficiency features consistent hyperkinesis, variable epilepsy, and has a role in dorsal

and ventral neuronal developmental pathways. neurogenetics. 2013;14:99–111.

31. Runge K, Mathieu R, Bugeon S, Lafi S, Beurrier C, Sahu S, et al. Disruption of NEUROD2

causes a neurodevelopmental syndrome with autistic features via cell-autonomous defects in

forebrain glutamatergic neurons. Molecular Psychiatry. 2021;26:6125–48.

32. Doan RN, Bae B-I, Cubelos B, Chang C, Hossain AA, Al-Saad S, et al. Mutations in Human

Accelerated Regions Disrupt Cognition and Social Behavior. Cell. 2016;167:341-354.e12.

33. Köhler S, Gargano M, Matentzoglu N, Carmody LC, Lewis-Smith D, Vasilevsky NA, et al.

The Human Phenotype Ontology in 2021. Nucleic Acids Research. 2021;49:D1207–17.

34. Carbon S, Ireland A, Mungall CJ, Shu S, Marshall B, Lewis S. AmiGO: online access to

ontology and annotation data. Bioinformatics. 2009;25:288–9.

35. Jassal B, Matthews L, Viteri G, Gong C, Lorente P, Fabregat A, et al. The reactome

pathway knowledgebase. Nucleic Acids Research. 2019;

36. Kanehisa M, Sato Y, Kawashima M, Furumichi M, Tanabe M. KEGG as a reference

resource for gene and protein annotation. Nucleic Acids Research. 2016;44:D457–62.

37. Köhler S, Doelken SC, Mungall CJ, Bauer S, Firth H v., Bailleul-Forestier I, et al. The

Human Phenotype Ontology project: linking molecular biology and disease through phenotype

data. Nucleic Acids Research. 2014;42:D966–74.

38. Girskis KM, Stergachis AB, DeGennaro EM, Doan RN, Qian X, Johnson MB, et al. Rewiring

of human neurodevelopmental gene regulatory programs by human accelerated regions.

Neuron. 2021;109:3239-3251.e7.

39. Kita R, Venkataram S, Zhou Y, Fraser HB. High-resolution mapping of *cis* -regulatory

variation in budding yeast. Proceedings of the National Academy of Sciences.

2017;114:E10736–44.

40. Fraser HB. Genome-wide approaches to the study of adaptive gene expression evolution.

BioEssays. 2011;33:469–77.

41. Adams KL, Riparini G, Banerjee P, Breur M, Bugiani M, Gallo V. Endothelin-1 signaling

maintains glial progenitor proliferation in the postnatal subventricular zone. Nature

Communications. 2020;11:2138.

42. Kanton S, Boyle MJ, He Z, Santel M, Weigert A, Sanchís-Calleja F, et al. Organoid single-

cell genomic atlas uncovers human-specific features of brain development. Nature.

2019;574:418–22.

43. Vidovic M, Chen M-M, Lu Q-Y, Kalloniatis KF, Martin BM, Tan AHY, et al. Deficiency in

Endothelin Receptor B Reduces Proliferation of Neuronal Progenitors and Increases Apoptosis

in Postnatal Rat Cerebellum. Cellular and Molecular Neurobiology. 2008;28:1129–38.

44. Shinohara H, Udagawa J, Morishita R, Ueda H, Otani H, Semba R, et al. Gi2 Signaling

Enhances Proliferation of Neural Progenitor Cells in the Developing Brain. Journal of Biological

Chemistry. 2004;279:41141–8.

45. Eichmüller OL, Corsini NS, Vértesy Á, Morassut I, Scholl T, Gruber V-E, et al. Amplification

of human interneuron progenitors promotes brain tumors and neurological defects. Science

(1979). 2022;375.

46. Gkini V, Namba T. Glutaminolysis and the Control of Neural Progenitors in Neocortical

Development and Evolution. The Neuroscientist. 2022;107385842110690.

47. Clark HM, Duffy JR, Whitwell JL, Ahlskog JE, Sorenson EJ, Josephs KA. Clinical and

imaging characterization of progressive spastic dysarthria. European Journal of Neurology.

2014;21:368–76.

48. Alkhayat AH, Kraemer SA, Leipprandt JR, Macek M, Kleijer WJ, Friderici KH. Human -

Mannosidase cDNA Characterization and First Identification of a Mutation Associated with

Human -Mannosidosis. Human Molecular Genetics. 1998;7:75–83.

49. Cathey SS, Sarasua SM, Simensen R, Pietris K, Kimbrell G, Sillence D, et al. Intellectual

functioning in alpha-mannosidosis. JIMD Reports. 2019;50:44–9.

50. Blomqvist M, Smeland MF, Lindgren J, Sikora P, Riise Stensland HMF, Asin-Cayuela J. β-

Mannosidosis caused by a novel homozygous intragenic inverted duplication in *MANBA*.

Molecular Case Studies. 2019;5:a003954.

51. Hudson RR, Kreitman M, Aguadé M. A Test of Neutral Molecular Evolution Based on

Nucleotide Data. Genetics. 1987;116:153–9.

52. Zhang H, Zhang F, Yu Y, Feng L, Jia J, Liu B, et al. A Comprehensive Online Database for

Exploring ~20,000 Public Arabidopsis RNA-Seq Libraries. Molecular Plant. 2020;13:1231–3.

53. Field AR, Jacobs FMJ, Fiddes IT, Phillips APR, Reyes-Ortiz AM, LaMontagne E, et al.

Structurally Conserved Primate LncRNAs Are Transiently Expressed during Human Cortical

Differentiation and Influence Cell-Type-Specific Genes. Stem Cell Reports. 2019;12:245–57.

54. Zhu Y, Sousa AMM, Gao T, Skarica M, Li M, Santpere G, et al. Spatiotemporal

transcriptomic divergence across human and macaque brain development. Science (1979).

2018;362.

55. Benito-Kwiecinski S, Giandomenico SL, Sutcliffe M, Riis ES, Freire-Pritchett P, Kelava I, et

al. An early cell shape transition drives evolutionary expansion of the human forebrain. Cell.

2021;184:2084-2102.e19.

56. Khrameeva E, Kurochkin I, Han D, Guijarro P, Kanton S, Santel M, et al. Single-cell-

resolution transcriptome map of human, chimpanzee, bonobo, and macaque brains. Genome

Research. 2020;30:776–89.

57. Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, et al. STAR: ultrafast

universal RNA-seq aligner. Bioinformatics. 2013;29:15–21.

58. Broad Institute. Picard Toolkit. 2019.

59. Givanna H Putri SAPTPJEPFZ. Analysing high-throughput sequencing data in Python with

HTSeq 2.0. ArXiv. 2021;

60. Li B, Ruotti V, Stewart RM, Thomson JA, Dewey CN. RNA-Seq gene expression estimation

with read mapping uncertainty. Bioinformatics. 2010;26:493–500.

61. Zhu A, Ibrahim JG, Love MI. Heavy-tailed prior distributions for sequence count data:

removing the noise and preserving large differences. Bioinformatics. 2019;35:2084–92.

62. Supek F, Bošnjak M, Škunca N, Šmuc T. REVIGO Summarizes and Visualizes Long Lists of

Gene Ontology Terms. PLoS ONE. 2011;6:e21800.

63. Wolf FA, Angerer P, Theis FJ. SCANPY: large-scale single-cell gene expression data

analysis. Genome Biology. 2018;19:15.

64. Cao J, O'Day DR, Pliner HA, Kingsley PD, Deng M, Daza RM, et al. A human cell atlas of

fetal gene expression. Science (1979). 2020;370.

818