Grace-AKO: A Novel and Stable Knockoff Filter for Variable Selection Incorporating Gene Network Structures

Peixin Tian[1], Yiqian Hu[1], Zhonghua Liu[1]*, Yan Dora Zhang[1,2,*]

[1]Department of Statistics and Actuarial Science, The University of Hong Kong, Pokfulam Hong Kong SAR, China.

[2]Centre for PanorOmic Sciences, Li Ka Shing Faculty of Medicine, The University of Hong Kong, Hong Kong SAR, People's Republic of China

Abstract

Motivation: Variable selection is a common statistical approach to detecting genes associated with clinical outcomes of scientific interest. There are thousands of genes in genomic studies, while only a limited number of individual samples are available. Therefore, it is important to develop a method to identify genes associated with outcomes of interest that can control finite-sample false discovery rate (FDR) in high-dimensional data settings.

Results: This article proposes a novel method named Grace-AKO for graph-constrained estimation (Grace), which incorporates aggregation of multiple knockoffs (AKO) with the network-constrained penalty. Grace-AKO can control FDR in finite sample settings and improve model stability simultaneously. Simulation studies show that Grace-AKO has better performance in FDR control in finite sample settings than the original Grace model. We apply Grace-AKO to the prostate cancer data in the The Cancer Genome Atlas (TCGA) program by incorporating prostate-specific antigen (PSA) pathways in the Kyoto Encyclopedia of Genes and Genomes (KEGG) as the prior information. Grace-AKO finally identifies 47 candidate genes associated with PSA level, and more than 75% of the detected genes can be validated.

Availability and implementation: We developed an R package for Grace-AKO available at: https://github.com/mxxptian/GraceAKO

---

*Contributed equally

1 Introduction

Identifying genes and pathways associated with human diseases is a primary focus for advancing the scientific understanding in genomic studies, for which extensive clinical experiments and genetic counseling are required (Katsevich and Sabatti, 2019). A major challenge is that genomic data is usually high-dimensional but with a limited sample size. Currently, there are rich public genomic data, such as the Kyoto Encyclopedia of Genes and Genomes (KEGG, https://www.genome.jp/kegg/), which provides gene-regulatory pathways including biological regulatory relationships between genes or gene products. These gene-regulatory pathways form a network that can be represented as a graphical structure, where the vertices are genes or gene products, and the edges are the gene-regulatory pathways. Inspired by the nature of genomic data, graph-constrained estimation (Grace) provides a new insight on variable selection for genomic data, and accounts for the graphical structures. The predictors in the network are gene-expression data linked by gene-regulatory pathways for a specific clinical outcome, which can also be called graph-structured covariates (Li and Li, 2010). These graphical structures, such as gene-regulatory pathways, can improve the sensitivity of detecting pathways (Rahnenführer et al., 2004). Furthermore, Li and Li (2008) developed a Grace model, network-constrained regularization procedure by encoding the graphical structures into a Laplacian matrix to incorporate this kind of prior biological information into regression analysis. Network-constrained regularization procedure includes the Lasso (Tibshirani, 1996), and the elastic net regulation procedure (Zou and Hastie, 2005) as special cases. According to Katsevich and Sabatti (2019), the optimal policy for this variable selection effort would be to identify significant relevant genes and provide error control for these discoveries (both genes and variants). However, these conventional regression approaches only control false discovery rate (FDR) asymptotically with no guarantee in finite-sample FDR control.

To control the finite-sample FDR in variable selection, fixed-X knockoffs filter and model-X knockoffs filter were proposed as selection procedures (Barber and Candès, 2015; Candes et al., 2018). Model-X knockoffs, as opposed to fixed-X knockoffs, regarded the original variables

as random and relied on the specific stochastic properties of the linear model, thus extending knockoff inference to high-dimensional data (Candes et al., 2018). The salient idea of knockoff inference is to generate knockoff variables by resembling the correlation structure of the original variables without considering the response variable (conditionally on the original variables) (Candes et al., 2018). These knockoff variables are applied to control finite-sample FDR served as negative controls so that the original variables are selected if they are considerably more connected with the response variable than their knockoff variables. However, due to the randomness of model-X knockoffs generation, model-X knockoff performance is unstable. To address this limitation, multiple knockoffs were proposed to allow a more stable finite-sample FDR control and more reproducible results (Emery and Keich, 2019; Gimenez and Zou, 2019; Nguyen et al., 2020). In particular, statistical aggregation is a typical statistical approach to solving instability by aggregating model-X knockoffs inference. Aggregation of multiple knockoffs (AKO) was proposed by Nguyen et al. (2020), which rested on a reformulation of model-X knockoffs to introduce an intermediate feature statistic. It borrowed the idea of Meinshausen et al. (2009) to replicate model-X knockoff procedure multiple times, and then performed statistical aggregation to generate new intermediate feature statistics. Hence, it is more stable than the model-X knockoff procedure.

In this article, we propose a novel method, graph-constrained estimation with aggregation of multiple knockoffs (Grace-AKO). Specifically, the main procedure of Grace-AKO is as follows. First, we generate model-X knockoffs according to the correlation structure of the graph-structured covariates (Candes et al., 2018) and encode graphical structures into a Laplacian matrix (Chung, 1997). Second, we simultaneously fit the graph-structured covariates and their model-X knockoffs into the network-constrained regularization procedure to compute multiple feature statistics: Lasso coefficient-differences (LCDs). Third, we repeat the above procedures multiple times and employ a statistical aggregation approach to transform the LCDs into new intermediate feature statistics, Aggregated Grace Coefficients (AGCs). Fourth, we conduct the Benjamini-Hochberg procedure (Benjamini and Hochberg, 1995) on the multiple AGCs to select the candidate variables. In our simulation studies, we demonstrate that Grace-AKO can accurately control fdr in finite sample settings and has consistent performance, allowing for better repeatability of results. We further investigate a prostate cancer data set from The Can-

cer Genome Atlas (TCGA) program using Grace-AKO, which identifies 47 candidate genes, of which 75% can be validated by literature.

The remainder of this article is organized as follows. In Section 2, we describe the method of Grace-AKO under the linear regression framework. In Section 3, we assess the performance of Grace-AKO using simulation studies. In Section 4, we apply Grace-AKO to a prostate cancer data set from the TCGA program by incorporating KEGG pathways as prior information. In Section 5, we briefly summarize our method.

2 Method

In genomic study, we usually apply a prediction model to identify genes and pathways associated with the trait of interest by linking high-dimensional data (e.g., microarray gene-expression data) to the complex traits. Consider the following linear model where $\boldsymbol{X}$ is a $n \times p$ design matrix with n observations and p predictors, and $\mathbf{y}$ is the response:

$$\mathbf{y} = \mathbf{X}\boldsymbol{\beta} + \boldsymbol{\epsilon}, \tag{1}$$

where design matrix $\boldsymbol{X} = (\boldsymbol{x}_1, \cdots, \boldsymbol{x}_p)$ in which $\mathbf{x}_j = (\mathrm{x}_{1j}, \mathrm{x}_{2j}, \cdots, \mathrm{x}_{nj})$, $j = 1, \cdots, p$ represents a vector of the graph-structured covariate from genomic data (e.g., gene-expression data), and the coefficient $\boldsymbol{\beta}$ represents the contribution of the graph-structured covariates to the trait of interest, and $\boldsymbol{\epsilon}$ is a vector of random errors. We further assume that the response is centred and the predictors are standardized,

$$\sum_{i=1}^{n} \mathrm{y}_i = 0, \quad \sum_{i=1}^{n} \mathrm{x}_{ij} = 0 \quad \text{and} \quad \sum_{i=1}^{n} \mathrm{x}_{ij}^2 = 1, \quad \text{for } j \in [p],$$

where $[p]$ denotes the set including $\{1, 2, \cdots, p\}$. The main goal is to estimate $\boldsymbol{\beta} = (\beta_0, \beta_1, \cdots, \beta_p)$ and select the predictors with nonzero contribution. The regulatory relationships of biological networks for some complex traits, represented as graphical structures between genes or gene products in genomic studies, shed light on underlying biological knowledge, where the covariates are the graph's nodes and the edges indicate functional relationships between two genes. Furthermore, in such a graphical structure, the genes are linked by edges with certain probabilities, where the edge probability is interpreted as a weight in an undirected graph to form a weighted

graph (Li and Li, 2010). To incorporate the prior information about the biological networks, Li and Li (2008) proposed a network-constrained regularization criterion:

$$L(\lambda_1, \lambda_2, \boldsymbol{\beta}) = (\mathbf{y} - \boldsymbol{X}\boldsymbol{\beta})^T(\mathbf{y} - \boldsymbol{X}\boldsymbol{\beta}) + \lambda_1||\boldsymbol{\beta}||_1 + \lambda_2\boldsymbol{\beta}^T\boldsymbol{L}\boldsymbol{\beta}, \tag{2}$$

where $\boldsymbol{L}$ is a non-negative Laplacian matrix of a weighted graph containing biological networks information, and $|| \cdot ||_1$ indicates the L_1 norm, and λ_1 is the Lasso penalty (Tibshirani, 1996), and λ_2 is the penalty for the Laplacian matrix $\boldsymbol{L}$. Specifically, the Laplacian matrix was first introduced by Chung (1997), which included numerous properties of the graph by its consistent set of the eigenvalues or spectrum. When p is large, the model (1) is treated as 'sparse', in which most of coefficients β are zero (Li and Li, 2010). In equation (2), the L_1 norm deals with sparse matrices, and $\boldsymbol{\beta}^T\boldsymbol{L}\boldsymbol{\beta}$ induces a smooth solution of coefficients of the graph-structured covariates. Additionally, $\boldsymbol{L}$ depicts the graphical structure assuming set V including vertices corresponding to the graph-structured covariates and a weighted edge $w(u, v)$, where u and v denote the graph-structured covariates in the graphical structure, and the degree of vertex v is represented as $d_v = \sum_{u \sim v} w(u, v)$. In the genomic data, $w(u, v)$ quantifies the uncertainty of the edge between two vertices, such as the probability of an edge connecting two graph-structured covariates when the graphical structure is constructed from data. Motivated by Chung (1997), we apply the normalized $\boldsymbol{L}$ given in Li and Li (2008):

$$L(u, v) = \begin{cases} 1 - w(u, v)/d_u, & \text{if } u = v \text{ and } d_u \neq 0; \\ -w(u, v)/\sqrt{d_u d_v}, & \text{if } u \text{ and } v \text{ are adjacent}; \\ 0, & \text{otherwise}. \end{cases} \tag{3}$$

Therefore, we can rewrite the second penalty term $\boldsymbol{\beta}^T\boldsymbol{L}\boldsymbol{\beta}$ of equation (2) as follows (Li and Li, 2008):

$$\boldsymbol{\beta}^T\boldsymbol{L}\boldsymbol{\beta} = \sum_{u \sim v}(\frac{\beta_u}{\sqrt{d_u}} - \frac{\beta_v}{\sqrt{d_v}})^2 w(u, v).$$

The network-constrained regularization procedure incorporate the information in the known biological network for variable selection. However, it does not guarantee false discovery rate (FDR) control in finite sample settings. The network-constrained regularization procedure suffers from identifying numerous false positive discoveries when p is large and the number of

samples n is limited. To address this issue, we introduce a kind of multiple knockoffs to control finite-sample FDR. The knockoff filter procedure was first proposed in Barber and Candès (2015), and Candes et al. (2018) further proposed the model-X knockoff filter to extend its application to high-dimensional data. Model-X knockoffs, $\tilde{\boldsymbol{X}}$ are generated from the original data and retain the same data structure as the originals $\boldsymbol{X}$, in which $\boldsymbol{X}$ and $\tilde{\boldsymbol{X}}$ are pairwise exchangeable. We summarize the properties of model-X knockoffs as (Candes et al., 2018):

(1) Swapping the locations of related elements, $\mathbf{x}_j$ and $\tilde{\mathbf{x}}_j$, would not change the joint distribution of $(\boldsymbol{X}, \tilde{\boldsymbol{X}})$ conditional on $\mathbf{y}$.

(2) Once the original covariates $\boldsymbol{X}$ are known, their model-X knockoffs, $\tilde{\boldsymbol{X}}$ provides no extra information on the response variable $\mathbf{y}$.

The knockoffs filter is a cheap method to control finite-sample FDR since it does not require strong assumptions about the design matrix $\boldsymbol{X}$. Due of the random nature of model-X knockoffs sampling, however, the outcome would be unstable and cannot be guaranteed to be reproduced. To increase the stability, multiple knockoffs approaches were developed, and aggregation of multiple knockoffs (AKO) is one of them (Nguyen et al., 2020). In this article, we propose a novel approach, namely Grace-AKO by introducing the statistical aggregation idea of AKO to achieve finite-sample FDR control and stability improvement. Grace-AKO takes the advantages of the network-constrained penalty and AKO to account for biological network information with finite-sample FDR control simultaneously, which contains four major steps.

First, we generate model-X knockoffs, $\tilde{\boldsymbol{X}}$ from the original data matrix $\boldsymbol{X}$ using R package 'knockoff' (Candes et al., 2018). $(\boldsymbol{X}, \tilde{\boldsymbol{X}})$ is further regarded as a new design matrix of the predictors. Based on the properties of model-X knockoffs whose correlation structure is consistent with that of the original variables, we generate a new normalized Laplacian matrix $\boldsymbol{L}$.

Second, the response variable $\boldsymbol{y}$ and the new predictors $(\boldsymbol{X}, \tilde{\boldsymbol{X}})$ are fitted in equation (2). Motivated by Li and Li (2008), a natural solution of Grace-AKO is equivalent to the following optimization problem:

$$\hat{\boldsymbol{\beta}} = \text{argmin}_{\boldsymbol{\beta}}\{\|\mathbf{y} - (\boldsymbol{X}, \tilde{\boldsymbol{X}})(\boldsymbol{\beta}, \tilde{\boldsymbol{\beta}})^T\|_2\}, \tag{4}$$

subject to:

$$(1 - \alpha) \sum_{j=1}^{p} \|\beta_j\|_1 + \alpha \sum_{u \sim v} (\frac{\beta_u}{\sqrt{d_u}} - \frac{\beta_v}{\sqrt{d_v}})^2 w(u, v) \leq t, \tag{5}$$

where $\alpha = \lambda_2/(\lambda_1 + \lambda_2)$, and t is a constant value, and $(\boldsymbol{\beta}, \tilde{\boldsymbol{\beta}})^T$ is a vector of the new coefficients. Furthermore, in the function $(1 - \alpha) \sum_{j=1}^{p} \|\beta_j\|_1 + \alpha \sum_{u \sim v} (\frac{\beta_u}{\sqrt{d_u}} - \frac{\beta_v}{\sqrt{d_v}})^2 w(u, v)$, $\| \cdot \|_1$ deals with sparse data matrix consistent with its function in the network-constrained penalty, and the second term of it penalizes the weighted sum of the squares of the difference of coefficients between the graph-structured covariates, which is scaled by the degree of the associated vertices in the network. Specifically, when two genes are connected, it is expected that their coefficients would be similar rather than identical, which is accomplished by applying the second term of the penalty (Li and Li, 2008).

To identify the relevant variables, we compute the Lasso coefficient-difference (LCD) as the feature statistic to measure the evidence against null hypothesis ($\beta_j = 0$) (Candes et al., 2018):

$$W_j = |\hat{\beta}_j| - |\hat{\beta}_{j+p}|, \tag{6}$$

where $\hat{\beta}_j$, and $\hat{\beta}_{j+p}$ are the estimated coefficients of $\mathbf{x}_j$ and $\tilde{\mathbf{x}}_j$, respectively. Due to the symmetric distribution of W_j under the null, W_j equally takes on positive and negative values (Candes et al., 2018). Moreover, a large positive value of W_j suggests that the distribution of $\mathbf{y}$ is statistically dependent on $\mathbf{x}_j$ and that there is a strong probability that $\mathbf{x}_j$ is the relevant gene linked with the response $\mathbf{y}$.

Third, following Nguyen et al. (2020), Grace-AKO transforms the feature statistic W_j to a new intermediate feature statistic q_j:

$$q_j = \begin{cases} \frac{1 + \#\{k : W_k \leq -W_j\}}{p}, & W_j > 0; \\ 1, & W_j < 0. \end{cases} \tag{7}$$

Then, we repeat the preceding steps B times, including the generation of knockoffs and the calculation of the intermediate feature statistic q_j, to generate a $B \times p$ matrix of the intermediate feature statistics. Then, we provide a new feature statistic, Aggregated Grace Coefficient (AGC), which is derived by applying the quantile aggregation algorithm (Meinshausen et al.,

2009) to the $B \times p$ matrix:

$$\bar{q}_j = \min\{1, \frac{Q_\gamma(\{q_j^{(b)} : b = \{1, 2, \cdots, B\})}{\gamma}\}, \quad (8)$$

where γ is the quantile point, and $Q(\cdot)$ denotes the quantile function, and B is the pre-specified replication times.

Fourth, we apply the multiple testing procedure, the Benjamini-Hochberg procedure (BH) (Benjamini and Hochberg, 1995) to the AGCs, $\bar{q} = (\bar{q}_1, \bar{q}_2, \cdots, \bar{q}_p)$ to compute a data-dependent threshold:

$$t_{BH} = \max\{k : \bar{q}_{(k)} \leq \frac{k}{\alpha p}\},$$

where α is the user-specified nominal FDR level. We finally choose the candidate variables satisfying the following requirement:

$$\hat{S} = \{j \in [p] : \bar{q}_{(j)} \leq \bar{q}_{(\hat{t}_{BH})}\}.$$

In this article, we measure the performance using the modified false discovery rate (mFDR):

$$\text{mFDR} = \mathbb{E}\Big[\frac{|\{j \in \hat{S} \cap S_0\}|}{|\hat{S}| + 1/\alpha}\Big],$$

where $S_0 = \{j \in [p] : \beta_j = 0\}$ contains predictors that have no effect on the trait of interest. The implementation details are available in Algorithm 1.

Algorithm 1: Algorithm of Grace-AKO

Data: $(\boldsymbol{X}, \boldsymbol{Y}, \gamma, q)$: $\boldsymbol{X}$-original data, $\boldsymbol{Y}$-response variable, B-loop times, γ-pre-specified quantiles, q-pre-specified FDR level.

Result: Variable selection set: $\hat{S}$.

while $b \leq B$ **do**

> Draw knockoffs variables $\tilde{\boldsymbol{X}}$;
> Fit $(\boldsymbol{X}, \tilde{\boldsymbol{X}})$ into Grace model;
> Calculate W_j based on (6); Calculate new statistics q_j based on (7).

Do aggregation by quantile function (8) with a matrix of q_j and γ to get AGCs;
Use BH procedure to compute threshold: $t_{BH} = max\{k : \bar{q}_{(k)} \leq \frac{k}{\alpha p}\}$;
Variable selection set: $\hat{S} = \{j \in [p] : \bar{q}_{(j)} \leq \bar{q}_{(t_{BH})}\}$.

As an illustration in Candes et al. (2018), the primary objective of the knockoff filter pro-

cedure was to build an as permissive as possible data-dependent threshold. The threshold can provide a controllable estimation of FDR to ensure model-X knockoffs property. Nguyen et al. (2020) further established an independence or positive correlation between the intermediate feature statistics and the original feature statistics in Candes et al. (2018) to replace the originals. Specifically, AKO still provided a data-dependent threshold based on the specification in Candes et al. (2018) which is also carried on by Grace-AKO.

3   Simulation Studies

In this section, we performed a wide range of simulation studies to evaluate our proposed method, Grace-AKO, and compared it to the network-constrained regularization procedure (namely Grace) (Li and Li, 2008). We supposed that there were 10 transcription factors (TFs) and each regulates 10 genes. The graphical structure included g unconnected regulatory modules with p genes in total and edges linked each of the TFs and 10 genes regulated. Here we denoted the first 44 genes related to the response variable $\mathbf{y}$. We assumed that the data were simulated from the following settings:

(1) $\mathbf{y} = \mathbf{X}\boldsymbol{\beta} + \boldsymbol{\epsilon}$, where
$$\boldsymbol{\beta} = (5, \overbrace{\frac{5}{\sqrt{10}}, \cdots, \frac{5}{\sqrt{10}}}^{10}, -5, \overbrace{\frac{-5}{\sqrt{10}}, \cdots, \frac{-5}{\sqrt{10}}}^{10}, 3, \overbrace{\frac{3}{\sqrt{10}}, \cdots, \frac{3}{\sqrt{10}}}^{10}, -3, \overbrace{\frac{-3}{\sqrt{10}}, \cdots, \frac{-3}{\sqrt{10}}}^{10}, 0, \cdots 0),$$
and $\boldsymbol{\epsilon}$ is generated from $N(0, \boldsymbol{\sigma}^2)$;

(2) The noise level is denoted as $\sigma^2 = \sum_u^p \beta_u^2 / 4$;

(3) For each expression level TF, X is drawn from normal distribution: $\boldsymbol{X}_g \sim N(0, 1)$, and conditional on the TF, the expression levels of genes which regulated to the specific TF are drawn from a conditional normal distribution with correlation of 0.7.

We set $n = (100, 200, 300)$, $g = (10, 20, 40, 60)$ and $p = (110, 220, 440, 660)$, where g represents the total number of unconnected regulatory modules, to simulate 12 settings in total. As the empirical research in Nguyen et al. (2020) demonstrated, performance was consistent and robust once the iteration B approached 25 times. In the meanwhile, FDR could almost be controlled under the pre-specified level. Therefore, we fixed $B = 25$, and $\gamma = 0.1$ in all scenarios. Additionally, we specified the FDR control level α at 0.1. According to Li and Li (2008), Grace

performed better than the Lasso (Tibshirani, 1996) and the elastic net (Zou and Hastie, 2005) in term of the combination of $\|\cdot\|_1$ and $\|\cdot\|_2$ and the Laplacian matrix $\boldsymbol{L}$. We thus focused on the comparison of Grace-AKO and Grace in our simulation studies. We assessed the regression performance using the average values of mFDR, standard errors (calculated by Monte Carlo method), and the number of selected variables.

To tune the parameters, we applied 10-fold cross-validation. The values of tuning parameter $\lambda_{\boldsymbol{L}}$ were specified from the range of 0.1 to 2.0 with a step size of 0.1, which ensured that the $\boldsymbol{L}$ matrix was non-negative. Moreover, the values of λ_1 were drawn from the range of 110 to 200 with a step size of 5. The validation set of λ_2 ranged from 1 to 10 with a step size of 1. Each simulation setting was repeated for 30 times.

As Table 1 showed, the mFDR values of Grace-AKO were controlled at the pre-specified level, and Grace could not control mFDR in finite sample settings. Furthermore, as the data dimension p increased with a fixed sample size n, the mFDR for Grace increased which even reached at 0.67. However, Grace-AKO could always control the mFDR under 0.1. Moreover, the standard errors for Grac-AKO were alway smaller than Grace's, which indicated that Grace-AKO could also improve the stability. In Figure 1, we depicted a scatter plot for the findings when $n = 300$. The upper figure's black line represented the pre-specified level. Grace, in particular, had inflated power at cost of high mFDR.

Table 1: The mean and standard errors (in brackets) of modified false discovery rate (mFDR) over 30 replications for Grace-AKO v.s. Grace model with a pre-specified mFDR level at 0.1.

p	$n = 100$		$n = 200$		$n = 300$	
	Grace-AKO	**Grace**	**Grace-AKO**	**Grace**	**Grace-AKO**	**Grace**
10	0.01 (0.018)	0.15 (0.032)	0.03 (0.032)	0.22 (0.049)	0.03 (0.025)	0.26 (0.053)
20	0.02 (0.027)	0.26 (0.045)	0.02 (0.023)	0.37 (0.045)	0.03 (0.040)	0.43 (0.010)
40	0.03 (0.030)	0.24 (0.062)	0.04 (0.030)	0.48 (0.040)	0.06 (0.047)	0.61 (0.030)
60	0.04 (0.026)	0.40 (0.053)	0.03 (0.026)	0.55 (0.032)	0.06 (0.034)	0.67 (0.018)

Figure 1: Figures of mFDR and Power for Grace-AKO v.s Grace model when $n = 300$. The upper figure plots mFDR and lower figure plots power. The red circle represents our new method, Grace-AKO and the blue one represents Grace. Each setting has been replicated for 30 times, and each dot represents one trail. The black dashed line is the pre-specified mFDR level which equals to 0.1.

As shown in Table 2, the number of selected variables varied significantly between these two models. The number of true variables was fixed at 44. When n and p increased, the number of variables selected by Grace increased. Notably, when $n = 300$ and $p = 60$, Grace identified 138 candidate variables when the number of true variables was 44. It demonstrated that Grace could identify many false positive variables in finite sample settings. Grace-AKO had relatively more stable performance under all the data settings, which could correctly capture the true candidate genes. This is the reason why, in Figure 1, Grace's power was inflated while Grace-AKO performed relatively more conservatively.

Table 2: The average numbers of variables selected for Grace-AKO v.s. Grace over 30 replications with a pre-specified mFDR level at 0.1. The number of true variables is 44.

p	$n = 100$		$n = 200$		$n = 300$	
	Grace-AKO	**Grace**	**Grace-AKO**	**Grace**	**Grace-AKO**	**Grace**
10	24	45	33	56	37	62
20	25	52	32	71	37	83
40	24	59	34	90	39	118
60	24	63	32	101	39	138

4 Application to the TCGA Prostate Cancer Data

To demonstrate the usefulness of Grace-AKO, we applied it to a gene-expression data of prostate-specific antigen (PSA) level from the The Cancer Genome Atlas (TCGA) program. The TCGA program is a landmark cancer genomics program with over 20,000 primary cancers and standard samples spanning 33 cancer types (Koboldt et al., 2012). Additionally, the Kyoto Encyclopedia of Genes and Genomes (KEGG), as a public database, has enabled a profound insight into the graphical structures of genes and various clinical outcomes (Kanehisa and Goto, 2000). It contributes to understanding various aspects of biological systems and pathway information. Prostate cancer is the most common malignancy in mid-aged males and the second leading malignancy (Stangelberger et al., 2008). This external graphical structure is represented by a penalty weight matrix, which is the Laplacian matrix $\boldsymbol{L}$ constructed in equation (3). Wang et al. (2018) indicated that metastatic prostate cancer remained incurable even in patients who finished intensive multimodal therapy. It is an urgent challenge to propose a novel approach for disease management via identifying prognostic determinant genes. Moreover, Reiner et al. (2003) indicated that the statistical significance of differential expression might require abundant experiments, and the probability of type I error increased as the multiple hypotheses were tested. Katsevich and Sabatti (2019) shown that identifying the significant variables that might promote replicability of the results while dealing with a comprehensive collection of possible explanatory variables. In this article, we thus were more concerned with the accuracy of gene selection in tumor investigations (assessed by mFDR) than with the sensitivity of detection (assessed by power).

We first removed the samples with missing measurements and then encoded PSA pathways (Kim et al., 2018) from the KEGG to construct the normalized Laplacian matrix for Grace

and Grace-AKO. We finally obtained the data with sample size $n = 339$ and dimension $p = 5,947$. We denoted the PSA level as our response. To reduce computational demands, we first performed variable screening via correlation learning following Fan and Lv (2008). The final sample size and dimension were $n = 339$ and $p = 600$, respectively, consisting of the data structure in simulation studies. We then standardized the explanatory variables and response variables. The parameter λ_1 ranges from 1 to 40 length out as 4, and λ_2 is among 1, 2, 3, 4, and 5. We denote the target FDR level at 0.2 and $B = 25$ for the AKO procedure. The quantile point is $\gamma = 0.3$. For a fair comparison, we both conducted 10-fold cross-validation to select the tuning parameters. We showed the details in Figure 2.

There were 90 genes selected by Grace in total, and 47 genes selected by Grace-AKO. Li and Li (2010) indicated that Grace might lose some accuracy when the coefficients' signs are different. After checking the intersection of variable selection sets, we found that the genes selected by Grace-AKO were a subset of Grace's. However, due to the inflated FDR in the simulation studies, the results of Grace might identify some false discoveries. Moreover, previous studies confirmed 35 out of 47 candidate genes for Grace-AKO as true associated genes. Some details of them were listed in Table 3. For example, IL9 was recently assigned as an essential gene in tumor immunity, and AK6 was already regarded as a biomarker in prostate cancer treatments (Liu et al., 2015; Wan et al., 2020). HLA-DRB5 was highly related to MHC-II genes, which were the related genes of prostate cancer, and so was the CHRNB2 (Axelrod et al., 2019). Moreover, a high concentration of IFNA2 was related to advanced prostate carcinoma (Erb et al., 2013). PLK1 was over-expressed in many cancers, including prostate cancer, and scientists found that translation of the PP2A-PLK1 SDL interaction caused the expression of PLK1 and PP2A, which were commonly regarded as a biomarker in the cancer cells (Cunningham et al., 2016). Melloy (2019) indicated the percentage of the case with alteration of ANAPC7 achieved beyond 5.21 percentage points. The results demonstrated that the APC/C had a profound effect on cancer survival. MYC in 2010 had already been confirmed to be affected by the loss of the tumor in Koh et al. (2010).

Furthermore, there were also some subnetworks in our findings. H2BC9, H2AC14, and H2AC7 consisted of a small subnetwork. Moreover, IFNA7, IL9, IL5RA, IFNA2, PDGFB, and KIT comprised a subnetwork of validated pathways. Except for KIT, the remaining genes

Figure 2: Flowchart of Grace-AKO. These two procedures are differentiated by colors, in which the green one is Grace and the blue one is Grace-AKO. First, we encode the Laplacian matrix account for the network structure and sample model-X knockoffs. Second, we compute the feature statistics LCDs. Third, we aggregate the LCDs to get AGCs. Fourth, we select the variables whose AGCs satisfy the requirements.

were investigated as possible prostate cancer therapy genes. Additionally, there was a pathway between CAMK4 and PRKACF, where CAMKK2 was a significant androgen receptor target for prostate cancer tumor growth, according to Dadwal et al. (2018). MYC was linked with RXRA. In Ray et al. (2020), RXRA, which was discovered as a novel target of miR-191, was

conserved in a cell line derived from radio recurrent prostate cancer.

Table 3: Genes Selected by Grace-AKO in the application of prostate cancer and PSA pathways.

Genes Name	CHR	Genes Expression
ANAPC7	12	Ubiquitous expression in testis (RPKM 11.7), esophagus (RPKM 9.0) and 25 other tissues
AK6	5	Ubiquitous expression in testis (RPKM 18.6), adrenal (RPKM 17.2) and 25 other tissues
CHRNB2	1	Biased expression in brain (RPKM 9.5), adrenal (RPKM 0.7) and 1 other tissue
COL4A5	X	Broad expression in endometrium (RPKM 21.7), skin (RPKM 10.8) and 20 other tissues
PRLR	5	Biased expression in placenta (RPKM 18.9), endometrium (RPKM 11.6) and 8 other tissues
CAMK4	5	Broad expression in brain (RPKM 7.0), lymph node (RPKM 3.7) and 21 other tissues
CHRNB2	1	Biased expression in brain (RPKM 9.5), adrenal (RPKM 0.7) and 1 other tissue
PPP2R2D	10	Ubiquitous expression in pancreas (RPKM 6.3), ovary (RPKM 5.9) and 25 other tissues
IFNA13	9	-
COL1A1	17	Biased expression in gall bladder (RPKM 850.7), urinary bladder (RPKM 497.1) and 11 other tissues
RBPJ	4	Ubiquitous expression in placenta (RPKM 17.3), endometrium (RPKM 16.0) and 25 other tissues
PARD6G	18	Broad expression in skin (RPKM 15.6), esophagus (RPKM 5.7) and 15 other tissues
ZNF766	19	Ubiquitous expression in testis (RPKM 7.0), thyroid (RPKM 4.6) and 25 other tissues
PRKACG	9	-
SPDYE4	17	Restricted expression toward testis (RPKM 1.5)
PRKACA	19	Ubiquitous expression in heart (RPKM 62.0), adrenal (RPKM 43.9) and 25 other tissues
HLA-DRB5	6	Broad expression in lung (RPKM 275.5), lymph node (RPKM 163.3) and 14 other tissues
ZNF671	19	Ubiquitous expression in spleen (RPKM 4.7), lymph node (RPKM 4.6) and 25 other tissues
ZNF492	19	Broad expression in testis (RPKM 1.4), bone marrow (RPKM 0.9) and 16 other tissues
MAPK3	16	Ubiquitous expression in small intestine (RPKM 45.8), colon (RPKM 42.8) and 25 other tissues
ZNF461	19	Ubiquitous expression in testis (RPKM 1.8), thyroid (RPKM 1.6) and 25 other tissues
CCNE1	19	Biased expression in placenta (RPKM 16.5), bone marrow (RPKM 11.3) and 12 other tissues
CCR3	3	-
ZNF251	8	Ubiquitous expression in endometrium (RPKM 6.9), thyroid (RPKM 6.8) and 25 other tissues
CSH1	17	Restricted expression toward placenta (RPKM 9553.7)
ACTG1	17	Ubiquitous expression in ovary (RPKM 1227.2), esophagus (RPKM 970.4) and 25 other tissues
TH	11	Restricted expression toward adrenal (RPKM 42.8)
ANF853	-	-
CDC27	17	Ubiquitous expression in thyroid (RPKM 20.3), testis (RPKM 11.6) and 25 other tissues
H2AC14	6	-
H2AC7	6	-
IL9	5	-
IL5RA	3	Biased expression in lung (RPKM 1.2), prostate (RPKM 0.7) and 13 other tissues
IFNA7	9	-
IFNA2	9	-
FARR2	-	-
ZNF248	10	Ubiquitous expression in endometrium (RPKM 2.8), thyroid (RPKM 2.7) and 24 other tissues
IFNA17	9	-
ZNF331	19	Broad expression in adrenal (RPKM 22.0), placenta (RPKM 9.5) and 20 other tissues
ZNF473	19	Broad expression in testis (RPKM 14.8), spleen (RPKM 2.3) and 21 other tissues
ZNF713	7	Broad expression in brain (RPKM 2.1), testis (RPKM 1.9) and 25 other tissues
COL9A1	6	Biased expression in prostate (RPKM 6.1), testis (RPKM 1.3) and 6 other tissues
H2BC9	6	-
PDGFB	22	Broad expression in placenta (RPKM 18.6), fat (RPKM 12.7) and 23 other tissues
MYC	8	Ubiquitous expression in gall bladder (RPKM 49.6), esophagus (RPKM 44.6) and 25 other tissues

Note: CHR stands for chromosome.

5 Conclusions

This article introduces Grace-AKO to perform the variable selection by incorporating the network-constrained penalty (Li and Li, 2008) and AKO (Nguyen et al., 2020). In contrast to the conventional variable selection process, Grace-AKO applies a normalized Laplacian matrix to encode the graphical structures between the potential genes or the gene products. It applies the L_1 penalty to selected variables and the L_2 penalty to degree-scaled differences of coefficients concerning the graphical structure. Moreover, our proposed Grace-AKO guarantees of FDR control in finite sample settings by identifying variable employing multiple knockoff variables. In finite sample settings, Grace-AKO solves the instability of model-X knockoffs by incorporating an aggregation process and proposing a new feature statistics AGC. We assessed the performance under a wide range of simulation settings, and the simulation studies demonstrated that Grace-AKO had better performance in finite-sample FDR control. Furthermore, the proposed general framework for variable selection with finite-sample FDR control can be broadly extended to existing penalties (e.g., the Lasso penalty and the elastic net penalty).

References

Axelrod, M. L., Cook, R. S., Johnson, D. B., and Balko, J. M. (2019). Biological consequences of MHC-II expression by tumor cells in cancer. *Clin Cancer Res*, 25(8):2392–2402.

Barber, R. F. and Candès, E. J. (2015). Controlling the false discovery rate via knockoffs. *The Annals of Statistics*, 43(5):2055–2085.

Benjamini, Y. and Hochberg, Y. (1995). Controlling the false discovery rate: a practical and powerful approach to multiple testing. *Journal of the Royal statistical society: series B (Methodological)*, 57(1):289–300.

Candes, E., Fan, Y., Janson, L., and Lv, J. (2018). Panning for gold:'model-x'knockoffs for high dimensional controlled variable selection. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, 80(3):551–577.

Chung, F. R. K. (1997). *Spectral Graph Theory*. American Mathematical Society.

Cunningham, C. E., Li, S., Vizeacoumar, F. S., Bhanumathy, K. K., Lee, J. S., Parameswaran, S., Furber, L., Abuhussein, O., Paul, J. M., McDonald, M., Templeton, S. D., Shukla, H., El Zawily, A. M., Boyd, F., Alli, N., Mousseau, D. D., Geyer, R., Bonham, K., Anderson, D. H., Yan, J., Yu-Lee, L.-Y., Weaver, B. A., Uppalapati, M., Ruppin, E., Sablina, A., Freywald, A., and Vizeacoumar, F. J. (2016). Therapeutic relevance of the protein phosphatase 2a in cancer. *Oncotarget*, 7(38):61544–61561.

Dadwal, U. C., Chang, E. S., and Sankar, U. (2018). Androgen receptor-CaMKK2 axis in prostate cancer and bone microenvironment. *Front Endocrinol (Lausanne)*, 9:335.

Emery, K. and Keich, U. (2019). Controlling the FDR in variable selection via multiple knockoffs. *arXiv e-prints*, page arXiv:1911.09442.

Erb, H. H., Langlechner, R. V., Moser, P. L., Handle, F., Casneuf, T., Verstraeten, K., Schlick, B., Schäfer, G., Hall, B., Sasser, K., et al. (2013). Il6 sensitizes prostate cancer to the antiproliferative effect of ifnα2 through irf9. *Endocrine-related cancer*, 20(5):677.

Fan, J. and Lv, J. (2008). Sure independence screening for ultrahigh dimensional feature space. *Journal of The Royal Statistical Society Series B-statistical Methodology*, 70(5):849–911.

Gimenez, J. R. and Zou, J. (2019). Improving the stability of the knockoff procedure: Multiple simultaneous knockoffs and entropy maximization. *Proceedings of the Twenty-Second International Conference on Artificial Intelligence and Statistics*, 89:2184–2192.

Kanehisa, M. and Goto, S. (2000). KEGG: kyoto encyclopedia of genes and genomes. *Nucleic Acids Res*, 28(1):27–30.

Katsevich, E. and Sabatti, C. (2019). Multilayer knockoff filter: Controlled variable selection at multiple resolutions. *The annals of applied statistics*, 13(1):1.

Kim, I., Choi, S., and Kim, S. (2018). BRCA-pathway: a structural integration and visualization system of TCGA breast cancer data on KEGG pathways. *BMC Bioinformatics*, 19(1):42.

Koboldt, D., Fulton, R., McLellan, M., Schmidt, H., Kalicki-Veizer, J., McMichael, J., Fulton, L., Dooling, D., Ding, L., Mardis, E., et al. (2012). Comprehensive molecular portraits of human breast tumours. *Nature*, 490(7418):61–70.

Koh, C. M., Bieberich, C. J., Dang, C. V., Nelson, W. G., Yegnasubramanian, S., and De Marzo, A. M. (2010). MYC and prostate cancer. *Genes Cancer*, 1(6):617–628.

Li, C. and Li, H. (2008). Network-constrained regularization and variable selection for analysis of genomic data. *Bioinformatics*, 24(9):1175–82.

Li, C. and Li, H. (2010). Variable selection and regression analysis for graph-structured covariates with an application to genomics. *The annals of applied statistics*, 4(3):1498–1516.

Liu, C., Zhang, L., Huang, Y., Lu, K., Tao, T., Chen, S., Zhang, X., Guan, H., Chen, M., and Xu, B. (2015). MicroRNA-328 directly targets p21-activated protein kinase 6 inhibiting prostate cancer proliferation and enhancing docetaxel sensitivity. *Mol Med Rep*, 12(5):7389–7395.

Meinshausen, N., Meier, L., and Bühlmann, P. (2009). P-values for high-dimensional regression. *Journal of the American Statistical Association*, 104(488):1671–1681.

Melloy, P. (2019). The anaphase-promoting complex: a key mitotic regulator associated with somatic mutations occurring in cancer. *Genes, Chromosomes and Cancer*, 59(3):189–202.

Nguyen, T.-B., Chevalier, J.-A., Thirion, B., and Arlot, S. (2020). Aggregation of multiple knockoffs. In *International Conference on Machine Learning*, pages 7283–7293. PMLR.

Rahnenführer, J., Domingues, F. S., Maydt, J., and Lengauer, T. (2004). Calculating the statistical significance of changes in pathway activity from gene expression data. *Statistical applications in genetics and molecular biology*, 3(1).

Ray, H., a. . J. R., Haughey, C., Hoey, C., Jeon, J., Murphy, R., Dura-Perez, L., McCabe, N., Downes, M., Jain, S., Boutros, P. C., Mills, I. G., and Liu, S. K. (2020). miR-191 promotes radiation resistance of prostate cancer through interaction with RXRA. *Cancer Letters*, 473:107–117.

Reiner, A., Yekutieli, D., and Benjamini, Y. (2003). Identifying differentially expressed genes using false discovery rate controlling procedures. *Bioinformatics*, 19(3):368–375.

Stangelberger, A., Waldert, M., and Djavan, B. (2008). Prostate cancer in elderly men. *Rev Urol*, 10(2):111–119.

Tibshirani, R. (1996). Regression shrinkage and selection via the Lasso. *Journal of the Royal Statistical Society. Series B (Methodological)*, 58(1):267–288.

Wan, J., Wu, Y., Ji, X., Huang, L., Cai, W., Su, Z., Wang, S., and Xu, H. (2020). IL-9 and IL-9-producing cells in tumor immunity. *Cell Commun Signal*, 18(1):50.

Wang, G., Zhao, D., Spring, D. J., and DePinho, R. A. (2018). Genetics and biology of prostate cancer. *Genes Dev*, 32(17-18):1105–1140.

Zou, H. and Hastie, T. (2005). Regularization and variable selection via the elastic net. *Journal of the Royal Statistical Society. Series B (Statistical Methodology)*, 67(2):301–320.