

Cross-neutralization of Omicron BA.1 against BA.2 and BA.3 SARS-CoV-2

Jing Zou¹, Chaitanya Kurhade¹, Hongjie Xia¹, Mingru Liu², Xuping Xie^{1,#}, Ping Ren^{2,#},
Pei-Yong Shi^{1,3,4,5,6,#},

¹Department of Biochemistry and Molecular Biology, University of Texas Medical Branch,
Galveston TX, U.S.A.

²Department of Pathology, University of Texas Medical Branch, Galveston TX, U.S.A.

³Sealy Institute for Drug Discovery, University of Texas Medical Branch, Galveston, TX,
U.S.A.

⁴Institute for Human Infection and Immunity, University of Texas Medical Branch, Gal-
veston, TX, U.S.A.

⁵Institute for Translational Sciences, University of Texas Medical Branch, Galveston, TX,
U.S.A.

⁶Sealy Institute for Vaccine Sciences, University of Texas Medical Branch, Galveston,
TX, U.S.A.

#Correspondence: X.X. (xuxie@utmb.edu), P.R. (piren@utmb.edu) or P.-Y.S.
(peshi@UTMB.edu)

Abstract

The Omicron SARS-CoV-2 has three distinct sublineages, among which sublineage BA.1 is responsible for the initial Omicron surge and is now being replaced by BA.2 world-wide, whereas BA.3 is currently at a low frequency. The ongoing BA.1-to-BA.2 replacement underscores the importance to understand the cross-neutralization among the three Omicron sublineages. Here we tested the neutralization of BA.1-infected human sera against BA.2, BA.3, and USA/WA1-2020 (a strain isolated in late January 2020). The BA.1-infected sera neutralized BA.1, BA.2, BA.3, and USA/WA1-2020 SARS-CoV-2s with geometric mean titers (GMTs) of 445, 107, 102, and 16, respectively. Thus, the neutralizing GMTs against heterologous BA.2, BA.3, and USA/WA1-2020 were 4.2-, 4.4-, and 28.4-fold lower than the GMT against homologous BA.1, respectively. These findings have implications in COVID-19 vaccine strategy.

Main text

Since the emergence of severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) in late 2019, the virus has evolved to increase viral transmission and immune evasion. The World Health Organization (WHO) has so far designated 5 variants of concern (VOC), including Alpha, Beta, Gamma, Delta, and Omicron. The most recently emerged Omicron variant has 3 distinct sublineages: BA.1, BA.2, and BA.3. BA.1 was first identified in South Africa in November 2021. BA.1 and its derivative BA.1.1 (containing an extra R346K substitution in the spike of BA.1) caused the initial surges of Omicron around the world. Subsequently, the frequency of BA.2 increased steeply, replacing BA.1 in many parts of the world. In the USA, the frequency of BA.2 increased from 0.4% to 54.9% between January 22 and March 24, 2022 (<https://covid.cdc.gov/covid-data-tracker/#variant-proportions>). Compared with BA.1, BA.2

did not seem to cause more severe disease¹, but may increase viral transmissibility by ~30%². As of March 30, 2022, the frequency BA.3 remained low in the GISAID database (<https://www.gisaid.org/>). All three sublineages of Omicron could significantly evade vaccine-elicited neutralization, among which BA.3 exhibited the greatest reduction^{3,4}. In addition, Omicron BA.1 could efficiently evade non-Omicron SARS-CoV-2 infection-elicited neutralization⁵. The increased transmissibility and immune evasion of the Omicron variant may be responsible for the replacement of VOC from the previous Delta to the current Omicron. Since many unvaccinated individuals were infected by BA.1 during the initial Omicron surge, it is important to examine the cross-neutralization of BA.1 infection against BA.2, BA.3, and other variants. Such laboratory information is essential to guide vaccine strategy and public health policy.

To examine the cross-neutralization among the three Omicron sublineages, we collected 20 human sera from unvaccinated patients who were infected with Omicron BA.1 (**Table 1**). The genotype of infecting virus was verified for each patient by Sanger sequencing. The sera were collected on day 8 to 62 after positive RT-PCR test. The serum panel was measured for neutralization against four recombinant SARS-CoV-2s (**Figure 1A**): USA/WA1-2020 (wild-type) and three chimeric USA/WA1-2020 bearing the full-length spike protein from Omicron BA.1 (GISAID EPI_ISL_6640916), BA.2 (GISAID EPI_ISL_6795834.2), or BA.3 (GISAID EPI_ISL_7605591). The spike proteins of the three Omicron sublineages have distinct amino acid mutations, deletions, and insertions (**Figure 1A**). To facilitate neutralization testing, an mNeonGreen (mNG) reporter was engineered into the four viruses, resulting in wild-type, BA.1-, BA.2-, and BA.3-spike mNG SARS-CoV-2s. The construction and characterization of the four mNG SARS-CoV-2s were recently reported⁴. Using an mNG-based fluorescent focus-reduction neutralization test (FFRNT), we determined the neutralizing geometric mean titers (GMTs) of the sera against

wild-type, BA.1-, BA.2-, and BA.3-spike mNG SARS-CoV-2s to be 445, 107, 102, and 16, respectively (**Fig. 1B**). Thus, the neutralizing GMTs against heterologous BA.2-spike, BA.3-spike, and wild-type viruses were 4.2-, 4.4-, and 28.4-fold lower than the GMT against the homologous BA.1-spike virus, respectively (**Fig. 1B**). Consistently, all sera neutralized BA.1-spike virus at neutralizing titers of ≥ 80 , whereas 12 out of 20 sera did not neutralize the wild-type USA/WA1-2020 (defined as 10 for plot and calculation purposes; **Fig. 1B** and **Table 1**). Notably, 2 sera neutralized BA.2-spike virus more efficiently than the BA.1-spike virus (indicated by symbol * in **Fig. 1C**). Collectively, the results support two conclusions. First, BA.1 infection elicited similar levels of cross-neutralization against BA.2 and BA.3, although at a decreased efficiency that was 4.2- to 4.4-fold lower than that against BA.1. This result is in contrast with the neutralization results from vaccinated sera (collected at 1 month after three doses of Pfizer/BioNTech's BNT162b2 vaccine) which neutralized BA.1 and BA.2 much more efficiently than BA.3⁴. Second, the neutralization of BA.1-infected sera against USA/WA1-2020 were 6.7- and 6.4-fold lower than that against Omicron BA.2 and BA.3, respectively. The results indicate the antigenic distinctions among different variant spikes, which must be carefully considered when deciding to switch the vaccine sequence to new variants⁶. If future variants are Omicron decedents, switch vaccine sequence to an Omicron spike is conceptually attractive.

Emerging evidence supports a vaccine booster strategy to minimize the health risk of the ongoing Omicron surge. First, 2 doses of BNT162b2 vaccine are inefficient to elicit robust neutralization against Omicron variant, whereas 3 doses of BNT162b2 produces robust neutralization against Omicron. Although Omicron-neutralizing activity remains robust for up to 4 months³, the durability of such neutralization beyond 4 months after dose 3 remains to be determined. The latter result, together with the real-world vaccine effectiveness, are required to guide the timing of dose 4 vaccine. Second, non-Omicron

SARS-CoV-2 infection does not elicit robust neutralization against Omicron variant⁵, suggesting that previously infected individuals should be vaccinated to mitigate the health threat from Omicron. The cross-neutralization of BA.1-infected sera against BA.2 and BA.3 suggests the recent BA.1-infected individuals are likely to be protected against the ongoing BA.2 surge. Third, vaccine-mediated T cell immunity and non-neutralizing antibodies that mediate antibody-dependent cytotoxicity could also confer protection against severe COVID-19. After vaccination or infection, the majority of T cell epitopes are highly preserved against Omicron spikes⁷. In agreement with this notion, 3 doses of BNT162b2 conferred efficacy against Omicron disease, but the protection wanes over time, with overall efficacy remaining high up to 6 months after dose 3⁸⁻¹². The real-world vaccine effectiveness and laboratory studies will guide vaccine booster strategy to achieve optimal breadth and duration of protection.

Methods

Recombinant Omicron spike mNG SARS-CoV-2s. The construction and characterization of recombinant Omicron BA.1-, BA.2-, and BA.3-spike mNG SARS-CoV-2s were recently reported⁴. The BA.1, BA.2, and BA.3 spike sequences were derived from GISAID EPI_ISL_6640916, EPI_ISL_6795834.2, and EPI_ISL_7605591, respectively. Passage 1 (P1) virus stocks were produced from infectious cDNA clones of corresponding viruses^{13,14}. The P1 viruses were used for neutralization testing throughout the study. The spike gene from each P1 virus was sequenced to ensure no undesired mutations. Equivalent specific infectivities, defined by the genomic RNA-to-FFU (fluorescent focus-forming unit)

ratios, were confirmed for individual recombinant P1 virus stocks, as previously reported⁴.

Serum specimens. The research protocol regarding the use of human serum specimens was reviewed and approved by the University of Texas Medical Branch (UTMB) Institutional Review Board (IRB number 20-0070). The de-identified human sera from unvaccinated patients who were infected by Omicron sublineage BA.1 were heat-inactivated at 56°C for 30 min before neutralization testing. The genotype of infecting virus was verified by the molecular tests with FDA's Emergency Use Authorization and Sanger sequencing. The serum information is presented in **Table 1**.

Fluorescent focus reduction neutralization test. Neutralization titers of sera were measured by fluorescent focus reduction neutralization test (FFRNT) using the USA-WA1/2020, BA.1-, BA.2-, and BA.3-spike mNG SARS-CoV-2s. The FFRNT protocol was reported previously⁵. Briefly, Vero E6 cells were seeded to 96-well plates at 2.5×10^4 per well (Greiner Bio-one™). On the following day, heat-inactivated sera were 2-fold serially diluted in culture medium with the first dilution of 1:20 (final dilutions ranging from 1:20 to 1:20,480). The diluted serum was incubated with 100-150 FFUs of indicated mNG SARS-CoV-2s at 37°C for 1 h. Afterwards, the serum-virus mixtures were loaded onto the pre-seeded Vero E6 cell monolayer in 96-well plates. After 1 h infection, the inoculum was aspirated and overlay medium (100 µl supplemented with 0.8% methylcellulose) was added to each well. After incubating the plates at 37°C for 16-18 h, raw images of mNG

foci were acquired using Cytation™ 7 (BioTek). The foci in each well were counted and normalized to the no-serum-treated controls to calculate infection rates. The FFRNT₅₀ value was defined as the minimal serum dilution that suppressed >50% of fluorescent foci. The neutralization titer of each serum was determined in duplicates, and the geometric mean was presented. FFRNT₅₀ of <20 was treated as 10 for plot purpose and statistical analysis. **Tables 1** summarizes the FFRNT₅₀ results.

Statistics. The nonparametric Wilcoxon matched-pairs signed rank test was used to analyze the statistical significance in **Figure 1B**.

Data availability

The data that support the findings of this study are available from the corresponding authors upon request.

REFERENCES

1. Nicole Wolter, Waasila Jassat, DATCOV-Gen author group, Anne von Gottberg, Cohen C. Clinical severity of Omicron sub-lineage BA.2 compared to BA.1 in South Africa. MedRxiv 2022: <https://doi.org/10.1101/2022.02.17.22271030>.
2. Frederik Plesner Lyngse, Carsten Thure Kirkeby, Matthew Denwood, et al. Transmission of SARS-CoV-2 Omicron VOC subvariants BA.1 and BA.2: Evidence from Danish Households. BioRxiv 2022:doi: <https://doi.org/10.1101/2022.01.28.22270044>.
3. Xia H, Zou J, Kurhade C, et al. Neutralization and durability of 2 or 3 doses of the BNT162b2 vaccine against Omicron SARS-CoV-2. Cell Host Microbe 2022.
4. Kurhade C, Zou J, Xia H, et al. Neutralization of Omicron BA.1, BA.2, and BA.3 SARS-CoV-2 by 3 doses of BNT162b2 vaccine. BioRxiv 2022:<https://biorxiv.org/cgi/content/short/2022.03.24.485633v1>.
5. Zou J, Xia H, Xie X, et al. Neutralization against Omicron SARS-CoV-2 from previous non-Omicron infection. Nat Commun 2022;13:852.
6. Liu Y, Liu J, Zou J, et al. Distinct neutralizing kinetics and magnitudes elicited by different SARS-CoV-2 variant spikes. bioRxiv 2021.

7. Redd AD, Nardin A, Kared H, et al. Minimal Crossover between Mutations Associated with Omicron Variant of SARS-CoV-2 and CD8(+) T-Cell Epitopes Identified in COVID-19 Convalescent Individuals. *mBio* 2022:e0361721.
8. Ferdinands JM, Rao S, Dixon BE, et al. Waning 2-Dose and 3-Dose Effectiveness of mRNA Vaccines Against COVID-19-Associated Emergency Department and Urgent Care Encounters and Hospitalizations Among Adults During Periods of Delta and Omicron Variant Predominance - VISION Network, 10 States, August 2021-January 2022. *MMWR Morb Mortal Wkly Rep* 2022;71:255-63.
9. Chemaitelly H, Abu-Raddad LJ. Waning effectiveness of COVID-19 vaccines. *Lancet* 2022;399:771-3.
10. Tartof SY, Slezak JM, Puzniak L, et al. BNT162b2 (Pfizer–Biontech) mRNA COVID-19 Vaccine Against Omicron-Related Hospital and Emergency Department Admission in a Large US Health System: A Test-Negative Design. SSRN 2022:<https://ssrn.com/abstract=4011905> or <http://dx.doi.org/10.2139/ssrn>.
11. Andrews N, Stowe J, Kirsebom F, et al. Covid-19 Vaccine Effectiveness against the Omicron (B.1.1.529) Variant. *N Engl J Med* 2022.
12. Agency UHS. COVID-19 vaccine surveillance report – Week 9, 3 March 2022. 2022:https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/1058464/Vaccine-surveillance-report-week-9.pdf.
13. Xie X, Muruato A, Lokugamage KG, et al. An Infectious cDNA Clone of SARS-CoV-2. *Cell Host Microbe* 2020;27:841-8 e3.
14. Xie X, Lokugamage KG, Zhang XV, M.N. , Muruato AE, Menachery VD, Shi P-Y. Engineering SARS-CoV-2 using a reverse genetic system. *Nature Protocols* 2021;16:1761-84.

Acknowledgments

We thank colleagues at the University of Texas Medical Branch (UTMB) for helpful discussions. We thank Michael L O'Rourke from the Information System Department at UTMB for assisting with electronic medical record systems. P.-Y.S. was supported by NIH grants HHSN272201600013C, AI134907, AI145617, and UL1TR001439, and awards from the Sealy & Smith Foundation, the Kleberg Foundation, the John S. Dunn Foundation, the Amon G. Carter Foundation, the Gilson Longenbaugh Foundation, and the Summerfield Robert Foundation.

Author contributions

Conceptualization, X.X., P.R., P.-Y.S.; Methodology, J.Z., C.K., M.L., X.X., P.R., P.-Y.S.; Investigation, J.Z., C.K., M.L., X.X., P.R., P.-Y.S.; Resources, C.K., M.L., X.X., P.R., P.-Y.S.; Data Curation, J.Z., C.K., X.X., P.R., P.-Y.S.; Writing-Original Draft, J.Z., H.X., X.X., R.R., P.-Y.S.; Writing-Review & Editing, X.X., P.R., P.-Y.S.; Supervision, X.X., P.R., P.-Y.S.; Funding Acquisition, P.-Y.S.

Competing interests

X.X. and P.-Y.S. have filed a patent on the reverse genetic system. J.Z., C.K., X.X., and P.-Y.S. received compensation from Pfizer for COVID-19 vaccine development. Other authors declare no competing interests.




Figure 1. Cross-neutralization of human sera from unvaccinated individuals who were infected with Omicron BA.1 SARS-CoV-2. (A) Omicron BA.1-, BA.2-, and BA.3-spike mNG SARS-CoV-2s. The full-length spike gene from Omicron BA.1, BA.2, or BA.3 was engineered into an mNG USA-WA1/2020 SARS-CoV-2. The mNG gene was engineered at the open-reading-frame-7 of the viral genome. Amino acid mutations, deletions, and insertions (Ins) are indicated for BA.1, BA.2, and BA.3 spikes in reference to the USA-WA1/2020 spike. L: leader sequence; ORF: open reading frame; NTD: N-terminal domain of S1; RBD: receptor binding domain of S1; S: spike glycoprotein; S1: N-terminal furin cleavage fragment of S; S2: C-terminal furin cleavage fragment of S; E: envelope protein; M: membrane protein; N: nucleoprotein; UTR: untranslated region. (B) Scatterplot of neutralization titers. A panel of 20 human sera collected from Omicron BA.1-infected individuals were tested for the 50% fluorescent focus-reduction neutralization titers (FFRNT₅₀) against recombinant USA-WA1/2020, Omicron BA.1-, BA.2-, and BA.3-spike mNG SARS-CoV-2s. The neutralization titer for each virus was determined in duplicates. The serum information and FFRNT₅₀ values are summarized in **Table 1**. Each data point represents the geometric mean FFRNT₅₀ obtained with a serum specimen against the indicated virus. The bar heights and the numbers above indicate geometric mean titers (GMTs). Error bars indicate the 95% confidence interval (CI) of the GMTs. Statistical analysis was performed using the Wilcoxon matched-pairs signed-rank test. *P* values of the GMT against BA.1-spike and USA-WA1/2020, BA.2-spike, or BA.3 spikes

are all <0.0001 . (C) FFRNT₅₀ values with connected lines for individual sera. Two sera exhibiting slightly higher FFRNT₅₀s against BA.2 virus than that against BA.1-spike SARS-CoV-2 are indicated by symbol * (serum ID 1 and 2 in **Table 1**).

Table 1. Serum information and FFRNT₅₀ values

Serum ID	Age (years)	Gender (F/M)	Ethnicity	*FFRNT ₅₀				Serum collection time (days post positive RT-PCR test)
				USA-WA1/2020	BA.1-spike virus	BA.2-spike virus	BA.3-spike virus	
1	29	M	Hispanic	^10	80	160	20	26
2	41	F	White	10	80	113	40	33
3	84	M	White	14	113	40	14	21
4	14	M	Black	10	113	28	28	16
5	35	F	Black	10	160	40	40	16
6	28	F	Black	10	160	20	20	28
7	9	F	Hispanic	10	160	57	57	43
8	64	M	Black	10	160	80	20	40
9	84	M	White	20	226	160	226	40
10	5	F	Hispanic	10	320	28	40	26
11	1	F	Hispanic	10	320	113	160	56
12	57	F	White	10	453	28	80	35
13	75	M	Hispanic	160	453	10	20	17
14	26	F	Hispanic	20	640	453	160	62
15	55	M	White	10	905	40	113	29
16	74	F	^{\$} NA	10	1280	226	320	8
17	71	F	Hispanic	28	2560	453	640	32
18	63	M	Hispanic	113	2560	1280	1810	15
19	78	F	Hispanic	10	5120	640	1280	16
20	84	M	White	40	14482	2560	2560	13
[#] GMT	34	-	-	16	445	107	102	25
[†] 95% CI	20-58	-	-	11-24	225-881	54-215	48-218	20-32

*Individual FFRNT₅₀ value is the geometric mean of duplicate plaque assay results.

^FFRNT₅₀ of <20 was treated as 10 for plot purpose and statistical analysis.

^{\$}NA: not available.

[#]Geometric mean neutralizing titers (GMT).

[†]95% confidence interval (95% CI) for the GMT.