OTTERS: A powerful TWAS framework leveraging summary-level reference data

Qile Dai[1,12], Geyu Zhou[2], Hongyu Zhao[2,3], Urmo Võsa[4], Lude Franke[5,6], Alexis Battle[7], Alexander Teumer[8], Terho Lehtimäki[9], Olli Raitakari[10,11], Tõnu Esko[4], eQTLGen Consortium, Michael P. Epstein[12*], Jingjing Yang[12*]

1. Department of Biostatistics and Bioinformatics, Emory University School of Public Health, Atlanta, Georgia 30322, United States of America.
2. Program of Computational Biology and Bioinformatics, Yale University, New Haven, Connecticut 06511, United States of America.
3. Department of Biostatistics, Yale School of Public Health, New Haven, CT 06520, United States of America.
4. Estonian Genome Centre, Institute of Genomics, University of Tartu, 50090 Tartu, Estonia.
5. Department of Genetics, University of Groningen, University Medical Center Groningen, 9700 RB Groningen, Netherlands.
6. Oncode Institute, 3521 AL Utrecht, Netherlands.
7. Department of Computer Science, and Departments of Biomedical Engineering, Johns Hopkins University, Baltimore, Maryland 21218, United States of America
8. Institute for Community Medicine, University Medicine Greifswald, 17489 Greifswald, Germany.
9. Department of Clinical Chemistry, Fimlab Laboratories, and Finnish Cardiovascular Research Center, Tampere, Faculty of Medicine and Health Technology, Tampere University, Tampere 33520, Finland.
10. Centre for Population Health Research, and Research Centre of Applied and Preventive Cardiovascular Medicine, University of Turku, 20500 Turku, Finland.
11. Department of Clinical Physiology and Nuclear Medicine, Turku University Hospital, 20521 Turku, Finland.
12. Center for Computational and Quantitative Genetics, Department of Human Genetics, Emory University School of Medicine, Atlanta, Georgia 30322, United States of America.

*Correspondence Authors: M.P.E. (mpepste@emory.edu) and J.Y. (jingjing.yang@emory.edu)

Abstract

Most existing TWAS tools require individual-level eQTL reference data and thus are not applicable to summary-level reference eQTL datasets. The development of TWAS methods that can harness summary-level reference data is valuable to enable TWAS in broader settings and enhance power due to increased reference sample size. Thus, we develop a TWAS framework called OTTERS (Omnibus Transcriptome Test using Expression Reference Summary data) that adapts multiple polygenic risk score (PRS) methods to estimate eQTL weights from summary-level eQTL reference data and conducts an omnibus TWAS. We show that OTTERS is a practical and powerful TWAS tool by both simulations and application studies.

Keywords:

Transcriptome-wide association study; Summary-level eQTL reference data; PRS method; GWAS; UK Biobank; eQTLGen; Cardiovascular disease

Transcriptome-wide association study (TWAS) is a valuable analysis strategy for identifying genes that influence complex traits and diseases through genetic regulation of gene expression[1–5]. Researchers have successfully deployed TWAS analyses to identify risk genes for complex human diseases, including Alzheimer's disease[6–8], breast cancer[9–11], ovarian cancer[12,13], and cardiovascular disease[14,15]. A typical TWAS consists of two separate stages. In Stage I, TWAS acquires individual-level genetic and expression data from relevant tissues available in a reference dataset like the Genotype-Tissue Expression (GTEx) project[16,17] or the North American Brain Expression Consortium (NABEC)[18], and trains multivariable regression models on the reference data treating gene expression as outcome and SNP genotype data (typically cis-SNPs nearby the test gene) as predictors to determine genetically regulated expression (GReX). After

Stage I that uses the GReX regression models to estimate effect sizes of SNP predictors that, in

the broad sense, are expression quantitative trait loci (eQTLs), Stage II of TWAS proceeds by

using these trained eQTL effect sizes to impute GReX within an independent GWAS of a complex

human disease or trait. One can then test for association between the imputed GReX and

phenotype, which is equivalent to a gene-based association test taking these eQTL effect sizes

as corresponding test SNP weights[19–21].

For Stage I of TWAS, a variety of training tools exist for fitting GReX regression models

using reference expression and genetic data, including PrediXcan[19], FUSION[20], and TIGAR[22].

While these methods all employ different techniques for model fitting, they all require individual-

level reference expression and genetic data to estimate eQTL effect sizes for TWAS. Therefore,

these methods cannot be applied to emerging reference summary-level eQTL results such as

those generated by the eQTLGen[23] and CommonMind[24] consortia, which provide eQTL effect

sizes and p-values relating individual SNPs to gene expression. The development of TWAS

methods that can utilize such summary-level reference data is valuable to permit applicability of

the technique to broader analysis settings. Moreover, as TWAS power increases with increasing

reference sample size[25], TWAS using summary-level reference datasets can lead to enhanced

performance compared to using individual-level reference datasets since the sample sizes of the

former often are considerably larger than the latter. For example, the sample size of the summary-

based eQTLGen reference sample is 31,684 for blood, whereas the sample size of the individual-

level GTEx V6 reference is only 338 for the same tissue. Consequently, TWAS analysis

leveraging the summary-based eQTLGen dataset as reference likely can provide novel insights

into genetic regulation of complex human traits.

In this work, we propose a framework that can use summary-level reference data to train

GReX regression models required for Stage I of TWAS analysis. Our method is motivated by a

variety of published polygenic risk score (PRS) methods[26–31] that can predict phenotype in a test

dataset using summary-level SNP effect-size estimates and p-values based on single SNP tests

from an independent reference GWAS. We can adapt these PRS methods for TWAS since eQTL

effect sizes are essentially SNP effect sizes resulting from a reference "GWAS" of gene

expression. Thus, our predicted GReX in Stage II of TWAS is analogous to the PRS constructed

based on training GWAS summary statistics of single SNP-trait association. Here, we adapt four

representative summary-data based PRS methods — P-value Thresholding with linkage

disequilibrium (LD) clumping (*P+T*)[26], frequentist LASSO[32] regression based method *lassosum*[27],

nonparametric Bayesian Dirichlet Process Regression (DPR) model[33] based method *SDPR*[29],

and Bayesian multivariable regression model based method with continuous shrinkage (CS)

priors *PRS-CS*[28] for TWAS analysis. We apply each of these PRS methods to first train eQTL

effect sizes based on a multivariable regression model from summary-level reference eQTL data

(Stage I), and subsequently use these eQTL effect sizes (i.e., eQTL weights) to impute GReX

and then test GReX-trait association in an independent test GWAS (Stage II).

As we will show, the PRS method with optimal performance for TWAS depends on the

underlying genetic architecture for gene expression. Since the genetic architecture of expression

is unknown apriori, we maximize the performance of TWAS over different possible architectures

by proposing a novel TWAS framework called OTTERS (**O**mnibus **T**ranscriptome **T**est using

Expression **R**eference **S**ummary data). OTTERS first constructs individual TWAS tests and p-

values using eQTL weights trained by each of the PRS techniques outlined above, and then

calculates an omnibus test p-value using the aggregated Cauchy association test[34] (ACAT-O)

with all individual TWAS p-values (Figure 1). OTTERS is applicable to both summary-level and

individual-level test GWAS data within Stage II TWAS analysis.

In subsequent sections, we first describe how to use the PRS methods on summary-level

reference eQTL data in Stage I TWAS, and then describe how we can use the resulting eQTL

weights to perform Stage II TWAS using OTTERS. We then evaluate the performance of

individual PRS methods and OTTERS using simulated expression and real genetic data based

on patterns observed in real datasets. Interestingly, when we assume individual-level reference

data are available, we observe that OTTERS outperforms the popular FUSION[20] approach across

all simulation settings considered. Many of the individual PRS methods also outperform FUSION

in these settings. We then apply OTTERS to blood eQTL summary-level data (n=31,684) from

the eQTLGen consortium[23] and GWAS summary data of cardiovascular disease from the UK

Biobank (UKBB)[35]. By comparing OTTERS results to those of FUSION[20] using individual-level

GTEx reference data of whole blood tissue, we demonstrate that OTTERS using large summary-

level reference datasets and multiple gene expression imputation models can successfully reveal

potential risk genes missed by FUSION based on smaller individual-level reference datasets and

only one model. Finally, we conclude with a discussion.

Results

Method Overview

For the standard two-stage TWAS approach, Stage I estimates a GReX imputation model

using individual-level expression and genotype data available from a reference dataset, and then

Stage II uses the eQTL effect sizes from Stage I to impute gene expression (GReX) in an

independent GWAS and test for association between GReX and phenotype. GReX for test

samples can be imputed from individual-level genotype data and eQTL effect size estimates.

When individual-level GWAS data are not available, one can instead use summary-level GWAS

data for TWAS by applying the TWAS Z-score statistics proposed by FUSION[20] and S-

PrediXcan[36] (see details in Methods).

Since eQTL summary data are analogous to GWAS summary data where gene

expression represents the phenotype, we can follow the idea from PRS methods to estimate the

eQTL effect sizes based on a multivariable regression model using only marginal least squared

effect estimates and p-values (based on a single variant test) from the eQTL summary data as

well as a reference LD panel from samples of the same ancestry[26–29]. Although all PRS methods

are applicable to TWAS Stage I, we only consider four representative methods — *P+T*[26],

Frequentist lassosum[27], *Nonparametric Bayesian SDPR*[29]*, Bayesian PRS-CS*[28] (see details in

Methods)*.

As shown in Figure 1, OTTERS first trains GReX imputation models per gene *g* using *P+T,*

lassosum, SDPR, and *PRS-CS* methods that each infers cis-eQTL weights using cis-eQTL

summary data and an external LD reference panel of the same ancestry (Stage I). Once we derive

cis-eQTL weights for each training method, we can impute the respective GReX using that

method and perform the respective gene-based association analysis in the test GWAS dataset.

We thus derive a set of TWAS p-values for gene *g,* one per training method. We then use these

TWAS p-values to create an omnibus test using the ACAT-O[34] approach that employs a Cauchy

distribution for inference (see details in Supplemental Methods). We refer to the p-value derived

from ACAT-O test as the OTTERS p-value. The ACAT-O[34] approach has been widely used in

hypothesis testing to combine multiple testing methods for the same hypothesis[37–39], which has

been shown as an effective approach to leverage different test methods to increase the power

while still managing to control for type I error. Adding TWAS p-values based on additional PRS

methods to the ACAT-O test can possibly improve the power further at the cost of additional

computation.

Simulation Study

We used real genotype data from 1894 whole genome sequencing (WGS) samples from

the Religious Orders Study and Rush Memory and Aging Project (ROS/MAP) cohort[40,41] and

Mount Sinai Brain Bank (MSBB) study[42] for simulation. We divided 14,772 genes into five

groups according to gene length, and randomly selected 100 genes from each group (500

genes in total). We randomly split samples into 568 training (30%) and 1326 testing samples

(70%) to mimic a relatively small sample size in the real reference panel for training gene

expression imputation models.. From the real genotype data, we simulated 6 scenarios with 2

different proportions of causal cis-eQTL, $p_{causal} = (0.001, 0.01)$, as well as 3 different

proportions of gene expression variance explained by causal eQTL, $h_e^2 = (0.01, 0.05, 0.1)$.

We generated gene expression of gene g (E_g) using the multivariable regression model

$E_g = X_g w + \epsilon_g$, where X_g represents the standardized genotype matrix of the randomly

selected causal eQTL of gene g, $\epsilon_g \sim N(0, (1 - h_e^2)I)$. We generated the eQTL effect sizes w

from $N(0,1)$ and then re-scaled these effects to ensure that the expression variance explained

by all causal variants is h_e^2. We generated 10 replicates of gene expression per scenario. For

each simulated gene expression, we then generated 10 sets of GWAS Z-scores to perform a

total of 50,000 TWAS simulations. We generated the GWAS Z-scores from a multivariate

normal distribution with $Z \sim MVN\left(\Sigma_g w \sqrt{n_{gwas} h_p^2}, \Sigma_g\right)$[38], where w is the true causal eQTL

effect sizes, Σ_g is the correlation matrix of the standardized genotype X_g from test samples,

n_{gwas} is the assumed GWAS sample size, and h_p^2 denotes the amount of phenotypic variance

explained by simulated GReX=$X_g w$ (see Methods). We set $h_p^2 = 0.025$. To calibrate power, we

considered n_{gwas} = (200K, 300K, 400K, 500K) for scenarios with h_e^2 = 0.01, n_{gwas} = (25K, 50K,

75K, 100K) for scenarios with h_e^2 = 0.05, and n_{gwas} = (10K, 20K, 30K, 40K) for scenarios with

$h_e^2 = 0.1$.

In Stage I of our TWAS analysis, we applied *P+T (0.001)*, *P+T (0.05)*, *lassosum*, *SDPR*,

and *PRS-CS* methods to estimate eQTL weights using eQTL summary data and the reference

LD of training samples. In Stage II of the TWAS, we used the estimated eQTL weights and the

simulated GWAS Z-scores to conduct a gene-based association test. In addition to gene-based

association tests based on eQTL weights per training method, we further constructed the

corresponding OTTERS p-values. We evaluated the performance of the training methods with

test samples, comparing test R^2 that was the squared Pearson correlation coefficient between

imputed GReX and simulated gene expression. We evaluated TWAS power given by the

proportion of 50,000 repeated simulations with TWAS p-value $< 2.5 \times 10^{-6}$ (genome-wide

significance threshold adjusting for testing 20K independent genes). We only considered genes

with GReX test $R^2 > 0.01$ in the power analysis.

As shown in Figure 2, we demonstrated that the Stage I training method with optimal test

R^2 and TWAS power depended on the underlying genetic architecture of gene expression (p_{causal})

as well as gene expression heritability (h_e^2). In situations where true cis-eQTLs were sparse

(p_{causal} = 0.001) and the gene expression heritability was small (h_e^2 = 0.01), *P+T (0.05)* method

performed the best with the highest TWAS power among all individual methods. When gene

expression heritability is low (h_e^2 = 0.01), the power of *P+T (0.001)* and *lassosum* methods were

shown as the lowest for considering only genes with test $R^2 > 0.01$. When gene expression

heritability increased (h_e^2 = 0.05 or 0.1) within this sparse eQTL model, *P+T (0.001)* and PRS-CS

were generally the optimal methods*.* For a less sparse model with p_{causal} = 0.01, SDPR and PRS-

CS generally performed best among the individual methods. Relative to individual methods, we

found that combining the TWAS p-values based on the four PRS training methods together for

analysis in our OTTERS framework obtained the highest power across all scenarios.

To evaluate the type I error of the individual PRS methods along with OTTERS, we picked

one simulated replicate per gene from the scenario with $h_e^2 = 0.1$ and $p_{causal} = 0.001$,

simulated 2×10^3 phenotypes from $N(0,1)$, and permuted the eQTL weights for TWAS to

perform a total of 10^6 null simulations. OTTERS was shown well calibrated in the tails of the

distribution as shown by quantile-quantile (Q-Q) plots of TWAS p-values in Figure S1. We also

observed that OTTERS had well-controlled type I error for stringent significance levels between

10^{-4} and 2.5×10^{-6} (Table S1), which are typically utilized in TWAS. For more modest

significance thresholds ($\alpha = 10^{-2}$) , we noted that OTTERS had a slightly inflated type-I error rate.

This modest inflation is consistent with the findings of the original ACAT-O work, which showed

that the Cauchy-distribution-based approximation that ACAT-O employs may not be accurate for

larger p-values when correlation among tests is strong[34]. This suggests that modest OTTERS p-

values may be interpreted with caution.

We also compared the performance of our individual PRS training methods to those of

FUSION assuming individual-level reference data were available for the latter method to train

GReX models. As shown in Figure 2A, we interestingly observed that our training methods yielded

similar or improved test R^2 compared to FUSION in this situation, with *SDPR* and *PRS-CS*

outperforming FUSION across all simulation settings. Comparing TWAS power, we found that

OTTERS outperformed FUSION by a considerable margin in our simulations (Figure 2B). These

simulation results suggest that, while we developed OTTERS based on PRS training methods to

handle summary-level reference data, OTTERS can still improve TWAS power when individual-

level reference data are available. This is likely because OTTERS accounts for multiple possible

models of genetic architectures of gene expression assumed by the different PRS training

methods.

GReX Imputation Accuracy in GTEx V8 Blood Samples

To evaluate the imputation accuracy of *P+T (0.001)*, *P+T (0.05)*, *lassosum*, *SDPR*, and

PRS-CS methods in real data, we applied these training methods to summary-level eQTL

reference data from the eQTLGen consortium[23] with n=31,684 blood samples, to train GReX

imputation models for 16,699 genes. For test data, we downloaded the transcriptomic data of 315

blood tissue samples that are in GTEx V8 but were not part of GTEx V6 (as GTEx V6 samples

contributed to the reference eQTLGen consortium summary data). For these 315 samples, we

compared imputed GReX to observed expression levels. We considered trained imputation

models with test $R^2 > 0.01$ as "valid" models, as suggested by previous TWAS methods[20,43]. We

also compared imputation accuracy of these five training models to those using FUSION based

on a smaller individual-level training dataset (individual-level GTEx V6 reference dataset; see

Methods). For such models, we compared the test R^2 for genes that had test $R^2 > 0.01$ by at least

one training method.

We observed that *PRS-CS* obtained the most "valid" GReX imputation models with test

$R^2 > 0.01$. Among 16,699 tested genes, *PRS-CS* obtained "valid" GReX imputation models for

10,337 genes, compared to 9,816 genes by *P+T (0.001)* (5.0% less valid genes than PRS-CS),

9,662 genes by *P+T (0.05)* (6.5% less), 8,718 genes by *lassosum* (15.7% less), 9,670 genes by

SDPR (6.5% less), and 4,704 genes by FUSION (54.5% less) (Table 1). Among the "valid" GReX

imputation models obtained by each method, the ones trained by *PRS-CS* have the highest

median test R^2. The *P+T (0.001)* method obtained the second most "valid" GReX imputation

models with the second largest median test R^2, as compared to *P+T (0.05)*, *lassosum*, and *SDPR*

(Table 1). We note that the performance of PRS-CS method was not sensitive to the global

shrinkage parameter (Figure S2).

By comparing test R^2 per "valid" GReX imputation model by *PRS-CS* versus the other

methods (Figure 3), we observed that *PRS-CS* had the best overall performance for imputing

GReX as it provided the most "valid" models with higher GReX imputation accuracy compared to

P+T methods, *lassosum*, *SDPR*, and FUSION. Comparing the test R^2 among the other four

training methods, we observed that these two *P+T* methods obtained similar test R^2 per "valid"

model. Meanwhile, the test R^2 per valid model varied widely among the *P+T* methods, *lassosum*,

and *SDPR* (Figure S3), suggesting that none of these four were optimal across all genes and their

performance likely depended on the underlying unknown genetic architecture. These results are

consistent with our simulation results.

TWAS of Cardiovascular Disease

Using the eQTL weights trained by *P+T (0.001), P+T (0.05), lassosum, SDPR, and PRS-*

CS methods with the eQTLGen[23] reference data and reference LD from GTEx V8 WGS

samples[44], we applied our OTTERS framework to the summary-level GWAS data of

Cardiovascular Disease from UKBB (n=459,324, case fraction = 0.319)[35] (see Methods). We

performed TWAS of cardiovascular disease for 16,678 genes. First, for each gene, we obtained

TWAS p-values per individual training method (*P+T (0.001), P+T (0.05), lassosum, SDPR, and*

PRS-CS). Second, we performed genomic control[45] for TWAS test statistics generated under

each specific training model, by scaling all test statistics to ensure that the median test p-value

equals to 0.5. Last, we only considered genes with test GReX $R^2 > 0.01$ by at least one PRS

training method in additional GTEx V8 samples in the follow-up ACAT-O test. We combined the

adjusted p-values across all five training models using ACAT-O to obtain our OTTERS test

statistics and p-values. Genes with OTTERS p-values $< 2.998 \times 10^{-6}$ (Bonferroni corrected

significance level) were identified as significant TWAS genes for cardiovascular risk.

In total, we identified 40 significant TWAS genes by using OTTERS. To identify

independently significant TWAS genes, we calculated the R^2 (squared correlation) between the

GReX predicted by *PRS-CS* for of each pair of genes. For a pair of genes with the predicted

GReX $R^2 > 0.5$, we only kept the gene with the smaller TWAS p-value as the independently

significant gene. OTTERS obtained 38 independently significant TWAS genes (Table 2, Figure

3B), compared to 17 independently significant genes by *P+T (0.001)*, 11 by *P+T (0.05)*, 10 by

lassosum, 41 by *SDPR*, and 12 by *PRS-CS*. Among these 38 independent TWAS risk genes

identified by OTTERS, gene *RP11-378A13.1* (OTTERS p-value = 9.78×10^{-9}) was not within 1

MB of any known GWAS risk loci with genomic-control corrected p-value $< 5 \times 10^{-8}$ in the

UKBB summary-level GWAS data. This novel risk gene *RP11-378A13.1* was also identified to

be a significant TWAS risk gene in blood tissue for systolic blood pressure, high cholesterol, and

cardiovascular disease by FUSION[1].

We compared our OTTERS results with the TWAS results shown on TWAS hub (see

Web Resource) obtained by FUSION using the same UKBB GWAS summary data of

cardiovascular disease but using a smaller individual-level reference expression dataset from

GTEx V6 (whole blood tissue, n=338). Of the 38 independent genes that OTTERS identified

from TWAS with eQTLGen reference data of whole blood, FUSION only identified 8 of these

genes (*CLCN6b, PSRC1, RP11-378A13.1, CAMK1D, SIDT2, MTHFSD, NTN5, OPRL1*) when

using the GTEx V6 reference data of the same tissue. FUSION did identify 13 additional

OTTERS genes (*NPPA, CPEB4, NT5C2, TNNT3, C11orf49, CSK, FES, MBTPS1, ACE, MRI1,*

HAUS8, RPL28, CTSZ), when considering all available tissue types in GTEx V6 reference data.

These genes were identified by FUSION when considering the GTEx V6 reference data of

artery, thyroid, adipose visceral, and nerve tibial tissues. For example, the most significant gene

FES (OTTERS p-value = 2.87×10^{-32}) was identified by FUSION using GTEx reference data of

artery tibial, thyroid, and adipose visceral omentum tissues, and was also identified as a TWAS

risk gene for high blood pressure, which is strongly related to cardiovascular disease[46].

Our OTTERS method also identified 17 novel risk genes (*LINC01093, SERPINB6,*

CARMIL1, ZSCAN12P1, HCG4P7, HCG4P3, HLA-S, PSPHP1, LPL, PTP4A3, SLCO3A1,

RALBP1, SULT2B1, EDN3, ZBTB46, FAM3B, MX1) that were not detected by FUSION using

GTEx V6 data, where *EDN3* (Endothelin 3, a member of the endothelin family) was shown to be

active in the cardiovascular system and play an important role in the maintenance of blood

pressure or generation of hypertension[47].

By comparing OTTERS results with the ones obtained by individual methods (Table 2;

Figure 4; Figure S4), we found that all individual methods contributed to the OTTERS results. For

example, the novel risk gene *LINC01093* was only identified by *lassosum*, while genes *CPEB4,*

SIDT2, and *ACE* were only detected by *PRS-CS* and *SDPR* and the novel risk gene *EDN3* was

only identified by the *P+T* methods. To better understand the differences among individual

methods, we plotted the eQTL weights estimated by *P+T (0.001)*, *P+T (0.05)*, *lassosum*, *SDPR*,

and *PRS-CS* for three example genes that were only detected by one or two individual methods

(Figures S5-S7). For these genes, we plotted the eQTL weights produced by each method with

such weights color coded with respect to $-log_{10}$ (GWAS p-values) from the UKBB GWAS

summary statistics and shape coded with respect to the direction of UKBB GWAS Z-score

statistics. Generally, significant TWAS p-values would be obtained by methods that obtained

eQTL weights with relatively large magnitude for SNPs with relatively more significant GWAS p-

values.

In Figure S5, we showed the eQTL weights for gene *SIDT2*, which was a significant risk

gene identified by both *PRS-CS* and *SDPR*, and had p-values $< 10^{-4}$ by other methods.

Compared to *lassosum*, *SDPR* had more significant GWAS SNPs colocalized with eQTLs

having relatively large weights in the test region, and *PRS-CS* had more non-significant GWAS

SNPs colocalized with eQTLs having zero weights. Compared to the P+T methods, SDPR and

PRS-CS based on a multivariate regression model modeled LD among all test SNPs, and thus

estimated eQTL weights leading to significant TWAS findings. In Figure S6, we provided the

results of gene *EDN3*, which was only identified by *P+T* methods (p-values $\leq 9.15 \times 10^{-8}$).

Compared to *P+T* methods, *SDPR* (p-value = 5.9×10^{-3}) and *PRS-CS* (p-value = 0.0158) had

fewer significant GWAS SNPs colocalized with eQTLs that had relatively large weights in the

test region, while *lassosum* (p-value = 8.6×10^{-6}) assigned relatively large weights to more

non-significant GWAS SNPs. In Figure S7, we provided results for gene *LINC01093*, which was

only identified by *lassosum*. For this gene, *SDPR* and *PRS-CS* estimated near-zero weights for

most test SNPs with significant GWAS p-values in the test region. Most significant GWAS SNPs

did not have eQTL test p-values < 0.001 or 0.05, and were thus filtered out by *P+T* methods.

lassosum was the only method that produced relatively large eQTL weights that co-localized

with GWAS significant SNPs.

These results were consistent with our simulation study results, demonstrating that the performance of different individual methods depended on the underlying genetic architecture. We do note that there were a handful of genes identified by an individual method that were not significant using OTTERS (Table S2). Nonetheless, the omnibus test borrows strength across all individual methods, thus generally achieves higher TWAS power and identifies the group of most robust TWAS risk genes.

By examining the Q-Q plots of TWAS p-values, we observed a moderate inflation for all methods (Figure S8). Such inflation in TWAS results is not uncommon[48–50], which could be due to similar inflation in the GWAS summary data and not distinguishing the pleiotropy and mediation effects for considered gene expression and phenotype of interest[51] (Figure S9). We also observed a notable inflation in the GWAS p-values of cardiovascular disease from UKBB (Figure S9), as we estimated the LD score regression[52] intercept to be 1.1 from the GWAS summary data.

We did not consider directly comparing to FUSION in our above TWAS analyses of cardiovascular disease since we used the summary-level reference data eQTLGen. However, to assess the performance of OTTERS and FUSION in a real study where individual-level reference data are available, we performed an additional TWAS analysis of cardiovascular disease in the UK Biobank using the GTEx V8 data of 574 whole blood samples as the reference data. We trained OTTERS Stage I using cis-eQTL summary statistics obtained from these 574 GTEx V8 whole blood samples and reference LD from GTEx V8 WGS samples, and trained FUSION models using individual-level genotype data and gene expression data of the same 574 whole blood samples.

We tested TWAS association for 19,653 genes and identified genes with TWAS p-values $< 2.53 \times 10^{-6}$ (Bonferroni corrected significance level) as significant TWAS genes. Training $R^2 >$

0.01 was used to select "valid" GReX imputation models for TWAS (Figure S10). To identify independently significant TWAS genes, we calculated the training R^2 between the GReX predicted by lassosum for of each pair of genes, since lassosum had the best training R^2 (Figure S10). For a pair of genes with the predicted GReX $R^2 > 0.5$, we only kept the gene with the smaller TWAS p-value as the independently significant gene. As a result, OTTERS obtained 34 independently significant TWAS genes, while FUSION identified 21 independently significant TWAS genes (Figure S11). A total of 14 genes were identified by both FUSION and OTTERS (Table S3).

These results demonstrate the advantages of OTTERS for using multiple PRS training methods to account for the unknown genetic architecture of gene expression, which is consistent in our simulation results. These results also showed the advantage of using eQTL summary data with a larger training sample size, as more independently significant TWAS genes were identified by using the eQTLGen summary reference data (38 vs. 34), even with a more stringent rule (test instead of training $R^2 > 0.01$) applied to select test genes with "valid" GReX imputation models.

Computational Time

The computational time per gene of different PRS methods depends on the number of test variants considered for the target gene. Thus, we calculated the computational time and memory usage for 4 groups of genes whose test variants were <2000, between 2000 and 3000, between 3000 and 4000, and >4000, respectively. Among all tested genes in our real studies, the median number of test variants per gene is 3152, and the proportion of genes in each group is 10.3%, 33.4%, 34.5%, and 21.8%, respectively. For each group, we randomly selected 10 genes on Chromosome 4 to evaluate the average computational time and memory usage per gene. We benchmarked the computational time and memory usage of each method on one Intel(R) Xeon(R) processor (2.10 GHz). The evaluation was based on 1000 MCMC iterations for SDPR and PRS-CS (default) without parallel computation (Table S4). We showed that *P+T* and *lassosum were*

378 computationally more efficient than *SDPR* and *PRS-CS*, whose speed were impeded by the need

379 of MCMC iterations. Between the two Bayesian methods, SDPR implemented in C++ uses

380 significantly less time and memory than PRS-CS implemented in Python.

Discussion

Our OTTERS framework represents an omnibus TWAS tool that can leverage summary-

level expression and genotype results from a reference sample, thereby robustly expanding the

use of TWAS into more settings. To this end, we adapted and evaluated five different PRS

methods assuming different underlying genetic models, including the relatively simple method

P+T[26] with two different p-value thresholds (0.001 and 0.05), the frequentist method *lassosum*[27],

as well as the Bayesian methods *PRS-CS*[28] and *SDPR*[29] within our omnibus test for optimal

inference. We note that additional PRS methods such as MegaPRS[30] or PUMAS[31] could also be

implemented as additional OTTERS Stage I training methods. Higher TWAS power might be

obtained by adding more PRS methods in OTTERS Stage I, with additional computation cost. We

also note that the existing SMR-HEIDI[53] method, which uses summary-level data from GWAS and

eQTL studies to test for possible causal genetic effects of a trait of interest that were mediated through

gene expression, could also be used as an alternative method besides TWAS. However, the SMR

method generally restricts eQTL for consideration, excluding those where the eQTL p-values

larger than a certain threshold, e.g., 0.05.

In simulation studies, we demonstrated that the performance of each of these five PRS

methods depended substantially on the underlying genetic architecture for gene expression, with

P+T methods generally performing better for sparse architecture whereas the Bayesian methods

performing better for denser architecture. Consequently, since genetic architecture of gene

expression is unknown apriori, we believe this justifies the use of the omnibus TWAS test

implemented in OTTERS for practical use as this test had near-optimal performance across all

simulation scenarios considered. While we developed our methods with summary-level reference

data in mind, we note that our prediction methods and OTTERS perform well (in terms of

imputation accuracy and power) relative to existing TWAS methods like FUSION when individual-

level reference data are available.

In our real data application using UKBB GWAS summary-level data, we compared

OTTERS TWAS results using reference eQTL summary data from eQTLGen consortium to

FUSION TWAS results using a substantially smaller individual-level reference dataset from GTEx

V6. OTTERS identified 13 significant TWAS risk genes that were missed by FUSION using

individual-level GTEx V6 reference data of blood tissue, suggesting that the use of larger

reference datasets like eQTLGen in TWAS can identify novel findings. Interestingly, the genes

missed by FUSION were instead detected using individual-level GTEx reference data of other

tissue types that are more directly related to cardiovascular disease. By comparing OTTERS to

FUSION when the same individual-level GTEx V8 reference data of whole blood samples were

used, we still observed that OTTERS identified more risk genes than FUSION, which we believe

is due to the former method accounting for the unknown genetic architecture of gene expression

by using multiple regression methods to train GReX imputation models. These applied results

were consistent with our simulation results.

Among all individual methods, *P+T* is the most computationally efficient method. The

Bayesian methods *SDPR* and *PRS-CS* require more computation time than the frequentist

method *lassosum* as the former set of methods require a large number of MCMC iterations for

model fit. By comparing the performance of these five methods in terms of the imputation accuracy

and TWAS power in simulations and real applications, we conclude that none of these methods

were optimal across different genetic architectures. We found that all methods provided distinct

and considerable contributions to the final OTTERS TWAS results. These results demonstrate

the benefits of OTTERS in practice, since OTTERS can combine the strength of these individual

methods to achieve the optimal performance.

To enable the use of OTTERS by the public, we provide an integrated tool (see Availability

of data and materials) to: (1) Train GReX imputation models (i.e., estimate eQTL weights in Stage

I) using eQTL summary data by *P+T*, *lassosum*, *SDPR*, and *PRS-CS*; (2) Conduct TWAS (i.e.,

testing gene-trait association in Stage II) using both individual-level and summary-level GWAS

data with the estimated eQTL weights; and (3) Apply ACAT-O to aggregate the TWAS p-values

from individual training methods. Since the existing tools for *P+T*, *lassosum*, *SDPR*, and *PRS-CS*

were originally developed for PRS calculations, we adapted and optimized them for training GReX

imputation models in our OTTERS tool. For example, we integrate TABIX[54] and PLINK[55] tools in

OTTERS to extract input data per target gene more efficiently. We also enable parallel

computation in OTTERS for training GReX imputation models and testing gene-trait association

of multiple genes.

The OTTERS framework does have its limitations. First, training GReX imputation models

by all individual methods on average cost ~20 minutes for all 5 training models per gene, which

might be computationally challenging for studying eQTL summary data of multiple tissue types

and for ~20K genome-wide genes. Users might consider prioritize *P+T(0.001)*, *lassosum,* and

SDPR training methods as these three provide complementary results in our studies. Second, the

currently available eQTL summary statistics are mainly derived from individuals of European

descent. Our OTTERS trained GReX imputations model based on these eQTL summary

statistics, and the resulting imputed GReX could consequently have attenuated cross-population

predictive performance[56]. This might limit the transferability of our TWAS results across

populations. Third, our OTTERS cannot provide the direction of the identified gene-phenotype

associations, which should be referred to the sign of the TWAS Z-score statistic per training

method. Last, even though the method applies to integrate both cis- and trans- eQTL with GWAS

data, the computation time and availability of summary-level trans-eQTL reference data are still

the main obstacles. Our current OTTERS tool only considers cis-eQTL effects. Extension of

OTTERS to enable cross-population TWAS and incorporation of trans-eQTL effects is part of our

ongoing research but out of the scope of this work.

Our novel OTTERS framework using large-scale eQTL summary data has the potential to

identify more significant TWAS risk genes than standard TWAS tools that use smaller individual-

level reference transcriptomic data and use only a single regression method for training GReX

imputation models. This tool provides the opportunity to leverage not only available public eQTL

summary data of various tissues for conducting TWAS of complex traits and diseases, but also

the emerging summary-level data of other types of molecular QTL such as splicing QTLs,

methylation QTLs, metabolomics QTLs, and protein QTLs. For example, OTTERS could be

applied to perform proteome-wide association studies using summary-level reference data of

genetic-protein relationships such as those reported by the SCALLOP consortium[57], and

epigenome-wide association studies using summary-level reference data of methylation-

phenotype relationships reported by Genetics of DNA Methylation Consortium (GoDMC) (see

Web Resources). OTTERS would be most useful for the broad researchers who only have access

to summary-level QTL reference data and summary-level GWAS data. The feasibility of

integrating summary-level molecular QTL data and GWAS data makes our OTTERS tool valuable

for wide application in current multi-omics studies of complex traits and diseases.

Methods

Traditional Two-Stage TWAS Analysis

Stage I of TWAS estimates a GReX imputation model using individual-level expression

and genotype data available from a reference dataset. Consider the following GReX imputation

model from n individuals and m SNPs (multivariable regression model assuming linear additive

genetic effects) within the reference dataset:

$$E_g = Xw + \epsilon, \ \epsilon \sim N(0, \sigma_\epsilon^2 I). \qquad \text{(Equation 1)}$$

Here, E_g is a vector representing gene expression levels of gene g, X is an $n \times m$ matrix of

genotype data of SNP predictors proximal or within gene g, w is a vector of genetic effect sizes

(referred to as a broad sense of eQTL effect sizes), and ϵ is the error term. Here, we consider

only cis-SNPs within 1 MB of the flanking 5' and 3' ends as genotype predictors that are coded

within X[19,20,22]. Once we configure the model in Equation 1, we can employ methods like

PrediXcan, FUSION, and TIGAR to fit the model and obtain estimates of eQTL effect sizes ($\hat{w}$).

Stage II of TWAS uses the eQTL effect sizes ($\hat{w}$) from Stage I to impute gene expression

(GReX) in an independent GWAS and then test for association between GReX and phenotype.

Given individual-level GWAS data with genotype data X_{new} and eQTL effect sizes ($\hat{w}$) from Stage

I, the GReX for X_{new} can be imputed by $\widehat{GReX} = X_{new} \, \hat{w}$. The follow-up TWAS would test the

association between $\widehat{GReX}$ and phenotype Y based on a generalized linear regression model,

which is equivalent to a gene-based association test taking $\hat{w}$ as test SNP weights. When

individual-level GWAS data are not available, one can apply FUSION and S-PrediXcan test

statistics to summary-level GWAS data as follows:

$$Z_{g,FUSION} = \frac{\sum_{j=1}^{J}(\hat{w}_j Z_j)}{\sqrt{\hat{w}' V \hat{w}}}, \quad Z_{g,S-PrediXcan} = \frac{\sum_{j=1}^{J}(\hat{w}_j \hat{\sigma}_j Z_j)}{\sqrt{\hat{w}' V \hat{w}}} \quad \text{(Equation 2)}$$

where Z_j is the single variant *Z-score* test statistic in GWAS for the j^{th} SNP, $j = 1, \dots, J$, for all test

SNPs that have both eQTL weights with respect to the test gene g and GWAS Z-scores; $\hat{\sigma}_j$ is

the genotype standard deviation of the j^{th} SNP; and V denotes the genotype correlation matrix in

FUSION Z-score statistic and genotype covariance matrix in S-PrediXcan Z-score statistic of the

test SNPs. In particular, $\hat{\sigma}_j$ and V can be approximated from a reference panel with genotype

data of samples of the same ancestry such as those available from the 1000 Genomes

Project[58]. If $\hat{w}$ are standardized effect sizes estimated assuming standardized genotype X and

gene expression E_g in Equation 1, FUSION and S-PrediXcan Z-score statistics are equivalent[13].

Otherwise, the S-PrediXcan Z-score should be applied to avoid false positive inflation.

TWAS Stage I Analysis using Summary-Level Reference Data

We now consider a variation of TWAS Stage I to estimate cis-eQTL effect sizes $\hat{w}$ based

on a multivariable regression model (Equation 1) from summary-level reference data. We assume

that the summary-level reference data provide information on the association between a single

genetic variant j ($j = 1, \dots, m$) and expression of gene *g*. This information generally consists of

effect size estimates ($\tilde{w}_j, j = 1, \dots, m$) and p-values derived from the following single variant

regression models:

508
$$E_g = X_j w_j + \epsilon, \ \epsilon \sim N(0, \sigma_\epsilon^2 I), j \ = \ 1, \dots, m. \quad \text{(Equation 3)}$$

Here, X_j is an $n \times 1$ vector of genotype data for genetic variant j. Since eQTL summary data are

analogous to GWAS summary data where gene expression represents the phenotype, we can

estimate the eQTL effect sizes $\hat{w}$ using marginal least squared effect estimates ($\tilde{w}_j, j = 1, \dots, m$)

and p-values from the QTL summary data as well as reference linkage disequilibrium (LD)

information of the same ancestry[26–29]. Although all PRS methods apply to the TWAS Stage I

framework, we only consider four representative methods as follows:

P+T: The *P+T* method selects eQTL weights by LD-clumping and P-value Thresholding[26].

Given threshold P_T for p-values and threshold R_T for LD R^2, we first exclude SNPs with marginal

p-values from eQTL summary data greater than P_T or strongly correlated (LD R^2 greater than R_T)

with another SNP having a more significant marginal p-value (or Z-score statistic value). For the

remaining selected test SNPs, we use marginal standardized eQTL effect sizes from eQTL

summary data as eQTL weights for TWAS in Stage II. We considered $R_T = 0.99$ and $P_T =$

$(0.001, 0.05)$ in this paper and implemented the *P+T* method using PLINK 1.9[55] (see Web

Resources). We denote the *P+T* method with P_T equal to 0.001 and 0.05 as *P+T (0.001)* and

P+T (0.05), respectively.

Frequentist lassosum: With standardized $\boldsymbol{E_g}$ and **X**, we can show that the marginal least

squared eQTL effect size estimates from the single variant regression model (Equation 3) is $\widetilde{\boldsymbol{w}} =$

$\boldsymbol{X}^T\boldsymbol{E_g}/n$ and that the LD correlation matrix is $\boldsymbol{R} = \boldsymbol{X}^T\boldsymbol{X}/n$. That is,

527                              $$\boldsymbol{X}^T\boldsymbol{E_g} = n\widetilde{\boldsymbol{w}} \text{ and } \boldsymbol{X}^T\boldsymbol{X} = n\boldsymbol{R}. \quad \text{(Equation 4)}$$

By approximating $n\boldsymbol{R}$ by $n\boldsymbol{R_s}$ ($\boldsymbol{R_s} = (1-s)\boldsymbol{R_r} + s\boldsymbol{I}$ with a tuning parameter $0 < s < 1$, a reference

LD correlation matrix $\boldsymbol{R_r}$ from an external panel such as one from the 1000 Genomes Project[58],

and an identity matrix $\boldsymbol{I}$) in the LASSO[32] penalized loss function, the frequentist *lassosum*

method[27] can tune the LASSO penalty parameter and *s* using a pseudovalidation approach and

then solve for eQTL effect size estimates $\widehat{\boldsymbol{w}}$ by minimizing the approximated LASSO loss function

requiring no individual-level data (see details in Supplemental Methods).

Bayesian SDPR: Bayesian DPR method[33] as implemented in TIGAR[22] estimates $\widehat{\boldsymbol{w}}$ for the

underlying multivariable regression model in Equation 1 by assuming a normal prior $N(0, \sigma_w^2)$ for

w_j and a Dirichlet process prior[59] $DP(H, \alpha)$ for σ_w^2 with base distribution H and concentration

parameter α. *SDPR*[29] assumes the same DPR model but can be applied to estimate the eQTL

effect sizes $\widehat{\boldsymbol{w}}$ using only eQTL summary data (see details in Supplemental Methods).

Bayesian PRS-CS: The *PRS-CS* method[28] assumes the following normal prior for w_j and

non-informative scale-invariant Jeffreys prior on the residual variance σ_ϵ^2 in Equation 1

$$w_j \sim N\left(0, \frac{\sigma_\epsilon^2}{n}\psi_j\right), \;\; p(\sigma_\epsilon^2) \propto \sigma_\epsilon^2; \;\; \psi_j \sim Gamma(a, \delta_j), \delta_j \sim Gamma(b, \phi),$$

where local shrinkage parameter ψ_j has an independent gamma-gamma prior and ϕ is a global-

shrinkage parameter controlling the overall sparsity of $\boldsymbol{w}$. *PRS-CS* sets hyper parameters $a = 1$

and $b = 1/2$ to ensure the prior density of w_j to have a sharp peak around zero to shrink small

effect sizes of potentially false eQTL towards zero, as well as heavy, Cauchy-like tails which

asserts little influence on eQTLs with larger effects. Posterior estimates $\hat{w}$ will be obtained from

eQTL summary data (i.e., marginal effect size estimates $\tilde{w}$ and p-values) and reference LD

correlation matrix R by Gibbs Sampler (see details in Supplemental Methods). We set ϕ as the

square of the proportion of causal variants in the simulation and as 10^{-4} per gene in the real data

application.

OTTERS Framework

As shown in Figure 1, OTTERS first trains GReX imputation models per gene *g* using *P+T,*

lassosum, SDPR, and *PRS-CS* methods that each infers cis-eQTLs weights using cis-eQTL

summary data and an external LD reference panel of similar ancestry (Stage I). Once we derive

cis-eQTLs weights for each training method, we can impute the respective GReX using that

method and perform the respective gene-based association analysis in the test GWAS dataset

using the formulas given in Equation 2 (Stage II). We thus derive a set of TWAS p-values for gene

g; one p-value for each training model that we applied. We then use these TWAS p-values to

create an omnibus test using the ACAT-O[34] approach that employs a Cauchy distribution for

inference (see details in Supplemental Methods). We refer to the p-value derived from ACAT-O

test as the OTTERS p-value.

Marginal eQTL Effect Sizes

In practice of training GReX imputation models using reference eQTL summary data, the

marginal standardized eQTL effect sizes were approximated by $\tilde{w}_j \approx Z_j/\sqrt{median(n_{g,j})}$, where

Z_j denotes the corresponding eQTL Z-score statistic value by single variant test and

$median(n_{g,j})$ denotes the median sample size of all cis-eQTLs for the target gene g. The median

cis-eQTL sample size per gene was also taken as the sample size value required by *lassosum,*

SDPR, and *PRS-CS* methods, for robust performance. Since summary eQTL datasets (e.g.,

eQTLGen) were generally obtained by meta-analysis of multiple cohorts, the sample size per test

SNP could vary across all cis-eQTLs of the test gene. The median cis-eQTL sample size ensures

a robust performance for applying those eQTL summary data based methods.

LD Clumping

We performed LD-clumping with R_T=0.99 for all individual methods in both simulation and

real studies. Using PRS-CS as an example, we also showed that LD-clumping does not affect the

GReX imputation accuracy compared to no clumping in the real data testing (Figure S12).

LD Blocks for lassosum, PRS-CS, and SDPR

LD blocks were determined externally by ldetect[60] for *lassosum* and *PRS-CS*, while

internally for *SDPR* which ensure that SNPs in one LD block do not have nonignorable correlation

($R^2 > 0.1$) with SNPs in other blocks.

Simulate GWAS Z-score

Given gene expression $\boldsymbol{E_g}$ simulated from the multivariate regression model $\boldsymbol{E_g} = \boldsymbol{X_g}\boldsymbol{w} +$

$\boldsymbol{\epsilon_g}$ with standardized genotype matrix $\boldsymbol{X_g}$ and $\boldsymbol{\epsilon_g} \sim N(0, (1 - h_e^2)\boldsymbol{I}$, we assume GWAS phenotype

data of n_{gwas} samples are simulated from the following linear regression model

584
$$\boldsymbol{Y} = h_p(\boldsymbol{X_g}\boldsymbol{w}) + \boldsymbol{\epsilon_p}, \qquad \boldsymbol{\epsilon_p} \sim N(0, \boldsymbol{I}).$$

585   Conditioning on true genetic effect sizes, the GWAS Z-score test statistics of all test SNPs will

586   follow a multivariate normal distribution, $MVN\left(\boldsymbol{\Sigma_g}\boldsymbol{w}\sqrt{n_{gwas}h_p^2}, \boldsymbol{\Sigma_g}\right)$, where $\boldsymbol{\Sigma_g}$ is the correlation

587   matrix of the standardized genotype X_g from test samples, and h_p^2 denotes the amount of

588   phenotypic variance explained by simulated GReX=$\boldsymbol{X_g}\boldsymbol{w}$[38]. Thus, for a given GWAS sample size,

589   we can generate GWAS Z-score statistic values from this multivariate normal distribution.

 FUSION using Individual-level Reference Data

To train GReX imputation models by FUSION with individual-level reference data, we

trained Best Linear Unbiased Predictor (BLUP) model[61], Elastic-net regression[62], LASSO

regression[32], and single best eQTL model as implemented in the FUSION tool (see Web

Resource). Default settings were used to train GReX imputation models by FUSION in our

simulation and real studies. LASSO regression was performed only for genes with positive

estimated expression heritability. The eQTL weights of the best trained GReX imputation model

will be used to conduct TWAS by FUSION.

GTEx V8 Dataset

GTEx V8 dataset (dbGaP phs000424.v8.p2) contains comprehensive profiling of WGS

genotype data and RNA-sequencing (RNA-seq) transcriptomic data across 54 human tissue

types of 838 donors. The GTEx V8 WGS genotype data of all samples were used to construct

reference LD in our studies. The GTEx V6 RNA-seq data of whole blood samples were used to

train GReX imputation models by FUSION, and the GTEx V8 RNA-seq data of additional whole

blood samples (n=315) were used to test GReX imputation accuracy in our studies. GTEx V8

RNA-seq data of all whole blood samples (n=574) were also used as reference data for

comparing the performance of OTTERS and FUSION.

eQTLGen Consortium Dataset

The eQTLGen consortium[23] dataset was generated based on meta-analysis across 37

individual cohorts (n=31,684) including GTEx V6 as a sub-cohort. eQTLGen samples consist of

25,482 blood (80.4%) and 6,202 peripheral blood mononuclear cell (19.6%) samples. We

considered SNPs with minor allele frequency (MAF) > 0.01, Hardy–Weinberg P value >0.0001,

call rate >0.95, genotype imputation r^2 > 0.5 and observed in at least 2 cohorts[23]. We only

considered cis-eQTL (within $\pm$1MB around gene transcription start sites (TSS)) with test sample

size > 3000. As a result, we used cis-eQTL summary data of 16,699 genes from eQTLGen to

train GReX imputation models for use in OTTERS in this study.

<u>UK Biobank GWAS Data of Cardiovascular Disease</u>

Summary-level GWAS data of Cardiovascular Disease from UKBB (n=459,324, case

fraction = 0.319)[35] were generated by BOLT-LMM based on the Bayesian linear mixed model per

SNP[63] with assessment centered, sex, age, and squared age as covariates. Although BOLT-LMM

was derived based on a quantitative trait model, it can be applied to analyze case-control traits

and has well-controlled false positive rate when the trait is sufficiently balanced with case fraction

$\geq$ 10% and samples are of the same ancestry . The tested dichotomous cardiovascular disease

phenotype includes a list of sub-phenotypes: hypertension, heart/cardiac problem, peripheral

vascular disease, venous thromboembolic disease, stroke, transient ischaemic attack (tia),

subdural haemorrhage/haematoma, cerebral aneurysm, high cholesterol, and other

venous/lymphatic disease.

Data Availability

eQTLGen consortium data are available from their portal website (https://www.eqtlgen.org). UK

Biobank summary-level GWAS data are available through the Alkes Group

(https://alkesgroup.broadinstitute.org/UKBB). Individual-level GTEx reference data are available

through dbGap (Accession phs000424.v8.p2). ROS/MAP/MSBB WGS data used in our

simulation studies are available through Synapse with data access application

(https://www.synapse.org/#!Synapse:syn10901595). All source code and scripts used in this

study are available through OTTERS Github page (https://github.com/daiqile96/OTTERS).

Ethics Approval

All data used in this study are de-identified genotype data and summary level eQTL and GWAS data. ROS/MAP genotype data were collected with ethics approval from the IRB at Rush University and all participants consented to participate.

Acknowledgements

The authors thank Dr. Greg Gibson from Georgia Tech for his insightful comments and discussion that help the development and improve the quality of this manuscript. This work was supported by National Institutes of Health grant awards R35GM138313 (QD, JY), RF1AG071170 (QD, MPE), and Estonian Research Council Grant PUT (PRG1291) for TE. NIH/NIA grants P30AG10161, R01AG15819, R01AG17917, R01AG30146, R01AG36836, R01AG56352, U01AG32984, U01AG46152, U01AG61356, the Illinois Department of Public Health, the Translational Genomics Research Institute support the generation of the ROS/MAP data led by ASB, PLDJ and DAB. The Young Finns Study has been financially supported by the Academy of Finland: grants 322098, 286284, 134309 (Eye), 126925, 121584, 124282, 255381, 256474, 283115, 319060, 320297, 314389, 338395, 330809, and 104821, 129378 (Salve), 117797 (Gendi), and 141071 (Skidi), the Social Insurance Institution of Finland, Competitive State Research Financing of the Expert Responsibility area of Kuopio, Tampere and Turku University Hospitals (grant X51001), Juho Vainio Foundation, Paavo Nurmi Foundation, Finnish Foundation for Cardiovascular Research, Finnish Cultural Foundation, The Sigrid Juselius Foundation, Tampere Tuberculosis Foundation, Emil Aaltonen Foundation, Yrjö Jahnsson Foundation, Signe and Ane Gyllenberg Foundation, Diabetes Research Foundation of Finnish Diabetes Association, EU Horizon 2020 (grant 755320 for TAXINOMISIS and grant 848146 for To Aition), European Research Council (grant 742927 for MULTIEPIGEN project), Tampere University Hospital Supporting Foundation, and Finnish Society of Clinical Chemistry and the Cancer Foundation Finland. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Authors' contributions

QD conducted data analysis and drafted the manuscript; JY and MPE conceptualized and led the

project, and edited the manuscript; GZ and HZ consulted data analysis and edited the manuscript;

UV, LF, AB, AT, TL, OR, TE, contributed reference eQTL summary data and edited the

manuscript; eQTLGen Consortium contributed eQTLGen summary data.

Competing interests

The authors declare no competing interests and consent for publication.

Materials & Correspondence

Correspondence and material requests should be sent to M.P.E. (mpepste@emory.edu) and

J.Y. (jingjing.yang@emory.edu).

Description of Supplemental Data

Supplemental data include 12 figures, 4 tables, and supplemental methods.

Web Resources

ACAT: https://github.com/yaowuliu/ACAT

eQTLGen: https://www.eqtlgen.org

FUSION: http://gusevlab.org/projects/fusion

GoDMC: http://www.godmc.org.uk

lassosum: https://github.com/tshmak/lassosum

OTTERS: https://github.com/daiqile96/OTTERS

PLINK: https://www.cog-genomics.org/plink

PRS-CS: https://github.com/getian107/PRScs

682   SDPR: https://github.com/eldronzhou/SDPR

683   TWAS hub: http://twas-hub.org

686 **Tables**

687 **Table 1. Test R^2 in 315 whole blood tissue samples from GTEx V8.**

	P+T(0.001)	P+T(0.05)	lassosum	SDPR	PRS-CS	FUSION[b]
# Genes with $R^2 > 0.01$	9,816	9,662	8,718	9,670	10,337	4,704
Median R^{2}[a]	0.044	0.0430	0.0416	0.0418	0.0517	0.0367

a: Median R^2 among genes with test $R^2 > 0.01$ per method.

b: FUSION was trained on GTEx V6 blood samples while all other training methods were trained using eQTLGen summary statistics (n=31,684) and reference LD from GTEx V8 samples.

Table 2. Independent TWAS risk genes of cardiovascular disease identified by OTTERS.
Reference eQTL summary data from eQTLGen consortium and GWAS summary data from
UKBB were used. The corresponding TWAS p-values by 5 individual PRS methods and
OTTERS are shown in the table with significant p-values in bold, and those for genes with test
GReX $R^2 \leq 0.01$ were shown as a dash.

CHROM	ID	OTTERS	P+T(0.001)	P+T(0.05)	lassosum	SDPR	PRS-CS
1	CLCN6[a]	**5.75E-15**	**4.94E-09**	**5.40E-08**	**8.77E-09**	**1.19E-15**	**1.43E-09**
1	NPPA[b]	**4.32E-08**	**1.55E-08**	**2.14E-07**	-	-	6.71E-06
1	PSRC1[a]	**8.37E-20**	**5.68E-08**	**8.46E-07**	**6.26E-11**	**1.67E-20**	**1.41E-12**
2	RP11-378A13.1[a]	**9.78E-09**	3.97E-02	4.98E-02	1.62E-05	**1.96E-09**	1.15E-04
4	LINC01093[c]	**2.57E-09**	9.85E-02	5.31E-02	**5.13E-10**	1.08E-02	2.41E-02
5	CPEB4[b]	**3.05E-14**	1.26E-02	2.05E-02	2.70E-05	**6.05E-15**	**1.60E-07**
6	SERPINB6[c]	**1.47E-07**	2.12E-01	2.24E-01	7.56E-03	**2.95E-08**	7.53E-04
6	CARMIL1[c]	**9.23E-09**	5.34E-03	3.41E-03	4.15E-03	**1.85E-09**	1.72E-03
6	ZSCAN12P1[c]	**1.84E-08**	6.00E-01	5.75E-01	4.62E-01	**3.67E-09**	3.10E-01
6	HCG4P7[c]	**8.93E-50**	3.70E-01	3.69E-01	2.30E-01	**1.79E-50**	7.26E-01
6	HCG4P3[c]	**5.33E-20**	4.20E-01	4.05E-01	5.03E-04	**1.07E-20**	2.42E-03
6	HLA-S[c]	**4.57E-07**	7.13E-01	7.31E-01	3.02E-01	**9.14E-08**	2.33E-01
7	PSPHP1[c]	**1.21E-09**	2.17E-01	2.26E-01	9.65E-03	**2.43E-10**	1.10E-01
8	LPL[c]	**5.73E-07**	1.78E-03	3.26E-03	4.44E-02	**1.15E-07**	1.05E-04
8	PTP4A3[c]	**1.28E-06**	8.13E-02	8.33E-02	6.23E-05	**2.58E-07**	1.67E-03
10	CAMK1D[a]	**2.51E-09**	3.83E-02	4.97E-02	1.23E-03	**5.03E-10**	4.97E-05
10	NT5C2[b]	**1.21E-07**	**1.69E-06**	**2.92E-06**	1.64E-05	**3.15E-07**	**2.69E-08**
11	TNNT3[b]	**1.67E-10**	**1.09E-06**	3.33E-06	**2.03E-09**	**3.40E-11**	**4.01E-07**
11	C11orf49[b]	**2.28E-06**	**8.55E-07**	**1.78E-06**	5.44E-05	-	2.93E-04
11	SIDT2[a]	**7.26E-09**	6.14E-05	1.33E-04	3.66E-05	**1.46E-09**	**3.81E-07**
15	CSK[b]	**2.30E-09**	**1.70E-07**	**2.15E-06**	**7.41E-10**	**2.80E-09**	**2.17E-09**
15	FES[b]	**2.87E-32**	**4.78E-08**	**1.23E-06**	**9.13E-24**	**5.75E-33**	**1.94E-15**
15	SLCO3A1[c]	**3.78E-08**	1.85E-02	3.15E-02	4.65E-05	**7.57E-09**	1.14E-03
16	MBTPS1[b]	**5.80E-08**	2.62E-01	3.05E-01	9.15E-04	**1.16E-08**	2.34E-03
16	MTHFSD[a]	**4.65E-07**	5.16E-02	5.94E-02	1.65E-02	**9.30E-08**	3.20E-03
17	ACE[b]	**9.42E-07**	**4.93E-06**	**1.03E-05**	**4.23E-06**	**9.66E-07**	**2.68E-07**
18	RALBP1[c]	**1.40E-06**	1.48E-01	1.54E-01	2.12E-04	**2.81E-07**	5.55E-03
19	MRI1[b]	**8.38E-09**	8.34E-03	1.60E-02	7.79E-03	**1.68E-09**	2.65E-03
19	HAUS8[b]	**1.60E-07**	**4.41E-08**	**1.38E-07**	**1.67E-06**	**1.42E-06**	3.29E-05
19	SULT2B1[c]	**2.32E-06**	**7.73E-07**	-	-	2.97E-02	1.10E-02
19	NTN5[a]	**9.03E-10**	**2.75E-08**	**1.16E-07**	6.23E-06	**1.85E-10**	**9.73E-09**
19	RPL28[b]	**3.76E-07**	7.33E-02	1.16E-01	6.64E-03	**7.52E-08**	4.23E-03
20	CTSZ[b]	**3.32E-09**	2.57E-02	1.99E-02	**3.40E-09**	**8.25E-10**	1.04E-01
20	EDN3[c]	**1.29E-07**	**3.61E-08**	**9.15E-08**	8.60E-06	5.90E-03	1.58E-02
20	ZBTB46[c]	**1.07E-06**	**2.83E-07**	8.35E-06	-	1.81E-03	1.27E-05
20	OPRL1[a]	**5.84E-07**	**3.44E-07**	**2.69E-06**	1.85E-03	5.51E-05	**1.90E-07**
21	FAM3B[c]	**1.08E-10**	2.28E-02	2.58E-02	8.07E-06	**2.17E-11**	1.04E-05
21	MX1[c]	**6.04E-22**	4.36E-01	3.83E-01	**3.16E-07**	**1.21E-22**	1.24E-03

a: Risk gene of UKBB cardiovascular disease in TWAS-hub identified using GTEx whole blood tissue.

b: Risk genes of UKBB cardiovascular disease in TWAS-hub identified using other GTEx tissue types.

c: Novel risk gene

 Figure Titles and Legends

Figure 1. OTTERS framework.

OTTERS estimates cis-eQTL weights from eQTL summary data and reference LD panel using

four imputation models (Stage I), and conducts ACAT-O test to combine gene-based

association test p-values from individual methods with individual/summary level test GWAS data

(Stage II).

Figure 2. Test R^2 (A) and TWAS power (B) comparison in simulation studies

Various proportions of true causal cis-eQTL $p_{causal} = (0.001, 0.01)$ and gene expression

heritability $h_e^2 = (0.01, 0.05, 0.1)$ were considered in the simulation studies. The GWAS sample

size was chosen with respect to h_e^2 values. The proportion of phenotype variance explained by

gene expression (h_p^2) was set to be 0.025. TWAS was conducted using simulated GWAS Z-

scores.

Figure 3. Test R^2 by *PRS-CS* versus *P+T(0.001), P+T(0.05), lassosum, SDPR, FUSION.*

Test R^2 by *PRS-CS* versus *P+T(0.001) (A), P+T(0.05) (B), lassosum (C), SDPR (D), and FUSION*

(E) with 315 GTEx V8 test samples, with different colors denoting whether test $R^2 > 0.01$ only by

PRS-CS (red), only by the y axis method (green), or both methods (blue). Genes with test $R^2 >$

0.01 by at least one method were included in the plot.

Figure 4. Manhattan plot of TWAS results by OTTERS.

Manhattan plot of TWAS results by OTTERS using GWAS summary-level statistics of

cardiovascular disease and imputation models fitted based on eQTLGen summary statistics.

Independently significant TWAS risk genes are labeled.

eQTLGen Consortium Author List

Mawussé Agbessi[1], Habibul Ahsan[2], Isabel Alves[1], Anand Kumar Andiappan[3], Wibowo Arindrarto[4], Philip Awadalla[1], Alexis Battle[5,6], Frank Beutner[7], Marc Jan Bonder[8,9], Dorret I. Boomsma[10], Mark W. Christiansen[11], Annique Claringbould[8,12], Patrick Deelen[8,13,12,14], Tõnu Esko[15], Marie-Julie Favé[1], Lude Franke[8,12], Timothy Frayling[16], Sina A. Gharib[11,17], Greg Gibson[18], Bastiaan T. Heijmans[4], Gibran Hemani[19], Rick Jansen[20], Mika Kähönen[21], Anette Kalnapenkis[15], Silva Kasela[15], Johannes Kettunen[22], Yungil Kim[23,5], Holger Kirsten[24], Peter Kovacs[25], Knut Krohn[26], Jaanika Kronberg[15], Viktorija Kukushkina[15], Zoltan Kutalik[27], Bernett Lee[3], Terho Lehtimäki[28], Markus Loeffler[24], Urko M. Marigorta[18,29,30], Hailang Mei[31], Lili Milani[15], Grant W. Montgomery[32], Martina Müller-Nurasyid[33,34,35], Matthias Nauck[36,37], Michel G. Nivard[38], Brenda Penninx[20], Markus Perola[39], Natalia Pervjakova[15], Brandon L. Pierce[2], Joseph Powell[40], Holger Prokisch[41,42], Bruce M. Psaty[11,43], Olli T. Raitakari[44], Samuli Ripatti[45], Olaf Rotzschke[3], Sina Rüeger[27], Ashis Saha[5], Markus Scholz[24], Katharina Schramm[46,34], Ilkka Seppälä[28], Eline P. Slagboom[4], Coen D.A. Stehouwer[47], Michael Stumvoll[48], Patrick Sullivan[49], Peter A.C. 't Hoen[50], Alexander Teumer[51], Joachim Thiery[52], Lin Tong[2], Anke Tönjes[48], Jenny van Dongen[10], Maarten van Iterson[4], Joyce van Meurs[53], Jan H. Veldink[54], Joost Verlouw[53], Peter M. Visscher[32], Uwe Völker[55], Urmo Võsa[8,15], Harm-Jan Westra[8,12], Cisca Wijmenga[8], Hanieh Yaghootkar[16,56,57], Jian Yang[32,58], Biao Zeng[18], Futao Zhang[32]

Author list is ordered alphabetically

1. Computational Biology, Ontario Institute for Cancer Research, Toronto, Canada
2. Department of Public Health Sciences, University of Chicago, Chicago, United States of America
3. Singapore Immunology Network, Agency for Science, Technology and Research, Singapore, Singapore
4. Leiden University Medical Center, Leiden, The Netherlands
5. Department of Computer Science, Johns Hopkins University, Baltimore, United States of America
6. Departments of Biomedical Engineering, Johns Hopkins University, Baltimore, United States of America
7. Heart Center Leipzig, Universität Leipzig, Leipzig, Germany
8. Department of Genetics, University Medical Centre Groningen, Groningen, The Netherlands
9. European Molecular Biology Laboratory, Genome Biology Unit, 69117 Heidelberg, Germany
10. Netherlands Twin Register, Department of Biological Psychology, Vrije Universiteit Amsterdam, Amsterdam Public Health research institute and Amsterdam Neuroscience, the Netherlands
11. Cardiovascular Health Research Unit, University of Washington, Seattle, United States of America
12. Oncode Institute
13. Genomics Coordination Center, University Medical Centre Groningen, Groningen, The Netherlands
14. Department of Genetics, University Medical Centre Utrecht, P.O. Box 85500, 3508 GA, Utrecht, The Netherlands
15. Estonian Genome Center, Institute of Genomics, University of Tartu, Tartu 51010, Estonia
16. Genetics of Complex Traits, University of Exeter Medical School, Royal Devon & Exeter Hospital, Exeter, United Kingdom
17. Department of Medicine, University of Washington, Seattle, United States of America
18. School of Biological Sciences, Georgia Tech, Atlanta, United States of America

19. MRC Integrative Epidemiology Unit, University of Bristol, Bristol, United Kingdom

20. Amsterdam UMC, Vrije Universiteit, Department of Psychiatry, Amsterdam Public Health research institute and Amsterdam Neuroscience, The Netherlands

21. Department of Clinical Physiology, Tampere University Hospital and Faculty of Medicine and Health Technology, Tampere University, Tampere, Finland

22. University of Helsinki, Helsinki, Finland

23. Genetics and Genomic Science Department, Icahn School of Medicine at Mount Sinai, New York, United States of America

24. Institut für Medizinische InformatiK, Statistik und Epidemiologie, LIFE – Leipzig Research Center for Civilization Diseases, Universität Leipzig, Leipzig, Germany

25. IFB Adiposity Diseases, Universität Leipzig, Leipzig, Germany

26. Interdisciplinary Center for Clinical Research, Faculty of Medicine, Universität Leipzig, Leipzig, Germany

27. Lausanne University Hospital, Lausanne, Switzerland

28. Department of Clinical Chemistry, Fimlab Laboratories and Finnish Cardiovascular Research Center-Tampere, Faculty of Medicine and Health Technology, Tampere University, Tampere, Finland

29. Integrative Genomics Lab, CIC bioGUNE, Bizkaia Science and Technology Park, Derio, Bizkaia, Basque Country, Spain

30. IKERBASQUE, Basque Foundation for Science, Bilbao, Spain

31. Department of Medical Statistics and Bioinformatics, Leiden University Medical Center, Leiden, The Netherlands

32. Institute for Molecular Bioscience, University of Queensland, Brisbane, Australia

33. Institute of Genetic Epidemiology, Helmholtz Zentrum München - German Research Center for Environmental Health, Neuherberg, Germany

34. Department of Medicine I, University Hospital Munich, Ludwig Maximilian's University, München, Germany

35. DZHK (German Centre for Cardiovascular Research), partner site Munich Heart Alliance, Munich, Germany

36. Institute of Clinical Chemistry and Laboratory Medicine, Greifswald University Hospital, Greifswald, Germany

37. German Center for Cardiovascular Research (partner site Greifswald), Greifswald, Germany

38. Department of Biological Psychology, Faculty of Behaviour and Movement Sciences, VU, Amsterdam, The Netherlands

39. National Institute for Health and Welfare, University of Helsinki, Helsinki, Finland

40. Garvan Institute of Medical Research, Garvan-Weizmann Centre for Cellular Genomics, Sydney, Australia

41. Institute of Neurogenomics, Helmholtz Zentrum München, Neuherberg, Germany

42. Institute of Human Genetics, Technical University Munich, Munich, Germany

43. Kaiser Permanente Washington Health Research Institute, Seattle, WA, United States of America

44. Centre for Population Health Research, Department of Clinical Physiology and Nuclear Medicine, Turku University Hospital and University of Turku, Turku, Finland

45. Statistical and Translational Genetics, University of Helsinki, Helsinki, Finland

46. Institute of Genetic Epidemiology, Helmholtz Zentrum München - German Research Center for Environmental Health, Neuherberg, Germany

47. Department of Internal Medicine and School for Cardiovascular Diseases (CARIM), Maastricht University Medical Center, Maastricht, The Netherlands

48. Department of Medicine, Universität Leipzig, Leipzig, Germany

49. Department of Medical Epidemiology and Biostatistics, Karolinska Institutet, Stockholm, Sweden

50. Center for Molecular and Biomolecular Informatics, Radboud Institute for Molecular Life
Sciences, Radboud University Medical Center Nijmegen, Nijmegen, The Netherlands
51. Institute for Community Medicine, University Medicine Greifswald, Greifswald, Germany
52. Institute for Laboratory Medicine, LIFE – Leipzig Research Center for Civilization Diseases,
Universität Leipzig, Leipzig, Germany
53. Department of Internal Medicine, Erasmus Medical Centre, Rotterdam, The Netherlands
54. UMC Utrecht Brain Center, University Medical Center Utrecht, Department of Neurology,
Utrecht University, Utrecht, The Netherlands
55. Interfaculty Institute for Genetics and Functional Genomics, University Medicine Greifswald,
Greifswald, Germany
56. School of Life Sciences, College of Liberal Arts and Science, University of Westminster, 115
New Cavendish Street, London, United Kingdom
57. Division of Medical Sciences, Department of Health Sciences, Luleå University of Technology,
Luleå, Sweden
58. Institute for Advanced Research, Wenzhou Medical University, Wenzhou, Zhejiang 325027,
China
832

References

1. Mancuso, N. *et al.* Integrating Gene Expression with Summary Association Statistics to Identify Genes Associated with 30 Complex Traits. *Am J Hum Genet* **100**, 473–487 (2017).

2. Gusev, A. *et al.* Transcriptome-wide association study of schizophrenia and chromatin activity yields mechanistic disease insights. *Nat Genet* **50**, 538–548 (2018).

3. Mancuso, N. *et al.* Large-scale transcriptome-wide association study identifies new prostate cancer risk regions. *Nat Commun* **9**, 4079 (2018).

4. Wainberg, M. *et al.* Opportunities and challenges for transcriptome-wide association studies. *Nat Genet* **51**, 592–599 (2019).

5. Strunz, T., Lauwen, S., Kiel, C., Hollander, A. den & Weber, B. H. F. A transcriptome-wide association study based on 27 tissues identifies 106 genes potentially relevant for disease pathology in age-related macular degeneration. *Sci Rep* **10**, 1584 (2020).

6. Raj, T. *et al.* Integrative transcriptome analyses of the aging brain implicate altered splicing in Alzheimer's disease susceptibility. *Nat Genet* **50**, 1584–1592 (2018).

7. Hao, S., Wang, R., Zhang, Y. & Zhan, H. Prediction of Alzheimer's Disease-Associated Genes by Integration of GWAS Summary Data and Expression Data. *Frontiers in Genetics* **9**, (2019).

8. Luningham, J. M. *et al.* Bayesian Genome-wide TWAS Method to Leverage both cis- and trans-eQTL Information through Summary Statistics. *The American Journal of Human Genetics* **107**, 714–726 (2020).

9. Hoffman, J. D. *et al.* Cis-eQTL-based trans-ethnic meta-analysis reveals novel genes associated with breast cancer risk. *PLoS Genet* **13**, e1006690 (2017).

10. Wu, L. *et al.* A transcriptome-wide association study of 229,000 women identifies new candidate susceptibility genes for breast cancer. *Nat Genet* **50**, 968–978 (2018).

11. Bhattacharya, A. *et al.* A framework for transcriptome-wide association studies in breast cancer in

diverse study populations. *Genome Biol* **21**, 42 (2020).

12. Gusev, A. *et al.* A transcriptome-wide association study of high-grade serous epithelial ovarian

cancer identifies new susceptibility genes and splice variants. *Nat Genet* **51**, 815–823 (2019).

13. Parrish, R. L., Gibson, G. C., Epstein, M. P. & Yang, J. TIGAR-V2: Efficient TWAS tool with

nonparametric Bayesian eQTL weights of 49 tissue types from GTEx V8. *Human Genetics and*

Genomics Advances **3**, 100068 (2022).

14. Thériault, S. *et al.* Genetic Association Analyses Highlight IL6, ALPL, and NAV1 As 3 New

Susceptibility Genes Underlying Calcific Aortic Valve Stenosis. *Circulation: Genomic and Precision*

Medicine **12**, e002617 (2019).

15. Zhu, Z. *et al.* Genetic overlap of chronic obstructive pulmonary disease and cardiovascular disease-

related traits: a large-scale genome-wide cross-trait analysis. *Respiratory Research* **20**, 64 (2019).

16. Lonsdale, J. *et al.* The Genotype-Tissue Expression (GTEx) project. *Nat Genet* **45**, 580–585 (2013).

17. THE GTEX CONSORTIUM. The GTEx Consortium atlas of genetic regulatory effects across human

tissues. *Science* **369**, 1318–1330 (2020).

18. Gibbs, J. R. *et al.* Abundant Quantitative Trait Loci Exist for DNA Methylation and Gene Expression in

Human Brain. *PLOS Genetics* **6**, e1000952 (2010).

19. Gamazon, E. R. *et al.* A gene-based association method for mapping traits using reference

transcriptome data. *Nat Genet* **47**, 1091–1098 (2015).

20. Gusev, A. *et al.* Integrative approaches for large-scale transcriptome-wide association studies. *Nat*

Genet **48**, 245–252 (2016).

21. Tang, S. *et al.* Novel Variance-Component TWAS method for studying complex human diseases with

applications to Alzheimer's dementia. *PLOS Genetics* **17**, e1009482 (2021).

22. Nagpal, S. *et al.* TIGAR: An Improved Bayesian Tool for Transcriptomic Data Imputation Enhances

Gene Mapping of Complex Traits. *The American Journal of Human Genetics* **105**, 258–266 (2019).

23. Võsa, U. *et al.* Large-scale cis- and trans-eQTL analyses identify thousands of genetic loci and

polygenic scores that regulate blood gene expression. *Nat Genet* **53**, 1300–1310 (2021).

24. The CommonMind Consortium (CMC) *et al.* Large eQTL meta-analysis reveals differing patterns

between cerebral cortical and cerebellar brain regions. *Sci Data* **7**, 340 (2020).

25. Cao, C. *et al.* Power analysis of transcriptome-wide association study: Implications for practical

protocol choice. *PLOS Genetics* **17**, e1009405 (2021).

26. Purcell, S. M. *et al.* Common polygenic variation contributes to risk of schizophrenia and bipolar

disorder. *Nature* **460**, 748–752 (2009).

27. Mak, T. S. H., Porsch, R. M., Choi, S. W., Zhou, X. & Sham, P. C. Polygenic scores via penalized

regression on summary statistics. *Genetic Epidemiology* **41**, 469–480 (2017).

28. Ge, T., Chen, C.-Y., Ni, Y., Feng, Y.-C. A. & Smoller, J. W. Polygenic prediction via Bayesian regression

and continuous shrinkage priors. *Nat Commun* **10**, 1776 (2019).

29. Zhou, G. & Zhao, H. A fast and robust Bayesian nonparametric method for prediction of complex

traits using summary statistics. *PLoS Genet* **17**, e1009697 (2021).

30. Zhang, Q., Privé, F., Vilhjálmsson, B. & Speed, D. Improved genetic prediction of complex traits from

individual-level data or summary statistics. *Nat Commun* **12**, 4192 (2021).

31. Zhao, Z. *et al.* PUMAS: fine-tuning polygenic risk scores with GWAS summary statistics. *Genome

Biology* **22**, 257 (2021).

32. Tibshirani, R. Regression Shrinkage and Selection Via the Lasso. *Journal of the Royal Statistical

Society: Series B (Methodological)* **58**, 267–288 (1996).

33. Zeng, P. & Zhou, X. Non-parametric genetic prediction of complex traits with latent Dirichlet process

regression models. *Nat Commun* **8**, 456 (2017).

34. Liu, Y. *et al.* ACAT: A Fast and Powerful p Value Combination Method for Rare-Variant Analysis in

Sequencing Studies. *The American Journal of Human Genetics* **104**, 410–421 (2019).

35. Loh, P.-R., Kichaev, G., Gazal, S., Schoech, A. P. & Price, A. L. Mixed-model association for biobank-

scale datasets. *Nat Genet* **50**, 906–908 (2018).

36. Gamazon, E. R. *et al.* A gene-based association method for mapping traits using reference

transcriptome data. *Nat Genet* **47**, 1091–1098 (2015).

37. Li, X. *et al.* Dynamic incorporation of multiple in silico functional annotations empowers rare variant

association analysis of large whole-genome sequencing studies at scale. *Nat Genet* **52**, 969–983

(2020).

38. Feng, H. *et al.* Leveraging expression from multiple tissues using sparse canonical correlation

analysis and aggregate tests improves the power of transcriptome-wide association studies. *PLOS*

Genetics **17**, e1008973 (2021).

39. Wang, T., Ionita-Laza, I. & Wei, Y. Integrated Quantile RAnk Test (iQRAT) for gene-level associations.

arXiv:1910.10102 [stat] (2020).

40. Bennett, D. A., Schneider, J. A., Arvanitakis, Z. & Wilson, R. S. OVERVIEW AND FINDINGS FROM THE

RELIGIOUS ORDERS STUDY. *Curr Alzheimer Res* **9**, 628–645 (2012).

41. Bennett, D. A. *et al.* Religious Orders Study and Rush Memory and Aging Project. *J Alzheimers Dis* **64**,

S161–S189 (2018).

42. Wang, M. *et al.* The Mount Sinai cohort of large-scale genomic, transcriptomic and proteomic data

in Alzheimer's disease. *Sci Data* **5**, 180185 (2018).

43. Bhattacharya, A., Li, Y. & Love, M. I. MOSTWAS: Multi-Omic Strategies for Transcriptome-Wide

Association Studies. *PLOS Genetics* **17**, e1009398 (2021).

44. Battle, A., Brown, C. D., Engelhardt, B. E. & Montgomery, S. B. Genetic effects on gene expression

across human tissues. *Nature* **550**, 204–213 (2017).

45. Devlin, B., Roeder, K. & Wasserman, L. Genomic Control, a New Approach to Genetic-Based

Association Studies. *Theoretical Population Biology* **60**, 155–166 (2001).

46. Fuchs, F. D. & Whelton, P. K. High Blood Pressure and Cardiovascular Disease. *Hypertension* **75**, 285–

292 (2020).

47. Masaki, T. The endothelin family: an overview. *J Cardiovasc Pharmacol* **35**, S3-5 (2000).

48. Xue, H., Pan, W. & Initiative, for the A. D. N. Some statistical consideration in transcriptome-wide

association studies. *Genetic Epidemiology* **44**, 221–232 (2020).

49. Liu, A. E. & Kang, H. M. Meta-imputation of transcriptome from genotypes across multiple datasets

by leveraging publicly available summary-level data. *PLOS Genetics* **18**, e1009571 (2022).

50. Yang, Y. *et al.* CoMM-S2: a collaborative mixed model using summary statistics in transcriptome-

wide association studies. *Bioinformatics* **36**, 2009–2016 (2020).

51. Yuan, Z. *et al.* Testing and controlling for horizontal pleiotropy with probabilistic Mendelian

randomization in transcriptome-wide association studies. *Nat Commun* **11**, 3861 (2020).

52. Bulik-Sullivan, B. K. *et al.* LD Score regression distinguishes confounding from polygenicity in

genome-wide association studies. *Nat Genet* **47**, 291–295 (2015).

53. Zhu, Z. *et al.* Integration of summary data from GWAS and eQTL studies predicts complex trait gene

targets. *Nat Genet* **48**, 481–487 (2016).

54. Li, H. Tabix: fast retrieval of sequence features from generic TAB-delimited files. *Bioinformatics* **27**,

718–719 (2011).

55. Chang, C. C. *et al.* Second-generation PLINK: rising to the challenge of larger and richer datasets.

Gigascience **4**, 7 (2015).

56. Keys, K. L. *et al.* On the cross-population generalizability of gene expression prediction models. *PLOS*

Genetics **16**, e1008927 (2020).

57. Folkersen, L. *et al.* Genomic and drug target evaluation of 90 cardiovascular proteins in 30,931

individuals. *Nat Metab* **2**, 1135–1148 (2020).

58. Auton, A. *et al.* A global reference for human genetic variation. *Nature* **526**, 68–74 (2015).

59. Lijoi, A., Prünster, I. & Walker, S. G. On Consistency of Nonparametric Normal Mixtures for Bayesian

Density Estimation. *Journal of the American Statistical Association* **100**, 1292–1296 (2005).

60. Berisa, T. & Pickrell, J. K. Approximately independent linkage disequilibrium blocks in human

populations. *Bioinformatics* **32**, 283–285 (2016).

61. Robinson, G. K. That BLUP is a Good Thing: The Estimation of Random Effects. *Statistical Science* **6**,

15–32 (1991).

62. Zou, H. & Hastie, T. Regularization and variable selection via the elastic net. *Journal of the Royal*

Statistical Society: Series B (Statistical Methodology) **67**, 301–320 (2005).

63. Loh, P.-R. *et al.* Efficient Bayesian mixed-model analysis increases association power in large

cohorts. *Nat Genet* **47**, 284–290 (2015).

TWAS of Cardiovascular Disease by OTTERS

Test R^2