1 Responses of cortical neurons to intracortical microstimulation in

2 awake primates

Running Title: Responses of cortical neurons to ICMS in vivo

Richy Yun[1,4,5]*, Jonathan H. Mishler[1,4,5]*, Steve I. Perlmutter[2,4,5], Rajesh P. N. Rao[3,4], Eberhard
E. Fetz[1,2,4,5]

Departments of [1]Bioengineering, [2]Physiology and Biophysics, [3]Allen School for Computer
Science and Engineering, University of Washington
[4]Center for Neurotechnology, University of Washington
[5]Washington National Primate Research Center, University of Washington
*Authors contributed equally
The authors declare no competing financial interests.
Corresponding author: Richy Yun, rjyun@uw.edu

13 Acknowledgements

14 We thank Larry Shupe for programming and software support and Rebekah Schaefer for

15 assistance with animal care, handling, training, and surgery. We also thank Irene Rembado and

16 Nikolai Dembrow for helpful discussion. This work was supported by the National Institutes of

17 Health (NS012542, RR00166, and NS118781) and the National Science Foundation (EEC-

18 1028725).

Abstract

Intracortical microstimulation (ICMS) is commonly used in many experimental and clinical

paradigms; however, its effects on the activation of neurons are still not completely understood.

To document the responses of cortical neurons in non-human primates to stimulation, we

recorded single unit activity while delivering single-pulse stimulation via Utah arrays implanted

in primary motor cortex of three macaque monkeys. Stimuli between 5-50 μA delivered to single

channels reliably evoked spikes in neurons recorded throughout the array with delays of up to 12

milliseconds. ICMS pulses also induced a period of inhibition lasting up to 150 ms that typically

followed the initial excitatory response. Higher current amplitudes led to a greater probability of

evoking a spike and extended the duration of inhibition. The likelihood of evoking a spike in a

neuron was dependent on the spontaneous firing rate as well as the delay between its most recent

spike time and stimulus onset. Repetitive stimulation often modulated both the probability of

evoking spikes and the duration of inhibition, although high frequency stimulation in particular

was more likely to change both responses. Comparisons between stimuli that evoked a spike and

stimuli that did not revealed that the excitatory and inhibitory responses were independent of one

another on a trial-by-trial basis; however, their changes over time were frequently positively or

negatively correlated. Our results document the complex dynamics of cortical neural responses

to electrical stimulation that need to be considered when utilizing ICMS for scientific and

clinical applications.

Significance statement

Intracortical microstimulation (ICMS) is commonly used to probe the cortex, and previous

studies have characterized the responses of single neurons to ICMS. However, these studies

typically explored the averaged effects of ICMS throughout each experimental session, rather

than by a trial-by-trial basis for each stimulation pulse. By shifting the approach, we explored the

dependence of neural responses to ICMS on the spontaneous neural activity as well as the

dynamics of responses over time due to repetitive stimulation. Our results highlight how the

responses of neurons to ICMS are likely the result of interactions between local excitatory and

inhibitory cortical circuits. These results will help inform the design of ICMS for both basic

research and clinically relevant stimulation protocols.

Introduction

Intracortical microstimulation (ICMS) is widely used for interfacing with the brain in both basic

and clinical research, from inducing plasticity to employing sensory neuroprostheses in various

animal models (Flesher et al., 2016; Hartmann et al., 2016; Jackson & Fetz, 2011; Lebedev &

Nicolelis, 2017). The applicability of ICMS arises from the fact that it has the highest spatial and

temporal specificity of all clinically applicable cortical stimulation techniques (Sejnowski et al.,

2014). However, the circuit mechanisms that drive the responses of neurons following ICMS,

and the ways in which other factors such as timing and stimulation frequency affect the stimulus

responses are not fully understood.

ICMS was originally thought to activate neural elements around the electrode tip. Regions closer

to the tip would have higher activation in a sphere with an isotropic gradient, and the volume

would grow with increasing current amplitude (Ranck, 1975; Stoney et al., 1968; Tehovnik et al.,

2006). However, it is now accepted that ICMS predominantly excites axons near the electrode tip

that transsynaptically excite neurons up to several millimeters away, favoring pathways with

similar functional properties (Gustafsson & Jankowska, 1976; Hao et al., 2016; Histed et al.,

2009; Lesser et al., 2008; Logothetis et al., 2010; McIntyre & Grill, 2000). Additionally, the

effects of ICMS are not limited to excitation, and includes a long-lasting inhibitory response that

is commonly attributed to GABAergic synapses (Berman et al., 1991; Butovas & Schwarz, 2003;

Hao et al., 2016).

Single neuron responses to ICMS are dynamic and can be modulated with repeated stimulation.

The changes, in part, also depend on stimulus frequency. In particular, the excitation of neurons

generally decreases over time and becomes more localized with higher frequencies (Dadarlat et

al., 2019; Michelson et al., 2019). However, the reported frequency ranges and timescales are

variable, and the driving processes remain unclear. The changes over time are often attributed to

short-term synaptic plasticity or intrinsic plasticity of neurons which both depend on the

frequency and pattern of stimulation (Abbott & Regehr, 2004; Citri & Malenka, 2008; Zucker &

Regehr, 2002).

Altogether, these studies demonstrate that the effects of ICMS are not restricted to regions

proximal to the electrode tip, and that responses consist of interplay between excitation and

inhibition (Borchers et al., 2012; Griffin et al., 2011; Logothetis et al., 2010). Despite our

increasing understanding of how ICMS activates cortical circuits, several significant questions

remain. How does the background neuronal activity, including firing rate and previous spike

time impact the stimulus response? How do the responses change over time as a function of both

the frequency of stimulation and proximity to stimulation site? Is the inhibitory response coupled

to the excitatory response, or are they independent?

We addressed the questions above by delivering ICMS and examining responses of single

neurons in primary motor cortex (M1) of three macaque monkeys with chronically implanted

Utah arrays. Single-pulse ICMS was delivered to one channel for up to 20 minutes while the

spikes of single neurons were simultaneously recorded from all other electrode channels. We

tracked the probability of evoking spikes as well as the duration of the evoked inhibition and

varied both the sites and frequency of the stimulation between sessions. Our results expand upon

previous findings by characterizing the dependencies of the neuronal responses to background

neuronal activity, distance, and stimulation frequency, and exploring the interactions between the

excitatory and inhibitory responses.

Methods and Materials

Experimental Design

Implants and surgery

Three pigtail macaque monkeys (*Macaca nemestrina*) were unilaterally (right hemisphere) or bilaterally implanted with 96-channel Utah microelectrode arrays (Blackrock Microsystems; 10x10, 400μm inter-electrode distance, 1.5 mm depth, Iridium oxide) in the hand region of M1. Sterile surgeries were performed under isoflurane anesthesia and aseptic conditions with continuous monitoring of all vitals. Animals received postoperative courses of analgesics and antibiotics following each surgery. All procedures conformed to the National Institutes of Health *Guide for the Care and Use of Laboratory Animals* and were approved by the University of Washington Institutional Animal Care and Use Committee.

Implantation of the arrays was guided via stereotaxic coordinates. A 1.5 cm wide square craniotomy centered at 4 mm lateral of Bregma was performed to expose the dura. Three sides of the exposed dura were cut to expose the cortex, after which a Utah microelectrode array was implanted. Two reference wires were inserted under the dura and two were inserted between the dura and the skull. The dura was then sutured around the implant, and the bone flap from the craniotomy was reattached to the skull with a titanium strap and titanium skull screws. A second, smaller titanium strap was screwed onto the skull to secure the wire bundle that connected the array to a connector "pedestal" that was also secured to the skull with eight skull screws. The skin incision was then sutured around the pedestal base.

To facilitate the chronic recording of neuronal activity, the monkeys were also implanted with

halos made with 3/8" aluminum bars in an egg-shaped oval that was 17 cm long and 15.3 cm

wide. Four titanium straps were affixed to the skull via titanium bone screws. Two of the straps

were implanted bilaterally over the occipital ridge, and two were bilaterally implanted

temporally. After the plates integrated with the skull for 6 weeks, the halo was secured to the

skull with four pins, each of which were seated in one of the four plates.

Electrophysiology

Stimulation and recording of single unit activity were conducted with one of three systems: 1)

Neurochip3 (custom bidirectional brain-computer interface developed in our laboratory (Shupe

et al., 2021), 32 channels, 20 kHz sampling rate), 2) Neural Interface Processor (Ripple Neuro,

96 channels, 30 kHz sampling rate), or 3) RZ2 BioAmp Processor, PZ5 NeuroDigitizer

Amplifier, and IZ2 Electrical Stimulator (Tucker-Davis Technologies, 96 channels, 25 kHz

sampling rate).

Experimental Setup

The monkeys were trained to calmly sit in a primate chair while periodically receiving an apple

smoothie reward without performing a task (Figure 1). Each session included a prestimulus

epoch lasting between 5 and 10 minutes and a stimulus epoch lasting between 5 and 20 minutes.

During the stimulus epoch we delivered tonic or Poisson distributed single pulse stimulation

(cathodal, biphasic, 200μs phase width) to a single channel at rates between 1-20 Hz. For testing

the effects of current amplitude, a range of 2-50 μA was used. Current amplitude was fixed at 15

μA for all other experiments and analyses. The stimulation frequency was fixed during the

stimulus epoch for each session.

<u>Data analysis</u>

Evoked spike acquisition

Spikes were sorted using two time-amplitude windows, initially online and subsequently

confirmed offline. Stimulus artifacts lasted around 1.1-1.6 ms. Spikes were frequently detected

immediately following the artifact (Figure 2A). The timing of evoked spikes was found by

calculating the peristimulus time histogram (PSTH, 0.5 ms bin widths) of spikes in the window

from -20 to 20 ms from the time of stimulation. To isolate the evoked spikes from the

spontaneous activity, we defined upper and lower thresholds in the PSTH as the histogram mean

plus or minus 2 times the standard deviation from -20 to -2 ms. We then found the largest peak in

the PSTH from 1 to 15 ms after stimulation that was larger than the upper threshold and tracked

adjacent bins in both directions until we reached the lower threshold on both sides. All spikes

occurring within this window were denoted as stimulus-evoked spikes (Figure 2B). If no peak

was greater than the threshold the spike was not considered to have been evoked by stimulation.

The probability of evoked spikes was calculated as the number of evoked spikes divided by the

number of total stimuli. For any analysis over time, the evoked spike probability was calculated

for stimuli within overlapping 30 second bins with 1 second steps.

We tracked a total of 602 units, 17 of which were exclusively used to analyze the effects of

stimulus timing on the probability of evoking a spike since the analysis required high firing

and/or stimulation rates. For these experiments, stimulation was sometimes delivered to multiple

channels in the same experiment. All units were used for all other analyses.

Inhibitory response acquisition

The inhibitory response was measured using the PSTH in previous studies (Butovas & Schwarz, 2003; Hao et al., 2016). While evoked spike timing can easily be determined with the PSTH due to their high probability and narrow time window, the inhibitory response depends on a broad window with sparse activity, particularly for units with low firing rate. As a result, a large number of stimuli is required to reliably detect inhibition in the PSTH. Therefore, rather than using the PSTH, we measured the duration of inhibition by removing all evoked spikes and calculating the time between the onset of stimulation and the next spontaneous spike (Figure 2C). Inhibition was deemed to be stronger when the delay from stimulation onset to the next spontaneous spike was longer, giving us a measure directly comparable to the PSTH but with much higher temporal resolution. We discarded any stimuli for which the subsequent stimuli occurred before the next spontaneous spike. For any analysis over time, we used the median inhibition duration of stimuli within overlapping 30 second bins with 1 second steps.

Evoked spike probability dependencies

To calculate the spontaneous firing rate for each unit over time, we disregarded the times from each stimulus onset to the next spontaneous spike. This effectively removed the stimulus response from the firing rate calculation, providing us with an independent measure of spontaneous activity.

The autocorrelation histograms for the 17 units in which we characterized the evoked spike probability as a function of the time delay between their previous spike times and stimulation onsets were calculated from their respective prestimulus epochs. The histograms were binned between 0 to 50 ms in non-overlapping 1 ms bins.

The dependencies of evoked spike probabilities on the timing of stimulation were calculated by

first measuring the delays between each stimulus and the preceding spike. We separated this into

two groups corresponding to whether the preceding spike was spontaneous or evoked. Stimuli

that did not have a spike following the previous stimulus were discarded from the analysis. To

account for the transmission delay from the stimulated site to the evoked spike, we added the

average evoked spike latency for each unit to the delays. We then calculated the probability of

evoking a spike for stimuli with delays from 0 to 50 ms with 1 ms time bins, then applied a

moving average with a 5 ms window. Since the average spike latencies were added to each

delay, time bins that were less than the average evoked spike latency were not included in

correlation calculations.

Changes over time

Pairwise correlations and their statistical significance between firing rate, evoked spike

probability, and inhibition duration over time were calculated using the Pearson correlation

coefficient r:

$$r = \frac{\sum (x_i - \bar{x})(y_i - \bar{y})}{\sqrt{\sum (x_i - \bar{x})^2 \sum (y_i - \bar{y})^2}} \tag{1}$$

where $\bar{x}$ is the mean of the x-variable, and $\bar{y}$ is the mean of the y-variable. Firing rate was

calculated by removing all times between each stimulus onset to the first spontaneous spike (the

inhibitory period) to remove any confounding affects between firing rate and the inhibitory

response.

Linear and exponential fits were performed on binned evoked spike probabilities and inhibition

durations to determine changes over time due to repetitive stimulation:

Linear	$y = a * t + b$	(2)
Exponential	$y = a + b * (1 - c)^t$	(3)

where y is either the evoked spike probability or inhibition duration, t is time and a, b, and c are

the fitted variables. Changes were denoted to occur if the analysis of variance (ANOVA) F-

statistic resulted in $p<0.05$. The sign of the linear fit slope (a in Equation 2) or the sign of the

exponent base (b in Equation 3) of the exponential fit determined whether the changes were

classified as increasing or decreasing. Neuron-stimulation site pairs with less than 3% average

probability of evoking spikes were disregarded for analyses over time due to their inconsistency.

The changes over time in spikes and inhibition were designated to be correlated if their

correlation had a p-value less than 0.05.

Statistical analysis

Student's t-test was used for statistical comparisons between groups when the data were

determined to be normally distributed. Two-sided Wilcoxon signed-rank test (*signrank*,

MATLAB) was used to confirm normality and for nonparametric data. Paired t-test or two-sided

Wilcoxon rank-sum (*ranksum*, MATLAB) tests were used where appropriate. Fisher's exact test

(*fishertest*, MATLAB) or two-way analysis of variance (ANOVA) (*anova2*, MATLAB) was

used to compare categorical data. The Pearson correlation was used for all correlation tests. P-

values for significance and tests used are reported in individual analyses.

Results

Evoked spikes and inhibitory response

We found that ICMS elicited a brief excitatory response followed by a longer inhibition.

Electrodes on the Utah array typically showed evoked spikes occurring 1.5 to 10 ms after single-

pulse stimulation. The inhibition typically followed the excitatory response and was observed as

the suppression of firing in the PSTH for 5 - 100 ms, although in some instances it lasted up to

200 ms. We also observed that stimulus-evoked inhibition could occur in the absence of the

excitatory response.

To ensure that stimulation arriving during inhibition was not affecting the stimulus response, we

delivered trains of two or three pulses with each subsequent pulse timed to occur during the

inhibitory response of the previous stimulus. Single, double, and triple pulse stimuli were

delivered to a single channel, randomly interleaved at 2 Hz. Our results across 5 different

sessions show that stimuli delivered during the inhibitory response were able to reliably evoke

spikes comparable to when stimuli was delivered at other times, as previously reported (Butovas

& Schwarz, 2003) (Figure 3, left). Furthermore, each stimulus pulse "reset" the inhibitory

response such that the duration of inhibition was the same following each pulse train (Figure 3,

right).

Effects of distance from stimulation site and stimulus amplitude

Evoked spikes occurred with greater probability and less variable latencies for units close to the

stimulus electrode than for more distant units. The probability of evoking a spike in electrodes

<1 mm from the stimulus site was significantly greater than for further sites (p=1.3e-14). In

addition, closer sites on average had evoked spikes that occurred at shorter, and less variable

latencies (p=8.5e-29), suggesting the presence of mono- and polysynaptic activation (Figure 4A).

The duration of inhibition had a similar trend: recording sites <1 mm from the stimulus site

tended to show stronger inhibition compared to further sites (p=0.003) (Figure 4B). The

probability of evoking spikes and the duration of inhibition increased sigmoidally with the

stimulus amplitude for all responsive units (Figure 5). The sigmoid curves were always steep: a

change of 10 to 20 μA in stimulus intensity generated the difference between 5% and 95% of the

maximum value for both evoked spike probability and inhibition duration.

<u>Spontaneous activity affects stimulus responses</u>

In addition to stimulation current and separation of the recording and stimulation sites, we found

two other dependencies that affect the response. One is the spontaneous firing rate of the

recorded units. Both the evoked spike probability and inhibition duration had statistically

significant correlations with spontaneous firing rate over time. Table 1 documents the number of

units with uncorrelated, positively correlated, or negatively correlated evoked spike probability

and inhibition duration with spontaneous firing rate. A slight majority of units tested (300/585,

51%) had evoked spike probabilities that were positively correlated with firing rate and

inhibition duration that were negatively correlated with firing rate. We additionally performed a

2-way ANOVA to determine whether the two relationships were dependent on one another but

found no significant relationship (p=0.11). Units with positively correlated evoked spike

probabilities and spontaneous firing rate were typically farther from the stimulated site than units

with a negatively correlated relationship (Figure 6C). In contrast, units with positively correlated

inhibition duration and spontaneous firing rate were typically closer to the stimulated site (Figure

6C).

The second dependency was the timing of stimuli relative to the most recent spike. We analyzed

17 units (2 from monkey S, 3 from monkey K, and 12 from monkey J). In 14 of the 17 units (2

from monkey S, 3 from monkey K, and 9 from monkey J), the probability of evoking a spike

varied as a function of the time between the onset of stimulation and the most recent spontaneous

spike. For these 14 units the probability was significantly positively correlated with the unit's

autocorrelogram in the absence of stimulation (Figure 7, left and middle). The probability

distributions were the same even if the most recent spike was evoked by stimulation rather than

being spontaneous. The inhibitory response did not depend on the timing of pre-stimulus spikes.

In 3 of the 17 tested units the probability of evoking a spike did not depend on the timing

between the previous spike and stimulus onset. In all 3 of these units the probability of evoking a

spike was consistently high (Figure 7, right), which suggests that the stimulation amplitude was

large enough to evoke spikes regardless of other properties.

<u>Repetitive stimulation changes stimulus response over time</u>

Repetitive microsimulation has been shown to modulate the responses of units to ICMS

(Dadarlat et al., 2019; Michelson et al., 2019; Zucker & Regehr, 2002). We documented the

effects of repetitive ICMS over the stimulus period ranging from 5 to 20 minutes on the evoked

spike probability and the duration of inhibition by delivering stimuli at between 1 and 20 Hz.

The probability of evoking a spike often increased or decreased over time, following a linear or

exponential trend over the course of the session to an asymptote (Figure 8A).

The evoked spike probability was significantly more likely to change with higher frequencies of

stimulation, with a clear trend over successive 5 Hz bins (Figure 8B). Of the changes, high

frequencies (15-20 Hz) were likely to cause more decreases in evoke spike probability compared

to lower frequencies. Changes in inhibition duration over time were also more likely to occur at

higher frequencies, but there was no clear frequency dependence in the direction of change.

(Figure 8B)

We also observed changes depending on the distance from the recording site to the stimulated

site (Figure 8C). The probability of evoking spikes was significantly more likely to change the

closer the units were to the stimulated site. Sites further away tended to be more likely to have

decreased probability over time, but this was not statistically significant (p>0.05 for all pairwise

comparisons). Similarly, inhibition duration also had a higher likelihood of changing over time

in sites closer to the stimulated site. The changes tended to be more likely to decrease further

away, but this was also not statistically significant (p>0.05 for all pairwise comparisons).

Finally, in 72/585 (12%) of the tested units, we also observed changes over time in the latency of

the evoked spikes (Figure 9A). Latency changes typically occurred in units that were recorded on

electrodes closer to the stimulation site (Figure 9B), and more commonly occurred with high

frequency stimulation (Figure 9C). Of the units with changes in their evoked spike latency, 51/72

(71%) units had increasing latencies and 21/72 (29%) had decreasing latencies.

<u>Relationship between evoked spikes and inhibition</u>

Since evoked spikes and inhibition often both exhibited changes over time, we sought to

determine whether they were directly related on a trial-by-trial basis. Individual stimuli in each

experiment were divided into two categories: those that evoked spikes and those that did not. For

the unit in Figure 10A the PSTHs of the two classifications of stimuli are very similar except for

the evoked spike peak. We found no statistical pairwise difference between the inhibition

induced by stimuli that evoked spikes compared to the stimuli that did not evoke spikes (585

units, Figure 10B).

Although evoked spikes and inhibition seemed to be independently activated on a trial-by-trial

basis, the mechanisms underlying the changes over time could nevertheless be related. We

analyzed the correlation between the probability of evoking a spike and duration of inhibition

over time to determine whether they were positively or negatively correlated. Correlations with

$p<0.05$ were considered significant, and all other instances were denoted to be uncorrelated. Of

the 585 units tested, 27% (161 units) had positively correlated changes in the probability of

evoking spikes and the duration of inhibition, 50% (292 units) had negatively correlated changes,

and 23% (132 units) were uncorrelated. Units with positively correlated evoked spike

probabilities and inhibition duration tended to be closer to the stimulated site (Figure 10C).

<u>Spike type does not correlate with stimulus response properties</u>

To determine whether the cell type of the recorded units influenced their responses to ICMS, we

used both the spike waveform and firing rate to classify each neuron as fast-spiking (FS) or

regular-spiking (RS). Although previous studies have used the spike width and the inter-spike

interval to differentiate between the two (Connors & Gutnick, 1990; McCormick et al., 1985),

we used a proxy measure of the area under the action potential curve divided by the trough

magnitude, which resulted in a more continuous distribution.

Figures 11A and B show an example of the two different spike waveforms, as well as the

distribution of their action potential areas and firing rates, respectively. We separated the units

into two groups based on their defined spike width shown at the dotted line in Figure 11B.

Around 10% of all units fell to the left of the line and were denoted to be FS; the rest were

denoted to be RS. We found that the putative cell type did not correlate with the distribution of

distances from the stimulated site, whether a spike could be evoked, the probability of evoking a

spike, the duration of inhibition, or how any measure changed over time (Figures 11C-G). Thus,

all results reported herein are independent of the type of neuron recorded.

Discussion

<u>Comparisons to previous studies</u>

ICMS has been shown to predominantly activate neurons transsynaptically (Butovas & Schwarz,

2003; Hussin et al., 2015; Klink et al., 2017). We observed in our study that spike latencies

fluctuated more than would occur with antidromic activation, and stimuli that were delivered

within 1 millisecond after a spike were still able to evoke spikes. Since antidromic activation

would result in collision and an absence of a spike at such short latencies, this suggests that the

recorded units were predominantly activated orthodromically. Also consistent with previous

experiments, we saw evoked spikes in units recorded up to 4.5 mm away from the stimulation

site, which suggests that ICMS activates a distributed population rather than only a concentrated

sphere of neurons around the electrode tip (Butovas & Schwarz, 2003; Hao et al., 2016; Histed et

al., 2009).

Although the excitatory response was directly measurable in our experiments, the subsequent

inhibitory response manifested as a lack of spikes. Butovas et al. concluded that a similar

inhibitory response to ICMS was likely caused by $GABA_B$ receptors, which they confirmed with

a follow-up study with pharmacological blocks (Butovas et al., 2006; Butovas & Schwarz,

2003). GABAergic inhibition would also explain the similarity in the sigmoidal curves of evoked

spikes and inhibition with stimulus amplitude in our experiments – if more excitatory neurons

are excited by the higher intensity stimulation, more inhibitory neurons will be activated via

feedforward and feedback circuits, thereby increasing the amount of inhibition (Figure 12). A

lack of relationship between inhibition and distance to stimulation site as well as the presence of

inhibition in units without evoked spikes is likely due to the high connectivity of interneurons

compared to principal cells (Isaacson & Scanziani, 2011; Matsumura et al., 1996; Rudy et al.,

2013; Tremblay et al., 2016). Interestingly, Hao et al. found that inhibition decreased as a

function of distance, which we did not find (Hao et al., 2016). Potential reasons for this

discrepancy could be the differences in how we measured inhibition and the stimulus amplitude.

However, we found that inhibition typically lasted between 5-100 ms and rarely over 100 ms,

which is significantly shorter than the average time constants of $GABA_B$ inhibitory postsynaptic

potentials of 150-200 ms (Bettler et al., 2004; Connors et al., 1988). Therefore, $GABA_A$

mediated inhibition is a more likely candidate to explain our results; previous studies have shown

that $GABA_A$ is involved in recurrent polysynaptic inhibition, which we are likely activating via

ICMS (Silberberg & Markram, 2007; Zhu et al., 2011). The different animal models and

recorded cortical region in these studies may account for these discrepancies.

ICMS activates long horizontal fibers to both feedforward inhibitory networks and the recorded

excitatory neurons (Figure 12). These afferent fibers have stronger excitatory connections to the

inhibitory interneurons than the principal cells, particularly in layer 2/3, which may explain why

we sometimes observe an inhibitory response without any excitatory response (Adesnik &

Scanziani, 2010; Cruikshank et al., 2007; Helmstaedter et al., 2008). This, coupled with the fact

that inhibitory neurons often target somatic or perisomatic compartments (Isaacson & Scanziani,

2011; Rudy et al., 2013; Tremblay et al., 2016), suggests that the observed inhibition may be

initially activated via feedforward circuitry, and is subsequently followed by the excitatory

response. The excited principal cells may then activate feedback circuitry that contributes to the

inhibitory response (Figure 12).

<u>Stimulus response depends on network activity and intrinsic membrane properties</u>

Previous research has shown that stimulus responses depend on network activity both *in vitro*

(Kumar et al., 2016; Weihberger et al., 2013) and *in vivo* (Kara et al., 2002). Our findings were

consistent with these results; the probability of evoking a spike was often positively correlated

with spontaneous firing rate, whereas the duration of inhibition was often negatively correlated

with firing rate.

English et al. demonstrated that the transmission probability for postsynaptic spikes of inhibitory

neurons in the hippocampus *in vivo* is a function of the timing between the previous postsynaptic

spike and presynaptic spike (English et al., 2017). Moreover, that study found that this

dependency was independent of whether the previous postsynaptic spike was spontaneous or

evoked, which suggested that intrinsic properties of the postsynaptic membrane were responsible

for the dependence. Our results extend these outcomes by demonstrating that the timing of the

previous spike affects not only the transmission probability for spontaneous presynaptic spikes,

but also the stimulus-evoked spikes in cortical neurons. Furthermore, we found that the

probability distributions for evoking a spike as a function of the timing between the previous

spike time and stimulation onset and the spike train autocorrelation were often significantly

positively correlated, which further reinforces the notion that this dependency reflects the

intrinsic properties of the recorded units.

Altogether, our results reveal that the stimulus response has at least two dependencies other than

stimulus amplitude and distance from stimulus site: the intrinsic membrane properties of the

recorded neurons and the activity of the network. However, a sufficiently large stimulus current

may saturate the responses and overcome these dependencies.

Repetitive stimulation modulates stimulus responses

Michelson et al. showed that the number of neurons activated by electrical stimulation

diminished over time with higher frequencies measured with calcium imaging in a 407 μm x 407

μm window (Michelson et al., 2019). Their study showed that changes began at regions distal

from the stimulated site when delivering stimuli with frequencies greater than 10 Hz. These

results are consistent with our study across the larger spatial field (4x4 mm) of the Utah array, as

evoked spikes were more likely to be diminished with higher frequency of stimulation. The

changes we observed occurred over much longer timescales and continuously for up to 20

minutes. However, we also observed that in some units the probability of evoking a spike

increased over time even at longer distances and higher frequencies, which cannot be fully

explained by a diminishing region of activation.

Although we did not explicitly measure the duration of changes induced by repetitive

stimulation, we observed that they typically lasted less than 2 minutes. Due to the short-lived

nature of the induced changes, various mechanisms of short-term synaptic plasticity such as

vesicle depletion and facilitation by calcium influx (Citri & Malenka, 2008; Zucker & Regehr,

2002) may best explain our results. Evidence suggests that different forms of short-term

plasticity exist for synaptic connections between different cell-types (Blackman et al., 2013;

Losonczy et al., 2002). Beyond these differences, previous *in vitro* work by Markram showed

that synaptic connections between pyramidal neurons of the same morphological class and

interneurons had similar facilitating and depressing characteristics, but with different time

courses (Markram et al., 1998).

Together, these results may explain why the frequency-dependent changes that we measured

were different for each unit. Although we did not discern any differences between regular and

fast spiking neurons for any measure, there are limitations in such cell type classifications with

extracellular recordings. Furthermore, we also observed changes in the latency of evoked spikes

due to repetitive stimulation, which has previously been shown to occur in the presence of short-

term plasticity (Boudkkazi et al., 2007). Future studies with specific differentiation between cell

and synapse types may shed more light on whether cell-type specific differences account for the

variability across spikes.

<u>Excitation and inhibition are independently activated but modulated together</u>

<u>within an interconnected network</u>

The balance between excitation and inhibition within the cortex is a much-studied topic and is

highly relevant to neural computation. Though the examined network size, location, and synaptic

connections vary greatly, the strong consensus is that excitation and inhibition are generally

comodulated (Chen, 2004; Haider, 2006; Isaacson & Scanziani, 2011; Rubin et al., 2017;

Turrigiano, 2012; Xue et al., 2014). We found that the probability of evoking a spike and the

duration of inhibition were frequently positively or negatively correlated over time, though they

were independent on a trial-by-trial basis. Units close to the stimulated site typically had

positively correlated evoked spike probability and inhibition duration, both of which were

negatively correlated with firing rate. Units far from the stimulation channel, however, had

positively correlated evoked spike probabilities and firing rates, which were both negatively

correlated with inhibition duration.

The effect of distance can be explained by the fact that sites closer to stimulation are more likely

to be activated by ICMS (Butovas & Schwarz, 2003; Hao et al., 2016). Due to the feedforward

and feedback inhibitory circuitry, if the total excitation increased or decreased due to short-term

plasticity, the inhibition should change in a positively correlated manner. Sites far from

stimulation, however, are not activated as comprehensively and are thus less likely to be

susceptible to short-term plasticity. Similarly, the negative correlations in evoked spike

probability and inhibition at these far sites are likely due to network dynamics, whereas the

positive correlations in closer sites are likely due changes in short-term plasticity caused by

direct activation via ICMS.

References

Abbott, L. F., & Regehr, W. G. (2004). Synaptic computation. *Nature*, *431*(October), 796–803.

Adesnik, H., & Scanziani, M. (2010). Lateral competition for cortical space by layer-specific

horizontal circuits. *Nature*, *464*(7292), 1155–1160.

Berman, N. J., Douglas, R. J., Martin, K. A. C., & Whitteridge, D. (1991). Mechanisms of

inhibition in cat visual cortex. *Journal of Physiology*, *440*, 697–722.

Bettler, B., Kaupmann, K., Mosbacher, J., & Gassmann, M. (2004). Molecular structure and

physiological functions of GABAB receptors. *Physiological Reviews*, *84*(3), 835–867.

Blackman, A. V., Abrahamsson, T., Costa, R. P., Lalanne, T., & Sjöström, P. J. (2013). Target-

cell-specific short-term plasticity in local circuits. *Frontiers in Synaptic Neuroscience*, *5*(DEC),

1–13.

Borchers, S., Himmelbach, M., Logothetis, N., & Karnath, H. (2012). Direct electrical

stimulation of human cortex — the gold standard for mapping brain functions? *Nature Reviews*

Neuroscience, *13*(January), 63–70.

Boudkkazi, S., Carlier, E., Ankri, N., Caillard, O., Giraud, P., Fronzaroli-Molinieres, L., &

Debanne, D. (2007). Release-dependent bariations in synaptic latency: A putative code for short-

and long-term synaptic dynamics. *Neuron*, *56*(6), 1048–1060.

Butovas, S., Hormuzdi, S. G., Monyer, H., & Schwarz, C. (2006). Effects of electrically coupled

inhibitory networks on local neuronal responses to intracortical microstimulation. *Journal of*

Neurophysiology, *96*(3), 1227–1236.

Butovas, S., & Schwarz, C. (2003). Spatiotemporal rffects of microstimulation in rat neocortex:

A parametric study using multielectrode recordings. *Journal of Neurophysiology*, *90*(5), 3024–

3039.

Chen, R. (2004). Interactions between inhibitory and excitatory circuits in the human motor

cortex. *Experimental Brain Research*, *154*(1), 1–10.

Citri, A., & Malenka, R. C. (2008). Synaptic plasticity: Multiple forms, functions, and

mechanisms. *Neuropsychopharmacology*, *33*(1), 18–41.

Connors, B. W., & Gutnick, M. J. (1990). Intrinsic firing patterns of diverse neocortical neurons.

Trends in Neurosciences, *13*(3), 99–104.

Connors, B. W., Malenka, R. C., & Silva, L. R. (1988). Two inhibitory postsynaptic potentials,

and GABAA and GABAB receptor-mediated responses in neocortex of rat and cat. *The Journal*

of Physiology, *406*(1), 443–468.

Cruikshank, S. J., Lewis, T. J., & Connors, B. W. (2007). Synaptic basis for intense

thalamocortical activation of feedforward inhibitory cells in neocortex. *Nature Neuroscience*,

10(4), 462–468.

Dadarlat, M. C., Sun, Y., & Stryker, M. P. (2019). Widespread activation of awake mouse cortex

by electrical stimulation. *International IEEE/EMBS Conference on Neural Engineering, NER*,

2019-March, 1113–1117.

English, D. F., McKenzie, S., Evans, T., Kim, K., Yoon, E., & Buzsáki, G. (2017). Pyramidal

cell-interneuron circuit architecture and dynamics in hippocampal networks. *Neuron*, *96*(2), 505-

520.e7.

Flesher, S. N., Collinger, J. L., Foldes, S. T., Weiss, J. M., Downey, J. E., Tyler-Kabara, E. C.,

Bensmaia, S. J., Schwartz, A. B., Boninger, M. L., & Gaunt, R. A. (2016). Intracortical

microstimulation of human somatosensory cortex. *Science Translational Medicine*, *8*(361), 1–11.

Griffin, D. M., Hudson, H. M., Belhaj-Saif, A., & Cheney, P. D. (2011). Hijacking cortical motor

output with repetitive microstimulation. *Journal of Neuroscience*, *31*(37), 13088–13096.

Gustafsson, B., & Jankowska, E. (1976). Direct and indirect activation of nerve cells by

electrical pulses applied extracellularly. *The Journal of Physiology*, *258*(1), 33–61.

Haider, B. (2006). Neocortical network activity in vivo is generated through a dynamic balance

of excitation and inhibition. *Journal of Neuroscience*, *26*(17), 4535–4545.

Hao, Y., Riehle, A., & Brochier, T. G. (2016). Mapping horizontal spread of activity in monkey

motor cortex using single pulse microstimulation. *Frontiers in Neural Circuits*, *10*(December),

1–16.

Hartmann, K., Thomson, E. E., Zea, I., Yun, R., Mullen, P., Canarick, J., Huh, A., & Nicolelis,

M. A. L. (2016). Embedding a panoramic representation of infrared light in the adult rat

somatosensory cortex through a sensory neuroprosthesis. *Journal of Neuroscience*, *36*(8).

Helmstaedter, M., Staiger, J. F., Sakmann, B., & Feldmeyer, D. (2008). Efficient recruitment of

layer 2/3 interneurons by layer 4 input in single columns of rat somatosensory cortex. *Journal of*

Neuroscience, *28*(33), 8273–8284.

Histed, M. H., Bonin, V., & Reid, R. C. (2009). Direct activation of sparse, distributed

populations of cortical neurons by electrical microstimulation. *Neuron*, *63*(4), 508–522.

Hussin, A. T., Boychuk, J. A., Brown, A. R., Pittman, Q. J., & Campbell Teskey, G. (2015).

Intracortical microstimulation (ICMS) activates motor cortex layer 5 pyramidal neurons mainly

transsynaptically. *Brain Stimulation*, *8*(4), 742–750.

Isaacson, J. S., & Scanziani, M. (2011). How inhibition shapes cortical activity. *Neuron*, *72*(2),

231–243.

Jackson, A., & Fetz, E. E. (2011). Interfacing with the computational brain. *IEEE Transactions*

on Neural Systems and Rehabilitation Engineering.

Kara, P., Pezaris, J. S., Yurgenson, S., & Reid, R. C. (2002). *The spatial receptive field of*

thalamic inputs to single cortical simple cells revealed by the interaction of visual and electrical

stimulation. *99*(25), 1–6.

Klink, P. C., Dagnino, B., Gariel-Mathis, M. A., & Roelfsema, P. R. (2017). Distinct

feedforward and feedback effects of microstimulation in visual cortex reveal neural mechanisms

of texture segregation. *Neuron*, *95*(1), 209-220.e3.

Kumar, S. S., Wülfing, J., Okujeni, S., Boedecker, J., Riedmiller, M., & Egert, U. (2016).

Autonomous optimization of targeted stimulation of neuronal networks. *PLoS Computational*

Biology, *12*(8), 1–22.

Lebedev, M. A., & Nicolelis, M. A. L. (2017). Brain-machine interfaces: From basic science to

neuroprostheses and neurorehabilitation. *Physiological Reviews*, *97*(2), 767–837.

Lesser, R. P., Lee, H. W., Webber, W. R. S., Prince, B., Crone, N. E., & Miglioretti, D. L.

(2008). Short-term variations in response distribution to cortical stimulation. *Brain*, *131*(6),

1528–1539.

Logothetis, N. K., Augath, M., Murayama, Y., Rauch, A., Sultan, F., Goense, J., Oeltermann, A.,

& Merkle, H. (2010). The effects of electrical microstimulation on cortical signal propagation.

Nature Neuroscience, *13*(10), 1283–1291.

Losonczy, A., Zhang, L., Shigemoto, R., Somogyi, P., & Nusser, Z. (2002). Cell type

dependence and variability in the short-term plasticity of EPSCs in identified mouse

hippocampal interneurones. *Journal of Physiology*, *542*(1), 193–210.

Markram, H., Wang, Y., & Tsodyks, M. (1998). Differential signaling via the same axon of

neocortical pyramidal neurons. *Proceedings of the National Academy of Sciences of the United*

States of America, *95*(9), 5323–5328.

Matsumura, M., Chen, D., Sawaguchi, T., Kubota, K., & Fetz, E. E. (1996). Synaptic interactions

between primate precentral cortex neurons revealed by spike-triggered averaging of intracellular

membrane potentials in vivo. *Journal of Neuroscience*, *16*(23), 1–11.

McCormick, D. A., Connors, B. W., Lighthall, J. W., & Prince, D. A. (1985). Comparative

electrophysiology of pyramidal and sparsely spiny stellate neurons of the neocortex. *Journal of*

Neurophysiology, *54*(4), 782–806.

McIntyre, C. C., & Grill, W. M. (2000). Selective microstimulation of central nervous system

neurons. *Annals of Biomedical Engineering*, *28*(3), 219–233.

Michelson, N. J., Eles, J. R., Vazquez, A. L., Ludwig, K. A., & Kozai, T. D. Y. (2019). Calcium

activation of cortical neurons by continuous electrical stimulation: Frequency dependence,

temporal fidelity, and activation density. *Journal of Neuroscience Research*, *97*(5), 620–638.

Ranck, J. B. (1975). Which elements are excited in electrical stimulation of mammalian central

nervous system: A review. *Brain Research*, *98*(3), 417–440.

Rubin, R., Abbott, L. F., & Sompolinsky, H. (2017). Balanced excitation and inhibition are

required for high-capacity, noise-robust neuronal selectivity. *Proceedings of the National*

Academy of Sciences of the United States of America, *114*(44), E9366–E9375.

Rudy, B., Fishell, G., Lee, S., & Hjerling-Leffler, J. (2013). Three groups of interneurons

account for nearly 100 % of Neocortical GABAergic Neurons. *Dev Neurobiol*, *71*(1), 45–61.

Sejnowski, T. J., Churchland, P. S., & Movshon, J. A. (2014). Putting big data to good use in

neuroscience. *Nature Neuroscience*, *17*(11), 1440–1441.

Shupe, L. E., Miles, F. P., Jones, G., Yun, R., Mishler, J., Rembado, I., Murphy, R. L.,

Perlmutter, S. I., & Fetz, E. E. (2021). Neurochip3: An autonomous multichannel bidirectional

brain-computer interface for closed-loop activity-dependent stimulation. *Frontiers in*

Neuroscience, *15*(August), 1–15.

Silberberg, G., & Markram, H. (2007). Disynaptic inhibition between neocortical pyramidal cells

mediated by martinotti cells. *Neuron*, *53*(5), 735–746.

Stoney, S. D., Thompson, W. D., & Asanuma, H. (1968). Excitation of pyramidal tract cells by

intracortical microstimulation: effective extent of stimulating current. *Journal of*

Neurophysiology, *31*(5), 659–669.

Tehovnik, E. J., Tolias, A. S., Sultan, F., Slocum, W. M., & Logothetis, N. K. (2006). Direct and

indirect activation of cortical neurons by electrical microstimulation. *Journal of*

Neurophysiology, *96*(2), 512–521.

Tremblay, R., Lee, S., & Rudy, B. (2016). GABAergic interneurons in the neocortex: From

cellular properties to circuits. *Neuron*, *91*(2), 260–292.

Turrigiano, G. (2012). Homeostatic synaptic plasticity: Local and global mechanisms for

stabilizing neuronal function. *Cold Spring Harbor Perspectives in Biology*, *4*(1), a005736.

Weihberger, O., Okujeni, S., Mikkonen, J. E., & Egert, U. (2013). Quantitative examination of

stimulus-response relations in cortical networks in vitro. *Journal of Neurophysiology*, *109*(7),

1764–1774.

Xue, M., Atallah, B. V., & Scanziani, M. (2014). Equalizing excitation-inhibition ratios across

visual cortical neurons. *Nature*, *511*(7511), 596–600.

Zhu, J., Jiang, M., Yang, M., Hou, H., & Shu, Y. (2011). Membrane potential-dependent

modulation of recurrent inhibition in rat neocortex. *PLoS Biology*, *9*(3).

Zucker, R. S., & Regehr, W. G. (2002). Short-term synaptic plasticity. *Annual Review of*

Physiology, *64*(4), 355–405.

Tables

<u>Table 1. Evoked spike probability and inhibition duration correlations with</u>

<u>spontaneous firing rate.</u>

		Inhibition duration and spontaneous firing rate			
		Uncorrelated	Positive Correlation	Negative Correlation	Total
Evoked spike probability and spontaneous firing rate	Uncorrelated	10	18	37	65
	Positive Correlation	86	77	300	463
	Negative Correlation	12	13	32	57
	Total	108	108	369	585

Figures

Figure 1 – Experimental setup and timeline

Macaques calmly sat in a chair receiving apple smoothie reward through the experiment.

Cathodic, 200us phase width, single-pulse ICMS was delivered to one channel of the Utah array

in primary motor cortex while unit responses were recorded across the array. Each session

consisted of a prestimulus and stimulus epoch.

<u>Figure 2 – Detection of evoked spikes and inhibition</u>

A. Example of filtered data trace. The inset shows a stimulus followed by an evoked spike after

3.5 ms. **B.** Defining evoked spikes. A PSTH with 0.5 ms bins was generated. Peaks after the time

of stimulation greater than the upper threshold (mean+2 standard deviations of -20 to -2ms in the

PSTH) down to the lower threshold (mean-2 standard deviations) were called evoked spikes. **C.**

Defining inhibition. Top: PSTH. Bottom: corresponding raster plot. The timing of the first spike

that is not an evoked spike was recorded following each stimulus. The red line shows the median

inhibition averaged over 30 second bins with 1 second steps.

<u>Figure 3 – Stimulation during inhibition</u>

Two PSTH examples with 1 ms bins of double- and triple-pulse stimulation in which subsequent

pulses arrive during the inhibitory response of the previous pulse (left). Spikes were readily

evoked even when stimulating during the inhibitory response. Aligning the PSTHs to the final

stimulus pulse (right) shows that the inhibition restarts at each stimulus pulse. Each condition

consisted of 1500 stimuli.

<u>Figure 4 – Effect of distance from stimulus site</u>

A. Evoked spike probability with respect to distance from the stimulated site. **B.** inhibition

duration with respect to distance from the stimulated site. **C.** Histogram of evoked spike timings

split into sites close (<1mm) to the stimulated site and all other sites. The line shows the cubic

interpolated moving average over 3 bins.

<u>Figure 5 – Effect of stimulus amplitude</u>

A. An example raster plot of a spike over different stimulus current amplitudes delivered for five

minutes each at 2 Hz. Blue dots represent evoked spikes and the red line shows the median of the

inhibition duration binned every 30 seconds with 1 second steps. **B.** An example evoked spike

probability as a function of amplitude and **c)** inhibition duration as a function of stimulus

amplitude. The dashed lines are fitted sigmoidal curves using least-squares regression.

<u>Figure 6 – Probability of evoking a spike and inhibition duration are related to</u>

<u>spontaneous firing rate</u>

A. An example of a neuron with positively correlated firing rate (black) and evoked spike

probability (blue) over 10 minutes. The rate and probabilities are averaged over 30 second bins

with 1 second steps. **B.** The same unit had a negatively correlated firing rate (black) and

inhibition duration (blue). The inhibition duration is also averaged over 30 second bins with 1

second steps. **C.** Distance from the stimulated site for units with uncorrelated, positively

correlated, and negatively correlated evoked spike probabilities (left) or inhibition duration

(right) with firing rate.

<u>Figure 7 – Probability of evoking a spike is dependent on the timing of stimulus</u>

Three examples of spike autocorrelations (Auto Cor), and probability of a stimulus evoking a

spike relative to timing from the most recent spontaneous (Spont) and evoked (Evoked) spike.

Top and middle show two different autocorrelation waveforms with correlated evoked spike

probability. They are not aligned in the bottom example; this typically occurs when the

probability of evoking a spike is high. All traces show a moving average using 5 ms bins with a 1

ms step size. R_s is the correlation coefficient between Auto Cor and Spont, R_e the correlation

coefficient between Auto Cor and Evoked, and p_s and p_e are the corresponding p-values of

correlation.

<u>Figure 8 – Changes in evoked spike probability and inhibition duration with</u>

<u>repetitive stimulation</u>

A. An example of changes in the likelihood of evoking spikes at different sites across the array

with 10 Hz repetitive stimulation at one site. Insets show examples of a decrease and increase of

evoked spike probability over time. **B.** The likelihood of evoked spike probability (left) and

inhibition duration (right) changing over time (Change), and the likelihood that the changes are

decreasing (Decrease) dependent on stimulation frequency. Labeled p-values show all

statistically significant differences. Evoked spike probability is more likely to change over time

with higher stimulation frequencies, and the changes are more likely to be decreasing. Inhibition

is also more likely to change with higher stimulation frequency. **C.** Likelihoods of changes

dependent on the distance of the recorded site from the stimulated site. Labeled p-values show all

statistically significant differences. It is less likely for the probability of evoked spikes at sites

closer to the stimulated site to change over time.

<u>Figure 9 – Changes in evoked spike latency with repetitive stimulation</u>

A. An example of evoked spike latency changing over time. Evoked spikes that had changes in

their latency over time were **(B)** typically at sites closer to the stimulated site and **(C)** occurred

when delivering stimulation at a higher frequency.

<u>Figure 10 – Relationship between evoked spikes and inhibition</u>

A. Example PSTH with 1 ms bins following stimuli that evoked spikes and those that did not

demonstrating similar inhibitory response. Each condition consisted of 1500 stimuli. Smoothed

(2 ms wide gaussian moving window) PSTH of the two different stimulation classifications.

Note the inhibition is extremely similar for both. **B.** A comparison of the inhibition strength in

the two different classifications for 470 units. There was no statistically significant pairwise

difference between the two groups. **C.** Comparisons of distance from the stimulus site of units

with uncorrelated, positively correlated, and negatively correlated evoked spike probability and

inhibition duration.

Figure 11 – Comparisons between regular spiking and fast spiking neurons

A. Example of regular spiking (RS) and fast spiking (FS) waveforms. **B.** Distribution of

measured area under the curve of all spikes. Vertical dotted line indicates the classification

boundary – fast spiking falls to the left and regular spiking to the right. **C.** Spike width

distribution of all recorded spikes and spikes that were evoked by stimuli. **D.** Distance from the

stimulated site of evoked spikes grouped by spike width (p=0.03). **E.** Evoked spike probability

distribution (left) and inhibition duration distribution (right) of RS and FS neurons. **F.** Spike

width distributions of evoked spike probability change vs no change over time (left) and

inhibition duration change vs no change over time (right). **G.** Spike width distributions of evoked

spike probability decrease vs increase over time (left) and inhibition duration decrease vs

increase over time (right).

<u>Figure 12 – Stimulation response circuitry</u>

Model of ICMS activated circuitry that generates excitatory and inhibitory responses through

feedforward and feedback mechanisms. Stimulation activates axons projecting to the recording

site. Sites closer to the stimulated site have more complete activation compared to sites further

from the stimulated site. Possible direct connections from the stimulated site to the far site would

also be sparsely activated.