Temporal and Spatial Resolution of a Protein Quake that Activates Hydrogen Tunneling in Soybean Lipoxygenase

Jan Paulo T. Zaragoza,[1,2,#] Adam R. Offenbacher,[1,2,4,#] Shenshen Hu,[1,2] Christine L. Gee,[1,3] Zachary M. Firestein,[2] Natalie Minnetian,[2] Zhenyu Deng,[2] Flora Fan,[2] Anthony T. Iavarone,[1,2] and Judith P. Klinman[1,2,3*]

[1]California Institute for Quantitative Biosciences, University of California Berkeley; Berkeley, California 94720, United States.

[2]Department of Chemistry, University of California Berkeley; Berkeley, California 94720, United States.

[3]Department of Molecular and Cell Biology, University of California Berkeley; Berkeley, California 94720, United States.

[4]Department of Chemistry, East Carolina University; Greenville, North Carolina 27858, United States.

[#]These authors contributed equally.

*Corresponding author. Email: klinman@berkeley.edu

Abstract: The enzyme soybean lipoxygenase provides a prototype for deep tunneling mechanisms in hydrogen transfer catalysis. This work combines room temperature X-ray studies with extended hydrogen deuterium exchange experiments to detect a radiating cone of aliphatic side chains that extends from the iron active site of SLO to the protein-solvent interface. Employing eight variants of SLO, nanosecond fluorescence Stokes shifts have been measured using a probe appended to the identified surface loop. We report a remarkable identity of the enthalpies of activation for the Stokes shifts decay rates and the millisecond C-H bond cleavage step that is restricted to side chain mutants within the identified thermal network. While the role of dynamics in enzyme function has been predominantly attributed to a rapidly equilibrating protein conformational landscape, these new data implicate a rapid and cooperative protein quake as the origin of the thermal activation of SLO. A mechanism of catalysis is presented that combines a distributed conformational landscape with a long range and site-specific thermal quake.

The contribution of protein scaffold motions to the modulation of reaction barrier height and shape remains a major unresolved component in our ability to describe the physical properties underlying enzyme catalysis. There is a growing acceptance of the role of protein conformational sampling and selection in the determination of reaction rate and regulation of enzyme catalysis, formulated in the context of a Boltzmann distribution of thermodynamically equilibrated protein sub-states that are primarily populated by low energy configurations accessible near room temperature (1-3). As expected from the anisotropic topology of almost all folded proteins, site specificity in the achievement of functionally relevant protein sub-states is apparent from both computational analyses (4) and room temperature X-ray crystallographic measurements (5-7). Though more elusive, the involvement of real-time couplings of protein dynamical motions to the chemical steps of catalysis has also been addressed through the application of transition path sampling (8-10). This approach leads to the computation of local atomic motions that accompany reaction barrier crossings, but has not yet been adopted to include more widespread and remote protein motions.

A related but somewhat different context for our understanding of the role of the protein scaffold in catalysis comes from the central role of thermal activation among the vast majority of enzyme catalyzed reactions, with only a small, albeit important, handful of enzymes requiring photochemical initiation (e.g., bacterial reaction centers (11, 12), protochlorophyllide oxidoreductase (13), DNA photolyase and cryptochromes (14), and fatty acid photodecarboxylase (15)). The temperature dependence of enzyme-catalyzed reaction, in certain instances, has been analyzed in terms of changes in heat capacity ($\Delta C^{\ddagger}_p$), attributed to a reduction in active site flexibility at the point of barrier crossings (16, 17). Temperature dependent hydrogen-deuterium exchange coupled to mass spectrometry (TDHDX-MS) has recently emerged as a broadly accessible experimental tool to interrogate heat induced changes in local protein flexibility, which when coupled to function-altering protein mutations provides spatial resolution of activated networks that are found to lead from protein-solvent interfaces to enzyme active sites (18-22). While the thermal activation of enzymes can be rationalized as a redistribution of equilibrated protein states toward increasingly active configurations with increasing temperature, the growing identification of reaction specific thermal networks that are distinctive from each other and independent of protein scaffold conservation points toward scaffold-embedded heat conduits as a source of dynamical excitation (23-30).

The concept of protein quakes arose initially during studies of the photodissociation of CO from myoglobin and remains an active area of research (31-33). With the development of time-resolved X-ray scattering and free-electron laser (XFEL) techniques, it has become possible to monitor the spatial and time dependencies of anisotropic heat dissipation from a protein interior to the solvent (34-36). In the present work, we turn this concept around to focus on the role of anisotropic heat movement inwards, from a protein solvent interface toward the site of substrate binding and catalysis. It has been hypothesized that the transmission of thermal energy from the solvent bath to the protein surface provides the energy required to overcome the chemical reaction barrier (3, 37, 38). Addressing this question has been greatly facilitated by the large body of emergent evidence for quantum mechanical tunneling in enzymatic hydrogen transfer reactions.

The adaptation of Marcus-like models for long range electron transfer to C–H cleavage reactions provides a powerful formalism that can accommodate the multi-dimensional properties of hydrogen tunneling at room temperature (39-45). In particular, the resulting separation of the protein environmental reorganization terms that transiently create the energetic degeneracy and internuclear distances that enable effective H-tunneling from the inherently temperature-independent probability of wave function overlap refocuses structure-function-dynamics studies on the protein scaffold itself and allows its deconvolution from the primary chemical bond cleavage event.

Soybean lipoxygenase (SLO) emerged several decades ago as a paradigmatic system for exploring the properties of thermally activated tunneling in biology (46). SLO catalyzes the production of fatty acid hydroperoxides at (Z,Z)-1,4-pentadienoic moieties via a rate-limiting proton coupled electron (net hydrogen atom) transfer from substrate to an $Fe^{III}(OH)$ cofactor (Fig. 1A). The physiological substrate 9,12-(Z,Z)-octadecadienoic acid (linoleic acid, LA) is converted to 13-(S)-hydroper-oxy-9,11-(Z,E)-octadecadienoic acid (13-(S)-HPOD). The resulting oxygenated lipids and their byproducts are essential for signaling structural and metabolic changes in the cell, including seed growth and plant development (47). Native SLO catalysis exhibits a fairly weak temperature dependence ($E_a \sim 2.0$ kcal/mol), a small Arrhenius prefactor ($A_H < 10^5$ s^{-1}), and a greatly inflated kinetic isotope effect (KIE) on k_{cat} ($^D k_{cat} = 81$) (40, 48) that rises to $^D k_{cat} = 500 - 700$ for an enzyme variant that has been mutated at two hydrophobic side chains surrounding the reactive carbon of bound substrate (49). The aggregate structural and kinetic properties of SLO played a key role in the development of vibronically non-adiabatic analytical expressions that are now routinely applied to explain deep tunneling reactions that occur near room temperature (40, 42-44, 50, 51). SLO was also the first protein to be subjected to a combined analysis of the time-, temperature- and mutation-dependent properties of hydrogen deuterium exchange as analyzed by mass spectrometry (19). The resulting identification of a site selective thermal pathway, originating at a solvent-exposed loop and extending toward the buried iron cofactor and substrate binding site, provides a unique opportunity to explore the structural and dynamical features that control C-H activation in enzyme catalysis. Evidence is presented herein for a long-range protein quake as the source of thermal activation during the rate limiting hydrogen transfer step of SLO.

Fig. 1. (**A**) Rate-limiting hydrogen transfer reaction from linoleic acid (LA) catalyzed by an Fe^{III}(OH) cofactor in SLO (46). (**B**) X-ray structure of SLO (PDB:3PZW) depicting the active site Fe, and the network of residues (pink) and thermally activated surface loop (residues 317-334, in dark blue-green) identified from previous HDX-MS experiments (Ref. 19). Inset: The bound linoleic acid substrate (gray) is modeled from previous calculations (51) and overlaid in this structure. Leu754, which is outside the identified network, is shown in purple. The peptide 317-334 is labeled in blue-green.

Results

The catalytically linked thermal network in SLO extends beyond the TDHDX-MS defined pathway. The originally defined thermal pathway for SLO (19) was found to be comprised of three regions: (i) the substrate binding site in proximity to the Fe^{III}(OH) cofactor, (ii) an intermediate region comprised of Ile552 and Val750, and (iii) a loop that interacts directly with

solvent (Fig. 1B). The aliphatic residues Ile552 and Val750 are positioned to act as mediators between the active site side chains Leu546 and Ile553 and the residues that extend to the surface loop. Of particular interest, neither Ile552 nor Val750 is in direct contact with the reactive carbon, raising the question of their importance and impact to catalytic proficiency. To address this issue, we examined the effect of alanine substitution at Ile552 and Val750, obtaining a suite of kinetic parameters (Table 1). While the magnitude of the difference in the enthalpy of activation for the reaction with H- vs. D-labeled LA substrate, ΔE_a, is reduced for V750A, its value is within the experimental error of WT and the value of ΔE_a for I552A is unchanged. An increase in ΔE_a has been shown to map to a loss of precision in the local positioning of the reactive carbon (C11 of the LA substrate) for SLO (52), as well as for a wide range of H-transfer enzymes (53). In accordance with this behavior, evolutionary selection was recently applied to a primitive form of dihydrofolate reductase (DHFR), showing a progressive decrease in ΔE_a as the evolved enzyme approached the activity of native DHFRs (54). From this study of I552A and V750A, we conclude that ΔE_a is largely unaffected, with the impact of such mutations appearing primarily as a reduction in the enthalpy of activation ($E_a(H)$) that is accompanied by reduction of the first-order rate constant (k_{cat}) at 30 °C. The changes in $E_a(H)$ suggest a perturbation (loosening) of structure away from the configuration that is optimal for catalytic turnover in WT, with the impact of I552A exceeding that of V750A.

Table 1. Comparative kinetic parameters for Ala variants at positions 552 and 750 within the thermal network of SLO[a]

SLO variant	k_{cat}, s^{-1}	$^D k_{cat}$	E_a, kcal/mol	ΔE_a, kcal/mol	Reference
WT	297 (10)	81 (5)	2.1 (0.2)	0.9 (0.2)	Ref. (40)
I552A	81 (2)[b]	60 (2)[b]	-0.3 (0.2)	1.2 (0.5)	This work[a]
V750A	218 (8)[b]	69 (4)[b]	1.0 (0.4)	0.3 (0.4)	This work[a]

[a]Temperature dependent kinetics data had been reported (55) but were not previously analyzed for E_a and ΔE_a. The full range of additional mutants formerly characterized in this fashion is summarized in Table S1.

[b]For comparison under similar protein and substrate purification procedures (19), data for WT were recollected using the enzyme preparation of the present study, generating k_{cat} = 359 (7) s^{-1}, $^D k_{cat}$ = 57 (2), E_a = 2.4 (0.2) kcal/mol, and ΔE_a = 1.1 (0.1) kcal/mol.

With the goal of more fully understanding the contribution of residues Ile552 and Val750 to the thermal pathway in SLO, additional independent experimental directions were pursued. First, TDHDX-MS was extended to I552A, resulting in the same set of non-overlapping peptides as obtained for WT and other variants (19, 56) (*cf.* fig. S1). The majority of the I552A-derived peptides display nearly identical exchange behavior to WT across the 5 temperatures (10, 20, 25, 30, and 40 °C) analyzed (*cf.* Data S1). The peptide of particular interest is 317-334 that represents the surface loop that was previously shown (19) to undergo correlated alterations in the activation energy for HDX $E_a(k_{HDX})$ and $E_a(k_{cat})$ while interrogating the variants I553G and L546A in relation to WT. The new data for I552A indicate an $E_a(k_{HDX})$ = 3.2(3.7) kcal/mol, within experimental error

of zero, analogous to its $E_a(k_{cat})$ = -0.3(0.2) kcal/mol) (Table S2). As shown in Fig. 2 (red line) the new data for I552A fall on the original correlation of $E_a(k_{HDX})$ and $E_a(k_{cat})$, in support of the inclusion of Ile552 within the defined thermal pathway. We note that mutations at Leu754 fall off the line correlating $Ea(k_{HDX})$ to $Ea(k_{cat})$ (Fig. 2), and this feature serves as a valuable control for time and temperature dependent Stokes Shift decay analyses presented below. We also noted an unusually large perturbation from I552A on the TDHDX-MS of peptides 541-554 and 555-565 residing near the inferred substrate portal, with peptide 541-554 of I552A indicating a decrease in $E_a(k_{HDX})$ from 23.7(2.0) kcal/mol for WT to 5.64(0.5) kcal/mol (fig. S2); these data are consistent with an earlier report of an increase in k_{off} values for substrate release in I552A, attributed to a loosening of the substrate portal that is adjacent to the active site (55).

Fig. 2. Temperature-dependent HDX-MS analysis. Relationship between the HDX enthalpic barrier (E_a for $k_{HDX(avg)}$) at peptide 317-334 and the enthalpic barrier for hydrogen tunneling (E_a for k_{cat}). New data for I552A are shown and compared with data for other mutants from Ref. (56).

Room temperature (RT, 277 – 300 K) X-ray studies have been shown to reveal multiple, weakly populated sub-states within a protein conformational ensemble (5). The multi-conformer models obtained from qFit (57) uncover local anisotropic protein dynamics information that is supported by NMR measurements, and reflects multiscale motions of the protein backbone and side chains (7). Close inspection of the RT electron densities for WT SLO published earlier indicated a number of alternate side chain conformers for key active site residues (19). Herein we first applied RT X-ray studies (T = 300K) to the activity altering variants, I553G and L546A (56),

finding that the I553G variant decreases the conformational space accessed by its proximal Leu546 side chain (fig. S3). The L546A mutation, on the other hand, leaves Ile553 unperturbed while decreasing the electron density of an alternate conformer for a second proximal residue Ile552. With this observation, we proceeded to examine the impact of both I552A and V750A mutations on nearby residues via RT X-ray structural analysis (Table S3). It is expected that changes in side chain conformations following site specific mutagenesis may radiate throughout the protein, however the impacts of I552A and V750A are found to be concentrated within a region that neighbors the TDHDX-MS detected thermal pathway (Fig. 3). The I552A and V750A mutations show reciprocal differences in alternate conformers at the two positions. Specifically, V750A leads to a loss of alternate conformer in Ile552, while I552A results in a different configuration of the Val750 side chain from that of WT. Additionally, I552A and V750A mutations result in a loss of alternate conformers for the adjacent Leu546 and the more remote Ile746 and Leu262 side chains (Fig. 3A).

Fig. 3. Room temperature X-ray structure analysis. (A) Impact of I552A and V750A mutation on the 300 K X-ray structure electron densities of their respective side chains as well as those of Leu546, Ile746 and Leu262. **(B)** Connectivity of residues from the Fe active site to the surface loops (loop 1: 317-334 (blue-green), loop 2: 284-299 (brown)) in the 300 K X-ray structure of WT SLO (PDB:5T5V). Multiple side chain conformers along the thermal network are represented in pink and purple.

The above analyses provide strong support for the role of both Ile552 and Val750 as mediators between the active site side chains of SLO and an extended thermal network that terminates at the protein-water interface. As a guide for visualizing the entire network, side chain interactions have been traced from Leu546 to the surface loops comprised primarily by the region of protein represented by amino acids 317-334 and to a lesser extent the region 284-299, a second peptide shown to undergo correlated motions in the Ea for HDX-MS and catalysis (19) (Fig. 3B). Using a cut-off of 4 Å for van der Waals interactions between adjacent side chains, an extended network of amino acids is detected beyond the TDHDX-MS derived pathway. In particular, Ile552 is seen to act as a central lynchpin, exhibiting steric interactions with Leu546, Val750, Leu262, and Ile746. As shown in Fig. 3B, a clearly defined, cone-shaped region of protein with activity-related alternate conformers connects the active site to the solvent exposed loop. The abundance of hydrophobic residues in this extended network led us to consider other possible clustering of hydrophobic residues in SLO. It has been proposed that the sidechains of Ile, Leu, and Val often form hydrophobic clusters (58, 59) which serve as cores of stability and prevent intrusion of water molecules. The hydrophobic clusters in SLO (fig. S4) were computed using a CSU algorithm (60). Of considerable interest for how thermal networks may be constructed, the largest calculated hydrophobic cluster in SLO is comprised of 32 residues and includes the Ile839 ligand to the Fe center, the active site residues that are in contact of the linoleic acid substrate (Ile553, Leu546, Leu754) and all of the residues that are experimentally assigned to a thermal activation pathway.

Demonstration of a dynamically activated thermal network in SLO that stretches from the protein/solvent interface to the active site. To examine dynamical motions within the thermal loop in SLO, the fluorescent probe 6-bromoacetyl-2-dimethylaminonaphthalene (Badan, BD) was incorporated at position 322 at the tip of the identified thermal network (Fig. 4A) and at a remote control position 596 via thiol modification (61). Acetyl-2-dimethylaminonaphthalene-based dyes such as Badan are used as sensors to probe changes in local solvation dynamics in proteins following photo-excitation (62). The main mechanism for fluorescence modulation is through interactions of the surrounding solvent with the electronically excited state of the probe, which in the case of Badan leads to polarization of an N,N-dimethylamino group and a carbonyl substituent. Nearby Trp residues in proteins have been shown to quench Badan fluorescence via electron transfer (63). In Q322C-BD SLO, the nearest Trp residue, Trp340, to the Badan probe at position 322 is 30-40 Å away. Molecular dynamics simulations of Q322C-BD show that this distance does not change significantly (fig. S5), making artifactual quenching of the Badan probe at the 322 position an unlikely complication. In an earlier preliminary work (61), the temperature dependence of the Badan Stokes shift decay was observed to be negligible at a control loop at position 596. When the probe was appended to the end of the identified thermal loop at position 322, the enthalpies of activation for Stokes shifts matched that for catalytic C-H activation $E_a(k_{cat})$. The striking observation of an exact correspondence between the enthalpic barrier for a millisecond catalytic step and for a nanosecond Stokes shift decay process suggested a direct coupling of distal protein motions surrounding the Badan probe to active site motions controlling catalysis in the WT enzyme.

In this work we pursue Stokes shift measurements with SLO using a wide range of enzyme variants with distinctive trends in $E_a(k_{cat})$ and $E_a(k_{HDX})$. Four of the variants chosen for study are either in contact with bound substrate and/or with each other: I553G, I553A, L546A, and I552A (Fig. 2A). Substitution of residues at positions 553 and 546 in the substrate binding pocket leads to perturbations that either increase (L546A), decrease (I553G), or retain (I553A) the same $E_a(k_{cat})$ as WT (40, 64) (Table S1). Ile552, which is located proximal to Leu546, but is not in direct contact with substrate, produces a decreased $E_a(k_{cat})$ relative to WT (Table 1). The L754A variant was also investigated as internal control. While Leu754 is positioned on an opposite face of the reactive carbon of substrate from Leu546, and shows an elevated enthalpic barrier for catalysis, TDHDX-MS measurements (56) indicated little or no change relative to WT on $E_a(k_{HDX})$, in support of Leu754 residing outside the pathway for thermal activation (*cf.* Fig. 2). The impacts of L754A on rate and catalytic activation energy, in turn, have been attributed to an expanded active site that incorporates additional water molecules, while altering the active site microenvironment and Fe active site geometry (56). Stokes shift controls involving double mutations with L754A as the parent were also interrogated, involving I553A/L754A and L546A/L754A that exhibit some of the largest observed increases in activation enthalpy for catalysis relative to WT (65) (Table S1). The full set of eight SLO variants interrogated display $E_a(k_{cat})$ values from *ca.* 0 to 10 kcal/mol, providing a broad range of catalytic properties (Table S1) from which to test the correspondence between the magnitude of activation energies for Stokes shifts decay rates and for the rates of catalysis.

Each of the above-described variants was mutated at Gln322 to cysteine and reacted with the thiol-reactive alkyl bromide group of Badan (BD) according to protocols established with WT SLO (61) (Fig. 4A). The resulting bioconjugates were characterized by electrospray ionization mass spectrometry (ESI-MS) to confirm 1:1 adduct formation (Table S4). To verify the fidelity of the probe attachment to Q322C, liquid chromatography-tandem mass spectrometry (LC-MS/MS) was performed on the pepsin digests of each SLO-BD conjugates (fig. S6), corroborating the modification at position 322C, with a mass addition (+211 Da) corresponding to the 6-acetyl-2-dimethylaminonaphthalene group. In all instances, the Cys mutation and attachment of Badan at Gln322 position yields proteins with almost identical kinetic parameters (k_{cat}, E_a) to the parent SLO (Table S5), making these constructs well-suited for probing protein dynamics.

Fig. 4. Temperature-dependent nanosecond fluorescence Stokes shift analysis of a fluorescent probe on a remote surface loop in SLO. (**A**) Location of the mutated active site residues Ile553, Leu546, Ile552, and Leu754 in relation to the thermally-activated loop (317-334, blue-green), and the Badan-labeled Q322C residue (green). The Badan orientation is based on the optimized structure from Ref. (61). (**B**) Representative time-resolved emission spectra (TRES) at 30 ˚C, pH 9.0 for the L546A SLO mutant. (**C**) Representative time and temperature dependence of the Stokes shift decay rates of L546A. (**D**) Arrhenius plots for the nanosecond Stokes shift decays ($1/\tau_1$, $1/\tau_2$) of L546A.

Fig. 5. Comparative activation energies for the Stokes shift decay rates $1/\tau_2$ ($E_a(1/\tau_2)$) and for catalysis ($E_a(k_{cat})$). The red line indicates residues that are along the thermal network. Given the very close overlap between the datapoints for I552A and I553G, the datapoint for I552A is shown in magenta to allow distinction from I553G. The blue line represents the single mutant L754A together with two double mutants, I553A/L754A and L546A/L754A. In this case, the temperature dependence of the observed Stokes shift decays arises solely from the on-network mutations (I553A or L546A, labeled red).

Time-resolved emission spectra (TRES) were constructed from the steady state (fig. S8) and time-resolved fluorescence decay data upon excitation of Badan at 373 nm (See SI Materials and Methods). The time-dependent Stokes shift is attributed to solvent/environmental reorganization around the excited state probe, and is manifest as continuous red shifts in emission spectra until the steady state emission wavelength is reached (66-68). The TRES for the Q322C-BD labeled SLO mutants at 30 ˚C are shown in fig. S8, with total Stokes shifts varying from ~700-1800 cm^{-1} (Table S6). The Stokes shift decay curves (fig. S9) were fit to a biexponential function, indicating a fast ($\tau_1 \sim 0.5$ ns, ~60%) and a slower ($\tau_2 \sim 5$-8 ns, ~40%) component (Table S6). This bimodal behavior has been shown to be nearly universal and is typically referred to as the bimodality of the reorientational response of biological water (69). The shorter lifetime (τ_1) is assigned to the unconstrained (free) water in the biological water layer and τ_2 is ascribed to the nanosecond reorganization of protein associated with constrained (bound) water (69). Nanosecond Stokes shift decay experiments have been reported for a number of other enzyme systems that

include the analysis of a fluorescent probe attached to the surface of a protein tunnel in haloalkane dehalogenase (70) and to the active site of glutaminyl-tRNA synthetase (71). Molecular dynamics simulations support the view that water molecules hydrating the protein surface will undergo restricted translational and rotational motions (72, 73).

Arrhenius plots of the Stokes shift decay rates in SLO variants (fig. S10) reveal distinctive temperature dependences in the slower component (τ_2), but not the fast component (τ_1) (*cf*, Fig. 5 and fig. S11). In WT SLO, both τ_1 and τ_2 were seen to be temperature dependent (61), however temperature independent values for τ_1 appear to be the more general case. The previously reported viscosity independence of k_{cat} in SLO (65, 74) is consistent with a primary role for the hydration shell (related to τ_2) rather than the bulk solvent (related to τ_1) in the enzymatic activation barrier (33). The activation energies (E_a) for τ_2 in SLO range from ~0-4 kcal/mol (Table S8) and are similar to reported E_a values for the nanosecond dynamics of a solvent exposed tryptophan in a thermophilic ADH (75, 76) and the picosecond hydration dynamics around solvent-exposed tryptophan in Dpo4 (77), ApoMb (78), and SNase (79). The comparison of the experimental activation energies for Stokes shift decay rates ($1/\tau_2$) and for k_{cat} as a function of site-specific mutagenesis is presented in Fig. 5. The data indicate a remarkable 1:1 correspondence for WT SLO and four variants within the thermally activated network, I553G, I552A, I553A, and L546A (Fig. 5, red line), generating a line with slope of 1.0(0.1), $r^2 = 0.99$. The L754A mutant that lies outside the thermal network (56) (see Fig. 2, blue data point) acts as an internal control; in this case, the correlation between the temperature dependence of Stokes shifts and catalysis is displaced to the right (Fig. 5, blue line). Note that for the double mutants that contain both on-network and off-network mutations, I553A/L754A and L546A/L754A, the temperature dependence of the Stokes shift decay arises solely from the on-network mutation (I553A, L546A). With 3 data points, this second line of L754A-containing variants displays a slope of 0.7(0.1) and a correlation coefficient of $r^2 = 0.99$. The results with L754A mutation indicate that the equivalence of E_a values for catalysis and Stokes shift decay is restricted to residues that have been assigned to the thermal network.

In light of the unprecedented correspondence between the E_a values for Stokes shift decay (τ_2) and for k_{cat}, we considered it important to rule out unanticipated impacts from thermally induced protein destabilization and possible global unfolding in the SLO samples interrogated herein. Early cryogenic X-ray studies of hydrophobic side chain variants of apo-SLO had indicated virtually no change in protein structure relative to WT (49, 64), while subsequent advanced ENDOR studies of the SLO-substrate complex revealed a progressive elongation of hydrogen donor-acceptor distances as the volume of hydrophobic side chains was reduced (52, 80). The cryogenic X-ray behavior of SLO is consistent with similar studies of other enzyme systems that show little or no change to overall protein structure following a reduction in the size of interior hydrophobic side chains, with an incursion of water molecules into enlarged interior cavities seen only rarely (81-84). Differential scanning calorimetry (DSC) has been used to provide quantitative estimates of the impact of changes in hydrophobic side chain volume on protein unfolding

thermodynamics (ΔH°) relative to native enzyme (ΔΔH° = ΔH°(mutant) − ΔH°(WT)) (81, 85, 86). As summarized in Table S9 and fig. S12, DSC analysis of all 5 SLO variants located within the protein's thermal network yield values for ΔΔH° in the range of those found for hydrophobic mutations in the prototypic and much smaller enzyme lysozyme (81, 85, 86) and are within experimental error of each other (Table S9). The magnitude of the melting point values is also informative, with little difference seen among the values of WT, L546A and L546A/L754A (DM) together with a decrease by *ca.* 4 ˚C in the case of I553G and I552A. The high overall T_m values of SLO are similar to what was measured for a related isoform (87) consistent with the large size and compact global structure of this protein, and are 20 – 25 ˚C above the temperature ranges of the kinetic experiments reported herein. Of note, the decreases in T_m for I553G and I552A track with the decrease in their $E_a(k_{cat}) \sim 0$ kcal/mol (Table S1 and Fig. 5). Consistent with the TDHDX-MS evidence of enhanced flexibility at the surface loop (19) (Fig. 2), both variants I553G and I552A are concluded to undergo local destabilization at this region ($\Delta S° > 0$ and $\Delta H° > 0$). Restructuring of initially disordered ground state complexes ($\Delta S° < 0$ and $\Delta H° < 0$) may be expected as a prerequisite for the productive thermal activation of tunneling ($\Delta H^{\ddagger} > 0$), resulting in compensating temperature dependences and net experimental E_a values within error of zero. We conclude that the pattern seen for I553G and I552A in Fig. 5 is especially notable and in support of site-specific motions within the protein scaffold as the origin of corresponding E_a values for Stokes shift decay and catalytic rate constants.

Discussion

Room temperature X-ray crystallography of proteins has gained attention as a powerful tool in detecting amino acid side chain disorders that can be related to enzyme function (5, 6). In the present study we analyzed RT X-ray structures for single site mutants of SLO with demonstrated impacts on the properties of hydrogen tunneling from C11 of the linoleic acid substrate to the active site $Fe^{III}(OH)$ cofactor (19, 56) (Fig. 1A). Electron density analyses (fig. S3 and Fig. 3B) extend the thermal transfer pathway identified by TDHDX-MS (19) to an island of primarily hydrophobic side chains that reach from the protein/solvent interface to the active site (Fig. 3A). Both methods report on thermally-averaged protein sub-states that undergo functionally relevant responses to either mutation (RT X-ray studies) and/or to changes in temperature (TDHDX-MS). In the case of TDHDX-MS, the linear relationship between $E_a(k_{HDX})$ and $E_a(k_{cat})$ (56) (Fig. 2, red line) provided initial insights into the location and composition of a thermal network for enzyme catalysis in SLO. An important caveat of the TDHDX-MS derived correlation is the significant difference in the magnitudes of $E_a(k_{HDX})$ in relation to $E_a(k_{cat})$, as expected for inherently different processes that reflect local and fully reversible protein unfolding events ($E_a(k_{HDX})$) vs. the tuning of active site electrostatics and distances that promote hydrogenic wave function overlap between the donor and acceptor atoms ($E_a(k_{cat})$).

The presented nanosecond Stokes shift analyses differ in a fundamental way from the TDHDX-MS experiments, uncovering activation energies for a photo-induced environmental reorganization around a surface attached probe in SLO that are *identical* to the measured activation energies for cleavage of the C-H bond of substrate (Fig. 5). A limited number of previous studies

have shown corresponding enthalpies for distinct kinetic processes within a single system, that include a comparison of Stokes shifts and fluorescence anisotropy using the method of tryptophan scanning in DNA polymerase IV (77) and a comparison of the rate of solvent dielectric relaxation to changes in Mössbauer spectra following the photo-excitation of the carbon monoxide-heme complex in myoglobin (33). However, neither of these studies was able to address the physical origins and site specificity of heat activation in the context of an enzyme catalyzed bond cleavage reaction. The observation of the same enthalpy of activation among variants of SLO for two distinctive chemical processes that take place with rate constants that differ by *ca.* 10^6-fold and at a distance of at least 30 Å apart, provides compelling evidence for a shared protein quake that connects thermal activation at the protein/solvent interface to the primary bond cleavage step in SLO. The regional specificity of this heat flow, as first revealed from in depth studies of both TDHDX-MS and RT X-ray, illustrates the importance of anisotropic thermal conduits as a means of generating productive activated enzyme-substrate (ES) complexes that also avoid interference from disruptive protein motions that could slow down or eliminate catalysis. While the detection of outward directed protein quakes during the photoexcitation of protein-bound chromophores has been previously described (34), the data presented here implicate a functional and inward-directed protein quake in SLO that reaches from a localized protein surface toward a deeply buried active site.

Discussions of the role of protein dynamics in enzyme catalysis have been primarily focused on the role of conformational landscapes that are comprised of large numbers of thermodynamically equilibrated states, a sub-set of which catalyze the efficient conversion of substrate to product (88-91). One key question from the present work is the inter-relationship between a thermal quake from the solvent bath to an enzyme active site and the distributed conformational landscape. A revised working model of protein dynamics, Fig. 6, begins with the growing body of experimental evidence for protein embedded networks (18-22) that provide pathways for heat activation in thermally initiated enzyme reactions. The principle of conformational selection in protein function, as conceptualized by Frauenfelder for ligand binding dynamics in myoglobin (92), and developed by Kern and co-workers for catalyzed reactions (90, 91) can be readily integrated with this feature, via substrate induced shifts in the conformational ensemble of enzyme-sub-state complexes toward configurations that place the reactive components of an enzyme active site in close proximity to the region of protein capable of efficient heat transmission. We note that *both* pre-equilibrated protein sampling modes and long range thermal transfer processes arise via *transient* events that create energy states higher than the dominant enzyme ground state structures. A difference between these two components of protein dynamics is an expected much smaller deviation of activated states from ground states in the case of a distributed conformational landscape that gives rise to a wide range of protein configurations with energy differences close to RT. The model in Fig. 6 provides a context from which to rationalize the large and growing body of evidence from directed evolution (93-96) and high throughput screening (97) that regions distal to the active site play essential roles in enzyme catalysis (98).

Fig. 6. A revised working model of protein dynamics. (**A**) Two distinct types of protein motions contribute to enzyme catalysis. An initially formed ES complex first undergoes conformational selection to achieve refinement in the position of bound substrate relative to active site cofactor and surrounding protein side chains. The resulting family of ES_{active} states is positioned to take maximal advantage of an embedded site-specific network that provides transient activation via thermally activated protein quakes. (**B**) Left: Two-dimensional reaction coordinate illustrating contribution of protein scaffold dynamics to thermal activation in hydrogen tunneling enzymes. Right: Three-dimensional representation of the free energy landscape for catalysis, adapted from Ref. (1). As illustrated, the initial ES complex (blue) undergoes conformational sampling among a large number of possible ground state configurations (green and yellow) that are in rapid equilibrium near RT (y coordinate). Productive barrier crossings (red) occur when the transient alignment of substrate, active site side chains and/or cofactor coincides with a protein embedded thermal conduit (x coordinate).

A key question from this work concerns the different time scales for measured fluorescence Stokes shifts (nanoseconds) vs substrate turnover (milliseconds) that differ by *ca.* 10^6-fold. Protein motions are expected to vary greatly in rate and to be dependent on the forces and locality of the motions being studied and to the initiating events. With regard to thermal transfer through a protein, XFEL-based studies of heat evolution in a light activated protein illustrate structural changes that radiate out from the active site to solvent on a time scale of picoseconds (34). This time scale is consistent with experimental and computational studies of a number of well-studied model systems, e.g., myoglobin and albumin, that consistently indicate lifetimes in the range of 1 – 50 ps for heat flow in proteins over distances of tens of angstroms (99-104). None of these studies directly address the question of whether rapid, long distance and cooperative heat flow through a protein may deviate from a fully equilibrated process. In a recent study of IR-detected protein motions following photochemical excitation of a surface probe in bovine serum albumin, an out-of-equilibrium protein quake was proposed with properties of a Frochlich-like condensate comprised of coupled low frequency vibrations (105).

A simple but comprehensive formulation for the observation of a range of rate constants that arise from a shared thermal activation of the protein scaffold is shown in equation 1:

$$k_{obs} = P_{conf} \cdot k_{int} \qquad (1)$$

where the measured rate constant k_{obs} depends on both the primary thermal quake of the protein scaffold, $k_{intrinsic}$, and the probability that the resulting structural rearrangement is sufficient for the process undergoing measurement (according to the fraction of active conformers, P_{conf} = [E_{active}]/[E_{total}] (Fig. 6A). A related F_{conf} term had been earlier introduced into formal rate expressions for non-adiabatic hydrogen tunneling reactions that describe the deep tunneling reaction of SLO, in order to reconcile experimental rate constants that are significantly smaller than the computed tunneling rates (50, 106). In the case of a fully optimized enzyme catalyzed reaction, P_{conf} may be very small, a result of the dependence of productive barrier crossings on a multitude of transiently optimized electrostatic interactions and precise internuclear distances. Conformational sampling may play a key role in increasing the magnitude of P_{conf} into a range compatible with the typical millisecond turnover numbers of enzymes (Fig. 6A), however, in any case, the achievement of the barrier crossing states in enzymes is a comparatively rare event. The magnitude of P_{conf} can reasonably be expected to become much larger in the case of the Stokes shifts behavior of a surface-appended fluorophore, given the availability of a wide range of both protein and solvent interactions for the stabilization of excited state dipoles, as reflected in the observation of Stokes shifts within the picosecond to nanosecond regimes (61, 75-78).

A pictorial representation of Eq (1) is given in Figure 7, where a red cone is used to represent a shared site for heat transfer that originates at the protein-solvent interface and terminates at the active site (labeled 'thermal input'). The magnitude of the observed rate constants resulting from such scaffold activation is represented by the green cone at the top of the figure (labeled 'productive output'). In a separate enzyme system catalyzing hydride transfer from an alcohol substrate to a nicotimamide cofactor, microsecond FRET between an interior Trp residue and the nicotinamide ring of cofactor was reported to occur with an enthalpy of activation

equivalent to that of millisecond catalysis (107), and this behavior has been integrated into Figure 7 as arising from a value for P_{conf} that is greater than the catalytic hydride transfer, yet smaller than P_{conf} for the protein surface Stokes shifts measurements described herein. To summarize, the origin of the different timescales for each kinetic measurement arises from a common, thermally activated protein quake that is accompanied by a different probability for achieving the set of protein configurations capable of supporting the physical or chemical event under measurement, analogous to the entropic component of the familiar rate expressions describing chemical processes (108, 109). A similar formalism has been suggested in the context of solvent-slaved motions that modulate the internal movement of carbon monoxide within the interior of myoglobin (110).

Fig. 7. Time scales for productive responses to a long range protein quake. A model is presented that illustrates how different physical events with widely different rate constants can share a common activation energy. According to Eq (1) and the discussion in the text, a shared protein quake leads to a site-specific transfer of thermal activation from the protein/solvent interface to the active site on a picosecond time scale (indicated by the red cone in the above figure). The response (green cone, labeled productive output) is seen to vary from nanoseconds for Stokes shifts in SLO (present study) to microseconds for FRET behavior in the case of an alcohol dehydrogenase (Ref. (107)); for both processes, the frame of reference is enzymatic turnover on a millisecond time scale.

There has been a recent resurgence and growing interest in understanding and harnessing the role of quantum mechanics, including spin coherence, entanglement and tunneling, in biological and other macroscopic processes (111, 112). Among the unresolved questions is the

quest to determine and quantify the link, and possible synergy, between classical and quantum behavior, and the degree to which noise at the warm temperatures of biology influences the efficiency of quantum effects. The defining feature from the findings presented in this study is the primacy of the extended protein scaffold in determining the chemical transformations (in this case, hydrogen tunneling) that occur within local enzyme active sites. The presented model differs from previous textbook descriptions of enzyme catalysis that are rooted in the principle of "enhanced transition state binding" (113, 114) and have dominated the understanding (and teaching) of biological catalysis for over half a century. Though this particular study is focused on an enzyme reaction that proceeds via a deep tunneling hydrogen transfer (Fig. 6B, left) a combination of distributed protein conformational substates and embedded thermal networks within a protein scaffold is expected to play a key physical role in enzymatic rate enhancements, independent of the reaction catalyzed. The dominant role of the protein scaffold can be visualized through a three-dimensional energy plot (Fig. 6B, right). As illustrated, the primary coordinates for catalysis are an x-coordinate that represents site specific distribution of heat into the protein scaffold to reach the high energy states capable of barrier crossings ($ES^{\ddagger}$, labeled red) that acts in concert with an orthogonal distributed conformational landscape (y coordinate). The latter provides the high specificity with which a small sub-set of the conformational landscape can respond to local anisotropic heat transfer. This model incorporates a functionally linked thermal activation of embedded protein modes into the contributions of protein conformational selection induced by binding of either substrate (in the case of enzyme catalysis) or allosteric effector (in the case of enzyme regulation). The newly presented data and interpretation both emphasize the unique properties of enzyme reactivity relative to the reaction of small molecules in condensed phase and serve as a launching point to answering the long-sought goal of integrating the protein matrix into our understanding of the origins of naturally evolved enzymatic rate accelerations and our ability to carry over this knowledge into rational *de novo* design.

References

1.   S. J. Benkovic, G. G. Hammes, S. Hammes-Schiffer, Free-Energy Landscape of Enzyme Catalysis. *Biochemistry* **47**, 3317-3321 (2008).
2.   J. P. Klinman, A. Kohen, Hydrogen Tunneling Links Protein Dynamics to Enzyme Catalysis. *Annu. Rev. Biochem* **82**, 471-496 (2013).
3.   P. K. Agarwal, A Biophysical Perspective on Enzyme Catalysis. *Biochemistry* **58**, 438-449 (2019).
4.   A. Ramanathan, A. Savol, V. Burger, C. S. Chennubhotla, P. K. Agarwal, Protein Conformational Populations and Functionally Relevant Substates. *Acc. Chem. Res.* **47**, 149-156 (2014).
5.   J. S. Fraser *et al.*, Hidden alternative structures of proline isomerase essential for catalysis. *Nature* **462**, 669-673 (2009).
6.   J. S. Fraser *et al.*, Accessing protein conformational ensembles using room-temperature X-ray crystallography. *Proc. Natl. Acad. Sci.* **108**, 16247 (2011).
7.   R. B. Fenwick, H. van den Bedem, J. S. Fraser, P. E. Wright, Integrated description of protein dynamics from room-temperature X-ray crystallography and NMR. *Proc. Natl. Acad. Sci.* **111**, E445 (2014).
8.   D. Antoniou, S. D. Schwartz, Internal Enzyme Motions as a Source of Catalytic Activity: Rate-Promoting Vibrations and Hydrogen Tunneling. *J. Phys. Chem. B* **105**, 5553-5558 (2001).

9. D. Antoniou, J. Basner, S. Núñez, S. D. Schwartz, Computational and Theoretical Methods to Explore the Relation between Enzyme Dynamics and Catalysis. *Chem. Rev.* **106**, 3170-3187 (2006).

10. S. D. Schwartz, V. L. Schramm, Enzymatic transition states and dynamic motion in barrier crossing. *Nat. Chem. Biol.* **5**, 551 (2009).

11. M. H. Vos, F. Rappaport, J.-C. Lambry, J. Breton, J.-L. Martin, Visualization of coherent nuclear motion in a membrane protein by femtosecond spectroscopy. *Nature* **363**, 320-325 (1993).

12. H. Wang *et al.*, Protein Dynamics Control the Kinetics of Initial Electron Transfer in Photosynthesis. *Science* **316**, 747 (2007).

13. D. J. Heyes, M. Sakuma, S. P. de Visser, N. S. Scrutton, Nuclear Quantum Tunneling in the Light-activated Enzyme Protochlorophyllide Oxidoreductase *J. Biol. Chem.* **284**, 3762-3767 (2009).

14. A. Sancar, Structure and Function of DNA Photolyase and Cryptochrome Blue-Light Photoreceptors. *Chem. Rev.* **103**, 2203-2238 (2003).

15. D. J. Heyes *et al.*, Photochemical Mechanism of Light-Driven Fatty Acid Photodecarboxylase. *ACS Catal.* **10**, 6691-6696 (2020).

16. J. K. Hobbs *et al.*, Change in Heat Capacity for Enzyme Catalysis Determines Temperature Dependence of Enzyme Catalyzed Rates. *ACS Chem. Biol.* **8**, 2388-2393 (2013).

17. M. W. van der Kamp *et al.*, Dynamical origins of heat capacity changes in enzyme-catalysed reactions. *Nat. Commun.* **9**, 1177 (2018).

18. Z.-X. Liang, T. Lee, K. A. Resing, N. G. Ahn, J. P. Klinman, Thermal-activated protein mobility and its correlation with catalysis in thermophilic alcohol dehydrogenase. *Proc. Natl. Acad. Sci.* **101**, 9556 (2004).

19. A. R. Offenbacher *et al.*, Hydrogen–Deuterium Exchange of Lipoxygenase Uncovers a Relationship between Distal, Solvent Exposed Protein Motions and the Thermal Activation Barrier for Catalytic Proton-Coupled Electron Tunneling. *ACS Cent. Sci.* **3**, 570-579 (2017).

20. J. Zhang, Balsbaugh, J.L., Gao, S., Ahn, N.G., and Klinman, J.P., Hydrogen Deuterium Exchange Defines Catalytically Linked Regions of Protein Flexibility in the Catechol O-Methyltransferase Reaction. *Proc. Natl. Acad. Sci.* **117**, 10797 (2020).

21. S. Gao *et al.*, Hydrogen–Deuterium Exchange within Adenosine Deaminase, a TIM Barrel Hydrolase, Identifies Networks for Thermal Activation of Catalysis. *J. Am. Chem. Soc.* **142**, 19936-19949 (2020).

22. E. J. Thompson, A. Paul, A. T. Iavarone, J. P. Klinman, Identification of Thermal Conduits That Link the Protein–Water Interface to the Active Site Loop and Catalytic Base in Enolase. *J. Am. Chem. Soc.* **143**, 785-797 (2021).

23. Z. Wang *et al.*, Ultrafast Flash Thermal Conductance of Molecular Chains. *Science* **317**, 787 (2007).

24. L. Martínez, A. C. M. Figueira, P. Webb, I. Polikarpov, M. S. Skaf, Mapping the Intramolecular Vibrational Energy Flow in Proteins Reveals Functionally Important Residues. *J. Phys. Chem. Lett.* **2**, 2073-2078 (2011).

25. J. Helbing *et al.*, Temperature Dependence of the Heat Diffusivity of Proteins. *J. Phys. Chem. A* **116**, 2620-2628 (2012).

26. I. V. Rubtsov, A. L. Burin, Ballistic and diffusive vibrational energy transport in molecules. *J. Chem. Phys.* **150**, 020901 (2019).

27. M. Mizuno, Y. Mizutani, Role of atomic contacts in vibrational energy transfer in myoglobin. *Biophys. Rev.* **12**, 511-518 (2020).

28. H. Poudel, K. M. Reid, T. Yamato, D. M. Leitner, Energy Transfer across Nonpolar and Polar Contacts in Proteins: Role of Contact Fluctuations. *J. Phys. Chem. B* **124**, 9852-9861 (2020).

29. L. Valiño Borau, A. Gulzar, G. Stock, Master equation model to predict energy transport pathways in proteins. *J. Chem. Phys.* **152**, 045103 (2020).

30. R. Kozlowski, J. Zhao, R. B. Dyer, Acceleration of catalysis in dihydrofolate reductase by transient, site-specific photothermal excitation. *Proc. Natl. Acad. Sci.* **118**, e2014592118 (2021).

31. A. Ansari *et al.*, Protein states and proteinquakes. *Proc. Natl. Acad. Sci.* **82**, 5000 (1985).

32. O. Miyashita, J. N. Onuchic, P. G. Wolynes, Nonlinear elasticity, proteinquakes, and the energy landscapes of functional transitions in proteins. *Proc. Natl. Acad. Sci.* **100**, 12570 (2003).

33. H. Frauenfelder *et al.*, A unified model of protein dynamics. *Proc. Natl. Acad. Sci.* **106**, 5129 (2009).

34. D. Arnlund *et al.*, Visualizing a protein quake with time-resolved X-ray scattering at a free-electron laser. *Nat. Methods* **11**, 923-926 (2014).

35. J. G. Kim, T. W. Kim, J. Kim, H. Ihee, Protein Structural Dynamics Revealed by Time-Resolved X-ray Solution Scattering. *Acc. Chem. Res.* **48**, 2200-2208 (2015).

36. Y. Lee *et al.*, Ultrafast coherent motion and helix rearrangement of homodimeric hemoglobin visualized with femtosecond X-ray solution scattering. *Nat. Commun.* **12**, 3677 (2021).

37. R. C. Tolman, Statistical Merchanics Applied To Chemical Kinetics. *J. Am. Chem. Soc.* **42**, 2506-2528 (1920).
38. P. K. Agarwal, Role of Protein Dynamics in Reaction Rate Enhancement by Enzymes. *J. Am. Chem. Soc.* **127**, 15248-15256 (2005).
39. M. J. Knapp, J. P. Klinman, Environmentally coupled hydrogen tunneling. *Eur. J. Biochem.* **269**, 3113-3121 (2002).
40. M. J. Knapp, K. Rickert, J. P. Klinman, Temperature-Dependent Isotope Effects in Soybean Lipoxygenase-1: Correlating Hydrogen Tunneling with Protein Dynamics. *J. Am. Chem. Soc.* **124**, 3865-3874 (2002).
41. M. J. Sutcliffe, N. S. Scrutton, A new conceptual framework for enzyme catalysis. *Eur. J. Biochem.* **269**, 3096-3102 (2002).
42. E. Hatcher, A. V. Soudackov, S. Hammes-Schiffer, Proton-Coupled Electron Transfer in Soybean Lipoxygenase. *J. Am. Chem. Soc.* **126**, 5763-5775 (2004).
43. P. M. Kiefer, J. T. Hynes, Kinetic Isotope Effects for Nonadiabatic Proton Transfer Reactions in a Polar Environment. 1. Interpretation of Tunneling Kinetic Isotopic Effects. *J. Phys. Chem. A* **108**, 11793-11808 (2004).
44. S. Hammes-Schiffer, Hydrogen Tunneling and Protein Motion in Enzyme Reactions. *Acc. Chem. Res.* **39**, 93-100 (2006).
45. J. P. Klinman, A. R. Offenbacher, S. Hu, Origins of Enzyme Catalysis: Experimental Findings for C–H Activation, New Models, and Their Relevance to Prevailing Theoretical Constructs. *J. Am. Chem. Soc.* **139**, 18409-18427 (2017).
46. M. H. Glickman, J. S. Wiseman, J. P. Klinman, Extremely Large Isotope Effects in the Soybean Lipoxygenase-Linoleic Acid Reaction. *J. Am. Chem. Soc.* **116**, 793-794 (1994).
47. H. Porta, M. Rocha-Sosa, Plant Lipoxygenases. Physiological and Molecular Features. *Plant Physiol.* **130**, 15 (2002).
48. K. W. Rickert, J. P. Klinman, Nature of Hydrogen Transfer in Soybean Lipoxygenase 1: Separation of Primary and Secondary Isotope Effects. *Biochemistry* **38**, 12218-12228 (1999).
49. S. Hu *et al.*, Extremely Elevated Room-Temperature Kinetic Isotope Effects Quantify the Critical Role of Barrier Width in Enzymatic C–H Activation. *J. Am. Chem. Soc.* **136**, 8157-8160 (2014).
50. B. Salna, A. Benabbas, D. Russo, P. M. Champion, Tunneling Kinetics and Nonadiabatic Proton-Coupled Electron Transfer in Proteins: The Effect of Electric Fields and Anharmonic Donor–Acceptor Interactions. *J. Phys. Chem. B* **121**, 6869-6881 (2017).
51. P. Li, A. V. Soudackov, S. Hammes-Schiffer, Fundamental Insights into Proton-Coupled Electron Transfer in Soybean Lipoxygenase from Quantum Mechanical/Molecular Mechanical Free Energy Simulations. *J. Am. Chem. Soc.* **140**, 3068-3076 (2018).
52. A. R. Offenbacher, A. Sharma, P. E. Doan, J. P. Klinman, B. M. Hoffman, The Soybean Lipoxygenase–Substrate Complex: Correlation between the Properties of Tunneling-Ready States and ENDOR-Detected Structures of Ground States. *Biochemistry* **59**, 901-910 (2020).
53. J. P. Klinman, A. R. Offenbacher, Understanding Biological Hydrogen Transfer Through the Lens of Temperature Dependent Kinetic Isotope Effects. *Acc. Chem. Res.* **51**, 1966-1974 (2018).
54. P. Singh *et al.*, Evolution of the Chemical Step in Enzyme Catalysis. *ACS Catal.* **11**, 6726-6732 (2021).
55. S. Hu, A. R. Offenbacher, E. D. Lu, J. P. Klinman, Comparative kinetic isotope effects on first- and second-order rate constants of soybean lipoxygenase variants uncover a substrate-binding network. *J. Biol. Chem.* **294**, 18069-18076 (2019).
56. S. Hu *et al.*, Biophysical Characterization of a Disabled Double Mutant of Soybean Lipoxygenase: The "Undoing" of Precise Substrate Positioning Relative to Metal Cofactor and an Identified Dynamical Network. *J. Am. Chem. Soc.* **141**, 1555-1567 (2019).
57. D. A. Keedy, J. S. Fraser, H. van den Bedem, Exposing Hidden Alternative Backbone Conformations in X-ray Crystallography Using qFit. *PLoS Computat. Biol.* **11**, e1004507 (2015).
58. S. V. Kathuria, Y. H. Chan, R. P. Nobrega, A. Özen, C. R. Matthews, Clusters of isoleucine, leucine, and valine side chains define cores of stability in high-energy states of globular proteins: Sequence determinants of structure and stability. *Protein Sci.* **25**, 662-675 (2016).
59. S. Basak *et al.*, Networks of electrostatic and hydrophobic interactions modulate the complex folding free energy surface of a designed βα protein. *Proc. Natl. Acad. Sci.* **116**, 6806 (2019).
60. N. Ferruz, S. Schmidt, B. Höcker, ProteinTools: a toolkit to analyze protein structures. *Nucleic Acids Res.* **49**, W559-W566 (2021).

61. J. P. T. Zaragoza *et al.*, Detecting and Characterizing the Kinetic Activation of Thermal Networks in Proteins: Thermal Transfer from a Distal, Solvent-Exposed Loop to the Active Site in Soybean Lipoxygenase. *J. Phys. Chem. B* **123**, 8662-8674 (2019).
62. R. B. Macgregor, G. Weber, Estimation of the polarity of the protein interior by optical spectroscopy. *Nature* **319**, 70 (1986).
63. P. Pospíšil *et al.*, Fluorescence Quenching of (Dimethylamino)naphthalene Dyes Badan and Prodan by Tryptophan in Cytochromes P450 and Micelles. *J. Phys. Chem. B* **118**, 10085-10091 (2014).
64. M. P. Meyer, D. R. Tomchick, J. P. Klinman, Enzyme structure and dynamics affect hydrogen tunneling: The impact of a remote side chain (I553) in soybean lipoxygenase-1. *Proc. Natl. Acad. Sci.* **105**, 1146 (2008).
65. S. C. Sharma, J. P. Klinman, Kinetic Detection of Orthogonal Protein and Chemical Coordinates in Enzyme Catalysis: Double Mutants of Soybean Lipoxygenase. *Biochemistry* **54**, 5447-5456 (2015).
66. M. G. Badea, L. Brand, Time-resolved fluorescence measurements in *Methods in Enzymology*. (Academic Press, 1979), vol. 61, pp. 378-425.
67. J. R. Lakowicz, H. Cherek, G. Laczko, E. Gratton, Time-resolved fluorescence emission spectra of labeled phospholipid vesicles, as observed using multi-frequency phase-modulation fluorometry. *Biochim. Biophys. Acta Biomembr.* **777**, 183-193 (1984).
68. M. L. Horng, J. A. Gardecki, A. Papazyan, M. Maroncelli, Subpicosecond Measurements of Polar Solvation Dynamics: Coumarin 153 Revisited. *J. Chem. Phys.* **99**, 17311-17337 (1995).
69. N. Nandi, B. Bagchi, Dielectric Relaxation of Biological Water. *J. Phys. Chem. B* **101**, 10954-10961 (1997).
70. A. Jesenská *et al.*, Nanosecond Time-Dependent Stokes Shift at the Tunnel Mouth of Haloalkane Dehalogenases. *J. Am. Chem. Soc.* **131**, 494-501 (2009).
71. S. Guha *et al.*, Slow Solvation Dynamics at the Active Site of an Enzyme: Implications for Catalysis. *Biochemistry* **44**, 8940-8947 (2005).
72. V. Makarov, B. M. Pettitt, M. Feig, Solvation and Hydration of Proteins and Nucleic Acids: A Theoretical View of Simulation and Experiment. *Acc. Chem. Res.* **35**, 376-384 (2002).
73. S. Pal, S. Bandyopadhyay, Importance of Protein Conformational Motions and Electrostatic Anchoring Sites on the Dynamics and Hydrogen Bond Properties of Hydration Water. *Langmuir* **29**, 1162-1173 (2013).
74. M. H. Glickman, J. P. Klinman, Nature of Rate-Limiting Steps in the Soybean Lipoxygenase-1 Reaction. *Biochemistry* **34**, 14077-14092 (1995).
75. C. W. Meadows, J. E. Tsang, J. P. Klinman, Picosecond-Resolved Fluorescence Studies of Substrate and Cofactor-Binding Domain Mutants in a Thermophilic Alcohol Dehydrogenase Uncover an Extended Network of Communication. *J. Am. Chem. Soc.* **136**, 14821-14833 (2014).
76. C. W. Meadows, R. Ou, J. P. Klinman, Picosecond-Resolved Fluorescent Probes at Functionally Distinct Tryptophans within a Thermophilic Alcohol Dehydrogenase: Relationship of Temperature-Dependent Changes in Fluorescence to Catalysis. *J. Phys. Chem. B* **118**, 6049-6061 (2014).
77. Y. Qin, L. Wang, D. Zhong, Dynamics and mechanism of ultrafast water–protein interactions. *Proc. Natl. Acad. Sci.* **113**, 8424 (2016).
78. Y. Qin, L. Zhang, L. Wang, D. Zhong, Observation of the Global Dynamic Collectivity of a Hydration Shell around Apomyoglobin. *J. Phys. Chem. Lett.* **8**, 1124-1131 (2017).
79. M. Jia *et al.*, Determination of Protein Surface Hydration by Systematic Charge Mutations. *J. Phys. Chem. Lett.* **6**, 5100-5105 (2015).
80. M. Horitani *et al.*, 13C ENDOR Spectroscopy of Lipoxygenase–Substrate Complexes Reveals the Structural Basis for C–H Activation by Tunneling. *J. Am. Chem. Soc.* **139**, 1984-1997 (2017).
81. A. E. Eriksson *et al.*, Response of a Protein Structure to Cavity-Creating Mutations and Its Relation to the Hydrophobic Effect. *Science* **255**, 178-183 (1992).
82. M. Xue *et al.*, How internal cavities destabilize a protein. *Proc. Natl. Acad. Sci.* **116**, 21031 (2019).
83. B. C. Buer, J. L. Meagher, J. A. Stuckey, E. N. G. Marsh, Structural basis for the enhanced stability of highly fluorinated proteins. *Proc. Natl. Acad. Sci.* **109**, 4810 (2012).
84. M. D. Collins, G. Hummer, M. L. Quillin, B. W. Matthews, S. M. Gruner, Cooperative water filling of a nonpolar protein cavity observed by high-pressure crystallography and simulation. *Proc. Natl. Acad. Sci.* **102**, 16668 (2005).
85. A. E. Eriksson, W. A. Baase, B. W. Matthews, Similar Hydrophobic Replacements of Leu99 and Phe153 within the Core of T4 Lysozyme Have Different Structural and Thermodynamic Consequences. *J. Mol. Biol.* **229**, 747-769 (1993).

86. K. Takano *et al.*, Contribution of Hydrophobic Residues to the Stability of Human Lysozyme: Calorimetric Studies and X-ray Structural Analysis of the Five Isoleucine to Valine Mutants. *J. Mol. Biol.* **254**, 62-76 (1995).

87. P.-A. Brault *et al.*, Protein Micelles from Lipoxygenase 3. *Biomacromolecules* **3**, 649-654 (2002).

88. S. C. L. Kamerlin, A. Warshel, At the dawn of the 21st century: Is dynamics the missing link for understanding enzyme catalysis? *Proteins* **78**, 1339-1375 (2010).

89. V. C. Nashine, S. Hammes-Schiffer, S. J. Benkovic, Coupled motions in enzyme catalysis. *Curr. Opin. Chem. Biol.* **14**, 644-651 (2010).

90. K. A. Henzler-Wildman *et al.*, A hierarchy of timescales in protein dynamics is linked to enzyme catalysis. *Nature* **450**, 913 (2007).

91. E. Z. Eisenmesser *et al.*, Intrinsic dynamics of an enzyme underlies catalysis. *Nature* **438**, 117-121 (2005).

92. R. H. Austin, K. W. Beeson, L. Eisenstein, H. Frauenfelder, I. C. Gunsalus, Dynamics of ligand binding to myoglobin. *Biochemistry* **14**, 5355-5373 (1975).

93. F. H. Arnold, Directed Evolution: Bringing New Chemistry to Life. *Angew. Chem. Int. Ed.* **57**, 4143-4148 (2018).

94. R. Otten *et al.*, Rescue of conformational dynamics in enzyme catalysis by directed evolution. *Nature Commun.* **9**, 1314 (2018).

95. M. Wilding, N. Hong, M. Spence, A. M. Buckle, C. J. Jackson, Protein engineering: the potential of remote mutations. *Biochem. Soc. Trans.* **47**, 701-711 (2019).

96. J. W. Schafer, S. D. Schwartz, Directed Evolution's Influence on Rapid Density Fluctuations Illustrates How Protein Dynamics Can Become Coupled to Chemistry. *ACS Catal.* **10**, 8476-8484 (2020).

97. C. J. Markin *et al.*, Revealing enzyme functional architecture via high-throughput microfluidic enzyme kinetics. *Science* **373**, 411 (2021).

98. H. G. Saavedra, J. O. Wrabl, J. A. Anderson, J. Li, V. J. Hilser, Dynamic allostery can drive cold adaptation in enzymes. *Nature* **558**, 324-328 (2018).

99. G. Li, D. Magana, R. B. Dyer, Anisotropic energy flow and allosteric ligand binding in albumin. *Nat. Commun.* **5**, 3100 (2014).

100. T. P. Causgrove, R. B. Dyer, Picosecond Structural Dynamics of Myoglobin following Photolysis of Carbon Monoxide. *J. Phys. Chem.* **100**, 3273-3277 (1996).

101. D. M. Leitner, T. Yamato, "Mapping Energy Transport Networks in Proteins" in Reviews in Computational Chemistry, A. L. Parrill, K. B. Lipkowitz, Eds. (2018), vol. 31, pp. 63-113.

102. D. M. Leitner, Energy Flow in Proteins. *Annu. Rev. Phys. Chem.* **59**, 233-259 (2008).

103. X. Yu, D. M. Leitner, Heat flow in proteins: Computation of thermal transport coefficients. *J. Chem. Phys.* **122**, 054902 (2005).

104. X. Yu, D. M. Leitner, Vibrational Energy Transfer and Heat Conduction in a Protein. *J. Phys. Chem. B* **107**, 1698-1707 (2003).

105. I. Nardecchia *et al.*, Out-of-Equilibrium Collective Oscillation as Phonon Condensation in a Model Protein. *Phys. Rev. X* **8**, 031061 (2018).

106. S. C. Althorpe *et al.*, Non-adiabatic reactions: general discussion. *Faraday Discuss.* **195**, 311-344 (2016).

107. M. B. Vaughn, J. Zhang, T. G. Spiro, R. B. Dyer, J. P. Klinman, Activity-Related Microsecond Dynamics Revealed by Temperature-Jump Förster Resonance Energy Transfer Measurements on Thermophilic Alcohol Dehydrogenase. *J. Am. Chem. Soc.* **140**, 900-903 (2018).

108. J. D. Dunitz, Win some, lose some: enthalpy-entropy compensation in weak intermolecular interactions. *Chem. Biol.* **2**, 709-712 (1995).

109. B. Breiten *et al.*, Water Networks Contribute to Enthalpy/Entropy Compensation in Protein–Ligand Binding. *J. Am. Chem. Soc.* **135**, 15579-15584 (2013).

110. P. W. Fenimore, H. Frauenfelder, B. H. McMahon, F. G. Parak, Slaving: Solvent fluctuations dominate protein dynamics and functions. *Proc. Natl. Acad. Sci.* **99**, 16047 (2002).

111. Y. Kim *et al.*, Quantum Biology: An Update and Perspective. *Quantum Rep.* **3** (2021).

112. J. Cao *et al.*, Quantum biology revisited. *Sci. Adv.* **6**, eaaz4888.

113. L. Pauling, Nature of Forces between Large Molecules of Biological Interest. *Nature* **161**, 707-709 (1948).

114. L. Pauling, Molecular Architecture and Biological Reactions. *Chemical & Engineering News Archive* **24**, 1375-1377 (1946).

Acknowledgments:

We thank the members of the Klinman lab for the insightful discussions about the manuscript, specifically Dr. E. Thompson for assistance with programming and data analysis. **Funding:** This research is supported by funding from the National Institutes of Health grants to J.P.K. (GM118117), A.R.O. (GM113432 (F32)), and A.T.I. (1S10OD020062-01). J.P.T.Z. thanks the University of California President's Postdoctoral Program for a fellowship. Data collection at BL 8.3.1 at the Advanced Light Source is supported by the Director, Office of Science, Office of Basic Energy Sciences, of the U.S. Department of Energy under contract no. DE-AC02-05CH11231, UC Office of the President, Multicampus Research Programs, and Initiatives grant MR 15 32859, the Program Breakthrough Biomedical Research, which is partially funded by the Sandler Foundation, and the NIH P30GM124169. **Author contributions:** J.P.T.Z. designed the time-resolved fluorescence experiments and collected and analyzed the corresponding kinetics and mass spectrometry data. A.R.O. collected and analyzed TD-HDXMS, DSC, and enzyme kinetics data. A.R.O., S.H., and C.L.G. collected and analyzed the X-ray crystallography data. Z.M.F., N.M., F.F. and Z.D. provided assistance with data collection and analysis. A.T.I. collected and analyzed mass spectrometry data. J.P.K. conceived and supervised the project and acquired funding. J.P.T.Z., A.R.O., and J.P.K. wrote and revised the manuscript, with input from all authors. **Competing interests:** The authors declare no competing interest. **Data and materials availability:** Summary tables of all kinetic and thermodynamic parameters, as well as fluorescence and crystallography data measured for each mutant are included in the supplementary materials. The atomic models have been deposited at the Protein Data Bank with PDB IDs 7SOI (I552A SLO) and 7SOJ (V750A SLO).

Supplementary Materials:
Materials and Methods
Figs. S1 to S12
Tables S1 to S12
Data S1. HDX-MS deuterium uptake plots for all peptides of I552A and WT SLO
Data S2. Summary of HDX-MS data for I552A SLO at 10, 20, 25, 30, and 40 ˚C