Late maturation of executive control promotes conceptual development

Rebecca L. JACKSON[1], Matthew A. LAMBON RALPH[1] & Timothy T. ROGERS[2]

[1]MRC Cognition & Brain Sciences Unit

University of Cambridge

15 Chaucer Road

Cambridge

CB2 7EF

[2]524 WJ Brogden Hall

Department of Psychology

University of Wisconsin-Madison

Madison, Wisconsin

53706

Author Note

All data are available in the Supplementary Materials or can be generated with the code provided. All code is available at github.com/JacksonBecky/developmental. The study design and analyses were not preregistered.

This work was supported by a British Academy Postdoctoral Fellowship awarded to R.L.J. (no. pf170068), a programme grant to M.A.L.R. from the Medical Research Council (grant no. MR/R023883/1), an Advanced Grant from the European Research Council to M.A.L.R. (GAP: 670428) and Medical Research Council intramural funding (no. MC_UU_00005/18).

* <u>Correspondence to</u>:
Dr. Rebecca Jackson
MRC Cognition & Brain Sciences Unit
University of Cambridge
15 Chaucer Road
Cambridge
CB2 7EF
Rebecca.Jackson@mrc-cbu.cam.ac.uk
ttrogers@wisc.edu
Matt.Lambon-Ralph@mrc-cbu.cam.ac.uk

Abstract

Control processes are critical for the generation of task-appropriate behaviour across cognitive domains, yet children have a long developmental period with reduced executive control. Traditionally, this is viewed as a negative but necessary consequence of the time taken to learn control processes and develop the prefrontal cortex. Here, we exploit a recent computational model of controlled semantic cognition to formally test an alternative (yet perhaps complementary) view that a developmental period without control promotes conceptual knowledge acquisition. Our simulations show that maturational delay and anatomical connectivity conspire to promote conceptual learning. Learning conceptual structure necessitates a connection from control to more peripheral regions rather than the deep multimodal hub. Delayed control speeds conceptual learning without compromising conceptual representations, particularly when control connects to intermediate layers. These results are counterintuitive as delayed control onset can produce overall faster mastery of controlled behaviours. To assess whether delayed semantic control is also observed over development, we conducted a meta-analysis of the classic triadic matching task where participants decide which of two options best matches a third. Matching can be based on taxonomic or thematic relations, and when these conflict, participants must use task context to determine which relation to choose—that is, they must exert semantic control. Children aged 2 have the representations to identify both relations when distractors are unrelated, but when relations conflict, they cannot use context to guide their choices. Context-sensitivity develops later than conceptual structure and shows large increases around 6, a critical time for executive function across domains.

Keywords: executive control; semantic cognition; neural networks; development; learning.

Human thought is remarkably flexible. More than other mammals, humans can control the mental representations and behaviours that arise from moment to moment, avoiding prepotent responses and reconfiguring well-learned patterns to better suit immediate and long-term goals. These abilities depend upon the gradual maturation of the prefrontal cortex (PFC; Duncan, 2010; Fuster, 2015; Miller & Cohen, 2001; Smaers, Gómez-Robles, Parks, & Sherwood, 2017; Wood & Grafman, 2003), a brain structure that develops more slowly than other cortical regions within the human brain and more slowly in humans than the homologous structures of other mammals (Chugani & Phelps, 1986; Elston, Oga, & Fujita, 2009; Huttenlocher & Dabholkar, 1997). Slow developmental change is observed in structural (synaptogenesis, synaptic elimination, dendritic arborisation and myelination), functional (regional metabolism) and behavioural (speed of processing, working memory, inhibition and task switching) measures, with the PFC-mediated control system not fully maturing until past the age of twenty (Casey, Tottenham, Liston, & Durston, 2005; Davidson, Amso, Anderson, & Diamond, 2006; A. Diamond, 2002; A. Diamond & Doar, 1989; Mrzljak, Uylings, van Eden, & Judas, 1990; Veroude, Jolles, Croiset, & Krabbendam, 2013; Zelazo, Carlson, & Kesek, 2008). Thus, children undergo a long developmental period with comparatively little cognitive control.

The slow onset of control is often viewed as an unfortunate but necessary consequence of the PFC's complexity (e.g., Church, Bunge, Petersen, & Schlaggar, 2017; A. Diamond, 2002). Given constraints on the size of the head at birth, the requirement of growing out and pruning back connections from PFC to the rest of the brain, and other complex biological constraints, it may be that cognitive control simply requires a very long time to mature (Bunge, Dudukovic, Thomason, Chandan, & Gabrieli, 2002; Church et al., 2017; Crone, Donohue, Honomichl, Wendelken, & Bunge, 2006; Luna, Garver, Urban, Lazar, & Sweeney, 2004; Wilk & Morton, 2012). We consider an alternative though perhaps complementary view: that the slow maturation of control confers a functional advantage in the development of conceptual knowledge. Our proposal builds on prior arguments that a period without control may have positive functional

consequences for learning, which offset the costs incurred by the late development of flexible and adaptive behaviour (Chrysikou, Novick, Trueswell, & Thompson-Schill, 2011; Newport, 1990; Thompson-Schill, Ramscar, & Chrysikou, 2009). In this paper, we explain why conceptual knowledge acquisition in particular might benefit from a period with limited control, assess the hypothesis formally via simulations with a computational model of controlled semantic cognition, and then consider how the modelling results shed light on classic findings in studies of conceptual development.

Our proposal is motivated by the seemingly antagonistic relationship lying at the heart of controlled semantic cognition (Jackson, Rogers, & Lambon Ralph, 2021; Lambon Ralph, Jefferies, Patterson, & Rogers, 2017). On one hand, the human semantic system must accumulate information about the world across many disparate learning episodes, each providing only a partial glimpse into the environment's structure. Consider the widespread view that human concepts arise from learning bundles of correlated attributes in the environment (Keil, 1979; Murphy & Medin, 1985; Rogers & McClelland, 2004; Rosch, Mervis, Gray, Johnson, & Boyes-Braem, 1976). For instance, the concept *bird* might arise because there are sets of attributes that all tend to occur together in various instances involving birds: feathers, wings, beaks, hollow bones, the ability to fly, the ability to lay eggs, the name "bird," and so on. While these properties may all be true of the same items, such covariation is not transparently observable in everyday experience: a flying bird is not simultaneously laying an egg; the bird skeleton observed in science class does not possess feathers and does not fly; the word "bird" appearing in a novel typically does not appear together with an image of the bird, and so on. To learn that these various properties all inhere together within exemplars of the concept *bird*, the semantic system must be able to track sameness of kind across these different scenarios—that is, it must "know" that the bird observed flying overhead is the same kind of thing as the skeleton studied in science class, which in turn is the same kind of thing as the item described in a novel. Thus, the semantic system must *abstract* across different situational contexts, each providing only

a partial glimpse into an item's full complement of attributes, to form conceptual representations that fully express attribute covariances as they exist in the world and allow generalisation to new exemplars (Jackson et al., 2021; Lambon Ralph et al., 2017; Rogers & McClelland, 2004). On the other hand, to support everyday behaviour the semantic system must also produce internal representations and behaviours that are *context-sensitive,* so that information that comes to mind in the moment is suited to the goals of the immediate behavioural task. The features required to engage with a piano in different contexts, such as moving it and playing it, differ substantially (Saffran, 2000). In order to provide a context-appropriate output, only relevant features should be selected and allowed to bear on the output decision whilst other, often more dominant, features must be suppressed (Badre, Poldrack, Pare-Blagoev, Insler, & Wagner, 2005; Jefferies, 2013; Thompson-Schill, D'Esposito, Aguirre, & Farah, 1997; Wagner, Pare-Blagoev, Clark, & Poldrack, 2001). Thus, the semantic system must concurrently abstract across contexts to acquire representations that express conceptual structure (*representation*), yet also shape representations and behaviours to suit the immediate context (*control*).

In prior work, we showed that these opposing functional demands are best accomplished within an architecture like that shown in Figure 1, where representations across different modalities (e.g., vision, audition, language, action) mutually interact via a single deep, multimodal "hub" and where control operates by directly potentiating task-relevant modality-specific and immediately adjacent areas (corresponding to the presence of the black or green but not orange connections in Figure 1; Jackson et al., 2021). In this architecture, the "hub" units are always involved in propagating information between different "spokes" of the system, and so can learn the high-order covariance structure of the environment across different modalities and contexts—even though, for each training episode, the control system allows only a subset of an item's properties to become active. That is, when control operates on regions nearer the sensory-motor periphery, the distal representations that emerge in the multimodal hub are relatively invariant to context, and thus come to express the conceptual structure inherent across different learning episodes

regardless of the current contextual demands. Thus, the antagonistic processes of control and representation can co-exist in a semantic system with an architecture that promotes their relative functional separation. This organisational principle is further mimicked within the brain: PFC and lateral posterior temporal cortex support semantic control processes, whereas representation relies upon distributed spoke regions and the anterior temporal lobe multimodal hub (Jackson, 2021; Jefferies, 2013; Lambon Ralph et al., 2017). It also underlies the distinct behavioural profiles observed in *semantic dementia*—a gradual erosion of multimodal conceptual representations accompanying a progressive degradation of the anterior temporal lobe (Acosta-Cabronero et al., 2011; Patterson, Nestor, & Rogers, 2007; Warrington, 1975)—and *semantic aphasia*, a loss of the controlled, context-appropriate use of concepts following cerebrovascular accident to frontal or posterior temporal cortices (Jefferies, 2013; Jefferies & Lambon Ralph, 2006).

This paper considers whether delayed maturation of control might likewise promote better acquisition of conceptual structure, and whether such effects obviate or complement prior conclusions about the optimal anatomical structure of the network. Control signals are thought to promote the activation of task-relevant properties while suppressing task-irrelevant properties true of a given item (Cohen, Dunbar, & McClelland, 1990; Thompson-Schill et al., 1997; Waksom, Kumaran, Gordon, Rissman, & Wagner, 2014). For instance, when naming an item such as a toy train, control ensures that only a context-appropriate name is produced (e.g., "train" and not "vehicle," "toy," or "choo-choo"), and moreover that other associated information, such as action-plans (e.g., playing with the toy) or other irrelevant facts about the object ("It goes on tracks") do not drive behaviour. When only task-relevant properties become activated, it may be difficult to learn conceptual structure since such structure is latent in the distribution of properties apparent across many different contexts. Moreover, conceptual acquisition may be generally slowed, since the learner must inhibit all but a small subset of an item's properties in any given episode. Thus, a developmental period with limited control may

promote conceptual learning because the absence of control allows all properties associated with a stimulus, including those not relevant to the current task and not directly observed, to become activated simultaneously. Indeed, if delayed onset of control promotes conceptual abstraction, it may be less important for the control system to be anatomically insulated from the semantic network "hub," obviating conclusions from prior work. Alternatively, it may be that the architectural constraints identified in this prior work are sufficient to ensure good learning of conceptual structure regardless of the timing with which the control system develops, or that the developmental timing and architectural constraints conspire to best promote conceptual abstraction. To adjudicate these possibilities, we conducted simulations with the reverse-engineered model of semantic representation (Jackson et al., 2021), manipulating the rate at which the control system "matures" and the connectivity between the control and semantic representation networks, then measuring the effect of these factors on conceptual abstraction and training time. The simulations provide clear conclusions about both the development of control and the connectivity of control and representation regions.

They also make predictions about the time-course of semantic control over development. While a few developmental studies have touched on questions about semantic control (Blaye, Bernard-Peyron, Paour, & Bonthoux, 2006; Blaye & Jacques, 2009; Henderson, Clarke, & Snowling, 2011; Snyder & Munakata, 2010, 2013), most focus on the acquisition of semantic representations (e.g., Gershkoff-Stowe & Rakison, 2005). The second part of the paper thus tests the model predictions through a new literature review and meta-analysis of a classic task designed to test development of semantic knowledge, the triadic match-to-sample task in which children must decide which of two items is most similar to a third. While this task is not typically viewed as assessing semantic control, many studies have focused on manipulating both the nature of the information required for a successful match (taxonomic or thematic; Bauer & Mandler, 1989; Berger & Aguerra, 2010; Blanchet, Dunham, & Dunham, 2001; Greenfield & Scott, 1986; Smiley & Brown, 1979; Waxman & Namy, 1997) and the influence of instructional cues on behaviour

(e.g., Estes, Golonka, & Jones, 2011; Greenfield & Scott, 1986; Waxman & Namy, 1997). As an instructional cue is a form of task context, sensitivity to the cue—that is, the degree to which children can make different choices for the same stimulus depending upon the task instruction—indexes the ability to use control to shape semantic decisions. Our meta-analysis tracks changes in such sensitivity over development. Revisiting this extensive literature through the contemporary lens of controlled semantic cognition provides a powerful way to estimate the interplay of semantic control and representation processes across development.

Part 1: Simulations

Methods

We conducted simulations with the recent model of controlled semantic cognition from Jackson et al., (2021; see Figure 1), a deep variant of the hub-and-spokes framework in which the propagation of activation through the semantic representation network is constrained by inputs from a prefrontal "control" system. Items in the environment are represented as distributed patterns across three modality-specific input/output channels (e.g., vision, words, and action). From input in any given channel, the model learns to activate patterns in other channels appropriate to the current task context. For instance, this may simulate a picture naming task, if the model receives direct input via the "visual" channel for a given concept, and must activate the corresponding pattern for that concept in the "words" layer, without activating the associated pattern in the "action" layer. Alternatively, it could simulate production of an associated gesture given an object's name, if the model receives input in the "word" layer and must generate the associated pattern across "action" units, without activating "vision" units, and so on. Tasks are represented with units that locally indicate which modalities are involved—for instance, in a task taking input in M1 and producing output in M2, both the M1 and M2 units in the control layer are active. These then send connections directly to units across the deep hub-

and-spokes network. To understand how control influences the development of semantic representations in this setup, we manipulated (1) the timing and speed with which controlled semantic behaviour arises and (2) the connectivity between control units and the rest of the semantic network.

Figure 1. A. The model architecture. Sensory input is received and output generated, in three modality-specific sensory regions (black ovals) constituting the Spokes Layer. This layer is connected to an Intermediate Layer (green ovals) which in turn is connected to the Hub Layer (orange oval). A small number of sparse connections also directly connect the Spokes Layer to the Hub Layer. A context signal is received as input to three units forming a Control Layer (red oval). This context signal specifies the modality in which the sensory input is received and the modality in which the output should be generated. This Control Layer may be connected to all layers (full connectivity), or to the Spokes Layer (black arrow), Intermediate Layer (green arrow) or Hub Layer (orange arrow) only. B. The model environment. The 16 concepts are shown. Each concept is a row, and each feature is a column. If the feature is present the box is black, if absent the box is white. When control requirements are present, each concept can be presented in one of three modalities, and output required in one of three modalities, resulting in 144 different trials. Each modality has a surface similarity structure (orange, yellow or green boxes), which is orthogonal to a multimodal structure (red boxes) learnt through conceptual abstraction across modalities.

Construction of model environment. The model environment is the same as in Jackson et al., (2021), designed to assess how well each model layer can encode cross-modal similarity structure through conceptual abstraction across sensory modalities and contexts. It includes 16 concepts, each possessing a unique activation pattern across 12 feature units in each of the 3 modality-

specific information channels (see Figure 1.B.). The patterns were designed so that (1) the pairwise similarity structure encoded in a given modality was uncorrelated with the similarities encoded in each other modality, but with (2) a small number of features expressing common structure *across* the three modalities. This scheme captures the common insight that perceptual similarities in a given modality often differ both from the "surface" similarities in other modalities and from "deep" conceptual similarity structure—for instance, a pear and a lightbulb have similar shapes but are conceptually distinct and also have very different verbal descriptors and associated action plans. The inclusion of features that express the same structure across modalities ensures that the full multimodal structure differs from that apparent in any single channel, and captures the common intuition that conceptual structure emerges from patterns of covariation across stimulus modalities and task contexts. If the learned patterns of activation arising across a given layer express inter-item similarities highly correlated with the full crossmodal structure and independent of the current task context, that layer can be understood as encoding "deep" conceptual structure.

Architectural details. The architecture of the semantic network model (Figure 1) was shown in prior work to best promote both learning of the latent context-independent structure present across modalities and context-sensitive responding. We describe the key parameters briefly here; for the full rationale see Jackson et al. (2021). Twelve feature units in each modality (the "Spokes" layers) employed a sigmoid activation function and possessed a fixed bias of -3, thus adopting low activations in the absence of input. Each feature unit also received a direct positive connection from a corresponding input unit, with a fixed value of +6. Direct perception of an item's features was simulated by turning on the associated input units which, in combination with the negative biases, provided a total net input of +3 to the corresponding feature units. These feature units were reciprocally connected to two layers of hidden units, as shown in Figure 1. Each spoke was fully and reciprocally connected to a distinct set of 14 hidden units in the Intermediate Layer, and each such layer was reciprocally connected with a common Hub Layer

(including 18 units) using a sparsity of 0.6349. This sparsity level was used for consistency with the prior paper, where it was necessary to match the total number of connections across different network architectures. Thus, the Hub Layer is a multimodal hub mediating between all sensory modalities, whereas regions of the Intermediate Layer support relatively modality-specific processing. Additionally, a small number of 'shortcut connections' project directly from the spokes to the hub, bypassing the Intermediate Layer, analogous to sparse long-range connectivity observed in the temporal lobe. Each region also has self-connections at full sparsity in the hub and spoke layers and half sparsity within each Intermediate Layer region.

When provided with an item's partial input in one modality, the "mature" (i.e., fully trained) model must activate, across the Spoke Layer, only properties that are both true of the item and relevant to the current task context. Input and output patterns consist of the features of the relevant concept within a single modality (M1, M2, M3), resulting in 9 different 'tasks' as each of the three modalities may be the input or output modality (including mappings from one modality to itself, for instance in a task where the participant views a visual image and must "hold it in mind" after the stimulus vanishes, or where a word is repeated). All other input units are set to 0 and all other output units have targets of 0. The input and output modalities relevant to a given trial are specified by the context signal coded in a 'Control Layer' possessing 3 units, each corresponding to a single modality. A control unit is active when the associated modality is relevant to the current task as either input, output, or both. Trainable unidirectional weights from the control layer to the rest of the network thus allow the model to gate the flow of activity on the basis of the context signal so that only task-relevant properties become active in the spokes. The control connections can project to every layer of the semantic network (simulation 1), or selectively to the Spokes Layer, the first layer of hidden units, or the hub (simulation 2). In simulation 2, these connections were applied probabilistically, to equate the number of connections from the Control Layer to the rest of the model (matching the 18 connections possible in the Control to Hub Layer version).

Maturation of cognitive control. Within this framework we aimed to assess whether delayed maturation of control promotes better learning of conceptual (i.e., full context-insensitive, crossmodal) similarity structure and/or shorter time to mature performance. To test this, the model was trained for an initial *control-free* period during which the control units always adopted values of zero and the model was trained with targets on *all* of an item's associated properties across all modalities. This is the typical framework for conceptual models without control requirements; the models are trained to produce the full set of output features, receiving feedback on each feature. The same training was provided to the initial models assessing multimodal conceptual abstraction without context-sensitive behaviour in Jackson et al., (2021). This is analogous to rewarding a child for item-relevant responses regardless of their task relevance, such as rewarding either the answer "choo choo" or "train" (but not, for example, "moo" or "cow") when tasked with naming a train. *Controlled* training items were then introduced into the environment, with gradually increasing frequency, where control units indicating the current task were activated for each learning pattern and the model was trained to activate only those units true of the item and relevant to the task (e.g., only visual and word units for picture naming).

To understand how the onset of control impacted learning, we varied (a) the *maturational delay*, that is, duration of the training period without control, ranging from 0 to 5000 epochs in 1000 epoch increments, (b) the *transition function* describing how the frequency of controlled learning trials increased over time (linear or sigmoidal), and (c) the size of the *transition window* over which controlled trials ramped up from 0-100% of training trials (500, 1000, or 2000 epochs). The maturational delay indicates the number of epochs from the onset of training to the midpoint of the transition window (where there is an equal probability of selecting an item with and without control demands). Varying the transition window changed the number of epochs before the probability of sampling a controlled item shifted, but did not change the total number of shifts (which was always 10). We also trained a comparison model in which control was never

introduced, as a reference point for interpretation of the other models. Similarly, a model with the instant addition of full control requirements was included for comparison only (see Supplementary Materials 1). Note that that there are 48 total training patterns in the control-free period (inputs for 16 items in each of 3 modalities) whereas there are 144 total training patterns in the controlled period (9 different tasks for 16 items). Each epoch included 48 training patterns sampled stochastically from amongst all patterns (with and without control) according to their specified probabilities. This resulted in the same number of learning experiences (48) and weight updates (one) per epoch with or without control, allowing for a direct comparison of training time. Control trials were introduced by gradually changing the probability of sampling a control versus no control pattern from 0 to 1 according to either the linear or sigmoidal ramping function implemented across a transition window of 500, 1000, or 2000 epochs.

Simulation details. All simulations were conducted within the Light Efficient Network Simulator (LENS, version 2.63) software (Rohde, 1999) with identical parameters to Jackson et al., (2021), except that we sampled initial weights from a narrower uniform range (-0.2 to 0.2) to promote a higher level of consistency across runs. The models were fully recurrent continuous-time networks using a temporal integration constant of 0.25, with activity unfolding across 24 time-steps, and input provided in the first 12. Target values were applied throughout. Training used stochastic gradient descent to minimise cross-entropy error loss using a learning rate of .001, weight decay of .0001, and no momentum. All models were trained to the same performance criterion: the model must generate the correct, task- and item-specific pattern of activation across all output units to within 0.2 of their target values. This means that properties true of an item but irrelevant to the task must be correctly inhibited; properties appropriate to the task but not true of the item must likewise be inhibited; and properties both relevant to the task and true of the item must be activated, meeting the central challenges of controlled semantic cognition. Training halted when all output units met this criterion.

Assessment metrics. To understand how the connectivity and maturation of control impact simulated conceptual development, we considered two metrics. The first is *training time,* measured as the number of learning epochs required before the model generates the correct task-specific outputs for all *controlled* patterns in the environment. The second is the *conceptual abstraction score*, defined as the degree to which learned representations in the trained model capture the full similarity structure of the environment abstracted across items, learning episodes, contexts and modalities (Jackson et al., 2021; McRae, Cree, Seidenberg, & McNorgan, 2005; Rogers & McClelland, 2004). Good conceptual representations should express similarity amongst items in the environment based on their full complement of properties, regardless of which properties are observed or inferred in a given situation. Whilst behaviour must be context-sensitive, such context-invariant, multimodal conceptual representations are critical to allow past events to inform semantic behaviour through generalisation across tasks, events and sensory modalities (Jackson et al., 2021; Lambon Ralph et al., 2017; Lambon Ralph & Patterson, 2008). To measure conceptual abstraction, we first computed a *target conceptual similarity matrix* containing the pairwise correlations between all items in the model environment based on their full set of properties across modalities. For each hidden layer in a given model we also computed a *representational similarity matrix* containing pairwise correlations for all items in the evoked pattern of activation generated across units in the layer by each item input (at the final time point within a trial). Conceptual abstraction was defined as the Pearson correlation between the representational similarity matrix and the target matrix. We report this for the Hub Layer only since it always had the highest score, but values for the Intermediate Layer are reported in Supplementary Materials 2.

For each simulation condition we trained 80 models differing only in their weight initialisation to calculate the conceptual abstraction scores and 25 models to compute training time. The same sample sizes were used (here and throughout) as in Jackson et al., (2021) which are far higher than typical modelling investigations as large effect sizes are expected in simulation data. To

understand whether and how the delayed onset of control impacted these metrics, we first used an ANOVA to assess how they were affected by three factors governing the transition to fully controlled trials—maturational delay (i.e., duration of the period without control), the transition function governing the ramp-up of control, and the length of the transition window. We then assessed the nature of the effect of maturational delay on both metrics using independent-samples t-tests to contrast each delay period to the baseline condition in which the model was trained only with control trials from the onset of training (as in Jackson et al., 2021). Where necessary (determined using Levene's test) t-tests were corrected for unequal variance between the groups, and all p-values were Bonferroni-adjusted for 5 comparisons. For interpretation of the size of these effects, Cohen's d effect sizes and 95% confidence intervals are reported. For ANOVAs partial eta squared is reported. Code to run all simulations and process the results is available at github.com/JacksonBecky/developmental.

Results

Simulation 1: Effects of delayed control with full connectivity from control. The first simulations used an architecture with full connectivity from the control units throughout the network (all three connections shown in Figure 1) as previously explored in the prior work. These trainable weights allow the Control Layer to influence the flow of activation through the model at all layers. Precisely which units were influenced by control, and in what manner, was shaped solely by learning. The ANOVA showed that, in this model, the length of the maturational delay reliably influenced both conceptual abstraction ($F_{(5, 395)}$=81.031, p<.001, η^2=.788) and training time ($F_{(3.468, 83.228)}$= 177.248, p<.001, η^2=.881). Neither the transition function (conceptual abstraction score; $F_{(1, 79)}$=0.104, p=.748, η^2=.001; training time; $F_{(1, 24)}$=3.668, p=.067, η^2=.133), the transition window (conceptual abstraction score; $F_{(2, 158)}$=1.609, p=.203, η^2=.040; training time; $F_{(2, 48)}$=1.195, p=.312, η^2=.047), nor their interactions explained

significant additional variance in either metric, so the data were collapsed across these factors to further analyse the effects of maturational delay.

Figure 2. The effects of a maturational delay in learning control on conceptual abstraction, learning trajectory and training time. Control is added in full from the start, or after 1000, 2000, 3000, 4000 or 5000 epochs and the Control Layer has full connectivity to the rest of the model. Results are collapsed across the various protocols for gradually adding control. A. The effects of a maturational delay in control on the conceptual abstraction score. B. The effect of a maturational delay in control on the time taken to train the model. A&B. Bars signify the median value and the 25th and 75th percentile values. White asterisks signify a significant effect of the maturational delay in adding control in contrast to including control requirements from the start of training. Dashed lines allow easy comparison of each delay period to this baseline. C. The effects of a maturational delay in control on the evolution of the mean conceptual abstraction score across training. D. The effects of a maturational delay in control on the trajectory of the representation of the context signal, or control structure in the Hub Layer. C&D. The mean similarity of the representations in the Hub Layer to each structure is displayed after every

1000 epochs of training. Results for a model where control is never added is shown with a dashed line for comparison only as this model would never meet all core requirements of the human semantic system. A maturational delay results in greater representation of the conceptual than the contextual structure in the deep hub throughout training. In contrast, including control requirements throughout results in first learning the control structure in the deep hub before replacing it with the contextual structure. E&F. The effect of the developmental period without control on conceptual abstraction and training time are displayed for each of the different protocols and lengths for adding control (red = sigmoidal, blue = linear, lighter = control added over fewer epochs, darker = control added over more epochs). An additional instant protocol is shown for comparison (black). Error bars represent the standard error of the mean.

Figure 2.A. shows the distribution of conceptual abstraction scores across model runs at each level of maturational delay. Compared to the baseline condition (zero delay), a short maturational delay reliably improved conceptual abstraction (at 1000 epochs; $t(958)=2.758$, $p=.030$, CI=.015, .003; at 2000 epochs; $t(958)=4.792$, $p<.001$, CI=.008, .020); an intermediate delay showed no difference (3000 epochs; $t(958)=.867$, $p=1$, CI=-.003, .009); and longer delays showed significantly worse scores (4000 epochs, $t(958)=-5.430$, $p<.001$, CI=-.023, -.011; 5000 epochs, $t(958)=-13.015$, $p<.001$, CI=-.048, -.036). These effect sizes were, however, relatively small except for the most extreme delay (for improvements at 1000 and 2000, $d = 0.178$ and 0.309 respectively; for decrements at 4000 and 5000, $d = 0.351$ and 0.840 respectively). Thus, increasing the maturational delay produced an inverted U-shaped effect on the quality of the model's conceptual representations with modest initial improvement followed by modest reductions and significant degradation only at the longest lag.

Figure 2.B. shows the distribution of training time across model runs at each maturational delay level. Here increasing maturational delay was unambiguously and monotonically associated with shorter training times relative to baseline (1000 epochs, $t(298)=-1.349$, $p=.891$, CI=-541.81, 101.06, $d=.156$; 2000 epochs, $t(298)=-7.421$, $p<.001$, CI=-1415.36, -822.02, $d=.857$; 3000 epochs; $t(285.992)=-14.576$, $p<.001$, CI=-2426.90, -1849.43, $d=1.683$; 4000 epochs; $t(256.616)=-20.307$, $p<.001$, CI=-3029.77, -2494.09, $d=2.345$; 5000 epochs; $t(220.638)=-24.701$, $p<.001$, CI=-3403.65, -2900.66, $d=2.852$). Based on these Cohen's d values, a maturational delay of 2000 epochs or more produces a large or very large reduction of training time; indeed, a sufficiently long developmental period without control can reduce the training time by over a quarter.

Interim Discussion

Both patterns are counterintuitive. Recall that conceptual abstraction measures how well the model's learned representations capture a target conceptual structure, computed by considering all of an item's properties tabulated across all modalities, regardless of which modalities are relevant for the current task context. Prior to the onset of control, the model is trained to directly activate all such properties—thus one might expect that a longer maturational delay should produce monotonically *better* conceptual abstraction. Whilst small improvements in conceptual abstraction are possible with a short delay, increasing the delay further resulted in a *decrement* to conceptual abstraction. Overall, delaying control only induced small changes in conceptual abstraction, except where very long delays resulted in a large decrement. With regard to training time, recall that the criterion for mature performance is based solely on the model's ability to generate the correct output for the controlled, task-specific patterns. One might expect the fastest path to this objective would be to train the model with such patterns from the outset. Instead this condition produced the *longest* training time. The more we delayed the introduction of controlled training patterns, the shorter the overall training time to master these patterns. The addition of a moderate delay (between one third and one half of the subsequent learning time) provides a substantial reduction in training time whilst maintaining the same or a slightly reduced level of conceptual abstraction.

To better understand these effects we undertook some additional simulations and analyses. We first considered how the similarity structure of the hub representations evolved over time at each maturational lag. In each delay condition, we trained 80 independent models for 10,000 epochs (regardless of the performance criterion), stopping the training every 1,000 epochs and recording the patterns of activation evoked over the hub units for each item in every task. We then considered how the similarities amongst these learned representations correlated with two different target matrices: (a) the conceptual similarity matrix used to generate the conceptual

abstraction score, and (b) a *contextual similarity matrix* that indicates the degree to which different tasks activate overlapping units in the Control Layer irrespective of the item being processed. Thus, the contextual similarity matrix expresses the relevance of the various input and output modalities across trials. The conceptual and contextual similarity matrices are uncorrelated with one another, so their individual correlations with the similarities expressed in model representations indicates how much model representations are influenced by conceptual versus task structure.

Figure 2.C. shows evolving correlations with the target conceptual similarity matrix as the model learns, for each delay condition and as a comparison, for the model trained without control requirements (grey line). When control is present from the outset, correlations with conceptual structure increase gradually and monotonically. With delayed onset of control the model learns conceptual structure much more rapidly in initial epochs, but the introduction of controlled learning patterns significantly disrupts these trajectories, leading model representations to ultimately capture somewhat *less* conceptual structure with long delays than had control been present from the onset. This disruption is larger for longer maturational delays—the more the control-free model has mastered the conceptual structure, the greater the disruption to representations when control is introduced. All model representations recover somewhat from this disruption with moderate lags achieving similar levels of conceptual abstraction to those with control throughout. However, those arising under very late-developing control never "catch up" to acquire conceptual representations of comparable quality to the other models.

To understand why this disruption occurs, consider the most extreme delay, in which the model has learned good conceptual representations and is well on the way to producing the correct pattern of activation for each item across all three modalities—so given the input for *dog* in M1, true properties of *dog* in all modalities are activated and properties not true of *dog* are inactive. When a controlled pattern is introduced—say the model views the dog in M1 and must activate

its properties in M2 but not M3—the units in task-relevant modalities (M1 and M2) are already near the correct state and will generate little error. Likewise units in the irrelevant modality (M3) that are not true of dogs are near the correct state and generate little error. Thus, the error will mainly be driven by activation of units true of the dog but irrelevant to the task, and weight changes throughout the system will be driven to reduce the activation of these units.

Such changes could occur within the semantic representation network itself, or within the weights projecting from the active control units. If the disruption of conceptual structure arises via input from control, then different instances of the same task (mapping from M1 to M2) should disrupt conceptual representations in a similar way, since all such instances are represented in the same way in the Control Layer. Conversely instances of different tasks should disrupt the representations in different ways. Put differently, if the disruption to learned conceptual structure arises via inputs from control, the disrupted hub representations should begin to partially reflect the similarity structure of the task representations. Figure 2.D. shows just this pattern: as control is introduced, representational structure in the hub increases its correlations with task structure, just as correlations with conceptual structure decline. Correlations with task structure peak when the hub representations show the least amount of conceptual structure, and as conceptual structure recovers, the impact of task structure diminishes. This pattern suggests that the disruption of conceptual structure in the hub arises because control units directly impact the patterns of activation there, causing the network to represent a blend of conceptual and contextual structure.

Figure 3 shows the representational similarity structure in the hub at each time point across different lags. For reference, the bottom panels show the conceptual similarity (left), the contextual similarity (middle), and the similarities amongst output activations for all controlled training patterns (right). Where control is present throughout training, the hub initially learns a roughly equal blend of the conceptual and contextual similarities, and over time, the contextual

similarities gradually fade as the conceptual structure comes to dominate. Achieving a high level of conceptual abstraction relies on the functional specialisation of the hub for conceptual similarity, as distilling this crossmodal, cross-context structure is only possible in an area with connections to all sensory modalities. When control is delayed, the hub first encodes conceptual structure, and the longer the delay, the more structure it captures (i.e., similarity matrices increasingly resemble the context-independent crossmodal conceptual structure at bottom left). The introduction of control always produces a large initial effect, with task similarities very clearly apparent, changing the nature of the hub representations to reflect contextual similarity. Conceptual structure gradually re-emerges, yet changes later in learning are slow due to reductions in the effective learning rate with smaller error values. For shorter delays the conceptual structure in the hub is recovered to an equivalent level when the task criterion is reached, yet with longer delays conceptual abstraction never reaches the level observed when control is present from the start.

Figure 3. Visualising the model representations during learning across varying maturational delays in control. Control was present from the start of training or added around 1000, 2000, 3000, 4000 or 5000 epochs (dashed line). Control was added gradually using the sigmoidal protocol over 1000 epochs in the model with connectivity between the Control Layer and every other layer. Similarity matrices were constructed comparing activation in the Hub Layer across all the examples in each context. For comparison the target conceptual similarity matrix used to determine the conceptual abstraction score (i.e., the context-independent representation structure) is displayed, alongside the contextual similarity structure (reflecting the context signal only, which denotes the input and output modality of each trial) and the full structure (the combination of these contextual and conceptual structures). The Hub Layer specialises in the abstracted conceptual representation structure. Without control this functional specialisation is present in the hub from the start of training, whereas with control the context signal has a large initial effect on these deep representations.

These observations explain why long-delayed control degrades conceptual abstraction in the hub. Initially one might expect degraded abstraction to produce longer learning times as well: if the model does a poorer job of representing cross-item, cross-modality structure, then learning should generalise less well across related times, leading to slower overall learning. Yet we observed *faster* learning the longer control was delayed. To understand why such speeding may

occur, consider again what happens in the longest delay condition: the model comes close to mastering all properties for all items across all modalities prior to the onset of control. This in turn means that, for any given controlled-pattern, the majority of output units will already be in the correct states: given a task where the network views *dog* in M1 and must generate output in M2 but not M3, all M1 and M2 units will be near the correct active or inactive state, but so will many units in M3—namely, all those that are not properties of dogs. If each item activates half the properties in each domain, then learning prior to the onset of control will place 5/6 of the output units in the correct state for any task that maps from one domain to another. Even for tasks involving only a single modality (such as retaining a visual stimulus in mind when it has vanished), all units in the relevant modality and half the units in the other modalities—or 2/3 of output units in total—will be in the correct state. The sparser an item's properties are in each modality, the more units in irrelevant modalities will already be in the correct state as control trials are introduced. In other words, extensive experience with full patterns prior to control benefits performance for the majority of units in each controlled pattern.

Simulation 2: Effect of control connectivity. Simulation 1 showed that, when control connects to all layers of the semantic network, a maturational delay to control onset can modestly improve conceptual abstraction or greatly speed training time. With a long delay, the large reductions in training time come at the cost of significantly reduced conceptual abstraction. However, a moderate delay can result in much faster training, without large differences in conceptual abstraction. Subsequent analyses suggested that the disruption to conceptual structure reflects the influence of task structure on hub representations as control trials are introduced. However, previous findings have demonstrated the importance of avoiding direct connections between control and the hub (Jackson et al., 2021). Would connections restricted to shallower layers show the same pattern of changes with a maturational delay? Would the delay obviate the need to avoid a direct connection from control to the hub, or otherwise change the optimal location of this connection? To answer these questions we considered whether the benefits of delayed

control depend upon the locus of the connection from control. Simulation 2 replicated simulation 1 but restricted connections from the Control Layer so that they projected only to (1) the spokes, (2) the intermediate hidden layer, or (3) the hub. Additionally since the transition function and width of the transition window had no effect in simulation 1, we employed only a single transition protocol involving a sigmoidal function extended over 1000 epochs. 80 models were trained for each architecture at each maturational delay. As before we measured the influence of these factors on the model's conceptual abstraction score at the end of learning, and on the total training time.

Results are shown in Figure 4. Conceptual abstraction scores were influenced by the location of the control connection ($F(2, 158)=3414.917$, $p<.001$, $\eta^2=.977$), the maturational delay ($F(5, 395)=14.449$, $p<.001$, $\eta^2=.155$) and their interaction ($F(10, 790)=9.806$, $p<.001$, $\eta^2=.110$). Conceptual abstraction was always significantly worse when control connected to the Hub versus the Intermediate ($t(822.103)=-56.164$, $p<.001$, $CI=-.228, -.213$, $d=3.989$) or Spokes Layers ($t(746.774)=-61.795$, $p<.001$, $CI=-.240, -.225$, $d=3.99$), replicating Jackson et al., (2021). When control connected to the hub, maturational delay improved conceptual abstraction (at 1000 epochs; $t(158)=2.989$, $p=.016$, $CI=.012, .057$, $d=.473$; 2000 epochs; $t(158)=5.187$, $p<.001$, $CI=.037, .083$, $d=.820$; 3000 epochs; $t(158)=5.761$, $p<.001$, $CI=.042, .086$, $d=.911$; 4000 epochs; $t(158)=2.895$, $p=.022$, $CI=.011, .057$, $d=.458$; 5000 epochs; $t(158)=3.381$, $p=.005$, $CI=.016, .059$, $d=.535$), but never produced scores comparable to other connection conditions. Thus, even with maturational delay, conceptual abstraction is best promoted by architectures that prevent the direct impact of control on the hub.

Control connections to the Spoke versus Intermediate Layer produced reliably better conceptual abstraction ($t(934.139)=4.265$, $p<.001$, $CI=.006, .018$) but the effect size was small ($d=.275$). In neither case did maturational delay produce the inverted-U shape observed in simulation 1: both models showed equally good conceptual abstraction for delays up to 3000 epochs (Spokes Layer

connection; at 1000 epochs; $t(158)=-1.868$, $p=.318$, CI$=-.022$, .001; at 2000 epochs; $t(158)=-1.993$, $p=.240$, CI$=-.024$, .000; at 3000 epochs; $t(158)=-2.488$, $p=.069$, CI$=-.026$, -.003; Intermediate Layer connection; at 1000 epochs; $t(158)=-0.134$, $p=1$, CI$=-.014$, .013; at 2000 epochs; $t(158)=0.163$, $p=1$, CI$=.011$, .013; at 3000 epochs; $t(158)=-1.460$, $p=.731$, CI$=-.022$, -.003). As with simulation 1, longer lags produced reliable decrements in both cases, with only the longest lag producing a large effect size (contrast to baseline for connections to the Intermediate Layer: at 4000 epochs; $t(158)=-4.886$, $p<.001$, CI$=-.043$, -.018, $d=.773$; at 5000 epochs; $t(158)=-8.021$, $p<.001$, CI$=-.068$, -.041, $d=1.268$; for connections to the Spokes Layer: at 4000 epochs; $t(158)=-3.616$, $p=.002$, CI$=-.035$, -.010, $d=.572$; at 5000 epochs; $t(158)=-5.369$, $p<.001$, CI$=-.045$, -.021, $d=.849$).

Figure 4. Assessing the effects of a maturational delay in learning control on conceptual abstraction score and training time with control connections to the different layers. The Control Layer is connected to the Spokes Layer (black), Intermediate Layer (green) or Hub Layer (orange). Control is present from the start, or added gradually around 1000, 2000, 3000, 4000 or 5000 epochs (with a sigmoidal protocol over 1000 epochs). A. The effect on conceptual abstraction. B. The effect on training time. Bars signify the median value and the 25th and 75th percentile values. White asterisks signify a significant effect of the delay in adding control in contrast to the inclusion of control requirements from the start of training. Dashed lines allow easy comparison of each delay period to the baseline level where control is present from the start.

Training time also varied with control connection ($F_{(2, 48)}=40.169$, $p<.001$, $\eta^2=.626$),

maturational delay ($F_{(5, 120)}=32.659$, $p<.001$, $\eta^2=.576$), and their interaction ($F_{(6.153,}$

$_{147.668)}=3.612$, $p=.002$, $\eta^2=.131$; see Figure 4.B.). Across all connection loci delayed maturation

sped learning (for each contrast to baseline t(298)>4.2, p<0.001). However, the extent of the speeding was greater for the deeper connections (Hub Layer maximum d=2.221; Intermediate Layer maximum d=2.139) than the Spokes Layer (maximum d=1.106). As the connection to the Intermediate Layer also started with lower training times than the control projection to the hub, the fastest training occurred when control projected to the Intermediate Layer (contrast to Spokes Layer connection t(298)=-9.958, p<.001, CI=-1882.411, -1261.134, d=1.150; contrast to Hub Layer connection t(298)=-5.466, p<.001, CI=-1210.621, -569.632, d=.631). At all lags, a control connection to the Intermediate Layer resulted in the fastest training. With this connection locus, as in Simulation 1, moderate and long maturational delays decreased training times substantially (at 1000 epochs; t(48)=-1.901, p=.316, CI=-1494.258, 41.778, d=.316; at 2000 epochs; t(48)=-4.268, p<.001, CI=-2313.239, -831.721, d=1.207; at 3000 epochs; t(38.141)=-4.369, p<.001, CI=-2267.188, -831.532, d=1.236; at 4000 epochs; t(48)=-5.252, p<.001, CI=-2821.033, -1258.97, d=1.485; at 5000 epochs; t(31.551)=-7.563, p<.001, CI=-3177.669, -1842.971, d=2.139).Thus, with an intermediate control connection locus, a moderate maturational delay (3000-4000 epochs or around half the subsequent training time) results in greatly speeded training in the context of relatively preserved conceptual abstraction.

Interim Discussion

Simulation 2 demonstrates that maturational delay and system connectivity can conspire to promote faster overall learning with high levels of conceptual abstraction. As in prior work, direct connections from control to the hub impair its ability to uncover the latent conceptual structure across contexts and modalities even with the positive effects of a maturational delay. Unlike Simulation 1, when the absence of these direct connections allows the semantic hub to remain insulated from control inputs, maturational delay cannot improve conceptual abstraction further by protecting the hub from control signals early in learning. However, the positive effect of a maturational delay on training time is preserved in models with control connections to the

Spokes or Intermediate Layers. Whilst the effects of connecting to Intermediate versus Spokes layers appear similar when control is presented throughout training, these large improvements in speed were greater when control connected to the Intermediate Layer. Thus, the present results highlight the benefit of both a maturational delay in control and a control connection to intermediate regions. Whilst long delays result in compromised conceptual abstraction ability, moderate delays can have very large speeding effects with either fully or relatively preserved conceptual abstraction. The optimal length of the delay is around one third to one half of the subsequent learning time, depending whether some trade-off with conceptual abstraction is considered advantageous.

Part 2: Meta-analysis of Developmental Data

The simulations suggest that delayed onset of semantic control may ultimately promote more rapid acquisition of mature controlled semantic cognition. Does the literature contain evidence of such a delay? While the development of executive function (i.e., domain-general control) and of conceptual knowledge (i.e., semantic representation) have each been extensively studied, there is relatively little work focusing explicitly on the development of semantic control. There is, however, extensive research arising from an early controversy in conceptual development that can shed light on the central question of a hypothesised maturational delay in semantic control.

This controversy arose from Inhelder and Piaget's (1958) early view that children's conceptual representations first capture thematic associations that eventually give way to taxonomic relationships, and the subsequent competing view that early concepts primarily reflect taxonomic structure at either basic (Rosch et al., 1976) or even more general or abstract (Mandler & McDonough, 1993) levels. To adjudicate these possibilities, many studies have employed a task now known as *triadic match-to-sample*: the participant must decide which of two items best

matches a third sample item. For instance, a participant might judge which "goes better with" a picture of a horse: a saddle or a deer? Choice of the saddle indicates a thematic match, since horses and saddles occur together in many scenarios but are distinct kinds of things. Choice of the deer indicates a taxonomic match, since horses and deer don't often occur together in familiar scenarios but are similar kinds of things (i.e., four-legged land mammals).

Throughout the 1970's and 80's many such studies were conducted, with some favouring the view that taxonomic knowledge trumped thematic knowledge and others suggesting the reverse (e.g., Bauer & Mandler, 1989; N. W. Denney, 1974; Greenfield & Scott, 1986; Smiley & Brown, 1979). Eventually it became clear that the results depend critically upon the wording of the instructions (Estes et al., 2011; Lin & Murphy, 2001; Waxman & Namy, 1997)—that is, upon the task context. If asked to choose which item has the same name, children most often selected a taxonomic match. If asked which item "goes better with" the target, they more often selected the thematic match. In 2001, Lin and Murphy conducted a large triadic matching study with adult participants, assessing the effects of many different instructional contexts. The results revealed a gradation of choice preferences, with some contexts strongly eliciting taxonomic choices (e.g., "called by the same name"), others reliably eliciting thematic choices ("goes with"), and still others yielding somewhat ambiguous results. The latter findings indicate how the triadic matching task indexes semantic control when the choice options include both a taxonomic and thematic match. In such cases the semantic representation system yields information about precisely how each option matches the target, and the participant must use some representation of the current goal to select which kind of information is most relevant to the task. The instructional cue is a form of context, providing this goal representation. Successful semantic control is signalled by the ability to reliably choose the taxonomic option under one instructional context and the thematic option under another. Suppose, however, that a participant fails this task by choosing unsystematically. Such a failure might arise (a) because the participant is unable to use the control system to choose the correct kind of semantic information, or (b) because the

participant does not possess the semantic knowledge needed to choose correctly. These possibilities can be adjudicated by assessing whether the participant correctly matches a given item for triads where the second option is *unrelated* to the sample. If the participant is able to choose the correct match in the context of unrelated distractors, this indicates that they possess sufficient knowledge of the relevant semantic information to find the target under a given set of instructions. That is, selecting the match when the target appears with an unrelated item does not strongly tax semantic control. Thus, a signature of good semantic knowledge with underdeveloped semantic control would arise when a participant succeeds at a given kind of matching when the second option is unrelated to the target, but fails when the second option is related to the target in a manner irrelevant to the task context. With these ideas in mind we undertook a meta-analysis of the developmental literature on triadic matching for taxonomic and thematic knowledge, aiming to assess whether knowledge of taxonomic and thematic relations emerges prior to, or in parallel with, semantic control.

Methods

We conducted a literature review seeking match-to-sample studies employing either thematic or superordinate taxonomic matching or both. A match-to-sample task is defined as presentation of a single item as a probe or standard for comparison, and two or more targets requiring an active choice to decide which is related to the standard in some way (indicated via pointing, picking up an object, or a spoken response; preferential-looking paradigms were excluded). Stimuli could be pictures, words or objects. The measure of interest was the frequency of choosing the taxonomic or the thematic match; studies reporting reaction time or individual differences only were excluded. We then considered two kinds of studies:

   (1) *Low control*, where the target matched the sample either taxonomically or thematically, and remaining options were semantically unrelated to the sample. These trials thus indicated

whether the participant possessed taxonomic or thematic knowledge sufficient to choose the target without having to select what kind of semantic information was relevant for the task.

(2) High control, where all trials have both a thematic and taxonomic match but the instructional context indicates which kind of information should drive the selection. These trials thus require both the relevant semantic knowledge and the ability to select the correct kind of information for the task.

Studies were initially identified from two sources: the list of developmental studies of taxonomic and thematic structures in Mirman et al., (2017) and a Web of Knowledge search with any two of the terms 'taxonomic', 'thematic' and 'perceptual' or with the term 'noun-category bias', performed in January 2021 and including full papers from any year. This resulted in an initial screening of 383 studies. The references of all included studies were also screened for inclusion. Studies were excluded if participants were explicitly rewarded for taxonomic vs. thematic choices, or if the studies intentionally weighted the taxonomic and thematic responses to be stronger or weaker than each other. Studies could include additional response options which were factored out to assess only the contrast of interest. Studies with more than two choices were corrected to a chance level of 0.5. All studies assessed healthy participants without neurostimulation during childhood or young adulthood (studies with mean ages over 55 were excluded to avoid possible effects of ageing).

For the low control studies we did not further sort based on instructions as any semantic relationship cue should lead participants to choose the correct (taxonomic or thematic) target over an unrelated concept. For the high control studies we analysed the instructions provided to participants and selected only those previously identified as reliably promoting a thematic or taxonomic decision in adults (D. R. Denney & Moulton, 1976; Estes et al., 2011; Greenfield & Scott, 1986; Lin & Murphy, 2001; Markman & Hutchinson, 1984). Thematic contexts were those cued by 'goes with', 'goes best with' or 'goes together' instructions, which imply concepts are

found together in time or space. Taxonomic decisions were cued by the provision of a novel label for the standard (e.g., 'this is a dax') and the request to find another item in this category (e.g., 'find another dax'). The novel word acts as a basic noun which typically relates to the taxonomic status of the concept (referred to as the noun-category bias; Waxman, Senghas, & Benveniste, 1997). Studies using ambiguous instructions unlikely to strongly weight the context toward taxonomic vs. thematic cues in adults (e.g., choose 'the most related' or 'more similar' item) were excluded.

For each study the average proportion of taxonomic or thematic responses was recorded, as well as the standard deviation (where present) and the number of participants. Studies were chunked on the basis of age (i.e., studies with an average age between 12 and 24 months were grouped as one-year-olds, etc.) from ages one to seven and adults (ages 18-55). For low control conditions we found 18 taxonomic-match experiments consisting of 639 participants across 28 data points and 17 thematic-match experiments with 657 participants across 29 data points. For high control conditions, we found 20 experiments employing a clear thematic cue, consisting of 805 participants over 35 data points, and 22 experiments employing a clear taxonomic cue, consisting of 561 participants over 29 data points. Thus, data from 2662 participants contributed to the assessment of control and representation ability across development. See Supplementary Materials 3 for full details of the studies included at each age, including the data. The methods used throughout comply with APA ethical standards, as all data was secondary (or simulated).

Statistics were performed in SPSS (IBM, 2017) on the average proportions of taxonomic or thematic responses in each study. To determine whether each age group had evidence of a conceptual representation structure, the proportion of correct responses in the taxonomic-only and thematic-only conditions were compared to chance (0.5), using one-tailed one-sampled t-tests. To test for evidence of significant context effects, the proportion of taxonomic (vs. thematic) responding in studies with a taxonomic or thematic instruction was compared using

one-tailed between-samples t-tests (as the specific expectation was that a taxonomic cue should have greater taxonomic responding). Where possible, effect sizes (g) were calculated using Hedges correction for unequal sample sizes. To fully understand the distribution of responses and for graphical purposes, the response distribution across all studies was estimated based on the reported mean and standard deviation per study. The majority of studies reported the variance. In some cases this was reported separately for different conditions and then combined. Where variance was not reported, the maximum possible standard deviation was computed. This combination of reported and maximum standard deviations provides the best estimate of the real variance whilst being conservative. However, the same pattern of results is obtained if the maximum possible standard deviation is used for all studies. These analyses were performed using R Statistical Software (v3.5.2; R Core Team, 2021) using custom built code available at github.com/JacksonBecky/developmental. For each study, the results were simulated 1000 times by drawing a number of samples equal to the number of participants in the study from a normal distribution with the reported mean and standard deviation. These simulated data points were used to compute a grand mean (across all studies and simulation iterations, which is weighted by the number of participants), which is displayed in the plot. The .025 and .975 quantiles of this distribution were used to compute the 95% confidence intervals of the mean across studies. Where the confidence intervals for thematic or taxonomic over unrelated selections reliably exclude 0.5, this further supports the conclusion that this form of knowledge is present above chance level. Where the confidence intervals for the taxonomic and thematic cues reliably exclude each other, this further supports the presence of context-sensitive responding to differential cues.

Transparency & openness. We report how we determined our sample size, all data exclusions, all manipulations and all measures in all studies within this article. We follow JARS (Kazak, 2018). All data are available in the Supplementary Materials or can be generated with the code provided. All code is available at github.com/JacksonBecky/developmental. Data were generated using

LENS (v2.63; Rohde, 1999) and analysed using Matlab (The MathWorks, 2018), SPSS (v.25; IBM, 2017) and R Statistical Software (v3.5.2; R Core Team, 2021). The study design and analyses were not preregistered.

Results

Figure 5 shows the mean proportion of taxonomic (top end of y axis) or thematic (bottom end) choices at each age and for each experiment type. Studies with children as young as one showed chance behaviour, but by age two children chose the correct taxonomic or thematic match reliably more often than chance, so long as other options were semantically unrelated to the sample (taxonomic; $t(3)=5.207$, $p=.007$; thematic; $t(3)=2.98$, $p=.029$). When choices contained both a taxonomic and thematic match, two-year-olds chose equivalently (and near chance) regardless of the instructional context (taxonomic vs. thematic cues; $t(3)=2.97$, $p=.393$). Thus, by age two children appear to possess knowledge of both taxonomic and thematic relations, but are unable to use instructional context to determine the correct basis for matching. For ages 3 and 4, there were significant differences in responses to the two cue types, demonstrating the presence of some context-sensitivity (taxonomic vs. thematic cues; age 3; $t(13)=2.222$, $p=.045$; age 4; $t(17)=3.985$, $p<.001$). It is not clear whether these children can utilise both the taxonomic and thematic cues, or whether the (stronger) taxonomic cue is driving this difference alone (whilst thematically cued responding remains unsystematic). However, it is clear that this context-sensitivity is not yet at adult levels (taxonomic vs. thematic cues; age 5; $t(5)=3.38$, $p=.01$; age 6; $t(1)=23.286$, $p=.014$; age 7; $t(1)=26.353$, $p=.024$; adult; $t(10)=11.972$, $p<.001$). Context-sensitivity increases in strength throughout development (age 3; $g=1.082$; age 4; $g=1.748$, age 5; $g=2.171$; adult; $g=7.364$) and from age 5 onward children also display a preference for the thematic response when presented with a thematic cue. Context-sensitivity shows a large increase and approaches adult levels around the age of 6. Thus consistent with the model predictions,

children acquire the relevant semantic information and can deploy it to perform the matching task as young as 2, but cannot begin to make systematic use of contextual information to guide task-appropriate semantic choices until age 3, with large increases in context-sensitivity around 6 years.

Figure 5. The emergence of semantic control after semantic representation in our meta-analysis of triadic comparison. Top panel: Examples of a match-to-sample task illustrating the two low control conditions (taxonomic vs. unrelated and thematic vs. unrelated concepts) and the two high control conditions (both including a

taxonomic and a thematic match but differing in the instructional cue). The correct answers are highlighted in green. Bottom panel: Mean and bootstrapped confidence intervals indicating the proportion of times participants in different age groups chose the taxonomic (high proportion) or thematic (low proportion) target in each study type. Knowledge of taxonomic or thematic relationships is indicated by non-chance responding in the low control conditions (light grey and dark grey lines respectively, light or dark grey asterisks denote responding significantly better than chance, not all contrasts are possible). The ability to use the instructional cue to inform the choice of an appropriate match in high control conditions is indicated by the difference between the proportion of taxonomic versus thematic responding for the taxonomic (purple line) and thematic (orange line) cues. Significant context-sensitive effects indexing semantic control ability, are signified by the black asterisks.

General Discussion

Building on prior suggestions (Chrysikou et al., 2011; Newport, 1990; Thompson-Schill et al., 2009), we considered whether the delayed maturation of control might benefit development of controlled semantic cognition. Using a neural network architecture shown in prior work to promote both conceptual representation and semantic control, we assessed the effects of delayed control in scenarios where (1) control regions connect to the entire semantic representation network, or (2) control connects only to specific levels of the network. In both cases, maturational delay promoted conceptual development. With full connectivity from control, a short delay modestly improved the structure of conceptual representations while a longer delay degraded such structure, a U-shaped effect attributable to the presence of direct connections from control representations to the semantic hub. When control systems connected to shallower parts of the network, the model showed little cost to representational structure except at the very longest delays, and when control connected only to the multimodal hub, the model acquired poor-quality representations regardless of delay. Thus, anatomical connectivity that insulates the hub from the direct effects of control promotes good learning of conceptual structure even when control is substantially delayed. In all architectures, a maturational delay improved speed of learning. The fastest training was observed with both a maturational delay and control connected to intermediate model layers. The simulations thus suggest that delayed onset of control, combined with anatomical structure that insulates the semantic hub from direct control, leads to substantially faster mastery of controlled semantic cognition, and consequently that the slow

maturation of executive function may be adaptive in this regard. Consistent with this view, a new meta-analysis of the classic match-to-sample task showed that, while children as young as two years old possess taxonomic and thematic semantic information sufficient to choose a correct match from among unrelated distractors, they do not have the semantic control processes required to produce context-sensitive responses when taxonomic and thematic matches compete for selection. Subtle context-sensitivity arises from age 3 and increases across development, with reliable use of both instructional cues occurring around 6 years. These data thus suggest that maturation of semantic control lags behind acquisition of conceptual knowledge and representations, as predicted by the models.

Why does maturational delay speed learning? The simulations suggest two reasons why delayed onset of control may ultimately speed mature semantic cognition. First, prior to the onset of control, each learning experience can activate all known features of a concept, whereas with control, each experience activates only a subset of features. Training without control thus effectively increases the amount of exposure to each feature, producing faster learning and good approximation of cross-modal representational structure. Second, in allowing the different features of a concept to co-activate regardless of the current task, a period with limited control can help the system learn to use highly similar conceptual representations across tasks, rather than learning distinct mappings in different contexts. That is, delayed control may avoid a key challenge for controlled semantic cognition, namely the need to extract context-invariant knowledge whilst simultaneously producing context-sensitive responding. Instead, context-invariant representations may be acquired before there is a strong requirement for context-sensitive behaviour and the presence of a "base" conceptual representation may then allow the continued use of shared representations across different tasks later in development.

Importantly, these effects depend upon the anatomical structure of the overall network. The beneficial effect of delayed control on learning speed is greatest when control representations

connect to intermediate network layers, whilst acquisition of conceptual structure depends upon insulating the hub from direct influence of control. The observed differences cannot be simply due to the amount of experience with each feature, since this is matched across different control connections. Thus, anatomy and maturational delay can conspire to promote fast learning with good conceptual abstraction, by having control interface at an intermediate part of the semantic network that insulates the multimodal hub, and with the maturational delay allowing the full network to lay down base representations that approximate the cross-context structure of the environment before control begins to shape behaviour.

Implications for the development of controlled semantic cognition. Much prior work modelling conceptual development has focused on the emergence of representational structure rather than control. This research has been successful in explaining how sensitivity to multimodal environmental structure can explain a variety of developmental phenomena, such as the progressive differentiation of conceptual knowledge (Mandler & McDonough, 1993; Poulin-Dubois & Pauen, 2017; Rogers & McClelland, 2004; Saxe, McClelland, & Ganguli, 2019), the primacy of basic-level naming (Rogers & McClelland, 2004; Rogers & Patterson, 2007; Rosch et al., 1976), temporary overgeneralisation of frequent object names and properties (Mervis, 1987), and illusory correlations (Chapman, 1967; Rogers & McClelland, 2004). Indeed, the considerable success of this viewpoint may indicate that conceptual development proceeds without the disruptive influence of control. That is, the importance of control for conceptual development may have been neglected precisely because its onset is delayed until much of the core conceptual structure is learnt, as suggested by our meta-analysis of match-to-sample data.

Our results also suggest a new perspective on the development of taxonomic and thematic concepts—specifically, that patterns of developmental change reflect growing sensitivity to contextual information maintained by the maturing control system rather than an overall bias toward learning or deploying a particular kind of semantic information at a given age. Indeed,

changes in the salience of or bias toward taxonomic versus thematic knowledge cannot explain the full pattern of results, since our meta-analysis showed that, across ages, children are equally able to exploit either kind of information when distractors are unrelated to the target (and thus there is little need for control). Instead what changes across development is children's increasing sensitivity to task context. In this sense taxonomic and thematic relationships may be more similar than is often considered, with both being encoded within the same system of representation (Jackson, Hoffman, Pobric, & Lambon Ralph, 2015; Lambon Ralph et al., 2017) rather than in distinct systems as some have proposed (e.g., Mirman, 2017; Schwartz et al., 2011), and with control governing which contributes to behaviour in a given task. Our proposal is also consistent with prior work showing the difficulty that both children (Blaye et al., 2006; Blaye & Jacques, 2009; Mirman, 2017) and healthy adults (Landrigan & Mirman, 2018; Sachs et al., 2008) have in switching between taxonomic and thematic judgments, a task that requires shifting the task representation and hence the semantic criterion of a decision in real time.

Relation to other work on development of semantic control. The emergence of semantic representation prior to the onset of control, as well as the timing of each process, appear consistent with evidence from the few studies directly assessing semantic control in childhood. Whilst children under 5 years old perseverate on subcategories in semantic fluency tasks, older children can switch to a new subcategory when explicitly cued and then learn to employ this as an endogenous strategy (Snyder & Munakata, 2010). Around 5 to 6 years of age, children become sensitive to the presentation of stimuli in a set (Snyder & Munakata, 2013), and learn to switch flexibly between multiple meanings of a word (Blaye et al., 2006; Blaye & Jacques, 2009). Other work documents a qualitative change in semantic cognition around 6 years in which children cease to simply name objects and start to describe them (Funnell, Hughes, & Woodcock, 2006), perhaps reflecting flexible, context-sensitive behaviour. Research in the development of creativity provides another source of evidence, as such work often uses similar tasks to those employed in studies of semantic control (e.g. generating atypical uses of an object) and engages similar brain

areas (Cogdell-Brooke, Sowden, Violante, & Thompson, 2020; Jackson, 2021; Noonan, Jefferies, Visser, & Lambon Ralph, 2013). Children under 6 years old lack 'functional fixedness', performing better than older children or adults on tasks which typically lead to fixation on a particular use of an object in adults. For instance, after a demonstration where a box is used as a container, adults and older children are less likely to solve a problem requiring its use as a step than are younger children (T. C. German & Defeyter, 2000). Similarly, 5-year-olds outperform 7-year-olds at identifying the function of an object not being used for its original purpose (T. P. German & Johnson, 2002) and generating potential uses outside of the design parameters of an object in a functional fluency task (Defeyter, Avons, & German, 2007). These differences are understood to arise because younger children activate more information regarding the object and possess weaker selection processes—ideas that clearly relate to our proposal that early concepts develop with minimal influence from control.

Relation to research on cognitive control. The ideas motivating our approach are closely connected to a recent proposal about the computational trade-offs that motivate the need for control in the first place (Musslick & Cohen, 2021), which in turn relates to recent insights about multi-task learning from computer science. Briefly, any system that must learn mappings from various input to various output channels can do so either via separate, independent pathways that each acquire their own representations, or via a shared substrate that learns a common representation for different mappings (or some blend of these possibilities). When different input/output mappings share common structure, there is a computational benefit to employing a common representation, as learning about one task will promote generalisation to and thus more rapid learning of other tasks. This is the "multi-task learning" insight that allows for zero- or few-shot learning in a variety of AI applications, including machine translation (e.g., Johnson et al., 2017) and game-playing systems (e.g., Hu, Lerer, Peysakhovich, & Foerster, 2020). With a shared representation, however, the system can only compute one input/output mapping at a time, requiring a control system to determine which inputs and outputs should engage the common

representation in a given context. Moreover, different tasks must be performed in serial, as the effort to perform two different tasks in parallel will produce interference in the shared representation. Thus, capacity limits on multi-tasking arise precisely when different tasks share a representational substrate and control is needed to serialise the tasks and prevent interference. Conversely, if different input/output mappings share little structure, there is no learning benefit to using a common representation, and the system can as easily acquire separate, independent representations for each task. This avoids the need for seriality and control, since all mappings can be computed in parallel, but exerts a cost on learning since the representations in each pathway must be acquired separately. Thus the canonical distinction between controlled, serial processing on one hand, and automatic, parallel processing on the other, represent different optimisations of a trade-off between cost of learning on the one hand versus cost of requiring serial control on the other.

Through this lens, semantic cognition can be viewed as the ultimate multi-task learning problem. As argued in the Introduction, the discernment of conceptual structure requires sensitivity to information distributed across multiple representational modalities, and across different tasks and learning episodes. The benefit of such learning is precisely that emphasised by Musslick et al. (2021) for controlled systems: use of a common conceptual representation promotes broad generalisation across items, tasks and situations. Thus, semantic cognition is a key domain where cognitive control and serial processing are required—explaining, for instance, why it is difficult to simultaneously read a book and listen to a podcast, or understand what someone is saying while you are also speaking. This perspective suggests that tasks in other cognitive domains requiring controlled, serial processing may do so because they adopt a shared representation, in which case the delayed maturation of control may likewise benefit learning in those domains.

Finally, although semantic control may rely on differing neural substrates than domain-general control (Gao et al., 2021; Hodgson, Lambon Ralph, & Jackson, 2021; Jackson, 2021), it appears

to mature along a similar timeline. Growing sensitivity to instructional context in our literature

review falls within the critical period for executive function development from 3-7 years, and

parallels the development of related processes including inhibition of prior responses, the

continued development of frontal cortex and the myelination of frontal white matter tracts (A.

Diamond, 2002, 2013; Giedd et al., 1999; Klingberg, Vaidya, Gabrieli, Moseley, & Hedehus,

1999). Thus, the onset of controlled semantic cognition may mirror the maturation of executive

function more generally, sharing a similar timeline and recruiting the same developing neural

resources. Future work should thus consider whether delayed maturation of executive function

likewise benefits learning in other domains that also exploit shared representations.

References

Acosta-Cabronero, J., Patterson, K., Fryer, T. D., Hodges, J. R., Pengas, G., Williams, G., & Nestor, P. J. (2011). Atrophy, hypometabolism and white matter abnormalities in semantic dementia tell a coherent story. *Brain, 134*, 2025-2035.

Badre, D., Poldrack, R. A., Pare-Blagoev, E. J., Insler, R. Z., & Wagner, A. D. (2005). Dissociable controlled retrieval and generalized selection mechanisms in ventrolateral prefrontal cortex. *Neuron, 47*(6), 907-918. doi:10.1016/j.neuron.2005.07.023

Bauer, P. J., & Mandler, J. M. (1989). Taxonomies and triads: Conceptual organization in one- to two-year-olds. *Cognitive Psychology, 21*, 156-184.

Berger, C., & Aguerra, E. (2010). Dynamic categorization and slot-filler representaton in 4- and 6-year-old children. *International Journal of Psychology, 45*(2), 81-89.

Blanchet, N., Dunham, P. J., & Dunham, F. (2001). Differences in preschool children's conceptal strategies when thinking about animate entities and artifacts. *Developmental Psychology, 37*(6), 791-800.

Blaye, A., Bernard-Peyron, V., Paour, J. L., & Bonthoux, F. (2006). Categorical flexibility in children: distinguishing response flexibility from conceptual flexibility; the protracted development of taxonomic representation. *European Journal of Developmental Psychology, 3*, 163-188.

Blaye, A., & Jacques, S. (2009). Categorical flexibility in preschoolers: contributions of conceptual knowledge and executive control. *Developmental Science, 12*(6), 863-873.

Bunge, S. A., Dudukovic, N. M., Thomason, M. E., Chandan, J. V., & Gabrieli, J. D. E. (2002). Immature frontal lobe contributions to cognitive control in children: Evidence from fMRI. *Neuron, 33*(2), 301-311.

Casey, B. J., Tottenham, N., Liston, C., & Durston, S. (2005). Imaging the developing brain: what have we learned about cognitive development? *Trends in Cognitive Sciences, 9*, 104-110.

Chapman, L. J. (1967). Illusory correlation in observational report. *Journal of Verbal Learning and Verbal Behavior, 6*(1), 151-155.

Chrysikou, E. G., Novick, J. M., Trueswell, J. C., & Thompson-Schill, S. L. (2011). The other side of cognitive control: Can a lack of cognitive control benefit language and cognition? *Topics in Cognitive Science, 3*(2), 253-256.

Chugani, H. T., & Phelps, M. E. (1986). Maturational changes in cerebral function in infants determined by 18FDG positron emission tomography. *Science, 231*(4740), 840-843.

Church, J. A., Bunge, S. A., Petersen, S. E., & Schlaggar, B. L. (2017). Preparatory engagement of cognitive control networks increases late in childhood *Cerebral Cortex, 27*(3), 2139-2153.

Cogdell-Brooke, L. S., Sowden, P. T., Violante, I. R., & Thompson, H. E. (2020). A meta-analysis of functional magnetic resonance imaging studies of divergent thinking using activation likelihood estimation. *Human Brain Mapping, 41*, 5057-5077.

Cohen, J. D., Dunbar, K., & McClelland, J. L. (1990). On the control of automatic processes: A parallel distributed processing account of the Stroop effect. *Psychological Review, 97*(3), 332-361.

Crone, E. A., Donohue, S. E., Honomichl, R., Wendelken, C., & Bunge, S. A. (2006). Brain regions mediating flexible rule use during development. *The Journal of Neuroscience, 26*(43), 11239–11247.

Davidson, M. C., Amso, D., Anderson, L. C., & Diamond, A. (2006). Development of cognitive control and executive functions from 4 to 13 years: evidence from manipulations of memory, inhibition, and task switching. . *Neuropsychologia, 44*, 2037-2078.

Defeyter, M. A., Avons, S. E., & German, T. C. (2007). Developmental changes in information central to artifact representation: evidence from 'functional fluency' tasks. *Developmental Science, 10*(5), 538-546.

Denney, D. R., & Moulton, P. A. (1976). Conceptual preferences among preschool children. *Developmental Psychology, 12*, 509–513.

Denney, N. W. (1974). Evidence for developmental changes in categorization criteria for children and adults. *Human Development, 17*, 41-53.

Diamond, A. (2002). *Normal development of prefrontal cortex from birth to young adulthood: cognitive functions, anatomy, and biochemistry*. New York: Oxford University Press.

Diamond, A. (2013). Executive Functions. *Annual Review of Psychology, 64*, 135-168.

Diamond, A., & Doar, B. (1989). The performance of human infants on a measure of frontal cortex function, the delayed response task. *Developmental Psychobiology, 22*(3), 271-294.

Duncan, J. (2010). The multiple-demand (MD) system of the primate brain: mental programs for intelligent behaviour. *Trends in Cognitive Sciences, 14*(4), 172-179. doi:10.1016/j.tics.2010.01.004

Elston, G. N., Oga, T., & Fujita, I. (2009). Spinogenesis and pruning scales across functiona hierarchies. *The Journal of Neuroscience, 29*(10), 3271-3275.

Estes, Z., Golonka, S., & Jones, L. L. (2011). Thematic Thinking: The Apprehension and Consequences of Thematic Relations. In B. H. Ross (Ed.), *Psychology of Learning and Motivation: Advances in Research and Theory, Vol 54* (Vol. 54, pp. 249-294). San Diego: Elsevier Academic Press Inc.

Funnell, E., Hughes, D., & Woodcock, J. (2006). Age of acquisition for naming and knowing: A new hypothesis. *Quarterly Journal of Experimental Psychology, 59*(2), 268-295.

Fuster, J. M. (2015). *The prefrontal cortex* (5th ed.). London, England.: Academic Press Inc.

Gao, Z., Zheng, L., Chiou, R., Gouws, A., Krieger-Redwood, K., Wang, X., . . . Jefferies, E. (2021). Distinct and common neural coding of semantic and non-semantic control demands. *Neuroimage, 236*, 118230.

German, T. C., & Defeyter, M. A. (2000). Immunity to functional fixedness in young children. *Psychonomic Bulletin & Review, 7*(4), 707-712.

German, T. P., & Johnson, S. C. (2002). Function and the origins of the design stance. *Journal of Cognition and Development, 3*(3), 279-300.

Gershkoff-Stowe, L., & Rakison, D. H. (Eds.). (2005). *Building object categories in developmental time*. Sussex, UK: Psychology Press.

Giedd, J. N., Blumenthal, J., Jeffries, N. O., Castellanos, F. X., Liu, H., Zijdenbos, A., . . . Rapoport, J. L. (1999). Brain-development during chHdhood and adolescence: A longitudinal MRI study. *Nature Neuroscience, 2*, 861-863.

Greenfield, D. B., & Scott, M. S. (1986). Young children's preference for complementary pairs: Evidence against a shift to a taxonomic preference. *Developmental Psychology, 22*(1), 19-21.

Henderson, L. M., Clarke, P. J., & Snowling, M. J. (2011). Accessing and selecting word meaning in autism spectrum disorder. *The Journal of Child Psychology and Psychiatry, 52*(9), 964-973.

Hodgson, V. J., Lambon Ralph, M. A., & Jackson, R. L. (2021). Multiple dimensions underlying the functional organisation of the language network. *Neuroimage, 241*, 118444. doi:https://www.biorxiv.org/content/10.1101/2021.02.05.429902v2

Hu, H., Lerer, A., Peysakhovich, A., & Foerster, J. (2020). "Other-play" for zero-shot coordination. *arXiv*. doi:10.48550/arXiv.2003.02979

Huttenlocher, P. R., & Dabholkar, A. S. (1997). Regional differences in synaptogenesis in human cerebral cortex. *Journal of Comparative Neurology, 387*, 167-178.

IBM. (2017). IBM SPSS Statistics for Windows (Version 25.0). Armonk, NY.

Inhelder, B., & Piaget, J. (1958). *The growth of logical thinking: From childhood to adolescence* (A. P. S. Milgram, Trans.). New York: Basic Books.

Jackson, R. L. (2021). The neural correlates of semantic control revisited. *Neuroimage, 224*, 117444.

Jackson, R. L., Hoffman, P., Pobric, G., & Lambon Ralph, M. A. (2015). The nature and neural correlates of semantic association vs. conceptual similarity. *Cerebral Cortex, 25*(11), 4319-4333.

Jackson, R. L., Rogers, T. T., & Lambon Ralph, M. A. (2021). Reverse-engineering the cortical architecture for controlled semantic cognition. *Nature Human Behaviour, 5*, 774–786.

Jefferies, E. (2013). The neural basis of semantic cognition: Converging evidence from neuropsychology, neuroimaging and TMS. *Cortex, 49*(3), 611-625. doi:10.1016/j.cortex.2012.10.008

Jefferies, E., & Lambon Ralph, M. A. (2006). Semantic impairment in stroke aphasia versus semantic dementia: a case-series comparison. *Brain, 129*, 2132-2147.

Johnson, M., Schuster, M., Le, Q. V., Krikun, M., Wu, Y., Chen, Z., . . . Dean, J. (2017). Google's multilingual neural machine translation system: Enabling zero-shot translation. *Transactions of the Association for Computational Linguistics, 5*, 339-351.

Kazak, A. E. (2018). Editorial: Journal article reporting standards. *American Psychologist, 73*(1), 1-2.

Keil, F. C. (1979). *Semantic and conceptual development: An ontological perspective.* Cambridge, MA: Harvard University Press.

Klingberg, T., Vaidya, C. J., Gabrieli, J. D. E., Moseley, M. E., & Hedehus, M. (1999). Myelination and organization of the frontal white matter in children: A diffusion tensor MRI study. *Neuroreport, 10*(13), 2817-2821.

Lambon Ralph, M. A., Jefferies, E., Patterson, K., & Rogers, T. T. (2017). The neural and computational bases of semantic cognition. *Nature Reviews Neuroscience, 18*, 42-55.

Lambon Ralph, M. A., & Patterson, K. (2008). Generalization and differentiation in semantic memory - Insights from semantic dementia *Year in Cognitive Neuroscience 2008* (Vol. 1124, pp. 61-76). Oxford: Blackwell Publishing.

Landrigan, J., & Mirman, D. (2018). The cost of switching between taxonomic and thematic sentences. *Memory & Cognition, 46*(2), 191-203.

Lin, E. L., & Murphy, G. L. (2001). Thematic relations in adults' concepts. *Journal of Experimental Psychology-General, 130*(1), 3-28. doi:10.1037//0096-3445.130.1.3

Luna, B., Garver, K. E., Urban, T. A., Lazar, N. A., & Sweeney, J. A. (2004). Maturation of cognitive processes from late childhood to adulthood. *Child Development, 75*(5), 1357-1372.

Mandler, J. M., & McDonough, L. (1993). Concept formation in infancy. *Cognitive Development, 8*, 291-318.

Markman, E. M., & Hutchinson, J. E. (1984). Children's sensitivity to constraints on word meaning: Taxonomic versus thematic relations. *Cognitive Psychology, 16*, 1-27.

McRae, K., Cree, G. S., Seidenberg, M. S., & McNorgan, C. (2005). Semantic feature production norms for a large set of living and nonliving things. *Behavior Research Methods, 37*, 547-559.

Mervis, C. B. (1987). Child basic object categories and early lexical development. In U. Neisser (Ed.), *Concepts and conceptual development: Ecological and intellectual factors in categorization*. Cambridge, England: Cambridge University Press.

Miller, E. K., & Cohen, J. D. (2001). An integrative theory of prefrontal cortex function. *Annual Review of Neuroscience, 24*, 167-202.

Mirman, D. (2017). Taxonomic and thematic semantic systems. *Psychological Bulletin, 143*(5), 499-520.

Mrzljak, L., Uylings, H. B., van Eden, C. G., & Judas, M. (1990). Neuronal development in human prefrontal cortex in prenatal and postnatal stages. *Progress in Brain Research, 85*, 185-222.

Murphy, G. L., & Medin, D. L. (1985). The role of theories in conceptual coherence. *Psychological Review, 92*(3), 289-316.

Musslick, S., & Cohen, J. D. (2021). Rationalizing constraints on the capacity for cognitive control. *Trends in Cognitive Sciences, 25*(9), 757-775.

Newport, E. L. (1990). Maturational constraints on language learning. *Cognitive Science, 14*(1), 11-28.

Noonan, K. A., Jefferies, E., Visser, M., & Lambon Ralph, M. A. (2013). Going beyond inferior prefrontal involvement in semantic control: Evidence for the additional contribution of dorsal angular gyrus and posterior middle temporal cortex. *Journal of Cognitive Neuroscience, 25*(11), 1824-1850.

Patterson, K., Nestor, P. J., & Rogers, T. T. (2007). Where do you know what you know? The representation of semantic knowledge in the human brain. *Nature Reviews Neuroscience, 8*(12), 976-987.

Poulin-Dubois, D., & Pauen, S. (2017). The development of object categories: What, when, and how? In H. Cohen & C. Lefebvre (Eds.), *Handbook of Categorization in Cognitive Science* (pp. 653-671). Amsterdam: Elsevier Science.

Rogers, T. T., & McClelland, J. L. (2004). *Semantic cognition: A parallel distributed processing approach*. Cambridge, MA: MIT Press.

Rogers, T. T., & Patterson, K. (2007). Object categorization: Reversals and explanations of the basic-level advantage. *Journal of Experimental Psychology-General, 136*(3), 451-469. doi:10.1037/0096-3445.1363.451

Rohde, D. L. T. (1999). LENS: The light, efficient network simulator. *Technical Report CMU-CS-99-164, Carnegie Mellon University, Department of Computer Science, Pittsburgh, PA*.

Rosch, E., Mervis, C. B., Gray, W., Johnson, D., & Boyes-Braem, P. (1976). Basic objects in natural categories. *Cognitive Psychology, 8*, 382-439.

Sachs, O., Weis, S., Zellagui, N., Huber, W., Zvyagintsev, M., Mathiak, K., & Kircher, T. (2008). Automatic processing of semantic relations in fMRI: Neural activation during semantic priming of taxonomic and thematic categories. *Brain Research, 1218*, 194-205. doi:10.1016/j.brainres.2008.03.04S

Saffran, E. M. (2000). The organization of semantic memory: In support of a distributed model. *Brain and Language, 71*(1), 204-212.

Saxe, A. M., McClelland, J. L., & Ganguli, S. (2019). A mathematical theory of semantic development in deep neural networks. *Proceedings of the National Academy of Sciences of the United States of America, 116*(23), 11537-11546.

Schwartz, M. F., Kimberg, D. Y., Walker, G. M., Brecher, A., Faseyitan, O. K., Dell, G. S., . . . Coslett, H. B. (2011). Neuroanatomical dissociation for taxonomic and thematic knowledge in the human brain. *Proceedings of the National Academy of Sciences of the United States of America, 108*(20), 8520-8524. doi:10.1073/pnas.1014935108

Smaers, J. B., Gómez-Robles, A., Parks, A. N., & Sherwood, C. C. (2017). Exceptional Evolutionary Expansion of Prefrontal Cortex in Great Apes and Humans. *Current Biology, 27*(5), 714-720.

Smiley, S. S., & Brown, A. L. (1979). Conceptual preference for thematic or taxonomic relations: A nonmonotonic age trend from preschool to old age. *Journal of Experimental Child Psychology, 28*, 249-257.

Snyder, H. R., & Munakata, Y. (2010). Becoming self-directed: Abstract representations support endogenous Flexibility in children. *Cognition, 116*(2), 155–167.

Snyder, H. R., & Munakata, Y. (2013). So many options, so little control: Abstract representations can reduce selection demands to increase children's self-directed flexibility. *Journal of Experimental Child Psychology, 116*(3), 659–673.

Team, R. C. (2021). R: A language and environment for statistical computing. Vienna, Austria: R Foundation for Statistical Computing. Retrieved from https://www.R-project.org/

The MathWorks, I. (2018). MATLAB and Statistics Toolbox Release 2018a. Natick, Massachusetts, United States.

Thompson-Schill, S. L., D'Esposito, M., Aguirre, G. K., & Farah, M. J. (1997). Role of left inferior prefrontal cortex in retrieval of semantic knowledge: A reevaluation. *Proceedings of the National Academy of Sciences of the United States of America, 94*(26), 14792-14797.

Thompson-Schill, S. L., Ramscar, M., & Chrysikou, E. G. (2009). Cognition without control: When a little frontal lobe goes a long way. *Current Directions in Psychological Science, 18*(5), 259-263.

Veroude, K., Jolles, J., Croiset, G., & Krabbendam, L. (2013). Changes in neural mechanisms of cognitive control during the transition from late adolescence to young adulthood. *Developmental Cognitive Neuroscience, 5*, 63-70.

Wagner, A. D., Pare-Blagoev, E. J., Clark, J., & Poldrack, R. A. (2001). Recovering meaning: Left prefrontal cortex guides controlled semantic retrieval. *Neuron, 31*(2), 329-338. doi:10.1016/s0896-6273(01)00359-2

Waksom, M. L., Kumaran, D., Gordon, A. M., Rissman, J., & Wagner, A. D. (2014). Frontoparietal representations of task context support the flexible control of goal-directed cognition. *The Journal of Neuroscience, 34*(32), 10743-10755.

Warrington, E. K. (1975). Selective impairment of semantic memory. *Quarterly Journal of Experimental Psychology, 27*(NOV), 635-657. doi:10.1080/14640747508400525

Waxman, S. R., & Namy, L. L. (1997). Challenging the notion of a thematic preference in young children. *Developmental Psychology, 33*(3), 555-567.

Waxman, S. R., Senghas, A., & Benveniste, S. (1997). A cross-linguistic examination of the noun-category bias: Its existence and specificity in French- and Spanish-speaking preschool-aged children. *Cognitive Psychology, 32*, 183-218.

Wilk, H. A., & Morton, J. B. (2012). Developmental changes in patterns of brain activity associated with moment-to-moment adjustments in control. *Neuroimage, 63*, 475-484.

Wood, J. N., & Grafman, J. (2003). Human prefrontal cortex: processing and representational perspectives. *Nature Reviews Neuroscience, 4*, 139-147.

Zelazo, P. D., Carlson, S. M., & Kesek, A. (2008). *The development of executive function in childhood*. Cambridge, Mass.: MIT Press.