**Mapping whole-brain projections of anatomically-defined prefrontal neurons using combined 3D convolutional networks**

Michael W. Gongwer[1,2,3*], Cassandra B. Klune[1,2*], Alexander S. Enos[1], Rita Chen[1], Drew Friedmann[4] and Laura A. DeNardo[1,#]

Author Affiliations: [1]Department of Physiology, [2]Neuroscience Interdepartmental Program, [3]Medical Scientist Training Program, University of California Los Angeles, [4]Cajal Neuroscience, Seattle, WA

*Denotes equal contribution

#Correspondence: Dr. Laura DeNardo, 10833 Le Conte Ave, Center for Health Sciences, David Geffen School of Medicine at UCLA, Los Angeles, CA 90095 ldenardo@ucla.edu

**Abstract** Long-range axonal projections provide the foundation for functional connectivity between brain regions and are critical in the modulation of behavior. Descending projections from medial prefrontal cortex (mPFC) to various target regions regulate critical behavioral functions including decision making, social behavior and mood. While specific mPFC projections have distinct behavioral roles, individual mPFC projection neurons can also innervate multiple target regions. Yet how mPFC projection neurons divide their axons across the brain is poorly understood. In this study, we mapped the axon collaterals of mPFC neurons that project to nucleus accumbens (NAc), ventral tegmental area (VTA), or contralateral mPFC (cmPFC) in mice. We used tissue clearing and light sheet fluorescence microscopy to visualize the 3-D structure of axonal arbors across the intact brain. While machine learning can automate analysis of axons in images of cleared tissue, it is challenging to train a model that generalizes to all axonal structures because the appearance of axons varies by target region. In this study, we present DeepTraCE (Deep learning-based image Tracing with Combined-model Enhancement), a new strategy for axon segmentation and quantification in images of cleared

tissue. DeepTraCE is based on the deep-learning framework TRAILMAP; it achieves highly

accurate axon detection by combining multiple machine learning models that are each applied

to different brain regions. Using DeepTraCE, we find that cmPFC, NAc, and VTA-projecting

mPFC neurons represent largely separable classes with unique axon collaterals in cortical,

olfactory, and thalamic regions, respectively.

**Introduction** The medial prefrontal cortex (mPFC) is a vastly interconnected brain region that

controls complex cognitive and emotional functions including memory, decision making, social

interactions and mood[1–3]. mPFC acts through its axonal projections to exert top-down control,

biasing activity in downstream brain regions to promote adaptive responding to the

environment[4,5]. Recent studies have linked individual mPFC projections to discrete behavioral

functions[6–16]. On the other hand, many mPFC-dependent behaviors are known to involve

multiple downstream brain regions[17]. Taken together, these studies suggest that subtypes of

mPFC neurons may engage in specific behaviors by controlling small groups of downstream

regions. This functional organization may arise from classes of mPFC projection neurons that

send branched axon collaterals to specific sets of brain regions. Yet how mPFC neurons divide

up their axons across the brain is poorly understood (Figure 1A).

We mapped the brain-wide axonal collaterals of neurons in the prelimbic (PL) subregion

of mPFC that project to the contralateral PL (cPL), nucleus accumbens (NAc) or ventral

tegmental area (VTA). These projection classes are widely studied from a behavioral

perspective, but little is known about their connectivity at the whole-brain level. PL-cPL, PL-NAc

and PL-VTA neurons have distinct distributions across cortical layers, distinct genetic profiles,

and distinct functions[9,12,18]. PL-cPL connections are important for both hemispheric

synchronization and lateralization of mPFC activity; lateralized activity has been associated with

anxiety-like behavior[19] and stress responses in rodents[20–22], and psychiatric illness in humans[23].

PL-NAc neurons control reward seeking, social preference, threat avoidance, and strategy

shifting[9,13,24–26]. PL-VTA neurons respond to both positive and negative stimuli and are involved

in appetitive conditioning[15,27]. It is important to note that many of these studies used virally-

delivered neuronal activators (e.g. opsins[28] or DREADDS[29]) to determine the roles of projection-

defined mPFC neurons during behavior[7,9–13]. With this experimental design, activation of axon

collaterals can inadvertently influence animal behaviors. Our understanding of how PL-cPL, PL-

NAc and PL-VTA neurons control behavior is limited because we lack detailed maps of their

axon collaterals.

Traditional studies of axon collaterals have established an important foundation of

knowledge but have been limited by low throughput approaches. Anatomical mapping strategies

include injecting retrograde tracers into multiple target areas and measuring overlap in source

neurons[9,30]. Functional mapping strategies include recording from source neurons while

performing multisite antidromic stimulation in target regions[24,31]. While informative, these

approaches are biased – users predefine the target areas to investigate – and limited to small

numbers of regions. For most anatomical studies, researchers analyze connectivity manually in

2-dimensional (2-D) brain slices, a laborious process in which individual slices are difficult to

align to a reference brain. As a result, it is challenging to visualize the 3-D relationships of cells

across sections. RNA-sequencing based approaches[32] allow higher throughput examination of

collateralization, but the spatial resolution is limited by tissue micro-dissection volumes and

analyses are also limited to experimenter-defined sets of regions.

Tissue clearing methods[33,34] combined with light sheet fluorescence microscopy can

preserve 3-D circuit architecture in intact tissue without the need to align structures. Advances

in computer vision now allow researchers to train classifiers to automatically detect axons in

tissue volumes[35,36]. However, classifiers sometimes fail to identify all axons because axon

appearance varies across brain regions. Variance in axonal appearance is partly due to the

resolution limitations of most light sheet microscopes that are capable of visualizing entire

mouse brains in a timely fashion. Axons can appear thick and bright in sparsely innervated brain

regions near the surface of the brain, but thin and indistinct in densely innervated regions deep

in the tissue.

To overcome these hurdles, we developed DeepTraCE (Deep-learning based image

Tracing with Combined-model Enhancement), a novel strategy for axon segmentation in images

of cleared tissue. DeepTraCE builds upon a machine-learning based axon segmentation

pipeline called TRAILMAP[36]. After TRAILMAP trains several models to detect axons in 3-D,

DeepTraCE combines model outputs, applying different models to different brain regions

depending on the appearance of local axons. After segmenting the axons, DeepTraCE registers

the brains to a common coordinate framework and quantifies innervation density by brain

region.

In the present study, we used DeepTraCE in combination with viral circuit tracing, tissue

clearing, and light sheet microscopy to generated unbiased, whole-brain maps of PL-cPL, PL-

NAc and PL-VTA axon collaterals. First, we quantitatively compared projection patterns for the

three neuronal classes across all brain regions. While each class innervated dozens of brain

regions, we identified nearly 30 differentially innervated regions, most of which were located in

the cortex, subplate and thalamus. Next, we carefully examined each of those areas, identifying

striking topographic, layer-specific and region-specific innervation patterns across the three

classes. Finally, we performed unbiased analyses of the collateralization patterns, finding that

classes were largely separable based on their collateralization patterns. Overall, PL-cPL and

PL-VTA neurons were distinguished by their projections to cortex/subplate and

thalamus/hypothalamus/midbrain, respectively. PL-NAc neurons shared collateral targets with

both other classes, but also sent unique collaterals to groups of olfactory and limbic areas. Our

data support a model in which PL-cPL, PL-NAc and PL-VTA are separable neurons classes that

are defined by biased collateral outputs to groups of PL targets and that contain heterogeneity.

Indeed, we propose the existence of potential subclasses that can be investigated in future

studies.

**Results**

We hypothesized that classes of PL projection neuron collateralize to distinct subsets of target

regions (Figure 1A). To test this, we used a dual-virus approach to fluorescently label PL-cPL,

PL-NAc-, and PL-VTA neurons. In contrast to multi-site retrograde tracer injections, this

approach allows us to visualize their entire axonal arbors – including all the collaterals – in an

unbiased way.  Using light sheet imaging and machine learning analyses, we then quantified

innervation density across all brain regions and compared across neuronal types.

**Viral circuit mapping strategy**

We first injected an axon-terminal-transducing adeno-associated virus[37] expressing Cre

recombinase (AAVrg-hSyn-Cre) into cPL, NAc, or VTA. Next, into PL, we injected AAV

expressing Cre-dependent EYFP-tagged Channelrhodopsin-2 (AAV8-hSyn-DIO-ChR2-EYFP),

which traffics efficiently to axons (Figure 1B). We examined 2-D brain sections to determine the

location of retrogradely labeled cell bodies and get a broad overview of their projection patterns.

For each cell type, retrogradely labeled cell bodies had different but partially overlapping

locations across PL layers that aligned with previous literature[38]. PL-cPL neurons were

preferentially located in layer (L) 2–5a, PL-NAc neurons were distributed across L2–6, and PL-

VTA neurons were restricted to L5b (Figure S1A). Axon collaterals from PL-cPL neurons were

observed in cortex, striatum, and subplate regions. PL-NAc neurons collateralized broadly,

including to the cortex, thalamus, hypothalamus, amygdala, and claustrum. Collaterals of PL-

VTA neurons appeared in the thalamus and midbrain (Figure 1B). Overall, PL-cPL, PL-NAc and

PL-VTA neurons collateralize extensively to distinct but partially overlapping sets of brain

regions.

**DeepTraCE Workflow**

To map the brain-wide collateralization patterns in 3D, we used the same viral tracing strategy

and visualized axons in cleared intact brains. Using an iDISCO clearing variant, Adipo-Clear[39,40],

we immunostained intact brains for GFP and then rendered the tissue transparent. We then

imaged the hemispheres ipsilateral to the PL injection on a light sheet fluorescence microscope.

We used TRAILMAP[36], a machine learning pipeline that trains a 3D U-net framework to

automatically identify axons in cleared tissue (Figure 2A, Figure S2). TRAILMAP provides a pre-

trained model that users can fine tune to fit their samples. However, it was challenging to train a

model that could sufficiently generalize across the diversity of mPFC axonal structures in all of

their target regions; some axons were bright and delineated (e.g. in cortical areas) while others

were dense and indistinct (e.g. in the basolateral amygdala and thalamus).

To surmount this hurdle, DeepTraCE combines multiple trained models to optimize axon

detection throughout the brain. After importing 3-D image stacks, DeepTraCE fully segments

each brain using multiple models, registers each brain to a standard atlas, and generates a new

image in which pixel values in each region were extracted from the assigned model. Axons in

the resulting images are then thinned to single-pixel width using the TRAILMAP pipeline and the

number of axon-containing pixels are quantified for each brain region (Figures 2A).

**Validation of model combination method**

To determine how many models to combine, we examined raw images from the light sheet

microscope. Based on our visual inspection, we grouped brain regions into three main bins –

'delineated, 'marginally distinguished', or 'indistinct' – depending on the ability to identify the

structure of individual PL axons. We used TRAILMAP[36] to train three separate models that we

fine-tuned for each set of regions (Figures 2B,C; Figure S2A–C).

To validate the DeepTraCE model assignments, we performed quantitative comparisons

of model accuracy in representative brain regions from each category, using human annotations

as a reference. Human experts finely traced pixels containing axons in images with clearly

delineated axons and broadly estimated groups of pixels likely to contain axons in images with

marginally-distinguished and indistinct axons (Figures S2A–D). We segmented images using

each of the three models in addition to the original untuned TRAILMAP model that was trained

on serotonergic axons. We then calculated the resulting axonal labeling density with reference

to human labels. For reasons described below, common metrics such as F1 Score, precision,

and recall do not provide interpretable measures of segmentation accuracy with our method. To

validate our pipeline, we instead developed a distinction score based on labeling density in

pixels that humans labeled as axon-positive and axon-negative.

In light sheet microscopy of cleared tissue, the size of small fluorescent structures are

amplified such that a thin object such as an axon occupies several more pixels in the image

than it does in true biological space. As in TRAILMAP, we account for this overrepresentation by

thinning segmented axons to a single pixel width. However, because rapid imaging of intact

rodent brains requires lower optical resolution, it can be challenging to hand-label axons with

single pixel precision. This caveat to whole-brain light sheet microscopy approach prevents the

calculation of "true positive" and "true negative" rates based on human annotation. Instead, our

distinction score calculates the density of segmented axons in regions that humans estimated

as containing axons vs. not containing axons. A higher distinction score indicates better overlap

with human labels. When applied to our validation set, Models 1, 2, and 3 produced the highest

distinction scores in images containing delineated, marginally-distinguished and indistinct

axons, respectively (Figure S2F).

Previous studies used alternative approaches to overcome the limitations of a single-

model approach in whole-brain image segmentation. Some excluded deeper, more densely

innervated areas from analysis[41]. Others combined probability maps from different segmentation

methods by extracting the maximum probability for each pixel[42]. To compare DeepTraCE with

these alternative approaches, we calculated the distinction score across all human-annotated

images in the validation set using the original untuned TRAILMAP model (originally trained on

serotonin neurons), our best-trained single model, the maximum probability between our three

models, the sum of probabilities, and the DeepTraCE combined models. DeepTraCE had the

highest distinction score (Figure S2G), indicating that it performed the more accurate axon

segmentation from images.

**Whole-brain projection patterns of PL-cPL, PL-NAc and PL-VTA neurons**

We used DeepTraCE to determine the extent to which PL-cPL, PL-NAc and PL-VTA neurons

collateralize to other brain regions. We analyzed four brains from each class. After axon

segmentation from image volumes and registration to the brain atlas, DeepTraCE quantifies

axon innervation by brain region. Examination of thin optical sections revealed widespread

collateralization for all three classes. Overall, innervation patterns were consistent with reports

of general mPFC projection patterns[30,43,44]; mPFC axons were prominent in cortical association

areas, striatum, midline thalamus, claustrum, amygdala, hypothalamus and midbrain but largely

absent from hippocampus, sensory thalamus, and relatively sparse in primary sensory areas

(Supplemental Video). However, we observed striking differences in the innervation patterns of

the three projection classes, particularly in the cortex, olfactory areas, amygdala, thalamus and

midbrain (Figure 3A–C; Figure S3–5).

We generated dotograms[45] to visually summarize the inter-class distinctions and

observed regions with both quantitative and qualitative differences in innervation. To create the

dotogram, we averaged axon density across all samples for a given projection class and then

overlaid the samples in brain space. Dot color indicates the neuronal class and dot size

corresponds to the averaged axonal density within a voxel (Figure 3D). Compared to PL-VTA

neurons, PL-cPL and PL-NAc neurons more densely innervated cortical regions. On the other

hand, PL-VTA collaterals preferentially innervated thalamic (TH) and midbrain (MB) regions, a

subset of which also received collateral input from PL-NAc neurons. All three subtypes

innervated NAc and olfactory tubercle (OT), while PL-NAc neurons were the primary source of

collaterals to piriform cortex (PIR). These patterns were consistent with what we observed in 2D

brain sections (Figure 1B), indicating that DeepTraCE accurately captured meaningful class-

specific distinctions in collateral targets.

We next quantified regional innervation densities and plotted them as a heatmap. For

each brain, we normalized the number of axon-containing pixels in each region to the total

number of axon-containing pixels in the brain. We then sorted regions according to their

innervation density by projection classes; regions receiving more PL-cPL collaterals are on top

and those receiving more PL-VTA collaterals are in the bottom half of the heatmap (Figure 3E).

There was relatively little overlap between projection patterns of PL-cPL and PL-VTA neurons.

PL-NAc neurons, however, shared several projection targets with both other projection classes.

Brain-wide statistical comparisons confirmed the presence of 29 significantly differentially

innervated brain regions, the most notable of which were in the cortex, subplate, and thalamus

(Figure S6, Table 2, Table 3). We investigated these areas in more detail.

**Collateralization patterns in the cortex and subplate**

PL-cPL, PL-NAc and PL-VTA neurons all sent collaterals to agranular insula (AI), temporal

association areas (TEa), the parahippocampal region – including perirhinal (PERI), ectorhinal

(ECT) and entorhinal (ENT) cortices – and high order sensory areas (Figure 4A). Within these

regions, however, we observed interesting subclass-specific collateralization patterns.

Compared to PL-NAc and PL-VTA, PL-cPL neurons had the densest collaterals in the cortex.

PL-cPL axon collaterals were most prominent in AI, TEa, ECT, and PERI along with the

claustrum (CLA), which is part of the cortical subplate. Many regions were dually innervated by

PL-cPL and PL-NAc collaterals, suggesting PL-cPL and PL-NAc populations share common

subclasses. Indeed, gustatory (GU), visceral (VISC), ENTl, AIp, and high order visual (VISpl,

VISrl) and auditory areas (AUDd, AUDv) received equally dense collateral innervation from PL-

cPL and PL-NAc neurons, suggesting there may one or more classes of PL neurons that target

cPL, NAc, and GU, VISC, ENTl, AIp, and high order sensory areas. Compared to both PL-cPL

and PL-NAc, PL-VTA neurons sent few collaterals to the cortex, but with some exceptions. PL-

VTA collaterals were denser than the other classes in medial orbitofrontal cortex (ORBm). In

VISpm and AIv, the density of PL-VTA and PL-NAc collaterals was similar, but lower than PL-

cPL collateral density. ORBm, AIv, and VISpm may have privileged communication with PL-VTA

neurons (Figure 4A,B, Table 3).

Many PL target regions contain functional gradients. We therefore investigated whether

PL axons favor topographic locations in target regions and if there are distinctions between

projection classes. We plotted the anterior-posterior distribution of labeled axons within several

cortical areas and observed striking subclass-specific topographic trends. PL-NAc collaterals

were especially prominent in the posterior piriform cortex (PIR). Compared to PL-cPL PL-NAc,

PL-VTA collaterals had a tighter distribution in the anterior portion of AI (Figure 4C). Both PIR

and AI are both complex anatomical hubs with distinct functional zones along the A-P axis[46–48].

Preferential innervation by PL projection neurons may underlie some of the functional

specializations within these regions.

The neocortex is organized into layers that contain distinct neuronal types with different

morphology, physiology, and connectivity[38,49]. Long-range cortico-cortical axons may innervate

superficial or deep layers depending on the hierarchical relationship with the target region[38,49].

Compared to sensory and motor cortices, much less is known about the layer organization of

long-range mPFC connectivity, especially for specific projection classes. To investigate this, we

plotted PL-cPL, PL-NAc and PL-VTA collateral innervation density across layers in cortical

target areas (Fig. 4D). In ORB, all classes preferentially targeted superficial layers (L1, L2/3). In

AUD, TEa, ECT, and PERI, PL-cPL and PL-NAc collaterals preferentially targeted superficial

layers (L1, L2/3) while PL-VTA collaterals targeted the deep layers (L5, L6) (Figure 4D, Table

3). In ENT, axon collaterals were more prominent in deep layers for all 3 classes. These findings

suggest that PL-cPL, PL-NAc and PL-VTA may all participate in feedforward or feedback

connectivity depending on the target region. Further, while PL-cPL and PL-NAc collaterals had

similar layer-specific innervation patterns in all regions we quantified, PL-VTA collaterals had

opposing patterns in some of the regions, suggesting their role in intracortical communication is

distinct from the other two classes.

While all projection classes sent collaterals to the amygdalar complex, amygdalar nuclei

were differentially innervated. Most notably, the anterior basolateral amygdala (BLAa) was

robustly innervated by PL-cPL and PL-NAc neurons, but significantly less so by PL-VTA

neurons. Meanwhile, compared to PL-cPL neurons, PL-NAc neurons sent many more

collaterals to olfactory amygdalar areas, including anterior cortical amygdalar area (COAa) and

piriform amygdalar area (PAA) (Figure 5A–C). As PL-NAc collaterals were also more prominent

in posterior PIR compared to the other two classes, PL-NAc neurons may have privileged

control over areas dedicated to olfactory processing during cognition and emotional learning.

**Collateralization patterns in the thalamus**

PL projections to the thalamus play a key role in cognition, but whether and how classes of PL

neurons divide up their axons across thalamic nuclei remains poorly understood. Compared to

the other projection classes, PL-VTA axon collaterals were particularly dense in the thalamus

(Figure 1B, Fig. 3). PL-VTA collaterals were particularly notable in the medial (MED) and midline

thalamic nuclei (MTN) and the medial and lateral habenula (MHb, LHb). PL-NAc neurons also

densely collateralized in MHb, MED and MTN, suggesting PL-VTA and PL-NAc neurons may

share subclasses that project to VTA, NAc, MHb, MED and MTN. Interestingly, corticocortical

neurons are generally thought not to collateralize to the thalamus[38,49–51]. However, although

sparse, we observed PL-cPL collaterals in MED, ILM and MTN with varicosities reminiscent of

synaptic release sites (Figure 5D,E). These collaterals may be a unique feature of mPFC

neurons that can be further investigated with functional studies.

We also analyzed the topographic innervation of some of the most densely innervated

thalamic nuclei. PL axons favored anterior portions of MTN, posterior portions of LHb and were

relatively evenly distributed across the MED nuclei (Figure 5F). The topographic organization

was similar across subclasses, suggesting the location within PL, rather than projection class,

was the major determinant of axon location in thalamic regions.

**Unbiased assessment of brain-wide collaterals**

Because PL-cPL, PL-NAc and PL-VTA projection neurons all collateralize broadly, we

wondered if these neuronal classes could be separated based on their whole-brain projection

patterns. To unbiasedly assess differences between their whole-brain projection patterns, we

implemented two dimensionality reduction techniques. We first performed hierarchical

clustering, in which euclidean distance between individual brains based on axonal density in

each subregion is marked by a higher branch point on the graph. PL-VTA neurons formed a

distinct cluster, while PL-NAc and PL-cPL collaterals formed largely distinct but partially

overlapping clusters (Figure 6A). This aligns with our observation that the projection patterns of

PL-NAc neurons shared more overlap with PL-cPL than PL-VTA neurons (Fig. 3).

We used principal component analysis to assess differences in whole-brain projection

patterns for each cell type. We plotted the location of each brain along the axes of the first two

principal components, which account for 41.83% and 16.18% of the variance in the data,

respectively. We found that that PL-cPL and PL-VTA projection classes could be clearly

separated along the first principal component. PL-NAc brains were positioned in between, with

some overlap with PL-cPL brains (Figure 6B). To assess which distinguished the classes, we

plotted the weights of each brain region in contribution to the first principal component.

Positively weighted regions generally had the most innervation from PL-cPL collaterals including

TEa, PERI, and ECT, and negatively weighted regions were those with most innervation from

PL-VTA collaterals, including MTN and midbrain areas (Figure 6C). Together, these data

suggest that PL-cPL, PL-NAc and PL-VTA neurons represent mostly separable classes,

distinguished in large part by their collaterals to TEa/PERI/ECT, olfactory and limbic areas, and

thalamus and midbrain, respectively.

We next assessed the existence of potential connectivity motifs that represent groups of

regions that are highly co-innervated by PL projection neurons. Because PL-cPL, PL-NAc and

PL-VTA projection classes likely have some overlap, we performed this analysis on our entire

dataset (n=12 brains total, n=4 from each class). We calculated the Pearson R correlation

between axonal density for all pairs of brain regions. A higher R value represents increased

likelihood that two regions share PL collaterals. These R values were plotted on a heatmap and

sorted according to the Allen Brain Atlas (Figure 6D). Most cortical regions were highly

correlated with other cortical regions, parahippocampal regions, and subplate, but there were

some interesting exceptions. ORB cortex was highly correlated with olfactory areas including

taenia tecta (TT), dorsal peduncular area, and nucleus of the lateral olfactory tract (NLOT),

striatal regions including substantia innominata (SI) and magnocellular nucleus (MA). Since PL-

VTA neurons preferentially innervated ORBm (Figure 4), they are a likely source of these

shared collaterals. PIR was correlated with a distinct set of regions including COA, PAA, PPA,

and BLA. Given the preferential innervation of PL-NAc collaterals in these regions (Figure 4,5),

they are a likely source. These findings suggest the existence of two neuronal PL subclasses

with biased output connectivity: one that projects to VTA/ORB/TT/NLOT/SI/MA and another to

NAc/PIR/COA/PAA/PPA/BLA.

To further explore the existence of potential PL subclasses, we generated a network

diagram for strongly correlated region pairs (R>0.85 and p<0.05), with brain regions colored by

their weight in the first principal component (Figure 6E). This yielded two large networks of

prefrontal collateralization, along with small 'networks' of paired regions. One large network

comprised cortical and subplate regions that were densely innervated by PL-cPL and, to some

extent, PL-NAc collaterals. The other comprised regions in the thalamus, hypothalamus,

midbrain and hindbrain that are densely innervated by PL-VTA and, in some cases, by PL-NAc

collaterals. Each larger network contained one tightly interconnected subnetwork. Innervation

density was tightly correlated in a subset of PL-cPL collateral sites – ENTl, ECT, PERI, TEa,

AUDp, AUDv, VIS, VISrl, GU, AIp, MOp, SSs, CLA, and BLAa – suggesting these regions are

co-innervated by a subclass of PL-cPL neurons. The other subnetwork represents another likely

PL-VTA subclass that projects to five thalamic nuclei (RT, VENT, APN, SPF, TH) and the zona

incerta (ZI), a hypothalamic region (Figure 6E). Overall, our data support a biased connectivity

model for PL (Figure 1) in which PL-cPL, PL-NAc and PL-VTA are largely separable populations

with some overlap. Each class contains some unique subclasses defined by their projections to

small groups of downstream regions. PL-NAc appears to be the most heterogenous, with many

shared targets with the other classes.

**Discussion**

We defined the brain-wide collateral projections of PL-cPL, PL-NAc, and PL-VTA neurons.

These neurons have distinct, important behavioral functions, but their global connectivity

patterns remained unknown. Our data indicate that PL-cPL, PL-NAc, and PL-VTA neurons are

separable classes of projection neurons, potentially with distinct subclasses embedded within

them. PL-cPL neurons are distinguished by their collaterals to cortical association areas

including AI, TEa, PERI, ECT, CLA, ENT and high order visual and auditory areas. We also

identified unexpected PL-cPL collaterals in the thalamus. PL-VTA neurons collateralized to

specific regions in the thalamus –especially MTN, MED, and LHb – along with hypothalamus

and midbrain. PL-NAc neurons collateralized more broadly than the other classes, including to

many of the regions targeted by PL-cPL and PL-VTA neurons. While PL-NAc neurons shared

targets with both PL-cPL and PL-VTA neurons, prominent collaterals to posterior PIR, COA, and

PAA distinguished PL-NAc from the other classes (Figure 7A,B).

We quantified the brain wide collateralization patterns of these neurons using

DeepTraCE, a method we developed. DeepTraCE builds upon TRAILMAP by combining

multiple deep learning networks to accurately identify axons, which can differ in appearance

across brain regions. Variance in axonal appearance across the brain can arise due to

differences in axon morphology and density, due to differences in light penetration throughout

the tissue, and due to the resolution of the light sheet optics capable of imaging an intact mouse

brain in a few hours. By combining different models that are uniquely suited to axons with

distinct visual qualities, DeepTraCE allowed us to quantify the brain-wide axonal projection

patterns more accurately than previous methods (Figure S2D).

Our viral approach bulk labels many neurons at once, so the observed output

organization may represent the compound projections patterns of multiple subclasses within a

projection class. Indeed, recent single neuron reconstructions of prefrontal neurons indicate that

individual neurons project to approximately 5 brain regions[52]. Subclasses may be defined by

multiple factors including their layer of origin and transcriptional profile. PL-VTA and PL-cPL

were the most separable projection classes of the three and, consistent with previous studies,

we see that PL-VTA neurons reside exclusively in L5b,[53,30] while PL-cPL neurons reside mostly

in L2/3 and 5a[54,55] (Figure S1A). PL-NAc, which collateralized most broadly, spanned layers 2-6.

Consistent with this, cortico-striatal neurons in mPFC are more numerous than in other cortical

areas and are distributed more evenly across layers[38]. Layer of origin may therefore explain

some of the diversity in PL-NAc collaterals[44].

Transcriptomic heterogeneity has been observed within individual classes of prefrontal

projection neurons[18], suggesting they contain genetically-defined subpopulations. Cre driver

lines and transcriptome sequencing of projection-defined neurons further indicate that

genetically defined subclasses may exist that project to small groups of target regions[56]. For

example, Colgat2+ neurons in mPFC project to NAc, dorsal striatum (DS), ZI, and BLA[55]. Since

Colgat2+ axons do not innervate all targets of PL-NAc collaterals that we observed, Colgat2

may indicate a subset of PL-NAc neurons that collateralize to a specific subsets of PL-NAc

target regions. A recent sequencing study identified a set of genes including *Npr3, Otof, Syt6*

and *Tshz2* that are differentially expressed in classes mPFC projection neurons and may define

subclasses that project to combinations of BLA, DS, HY, NAc PAG and cPFC[18]. Our data

support a model in which PL projection classes are heterogeneous. Embedded subclasses may

influence behavior via simultaneous control of ~5–10 brain regions (Figure 7C).

PL-VTA neurons are distinguished in part by their dense collateral innervation of

thalamic nuclei including MED, MTN, SPA, PP, MHb and LHb. Separate studies found that

mPFC projections to VTA and thalamus both encode learned information about predictive cues

during adaptive decision making. While mPFC-VTA neurons can compare predictions with

outcomes encoded by VTA dopamine neurons[57], mPFC neurons projecting to the

paraventricular thalamus, part of MTN, can bias decision making towards predicting negative

outcomes[58]. Another PL-VTA collateral site, MD thalamus, also plays a key role in decision

making[59]. PL-VTA neurons reside in L5b. In contrast to L6 projections, which modulate thalamic

activity, L5 PL projections strongly activate thalamic neurons[53]. PL-VTA neurons may therefore

influence cognition by exerting powerful, simultaneous top-down control over a subset of

thalamic areas.

During social interactions, NAc, BLA, COA, TEA, PERI and ECT are activated in the

mouse brain[17]. We found that PL-NAc collaterals innervated all of these regions and previous

studies showed that PL-NAc and PL-BLA both directly regulate social behavior[9,60]. Therefore,

individual PL neurons may send divergent outputs to this set of regions to control social

behaviors.

fMRI studies in both humans and rodents indicate that prefrontal top-down control is

disrupted in numerous neuropsychiatric disorders[61–64]. For instance, disrupted prefrontal long-

range communication and decreased interhemispheric functional connectivity are core features

of pathophysiology in autism spectrum disorder (ASD)[65–67]. We find that PL-cPL and PL-NAc,

collateralize to several key regions disrupted in ASD including TEa, ECT, PERI, ENTl, BLA, AI,

and NAc[68]. Dysfunction in a subclass of prefrontal neurons collateralizing to these regions may

contribute to ASD pathogenesis.

Here, we present comprehensive projection maps of three PL projection classes as well

as a new analysis pipeline that allows more accurate axon segmentation from brain volumes.

However, it is important to note some limitations of our study. First, DeepTraCE is not

completely automated. It requires an initial time investment during which users assign machine

learning models to brain regions. Also, our viral labeling method does not distinguish between

axon terminals and axons of passage. Although we cannot claim that axons make direct

synaptic connections in target areas, many of the putative connections we observed have been

validated with electrophysiology[44,69,70] or rabies trans-synaptic tracing[47,71–74]. These caveats are

balanced by the ability to analyze the brain-wide projection patterns of defined neuronal classes

in a highly accurate, high throughput and unbiased way. Our data illuminate new organizational

properties of PL circuits. We propose that PL-cPL, PL-NAc and PL-VTA projection neurons

represent separable classes but are also heterogeneous populations. Each likely consists of

multiple neuronal subclasses that project to 5–10 brain regions, with PL-NAc being the most

heterogeneous of the three (Figure 7C). The many collaterals we observed are important

considerations for circuit dissection studies that use opto- or chemogenetics to activate

projection-defined neurons during behavior. Our observations can inform the design of new

studies that determine how mPFC controls behavior via multi-region influence.

**Methods**

**Animals**

Female and male C57B16/J mice (JAX Stock No. 000664) were group housed (2–5 per cage)

and kept on a 12 hr light cycle. All animal procedures followed animal care guidelines approved

by the University of California, Los Angeles Chancellor's Animal Research Committee.

**Surgery**

Mice were induced in 5% isoflurane in oxygen until loss of righting reflex and transferred to a stereotaxic apparatus where they were maintained under 2% isoflurane in oxygen. Mice were warmed with a circulating water heating pad throughout surgery and eye gel was applied to the animal's eyes. The mouse's head was shaved and prepped with three scrubs of alternating betadine and then 70% ethanol. Following a small skin incision, a dental drill was used to drill through the skulls above the injection targets. A syringe pump (Kopf, 693A) with a hamilton syringe was used for injections. Injections were delivered at a rate 75nL/min and the syringe was left in the brain for 7 minutes following injection. For collateralization mapping, 300uL of AAVrg-Ef1a-mCherry-IRES-Cre-WPRE (Addgene 55632-AAVrg, $1.7 \times 10^{13}$ vg/mL) was injected unilaterally into either left NAc (AP: 1.3, ML: -1.0, DV: 4.7), left VTA (AP: -3.3, ML: 0.4, DV: -4.5), or right PL (AP: 1.8, ML: -0.5, DV: -2.3). 200uL of AAV8-hSyn-DIO-hCHR2(H134R)-EYFP-WPRE ($1.7 \times 10^{13}$ vg/mL) was then injected into left PL (AP: 1.8, ML: -0.4, DV: -2.3). To control for Cre-independent expression of EYFP, control mice received a single injection of 200uL of AAV8-hSyn-DIO-hCHR2(H134R)-EYFP-WPRE into left PL. For topographical mapping of the origin of each projection type, mice were injected with 300uL of cholera toxin subunit B (ThermoFisher, C34775, C34776, C34778) of Fluorogold (SCBT, C223769) at the same coordinates for cPL, NAc and VTA listed above. For pain management mice received 5mg/kg carprofen diluted in 0.9% saline subcutaneously. Mice received one injection during surgery and daily injections for two days following surgery. Samples with mistargeted injection sites were excluded from analysis. Samples with obviously poor antibody penetration or distribution following tissue clearing were also excluded.

**Brain Slice Histology and Immunostaining** Mice were transcardially perfused with phosphate-buffered saline (PBS) followed by 4% paraformaldehyde (PFA) in PBS. Brains were

dissected, post-fixed in 4% PFA for 12–24h and placed in 30% sucrose for 24–48 hours. They

were then embedded in Optimum Cutting Temperature (OCT, Tissue Tek) and stored at -80°C

until sectioning. 60um floating sections were collected into PBS. Sections were washed

3x10min in PBS and then blocked in 0.3% PBST containing 10% normal donkey serum

(JacksonImmunoresearch, 17-000-121) for 2h. Sections were then stained with chicken anti-

GFP (AVES 1020 at 1:2000) in 0.3% PBST containing 3% donkey serum overnight at 4°C. The

following day, sections were washed 3x5min in PBS and then stained with secondary antibody

(JacksonImmunoresearch Cy2 donkey anti-chicken IgG(H+L) 703-225-155, 1:1000) in 0.3%

PBST containing 5% donkey serum for 2 hours at room temperature. Sections were then

washed 5 min with PBS, 15 min with PBS+DAPI (Thermofisher Scientific, D1306, 1:4000), and

then 5 min with PBS. Sections were mounted on glass slides using FluoroMount-G

(ThermoFisher, 00-4958-02) and then imaged at 10x with a Leica slide scanning microscope

(VT1200S).

**Brain Clearing** Mouse brains were collected and processed based on the published Adipo-

Clear protocol[39] with slight modifications. Mice were perfused intracardially with 20mL of PBS

(Gibco) followed by 4% paraformaldehyde (PFA, Electron Microscopy Sciences) on ice. Brains

were hemisected approximately 1mm past midline and postfixed overnight in 4% PFA at 4°C.

The following day, samples were dehydrated with a gradient of methanol (MeOH, Fisher

Scientific):B1n buffer (1:1,000 Triton X-100, 2% w/v glycine, 1:10,000 NaOH 10N, 0.02%

sodium azide) for 1 hour for each step (20%, 40%, 60%, 80%) on a nutator (VWR). Samples

were then washed with 100% MeOH 2x for 1hr each and then incubated in a 2:1

dicholoromethane (DCM):MeOH solution overnight. The following day, two washes of 1hr in

100% DCM were performed followed by three washes of 100% MeOH for 30min, 45min then

1hr. Samples were bleached for 4 hours in 5:1 $H_2O_2$/MeOH buffer. A cascade of MeOH/B1n

washes (80%, 60%, 40%, 20%) for 30min each rehydrated the samples followed by a 1h wash

in B1n buffer. 5%DMSO/0.3M Glycine/PTxWH permeabilized tissue for one hour and then again

for 2h with fresh solution. Samples were washed with PTxwH for 30min and then incubated in

fresh PTxwH overnight. The following day two more PTxwH washes lasted 1h then 2h. Samples

were incubated in primary GFP antibody (AVES Labs GFP 1020) at 1:2000 in PTxwH shaking

at 37°C for 11, washed in PTxwH 2x1h and then 2x2h, then for two days with at least one

PTxwH change per day while shaken at 37°C. Samples were then incubated in secondary

antibody (AlexaFluor 647, ThermoFisher Scientific) for 8 days shaken at 37°C. Samples were

washed in PTxwH 2x1h, then 2x2h, then 2 days with at least one PTxwH change per day while

shaken at 37°C. Samples were then washed in 1x PBS twice 1x1hr, 2x2hr and then overnight.

To dehydrate samples, a gradient of washes in MeOH:H2O (20%, 40%, 60% and 80%) were

conducted for 30min each, followed by 3x100% MeOH for 30min, 1h, then 1.5h. Samples were

incubated overnight in 2:1 DCM:MeOH on a nutator. The next day, samples were washed in

100% DCM 2x1h each. Samples were then cleared in 100% DBE. DBE was changed after 4h.

Samples were stored in DBE in a dark place at room temperature. Imaging took place at least

24h after clearing.

**Whole Brain Imaging** Brain samples were imaged on a light-sheet microscope

(Ultramicroscope II, LaVision Biotec) equipped with a sCMOS camera (Andor Neo) and a 2x/0.5

NA objective lens (MVPLAPO 2x) equipped with a 6 mm working distance dipping cap. Image

stacks were acquired at 0.8x optical zoom using Imspector Microscope v285 controller software.

We imaged using 488-nm (laser power 20%) and 640-nm (laser power 50%) lasers. The

samples were scanned with a step-size of 3 μm using the continuous light-sheet scanning

method with the included contrast adaptive algorithm for the 640-nm channel (20 acquisitions

per plane), and without horizontal scanning for the 488-nm channel.

**Model Training** Light sheet images of fluorescently labeled axons were segmented using the 3D U-net based machine learning pipeline TRAILMAP (Friedmann et al., 2020). We trained new models for segmentation of cortical axons using inference learning as described in the TRAILMAP pipeline (https://github.com/AlbertPun/TRAILMAP). Briefly, 120x120 pixel stacks of 100 images were selected for training, and axons and artifacts (pixels that could be mistakenly interpreted as axons) were hand-labeled in 3 or 4 images from each stack. 1-pixel-wide edges of axons were labeled using python to be given less weight in training of the 3D convolutional network, which accounts for slight variability in human annotation patterns.

Models 1, 2, and 3 were designed for axon segmentation in regions with delineated, marginally distinguished, and indistinct axons, respectively. Beginning with the published TRAILMAP model weights, trainings were iteratively performed. Model 2 was trained first for 7 sessions and represents our best-trained single model for use across the entire brain. From this, Model 1 for use in superficial regions was trained for 2 sessions and Model 3 was trained for 4 sessions for use in deep regions.

Each training session used between 6 and 20 hand-labeled image stacks, of which approximately 10-20% were set aside for use in model validation. Training was performed using between 20 and 120 steps per epoch and between 5 and 20 epochs per session. The best model from each session was selected by plotting loss in validation data across epoch, which was minimized to prevent overtraining. The weaknesses of the best model from each session were assessed to guide selection of training data for the next session. Sessions were continued until all visible axons in the desired regions were reliably detected by the model.

**Model Validation** We validated our analysis pipeline including image segmentation, scaling and axon thinning by selecting 120x120 pixel stacks of 100 images from regions assigned to the 'delineated', 'moderately distinguished', and 'indistinct' groups in brains that were not included in initial training or validation data sets. Two human experts annotated 2–4 images from each

stack, marking pixels likely to contain axons. In stacks with difficult-to-distinguish axons, such as

deep regions, experts used larger brush strokes to label broader groups of pixels likely to

contain axons (Figure S2D). One-pixel edges were then added to human annotations in python.

The raw image stacks were segmented as explained below. In brief, each stack was

segmented with Models 1, 2 and 3 using TRAILMAP and scaled in ImageJ to a 10um space.

Scaled images were skeletonized in python at 8 different thresholds for binarization, combined

in MATLAB, and pixel values were adjusted in ImageJ. Using the same quantification threshold

as used in all analyses (number of skeletonized pixels with intensity above 64 divided by total

number of pixels in the region), labeling density was calculated in human-annotated "axon-

positive" pixels (LP) and "axon-negative" pixels (LN), excluding edges to account for slight

variability in human labels. Distinction score was calculated from these values as (LP-

LN)/LN+(1-LP). While direct true/false positive and true/false negative rates cannot be

calculated due to the axon thinning process, which reduces the number of labeled pixels, LP

correlates with the true positive rate, LN correlates with the false positive rate, and 1-LP

correlates with the false negative rate. Thus, LP-LN in the numerator will increase distinction

score when better separation is obtained between true and false positives, and LN and 1-LP in

the denominator will decrease the distinction score if there is a higher rate of false positives or

false negatives.

Alternate model combination methods were performed in FIJI. Maximum probability

projections between all models were calculated by generating images with maximum pixel

values across all models for each cube then processing the cubes for validation as above.

Summation of probability maps between models was calculated by adding probability calculated

by each model, then dividing by 3 prior to processing the cubes for validation as above.

Distinction scores from alternate approaches were compared using repeated measures one-

way ANOVA with Dunnett's multiple comparisons test, comparing individual model selection

(DeepTraCE) with each other method. Statistical comparisons of distinguishment scores were

performed using GraphPad PRISM.

**DeepTraCE Analysis Pipeline**

*Image Segmentation & Registration*

Whole-brain image stacks from the 640nm channel from each brain were segmented

three times in TRAILMAP, once each using models 1, 2, and 3. Following segmentation, the

488nm autofluorescence channel and axon segmentations from each model were converted to

8-bit and scaled to 10um resolution in FIJI with scaling values of 0.40625 in the x and y

directions and 0.3 in the z direction using a bilinear interpolation algorithm. To improve image

registration, each scaled 488nm image was manually rotated in the x, y, and z planes using the

TransformJ ImageJ plugin (https://imagescience.org/meijering/software/transformj/) such that

the midline blood vessels visible were all visible in the same z plane. The same manual rotation

parameters were then applied to the three scaled model segmentation images from the

corresponding brain.

The scaled and manually rotated 488nm autofluorescence channel was registered using

elastix to the Gubra Lab LSFM atlas average template, which has annotations based on the

Allen CCF[75–77]. The same transformation was applied to the scaled and manually rotated model

segmentations using transformix. Image registration quality was manually verified by overlaying

the atlas image and the registered 488nm channels in ImageJ. Following segmentation and

registration of each model, the transformix images were converted to 8-bit .tif format in ImageJ.

When combining multiple probability maps, it is important that minimum and maximum pixel

values are comparable between images. Model 1 produced slightly lower maximum probability

values compared to the other two models. We corrected for this by brightening segmentations

from Model 1 by 34% to match pixel values obtained by the other models prior to combination

and thinning.

*Model Combination & Thinning*

Converted segmentations from the three models were then combined by generating a new image in which pixel values in each region were extracted from the model with best performance in that region (Figure 2B). The same regional model assignments for combination were used for all brains in the data set and are provided in Table 2. Following model combination, axon segmentations were thinned (or 'skeletonized') as described in the TRAILMAP pipeline[36]. In brief, images were binarized at 8 different thresholds from 20 to 90% of the maximum intensity value using python. Skeletons were combined in MATLAB by summing values from each skeleton. Small objects unlikely to be axons were removed by calculating connected components within the combined skeleton and removing objects less than 90 voxels in size. Combined skeletons with small objects removed were optimized for visualization and quantification using an ImageJ macro that multiplied each pixel value by 17.

*Axon Quantification*

Regional axon innervation was quantified in MATLAB (Mathworks) by counting the number of skeletonized pixels in each brain region above a threshold (64), then dividing this pixel count by the total number of pixels in a region. This regional pixel count was then divided by the total number of labeled pixels across the brain to normalize for differences in total fluorescence and viral expression. Regions were defined by a collapsed version of the LSFM atlas in which maximum granularity was balanced with the need to account for slight differences in registration which would lead to inaccurate quantification of small brain regions. This atlas was cropped on the anterior and posterior ends to match the amount of tissue visible in our data. Fiber tracts, ventricular systems, cerebellum, and olfactory bulb were excluded from analysis.

All statistical comparisons were performed in MATLAB and GraphPad Prism v9. For comparison of regional axon labeling between cell types, Two-way ANOVA with Tukey's

multiple comparison correction was performed on all brain regions. Anterior-posterior axon

distributions within regions were calculated in MATLAB by binning the whole-brain image into

100um voxels and calculating the percentage of segmented pixels within each voxel. Voxels

falling within a given region were summed across the medial-lateral and dorsal-ventral axis and

normalized for total fluorescence as above. The averaged summation of axon counts from a

given cell type was then averaged and plotted along with the standard error of the mean.

*Axon Visualization*

To visualize axons as shown in Figures 2, 4, and 5, Z-projections of raw light sheet data were

created in FIJI by scaling images to a 4.0625um space, virtually reslicing images in the coronal

plane, and performing maximum intensity z-projections of 100um depth followed by local

contrast enhancement. Axon segmentations were created by overlaying skeletonized and

registered axon segmentations from each sample of a cell type in a slightly different color and

virtually reslicing in the coronal plane. 3D projections were created in Imaris using a

representative sample from each cell type. Dotogram overlays (Figure 3) were created using

MATLAB. Images were binned into 100um voxels and the percentage of segmented pixels

within each voxel was calculated. Area of visible dot in the overlay corresponds with the

averaged labeling intensity within a voxel across a cell type. Outer dots represent the cell type

with the highest labeling intensity within that voxel.

*Computational Analyses*

Hierarchical clustering of regional axon quantifications was performed as described in Kebschull

et al., 2020[42]. Principal component analysis and network analyses were performed in MATLAB.

For network analysis, Pearson R correlations were calculated between quantified axons in each

brain region and each other brain region to generate a heatmap as shown in Figure 4D.

Samples from all labeled cell types were combined for this analysis, and regions with average

labeling density below 0.01 across all cell types were excluded from correlational analysis.

Regional correlations with a p value less than 0.05 and an R value greater than 0.85 were

plotted as a network diagram. Regions in this network diagram were color coded by their weight

within the first principal component from PCA.

**CODE AVAILABILITY**

Code, instructions and sample data available at https://github.com/DeNardoLab/DeepTraCE

**ACKNOWLEDGEMENTS**

We thank Justus Kebschull and Xiaoyin Chen for helpful comments on the manuscript. This

work was funded by an NIH K01MH116264 (L.A.D.), a Whitehall Foundation Research Grant

(L.A.D), a Klingenstein-Simons Foundation Grant (L.A.D.), a NARSAD Young Investigator

Award (L.A.D.), a Fay/Frank Seed Grant (L.A.D.), an NSERC Postgraduate Fellowship (C.B.K.),

and UCLA Medical Scientist Training Program (M.W.G.)

**AUTHOR CONTRIBUTIONS**

C.B.K, M.W.G. and L.A.D designed the research; C.B.K. performed the stereotaxic AAV

injections and histology; C.B.K. and M.W.G. performed imaging and data analysis; D.F. and

M.W.G. wrote DeepTraCE code; R.C. and A.S.E. performed image analysis, C.B.K, M.W.G. and

L.A.D. wrote the manuscript.

**DECLARATION OF INTERESTS**

None to declare.

**REFERENCES**

1.   Euston, D. R., Gruber, A. J. & McNaughton, B. L. The Role of Medial Prefrontal Cortex in

Memory and Decision Making. *Neuron* **76**, 1057–1070 (2012).

2.   Giustino, T. F. & Maren, S. The Role of the Medial Prefrontal Cortex in the Conditioning

and Extinction of Fear. *Front. Behav. Neurosci.* **9**, 1–20 (2015).

3.   Klune, C. B., Jin, B. & DeNardo, L. A. Linking mPFC circuit maturation to the

developmental regulation of emotional memory and cognitive flexibility. *Elife* **10**, 1–33

(2021).

4.   Matsuzaki, M. *et al.* Dendritic spine geometry is critical for AMPA receptor expression in

hippocampal CA1 pyramidal neurons. *Nat. Neurosci.* **4**, 1086–1092 (2001).

5.   Jayachandran, M. *et al.* Prefrontal Pathways Provide Top-Down Control of Memory for

Sequences of Events. *Cell Rep.* **28**, 640-654.e6 (2019).

6.   DeNardo, L. A., Berns, D. S., Deloach, K. & Luo, L. Connectivity of mouse

somatosensory and prefrontal cortex examined with trans-synaptic tracing. *Nat. Neurosci.*

**18**, 1687–1697 (2015).

7.   Otis, J. M. *et al.* Prefrontal cortex output circuits guide reward seeking through divergent

cue encoding. *Nature* **543**, 103–107 (2017).

8.   Siciliano, C. A. *et al.* A cortical-brainstem circuit predicts and governs compulsive alcohol

drinking. *Science.* (2019) doi:10.1126/science.aay1186.

9.   Murugan, M. *et al.* Combined Social and Spatial Coding in a Descending Projection from

the Prefrontal Cortex. *Cell* **171**, 1663-1677.e16 (2017).

10.  Vander Weele, C. M. *et al.* Dopamine enhances signal-to-noise ratio in cortical-brainstem

encoding of aversive stimuli. *Nature* **563**, 397–401 (2018).

11.    Warden, M. R. *et al.* A prefrontal cortex-brainstem neuronal projection that controls

687            response to behavioural challenge. *Nature* (2012) doi:10.1038/nature11617.

688    12.    Kim, C. K. *et al.* Molecular and Circuit-Dynamical Identification of Top-Down Neural

689            Mechanisms for Restraint of Reward Seeking. *Cell* (2017) doi:10.1016/j.cell.2017.07.020.

690    13.    Diehl, M. M. *et al.* Divergent projections of the prelimbic cortex bidirectionally regulate

691            active avoidance. *Elife* **9**, 1–13 (2020).

692    14.    Riga, D. *et al.* Optogenetic dissection of medial prefrontal cortex circuitry. *Frontiers in*

693            *Systems Neuroscience* (2014) doi:10.3389/fnsys.2014.00230.

694    15.    Ye, L. *et al.* Wiring and Molecular Features of Prefrontal Ensembles Representing

695            Distinct Experiences. *Cell* (2016) doi:10.1016/j.cell.2016.05.010.

696    16.    Mathis, V. P., Williams, M., Fillinger, C. & Kenny, P. J. Networks of habenula-projecting

697            cortical neurons regulate cocaine seeking. *Sci. Adv.* **7**, (2021).

698    17.    Kim, Y. *et al.* Mapping Social Behavior-Induced Brain Activation at Cellular Resolution in

699            the Mouse. *Cell Rep.* **10**, 292–305 (2015).

700    18.    Lui, J. H. *et al.* Differential encoding in prefrontal cortex projection neuron classes across

701            cognitive tasks. *Cell* **184**, 489-506.e26 (2021).

702    19.    Costa, N. S., Vicente, M. A., Cipriano, A. C., Miguel, T. T. & Nunes-de-Souza, R. L.

703            Functional lateralization of the medial prefrontal cortex in the modulation of anxiety in

704            mice: Left or right? *Neuropharmacology* **108**, 82–90 (2016).

705    20.    Stevenson, C. W., Halliday, D. M., Marsden, C. A. & Mason, R. Early life programming of

706            hemispheric lateralization and synchronization in the adult medial prefrontal cortex.

707            *Neuroscience* **155**, 852–863 (2008).

708    21.    Sullivan, R. M. & Gratton, A. Lateralized effects of medial prefrontal cortex lesions on

709           neuroendocrine and  autonomic stress responses in rats. *J. Neurosci.* **19**, 2834–2840

710           (1999).

711    22.    Lee, E. *et al.* Left brain cortical activity modulates stress effects on social behavior. *Sci.*

712           *Rep.* **5**, 13342 (2015).

713    23.    Grimm, S. *et al.* Imbalance between left and right dorsolateral prefrontal cortex in major

714           depression  is linked to negative emotional judgment: an fMRI study in severe major

715           depressive disorder. *Biol. Psychiatry* **63**, 369–376 (2008).

716    24.    Ishikawa, A., Ambroggi, F., Nicola, S. M. & Fields, H. L. Dorsomedial Prefrontal Cortex

717           Contribution to Behavioral and Nucleus Accumbens Neuronal Responses to Incentive

718           Cues. *J. Neurosci.* **28**, 5088 LP – 5098 (2008).

719    25.    Domingo-Rodriguez, L. *et al.* A specific prelimbic-nucleus accumbens pathway controls

720           resilience versus vulnerability to food addiction. *Nat. Commun.* **11**, 782 (2020).

721    26.    Cui, Q. *et al.* Dopamine receptors mediate strategy abandoning via modulation of a

722           specific prelimbic cortex–nucleus accumbens pathway in mice. *Proc. Natl. Acad. Sci.*

723           **115**, E4890 LP-E4899 (2018).

724    27.    Parker, J. G., Beutler, L. R. & Palmiter, R. D. The contribution of NMDA receptor

725           signaling in the corticobasal ganglia reward  network to appetitive Pavlovian learning. *J.*

726           *Neurosci.* **31**, 11362–11369 (2011).

727    28.    Zhang, F. *et al.* The microbial opsin family of optogenetic tools. *Cell* (2011)

728           doi:10.1016/j.cell.2011.12.004.

729    29.    Roth, B. L. DREADDs for Neuroscientists. *Neuron* (2016)

730           doi:10.1016/j.neuron.2016.01.040.

30.   Gabbott, P. L. A., Warner, T. A., Jays, P. R. L., Salway, P. & Busby, S. J. Prefrontal

Cortex in the Rat☐: Projections to Subcortical Autonomic , Motor , and Limbic Centers. *J.*

*Comp. Neurol.* **177**, 145–177 (2005).

31.   Ishikawa, A. & Nakamura, S. Ventral hippocampal neurons project axons simultaneously

to the medial prefrontal cortex and amygdala in the rat. *J. Neurophysiol.* **96**, 2134–2138

(2006).

32.   Kebschull, J. M. *et al.* High-Throughput Mapping of Single-Neuron Projections by

Sequencing of Barcoded RNA. *Neuron* **91**, 975–987 (2016).

33.   Chung, K. *et al.* Structural and molecular interrogation of intact biological systems. *Nature*

**497**, 332–337 (2013).

34.   Renier, N. *et al.* Mapping of Brain Activity by Automated Volume Analysis of Immediate

Early Genes. *Cell* **165**, 1789–1802 (2016).

35.   Berg, S. *et al.* ilastik: interactive machine learning for (bio)image analysis. *Nat. Methods*

**16**, 1226–1232 (2019).

36.   Friedmann, D. *et al.* Mapping mesoscale axonal projections in the mouse brain using a

3D convolutional network. *Proc. Natl. Acad. Sci.* (2020) doi:10.1073/pnas.1918465117.

37.   Tervo, D. G. R. *et al.* A Designer AAV Variant Permits Efficient Retrograde Access to

Projection Neurons. *Neuron* (2016) doi:10.1016/j.neuron.2016.09.021.

38.   Anastasiades, P. G. & Carter, A. G. Circuit organization of the rodent medial prefrontal

cortex. *Trends Neurosci.* **44**, 550–563 (2021).

39.   Chi, J., Crane, A., Wu, Z. & Cohen, P. Adipo-Clear: A Tissue Clearing Method for Three-

Dimensional Imaging of Adipose Tissue. *J. Vis. Exp.* (2018) doi:10.3791/58271.

40.    Chi, J. *et al.* Three-Dimensional Adipose Tissue Imaging Reveals Regional Variation in

Beige Fat Biogenesis and PRDM16-Dependent Sympathetic Neurite Density. *Cell Metab.*

**27**, 226-236.e3 (2018).

41.    Kramer, D. J. *et al.* Generation of a DAT-P2A-Flpo mouse line for intersectional genetic

targeting of dopamine neuron subpopulations. *Cell Rep.* **35**, 109123 (2021).

42.    Kebschull, J. M. *et al.* Cerebellar nuclei evolved by repeatedly duplicating a conserved

cell-type set. *Science.* **370**, (2020).

43.    DeNardo, L. A. *et al.* Temporal evolution of cortical ensembles promoting remote memory

retrieval. *Nat. Neurosci.* **22**, 460–469 (2019).

44.    Anastasiades, P. G., Boada, C. & Carter, A. G. Cell-Type-Specific D1 Dopamine

Receptor Modulation of Projection Neurons and Interneurons in the Prefrontal Cortex.

*Cereb. Cortex* **29**, 3224–3242 (2019).

45.    Oh, S. W. *et al.* A mesoscale connectome of the mouse brain. *Nature* **508**, 207–214

(2014).

46.    Ährlund-Richter, S. *et al.* A whole-brain atlas of monosynaptic input targeting four

different cell types in the medial prefrontal cortex of the mouse. *Nat. Neurosci.* **22**, 657–

668 (2019).

47.    Wang, L. *et al.* Cell-Type-Specific Whole-Brain Direct Inputs to the Anterior and Posterior

Piriform Cortex . *Frontiers in Neural Circuits* vol. 14 (2020).

48.    Poo, C., Agarwal, G., Bonacchi, N. & Mainen, Z. F. Spatial maps in piriform cortex during

olfactory navigation. *Nature* **601**, 595–599 (2022).

49.    Harris, K. D. & Shepherd, G. M. G. The neocortical circuit: Themes and variations. *Nat.*

*Neurosci.* **18**, 170–181 (2015).

50.    Shepherd, G. M. G. Corticostriatal connectivity and its role in disease. *Nat. Rev.*

777            *Neurosci.* **14**, 278–291 (2013).

778    51.    Shepherd, G. M. G. & Svoboda, K. Laminar and columnar organization of ascending

779            excitatory projections to layer 2/3 pyramidal neurons in rat barrel cortex. *J. Neurosci.* **25**,

780            5670–9 (2005).

781    52.    Winnubst, J. *et al.* Reconstruction of 1,000 Projection Neurons Reveals New Cell Types

782            and Organization  of Long-Range Connectivity in the Mouse Brain. *Cell* **179**, 268-281.e13

783            (2019).

784    53.    Collins, D. P., Anastasiades, P. G., Marlin, J. J. & Carter, A. G. Reciprocal Circuits

785            Linking the Prefrontal Cortex with Dorsal and Ventral Thalamic  Nuclei. *Neuron* **98**, 366-

786            379.e4 (2018).

787    54.    Anastasiades, P. G., Marlin, J. J. & Carter, A. G. Cell-Type Specificity of Callosally

788            Evoked Excitation and Feedforward Inhibition in the Prefrontal Cortex. *Cell Rep.* **22**, 679–

789            692 (2018).

790    55.    Nakayama, H., Ibañez-Tallon, I. & Heintz, N. Cell-Type-Specific Contributions of Medial

791            Prefrontal Neurons to Flexible Behaviors. *J. Neurosci.* **38**, 4490–4504 (2018).

792    56.    Gerfen, C. R., Paletzki, R. & Heintz, N. GENSAT BAC cre-recombinase driver lines to

793            study the functional organization of cerebral cortical and basal ganglia circuits. *Neuron*

794            **80**, 1368–1383 (2013).

795    57.    Park, J. & Moghaddam, B. Risk of punishment influences discrete and coordinated

796            encoding of reward-guided  actions by prefrontal cortex and VTA neurons. *Elife* **6**, (2017).

797    58.    Lucantonio, F. *et al.* Aversive stimuli bias corticothalamic responses to motivationally

798            significant cues. *Elife* **10**, (2021).

799    59.    Mitchell, A. S. The mediodorsal thalamus as a higher order thalamic relay nucleus

800           important for  learning and decision-making. *Neurosci. Biobehav. Rev.* **54**, 76–88 (2015).

801    60.    Huang, W. C., Zucca, A., Levy, J. & Page, D. T. Social Behavior Is Modulated by

802           Valence-Encoding mPFC-Amygdala Sub-circuitry. *Cell Rep.* **32**, 107899 (2020).

803    61.    Guadagno, A., Belliveau, C., Mechawar, N. & Walker, C.-D. Effects of Early Life Stress

804           on the Developing Basolateral Amygdala-Prefrontal Cortex Circuit: The Emerging Role of

805           Local Inhibition and Perineuronal Nets   . *Frontiers in Human Neuroscience*   vol. 15

806           (2021).

807    62.    Dong, M. *et al.* A failed top-down control from the prefrontal cortex to the amygdala in

808           generalized anxiety disorder: Evidence from resting-state fMRI with Granger causality

809           analysis. *Neurosci. Lett.* **707**, 134314 (2019).

810    63.    Clauss, J. A., Benningfield, M. M., Rao, U. & Blackford, J. U. Altered Prefrontal Cortex

811           Function Marks Heightened Anxiety Risk in Children. *J. Am. Acad. Child Adolesc.*

812           *Psychiatry* **55**, 809–816 (2016).

813    64.    Johnstone, T., van Reekum, C. M., Urry, H. L., Kalin, N. H. & Davidson, R. J. Failure to

814           Regulate: Counterproductive Recruitment of Top-Down Prefrontal-Subcortical Circuitry in

815           Major Depression. *J. Neurosci.* **27**, 8877 LP – 8884 (2007).

816    65.    Courchesne, E. & Pierce, K. Why the frontal cortex in autism might be talking only to

817           itself: local over-connectivity but long-distance disconnection. *Curr. Opin. Neurobiol.* **15**,

818           225–230 (2005).

819    66.    Li, Q. *et al.* Decreased interhemispheric functional connectivity rather than corpus

820           callosum volume as a potential biomarker for autism spectrum disorder. *Cortex* **119**, 258–

821           266 (2019).

822    67.    Yao, S., Becker, B. & Kendrick, K. M. Reduced Inter-hemispheric Resting State

823            Functional Connectivity and Its Association With Social Deficits in Autism  . *Frontiers in*

824            *Psychiatry*  vol. 12 (2021).

825    68.    Weston, C. S. E. Four Social Brain Regions, Their Dysfunctions, and Sequelae,

826            Extensively Explain Autism Spectrum Disorder Symptomatology. *Brain Sciences*  vol. 9

827            (2019).

828    69.    Collins, D. P., Anastasiades, P. G., Marlin, J. J. & Carter, A. G. Reciprocal Circuits

829            Linking the Prefrontal Cortex with Dorsal and Ventral Thalamic Nuclei. *Neuron* **98**, 366-

830            379.e4 (2018).

831    70.    McGarry, L. M. & Carter, A. G. Prefrontal Cortex Drives Distinct Projection Neurons in the

832            Basolateral Amygdala. *Cell Rep.* **21**, 1426–1433 (2017).

833    71.    Gehrlach, D. A. *et al.* A whole-brain connectivity map of mouse insular cortex. *Elife* **9**,

834            e55585 (2020).

835    72.    Beier, K. T. *et al.* Circuit Architecture of VTA Dopamine Neurons Revealed by Systematic

836            Input-Output Mapping. *Cell* **162**, 622–634 (2015).

837    73.    Ährlund-Richter, S. *et al.* A whole-brain atlas of monosynaptic input targeting four

838            different cell types in the medial prefrontal cortex of the mouse. *Nat. Neurosci.* (2019)

839            doi:10.1038/s41593-019-0354-y.

840    74.    Schwarz, L. A. *et al.* Viral-genetic tracing of the input-output organization of a central

841            noradrenaline circuit. *Nature* **524**, 88–92 (2015).

842    75.    Klein, S., Staring, M., Murphy, K., Viergever, M. A. & Pluim, J. P. W. Elastix: A toolbox for

843            intensity-based medical image registration. *IEEE Trans. Med. Imaging* (2010)

844            doi:10.1109/TMI.2009.2035616.

845    76.    Wang, Q. *et al.* The Allen Mouse Brain Common Coordinate Framework: A 3D Reference

846            Atlas. *Cell* **181**, 936-953.e20 (2020).

847    77.    Perens, J. *et al.* An Optimized Mouse Brain Atlas for Automated Mapping and

848            Quantification of Neuronal Activity Using iDISCO+ and Light Sheet Fluorescence

849            Microscopy. *Neuroinformatics* **19**, 433–446 (2021).

**Figure 1. Viral strategy for whole-brain collateralization mapping.**
(A) Potential mPFC efferent connectivity patterns.
(B) Schematic of stereotaxic injection strategy for labeling PL-cPL, PL-NAc, and PL-VTA neurons (top) and coronal brain sections showing distinct axonal collateralization patterns for each cell type (bottom). Scale bars, 200um. Abbreviations: CLA, claustrum; CP, caudate putamen; SSp, primary somatosensory cortex; TH, thalamus; LH, lateral hypothalamus; RE, nucleus reuniens; RSP, retrosplenial cortex, MEA, medial amygdala; COA, cortical amygdala; BLA, basolateral amygdala; ZI, zona incerta; LHb, lateral habenula.

**Figure 2. DeepTraCE workflow**
(A) Overview of DeepTraCE workflow
(B) Demonstration of segmentation using 3 different models followed by concatenation
(C) Overlay of raw data and axon segmentation using single model or DeepTraCE concatenated models
(D) DeepTraCE segmentation of superficial (left) and deep (right) brain structures for each cell type. Scale bars, 200um. Abbreviations: Isoctx, isocortex; OLF, olfactory areas; HPF, hippocampal formation; CTXsp, cortical subplate; CNU, cerebral nuclei; HY, hypothalamus; MB, midbrain; HB, hindbrain

**Figure 3. Visualization and quantification of brain-wide projection patterns of PL-cPL, PL-NAc and PL-VTA neurons.**
(A–C) Single 10um coronal optical sections of thinned axons registered to the standardized brain atlas. Each image represents 4 overlaid brains from each type.
(D) Dotogram overlay of 3 cell types. Innvervation by cPL-, NAc- and VTA-projecting PL neurons shown in yellow, blue and purple, respectively.
(E) Heatmap of relative labeling density (normalized to region volume and gross label content per brain) across 140 regions defined by the Allen Brain Atlas.
See supplemental figures 3-6 for related data.

**Figure 4. Region-specific collateralization patterns of PL-cPL, PL-NAc and PL-VTA neurons**

(A) Coronal view of 100um z-projections of raw 640nm (axons, white) and 488nm (autofluorescence, blue) channels from individual brains. Images show cell-type specific innervation patterns in anterior (left) and posterior (right) cortical areas.

(B) Quantification of labeled axons in 21 cortex and subplate regions (n=4/group).

(C) Visualizations of axonal innervation density along the anterior-posterior axis in select regions. PL-cPL, PL-NAc and PL-VTA neurons coded in yellow, blue and purple, respectively.

(D) Layer distributions of axonal innervation in select cortical target regions.

See Table 1,2 for abbreviations and detailed statistics. 2-way ANOVA with Tukey's post-hoc multiple comparison test. Descriptive statistics: mean ± S.E.M. *P<0.05, **P<0.01, ***P<0.001.

**Figure 5. Collateralization patterns of PL-cPL, PL-NAc and PL-VTA neurons in amygdala and thalamus**

(A) Raw images showing axons in the amygdalar complex.

(B) Quantification of labeled axon density in amygdala subregions (n=4/group).

(C) Quantification of axonal innervation density along the anterior-posterior axis of a given region. PL-cPL, PL-NAc and PL-VTA neurons coded in yellow, blue and purple, respectively.

(D) Raw images showing axons in thalamic nuclei.

(E) Quantification of labeled axons in thalamic nuclei (n=4/group).

(F) Quantification of axonal innervation density along the anterior- posterior axis of a given region. Arrows indicate varicosities reminiscent of synaptic release sites. PL-cPL, PL-NAc and PL-VTA neurons coded in yellow, blue and purple, respectively.

See Tables 1, 2 for abbreviations and detailed statistics. 2-way ANOVA with Tukey's post-hoc multiple comparison test. Descriptive statistics: mean ± S.E.M. *$P<0.05$, **$P<0.01$, ***$P<0.001$.

**Figure 6. Hierarchical clustering and PCA analysis reveals cell-type specific output organization**
(A) Dendrogram of hierarchical clustering of regional axon quantifications from 12 mice, colored by target region.
(B) Locations of individual mice projected in principal component (PC) space defined by the first two PCs (arbitrary PC units, cPL, *n*=4; NAc, *n*=4; VTA, *n*=4).
(C) Loadings for PC1 (arbitrary PC weight units)
(D) Heatmap of Pearson's R correlations between regional axon quantifications. PL-cPL, PL-NAc and PL-VTA neurons were all included in correlation calculation. Each square in the heatmap corresponds to the strength of correlation between two brain regions, with brighter values indicating stronger correlation. See Tables 1 for abbreviations.
(E) Network visualization of correlations from (D) with *P*<0.05 and R>0.85, colored by PC1 weights from (C). Default Cytoscape layout was used for visualization.

**Figure 7. Summary of Findings**

(A) Visualization of collateralization density in key targets of PL–cPL, PL–NAc, and PL–VTA neurons. Dot radius correlates with average normalized labeling density within a region.

(B) Schematic of whole-brain collateralization patterns of PL–cPL, PL–NAc, and PL–VTA neurons.

(C) Proposed model of prefrontal efferent connectivity patterns.

See Table 1 for abbreviations.

**Supplemental Figure 1. Validation of Cre-Dependent Virus and Injection Site Mapping. Related to Figure 1.**

(A) Layer distributions of prelimbic neurons retrogradely labeled via injections of CTB-594, CTB-647, and AAVrg-EGFP in cPL, NAc, and VTA, respectively.

(B) Representative images of target sites (cPL, NAc and VTA) visualized with Cholera Toxin Subunit B and Fluorogold.

(C) Representative Coronal Sections sections of brains injected with AAVrg-Cre and AAV-DIO-ChR2-EYPF (left) or AAV-DIO-ChR2-EFYP in the absence of Cre and quantification of fluorescence in PL.

(D) Raw fluorescence images from LSFM showing mPFC and quantifications of cell body numbers within ACC, PL and IL.

(E) Maps of starter cells in mPFC for each cell type

**Supplemental Figure S2. Validation of DeepTraCE Pipeline. Related to Figure 2.**

(A-C) Single 500x500um planes of raw axon data, human annotation, and skeletons from Models 1, 2, and 3 in images of delineated, marginally delineated, and indistinct axons. Blue stars indicate model fine-tuned for segmentation at corresponding tissue depth.

(D) Representative images of raw data, human annotation, and probability maps from each model for regions assigned to Models 1, 2, and 3. Blue stars indicate model selected for segmentation of this region in DeepTraCE.

(E) Equation for distinction score. Higher score indicates higher distinction between axon-positive and axon-negative region and lower rate of false positives and false negatives.

(F) Distinction score produced by models 1, 2, and 3 for images of delineated, marginally delineated, and indistinct axons. Repeated-measures one-way ANOVA (Delineated: F=8.503, p=0.0257; Marginal: F=7.753, p=0.0309; Indistinct: F=73.76, p=0.0001).

(G) Comparison of DeepTraCE with alternate segmentation and model combination strategies. Repeated measures one-way ANOVA with Dunnett's multiple comparisons test (F=14.46, p<0.0001). See Table 3 for detailed statistics.

Descriptive statistics: mean ± SEM. *P<0.05.

**Supplemental Figure S3. Whole-brain PL–cPL collateralization patterns. Related to Figure 3.**

Eleven fully annotated 10um coronal optical sections of skeletonized images registered to the standardized brain atlas. Each image represents 4 overlaid PL–cPL brains. Sections from Bregma values +1.6, +0.7, -2.0, -3.5, and -4.7 are the same as shown in Figure 3.

**Supplemental Figure S4. Whole-brain PL–NAc collateralization patterns. Related to Figure 3.**

Eleven fully annotated 10um coronal optical sections of skeletonized images registered to the standardized brain atlas. Each image represents 4 overlaid PL–NAc brains. Sections from Bregma values +1.6, +0.7, -2.0, -3.5, and -4.7 are the same as shown in Figure 3.

**Supplemental Figure S5. Whole-brain PL–VTA collateralization patterns. Related to Figure 3.**

Eleven fully annotated 10um coronal optical sections of skeletonized images registered to the standardized brain atlas. Each image represents 4 overlaid PL–VTA brains. Sections from Bregma values +1.6, +0.7, -2.0, -3.5, and -4.7 are the same as shown in Figure 2.

**Supplemental Figure S6. Whole-brain collateralization data and statistics ordered by Allen Brain Atlas ontology. Related to Figure 3. See also Table 2.**

(A) Heatmap from Figure 3 ordered by Allen Brain Atlas ontology.
(B) Heatmap of p values from 2-way ANOVA with Tukey's multiple comparison correction for all brain regions.
(C) Heatmap of uncorrected p values from unpaired t-tests between cPL and NAc, cPL and VTA, and NAc and VTA.

| | | | | |
|---|---|---|---|---|
| ACC | Anterior Cingulate Cortex | | mPFC | medial prefrontal cortex |
| AH | Anterior Hypothalamic Area | | MPN | Medial preoptic nucleus |
| AHN | Anterior Hypothalamic Nucleus | | MTN | Midline group of the dorsal thalamus |
| AId | Agranular Insula, dorsal part | | MY-mot | Medulla, motor related |
| AIp | Agranular Insula, posterior part | | MY-sen | Medulla, sensory related |
| AIv | Agranular Insula, ventral part | | NAc | Nucleus accumbens |
| ATN | Anterior Group of the dorsal thalamus | | NDB | Nucleus of the diagonal band |
| AUDd | Dorsal auditory area | | NLOT | Nucleus of the lateral olfactory tract |
| AUDp | Primary auditory area | | OLF | Olfactory tract |
| AUDv | Ventral auditory area | | ORBm | Orbital area, medial part |
| BLA | Basolateral amygdala | | OT | Olfactory tubercle |
| BLAa | Basolateral amygdala, anterior part | | P-mot | Pons, motor related |
| BLAv | Basolateral amygdala, ventral part | | P-Sat | Pons, behavioral state related |
| BNST | Bed nucleus of the stria terminalis | | P-Sen | Pons, sensory related |
| CLA | Claustrum | | PAA | Piriform-amygdalar area |
| cPL | Contralateral prelimbic area | | PAG | Periaqueductal grey |
| CNU | Cerebral nuclei | | PERI | Perirhinal area |
| COA | Cortical amygdalar area | | PH | Posterior hypothalamic nucleus |
| COAa | Cortical amygdalar area, anterior part | | PIR | Piriform area |
| COAp | Cortical amygdalar area, posterior part | | PL | Prelimbic area |
| CP | Caudate putamen | | POST | Postrhinal area |
| CTXsp | Cortical subplate | | PP | Peripeduncular nucleus |
| DP | Dorsal peduncular area | | PPN | Pedunculopontine nucleus |
| ECT | Ectorhinal cortex | | PRC | Precommisural nucleus |
| ENTl | Entorhinal area, lateral part | | PVHd | Paraventricular hypothalamic nucleus, descending division |
| ENTm | Entorhinal area, medial part | | PVR | Periventricular region |
| EP | Endopiriform nucleus | | RE | Nucleus of reuniens |
| GENd | Geniculate group, dorsal thalamus | | RSP | Retrosplenial cortex |
| GU | Gustatory areas | | RT | Reticular nucleus of the thalamus |
| HB | Hindbrain | | sAMY | Striatum-like amygdala |
| HPF | Hippocampal formation | | SCm | Superior colliculus, motor related |
| IC | Inferior colliculus | | SCs | Superior colliculus, senosry related |
| IGL | Intergeniculat leaflet of the lateral geniculate complex | | SNr | Substantia nigra, reticular part |
| IL | Infralimbic cortex | | SPA | Subparafascicular area |
| ILM | Intralaminar nucleus of the dorsal thalamus | | SPF | Subparafascicular nucleus |
| Isoctx | Isocortex | | SSp | Primary somatosensory area |
| LAT | Lateral geniculate nucleus | | SSs | Secondary somatosensory area |
| LGv | Ventral part of the lateral geniculate nucleus | | SubG | Subgeniculate nucleus |
| LH | Lateral hypothalamus | | TEa | Temporal association area |
| LHb | Lateral habenula | | TH | Thalamus |
| MA | Magnocellular Nucleus | | TR | Postpiriform transition area |
| MB | Midbrain | | VENT | Ventral group of the dorsal thalamus |
| MB-un | Midbrain-unlabeled | | VISC | Visceral area |
| MBO | Mammillary body | | VISp | Primary visual area |
| MeA | Medial amygdala | | VISpl | Posterolateral visual area |
| MED | Medial group of the dorsal thalamus | | VISpm | Posteromedial visual area |
| MHb | Medial habenula | | VISrl | Rostrolateral visual area |
| MOp | Primary motor area | | VTA | Ventral tegmental area |
| MOs | Secondary motor area | | ZI | Zona incerta |

**Table 1.** List of abbreviations

**Table 2.** Quantification of axon density by brain region. **Related to Figures 2,4,5.**

| Abbreviation | Allen Brain Atlas ID | Mean cPL | Mean NAc | Mean VTA | cPL Brain 1 | cPL Brain 2 | cPL Brain 3 | cPL Brain 4 | NAc Brain 1 | NAc Brain 2 | NAc Brain 3 | NAc Brain 4 | VTA Brain 1 | VTA Brain 2 | VTA Brain 3 | VTA Brain 4 | PC1 Weight | Corrected 2-way ANOVA: cPL vs NAc | Corrected 2-way ANOVA: cPL vs VTA | Corrected 2-way ANOVA: NAc vs VTA | Uncorrected p value: cPL vs NAc | Uncorrected p value: cPL vs VTA | Uncorrected p value: NAc vs VTA | Model Used |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| FRP | 'Frontal pole, cerebral cortex' | 0.01032 | 0.00481 | 0.00452 | 0.00550 | 0.01398 | 0.01145 | 0.01035 | 0.00472 | 0.00597 | 0.00237 | 0.00620 | 0.00811 | 0.00304 | 0.00172 | 0.00518 | 0.0750 | 0.0952 | 0.0972 | 0.9817 | 0.0322 | 0.0425 | 0.8619 | 1 |
| Mop | 'Primary motor area' | 0.00270 | 0.00183 | 0.00089 | 0.00226 | 0.00214 | 0.00258 | 0.00381 | 0.00126 | 0.00206 | 0.00156 | 0.00245 | 0.00067 | 0.00061 | 0.00105 | 0.00123 | 0.0229 | 0.2381 | 0.0266 | 0.0621 | 0.1111 | 0.0045 | 0.0208 | 1 |
| Mos | 'Secondary motor area' | 0.00556 | 0.00388 | 0.00251 | 0.00354 | 0.00541 | 0.00534 | 0.00795 | 0.00382 | 0.00534 | 0.00194 | 0.00442 | 0.00146 | 0.00268 | 0.00309 | 0.00281 | 0.0405 | 0.3783 | 0.0757 | 0.301 | 0.1967 | 0.0206 | 0.1393 | 1 |
| SSp | 'Primary somatosensory area' | 0.00254 | 0.00190 | 0.00076 | 0.00277 | 0.00102 | 0.00261 | 0.00375 | 0.00098 | 0.00168 | 0.00220 | 0.00275 | 0.00060 | 0.00027 | 0.00089 | 0.00089 | 0.0089 | 0.6433 | 0.0996 | 0.1073 | 0.3865 | 0.0210 | 0.0252 | 1 |
| SSs | 'Supplemental somatosensory area' | 0.00479 | 0.00343 | 0.00151 | 0.00300 | 0.00422 | 0.00544 | 0.00649 | 0.00192 | 0.00352 | 0.00397 | 0.00431 | 0.00062 | 0.00030 | 0.00039 | 0.00081 | 0.0540 | 0.3732 | 0.007 | 0.0209 | 0.1907 | 0.014 | 0.017 | 1 |
| GU | 'Gustatory areas' | 0.00683 | 0.00435 | 0.00079 | 0.00219 | 0.00935 | 0.00894 | 0.00685 | 0.00271 | 0.00613 | 0.00372 | 0.00483 | 0.00097 | 0.00012 | 0.00168 | 0.00038 | 0.0814 | 0.4279 | 0.0646 | 0.0225 | 0.2158 | 0.0113 | 0.0047 | 1 |
| VISC | 'Visceral area' | 0.00900 | 0.00577 | 0.00056 | 0.00418 | 0.01124 | 0.01248 | 0.00811 | 0.00324 | 0.00819 | 0.00579 | 0.00586 | 0.00072 | 0.00010 | 0.00055 | 0.00086 | 0.1073 | 0.3588 | 0.039 | 0.0265 | 0.1766 | 0.0039 | 0.0022 | 1 |
| AUDd | 'Dorsal auditory area' | 0.00916 | 0.00652 | 0.00162 | 0.00851 | 0.00647 | 0.00846 | 0.01319 | 0.00418 | 0.00415 | 0.00753 | 0.01021 | 0.00134 | 0.00109 | 0.00136 | 0.00269 | 0.0918 | 0.4494 | 0.0222 | 0.0818 | 0.2436 | 0.0022 | 0.0176 | 1 |
| AUDp | 'Primary auditory area' | 0.00639 | 0.00440 | 0.00098 | 0.00652 | 0.00552 | 0.00597 | 0.00753 | 0.00309 | 0.00302 | 0.00505 | 0.00644 | 0.00076 | 0.00029 | 0.00109 | 0.00177 | 0.0640 | 0.1875 | 0.0002 | 0.0412 | 0.0771 | 0.0001 | 0.0082 | 1 |
| AUDpo | 'Posterior auditory area' | 0.01004 | 0.00598 | 0.00236 | 0.01315 | 0.00414 | 0.01124 | 0.01164 | 0.00384 | 0.00275 | 0.00613 | 0.01120 | 0.00233 | 0.00119 | 0.00259 | 0.00332 | 0.0887 | 0.3652 | 0.0582 | 0.2769 | 0.1902 | 0.0097 | 0.1090 | 1 |
| AUDv | 'Ventral auditory area' | 0.00967 | 0.00685 | 0.00064 | 0.00631 | 0.01255 | 0.00966 | 0.01015 | 0.00582 | 0.00769 | 0.00742 | 0.00646 | 0.00072 | 0.00018 | 0.00062 | 0.00105 | 0.1091 | 0.2182 | 0.0111 | 0.0004 | 0.0828 | 0.0004 | 0.0000 | 1 |
| VISal | 'Anterolateral visual area' | 0.00698 | 0.00444 | 0.00280 | 0.01018 | 0.00179 | 0.00790 | 0.00806 | 0.00296 | 0.00280 | 0.00551 | 0.00649 | 0.00184 | 0.00164 | 0.00326 | 0.00445 | 0.0438 | 0.4822 | 0.1953 | 0.3828 | 0.2570 | 0.0724 | 0.1975 | 1 |
| VISam | 'Anteromedial visual area' | 0.01042 | 0.00439 | 0.00541 | 0.01345 | 0.00338 | 0.01371 | 0.01123 | 0.00234 | 0.00526 | 0.00283 | 0.00714 | 0.00097 | 0.00742 | 0.00462 | 0.00461 | 0.0523 | 0.172 | 0.2481 | 0.7329 | 0.0660 | 0.0954 | 0.4668 | 1 |
| VISl | 'Lateral visual area' | 0.00532 | 0.00276 | 0.00183 | 0.00720 | 0.00250 | 0.00539 | 0.00618 | 0.00231 | 0.00197 | 0.00359 | 0.00317 | 0.00161 | 0.00087 | 0.00231 | 0.00255 | 0.0363 | 0.1594 | 0.0711 | 0.2677 | 0.0555 | 0.0180 | 0.1324 | 1 |
| VISp | 'Primary visual area' | 0.00655 | 0.00255 | 0.00152 | 0.00887 | 0.00527 | 0.00681 | 0.00525 | 0.00160 | 0.00126 | 0.00361 | 0.00371 | 0.00106 | 0.00151 | 0.00171 | 0.00179 | 0.0515 | 0.0252 | 0.0174 | 0.3816 | 0.0097 | 0.0012 | 0.1751 | 1 |
| VISpl | 'Posterolateral visual area' | 0.01018 | 0.00620 | 0.00227 | 0.00974 | 0.01249 | 0.01301 | 0.00547 | 0.00570 | 0.00196 | 0.00632 | 0.01083 | 0.00163 | 0.00059 | 0.00504 | 0.00180 | 0.0976 | 0.3215 | 0.026 | 0.237 | 0.1640 | 0.0071 | 0.1043 | 1 |
| VISpm | 'posteromedial visual area' | 0.01562 | 0.00630 | 0.00488 | 0.01909 | 0.01145 | 0.01953 | 0.01241 | 0.00397 | 0.00621 | 0.00591 | 0.00909 | 0.00429 | 0.00563 | 0.00332 | 0.00628 | 0.1124 | 0.3652 | 0.0322 | 0.0243 | 0.536 | 0.0079 | 0.0030 | 2 |
| ACC | 'Anterior cingulate area' | 0.00889 | 0.00599 | 0.00649 | 0.00911 | 0.00832 | 0.00942 | 0.00870 | 0.00424 | 0.00662 | 0.00517 | 0.00794 | 0.00502 | 0.00827 | 0.00756 | 0.00512 | 0.0288 | 0.0676 | 0.1188 | 0.905 | 0.0142 | 0.0330 | 0.6832 | 1 |
| PL | 'Prelimbic area' | 0.00833 | 0.00361 | 0.00626 | 0.00983 | 0.00911 | 0.00500 | 0.00939 | 0.00386 | 0.00286 | 0.00276 | 0.00497 | 0.00538 | 0.00474 | 0.01011 | 0.00481 | 0.0248 | 0.2481 | 0.4899 | 0.2532 | 0.0087 | 0.2706 | 0.1059 | 2 |
| IL | 'Infralimbic area' | 0.00730 | 0.00156 | 0.00871 | 0.00833 | 0.00387 | 0.00315 | 0.01387 | 0.00083 | 0.00437 | 0.00073 | 0.00031 | 0.00462 | 0.00568 | 0.01835 | 0.00621 | -0.0088 | 0.1939 | 0.9366 | 0.2132 | 0.0728 | 0.7404 | 0.0775 | 1 |
| ORBl | 'Orbital area, lateral part' | 0.00185 | 0.00101 | 0.00220 | 0.00122 | 0.00213 | 0.00168 | 0.00236 | 0.00099 | 0.00088 | 0.00126 | 0.00092 | 0.00160 | 0.00156 | 0.00317 | 0.00248 | -0.0027 | 0.081 | 0.7356 | 0.1016 | 0.0206 | 0.4717 | 0.0236 | 1 |
| ORBm | 'Orbital area, medial part' | 0.00432 | 0.00196 | 0.00066 | 0.00086 | 0.00794 | 0.00508 | 0.00342 | 0.00176 | 0.00550 | 0.00018 | 0.00041 | 0.00792 | 0.00531 | 0.00958 | 0.00653 | -0.0257 | 0.4838 | 0.2851 | 0.2668 | 0.2311 | 0.1350 | 0.0128 | 1 |
| ORBvl | 'Orbital area, ventrolateral part' | 0.00152 | 0.00066 | 0.00234 | 0.00074 | 0.00237 | 0.00175 | 0.00121 | 0.00098 | 0.00141 | 0.00041 | 0.00086 | 0.00152 | 0.00154 | 0.00505 | 0.00126 | -0.0041 | 0.231 | 0.6965 | 0.2998 | 0.1115 | 0.4275 | 0.1276 | 1 |
| AId | 'Agranular insular area, dorsal part' | 0.00911 | 0.00647 | 0.00565 | 0.00561 | 0.01027 | 0.01129 | 0.00925 | 0.00657 | 0.00598 | 0.00692 | 0.00642 | 0.00661 | 0.00183 | 0.00714 | 0.00703 | 0.0490 | 0.2307 | 0.2082 | 0.8131 | 0.0800 | 0.1001 | 0.5490 | 1 |
| AIp | 'Agranular insular area, posterior part' | 0.01245 | 0.00844 | 0.00136 | 0.00639 | 0.01595 | 0.01851 | 0.00894 | 0.00798 | 0.00965 | 0.00818 | 0.00795 | 0.00144 | 0.00250 | 0.00121 | 0.00028 | 0.1394 | 0.4458 | 0.0579 | 0.0001 | 0.2141 | 0.0086 | 0.0000 | 1 |
| AIv | 'Agranular insular area, ventral part' | 0.01792 | 0.00801 | 0.00869 | 0.00974 | 0.01733 | 0.01863 | 0.01918 | 0.00803 | 0.01064 | 0.00506 | 0.00831 | 0.01009 | 0.00459 | 0.00738 | 0.01269 | 0.1059 | 0.0021 | 0.0195 | 0.9441 | 0.0003 | 0.0024 | 0.7573 | 1 |
| RSP | 'Retrosplenial area' | 0.01266 | 0.00652 | 0.00656 | 0.01183 | 0.01119 | 0.01794 | 0.00969 | 0.00442 | 0.00631 | 0.00630 | 0.00907 | 0.00556 | 0.00702 | 0.00622 | 0.00742 | 0.0702 | 0.0735 | 0.0812 | 0.9995 | 0.0243 | 0.0169 | 0.9769 | 1 |
| VISrl | 'Rostrolateral visual area' | 0.01076 | 0.00761 | 0.00122 | 0.01044 | 0.00882 | 0.01256 | 0.01123 | 0.00955 | 0.00592 | 0.00620 | 0.00879 | 0.00073 | 0.00115 | 0.00075 | 0.00223 | 0.1092 | 0.0893 | 0.0007 | 0.0069 | 0.0395 | 0.0000 | 0.0006 | 1 |
| TEa | 'Temporal association areas' | 0.01926 | 0.01368 | 0.00171 | 0.01369 | 0.01957 | 0.02109 | 0.02271 | 0.01250 | 0.01669 | 0.01120 | 0.01431 | 0.00205 | 0.00051 | 0.00222 | 0.00207 | 0.2168 | 0.1263 | 0.0048 | 0.0022 | 0.0509 | 0.0001 | 0.0001 | 1 |
| PERI | 'Perirhinal area' | 0.02741 | 0.02306 | 0.00318 | 0.02034 | 0.03022 | 0.03142 | 0.02765 | 0.02414 | 0.02394 | 0.01485 | 0.02934 | 0.00337 | 0.00139 | 0.00415 | 0.00378 | 0.3069 | 0.543 | 0.0032 | 0.0121 | 0.3086 | 0.0001 | 0.0006 | 1 |
| ECT | 'Ectorhinal area' | 0.03475 | 0.02265 | 0.00307 | 0.02323 | 0.03932 | 0.04729 | 0.03620 | 0.02182 | 0.02649 | 0.01640 | 0.02591 | 0.00337 | 0.00138 | 0.00435 | 0.00377 | 0.3937 | 0.1031 | 0.0074 | 0.0048 | 0.0393 | 0.0002 | 0.0002 | 1 |
| OLF | 'Olfactory areas' | 0.00543 | 0.00473 | 0.00065 | 0.00392 | 0.00521 | 0.00729 | 0.00528 | 0.00543 | 0.00630 | 0.00283 | 0.00435 | 0.00097 | 0.00022 | 0.00057 | 0.00084 | 0.0841 | 0.7817 | 0.9951 | 0.8488 | 0.5184 | 0.9282 | 0.5981 | 1 |
| AOB | 'Accessory olfactory bulb' | 0.00145 | 0.00150 | 0.00189 | 0.00261 | 0.00083 | 0.00182 | 0.00054 | 0.00141 | 0.00143 | 0.00134 | 0.00180 | 0.00132 | 0.00113 | 0.00338 | 0.00174 | -0.0048 | 0.9953 | 0.8076 | 0.7475 | 0.9370 | 0.5490 | 0.4784 | 1 |
| AON | 'Anterior olfactory nucleus' | 0.00507 | 0.00369 | 0.00619 | 0.00399 | 0.00493 | 0.00748 | 0.00389 | 0.00267 | 0.00625 | 0.00184 | 0.00399 | 0.00709 | 0.00177 | 0.01145 | 0.00444 | 0.0020 | 0.5561 | 0.8749 | 0.5615 | 0.3183 | 0.6351 | 0.3144 | 1 |
| TT | 'Taenia tecta' | 0.01730 | 0.01019 | 0.01122 | 0.01100 | 0.02028 | 0.02117 | 0.01675 | 0.00648 | 0.01989 | 0.00452 | 0.00987 | 0.01559 | 0.00646 | 0.01230 | 0.01054 | 0.0929 | 0.281 | 0.1877 | 0.9626 | 0.1352 | 0.0882 | 0.8004 | 1 |
| DP | 'Dorsal peduncular area' | 0.00404 | 0.00327 | 0.01098 | 0.00374 | 0.00450 | 0.00410 | 0.00382 | 0.00370 | 0.00606 | 0.00169 | 0.00162 | 0.01197 | 0.00211 | 0.02059 | 0.00924 | -0.0614 | 0.7655 | 0.3051 | 0.2557 | 0.4951 | 0.1194 | 0.0995 | 2 |
| PIR | 'Piriform area' | 0.00391 | 0.00785 | 0.00318 | 0.00270 | 0.00337 | 0.00574 | 0.00380 | 0.00782 | 0.00455 | 0.00889 | 0.01017 | 0.00261 | 0.00031 | 0.00545 | 0.00435 | 0.0196 | 0.0803 | 0.8479 | 0.0663 | 0.0279 | 0.5954 | 0.0294 | 2 |
| NLOT | 'Nucleus of the lateral olfactory tract' | 0.01308 | 0.01788 | 0.02135 | 0.00846 | 0.01715 | 0.01303 | 0.01365 | 0.01838 | 0.01471 | 0.01458 | 0.02387 | 0.02413 | 0.00929 | 0.0561 | 0.01582 | -0.0444 | 0.2801 | 0.4455 | 0.8478 | 0.1388 | 0.2213 | 0.5956 | 2 |
| COAa | 'Cortical amygdalar area, anterior part' | 0.00976 | 0.02289 | 0.02058 | 0.00732 | 0.01161 | 0.01186 | 0.00826 | 0.02531 | 0.02412 | 0.01576 | 0.02636 | 0.02079 | 0.00931 | 0.03451 | 0.01768 | -0.0757 | 0.0148 | 0.2449 | 0.9173 | 0.0027 | 0.0907 | 0.7031 | 2 |
| COAp | 'Cortical amygdalar area, posterior part' | 0.00503 | 0.00796 | 0.00482 | 0.00246 | 0.00561 | 0.00434 | 0.00770 | 0.00961 | 0.00625 | 0.00848 | 0.00751 | 0.00340 | 0.00350 | 0.00664 | 0.00544 | 0.0104 | 0.1549 | 0.9878 | 0.0687 | 0.0669 | 0.8854 | 0.0298 | 2 |
| PAA | 'Piriform-amygdalar area' | 0.00468 | 0.01235 | 0.00570 | 0.00236 | 0.00505 | 0.00452 | 0.00678 | 0.01221 | 0.01153 | 0.01111 | 0.01453 | 0.00945 | 0.00027 | 0.00786 | 0.00520 | 0.0121 | 0.0018 | 0.8922 | 0.0798 | 0.0007 | 0.6603 | 0.0213 | 2 |
| TR | 'Postpiriform transition area' | 0.00106 | 0.00379 | 0.00135 | 0.00049 | 0.00221 | 0.00088 | 0.00066 | 0.00292 | 0.00079 | 0.00542 | 0.00600 | 0.00079 | 0.00125 | 0.00280 | 0.00078 | -0.0016 | 0.2033 | 0.7021 | 0.3827 | 0.3827 | 0.0741 | 0.4350 | 2 |
| HPF | 'Hippocampal formation' | 0.00256 | 0.00205 | 0.00115 | 0.00310 | 0.00389 | 0.00198 | 0.00128 | 0.00178 | 0.00076 | 0.00294 | 0.00273 | 0.00125 | 0.00057 | 0.00280 | 0.00078 | 0.0147 | 0.7906 | 0.3253 | 0.6089 | 0.5287 | 0.1646 | 0.3598 | 2 |
| CA1 | 'Field CA1' | 0.00026 | 0.00039 | 0.00012 | 0.00042 | 0.00033 | 0.00006 | 0.00021 | 0.00017 | 0.00021 | 0.00082 | 0.00036 | 0.00017 | 0.00007 | 0.00017 | 0.00006 | 0.0012 | 0.7647 | 0.3444 | 0.3788 | 0.4916 | 0.1609 | 0.1647 | 2 |
| CA2 | 'Field CA2' | 0.00012 | 0.00011 | 0.00007 | 0.00019 | 0.00015 | 0.00002 | 0.00011 | 0.00003 | 0.00002 | 0.00016 | 0.00023 | 0.00013 | 0.00006 | 0.00007 | 0.00003 | 0.0003 | 0.9922 | 0.5502 | 0.7683 | 0.9263 | 0.3421 | 0.5105 | 2 |
| CA3 | 'Field CA3' | 0.00048 | 0.00052 | 0.00038 | 0.00066 | 0.00053 | 0.00047 | 0.00025 | 0.00036 | 0.00015 | 0.00081 | 0.00078 | 0.00088 | 0.00026 | 0.00034 | 0.00003 | 0.0014 | 0.9638 | 0.8742 | 0.8204 | 0.8056 | 0.6463 | 0.5767 | 2 |
| DG | 'Dentate gyrus' | 0.00054 | 0.00063 | 0.00028 | 0.00068 | 0.00089 | 0.00037 | 0.00021 | 0.00054 | 0.00026 | 0.00070 | 0.00101 | 0.00028 | 0.00028 | 0.00023 | 0.00023 | 0.0024 | 0.9122 | 0.5854 | 0.3604 | 0.7123 | 0.3273 | 0.1795 | 2 |
| ENTl | 'Entorhinal area, lateral part' | 0.01786 | 0.01652 | 0.00298 | 0.01434 | 0.02296 | 0.01590 | 0.01826 | 0.01573 | 0.01002 | 0.01589 | 0.02442 | 0.00223 | 0.00049 | 0.00434 | 0.00485 | 0.1888 | 0.923 | 0.003 | 0.0315 | 0.7149 | 0.0004 | 0.0050 | 1 |
| ENTm | 'Entorhinal area, medial part, dorsal zone' | 0.00444 | 0.00550 | 0.00109 | 0.00245 | 0.00550 | 0.00387 | 0.00593 | 0.00631 | 0.00308 | 0.00587 | 0.00675 | 0.00052 | 0.00016 | 0.00186 | 0.00184 | 0.0448 | 0.6447 | 0.036 | 0.0149 | 0.3907 | 0.0105 | 0.0033 | 1 |
| PAR | 'Parasubiculum' | 0.00360 | 0.00230 | 0.00057 | 0.00434 | 0.00602 | 0.00276 | 0.00128 | 0.00174 | 0.00020 | 0.00260 | 0.00466 | 0.00048 | 0.00034 | 0.00036 | 0.00111 | 0.0338 | 0.6366 | 0.1127 | 0.2953 | 0.3829 | 0.0265 | 0.1174 | 1 |
| POST | 'Postsubiculum' | 0.00526 | 0.00162 | 0.00101 | 0.00742 | 0.00633 | 0.00504 | 0.00225 | 0.00138 | 0.00056 | 0.00252 | 0.00130 | 0.00096 | 0.00148 | 0.00029 | 0.00148 | 0.0433 | 0.0828 | 0.058 | 0.4896 | 0.0224 | 0.0099 | 0.2659 | 1 |
| PRE | 'Presubiculum' | 0.00332 | 0.00244 | 0.00085 | 0.00496 | 0.00362 | 0.00122 | 0.00177 | 0.00221 | 0.00530 | 0.00086 | 0.00134 | 0.00032 | 0.00278 | 0.00118 | 0.00255 | 0.0278 | 0.7965 | 0.1255 | 0.3924 | 0.5353 | 0.0337 | 0.1804 | 1 |
| SUB | 'Subiculum' | 0.00191 | 0.00097 | 0.00030 | 0.00383 | 0.00191 | 0.00170 | 0.00034 | 0.00073 | 0.00265 | 0.00024 | 0.00086 | 0.00134 | 0.00026 | 0.00232 | 0.00024 | -0.0170 | 0.6095 | 0.216 | 0.5095 | 0.3579 | 0.0701 | 0.2605 | 1 |
| CTXsp | 'Cortical subplate' | 0.00645 | 0.00733 | 0.00430 | 0.00529 | 0.00467 | 0.00631 | 0.00954 | 0.00627 | 0.00576 | 0.00944 | 0.00786 | 0.00518 | 0.00232 | 0.00624 | 0.00347 | 0.0332 | 0.8028 | 0.3397 | 0.1008 | 0.5438 | 0.1733 | 0.0458 | 2 |
| CLA | 'Claustrum' | 0.02057 | 0.01108 | 0.00573 | 0.01686 | 0.02129 | 0.02119 | 0.02294 | 0.00904 | 0.01206 | 0.00999 | 0.01322 | 0.00765 | 0.00328 | 0.00865 | 0.00333 | 0.1849 | 0.0034 | 0.0006 | 0.0534 | 0.0011 | 0.0002 | 0.0202 | 2 |
| EP | 'Endopiriform nucleus' | 0.01060 | 0.00856 | 0.00404 | 0.00481 | 0.00830 | 0.01422 | 0.01505 | 0.00892 | 0.00751 | 0.00953 | 0.00830 | 0.00652 | 0.00056 | 0.00741 | 0.00066 | 0.0919 | 0.7182 | 0.1418 | 0.0636 | 0.4444 | 0.0543 | 0.0144 | 1 |
| LA | 'Lateral amygdalar nucleus' | 0.00465 | 0.00539 | 0.00119 | 0.00299 | 0.00298 | 0.00463 | 0.00799 | 0.00229 | 0.00483 | 0.00883 | 0.00559 | 0.00151 | 0.00085 | 0.00168 | 0.00071 | 0.0460 | 0.9121 | 0.1152 | 0.1004 | 0.6943 | 0.0284 | 0.0220 | 1 |
| BLAa | 'Basolateral amygdalar nucleus, anterior part' | 0.01635 | 0.01832 | 0.00516 | 0.01210 | 0.01462 | 0.01916 | 0.01953 | 0.01445 | 0.02162 | 0.01771 | 0.01948 | 0.00635 | 0.00491 | 0.00623 | 0.00316 | 0.1496 | 0.6986 | 0.0101 | 0.0023 | 0.4371 | 0.0012 | 0.0002 | 1 |
| BLAp | 'Basolateral amygdalar nucleus, posterior part' | 0.00540 | 0.00750 | 0.00335 | 0.00379 | 0.00745 | 0.00476 | 0.00561 | 0.00448 | 0.00739 | 0.00930 | 0.00883 | 0.00353 | 0.00182 | 0.00558 | 0.00246 | 0.0351 | 0.3332 | 0.2435 | 0.0553 | 0.1669 | 0.1192 | 0.0226 | 2 |
| BLAv | 'Basolateral amygdalar nucleus, ventral part' | 0.01065 | 0.01646 | 0.01265 | 0.00657 | 0.00923 | 0.01543 | 0.01277 | 0.01343 | 0.01600 | 0.02362 | 0.01502 | 0.02161 | 0.00292 | 0.02161 | 0.01106 | 0.0211 | 0.2346 | 0.8922 | 0.7073 | 0.1117 | 0.6612 | 0.4426 | 1 |
| BMAa | 'Basomedial amygdalar nucleus, anterior part' | 0.00816 | 0.01107 | 0.00131 | 0.00613 | 0.00862 | 0.00913 | 0.01239 | 0.01112 | 0.00547 | 0.01071 | 0.01071 | 0.00258 | 0.00058 | 0.00744 | 0.00837 | -0.0416 | 0.1435 | 0.1259 | 0.5264 | 0.0658 | 0.0472 | 0.2858 | 1 |
| BMAp | 'Basomedial amygdalar nucleus, posterior part' | 0.00906 | 0.00909 | 0.00746 | 0.00613 | 0.00862 | 0.00913 | 0.01239 | 0.01112 | 0.00547 | 0.01071 | 0.01071 | 0.00837 | 0.00085 | 0.00944 | 0.00922 | -0.0203 | 0.9999 | 0.6197 | 0.6111 | 0.9877 | 0.3695 | 0.3612 | 1 |
| PA | 'Posterior amygdalar nucleus' | 0.00341 | 0.00423 | 0.00281 | 0.00257 | 0.00252 | 0.00366 | 0.00490 | 0.00456 | 0.00205 | 0.00488 | 0.00544 | 0.00095 | 0.00285 | 0.00463 | 0.00485 | -0.0077 | 0.6753 | 0.8348 | 0.477 | 0.4148 | 0.5814 | 0.2619 | 1 |
| STR | 'Striatum' | 0.00418 | 0.00384 | 0.00478 | 0.00291 | 0.00416 | 0.00410 | 0.00554 | 0.00333 | 0.00570 | 0.00321 | 0.00313 | 0.00560 | 0.00390 | 0.00619 | 0.00343 | -0.0007 | 0.9136 | 0.7691 | 0.585 | 0.6993 | 0.5046 | 0.3412 | 2 |
| CP | 'Caudoputamen' | 0.00751 | 0.00468 | 0.00356 | 0.00565 | 0.00384 | 0.00568 | 0.01487 | 0.00196 | 0.00376 | 0.00918 | 0.00383 | 0.00330 | 0.00504 | 0.00261 | 0.00330 | 0.0441 | 0.629 | 0.3799 | 0.7873 | 0.3728 | 0.1715 | 0.5223 | 3 |
| STRv | 'Striatum ventral region' | 0.00222 | 0.00370 | 0.00421 | 0.00216 | 0.00152 | 0.00197 | 0.00325 | 0.00318 | 0.00354 | 0.00462 | 0.00346 | 0.00505 | 0.00256 | 0.00628 | 0.00294 | -0.0171 | 0.0527 | 0.21 | 0.8563 | 0.0226 | 0.0830 | 0.6076 | 3 |
| NAc | 'Nucleus accumbens' | 0.03172 | 0.01904 | 0.01628 | 0.02104 | 0.02803 | 0.02880 | 0.04760 | 0.01726 | 0.03056 | 0.01164 | 0.01670 | 0.01852 | 0.00857 | 0.02228 | 0.01577 | 0.2146 | 0.2737 | 0.1578 | 0.8488 | 0.1312 | 0.0604 | 0.5993 | 3 |
| OT | 'Olfactory tubercle' | 0.01534 | 0.01624 | 0.01574 | 0.00947 | 0.01682 | 0.02063 | 0.01443 | 0.01926 | 0.01738 | 0.01047 | 0.01786 | 0.02075 | 0.00659 | 0.02357 | 0.01205 | 0.0330 | 0.9528 | 0.9957 | 0.9928 | 0.7767 | 0.9325 | 0.9123 | 3 |
| LSx | 'Lateral septal complex' | 0.00467 | 0.00483 | 0.00858 | 0.00587 | 0.00367 | 0.00165 | 0.00749 | 0.00373 | 0.00672 | 0.00422 | 0.00465 | 0.00972 | 0.00526 | 0.00719 | 0.01214 | -0.0459 | 0.9932 | 0.1982 | 0.1655 | 0.9145 | 0.0943 | 0.0620 | 3 |
| sAMY | 'Striatum-like amygdalar nuclei' | 0.00788 | 0.01287 | 0.01276 | 0.00685 | 0.01106 | 0.00700 | 0.01020 | 0.01122 | 0.01304 | 0.01276 | 0.01447 | 0.01850 | 0.00835 | 0.01552 | 0.00868 | -0.0243 | 0.0522 | 0.4012 | 0.999 | 0.0182 | 0.1975 | 0.9690 | 3 |
| CEA | 'Central amygdalar nucleus' | 0.00197 | 0.00215 | 0.00389 | 0.00130 | 0.00157 | 0.00092 | 0.00409 | 0.00129 | 0.00243 | 0.00350 | 0.00137 | 0.00058 | 0.00250 | 0.00733 | 0.00216 | -0.0158 | 0.9835 | 0.4472 | 0.4431 | 0.8653 | 0.2394 | 0.2270 | 3 |
| MEA | 'Medial amygdalar nucleus' | 0.00868 | 0.01360 | 0.01262 | 0.00633 | 0.01050 | 0.00981 | 0.00808 | 0.01671 | 0.01585 | 0.01011 | 0.01172 | 0.00979 | 0.01287 | 0.01485 | 0.01298 | -0.0381 | 0.986 | 0.0699 | 0.8692 | 0.8374 | 0.0308 | 0.6283 | 3 |
| PAL | 'Pallidum' | 0.00643 | 0.00651 | 0.00757 | 0.00573 | 0.00617 | 0.00599 | 0.00785 | 0.00522 | 0.00925 | 0.00504 | 0.00654 | 0.00671 | 0.01072 | 0.00634 | 0.00650 | -0.0130 | 0.9972 | 0.6253 | 0.7526 | 0.9457 | 0.3649 | 0.4885 | 1 |
| GPe | 'Globus pallidus, external segment' | 0.00108 | 0.00103 | 0.00191 | 0.00116 | 0.00021 | 0.00027 | 0.00267 | 0.00083 | 0.00047 | 0.00255 | 0.00025 | 0.00136 | 0.00164 | 0.00094 | 0.00371 | -0.0142 | 0.9975 | 0.6005 | 0.5405 | 0.9490 | 0.3551 | 0.3092 | 3 |
| GPi | 'Globus pallidus, internal segment' | 0.00179 | 0.00205 | 0.00380 | 0.00232 | 0.00091 | 0.00059 | 0.00334 | 0.00130 | 0.00372 | 0.00238 | 0.00079 | 0.00351 | 0.00642 | 0.00119 | 0.00409 | -0.0302 | 0.9572 | 0.3259 | 0.4099 | 0.7857 | 0.1586 | 0.2122 | 3 |
| SI | 'Substantia innominata' | 0.00666 | 0.00669 | 0.00983 | 0.00458 | 0.00864 | 0.00638 | 0.00705 | 0.00519 | 0.01064 | 0.00459 | 0.00636 | 0.01099 | 0.00622 | 0.01407 | 0.00802 | -0.0206 | 0.9998 | 0.3209 | 0.3908 | 0.9851 | 0.1498 | 0.2040 | 1 |
| MA | 'Magnocellular nucleus' | 0.00934 | 0.00964 | 0.01713 | 0.00694 | 0.01143 | 0.00837 | 0.01061 | 0.00586 | 0.01698 | 0.00592 | 0.00979 | 0.01817 | 0.00789 | 0.02403 | 0.01845 | -0.0636 | 0.9937 | 0.1941 | 0.2645 | 0.9187 | 0.0684 | 0.1288 | 1 |
| MS | 'Medial septal nucleus' | 0.01031 | 0.01251 | 0.01892 | 0.01269 | 0.00671 | 0.01559 | 0.01471 | 0.01471 | 0.01134 | 0.01094 | 0.01307 | 0.02035 | 0.00879 | 0.02825 | 0.02238 | -0.0888 | 0.7149 | 0.1992 | 0.299 | 0.4451 | 0.0925 | 0.1232 | 3 |
| NDB | 'Diagonal band nucleus' | 0.02438 | 0.01392 | 0.01706 | 0.01910 | 0.02705 | 0.03107 | 0.02028 | 0.01546 | 0.02259 | 0.00727 | 0.01005 | 0.02375 | 0.01704 | 0.02149 | 0.00596 | 0.0895 | 0.119 | 0.3593 | 0.8217 | 0.0539 | 0.1830 | 0.5659 | 3 |
| TRS | 'Triangular nucleus of septum' | 0.00607 | 0.00625 | 0.01072 | 0.00810 | 0.00792 | 0.00610 | 0.00553 | 0.00307 | 0.00546 | 0.00820 | 0.00827 | 0.01588 | 0.00777 | 0.01075 | 0.01352 | -0.0564 | 0.9968 | 0.3617 | 0.3145 | 0.9422 | 0.1870 | 0.1479 | 3 |
| BST | 'Bed nuclei of the stria terminalis' | 0.00482 | 0.00590 | 0.00980 | 0.00419 | 0.00247 | 0.00267 | 0.00996 | 0.00456 | 0.01223 | 0.00434 | 0.00248 | 0.00518 | 0.01031 | 0.00937 | 0.01433 | -0.0609 | 0.9214 | 0.2096 | 0.4174 | 0.7110 | 0.1010 | 0.2227 | 3 |
| TH | 'Thalamus' | 0.00416 | 0.00595 | 0.00981 | 0.00589 | 0.00395 | 0.00315 | 0.00367 | 0.00661 | 0.00581 | 0.00725 | 0.00415 | 0.00667 | 0.01578 | 0.00532 | 0.01147 | -0.0840 | 0.1956 | 0.1862 | 0.3734 | 0.0934 | 0.0615 | 0.1708 | 3 |
| VENT | 'Ventral group of the thalamus' | 0.00135 | 0.00111 | 0.00403 | 0.00271 | 0.00071 | 0.00079 | 0.00159 | 0.00126 | 0.00111 | 0.00158 | 0.00047 | 0.00105 | 0.00037 | 0.01035 | 0.00484 | -0.0472 | 0.9067 | 0.5792 | 0.5254 | 0.6873 | 0.3192 | 0.2731 | 3 |
| SPF | 'Subparafascicular nucleus' | 0.00283 | 0.00297 | 0.00909 | 0.00384 | 0.00222 | 0.00241 | 0.00285 | 0.00522 | 0.00247 | 0.00373 | 0.00045 | 0.00024 | 0.02081 | 0.00267 | 0.01265 | -0.1037 | 0.991 | 0.4772 | 0.4953 | 0.9014 | 0.2361 | 0.2536 | 3 |
| SPA | 'Subparafascicular area' | 0.00574 | 0.00990 | 0.01948 | 0.00167 | 0.00213 | 0.01133 | 0.00782 | 0.02195 | 0.00855 | 0.00663 | 0.00248 | 0.01384 | 0.03372 | 0.01500 | 0.01536 | -0.1912 | 0.6841 | 0.2643 | 0.5108 | 0.4201 | 0.1081 | 0.2808 | 3 |
| PP | 'Peripeduncular nucleus' | 0.00696 | 0.00961 | 0.01347 | 0.01080 | 0.00767 | 0.00420 | 0.00517 | 0.00421 | 0.00385 | 0.01514 | 0.01523 | 0.01155 | 0.01349 | 0.01612 | 0.01273 | -0.0747 | 0.7505 | 0.0299 | 0.5444 | 0.4837 | 0.0103 | 0.2944 | 3 |
| GENd | 'Geniculate group, dorsal thalamus' | 0.00754 | 0.00406 | 0.00352 | 0.01348 | 0.00544 | 0.00432 | 0.00694 | 0.00024 | 0.00306 | 0.01096 | 0.00499 | 0.00325 | 0.00809 | 0.00136 | 0.00139 | -0.0249 | 0.5715 | 0.3381 | 0.9826 | 0.3301 | 0.1712 | 0.8626 | 3 |
| LAT | 'Lateral group of the dorsal thalamus' | 0.00148 | 0.00214 | 0.00282 | 0.00294 | 0.00131 | 0.00077 | 0.00090 | 0.00131 | 0.00126 | 0.00401 | 0.00200 | 0.00339 | 0.00438 | 0.00166 | 0.00184 | -0.0209 | 0.7088 | 0.3084 | 0.753 | 0.4468 | 0.1539 | 0.4894 | 3 |
| ATN | 'Anterior group of the dorsal thalamus' | 0.00131 | 0.00192 | 0.00338 | 0.00358 | 0.00049 | 0.00091 | 0.00071 | 0.00267 | 0.00242 | 0.00127 | 0.00261 | 0.00684 | 0.00127 | 0.00960 | 0.00361 | -0.0550 | 0.4581 | 0.2665 | 0.3576 | 0.2501 | 0.1004 | 0.1567 | 3 |
| MED | 'Medial group of the dorsal thalamus' | 0.00260 | 0.00765 | 0.01230 | 0.00358 | 0.00144 | 0.00160 | 0.00379 | 0.01861 | 0.01940 | 0.01736 | 0.00957 | 0.02262 | 0.00231 | 0.02262 | 0.02101 | -0.1269 | 0.0145 | 0.011 | 0.8869 | 0.0045 | 0.0253 | 0.0185 | 3 |
| MTN | 'Midline group of the dorsal thalamus' | 0.00703 | 0.01827 | 0.02518 | 0.00925 | 0.00693 | 0.00426 | 0.00769 | 0.01861 | 0.01940 | 0.01736 | 0.01769 | 0.02262 | 0.00231 | 0.02262 | 0.02101 | -0.264 | 0.0011 | 0.0023 | 0.8869 | 0.0001 | 0.0002 | 0.0185 | 3 |
| ILM | 'Intralaminar nuclei of the dorsal thalamus' | 0.00130 | 0.00339 | 0.00583 | 0.00155 | 0.00067 | 0.00047 | 0.00251 | 0.00403 | 0.00383 | 0.00411 | 0.00160 | 0.00327 | 0.00794 | 0.00261 | 0.00952 | -0.0629 | 0.078 | 0.1435 | 0.449 | 0.0332 | 0.0427 | 0.2259 | 3 |
| RT | 'Reticular nucleus of the thalamus' | 0.00068 | 0.00088 | 0.00293 | 0.00110 | 0.00043 | 0.00033 | 0.00084 | 0.00091 | 0.00063 | 0.00178 | 0.00019 | 0.00127 | 0.00703 | 0.00127 | 0.00259 | -0.0347 | 0.8598 | 0.373 | 0.4296 | 0.6127 | 0.1643 | 0.2070 | 3 |
| IGL | 'Intergeniculate leaflet of the lateral geniculate' | 0.00235 | 0.00204 | 0.00345 | 0.00634 | 0.00166 | 0.00085 | 0.00055 | 0.00070 | 0.00042 | 0.00539 | 0.00165 | 0.00713 | 0.00105 | 0.00303 | 0.00199 | -0.0326 | 0.983 | 0.8692 | 0.7762 | 0.8656 | 0.6276 | 0.5127 | 3 |
| LGv | 'Ventral part of the lateral geniculate complex' | 0.00121 | 0.00217 | 0.00472 | 0.00304 | 0.00098 | 0.00036 | 0.00046 | 0.00074 | 0.00030 | 0.00416 | 0.00349 | 0.00714 | 0.00468 | 0.00505 | 0.00199 | -0.0400 | 0.6992 | 0.0795 | 0.2565 | 0.4349 | 0.0290 | 0.1262 | 3 |
| SubG | 'Subgeniculate nucleus' | 0.00209 | 0.00259 | 0.00307 | 0.00563 | 0.00070 | 0.00144 | 0.00058 | 0.00168 | 0.00072 | 0.00465 | 0.00328 | 0.00197 | 0.00328 | 0.00427 | 0.00277 | -0.0231 | 0.9711 | 0.7496 | 0.9648 | 0.8244 | 0.4802 | 0.8061 | 3 |
| MHb | 'Medial habenula' | 0.00545 | 0.01275 | 0.01638 | 0.00659 | 0.00731 | 0.00412 | 0.00380 | 0.00799 | 0.00800 | 0.01713 | 0.01785 | 0.02449 | 0.01618 | 0.00520 | 0.01963 | -0.1293 | 0.1428 | 0.1435 | 0.7536 | 0.0447 | 0.0403 | 0.4893 | 3 |
| LHb | 'Lateral habenula' | 0.00186 | 0.00796 | 0.01700 | 0.00202 | 0.00209 | 0.00159 | 0.00172 | 0.00763 | 0.00132 | 0.01049 | 0.00340 | 0.02252 | 0.01277 | 0.00906 | 0.02637 | -0.1838 | 0.0681 | 0.0491 | 0.165 | 0.0103 | 0.0057 | 0.0627 | 3 |
| HY | 'Hypothalamus' | 0.00675 | 0.00902 | 0.01004 | 0.00727 | 0.00898 | 0.00605 | 0.00468 | 0.01100 | 0.01032 | 0.00692 | 0.00787 | 0.01024 | 0.01138 | 0.01106 | 0.00747 | -0.0376 | 0.2767 | 0.0919 | 0.7355 | 0.1384 | 0.0415 | 0.4707 | 2 |
| PVZ | 'Periventricular zone' | 0.00456 | 0.00829 | 0.00640 | 0.00492 | 0.00560 | 0.00484 | 0.00287 | 0.01253 | 0.00471 | 0.00599 | 0.00992 | 0.00637 | 0.00484 | 0.00819 | 0.00620 | -0.0162 | 0.2437 | 0.1861 | 0.6264 | 0.0958 | 0.0878 | 0.3639 | 1 |
| PVR | 'Periventricular region' | 0.00808 | 0.01182 | 0.01257 | 0.00811 | 0.00979 | 0.00933 | 0.00511 | 0.01623 | 0.01362 | 0.00877 | 0.00866 | 0.01219 | 0.01389 | 0.01507 | 0.00911 | -0.0504 | 0.2842 | 0.0834 | 0.9432 | 0.1322 | 0.0363 | 0.7548 | 1 |

| Abbrev | Name | | | | | | | | | | | | | | | | | | | | | | | | Group |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| AHN | 'Anterior hypothalamic nucleus' | 0.01526 | 0.01678 | 0.01698 | 0.02051 | 0.01631 | 0.01564 | 0.00859 | 0.02385 | 0.01946 | 0.01383 | 0.00996 | 0.02004 | 0.02256 | 0.01167 | 0.01362 | -0.0472 | 0.9228 | 0.8838 | 0.9987 | 0.7134 | 0.6484 | 0.9618 | 1 |
| MBO | 'Mammillary body' | 0.01285 | 0.01370 | 0.01591 | 0.01407 | 0.01684 | 0.01336 | 0.00712 | 0.02143 | 0.01526 | 0.01090 | 0.00722 | 0.01416 | 0.02398 | 0.01368 | 0.01182 | -0.0573 | 0.9708 | 0.664 | 0.8561 | 0.8240 | 0.4049 | 0.6097 | 1 |
| MPN | 'Medial preoptic nucleus' | 0.00810 | 0.01465 | 0.01282 | 0.00795 | 0.00949 | 0.00981 | 0.00515 | 0.02361 | 0.01476 | 0.01080 | 0.00941 | 0.00784 | 0.01875 | 0.01006 | 0.01464 | -0.0707 | 0.2505 | 0.2839 | 0.8947 | 0.0997 | 0.1251 | 0.6656 | 1 |
| PMd | 'Dorsal premammillary nucleus' | 0.01371 | 0.01921 | 0.01581 | 0.01097 | 0.01499 | 0.01536 | 0.01351 | 0.03642 | 0.02678 | 0.01017 | 0.00346 | 0.01256 | 0.02410 | 0.00599 | 0.02061 | -0.0582 | 0.7682 | 0.8747 | 0.9188 | 0.4972 | 0.6332 | 0.7060 | 1 |
| PMv | 'Ventral premammillary nucleus' | 0.01126 | 0.01673 | 0.01799 | 0.01082 | 0.02042 | 0.01211 | 0.00168 | 0.02832 | 0.01008 | 0.01269 | 0.01585 | 0.01970 | 0.00534 | 0.03259 | 0.01436 | -0.0532 | 0.6128 | 0.6172 | 0.9823 | 0.3637 | 0.3645 | 0.8627 | 1 |
| PVHd | 'Paraventricular hypothalamic nucleus, desce' | 0.00418 | 0.00998 | 0.01369 | 0.00478 | 0.00479 | 0.00238 | 0.00479 | 0.01523 | 0.01259 | 0.00907 | 0.00392 | 0.01227 | 0.01963 | 0.00692 | 0.01595 | -0.1307 | 0.1708 | 0.0732 | 0.6285 | 0.0539 | 0.0141 | 0.3762 | 2 |
| VMH | 'Ventromedial hypothalamic nucleus' | 0.00460 | 0.00998 | 0.00574 | 0.00613 | 0.00752 | 0.00345 | 0.00130 | 0.01399 | 0.00717 | 0.00736 | 0.01142 | 0.00475 | 0.00417 | 0.00969 | 0.00437 | -0.0067 | 0.1052 | 0.8264 | 0.1961 | 0.0471 | 0.5731 | 0.0922 | 2 |
| PH | 'Posterior hypothalamic nucleus' | 0.00500 | 0.01043 | 0.01144 | 0.00503 | 0.00454 | 0.00382 | 0.00663 | 0.01648 | 0.01287 | 0.00754 | 0.00483 | 0.00868 | 0.01832 | 0.00312 | 0.01563 | -0.0969 | 0.2428 | 0.2911 | 0.9706 | 0.0897 | 0.1147 | 0.8233 | 2 |
| LH | 'Hypothalamic lateral zone' | 0.00797 | 0.01147 | 0.01460 | 0.00632 | 0.01044 | 0.00854 | 0.00658 | 0.01216 | 0.01728 | 0.00866 | 0.00779 | 0.01711 | 0.01645 | 0.01558 | 0.00926 | -0.0693 | 0.3853 | 0.0567 | 0.5426 | 0.1886 | 0.0177 | 0.3079 | 2 |
| ZI | 'Zona incerta' | 0.00442 | 0.00514 | 0.01197 | 0.00849 | 0.00294 | 0.00291 | 0.00334 | 0.00327 | 0.00760 | 0.00666 | 0.00305 | 0.00480 | 0.02182 | 0.00818 | 0.01309 | -0.1141 | 0.9148 | 0.2533 | 0.3021 | 0.7002 | 0.1038 | 0.1286 | 3 |
| ME | 'Median eminence' | 0.00089 | 0.00240 | 0.00298 | 0.00310 | 0.00034 | 0.00012 | 0.00001 | 0.00325 | 0.00073 | 0.00397 | 0.00166 | 0.00056 | 0.00008 | 0.00853 | 0.00273 | -0.0226 | 0.3781 | 0.6146 | 0.9593 | 0.1979 | 0.3539 | 0.7915 | 1 |
| MB-un | 'Midbrain' | 0.00797 | 0.00832 | 0.01271 | 0.00985 | 0.00902 | 0.00627 | 0.00673 | 0.00708 | 0.00735 | 0.00990 | 0.00895 | 0.01248 | 0.01516 | 0.01078 | 0.01242 | -0.0632 | 0.9452 | 0.0216 | 0.0218 | 0.7605 | 0.0092 | 0.0080 | 3 |
| MB-sen | 'Midbrain, sensory related' | 0.01225 | 0.00951 | 0.01401 | 0.01252 | 0.01330 | 0.01256 | 0.01063 | 0.00446 | 0.00466 | 0.01453 | 0.01441 | 0.01822 | 0.01526 | 0.01391 | 0.00863 | -0.0156 | 0.6546 | 0.7039 | 0.4576 | 0.3842 | 0.4322 | 0.2458 | 3 |
| SCs | 'Superior colliculus, sensory related' | 0.00293 | 0.00343 | 0.00481 | 0.00443 | 0.00249 | 0.00427 | 0.00055 | 0.00243 | 0.00106 | 0.00553 | 0.00471 | 0.00659 | 0.00675 | 0.00348 | 0.00243 | -0.0255 | 0.9309 | 0.4372 | 0.649 | 0.7284 | 0.2357 | 0.3948 | 3 |
| IC | 'Inferior colliculus' | 0.00744 | 0.00504 | 0.00784 | 0.01098 | 0.01000 | 0.00300 | 0.00577 | 0.00363 | 0.00390 | 0.00671 | 0.00592 | 0.01000 | 0.00710 | 0.00655 | 0.00772 | -0.0139 | 0.5171 | 0.978 | 0.0878 | 0.2781 | 0.8468 | 0.0399 | 3 |
| MB-mot | 'Midbrain, motor related' | 0.00277 | 0.00322 | 0.00518 | 0.00520 | 0.00209 | 0.00197 | 0.00179 | 0.00197 | 0.00203 | 0.00647 | 0.00242 | 0.00546 | 0.00730 | 0.00387 | 0.00408 | -0.0378 | 0.9394 | 0.1643 | 0.3797 | 0.7487 | 0.0776 | 0.1958 | 3 |
| SNr | 'Substantia nigra, reticular part' | 0.00330 | 0.00406 | 0.00703 | 0.00419 | 0.00206 | 0.00292 | 0.00403 | 0.00446 | 0.00452 | 0.00193 | 0.00959 | 0.00854 | 0.00589 | 0.00409 | | -0.0465 | 0.6898 | 0.1059 | 0.1988 | 0.4285 | 0.0325 | 0.0865 | 2 |
| VTA | 'Ventral tegmental area' | 0.01588 | 0.02282 | 0.02446 | 0.02214 | 0.01372 | 0.01406 | 0.01359 | 0.02389 | 0.02954 | 0.02026 | 0.01757 | 0.02620 | 0.02905 | 0.02007 | 0.02252 | -0.1166 | 0.1765 | 0.0559 | 0.8721 | 0.0823 | 0.0247 | 0.6323 | 3 |
| SCm | 'Superior colliculus, motor related' | 0.00683 | 0.00580 | 0.01010 | 0.01162 | 0.00759 | 0.00419 | 0.00390 | 0.00319 | 0.00311 | 0.01020 | 0.00670 | 0.00900 | 0.01185 | 0.00816 | 0.01139 | -0.0551 | 0.9109 | 0.3287 | 0.1631 | 0.6928 | 0.1553 | 0.0658 | 3 |
| PAG | 'Periaqueductal gray' | 0.00646 | 0.00484 | 0.00763 | 0.01116 | 0.00518 | 0.00674 | 0.00276 | 0.00032 | 0.01464 | 0.00240 | 0.00183 | 0.01068 | 0.00188 | 0.01613 | | -0.0501 | 0.9036 | 0.9529 | 0.8358 | 0.6795 | 0.7763 | 0.5833 | 3 |
| PRC | 'Precommissural nucleus' | 0.00987 | 0.01800 | 0.02155 | 0.01151 | 0.00489 | 0.01678 | 0.00632 | 0.01295 | 0.02932 | 0.01685 | 0.01289 | 0.02043 | 0.02699 | 0.02293 | 0.01584 | -0.1312 | 0.2823 | 0.0403 | 0.7293 | 0.1369 | 0.0171 | 0.4634 | 3 |
| APN | 'Anterior pretectal nucleus' | 0.00044 | 0.00071 | 0.00223 | 0.00129 | 0.00016 | 0.00014 | 0.00018 | 0.00020 | 0.00018 | 0.00183 | 0.00063 | 0.00024 | 0.00536 | 0.00021 | 0.00312 | -0.0295 | 0.8469 | 0.436 | 0.5352 | 0.5912 | 0.2104 | 0.2886 | 3 |
| MPT | 'Medial pretectal area' | 0.00971 | 0.01661 | 0.01500 | 0.02077 | 0.01386 | 0.00153 | 0.00269 | 0.00807 | 0.00866 | 0.02371 | 0.02601 | 0.02722 | 0.02059 | 0.00372 | 0.00846 | -0.0780 | 0.5826 | 0.7486 | 0.9729 | 0.3394 | 0.4852 | 0.8305 | 3 |
| SNc | 'Substantia nigra, compact part' | 0.00215 | 0.00263 | 0.00455 | 0.00224 | 0.00281 | 0.00108 | 0.00246 | 0.00248 | 0.00265 | 0.00446 | 0.00092 | 0.00531 | 0.00743 | 0.00421 | 0.00125 | -0.0317 | 0.8326 | 0.2954 | 0.4562 | 0.5777 | 0.1230 | 0.2403 | 2 |
| PPN | 'Pedunculopontine nucleus' | 0.00224 | 0.00251 | 0.00705 | 0.00464 | 0.00177 | 0.00141 | 0.00515 | 0.00142 | 0.00093 | 0.00654 | 0.00115 | 0.01054 | 0.01128 | 0.00376 | 0.00263 | -0.0679 | 0.5213 | 0.2293 | 0.2827 | 0.8687 | 0.0902 | 0.1333 | 3 |
| IF | 'Interfascicular nucleus raphe' | 0.00119 | 0.00221 | 0.00231 | 0.00144 | 0.00020 | 0.00131 | 0.00180 | 0.00420 | 0.00170 | 0.00253 | 0.00041 | 0.00198 | 0.00300 | 0.00111 | 0.00313 | -0.0186 | 0.5213 | 0.2239 | 0.9942 | 0.2824 | 0.1056 | 0.9215 | 2 |
| IPN | 'Interpeduncular nucleus' | 0.00201 | 0.00155 | 0.00160 | 0.00344 | 0.00069 | 0.00250 | 0.00143 | 0.00177 | 0.00167 | 0.00202 | 0.00073 | 0.00071 | 0.00231 | 0.00029 | 0.00311 | -0.0030 | 0.7746 | 0.8932 | 0.9965 | 0.5117 | 0.6639 | 0.9421 | 3 |
| RL | 'Rostral linear nucleus raphe' | 0.00317 | 0.00707 | 0.01023 | 0.00433 | 0.00046 | 0.00411 | 0.00376 | 0.01370 | 0.00146 | 0.01056 | 0.00256 | 0.00346 | 0.01407 | 0.00130 | 0.01139 | -0.1171 | 0.4983 | 0.4249 | 0.8486 | 0.2596 | 0.2014 | 0.5991 | 3 |
| CLI | 'Central linear nucleus raphe' | 0.00369 | 0.00449 | 0.01023 | 0.00380 | 0.00002 | 0.00672 | 0.00421 | 0.00749 | 0.00047 | 0.00980 | 0.00019 | 0.00012 | 0.01665 | 0.00087 | 0.02327 | -0.1148 | 0.9567 | 0.5711 | 0.6607 | 0.7845 | 0.3134 | 0.3960 | 3 |
| DR | 'Dorsal nucleus raphe' | 0.00474 | 0.00492 | 0.00566 | 0.00466 | 0.00079 | 0.01234 | 0.00118 | 0.00200 | 0.00000 | 0.01756 | 0.00012 | 0.00057 | 0.00854 | 0.00026 | 0.01326 | -0.0385 | 0.9993 | 0.9737 | 0.9894 | 0.9729 | 0.8332 | 0.8940 | 3 |
| P | 'Pons' | 0.00734 | 0.00620 | 0.00703 | 0.00926 | 0.00952 | 0.00497 | 0.00563 | 0.00578 | 0.00595 | 0.00590 | 0.00715 | 0.00873 | 0.00873 | 0.00597 | 0.00470 | -0.0009 | 0.6552 | 0.9783 | 0.7315 | 0.3870 | 0.8485 | 0.4612 | 3 |
| P-sen | 'Pons, sensory related' | 0.00890 | 0.00705 | 0.00744 | 0.01356 | 0.00800 | 0.00303 | 0.01099 | 0.00570 | 0.00875 | 0.00631 | 0.00745 | 0.01043 | 0.00948 | 0.00548 | 0.00438 | 0.0079 | 0.7354 | 0.8569 | 0.969 | 0.4641 | 0.6099 | 0.8187 | 3 |
| NLL | 'Nucleus of the lateral lemniscus' | 0.00120 | 0.00205 | 0.00482 | 0.00154 | 0.00178 | 0.00104 | 0.00043 | 0.00220 | 0.00190 | 0.00173 | 0.00236 | 0.00674 | 0.00472 | 0.00561 | 0.00219 | -0.0385 | 0.1202 | 0.0585 | 0.1237 | 0.0424 | 0.0117 | 0.0299 | 3 |
| P-mot | 'Pons, motor related' | 0.00715 | 0.00560 | 0.00601 | 0.01348 | 0.00549 | 0.00578 | 0.00526 | 0.00377 | 0.00721 | 0.00616 | 0.00775 | 0.00945 | 0.00317 | 0.00365 | | -0.0053 | 0.7867 | 0.9038 | 0.9696 | 0.5205 | 0.6809 | 0.8190 | 3 |
| P-sat | 'Pons, behavioral state related' | 0.00428 | 0.00472 | 0.00696 | 0.00940 | 0.00289 | 0.00294 | 0.00189 | 0.00449 | 0.00312 | 0.00840 | 0.00289 | 0.00424 | 0.01323 | 0.00216 | 0.00820 | -0.0588 | 0.9766 | 0.6637 | 0.7144 | 0.8429 | 0.4045 | 0.4482 | 3 |
| MY | 'Medulla' | 0.00609 | 0.00642 | 0.00532 | 0.00421 | 0.00824 | 0.00342 | 0.00849 | 0.00608 | 0.00614 | 0.00492 | 0.00853 | 0.00724 | 0.00648 | 0.00408 | 0.00350 | -0.0159 | 0.9751 | 0.8849 | 0.647 | 0.8367 | 0.6502 | 0.3903 | 3 |
| MY-sen | 'Medulla, sensory related' | 0.00628 | 0.00647 | 0.00541 | 0.01334 | 0.00524 | 0.00190 | 0.00463 | 0.00744 | 0.00615 | 0.00726 | 0.00503 | 0.00779 | 0.00843 | 0.00366 | 0.00177 | -0.0081 | 0.9968 | 0.9539 | 0.818 | 0.9420 | 0.7791 | 0.5588 | 3 |
| MY-mot | 'Medulla, motor related' | 0.00330 | 0.00248 | 0.00307 | 0.00986 | 0.00159 | 0.00029 | 0.00144 | 0.00293 | 0.00137 | 0.00321 | 0.00240 | 0.00469 | 0.00449 | 0.00230 | 0.00080 | -0.0122 | 0.931 | 0.9952 | 0.8356 | 0.7277 | 0.9284 | 0.5797 | 3 |
| MY-sat | 'Medulla, behavioral state related' | 0.00477 | 0.00930 | 0.00784 | 0.00644 | 0.00638 | 0.00251 | 0.00376 | 0.01313 | 0.00412 | 0.00798 | 0.01197 | 0.01671 | 0.00644 | 0.00818 | 0.00003 | -0.0251 | 0.222 | 0.6946 | 0.9303 | 0.0931 | 0.4230 | 0.7280 | 3 |

**Table 3.** ANOVA Tables for Regional Axon Density Comparisons. Related to Figures 2,4,5.
**Whole-Brain 2-Way ANOVA (Figures 4, 5, S6)**

| ANOVA table | F (DFn, DFd) | P value |
|---|---|---|
| Cell Type x Brain Region | F (278, 1251) = 3.959 | P<0.0001 |
| Cell Type | F (2, 9) = 5.690 | P=0.0253 |
| Brain Region | F (139.0, 1251) = 20.28 | P<0.0001 |

**Figure 4D: ORB Layer-Specific Innervation**

| ANOVA table | F (DFn, DFd) | P value |
|---|---|---|
| Layer x Cell Type | F (6, 27) = 4.225 | P=0.0040 |
| Layer | F (1.539, 13.85) = 21.45 | P=0.0001 |
| Cell Type | F (2, 9) = 39.86 | P<0.0001 |

**Figure 4D: AUDs Layer-Specific Innervation**

| ANOVA table | F (DFn, DFd) | P value |
|---|---|---|
| Layer x Cell Type | F (8, 36) = 1.635 | P=0.1493 |
| Layer | F (1.215, 10.93) = 7.283 | P=0.0172 |
| Cell Type | F (2, 9) = 44.48 | P<0.0001 |

**Figure 4D: TEa Layer-Specific Innervation**

| ANOVA table | F (DFn, DFd) | P value |
|---|---|---|
| Layer x Cell Type | F (8, 36) = 2.466 | P=0.0305 |
| Layer | F (1.324, 11.92) = 4.447 | P=0.0484 |
| Cell Type | F (2, 9) = 45.49 | P<0.0001 |

**Figure 4D: ECT/PERI Layer-Specific Innervation**

| ANOVA table | F (DFn, DFd) | P value |
|---|---|---|
| Layer x Cell Type | F (6, 27) = 3.490 | P=0.0110 |
| Layer | F (1.118, 10.06) = 2.843 | P=0.1205 |
| Cell Type | F (2, 9) = 27.44 | P=0.0001 |

**Figure 4D: ENT Layer-Specific Innervation**

| ANOVA table | F (DFn, DFd) | P value |
|---|---|---|
| Layer x Cell Type | F (8, 36) = 5.357 | P=0.0002 |
| Layer | F (2.733, 24.60) = 38.73 | P<0.0001 |
| Cell Type | F (2, 9) = 11.15 | P=0.0037 |

**Figure S2G (Distinction Score)**
**Repeated measures ANOVA summary**

| | |
|---|---|
| F | 14.46 |
| P value | <0.0001 |

| Multiple comparisons (Tukey's) | Summary | Adjusted P Value |
|---|---|---|
| DeepTraCE vs untrained TRAILMAP mo | **** | <0.0001 |
| DeepTraCE vs Best Single Model | * | 0.0132 |
| DeepTraCE vs Max Probability of Models | *** | 0.0006 |
| DeepTraCE vs Sum of Models | * | 0.0211 |