Brain-wide projections and differential encoding of prefrontal neuronal classes underlying learned and innate threat avoidance

Michael W. Gongwer[1,2,3*], Cassandra B. Klune[1,2*], João Couto[4], Benita Jin[1,4] Alexander S. Enos[1], Rita Chen[1], Drew Friedmann[5] and Laura A. DeNardo[1,4#]

Author Affiliations: [1]Department of Physiology, [2]Neuroscience Interdepartmental Program, [3]Medical Scientist Training Program, [4]Deparatment of Neurobiology, University of California Los Angeles, [5]Cajal Neuroscience, Seattle, WA
*Denotes equal contribution
#Correspondence: Dr. Laura DeNardo, 10833 Le Conte Ave, Center for Health Sciences, David Geffen School of Medicine at UCLA, Los Angeles, CA 90095 ldenardo@ucla.edu

Abstract To understand how the brain produces behavior, we must elucidate the relationships between neuronal connectivity and function. The medial prefrontal cortex (mPFC) is critical for complex functions including decision-making and mood. mPFC projection neurons collateralize extensively, but the relationships between mPFC neuronal activity and brain-wide connectivity are poorly understood. We performed whole-brain connectivity mapping and fiber photometry to better understand the mPFC circuits that control threat avoidance. Using tissue clearing and light sheet fluorescence microscopy we mapped the brain-wide axon collaterals of populations of mPFC neurons that project to nucleus accumbens (NAc), ventral tegmental area (VTA), or contralateral mPFC (cmPFC) in mice. We present DeepTraCE, for quantifying bulk-labeled axonal projections in images of cleared tissue, and DeepCOUNT, for quantifying cell bodies. Anatomical maps produced with DeepTraCE aligned with known axonal projection patterns and revealed class-specific topographic projections within regions. During threat avoidance, cmPFC and NAc-projectors encoded conditioned stimuli, but only when action was required to avoid threats. mPFC-VTA neurons encoded learned but not innate avoidance behaviors. Together our results present new and optimized approaches for quantitative whole-brain analysis and indicate that anatomically-defined classes of mPFC neurons have specialized roles in threat avoidance.

Introduction The medial prefrontal cortex (mPFC) is a vastly interconnected brain region that controls complex cognitive and emotional functions including memory, decision making, social interactions and mood[1–3]. mPFC acts through its axonal projections to exert top-down control, biasing activity in downstream brain regions to promote adaptive behavioral responses in dynamic circumstances[4–9]. With the widespread use of opto- and chemogenetics to manipulate neural populations, a large number of studies have linked individual mPFC projections to discrete behavioral functions[5,6,8–16]. However, many mPFC-dependent behaviors are known to involve multiple downstream brain regions[17]. Recent work revealed the vast anatomical heterogeneity of mPFC projection neurons and their extensive axon collaterals[18]. Thus, there is a need to integrate anatomical findings with functional work to gain a deeper understanding of how classes of mPFC projection neurons contribute to behavior.

Many studies use retrograde viral approaches to activate, inhibit, or observe the activity of projection-defined populations[19–21]. In many cases, activation of axon collaterals beyond the projection of interest may inadvertently influence animal behaviors. Interpreting results of manipulation or observation of the somatic activity of virally- or genetically-targeted projection neurons therefore requires understanding their full projection patterns.

Traditional studies of axon collaterals have established an important foundation of knowledge but have been limited by low-throughput approaches. Anatomical mapping strategies include injecting retrograde tracers into multiple target areas and measuring overlap in source neurons[5,22]. Functional mapping strategies include recording from source neurons while performing multisite antidromic stimulation in target regions[23,24]. While informative, these approaches are biased in the sense that users predefine the target areas to investigate and are

limited to small numbers of regions. RNA-sequencing based approaches[25] allow higher
throughput examination of collateralization, but the spatial resolution is limited by tissue micro-
dissection volumes and analyses are also often biased to experimenter-defined sets of regions.
The Allen Connectivity Atlas[26] and the recent publication of a single neuron projectome of
mouse prefrontal cortex greatly improved our understanding of prefrontal projection patterns[18].
While highly informative, serial two-photon tomography and single neuron reconstruction require
specialized equipment and are computationally and time-intensive, making these approaches
impractical for most scientists to replicate.
On the other hand, tissue clearing[27,28] combined with light sheet fluorescence
microscopy (LSFM) is fast, easy and affordable and can preserve 3-D circuit architecture
without the need to align structures. These approaches are useful for comparing the projection
patterns of genetically, anatomically or behaviorally-defined neurons and generating new
hypotheses that can be tested using functional methods. For instance, users may be interested
in identifying novel circuits underlying behavior or in determining how genetic, environmental, or
pharmacological perturbations affect structural and functional connectivity. With commercial
lightsheet systems now readily available, these approaches are within reach for many labs. But
we need end-to-end open-source packages for performing whole-brain quantitative analyses.
To address this deficit, we developed DeepTraCE (Deep learning-based Tracing with Combined
Enhancement) for quantifying bulk-labeled fluorescent axons and DeepCOUNT for quantifying
fluorescently labeled cell bodies. These analysis pipelines build on the machine learning-based
image segmentation package TrailMap[29]. Together, they comprise a user-friendly, analysis
package for high-throughput, whole brain analysis based on tissue clearing and LSFM.
Advances in computer vision allow researchers to train classifiers to automatically detect
axons in tissue volumes[29,30]. However, classifiers sometimes fail to identify all axons because
axon appearance varies across brain regions. Axon morphology, including the extent and
symmetry of branching as well as the tortuousness of branches, varies by projection type[18].
Variance in axonal appearance is also partly due to the resolution limitations of most light sheet
microscopes that are capable of visualizing entire mouse brains in a timely fashion. Axons can
appear thick and bright in sparsely innervated brain regions near the surface of the brain, but
thin and indistinct in densely innervated regions deep in the tissue. To overcome these hurdles,
DeepTraCE allows users to combine multiple models that have been separately trained using
TrailMap to accurately identify axons with differing appearances. After segmenting the axons
using TrailMap models, DeepTraCE registers the brains to a common coordinate framework
and quantifies innervation density by brain region. Similarly, DeepCOUNT uses TrailMap to
detect cells and then quantifies cell bodies by brain region.
In this study, we used DeepTraCE, DeepCOUNT and fiber photometry to better
understand the mPFC circuits that control threat avoidance. Threat avoidance behavior is
critical for survival. While different brain regions have been implicated in innate and learned
avoidance, the prelimbic (PL) subregion of mPFC is integral to both[31–33]. But our understanding
of how PL circuits control threat avoidance remain incomplete. We focused on PL neurons that
project to the contralateral PL (cPL), nucleus accumbens (NAc) or ventral tegmental area (VTA).
PL-cPL, PL-NAc and PL-VTA neurons have distinct distributions across cortical layers, distinct
genetic profiles, and distinct functions[5,6,34]. We mapped the brain-wide collateral projections of
each neuronal class. DeepTraCE accurately detected axons throughout the brain,
outperforming other model combination methods. Next, we used DeepCOUNT and activity-
dependent genetic labeling to map brain-wide neuronal activation patterns after mice avoided
threats. We used these data to construct functional networks and found that PL was one of the
most highly connected nodes. Further, PL's functional targets were regions that are
preferentially targeted by axon collaterals of cPL, NAc or VTA projectors. Using fiber
photometry, we discovered class-specific activity during innate and learned avoidance.
Together, our work reveals that projection-defined PL classes have specialized roles in threat

avoidance. We also demonstrate the utility of combining common techniques like fiber
photometry with DeepTraCE and DeepCOUNT to understand the detailed structural and
functional connectivity of the cells being studied.
98
99   **Results**
100
DeepTraCE Workflow
We developed DeepTraCE, an open-source, end-to-end analysis pipeline for quantifying bulk
axonal projection patterns in cleared brains. DeepTraCE takes in raw images of fluorescently
labeled axons and then applies TrailMap[29], a machine learning pipeline that trains a 3D U-net
framework to automatically identify axons in cleared tissue (Figure 1A, Figure S1). TrailMap
provides a pre-trained model that users can fine tune to fit their samples. However, it was
challenging to train a model that could sufficiently generalize across the diversity of mPFC
axonal structures in all of their target regions; some axons were bright and delineated (e.g. in
cortical areas) while others were dense and indistinct (e.g. in the basolateral amygdala and
thalamus). To surmount this hurdle, DeepTraCE allows users to combine multiple trained
models to optimize axon detection throughout the brain. After importing 3-D image stacks,
multiple models trained using TrailMap fully segment each brain. DeepTraCE then registers
each brain to a standard atlas and generates a new image in which pixel values in each region
were extracted from the assigned model. Axons in the resulting images are then thinned to
single-pixel width[29] and the number of axon-containing pixels are quantified for each brain
region (Figure 1A).
117
Validation of model combination method
To determine how many models to combine, we examined raw images of fluorescently-labeled
axons taken with LSFM. We grouped brain regions into three main categories based on the
visual quality of PL axons (e.g. thick and delineated or dense and fuzzy). These categories
largely segregated with the physical location of brain regions in tissue volumes (Figure 1B). We
used TrailMap[29] to train three separate models that we fine-tuned for each set of regions (Figure
1B,C; Figure S1A–C).
To validate the DeepTraCE model assignments, we performed quantitative comparisons
of model accuracy in representative brain regions from each category, using human annotations
as a reference. Human experts traced pixels containing axons in image stacks from each group
of brain regions (Figures S1A–C). We segmented images using each of the three models in
addition to the original untuned TrailMap model that was trained on serotonergic axons[29]. We
then calculated the resulting axonal labeling density with reference to human labels.
In LSFM of cleared tissue, the size of small fluorescent structures is amplified such that
a thin object such as an axon occupies several more pixels in the image than it does in true
biological space. We account for this overrepresentation by thinning segmented axons to a
single pixel width[29]. However, because rapid imaging of intact rodent brains requires lower
optical resolution, it can be challenging in some regions to hand-label axons with single pixel
precision. This caveat to whole-brain light sheet microscopy approach prevents the calculation
of "true positive" and "true negative" rates based on human annotation. So, common metrics
such as F1 Score, precision, and recall do not provide interpretable measures of segmentation
accuracy. Instead, we calculated a 'distinction score' that measures the density of segmented
axons in regions that humans estimated as containing axons vs. not containing axons. A higher
distinction score indicates better overlap with human labels. When applied to our validation set,
Models 1, 2, and 3 produced the highest distinction scores in images from the brain regions
assigned to each model (Figure 1D).
Previous studies used alternative approaches to overcome the limitations of a single-
model approach in whole-brain image segmentation. Some excluded deeper, more densely

innervated areas from analysis[35]. Others combined probability maps from different segmentation
methods by extracting the maximum probability for each pixel[36]. To compare DeepTraCE with
these alternative approaches, we calculated the distinction score across all human-annotated
images in the validation set using the original TrailMap model that was trained on serotonin
neurons, our best-trained single model, the maximum probability between our three models, the
sum of probabilities, and the DeepTraCE combined models. DeepTraCE had the highest
distinction score (Figure 1E), indicating that it performed more accurate axon segmentation from
images.
154
Viral circuit mapping and brain clearing strategy
We mapped the brain-wide axon collaterals of PL neurons that project to cPL, NAc or VTA. Our
goals were 1) to demonstrate the utility of DeepTraCE and 2) to provide an important
anatomical resource for researchers studying mPFC function. Therefore, we used a commonly
employed dual-virus approach to fluorescently label populations of PL projection neurons[37]. In
contrast to multi-site retrograde tracer injections, this approach allows us to visualize the bulk
brain-wide collateralization patterns of each projection-defined class of neurons. We first
injected an axon-terminal-transducing adeno-associated virus[38] expressing Cre recombinase
(AAVrg-hSyn-Cre) into cPL, NAc, or VTA. Next, into PL, we injected AAV expressing Cre-
dependent EYFP-tagged Channelrhodopsin-2 (AAV8-hSyn-DIO-ChR2-EYFP), which traffics
efficiently to axons (Figure 2A-C). We used the iDISCO clearing variant, Adipo-Clear[39,40] to
immunostain intact brains for EYFP and render the tissue transparent. We then imaged the
hemispheres ipsilateral to the PL injection with LSFM.
In separate animals, we used confocal microscopy to examine the distribution of
retrogradely labeled cell bodies in brain sections (Figure S2A). Different retrograde tracers can
have preferences for particular neuronal types. For instance, while AAVrg favors cortical layer
(L)5 over L6, rabies virus has the opposite preference[41]. Cholera toxin subunit B (CTB) is
inefficient for labeling pyramidal tract neurons in L5[42]. To assess potential biases in our
retrograde labeling method, we analyzed the layer distribution for AAVrg-Cre-mCherry+ cells
and the combination of AAVrg-hSyn-Cre-mCherry and AAV8-hSyn-DIO-ChR2-EYFP (assessed
based on EYFP fluorescence). As a comparator, we measured the layer distribution of CTB+
cells in PL following injections into cPL and NAc. Consistent with previous reports, we did not
observe retrogradely labeled PL cells when we injected CTB into the VTA[42]. We determined
layer boundaries based on DAPI nuclear staining in combination with immunostaining for Ctip2,
a marker of subcerebral projection neurons located in L5b-6[43]. Layer boundaries and
thicknesses aligned with previous literature (Figure S2B)[11,44,45]. Consistent with previous
findings[45–47], PL-cPL and PL-NAc neurons were preferentially located in L2–5a and PL-VTA
neurons were restricted to L5b. We found no statistical differences in the distribution of cells
labeled with AAVrg alone, AAVrg+AAV8 or CTB (Figure S2A-C). As reported previously, we
observed few AAVrg-labeled cell bodies in L6[41,42]. While L6 neurons were likely
underrepresented in our datasets, we could still extract meaningful differences between classes.
186
Whole-brain projection patterns of PL-cPL, PL-NAc and PL-VTA neurons
We used DeepTraCE to determine the extent to which PL-cPL, PL-NAc and PL-VTA neurons
collateralize to other brain regions. Examination of thin optical sections revealed widespread
collateralization for all three classes. Overall, innervation patterns were consistent with reports
of general mPFC projection patterns[22,45,48]; mPFC axons were prominent in cortical association
areas, striatum, midline thalamus, claustrum, amygdala, hypothalamus and midbrain but largely
absent from hippocampus, sensory thalamus, and relatively sparse in primary sensory areas
(Supplemental Video; Figure 2D–F; Figure S3–5).
We generated dotograms[26] to visually summarize the inter-class distinctions. To create
the dotogram, we averaged axon density across all samples for a given projection class and

then overlaid the samples in brain space. Dot color indicates the neuronal class and dot size corresponds to the averaged axonal density within a voxel (Figure 2G). Compared to PL-VTA neurons, PL-cPL and PL-NAc neurons more densely innervated cortical areas, especially the temporal association area (TEa), ectorhinal area, and entorhinal cortex. On the other hand, PL-VTA collaterals preferentially innervated thalamic (TH) and midbrain (MB) regions, a subset of which also received collateral input from PL-NAc neurons. All three subclasses innervated NAc and olfactory tubercle (OT), while PL-NAc neurons were the primary source of collaterals to piriform cortex (PIR).

We next quantified regional innervation densities and plotted them as a heatmap. For each brain, we normalized the number of axon-containing pixels in each region to the total number of axon-containing pixels in the brain. We then sorted regions according to their innervation density by projection classes; regions receiving more PL-cPL collaterals are on top and those receiving more PL-VTA collaterals are in the bottom half of the heatmap (Figure 2H). While there was little overlap between projection patterns of PL-cPL and PL-VTA neurons, PL-NAc neurons shared several projection targets with both other classes. Brain-wide statistical comparisons confirmed the presence of 29 significantly differentially innervated brain regions, the most notable of which were in the cortex, subplate, and thalamus (Figure 2H, S6, Table 2, Table 3).

Because PL-cPL, PL-NAc and PL-VTA projection neurons all collateralize broadly, we wondered if these neuronal classes could be separated based on their whole-brain projection patterns. To test this, we used two dimensionality reduction techniques. We first performed hierarchical clustering, in which Euclidean distance between individual brains based on axonal density in each subregion is marked by a higher branch point on the graph. PL-VTA neurons formed a distinct cluster, while PL-NAc and PL-cPL collaterals formed partially overlapping clusters (Figure 2I). This aligns with our observation that the projection patterns of PL-NAc neurons shared more overlap with PL-cPL than PL-VTA neurons (Figure 2H).

We also used principal component analysis to assess differences in whole-brain projection patterns for each cell class. We plotted the location of each brain along the axes of the first two principal components. While PL-cPL and PL-VTA projection classes could be clearly separated along the first principal component, PL-NAc brains were positioned in between, with some overlap with PL-cPL brains (Figure 2J). To assess which projection patterns distinguished the classes, we plotted the weights of each brain region in contribution to the first principal component. Positively weighted regions generally had the most innervation from PL-cPL collaterals including TEa, PERI, and ECT, and negatively weighted regions were those with most innervation from PL-VTA collaterals, including MTN and midbrain areas (Figure 2H). These data show that while cPL and NAc-projectors have substantial overlap, PL-cPL and PL-VTA neurons represent separable classes, distinguished in large part by their collaterals to TEa/PERI/ECT, olfactory and limbic areas, and thalamus and midbrain, respectively. Together, the innervation patterns we observed were consistent bulk projection mapping[26,49,50] and single neuron reconstructions[18,51], indicating that DeepTraCE accurately captured meaningful class-specific distinctions in collateral targeting.

Layer-specific and topographic innervation patterns in the cortex and subplate
Whole-brain analysis of bulk labeled neuronal classes is well-suited for understanding the geometric organization of axonal projections *within* brain regions, which can have important functional implications. The neocortex is organized into layers that contain distinct neuronal types with different morphology, physiology, and connectivity[44,52]. Long-range cortico-cortical axons may innervate superficial or deep layers depending on the hierarchical relationship with the target region[44,52]. Compared to sensory and motor cortices, much less is known about the layer organization of long-range mPFC connectivity, especially for specific projection classes. To investigate this, we plotted PL-cPL, PL-NAc and PL-VTA collateral innervation density

across layers in cortical target areas (Fig. 3A-B). In ORB, all classes preferentially targeted
superficial layers (L1, L2/3). In AUD, TEa, ECT, and PERI, PL-cPL and PL-NAc collaterals
preferentially targeted superficial layers (L1, L2/3) while PL-VTA collaterals targeted the deep
layers (L5, L6) (Figure 3B, Table 3). In ENT, axon collaterals were more prominent in deep
layers for all 3 classes. These findings suggest that PL-cPL, PL-NAc and PL-VTA may all
participate in feedforward or feedback connectivity depending on the target region.
Many PL target regions contain functional gradients that are orthogonal to layer-defined
microcircuits. We therefore investigated whether PL classes project to topographically-defined
locations in target regions (Figure 3C). PL-NAc collaterals were especially prominent in the
posterior PIR. Compared to PL-cPL and PL-NAc, PL-VTA collaterals had a tighter distribution in
the anterior portion of agranular insula (AI). PIR and AI both have contain anatomical and
functional gradients[53,54]. Our data suggest that class-specific PL projections may be a
determining factor in these gradients.
While all projection classes sent collaterals to the amygdalar complex, amygdalar nuclei
were differentially innervated. Most notably, the anterior basolateral amygdala (BLAa) was
robustly innervated by PL-cPL and PL-NAc neurons, but less so by PL-VTA neurons.
Meanwhile, compared to PL-cPL neurons, PL-NAc neurons sent many more collaterals to
olfactory amygdalar areas, including anterior cortical amygdalar area (COAa) and piriform
amygdalar area (PAA) (Figure 3D,E). As PL-NAc collaterals were also more prominent in
posterior PIR, which was recently found to play a role in spatial cognition[53], PL-NAc neurons
may have privileged control over areas dedicated to olfactory processing during cognition and
emotional learning.
270
Whole-Brain Cell Counting Pipeline
The application of computer vision to light sheet data extends beyond axon tracing and can be
used to identify signatures of brain function. Expression of IEGs such as c-fos are frequently
used as a proxy of neural activity[55,56]. For instance, recent studies have used tissue clearing,
IEG-labeling, and LSFM to screen for changes neuronal activation following experiences with
drugs or fear learning[28,48,57,58]. Quantifying IEG-expressing cells on brain-wide scale can serve
as means to infer behaviorally-relevant changes in functional connectivity. However, currently
available open-source packages for whole-brain cellular quantification can be error prone,
especially in regions of particularly dense labeling.
As a companion to DeepTraCE, we developed DeepCOUNT (Deep-learning based
Counting of Objects via 3D U-Net pixel Tagging) (Figure 4A). We first trained a TrailMap model
to recognize fluorescently labeled cell bodies. This produces a probability map of which pixels
are most likely to contain a cell body. After thresholding the images to a desired probability
cutoff, they are registered to a standard brain, and DeepCOUNT uses a 3D maxima detection
strategy is used in combination with a connected component analysis to ensure each neuron is
represented by a single pixel. This single-pixel output can then be used to obtain regional cell
counts across the brain.
We compared DeepCOUNT performance to that of ClearMap[28], a commonly used cell
detection algorithm designed for tissue clearing and LSFM, and human observers. Two human
observers analyzed image volumes from six different brain regions. Cell counts produced with
DeepCOUNT were highly correlated with human observers and DeepCOUNT produced more
accurate counts than ClearMap, using human counts as the ground truth (Figure 4B,C).
293
Whole-brain threat avoidance networks
We used DeepCOUNT to map regions and functional connections involved in threat avoidance
at a whole-brain level. PL is required for threat avoidance behavior[32,59]. Previous studies
examining c-fos expression following active avoidance behavior analyzed only discrete regions
of interest[60–62], some of which are PL target sites. How PL coordinates threat avoidance-related

neural activity on a brain-wide scale is not understood. To begin to address this question, we
used DeepCOUNT and activity-dependent genetic labeling to map brain-wide neuronal
activation following a platform mediated avoidance (PMA)[32], a form of learned threat avoidance.
We labeled activated neuronal populations using TRAP2 mice (*Fos^icre-2A-ERT2*), in which
the c-fos promoter drives expression of tamoxifen-inducible Cre recombinase. We crossed
TRAP2 to the Ai14 Cre reporter line[48,63,64]. We then trained the *TRAP2;Ai14* mice in PMA and
TRAPed them after a threat avoidance retrieval session by pairing that experience with a
tamoxifen injection (Figure 4D, S7A). We used DeepCOUNT to quantify TRAPed cell density in
each brain region and plotted these data as a heatmap (Figure 4E). Principal component
analysis of whole-brain TRAPed cell counts separated PMA-trained and non-shocked control
animals along the first principal component (PC1) (Figure 4F). PC1 weights revealed that PMA-
activated brain regions were biased toward the cortex, hippocampal formation, and cortical
subplate (Figure 4E). We plotted TRAPed cell density across the brain as dotograms (Figure
4G) and compared cell counts between PMA and control animals in PL and several key targets.
Our DeepCOUNT screen identified regions known to be involved in threat avoidance (e.g. PL
and BLA). We also identified several regions with unknown roles in threat avoidance (e.g. the
TEa and postsubiculum (POST)) that will be of interest for future functional studies (Figure
4E,H).
To better understand the organization of PL circuits that control threat avoidance, we
assessed functional connectivity based on whole-brain TRAPing. We computed inter-regional
correlations for groups of PMA and control mice. This allowed us to identify sets of regions
where the numbers of TRAPed cells co-varied across mice (Figure S7B). Regions that co-vary
may constitute elements of a network engaged during learned threat avoidance. We generated
network graphs for each condition based on the strongest correlations (Pearson's r>0.9,
P<0.05, Figures 4I, S7C).
Our network analysis revealed PL as one of the mostly highly connected nodes in the
PMA group (Figure 4J). PL was functionally connected to several cortical areas (TEa, ORBl,
MOp, SSp, SSs), taenia tecta (TT), the lateral septal complex (LSx), claustrum (CLA), several
amygdalar regions (LA, BLAa, BMAa), and the median eminence (ME). Our neuroanatomical
analysis had already revealed that some these regions are preferentially innervated by PL-cPL
(CLA, TEa, TT) or PL-VTA (LSx). Compared to PL-VTA, PL-NAc and PL-cPL neurons send
denser projections to BLAa. Compared to PL-cPL neurons, PL-NAc and PL-VTA neurons send
denser projections to BMAa. To illustrate this, we colored the PL-connected nodes based on
their PCA weight from the analysis shown in Figure 2 (Figure 4I). These data suggest that all
three neuronal classes likely contribute to threat avoidance behavior and that they may act via
particular collateral projections.
335
Fiber Photometry During Platform-Mediated Avoidance
We investigated how different PL cell classes contribute to threat avoidance by using fiber
photometry to record population-level neural activity. Given the high overlap of collateral
projection targets for PL-cPL and PL-NAc neurons (Figures 2I,J), we further separated these
classes using an intersectional viral strategy[65] (Figure 5A). To record neuronal activity in PL-cPL
neurons that do not project to NAc, we injected AAVrg-Cre into cPL and AAVrg-Flp into NAc.
We then injected Cre-On Flp-Off GCaMP6M into PL. We switched the Cre and Flp injection
sites to record from NAc-projecting neurons that do not project to cPL. To record from VTA-
projecting neurons, we injected AAVrg-Cre into VTA and DIO-GCaMP6M into PL (Figure 5B).
Most GCaMP+ cell bodies from the intersectionally-defined cPL- and NAc-projecting neurons
were in superficial layers of PL, while cell bodies from VTA-projecting neurons were restricted to
the deeper layers (Figure 5C), as expected from our initial layer analysis (Figure S2A). We
observed few GCaMP+ axon terminals in NAc from the cPL-projecting group, and few GCaMP+

axon terminals in cPL from the NAc-projecting group (Figure 5D), suggesting our intersectional
strategy was efficient in separating these populations.
We recorded from PL while mice performed PMA across 3 days of training. By training
day 3, mice successfully avoided most shocks (Figure 5E). We first analyzed responses to the
conditioned tone. Interestingly, tone-evoked neural activity distinguished between epochs when
mice were off vs. on the safety platform (Figure 5F). cPL-projecting neurons had significantly
higher activity at the tone onset when the animal was off the platform. As the tone progressed,
activity in these neurons gradually decreased. We observed a similar trend in NAc-projecting
neurons. In contrast, PL-VTA cells lacked a tone onset response, but did have a gradual
decrease in activity during the conditioned tone (Figure 5G). These findings suggest neural
activity in response to a conditioned stimulus is modulated by whether threat requires action,
and that these patterns vary across projection classes.
Activity in all classes sharply increased upon foot shock onset (Figure 5H), and we found
no differences in peak shock response between the three classes (Figure 5I). NAc-projecting
neurons had significantly higher activity than the other two classes in the ten seconds preceding
a shock (Figure 5I), consistent with a larger role for those cells in risky exploration. Further,
while activity in PL-VTA neurons rapidly returned to baseline levels following the foot shock,
activity in PL-cPL and PL-NAc neurons had prolonged deviations from baseline for up to a
minute after the shock. Activity in NAc-projecting neurons remained elevated above baseline for
the forty seconds following a foot shock, while activity in cPL-projecting neurons decreased
steadily during the same period (Figure 5I,H). These data suggest that while all PL classes
encode aversive stimuli, prolonged post-shock activity in PL-cPL and PL-NAc neurons may play
specialized roles in action-outcome learning.
372
Prefrontal Neuron Classes Distinguish between Learned and Innate Avoidance
To further classify the effects of aversive learning on neural activity in PL cell classes, we
compared their activity during learned vs. innate threat avoidance. Using the same mice from
the PMA experiments, we recorded neural activity in the Elevated Zero Maze (EZM), in which
two quarters of an elevated ring are protected by walls and the other two quarters are open.
Mice innately avoid the open arms of this apparatus, where there is more perceived threat
potential.
We compared activity during entries and exits from the safe zone in each assay. In PMA,
population activity in all three neuronal classes increased prior to platform entries (Figure 6A). In
contrast, only NAc- and cPL-projecting neurons increased their activity prior to entry into the
closed arm of the EZM (Figure 6B). During exits from the safe zone, a form of risky exploration,
all three PL classes had similar increases in activity during PMA (Figure 6C). In innate
avoidance, however, exits from the closed arm were marked by a significantly larger response
in NAc-projecting neurons compared to the other two cell populations (Figure 6D). Together,
these data indicate that aversive learning engages VTA-projecting neurons to behavioral circuits
for threat avoidance and that PL-NAc neurons preferentially encode risky exploration.
A concern with comparing fiber photometry recordings across different classes of
neurons is that differences in signal between the classes could lead to exaggerated results in
one direction. While PL-VTA neurons had lower overall activity during risky exploration of the
shock bars and open arm, all three PL classes had similar activity during head dips on the EZM,
another form of exploratory behavior (Figure 6E). This suggests that PL-VTA neurons did not
simply have lower activity than the other classes during behavior, but instead, that these
differences are specific to situations when animals are navigating the environment. To further
corroborate this, we performed a separate analysis of signals normalized to the peak of the
average shock response for an animal, as the shock provides a consistent time-locked
response in each animal. All between-group statistical comparisons remained consistent upon

normalization to the shock response (Figure S8), suggesting the effects we observe are not
driven by differences in signal quality between the three classes.
401
Discussion
In this study, we integrated whole brain mapping with the observation of prefrontal neural
activity to better understand how mPFC controls threat avoidance behavior. We introduce
DeepTraCE and DeepCOUNT, two new open-source analysis pipelines that extend the utility of
TrailMap to quantify bulk axonal projection patterns and IEG labelling, respectively. We used
DeepTraCE to produce whole brain structural projection maps of three populations of mPFC
projection neurons: PL-cPL, PL-NAc, and PL-VTA. We next combined activity-dependent
genetic labeling with DeepCOUNT to relate the structural and functional organization of brain-
wide networks for threat avoidance. Finally, with whole-brain projection maps as a foundation,
we used intersectional viral targeting to separate the overlapping populations of PL-cPL and PL-
NAc neurons and then recorded the activity of PL-cPL, PL-NAc and PL-VTA neuronal
populations during learned and innate threat avoidance. Our study reveals PL class-specific
roles in threat avoidance and demonstrates the utility of DeepTraCE and DeepCOUNT for
linking high-throughput neuroanatomy with functional techniques to explore the reveal
mechanisms of complex behavior (Figure 7).
While we have gained invaluable insights into the complex anatomy of cortical neurons
from single neuron reconstructions[18,26,66], the equipment, computing power, expertise and time
required to reconstruct individual neurons are out of reach for many labs. Other high-throughput
techniques such as MAPseq[67] provide single cell resolution, but are costly and limited to a
predefined number of target regions. While bulk labelling approaches lose granularity of
individual neuron differences, DeepTraCE's high-throughput workflow combined with its reliance
on common viral techniques makes it a useful tool to tackle a variety of unanswered questions.
Capitalizing on mouse Cre-driver lines and floxed alleles, DeepTraCE can be used to analyze
the projection patterns of defined neuronal populations and how they can be altered by genetic
manipulations. Similarly, DeepTraCE and DeepCOUNT can be used to define differences in
structural and functional connectivity resulting from environmental insults such as chronic
stress. Given the sample size needed to pursue these questions, our pipeline's high-throughput
processing puts these experiments within practical reach when compared to single neuron
reconstruction. With DeepTraCE, in addition to simply quantifying relative densities of axonal
projections in target regions, users can assess within-region changes in layer-specific or
topographic targeting. This information can guide the design of subregion-targeted functional
studies. Experimenters can use the same viral-genetic strategies to drive expression of opsins
or GCaMP proteins in neuronal classes of interest and then implant optic fibers with targeting
informed by the anatomical maps.
It is important to also note the caveats associated with retrograde viruses, brain clearing
and LSFM. Compounds for retrograde labeling can have biases against projection classes[68].
AAVrg viruses have been shown to have preferential labelling of corticothalamic (CT) neurons in
L5, and biases against CT neurons in L6[41,42], while CTB showed the opposite pattern of layer
bias[42]. In line with this, we observed few AAVrg+ neurons in L6, but some CTB+ neurons
(Figure S2A). So we may have underestimated the contributions of L6 cells to the population
data for PL-cPL and PL-NAc neurons[22,44]. Like other retrograde labeling methods, the full range
of neuronal types that may be resistant to AAVrg is not well understood. Also, tissue clearing
does not always achieve complete transparency. This issue, together with resolution limits of
light sheet microscopes that can visualize entire mouse brains in a timely fashion, can produce
blurring of structures deep in the tissue. Therefore, estimates of axonal density may be less
accurate in deep brain structures. However, estimates of axon density are likely to be biased to
a similar degree for different neuronal classes in any given target structure. So, high-throughput

mapping using viral tracing, brain clearing and LSFM is well-suited for comparing inter-class
differences.
In our fiber photometry experiments, the intersectional viral strategy we used to parse
apart PL-cPL and PL-NAc populations relied on an AAVrg injection expressing Flp recombinase
to block GCaMP expression in projection populations. Our AAVrg-Flp injections did not span the
entirety of cPL or NAc. Therefore, we do not claim to have completely segregated these
populations but rather biased them towards each projection type. Finally, like most mapping
studies, we visualized axons but not synapses. Therefore, our DeepTraCE quantifications may
also include axons of passage. To mitigate this, our models were trained to exclude major white
matter tracts.
Despite these caveats, our brain wide projections maps of PL-cPL, PL-VTA, and PL-NAc
neurons aligned with known projection patterns[18,22,26,48–51] – underscoring the accuracy of
TrailMap[29] and DeepTraCE – and revealed unappreciated topographic innervation patterns.
Consistent with bulk tracing studies[48–50], we observed the densest PL projections in temporal
association cortices, claustrum, striatum, BLA, lateral hypothalamus, polymodal thalamic nuclei,
hypothalamus and midbrain. These bulk projection patterns are the sum of multiple classes.
Like sensory cortices, mPFC contains intratelencephalic (IT) neurons that project within the
cortex and pyramidal tract (PT) neurons, that project to subcortical areas. PL-cPL neurons fall in
the IT class, PL-VTA neurons are from the PT class, and PL-NAc contain members of both[44]. In
line with these classifications and single neuron reconstructions[18,44,51], we observed that PL-cPL
neurons were enriched in superficial layers (L2-5a) and distinguished by their collaterals to other
cortical association areas, subplate (CLA and BLA), and striatum. Consistent with the well-
described features of PT neurons[18,44,66], we observed retrogradely labeled PL-VTA neurons with
somas localized to L5b and axon collaterals in the thalamus, pons, PAG, striatum, and
neuromodulatory centers. Also consistent with prior work, PL-NAc neurons collateralized
broadly to both IT and PT targets [44]. The diversity of the collateral targets of PL-NAc neurons is
supported by single neuron reconstructions which further show a variety of subclasses targeting
only a few structures[18]. Overall, there was excellent alignment between our data and data
collected using methods that have higher spatial resolution. This underscores the accuracy of
DeepTraCE and its utility for capturing meaningful class-specific differences in connectivity in a
high-throughput manner.
We also observed unappreciated topography in how each class innervated target
regions. PL-NAc collaterals preferentially innervated the posterior PIR, which was recently found
to have a specialized role in spatial cognition[53]. Compared to PL-cPL and PL-NAc, PL-VTA
collaterals were restricted to the AI. PIR and AI are both complex anatomical hubs with distinct
functional zones along the rostro-caudal axis[53,69,70]. Preferential innervation by PL projection
neurons may underlie some of the functional specializations within these regions.
On a brain wide scale, counts of IEG-expressing cells can serve as means to identify
behaviorally-relevant activity and functional connectivity[48,57,58]. Previous studies examining c-fos
expression following active avoidance behavior analyzed only discrete regions of interest[60–62].
We used DeepCOUNT and TRAP2 mice to measure brain-wide neuronal activation following
threat avoidance. DeepCount recapitulated previous findings of activation in PL, BLA and OFC
following PMA[60]. We also found high numbers of TRAPed cells in POST and TEa, whose roles
in threat avoidance are unknown. POST is important for spatial learning[71–73]. During PMA
learning, prefrontal connections to this area may facilitate learning of the location of the safety
platform. Moreover, PL projection to TEa may facilitate long-term storage of the avoidance
memory as TEa is necessary for long term retrieval of cued fear[74]. Functional connectivity
analysis revealed the major functional targets of PL. Of these, TEa, CLA, and TT are highly
innervated by the collaterals of PL-cPL projection neurons, while LSx and BMAa are more
densely innervated by PL-NAc and PL-VTA collaterals. Only PL-NAc neurons have been

previously studied in active avoidance[9]. Our study highlights how whole brain anatomical and functional maps can be used together to identify novel behaviorally-relevant pathways.

We used fiber photometry to determine how each class of neuron encodes threat avoidance. We found that intersectionally-defined PL-NAc and PL-cPL neurons increased their activity in response to the conditioned tone in PMA, but only when mice were off the safety platform, suggesting those populations encode the predictability of threat. In contrast, PL-VTA neurons decreased their activity in response to the threat predictive cue. Single unit recordings haven shown that distinct populations are excited and inhibited by a conditioned tone during PMA[59]. Our results indicate that the projection targets we studied may be a factor that separates these populations.

PL-cPL connections are important for both hemispheric synchronization and lateralization of mPFC activity. While lateralization has been associated with anxiety-like behavior[75] and stress responses in rodents[76–78], little is known about the behavioral functions of PL-cPL neurons, especially in aversive behaviors. Our data show that PL-cPL neurons persistently tracked threat, threat predictive cues, and threat avoidance behaviors. In the future, studies manipulating these neurons will be required to understand whether these activity patterns are necessary for learned or innate avoidance.

We found that PL-NAc neurons encoded risky exploration. This is consistent with previous work showing that excitation of PL-NAc projections decreases avoidance behaviors[9]. However, individual neurons within this projection class can have heterogeneous functions. For instance, only some mPFC-NAc encoded foot shocks and promoted reward seeking under threat of punishment[6]. mPFC neurons projecting to the NAc shell play a larger role in suppression of reward seeking[79]. Given that our AAVrg targeted the NAc core, we likely captured a larger proportion of mPFC-NAc neurons involved in risk engagement rather than suppression. Our observation of PL-NAc neurons resembles the activity patterns seen in mPFC neurons projecting to dorsomedial striatum (DMS), which also increase their activity during conditioned tones and during risky exploration [80,81]. Indeed, we see that mPFC-NAc neurons send some axon collaterals in the DMS suggesting these populations overlap.

PL-VTA populations encoded learned but not innate threat avoidance behavior. Associative learning depends on the mPFC to encode predictive cues and on VTA to encode prediction errors[82]. PL-VTA neurons may regulate the separable actions of these two regions. To our knowledge, we are the first to directly study the role of mPFC-VTA neurons in threat avoidance. However, dopaminergic VTA cells projecting to mPFC have been widely studied and shown to robustly respond to aversive stimuli and facilitate associative learning[13,83–85]. mPFC-VTA neurons form synapses onto VTA cells projecting back to mPFC, including dopaminergic neurons[86] and may form part of a feedback loop influencing aversive learning.

In sum, we demonstrated the utility of pairing a standard approach like fiber photometry with whole-brain mapping to gain a detailed understanding of the relationships between function and connectivity. We show that PL-cPL, PL-NAc and PL-VTA neurons encode distinct features of innate and learned avoidance. Further, we mapped PL avoidance networks that that include discrete collateral targets of cPL, NAc and VTA projecting neurons. These whole-brain maps are an important resource for researchers studying the behavioral functions of mPFC. They can inform new experiments manipulating different PL pathways during threat avoidance and other disease-relevant behaviors. Further, our open-source analysis packages make brain wide anatomical approaches more accessible for integration into a variety of future research.

Methods

Animals Female and male C57B16/J mice (JAX Stock No. 000664) were group housed (2–5 per cage) and kept on a 12 hr light cycle. All animal procedures followed animal care guidelines approved by the University of California, Los Angeles Chancellor's Animal Research Committee.

Surgery Mice were induced in 5% isoflurane in oxygen until loss of righting reflex and transferred to a stereotaxic apparatus where they were maintained under 2% isoflurane in oxygen. Mice were warmed with a circulating water heating pad throughout surgery and eye gel was applied to the animal's eyes. The mouse's head was shaved and prepped with three scrubs of alternating betadine and then 70% ethanol. Following a small skin incision, a dental drill was used to drill through the skulls above the injection targets. A syringe pump (Kopf, 693A) with a hamilton syringe was used for injections. Injections were delivered at a rate 75nL/min and the syringe was left in the brain for 7 minutes following injection. For collateralization mapping, 300uL of AAVrg-Ef1a-mCherry-IRES-Cre-WPRE (Addgene 55632-AAVrg, 1.7×10^{13} vg/mL) was injected unilaterally into either left NAc (AP: 1.3, ML: -1.0, DV: 4.7), left VTA (AP: -3.3, ML: 0.4, DV: -4.5), or right PL (AP: 1.8, ML: -0.5, DV: -2.3). 200uL of AAV8-hSyn-DIO-hCHR2(H134R)-EYFP-WPRE (1.7×10^{13} vg/mL) was then injected into left PL (AP: 1.8, ML: -0.4, DV: -2.3). To control for Cre-independent expression of EYFP, control mice received a single injection of 200uL of AAV8-hSyn-DIO-hCHR2(H134R)-EYFP-WPRE into left PL. For fiber photometry from cPL and NAc-projectors, we injected AAVrg- Ef1a-mCherry-IRES-Cre-WPRE (same as above) or AAVrg-EF1a-mCherry-IRES-Flpo-WPRE (Addgene 55634-AAVrg, 1.7×10^{13} vg/mL) into either cPL or NAc, and after 2 weeks of expression injected AAV8-EF1a-Con-Foff 2.0-GCamp6m-WPRE into PL (Addgene 137120-AAV8). For recordings from PL-VTA neurons, we injected AAVrg- Ef1a-mCherry-IRES-Cre-WPRE (same as above) into VTA and AAV5-CAG-GCamp6m-WPRE (Addgene 100839-AAV5) into PL. 400uM optic fibers (Doric) were implanted into left PL (AP: 1.8, ML: -0.4, DV: -2.3) and sealed in place using Metabond (Patterson Dental Company, 5533559, 5533492, S371). For topographical mapping of the origin of each projection type, mice were injected with 300uL of cholera toxin subunit B (ThermoFisher, C34775, C34776, C34778) or Fluorogold (SCBT, C223769) at the same coordinates for cPL, NAc and VTA listed above. For pain management mice received 5mg/kg carprofen diluted in 0.9% saline subcutaneously. Mice received one injection during surgery and daily injections for two days following surgery. Samples with mistargeted injection sites were excluded from analysis. Samples with obviously poor antibody penetration or distribution following tissue clearing were also excluded.

Brain Slice Histology and Immunostaining Mice were transcardially perfused with phosphate-buffered saline (PBS) followed by 4% paraformaldehyde (PFA) in PBS. Brains were dissected, post-fixed in 4% PFA for 12–24h and placed in 30% sucrose for 24–48 hours. They were then embedded in Optimum Cutting Temperature (OCT, Tissue Tek) and stored at -80°C until sectioning. 60um floating sections were collected into PBS. Sections were washed 3x10min in PBS and then blocked in 0.3% PBST containing 10% normal donkey serum (JacksonImmunoresearch, 17-000-121) for 2h. Sections were then stained with chicken anti-GFP (AVES 1020 at 1:2000), rabbit anti-RFP (Rockland 600-401-379 at 1:2000) or rat anti-CTIP2 (Abcam ab18465, 1:200) in 0.3% PBST containing 3% donkey serum overnight at 4°C. The following day, sections were washed 3x5min in PBS and then stained with secondary antibody (JacksonImmunoresearch Cy2 donkey anti-chicken IgG(H+L) 703-225-155, 1:1000, Cy3 donkey anti-rabbit IgG(H+L) 711-005-152, 1:1000 or Alexa647 donkey anti-rat IgG(H+L) 702-605-150, 1:500) in 0.3% PBST containing 5% donkey serum for 2 hours at room temperature. Sections were then washed 5 min with PBS, 15 min with PBS+DAPI (Thermofisher Scientific, D1306, 1:4000), and then 5 min with PBS. Sections were mounted on glass slides using FluoroMount-G (ThermoFisher, 00-4958-02) and then imaged at 10x with a Leica STELLARIS confocal microscope.

Brain Clearing Mouse brains were collected and processed based on the published Adipo-Clear protocol[39] with slight modifications. Mice were perfused intracardially with 20mL of PBS (Gibco) followed by 4% paraformaldehyde (PFA, Electron Microscopy Sciences) on ice. Brains

were hemisected approximately 1mm past midline and postfixed overnight in 4% PFA at 4°C. The following day, samples were dehydrated with a gradient of methanol (MeOH, Fisher Scientific):B1n buffer (1:1,000 Triton X-100, 2% w/v glycine, 1:10,000 NaOH 10N, 0.02% sodium azide) for 1 hour for each step (20%, 40%, 60%, 80%) on a nutator (VWR). Samples were then washed with 100% MeOH 2x for 1hr each and then incubated in a 2:1 dichloromethane (DCM):MeOH solution overnight. The following day, two washes of 1hr in 100% DCM were performed followed by three washes of 100% MeOH for 30min, 45min then 1hr. Samples were bleached for 4 hours in 5:1 H2O2/MeOH buffer. A cascade of MeOH/B1n washes (80%, 60%, 40%, 20%) for 30min each rehydrated the samples followed by a 1h wash in B1n buffer. 5%DMSO/0.3M Glycine/PTxWH permeabilized tissue for one hour and then again for 2h with fresh solution. Samples were washed with PTxwH for 30min and then incubated in fresh PTxwH overnight. The following day two more PTxwH washes lasted 1h then 2h. Samples were incubated in primary GFP antibody (AVES Labs GFP 1020) at 1:2000 in PTxwH shaking at 37°C for 11, washed in PTxwH 2x1h and then 2x2h, then for two days with at least one PTxwH change per day while shaken at 37°C. Samples were then incubated in secondary antibody (AlexaFluor 647, ThermoFisher Scientific) for 8 days shaken at 37°C. Samples were washed in PTxwH 2x1h, then 2x2h, then 2 days with at least one PTxwH change per day while shaken at 37°C. Samples were then washed in 1x PBS twice 1x1hr, 2x2hr and then overnight. To dehydrate samples, a gradient of washes in MeOH:H2O (20%, 40%, 60% and 80%) were conducted for 30min each, followed by 3x100% MeOH for 30min, 1h, then 1.5h. Samples were incubated overnight in 2:1 DCM:MeOH on a nutator. The next day, samples were washed in 100% DCM 2x1h each. Samples were then cleared in 100% DBE. DBE was changed after 4h. Samples were stored in DBE in a dark place at room temperature. Imaging took place at least 24h after clearing.

Whole Brain Imaging Brain samples were imaged on a light-sheet microscope (Ultramicroscope II, LaVision Biotec) equipped with a sCMOS camera (Andor Neo) and a 2x/0.5 NA objective lens (MVPLAPO 2x) equipped with a 6 mm working distance dipping cap. Image stacks were acquired at 0.8x optical zoom using Imspector Microscope v285 controller software. For axons, we imaged using 488-nm (laser power 20%) and 640-nm (laser power 50%) lasers. The samples were scanned with a step-size of 3 µm using the continuous light-sheet scanning method with the included contrast adaptive algorithm for the 640-nm channel (20 acquisitions per plane), and without horizontal scanning for the 488-nm channel. For TRAP2-Ai14 brains, the 640-nm channel was imaged at 20% laser power without the contrast adaptive algorithm.

Model Training Light sheet images of fluorescently labeled axons were segmented using the 3D U-net based machine learning pipeline TRAILMAP (Friedmann et al., 2020). We trained new models for segmentation of cortical axons using inference learning as described in the TRAILMAP pipeline (https://github.com/AlbertPun/TRAILMAP). Briefly, 120x120 pixel stacks of 100 images were selected for training, and axons and artifacts (pixels that could be mistakenly interpreted as axons) were hand-labeled in 3 or 4 images from each stack. 1-pixel-wide edges of axons were labeled using python to be given less weight in training of the 3D convolutional network, which accounts for slight variability in human annotation patterns.

Models 1, 2, and 3 were designed for axon segmentation in regions with delineated, marginally distinguished, and indistinct axons, respectively. Beginning with the published TRAILMAP model weights, trainings were iteratively performed. Model 2 was trained first for 7 sessions and represents our best-trained single model for use across the entire brain. From this, Model 1 for use in superficial regions was trained for 2 sessions and Model 3 was trained for 4 sessions for use in deep regions.

Each training session used between 6 and 20 hand-labeled image stacks, of which approximately 10-20% were set aside for use in model validation. Training was performed using

between 20 and 120 steps per epoch and between 5 and 20 epochs per session. The best
model from each session was selected by plotting loss in validation data across epoch, which
was minimized to prevent overtraining. The weaknesses of the best model from each session
were assessed to guide selection of training data for the next session. Sessions were continued
until all visible axons in the desired regions were reliably detected by the model.
The TRAP2-Ai14 model was trained in the same manner. Initial weights were derived
from model that had been trained on fos$^+$ cells. From this, 6 sessions of training with 20 steps
per epoch and between 4 and 150 epochs per session were performed to generate the TRAP2-
Ai14 model.
661
Axon Model Validation We validated our analysis pipeline including image segmentation,
scaling and axon thinning by selecting 120x120 pixel stacks of 100 images from regions
assigned to the 'delineated', 'moderately distinguished', and 'indistinct' groups in brains that
were not included in initial training or validation data sets. Two human experts annotated 2–4
images from each stack, marking pixels likely to contain axons. In stacks with difficult-to-
distinguish axons, such as deep regions, experts used larger brush strokes to label broader
groups of pixels likely to contain axons (Figure S1B,C). One-pixel edges were then added to
human annotations in python.
The raw image stacks were segmented as explained below. In brief, each stack was
segmented with Models 1, 2 and 3 using TRAILMAP and scaled in ImageJ to a 10um space.
Scaled images were skeletonized in python at 8 different thresholds for binarization, combined
in MATLAB, and pixel values were adjusted in ImageJ. Using the same quantification threshold
as used in all analyses (number of skeletonized pixels with intensity above 64 divided by total
number of pixels in the region), labeling density was calculated in human-annotated "axon-
positive" pixels (LP) and "axon-negative" pixels (LN), excluding edges to account for slight
variability in human labels. Distinction score was calculated from these values as (LP-
LN)/LN+(1-LP). While direct true/false positive and true/false negative rates cannot be
calculated due to the axon thinning process, which reduces the number of labeled pixels, LP
correlates with the true positive rate, LN correlates with the false positive rate, and 1-LP
correlates with the false negative rate. Thus, LP-LN in the numerator will increase distinction
score when better separation is obtained between true and false positives, and LN and 1-LP in
the denominator will decrease the distinction score if there is a higher rate of false positives or
false negatives.
Alternate model combination methods were performed in FIJI. Maximum probability
projections between all models were calculated by generating images with maximum pixel
values across all models for each cube then processing the cubes for validation as above.
Summation of probability maps between models was calculated by adding probability calculated
by each model, then dividing by 3 prior to processing the cubes for validation as above.
Distinction scores from alternate approaches were compared using repeated measures one-
way ANOVA with Dunnett's multiple comparisons test, comparing individual model selection
(DeepTraCE) with each other method. Statistical comparisons of distinguishment scores were
performed using GraphPad PRISM.
694
TRAP2-Ai14 Model Validation To validate that cell counts produced by DeepCOUNT were
accurate, we selected 120x120x100 image cubes from several brain regions across multiple
brains not included in the training dataset. Cubes were scaled and processed in the same
manner as whole-brain data to produce a raw cell count from each cube. Two human experts
then manually counted cells in each cube. Average counts from the two human experts were
then compared to counts produced by DeepCOUNT as shown in Figure 4.
701
702 **DeepTraCE Analysis Pipeline**

Image Segmentation & Registration

Whole-brain image stacks from the 640nm channel from each brain were segmented three times in TRAILMAP, once each using models 1, 2, and 3. Following segmentation, the 488nm autofluorescence channel and axon segmentations from each model were converted to 8-bit and scaled to 10um resolution in FIJI with scaling values of 0.40625 in the x and y directions and 0.3 in the z direction using a bilinear interpolation algorithm. To improve image registration, each scaled 488nm image was manually rotated in the x, y, and z planes using the TransformJ ImageJ plugin (https://imagescience.org/meijering/software/transformj/) such that the midline blood vessels visible were all visible in the same z plane. The same manual rotation parameters were then applied to the three scaled model segmentation images from the corresponding brain.

The scaled and manually rotated 488nm autofluorescence channel was registered using elastix to the Gubra Lab LSFM atlas average template, which has annotations based on the Allen CCF[87–89]. The same transformation was applied to the scaled and manually rotated model segmentations using transformix. Image registration quality was manually verified by overlaying the atlas image and the registered 488nm channels in ImageJ. Following segmentation and registration of each model, the transformix images were converted to 8-bit .tif format in ImageJ. When combining multiple probability maps, it is important that minimum and maximum pixel values are comparable between images. Model 1 produced slightly lower maximum probability values compared to the other two models. We corrected for this by brightening segmentations from Model 1 by 34% to match pixel values obtained by the other models prior to combination and thinning. We provide a unified python pipeline for automating these steps in the supporting software repository.

Model Combination & Thinning

Converted segmentations from the three models were then combined by generating a new image in which pixel values in each region were extracted from the model with best performance in that region (Figure 2B). The same regional model assignments for combination were used for all brains in the data set and are provided in Table 2. Following model combination, axon segmentations were thinned (or 'skeletonized') as described in the TRAILMAP pipeline[29]. In brief, images were binarized at 8 different thresholds from 20 to 90% of the maximum intensity value using python. Skeletons were combined in MATLAB by summing values from each skeleton. Small objects unlikely to be axons were removed by calculating connected components within the combined skeleton and removing objects less than 90 voxels in size. Combined skeletons with small objects removed were optimized for visualization and quantification using an ImageJ macro that multiplied each pixel value by 17.

Axon Quantification

Regional axon innervation was quantified in MATLAB (Mathworks) by counting the number of skeletonized pixels in each brain region above a threshold (64), then dividing this pixel count by the total number of pixels in a region. This regional pixel count was then divided by the total number of labeled pixels across the brain to normalize for differences in total fluorescence and viral expression. Regions were defined by a collapsed version of the LSFM atlas in which maximum granularity was balanced with the need to account for slight differences in registration which would lead to inaccurate quantification of small brain regions. This atlas was cropped on the anterior and posterior ends to match the amount of tissue visible in our data. Fiber tracts, ventricular systems, cerebellum, and olfactory bulb were excluded from analysis.

All statistical comparisons were performed in MATLAB and GraphPad Prism v9. For comparison of regional axon labeling between cell types, Two-way ANOVA with Tukey's multiple comparison correction was performed on all brain regions. Anterior-posterior axon distributions within regions were calculated in MATLAB by binning the whole-brain image into

754 100um voxels and calculating the percentage of segmented pixels within each voxel. Voxels
755 falling within a given region were summed across the medial-lateral and dorsal-ventral axis and
756 normalized for total fluorescence as above. The averaged summation of axon counts from a
757 given cell class was then averaged and plotted along with the standard error of the mean.
758
Axon Visualization
To visualize axons as shown in Figures 1 and 3, Z-projections of raw light sheet data were
created in FIJI by scaling images to a 4.0625um space, virtually reslicing images in the coronal
plane, and performing maximum intensity z-projections of 100um depth followed by local
contrast enhancement. Axon segmentations were created by overlaying skeletonized and
registered axon segmentations from each sample of a cell type in a slightly different color and
virtually reslicing in the coronal plane. 3D projections were created in Imaris using a
representative sample from each cell type. Dotogram overlays (Figures 2 and 4) were created
using MATLAB. Images were binned into 100um voxels and the percentage of segmented
pixels within each voxel was calculated. Area of visible dot in the overlay corresponds with the
averaged labeling intensity within a voxel across a condition. Outer dots represent the cell type
with the highest labeling intensity within that voxel.
771
Computational Analyses
Hierarchical clustering of regional axon quantifications was performed as described in Kebschull
et al., 2020[36].
DeepCOUNT Analysis Pipeline
Image Segmentation & Registration
Whole-brain image stacks from the 640nm channel from each brain were segmented in
TRAILMAP using the TRAP2-Ai14 trained model. Images were registered using the 488nm
autofluorescence channel as described above. Transformed images were converted to 8-bit in
ImageJ.
781
782 *3D Maxima Detection & Single-Pixel Reduction*
783 MATLAB was used to identify 3D maxima of the transformed probability map. Connected
784 component analysis was then used to reduce any maxima that consisted of multiple pixels into a
785 single pixel per cell.
786
Cell Quantification
Regional TRAP$^+$ cell density was quantified in MATLAB (Mathworks) by counting the number of
labeled pixels (i.e. cells) in each brain region, then dividing this pixel count by the total number
of pixels in a region. This regional pixel count was then divided by the total number of detected
cells across the brain to normalize for differences in tamoxifen-induced recombination. Regions
were defined in the same way as described for the DeepTraCE pipeline.
793
794 **Functional Network Construction** Networks were constructed by thresholding inter-regional
795 correlations (Pearson's r≥0.9, *P*<0.05) for PMA and control groups. The nodes represent brain
796 region and the correlations above threshold were considered connections. Cytoscape software
797 was used to visualize networks. Node size was proportional to degree (number of connections).
798
4-Hydroxytamoxifen Preparation 4-hydroxytamoxifen (4-OHT; Sigma, Cat# H6278) was
dissolved at 20 mg mL^{-1} in ethanol by shaking at 37°C for 15 min and was then aliquoted and
stored at −20°C for up to several weeks. Before use, 4-OHT was redissolved in ethanol by
shaking at 37°C for 15 min, a 1:4 mixture of castor oil:sunflower seed oil (Sigma, Cat #s
259853 and S5007) was added to give a final concentration of 10 mg mL^{-1} 4-OHT, and the
ethanol was evaporated by vacuum under centrifugation. The final 10 mg mL^{-1} 4-OHT

solutions were always used on the day they were prepared. All injections were delivered
intraperitoneally (i.p.).
807
Behavioral assays For PMA experiments, mice were placed in an operant chamber with a
shock floor. The quarter of the floor furthest from the door of the chamber was covered in a
white plexiglass platform. Two odor pods with novel scents (vanilla, almond, coconut, or peanut
butter) were placed underneath the part of the shock floor not covered by the platform to
promote exploration.
For TRAP2 experiments, on training day, mice received 3 baseline tones (30s, 4000Hz,
75dB) followed by 9 tone-shock pairings (0.13mA shock, 2s, co-terminating), where mice could
learn to avoid the shock by entering the safety platform. Tones were separated by a random
interval between 80 and 150 seconds. The next day, mice received 6 tones with no shock. Mice
were injected with 4-OHT solution immediately following the retrieval session. Non-shock control
animals were placed in the operant chamber with no platform for five minutes on day 1, and on
day 2 were placed in the operant chamber for five minutes and injected with 4-OHT solution
immediately after this session.
For fiber photometry recordings during PMA, mice received 3 baseline tones on day 1,
followed by 12 tone-shock pairings on day 1 and 16 tone-shock pairings on days 2, and 3. For
EZM experiments, mice were placed on a custom-built elevated zero maze 24 inches in
diameter for 15 minutes. Both assays were recorded using a Point Grey Chameleon3 USB
camera (Teledyne FLIR).
826
827 **Fiber Photometry** Mice were habituated to the operant chamber and optic fiber for at least two
828 days prior to recording. A TDT RZ10x processor in combination with the TDT Synapse software
829 was used to simultaneously record the 405nm isosbestic channel and the 465nm signal channel
830 during behavior. For each mouse, the light output was adjusted such that the 465nm and 405nm
831 channel produced a signal of approximately 80mV as reported by the Synapse software.
832
Fiber Photometry Analysis Point-tracking of PMA videos were performed in DeepLabCut[90]
and behavior was analyzed using BehaviorDEPOT[91]. Elevated Zero Maze behavioral epochs
were annotated manually. Fiber photometry analysis was performed in MATLAB in a modified
version of an example provided by TDT. To align fiber photometry and behavioral data, TTL
pulses marking the beginning and end of each tone were aligned between the fiber photometry
signal and video frames. A lookup table was generated using linear interpolation between each
TTL pulse to identify which behavior frame lines up with each photometry frame. For alignment
of EZM fiber photometry data, a silent TTL pulse was generated every 30 seconds to be used
for alignment in the same manner.
Fiber photometry signal was down-sampled by a factor of 10 prior to analysis. To
account for potential movement artifacts and bleaching, the 405nm isosbestic control channel
was fit to the 465nm signal using the polyfit function, and this curve was then subtracted from
the 465nm signal. Z scores of this signal were calculated using a baseline period of -10 to 0
seconds relative to the tone for tone-aligned responses (i.e. tone and shock responses) and -20
to -15 seconds relative to epoch onset for all other behaviors (i.e. platform entries and exits,
closed arm entries and exits, head dips). The average of all traces for an individual animal was
calculated and used for analysis. To generate plots, each animal's average trace was smoothed
by averaging values from every 0.5 seconds (for time-locked tone and shock responses) or
using a moving average of 0.5 seconds (for all other traces), and the mean ± SEM of smoothed
traces across animals was displayed. To generate shock-normalized AUC values (provided in
Figure S8), the maximum shock response from an individual was calculated from the averaged
shock trace. All values in a trace were then divided by this value, and the resulting trace was
used to calculate the normalized AUC.

CODE AVAILABILITY
Code, instructions and sample data available at https://github.com/DeNardoLab/DeepTraCE
and https://github.com/jcouto/DeepTraCE/tree/gui

ACKNOWLEDGEMENTS
We thank Justus Kebschull and Xiaoyin Chen for helpful comments on the manuscript. This
work was funded by K01MH116264 (L.A.D.), 1R01MH127214-01A1 (L.A.D.), a Whitehall
Foundation Research Grant (L.A.D), a Klingenstein-Simons Foundation Grant (L.A.D.), a
NARSAD Young Investigator Award (L.A.D.), a Fay/Frank Seed Grant (L.A.D.), an NSERC
Postgraduate Fellowship (C.B.K.), T32MH073526-14 (M.W.G), and the UCLA Medical Scientist
Training Program (M.W.G.).

AUTHOR CONTRIBUTIONS
C.B.K, M.W.G. and L.A.D designed the research; C.B.K. performed the stereotaxic AAV
injections and PMA fiber photometry; C.B.K. and B.J. performed tissue clearing; C.B.K.,
M.W.G., and B.J. performed imaging and data analysis; D.F. and M.W.G. wrote DeepTraCE
and DeepCOUNT code; J.C. ported the pipeline to python and wrote the jupyter examples;
C.B.K, R.C., and A.S.E. performed histology and image analysis, A.S.E. and M.W.G performed
EZM fiber photometry; C.B.K, M.W.G. and L.A.D. wrote the manuscript.

DECLARATION OF INTERESTS
None to declare.

REFERENCES

1. Euston, D. R., Gruber, A. J. & McNaughton, B. L. The Role of Medial Prefrontal Cortex in Memory and Decision Making. *Neuron* **76**, 1057–1070 (2012).

2. Giustino, T. F. & Maren, S. The Role of the Medial Prefrontal Cortex in the Conditioning and Extinction of Fear. *Front. Behav. Neurosci.* **9**, 1–20 (2015).

3. Klune, C. B., Jin, B. & DeNardo, L. A. Linking mPFC circuit maturation to the developmental regulation of emotional memory and cognitive flexibility. *Elife* **10**, 1–33 (2021).

4. Jayachandran, M. *et al.* Prefrontal Pathways Provide Top-Down Control of Memory for Sequences of Events. *Cell Rep.* **28**, 640-654.e6 (2019).

5. Murugan, M. *et al.* Combined Social and Spatial Coding in a Descending Projection from the Prefrontal Cortex. *Cell* **171**, 1663-1677.e16 (2017).

6. Kim, C. K. *et al.* Molecular and Circuit-Dynamical Identification of Top-Down Neural Mechanisms for Restraint of Reward Seeking. *Cell* (2017) doi:10.1016/j.cell.2017.07.020.

7. Otis, J. M. *et al.* Prefrontal cortex output circuits guide reward seeking through divergent cue encoding. *Nature* (2017) doi:10.1038/nature21376.

8. Ye, L. *et al.* Wiring and Molecular Features of Prefrontal Ensembles Representing Distinct Experiences. *Cell* (2016) doi:10.1016/j.cell.2016.05.010.

9. Diehl, M. M. *et al.* Divergent projections of the prelimbic cortex bidirectionally regulate active avoidance. *Elife* **9**, 1–13 (2020).

10. Siciliano, C. A. *et al.* A cortical-brainstem circuit predicts and governs compulsive alcohol

901   drinking. *Science (80-.).* (2019) doi:10.1126/science.aay1186.

11. DeNardo, L. A., Berns, D. S., Deloach, K. & Luo, L. Connectivity of mouse somatosensory and prefrontal cortex examined with trans-synaptic tracing. *Nat. Neurosci.* **18**, 1687–1697 (2015).

12. Otis, J. M. *et al.* Prefrontal cortex output circuits guide reward seeking through divergent cue encoding. *Nature* **543**, 103–107 (2017).

13. Vander Weele, C. M. *et al.* Dopamine enhances signal-to-noise ratio in cortical-brainstem encoding of aversive stimuli. *Nature* **563**, 397–401 (2018).

14. Warden, M. R. *et al.* A prefrontal cortex-brainstem neuronal projection that controls response to behavioural challenge. *Nature* (2012) doi:10.1038/nature11617.

15. Riga, D. *et al.* Optogenetic dissection of medial prefrontal cortex circuitry. *Frontiers in Systems Neuroscience* (2014) doi:10.3389/fnsys.2014.00230.

16. Mathis, V. P., Williams, M., Fillinger, C. & Kenny, P. J. Networks of habenula-projecting cortical neurons regulate cocaine seeking. *Sci. Adv.* **7**, (2021).

17. Kim, Y. *et al.* Mapping Social Behavior-Induced Brain Activation at Cellular Resolution in the Mouse. *Cell Rep.* **10**, 292–305 (2015).

18. Gao, L. *et al.* Single-neuron projectome of mouse prefrontal cortex. *Nat. Neurosci.* **25**, 515–529 (2022).

19. Zhang, F. *et al.* The microbial opsin family of optogenetic tools. *Cell* (2011) doi:10.1016/j.cell.2011.12.004.

20. Roth, B. L. DREADDs for Neuroscientists. *Neuron* (2016) doi:10.1016/j.neuron.2016.01.040.

21. Dana, H. *et al.* High-performance calcium sensors for imaging activity in neuronal populations and microcompartments. *Nat. Methods* (2019) doi:10.1038/s41592-019-0435-6.

22. Gabbott, P. L. A., Warner, T. A., Jays, P. R. L., Salway, P. & Busby, S. J. Prefrontal Cortex in the Rat⬜: Projections to Subcortical Autonomic , Motor , and Limbic Centers. *J. Comp. Neurol.* **177**, 145–177 (2005).

23. Ishikawa, A. & Nakamura, S. Ventral hippocampal neurons project axons simultaneously to the medial prefrontal cortex and amygdala in the rat. *J. Neurophysiol.* **96**, 2134–2138 (2006).

24. Ishikawa, A., Ambroggi, F., Nicola, S. M. & Fields, H. L. Dorsomedial Prefrontal Cortex Contribution to Behavioral and Nucleus Accumbens Neuronal Responses to Incentive Cues. *J. Neurosci.* **28**, 5088 LP – 5098 (2008).

25. Kebschull, J. M. *et al.* High-Throughput Mapping of Single-Neuron Projections by Sequencing of Barcoded RNA. *Neuron* **91**, 975–987 (2016).

26. Oh, S. W. *et al.* A mesoscale connectome of the mouse brain. *Nature* **508**, 207–214 (2014).

27. Chung, K. *et al.* Structural and molecular interrogation of intact biological systems. *Nature* **497**, 332–337 (2013).

28. Renier, N. *et al.* Mapping of Brain Activity by Automated Volume Analysis of Immediate
Early Genes. *Cell* **165**, 1789–1802 (2016).

29. Friedmann, D. *et al.* Mapping mesoscale axonal projections in the mouse brain using a
3D convolutional network. *Proc. Natl. Acad. Sci.* (2020) doi:10.1073/pnas.1918465117.

30. Berg, S. *et al.* ilastik: interactive machine learning for (bio)image analysis. *Nat. Methods*
16, 1226–1232 (2019).

31. Capuzzo, G. & Floresco, S. B. Prelimbic and Infralimbic Prefrontal Regulation of Active
and Inhibitory Avoidance and Reward-Seeking. *J. Neurosci.* **40**, 4773–4787 (2020).

32. Bravo-Rivera, C., Roman-Ortiz, C., Brignoni-Perez, E., Sotres-Bayon, F. & Quirk, G. J.
Neural Structures Mediating Expression and Extinction of Platform-Mediated Avoidance.
J. Neurosci. (2014) doi:10.1523/JNEUROSCI.0191-14.2014.

33. Stern, C. A. J., Do Monte, F. H. M., Gazarini, L., Carobrez, A. P. & Bertoglio, L. J. Activity
in prelimbic cortex is required for adjusting the anxiety response level during the elevated
plus-maze retest. *Neuroscience* **170**, 214–222 (2010).

34. Lui, J. H. *et al.* Differential encoding in prefrontal cortex projection neuron classes across
cognitive tasks. *Cell* **184**, 489-506.e26 (2021).

35. Kramer, D. J. *et al.* Generation of a DAT-P2A-Flpo mouse line for intersectional genetic
targeting of dopamine neuron subpopulations. *Cell Rep.* **35**, 109123 (2021).

36. Kebschull, J. M. *et al.* Cerebellar nuclei evolved by repeatedly duplicating a conserved
cell-type set. *Science (80-.).* **370**, (2020).

37. Schwarz, L. A. *et al.* Viral-genetic tracing of the input-output organization of a central
noradrenaline circuit. *Nature* **advance on**, (2015).

38. Tervo, D. G. R. *et al.* A Designer AAV Variant Permits Efficient Retrograde Access to
Projection Neurons. *Neuron* (2016) doi:10.1016/j.neuron.2016.09.021.

39. Chi, J., Crane, A., Wu, Z. & Cohen, P. Adipo-Clear: A Tissue Clearing Method for Three-
Dimensional Imaging of Adipose Tissue. *J. Vis. Exp.* (2018) doi:10.3791/58271.

40. Chi, J. *et al.* Three-Dimensional Adipose Tissue Imaging Reveals Regional Variation in
Beige Fat Biogenesis and PRDM16-Dependent Sympathetic Neurite Density. *Cell Metab.*
27, 226-236.e3 (2018).

41. Sun, L. *et al.* Differences in neurotropism and neurotoxicity among retrograde viral
tracers. *Mol. Neurodegener.* **14**, 1–24 (2019).

42. Leow, Y. N. *et al.* Brain-wide mapping of inputs to the mouse lateral posterior
(LP/Pulvinar) thalamus–anterior cingulate cortex network. *J. Comp. Neurol.* **530**, 1992–
2013 (2022).

43. Arlotta, P. *et al.* Neuronal subtype-specific genes that control corticospinal motor neuron
development in vivo. *Neuron* **45**, 207–221 (2005).

44. Anastasiades, P. G. & Carter, A. G. Circuit organization of the rodent medial prefrontal
cortex. *Trends Neurosci.* **44**, 550–563 (2021).

45. Anastasiades, P. G., Boada, C. & Carter, A. G. Cell-Type-Specific D1 Dopamine
Receptor Modulation of Projection Neurons and Interneurons in the Prefrontal Cortex.

Cereb. Cortex **29**, 3224–3242 (2019).

46. McGarry, L. M. & Carter, A. G. Inhibitory gating of basolateral Amygdala inputs to the prefrontal cortex. *J. Neurosci.* **36**, 9391–9406 (2016).

47. Collins, D. P., Anastasiades, P. G., Marlin, J. J. & Carter, A. G. Reciprocal Circuits Linking the Prefrontal Cortex with Dorsal and Ventral Thalamic Nuclei. *Neuron* **98**, 366-379.e4 (2018).

48. DeNardo, L. A. *et al.* Temporal evolution of cortical ensembles promoting remote memory retrieval. *Nat. Neurosci.* **22**, 460–469 (2019).

49. Sesack, S. R., Deutch, A. Y., Roth, R. H. & Bunney, B. S. Topographical organization of the efferent projections of the medial prefrontal cortex in the rat: An anterograde tract-tracing study with Phaseolus vulgaris leucoagglutinin. *J. Comp. Neurol.* (1989) doi:10.1002/cne.902900205.

50. Vertes, R. P. Differential Projections of the Infralimbic and Prelimbic Cortex in the Rat. *Synapse* **51**, 32–58 (2004).

51. Winnubst, J. *et al.* Reconstruction of 1,000 Projection Neurons Reveals New Cell Types and Organization of Long-Range Connectivity in the Mouse Brain. *Cell* **179**, 268-281.e13 (2019).

52. Harris, K. D. & Shepherd, G. M. G. The neocortical circuit: Themes and variations. *Nat. Neurosci.* **18**, 170–181 (2015).

53. Poo, C., Agarwal, G., Bonacchi, N. & Mainen, Z. F. Spatial maps in piriform cortex during olfactory navigation. *Nature* **601**, 595–599 (2022).

54. Gehrlach, D. A. *et al.* A whole-brain connectivity map of mouse insular cortex. *Elife* **9**, e55585 (2020).

55. Sheng, M. & Greenberg, M. E. The regulation and function of c-fos and other immediate early genes in the nervous system. *Neuron* **4**, 477–485 (1990).

56. Franceschini, A., Costantini, I., Pavone, F. S. & Silvestri, L. Dissecting Neuronal Activation on a Brain-Wide Scale With Immediate Early Genes. *Front. Neurosci.* **14**, 569517 (2020).

57. Wheeler, A. L. *et al.* Identification of a Functional Connectome for Long-Term Fear Memory in Mice. *PLoS Comput. Biol.* **9**, e1002853 (2013).

58. Vetere, G. *et al.* Chemogenetic Interrogation of a Brain-wide Fear Memory Network in Mice. *Neuron* **94**, 363–374 (2017).

59. Diehl, M. M. *et al.* Active avoidance requires inhibitory signaling in the rodent prelimbic prefrontal cortex. *Elife* (2018) doi:10.7554/eLife.34657.

60. Bravo-Rivera, C., Roman-Ortiz, C., Montesinos-Cartagena, M. & Quirk, G. J. Persistent active avoidance correlates with activity in prelimbic cortex and ventral striatum. *Front. Behav. Neurosci.* (2015) doi:10.3389/fnbeh.2015.00184.

61. Nikolaev, E., Werka, T. & Kaczmarek, L. C-fos protooncogene expression in rat brain after long-term training of two-way active avoidance reaction. *Behav. Brain Res.* **48**, 91–94 (1992).

62. Duncan, G. E., Knapp, D. J. & Breese, G. R. Neuroanatomical characterization of Fos
induction in rat behavioral models of anxiety. *Brain Res.* **713**, 79–91 (1996).

63. Allen, W. E. *et al.* Thirst-associated preoptic neurons encode an aversive motivational
drive. *Science (80-.).* **357**, 1149–1155 (2017).

64. Madisen, L. *et al.* A robust and high-throughput Cre reporting and characterization
system for the whole mouse brain. *Nat. Neurosci.* **13**, 133–140 (2010).

65. Fenno, L. E. *et al.* Comprehensive Dual- and Triple-Feature Intersectional Single-Vector
Delivery of Diverse Functional Payloads to Cells of Behaving Mammals. *Neuron* (2020)
doi:10.1016/j.neuron.2020.06.003.

66. Economo, M. N. *et al.* Distinct descending motor cortex pathways and their roles in
movement. *Nature* **563**, 79–84 (2018).

67. Kebschull, J. M. *et al.* High-Throughput Mapping of Single-Neuron Projections by
Sequencing of Barcoded RNA. *Neuron* **91**, 975–987 (2016).

68. Senn, V. *et al.* Long-range connectivity defines behavioral specificity of amygdala
neurons. *Neuron* **81**, 428–437 (2014).

69. Ährlund-Richter, S. *et al.* A whole-brain atlas of monosynaptic input targeting four
different cell types in the medial prefrontal cortex of the mouse. *Nat. Neurosci.* **22**, 657–
668 (2019).

70. Wang, L. *et al.* Cell-Type-Specific Whole-Brain Direct Inputs to the Anterior and Posterior
Piriform Cortex . *Frontiers in Neural Circuits* vol. 14 (2020).

71. Peckford, G. *et al.* The effects of lesions to the postsubiculum or the anterior dorsal
nucleus of the thalamus on spatial learning in rats. *Behav. Neurosci.* **128**, 654–665
(2014).

72. Bett, D., Wood, E. R. & Dudchenko, P. A. The postsubiculum is necessary for spatial
alternation but not for homing by path integration. *Behav. Neurosci.* **126**, 237–248
(2012).

73. Bett, D. *et al.* The Postsubiculum and Spatial Learning: The Role of Postsubicular
Synaptic Activity and Synaptic Plasticity in Hippocampal Place Cell, Object, and Object-
Location Memory. *J. Neurosci.* **33**, 6928–6943 (2013).

74. Sacco, T. & Sacchetti, B. Role of secondary sensory cortices in emotional memory
storage and retrieval in rats. *Science (80-.).* **329**, 649–656 (2010).

75. Costa, N. S., Vicente, M. A., Cipriano, A. C., Miguel, T. T. & Nunes-de-Souza, R. L.
Functional lateralization of the medial prefrontal cortex in the modulation of anxiety in
mice: Left or right? *Neuropharmacology* **108**, 82–90 (2016).

76. Stevenson, C. W., Halliday, D. M., Marsden, C. A. & Mason, R. Early life programming of
hemispheric lateralization and synchronization in the adult medial prefrontal cortex.
Neuroscience **155**, 852–863 (2008).

77. Sullivan, R. M. & Gratton, A. Lateralized effects of medial prefrontal cortex lesions on
neuroendocrine and autonomic stress responses in rats. *J. Neurosci.* **19**, 2834–2840
(1999).

78. Lee, E. *et al.* Left brain cortical activity modulates stress effects on social behavior. *Sci.*

Rep. **5**, 13342 (2015).

79.   Piantadosi, P. T., Yeates, D. C. M. & Floresco, S. B. Prefrontal cortical and nucleus
accumbens contributions to discriminative conditioned suppression of reward-seeking.
Learn. Mem. **27**, 429–440 (2020).

80.   Loewke, A. C., Minerva, A. R., Nelson, A. B., Kreitzer, A. C. & Gunaydin, L. A.
Frontostriatal Projections Regulate Innate Avoidance Behavior. *J. Neurosci.* **41**, 5487–
5501 (2021).

81.   Kajs, B. L., Loewke, A. C., Dorsch, J. M., Vinson, L. T. & Gunaydin, L. A. Divergent
encoding of active avoidance behavior in corticostriatal and corticolimbic projections. *Sci.*
Rep. **12**, 1–11 (2022).

82.   Lak, A. *et al.* Dopaminergic and Prefrontal Basis of Learning from Sensory Confidence
and Reward Value. *Neuron* **105**, 700-711.e6 (2020).

83.   Kim, C. K. *et al.* Simultaneous fast measurement of circuit dynamics at multiple sites
across the mammalian brain. *Nat. Methods* **13**, 325–328 (2016).

84.   Puig, M. V., Antzoulatos, E. G. & Miller, E. K. Prefrontal dopamine in associative learning
and memory. *Neuroscience* **282**, 217–229 (2014).

85.   Gonzalez, M. C. *et al.* Medial prefrontal cortex dopamine controls the persistent storage
of aversive memories. *Front. Behav. Neurosci.* (2014) doi:10.3389/fnbeh.2014.00408.

86.   Sesack, S. R. & Carr, D. B. Selective prefrontal cortex inputs to dopamine cells:
implications for schizophrenia. *Physiol. Behav.* **77**, 513–517 (2002).

87.   Klein, S., Staring, M., Murphy, K., Viergever, M. A. & Pluim, J. P. W. Elastix: A toolbox for
intensity-based medical image registration. *IEEE Trans. Med. Imaging* (2010)
doi:10.1109/TMI.2009.2035616.

88.   Wang, Q. *et al.* The Allen Mouse Brain Common Coordinate Framework: A 3D Reference
Atlas. *Cell* **181**, 936-953.e20 (2020).

89.   Perens, J. *et al.* An Optimized Mouse Brain Atlas for Automated Mapping and
Quantification of Neuronal Activity Using iDISCO+ and Light Sheet Fluorescence
Microscopy. *Neuroinformatics* **19**, 433–446 (2021).

90.   Mathis, A. *et al.* DeepLabCut: markerless pose estimation of user-defined body parts with
deep learning. *Nat. Neurosci.* **21**, (2018).

91.   Gabriel, C. J. *et al.* Behavior DEPOT is a simple, flexible tool for automated behavioral
detection based on marker less pose tracking. *Elife* **11**, 1–33 (2022).

1094

Figure 1. DeepTraCE workflow

(A) Overview of DeepTraCE workflow

(B) Demonstration of segmentation using 3 different models followed by model combination

(C) Overlay of raw data and axon segmentation using single model or DeepTraCE concatenated models

(D) Distinction score produced by models 1, 2, and 3 for images from regions assigned to axon groups 1, 2, and 3. Repeated-measures one-way ANOVA with Benjamini, Krieger and Yekutieli post-hoc test. (Group 1: *F*=8.503, *P*=0.0257; Group 2: *F*=7.753, *P*=0.0309; Group 3: *F*=73.76, *P*=0.0001).

(E) Comparison of DeepTraCE with alternate segmentation and model combination strategies. Repeated measures one-way ANOVA with Benjamini, Krieger and Yekutieli post-hoc test. (*F*=14.46, *P*<0.0001). See Table 3 for detailed statistics.

Scale bars, 200um. Descriptive statistics: mean ± SEM. *P<0.05, **P<0.01, ***P<0.001. Distinction score: (LP-LN) / (LN+(1-LP)). LP = axon+ human label. LN = axon- human label.

Figure 2. Visualization and quantification of brain-wide projection patterns of mPFC

(A–C) Viral injection strategy and DeepTraCE segmentation of superficial (left) and deep (right) brain structures for each cell type. (D–F) Single 10um coronal optical sections of thinned axons registered to the standardized brain atlas. Represents 4 overlaid brains from each neuronal class. (G) Dotogram overlay of 3 cell types. Innervation by cPL-, NAc- and VTA-projecting PL neurons shown in yellow, blue and purple, respectively. (H) Left heatmap: relative labeling density (normalized to region volume and gross label content per brain) across 140 regions defined by the Allen Brain Atlas. Middle heatmap: P values from multiple comparisons of axon innervation density. Right heatmap: Loadings for PC1 (arbitrary PC weight units) (I) Dendrogram of hierarchical clustering of regional axon quantifications from 12 mice, colored by target region. (J) Locations of individual mice projected in principal component (PC) space defined by the first two PCs (arbitrary PC units, cPL, *n*=4; NAc, *n*=4; VTA, *n*=4). Abbreviations: Isoctx, isocortex; OLF, olfactory areas; HPF, hippocampal formation; CTXsp, cortical subplate; CNU, cerebral nuclei; HY, hypothalamus; MB, midbrain; HB, hindbrain. See Tables 1–3 and Figure S2-6 for related data.

Figure 3. Region-specific collateralization patterns of PL-cPL, PL-NAc and PL-VTA neurons

(A) Coronal view of 100um z-projections of raw 640nm (axons, white) and 488nm (autofluorescence, blue) channels from individual brains. Images show cell-type specific innervation patterns in anterior (left) and posterior (right) cortical areas. (B) Layer distributions of axonal innervation in select cortical target regions. (C) Visualizations of axonal innervation density along the anterior-posterior axis in select regions. (D) Raw images showing axons in the amygdalar complex. (E) Quantification of axonal innervation density along the anterior- posterior axis of a given region. PL-cPL, PL-NAc and PL-VTA neurons coded in yellow, blue and purple, respectively. N=4/group. Descriptive statistics are from 2-way ANOVA with Tukey's post-hoc multiple comparison test. See Table 1,3 for abbreviations and detailed statistics. Error bars: mean ± S.E.M. *P<0.05, **P<0.01.

Figure 4. Whole-brain IEG mapping of learned avoidance with DeepCOUNT.
(A) Overview of DeepCOUNT workflow (B) Overlay of raw data and axon segmentation using trained cell detection model.
(C) Upper: Correlation between human vs. DeepCOUNT or ClearMap neuronal counts in image volumes (Pearson's
R^2_{DC} = 0.99, P<0.0001; R^2_{CM} = 0.96, P<0.0001; Lower: Comparison of residual error values for DeepCOUNT and
ClearMap as compared to human annotations (P=0.022, n=6 brain regions; paired t-test). (D) Schematic of platform
mediated avoidance assay. TRAP2;Ai14 mice were trained in PMA and then TRAPed during a retrieval test the next day.
(E) Left heatmap: relative cell density (normalized to region volume and gross label content per brain) across 140 regions
defined by the Allen Brain Atlas. Right heatmap: Loadings for PC1 (arbitrary PC weight units). (F) Locations of individual
mice projected in principal component (PC) space defined by the first two PCs (arbitrary PC units, control, n=5; PMA,
n=4). (G) Dotogram overlay of control (grey) and PMA (green) conditions. Dots represent the density of TRAPed cells in
a given voxel. (H) Comparison of TRAPed cell density in 5 brain regions of naïve controls (grey) and PMA-trained mice
(green) (Student's t-test; *P<0.05, **P<0.01; control: n=5; PMA: n=4). (I) Network diagrams for control (left) and PMA-
trained (right) mice based on brain wide interregional correlations. Node size is proportional to degree. (J) Degree values
for top 20 most connected regions for each condition (PL outlined in red). See Figure S6 for related data.

Figure 5. Neuronal class-specific activity during threat cues and aversive stimuli

(A) GCaMP injection strategy. Left: to target cPL-projecting neurons, mice were injected with AAVrg-Cre in cPL and AAVrg-Flp in NAc, and Con-Foff-GCaMP6M in PL. Middle: to target NAc-projecting neurons, mice were injected with AAVrg-Flp in cPL and AAVrg-Cre in NAc, and Con-Foff-GCaMP6M in PL. Right: to target VTA-projecting neurons, mice were injected with AAVrg-Cre in VTA and DIO-GCaMP6M in PL. (B) Representative images of GCaMP expression and fiber placement sites in PL. Scale bar, 1mm. (C) Distribution of cPL-, NAc- and VTA-projecting GCaMP-expressing cells across cortical layers (F_{bin} = 28.3, $P<0.0001$, $F_{binxclass}$ = 34.09, $P<0.0001$; cPL: n=6, NAc: n=5, VTA: n =6; 2-way ANOVA). (D) Representative images of axon terminals in cPL and NAc using the intersectional viral targeting strategy shown in A. Scale bars, 100um. (E) Schematic of PMA assay and behavioral performance across sessions (F_{day} = 65.18, P<0.0001, F_{class} = 0.58, P=0.57, $F_{class\,x\,day}$ = 0.43, P=0.86, 2-way ANOVA). (F) GCaMP fluorescence in cPL-, NAc-, and VTA-projecting neurons. Signals are aligned to tone onset and separated by whether mouse is on (colored trace) or off (grey trace) the safety platform at the start of the tone. *$P<0.05$ for student's t-test comparing on vs. off platform activity in a given time window. (G) AUC analysis of Ca^{2+} signal for tone periods (0–10s) when mice were off the platform (F = 4.106, P=0.04, cPL: n=6, NAc: n=5, VTA: n =6; One-way ANOVA with Benjamini, Krieger and Yekutieli post-hoc test. (H) GCaMP fluorescence in cPL-, NAc- and VTA-projecting PL cells. Signals are aligned to shock onset and separated by whether mouse is on (colored trace) or off (grey trace) the safety platform. *$P<0.05$, **$P<0.01$ for student's t-test comparing on vs. off platform activity in a given time window. (I) Analysis of Ca^{2+} signals for circa-shock periods (Pre-shock AUC: F = 6.974, P=0.0079; Shock AUC: F = 1.868, P=0.19; Post-shock AUC (10–40s): F = 8.28, P=0.0042; Post-shock AUC (80–80s): F = 8.79, P=0.0034; cPL: n=6, NAc: n=5, VTA: n =6, One-way ANOVA with Benjamini, Krieger and Yekutieli post-hoc test).

Figure 6. Neuronal class-specific activity during learned vs. innate threat avoidance behavior
(A,B) GCaMP fluorescence aligned to safe-zone entry in learned (A) vs. innate (B) avoidance for cPL-projecting (yellow), NAc-projecting (blue) and VTA-projecting (purple) neurons. Inset plots show AUC analysis of Ca^{2+} signal aligned to avoidance (-5–5s) (F = 4.106, P=0.04, One-way ANOVA with Benjamini, Krieger and Yekutieli post-hoc test).
(C,D) GCaMP fluorescence aligned to onset of risky exploration in learned (C) vs. innate (D) avoidance for cPL-projecting (yellow), NAc-projecting (blue) and VTA-projecting (purple) neurons. Inset plots show AUC analysis of Ca^{2+} signal aligned to exploration onset (0–20s) (F = 4.106, P=0.04, One-way ANOVA with Benjamini, Krieger and Yekutieli post-hoc test).
(E) Analysis of Ca^{2+} signals during head dips (F = 0.64, P=0.55; PMA: cPL n=6, NAc n=5, VTA n=6; EZM: cPL n=5, NAc n=4, VTA n=3; One-way ANOVA with Benjamini, Krieger and Yekutieli post-hoc test.).

Figure 7. Summary of Findings

(A) Visualization of collateralization density in key targets of PL–cPL, PL–NAc, and PL–VTA neurons. Dot radius correlates with average normalized labeling density within a region.

(B) Schematic of whole-brain collateralization patterns of PL–cPL, PL–NAc, and PL–VTA neurons.

(C) Summary of PL efferent connectivity patterns.

(D) Visualization of activity levels in PL–cPL, PL–NAc, and PL–VTA neurons during aspects of threat avoidance. Dot radius correlates with average normalized signal intensity.

See Table 1 for abbreviations.