TITLE

Linking hubness, embryonic neurogenesis, transcriptomics and diseases in
human brain networks

AUTHORS

Ibai Diez*[1,2], Fernando Garcia-Moreno*[3,4,5], Nayara Carral-Sainz[6],
Sebastiano Stramaglia[7], Alicia Nieto-Reyes[8], Mauro D'Amato[5,9,10], Jesús
Maria Cortes[5,11,12], Paolo Bonifazi[5,11]

AFFILIATIONS

[1]Gordon Center for Medical Imaging, Department of Radiology, Massachusetts General Hospital, Harvard
Medical School, Boston, MA, USA.
[2]Athinoula A. Martinos Center for Biomedical Imaging, Department of Radiology, Massachusetts General
Hospital, Harvard Medical School, Charlestown, MA, USA.
[3]Achucarro Basque Center for Neuroscience, Scientific Park of the University of the Basque Country
(UPV/EHU), Leioa, Spain.
[4]Department of Neuroscience, Faculty of Medicine and Odontology, UPV/EHU, Barrio Sarriena s/n, Leioa,
Bizkaia, Spain.
[5]IKERBASQUE: The Basque Foundation for Science, Bilbao, Spain.
[6]Departamento de Ciencias de la Tierra y Física de la Materia Condensada, Facultad de Ciencias,
Universidad de Cantabria, Santander, Spain.
[7]Dipartimento Interateneo di Fisica, Università degli Studi di Bari Aldo Moro, and INFN, Sezione di Bari,
Italy.
[8]Departamento de Matemáticas, Estadística y Computación, Facultad de Ciencias, Universidad de
Cantabria, Santander, Spain.
[9]Department of Medicine and Surgery, LUM University, Casamassima, Italy.
[10]Gastrointestinal Genetics Lab, CIC bioGUNE - BRTA, Derio, Spain.
[11]Computational Neuroimaging Lab, Biocruces-Bizkaia Health Research Institute, Barakaldo, Spain.
[12]Department of Cell Biology and Histology, University of the Basque Country (UPV/EHU), Leioa, Spain.
28
29 ** These authors contributed equally to this work.*

30 *Corresponding author: paol.bonifazi@gmail.com*

31
SUMMARY

Understanding the architectural principles that shape human brain networks is a major

challenge for systems neuroscience. We hypothesize that the centrality of the different

brain circuits in the human connectome is a product of their embryogenic age, such that

early-born nodes should become stronger hubs than those born later. Using a human

brain segmentation based on embryogenic age, we observed that nodes' structural

centrality correlated with their embryogenic age, fully confirming our hypothesis.

Distinct trends were found at different resolutions on a functional level. The difference

in embryonic age between nodes inversely correlated with the probability of existence

of links and their weights. Brain transcriptomic analysis revealed strong associations

between embryonic age, structure-function centrality, and the expression of genes

related to nervous system development, synapse regulation and human neurological

diseases. Our results highlight two key principles regarding the wiring of the human

brain, "preferential age attachment" and "the older gets richer".

MAIN TEXT

The most characteristic anatomical property of brain networks is their organization

across multiple spatial scales. Micro-scale neurons are interconnected within local

circuits or microdomains, which are scaled up to the macro-scale over which long-range

connections allow distant neurons to communicate and different circuits to interact[1,2].

Consequently, a key challenge in fundamental and clinical neuroscience is to decipher

the rules of connectivity that shape brain networks in order to understand how the brain

works and how traumatic or neurological damage may affect brain functionality[3].

The general structure and function of the human brain, and its internal connectivity are

all the result of its developmental history, which is at the same time the product of

evolution[4].

As such, the brain follows developmental instructions to establish a blueprint of diverse

neural connections, the functioning of which can be later refined.

Previous studies have evaluated the development of brain networks at different stages

of maturation, particularly in relation to the appearance of key network structures like

central nodes[5]. However, there are still no detailed studies that fully relate the

connectivity of the adult human brain with the sequential (evolutionarily preserved)

neurogenesis of circuits.

Complex networks have emerged over the past two decades and provide a powerful

mathematical framework to quantify the complexity of brain circuitries.

Different studies that assessed the data from real-world networks' have highlighted that

small-world, scale-free or heavy-tailed distribution network organizations are

stereotypic topologies of many distinct domains([6,7]).

These key topologies have been identified at the structural-functional level in

microcircuits, and in meso- and macro-scale networks([8]).

A pioneering study([9]) showed that the principle "the rich gets richer" (a.k.a. "preferential

attachment") led to the development of the scale-free networks that are characterized

by highly connected rare nodes or hubs.

Inspired by this model([9]), here we aim to combine the heavy-tailed signatures widely

observed in brain networks' with the stereotypic, evolutionary-preserved sequential

neurogenesis in the developing brain. In this way, we will test the hypothesis that the

topology of brain networks could be shaped according to the rule that "the older gets

richer", i.e. the evolutionary older circuits or those generated earlier in embryogenesis

are those most central in the organization of the adult brain network([10]).

As a consequence, quantification of the hubness of brain circuits based on metrics of

centrality in complex networks should be correlated with embryogenic age. Our

hypothesis extrapolates on the macro-scale the earlier pioneering evidence that

GABAergic neuronal hubs were functionally identified in hippocampal circuits on a

micro-circuit level, based on early born GABAergic neurons in both developing and adult

murine circuits([11–13]).

To test our hypothesis, we differentiated and segmented human brain circuits according

to their *First neurogenic Time (FirsT)* following the criteria established in the methods,

i.e. the post-conception day (or embryonic day) on which the first neurons of the circuit

are generated in the corresponding developmental neuromere (Fig. 1A). The match

between developmental neuromeres and adult circuits, and the estimation of FirsT in

humans extrapolated from different animal species, is presented with underlying

literature and full references in the Method section. As such, a list was generated that

organized the human circuits according to their FirsT, from older to younger circuits

(Table 1). We identified 18 distinct macro-circuits (MACs) corresponding to 18 separated

fundamental developmental units for which a timing sequence based on their FirsT

could be defined. Other relevant circuits for which detailed developmental information

is known could not be segmented on MRI, such as the neocortical layers and thalamic

nuclei (see Supplementary Information).

We used dMRI and resting-state fMRI images acquired at 7T from N=184 healthy

subjects taken from the Human Connectome project to reconstruct structural-

functional brain networks (see Fig. 1B and Methods). Since the volume of the different

MACs spanned more than two orders of magnitude (see Supplementary Fig. 1B), the

MACs were also parcellated (see Methods) to obtain spatially segregated ROIs with a

volume comparable to the smallest circuits (e.g., the Locus Coeruleus -LC), each in the

range of several dozen voxels (see Supplementary Fig. 1A for the overall ROI volume

distribution and Table 1 for the number of ROIs per MAC).

Each of the 18 MACs had a corresponding FirsT, which was also assigned to all ROIs

obtained for a given MAC, and a total of 2,566 ROIs were defined in the brain. To explore

the possible existence of patterns of correlation between FirsT and brain circuit

connectivity at different spatial resolutions, eigenvector centrality (EC) was calculated

for each ROI and MAC (Fig. 1B; see Methods for the rationale behind the choice of

metrics). In addition, the brain maps of the transcriptome of 20787 coding-genes were

related to the brain maps of neurogenesis (FirsT) and nodes' hubness (Fig. 1C) in order

to obtain a complementary biological correlate to the brain connectivity model.

To quantify the centrality of each ROI and MAC in the brain networks, we calculated EC

on high spatial resolution networks composed of the 2,566 ROIs (referred to as high-

resolution structural and functional connectivity networks, SC^{HR} and FC^{HR}) and low

spatial resolution networks composed of the 18 MACs (referred to as low-resolution

structural and functional connectivity networks, SC^{LR} and FC^{LR}).

The high-resolution networks were calculated from the average of all the subjects,

based on probabilistic tractography and resting-state activity correlations (note that

only positive functional links with significant correlations, passing the Bonferroni

multiple-comparison correction, were considered; see Methods). The low-resolution

networks were calculated as the average of the links connecting the ROIs within the

MAC pairs in the SC^{HR} and FC^{HR}. Note that while the diagonal of the connectivity matrices

(self-connections) was taken to be zero in the high-resolution space, in the low-

resolution space it was not zero but rather, it represented the average strength (defined

as the sum of links' weights) of the ROIs within a given MAC (internal MAC connectivity,

Supplementary figure 3).

To test our main hypothesis, i.e. that the structural centrality of brain nodes is inversely

correlated with embryogenic age, such that the earlier the FirsT the higher the

centrality, we computed the Spearman's partial correlation (SP-pCorr) between the EC

and FirsT, eliminating the effect of the cofounder white matter volume per ROI or MAC

that also shows a gradient between early and late generated structures (Supplementary

Fig. 2). Similarly, we repeated the same calculation for the functional networks, in this

case removing the cofounder of grey matter volume per ROI or MAC (Supplementary

Fig. 2). The plots of the EC obtained for the SC and FC are summarized at both spatial

resolutions in Fig. 2A.

When looking at the high-resolution networks, the centrality of the SC^{HR} and FC^{HR} nodes

showed significant trends with FirsT, respectively with negative (SP-pCorr=-0.44,

p=0.001) and positive (SP-pCorr=0.34, p=0.01) correlations. Note that the negative

correlations observed for the SC^{HR} (physically constrained spatial domain) support the

leading hypothesis that early born brain regions (e.g., the LC and brainstem) have

greater "hubness" or centrality, we refer to this as "the older gets richer" principle. By

contrast, in the case of the FC^{HR} (functional domain related to time and dynamics) the

positive correlations observed between centrality and FirsT highlight the greater

functional centrality of late brain circuits (such as neocortical ones).

For low-resolution networks, we observed negative centrality correlations with respect

to FirsT in both the SC^{LR} (SP-pCorr =-0.81, p<0.001) and FC^{LR} (SP-pCorr =-0.63, p=0.007).

The latter result shows a clear inverted trend between functional networks in low and

high-resolution. The inverted trend between the time of birth of the 18 different MACs

and the EC of the SC^{LR} and FC^{LR} can be visualized on a brain surface (Fig. 3A). In order to

understand the inversion of the EC trend between the FC^{LR} and FC^{HR}, we first dissected

the contribution of the links between the ROIs belonging to the same MAC (since these

are not considered by the centrality measure in the low-resolution case; we refer to

them as internal links, *int-Links*, see Methods), by looking at the ROIs' strength (i.e. the

sum of the links' weights of the ROIs). In the high-resolution networks, the trend of the

average ROIs' strength as function of FirsT (Supplementary fig. 3A) did not change when

the *int-Links* were discarded, and it displayed similar trends as in the case of the ROIs'

EC shown in fig. 2A. Focusing next on the MACs' links (Supplementary fig. 3B) as function

of FirsT, we observed the presence of few but very strong links in earlier MACs, which

also displayed overall higher strength, with explain the negative trend between MACs'

strength and FirsT as observed for the EC.

We next checked whether the difference in the time of neurogenesis (ΔFirsT) might also

be related to the probability of the existence and the weight of the links (Fig. 2B). In the

SC^{HR} and FC^{HR}, where a connection was present in 5% and 6.5% of the overall node pairs

respectively, the link probability decreased as a function of ΔFirsT with a Spearman

correlation of -0.53 (p<0.01) and -0.93 ($p<10^{-10}$). Similarly, the link weight decreased

with a correlation of -0.81 (p<0.001) and -0.87 ($p<10^{-5}$). In the SC^{LR} and FC^{LR}, only the link

weights were considered as these matrices are dense and it decreased as a function of

ΔFirsT with a correlation of -0.50 (p<0.001) and -0.52 ($p<10^{-12}$). The plots of the link

weights (see Supp. Fig. 4 and Supp. Fig 5 for each MAC in which each plot represents the

links of a given MAC and its FirsT is highlighted as a vertical broken line) provide a

visualization of how the brain circuits generated at similar neurogenic time points tend

to connect more strongly, we refer to this as the "preferential age attachment"

principle. As an example, the structural connectivity of three representative MACs was

visualized over a linear time scale, relative to the earliest (Locus Coeruleus, 28 days),

latest (occipital cortex, FirsT 54 days) and intermediate FirsT (hippocampus, 40 days: Fig.

3B1-3). This visualization shows that the links' weights of the earliest MAC are much

higher than that of the later MACs, and for the case of the LC and hippocampus, it also

shows that brain circuits generated at a similar neurogenic time point are more strongly

connected.

To obtain a list of genes potentially associated with the brain circuits' FirsT and the brain

networks' centrality, post-mortem brain transcriptomic data from the Allen Human

Brain Atlas (AHBA) was used. Given the limited spatial sampling across brain regions and

donors, we used 90 distinct brain regions with corresponding unique FirsT

(supplementary fig. 6), on which gene expression was quantified, and average values of

structural and functional centrality was calculated from the englobed ROIs (see

Methods). Average Spearman correlations between the brain maps of expression of

20,737 protein coding genes with FirsT, structural and functional centrality was

computed (see scatter plots of Fig. 4A). First, the correlation of gene expression with

functional centrality displayed a positive linear trend with respect to FirsT (Pearson

correlation 0.90, p<0.01), while an opposite trend was observed for structural centrality

(Pearson correlation -0.72, p<0.01).This observation shows that the expression of a

given gene is similarly correlated to brain maps of FirsT and of functional centrality, in

agreement with the fact that FirsT and functional centrality are positively correlated in

the high resolution networks. On the contrary, the negative trend observed when gene

expression is correlated to FirsT and to structural centrality is also in agreement with the

fact that structural centrality and FirsT are inversely correlated. Further confirmation of

what was just described is given by the anti-correlation (Pearson correlation -0.52,

p<0.01) observed for gene expression in functional and structural centrality maps.

Next, we performed enrichment analysis on the 1,256 genes (6% of the protein coding

genes) that were found with a correlation >1.64 standard deviations from the

distribution obtained (r>0.43). Gene set enrichment analysis of biological process

showed that the genes overrepresented were involved in processes related to

development of nervous central system and neuron projection development, and

regulation of ion transport, membrane potential, synaptic plasticity and transmission,

cell projection organization, insulin secretion, and neurotransmitter levels (Fig. 4B). In

addition, different cell components were also overrepresented, such as anchoring

junction, voltage-gated potassium channel complexes, dendrites, distal axons, and

membranes of synaptic vesicle, pre and post synapses, and neurons (Fig. 4B). Cell type

signatures were highly over-represented for several midbrain neurotypes, PFC microglia

and astrocytes, and fetal neurons (Fig. 4B). Notably, enrichment of human diseases

showed overrepresentation of mental and neurodevelopmental disorders, memory

impairment, epilepsy, and other pathologies (Fig. 4B).

Given these final observations in relation to human diseases, next, we tested and

dissected the hypothesis that the genetic contribution to neurological diseases is

differentially related to the brain networks' centrality maps and neurogenesis.

Therefore, we studied the distribution of the correlation between gene expression, FirsT

(fig. 5 top plots), functional (fig. 5 middle plots) and structural (fig. 5 bottom plots)

networks' centrality (we refer to these last two cases as the gex-centrality distributions).

We expected that genes causally related to neurological disorders would show extreme

correlation values (Fig. 5). In particular, we focused on Autism Spectrum Disorder (ASD),

Parkinson's disease (PD), Alzheimer's disease (AD) and epilepsy, and we combined two

main human gene datasets (see Methods): GeneCards (www.genecards.org) and the

Genome-Wide   Association   Study   Catalogue   (GWAS   Catalogue;

https://www.ebi.ac.uk/gwas/). For each gene reported in the GWAS for a given disease,

GeneCards dataset provides a relevance score based on the reported literature linking

the gene to the disease (coloured dots in Fig. 5A) while statistical evidence of their causal

relationship to the risk of developing a disease is guaranteed by the GWAS data. The

genes associated with neurological diseases (via GWAS) were not spread evenly across

the gex-centrality and neurogenesis distributions (Fig. 5B) and they showed a significant

enrichment at the percentile tails (see black arrows in Fig. 5C), supporting our

hypothesis. Notably, this observation was typified by genes with a well-known role, such

as: *APOE*, *TREM2* and *SORL1* (AD); *SCN1A, CUX2, PCDH7* and *CPLX1* (epilepsy); *SNCA,*

GBA and *MAPT* (PD); or *MEF2C* and *CACNA1C* (ASD).

Discussion

In this work, we use the earliest time of neurogenesis (FirsT) reported for brain circuits

to study adult brain networks that were reconstructed from structural-functional

(resting-state) MRI with low (MAC level) and high (ROI level) spatial resolution. By

correlating centrality metrics of complex networks with the FirsT, we observed a

decreasing gradient of structural hubness in MACs and ROIs from those circuits

generated earliest to those that appear later. This observation confirms our main

working hypothesis that "older nodes get richer" in terms of hubness within brain

networks. In the functional domain, the centrality of the MACs follows the same trend,

while conversely, the ROIs that developed later have a higher centrality. Since the links

present between the ROIs within a MAC do not affect the overall trend of the ROIs

strength as function of neurogenesis, we showed how indeed it is the presence of

stronger connections between earlier MACs both in the functional and structural case

that leads to a negative trend between MACs' centrality and neurogenesis.

Our results are consistent with brain differences early and late in development. The

neurons generated earlier in the human brain mature in an environment favourable to

cell growth, migration and axonal pathfinding. By contrast, neurons born later

differentiate into a stiffer neuropil that has a dense extracellular matrix and thus, their

growth and axonal pathfinding is more severely restricted by prior developmental

maturation ([14]). In addition, the growth of the brain itself plays a significant role in the

formation of long-range connectivity. Whereas after four of weeks of gestation (GW)

when the LC neurons begin their genesis, the brain measures 3-5 mm in length, whereas

at eight GW when many cortical neurons are generated, the human foetal brain

measures 27-31 mm([15]). Thus, the formation of general structural connectivity is

affected by the timing of neurogenesis.

The "older gets richer" rule stems from the literature on the connectomics of birth-

dated populations, both for neuronal populations that we were able to trace through

MRI as well as for other smaller regions of the brain that cannot be distinguished at MRI

resolution. In the former case, the neurons appearing first in the brain are those of the

LC([16–19]) and the brainstem and midbrain motor neurons, including those of the

trigeminal mesencephalic nucleus([20]). These organize into small nuclei and although

most of them cannot be segmented by MRI (here we only considered the LC and Raphe

nuclei), our model predicts that they all are potentially relevant in terms of their

connectivity. Noradrenergic LC neurons establish connections that span the whole

brain([21]), and both LC and trigeminal nuclei contribute to pathologies such as AD when

they degenerate([22–24]).

Beyond the brainstem, pioneering studies on GABAergic hub cells([11]) in the developing

mouse hippocampus have shown how early born GABAergic neurons operationally and

morphologically represent hub cells([12,25]), also later identified in other structures([26]) and

in the adult mouse hippocampus([13]). Within the neocortex, where different neuronal

populations are generated over sequential neurogenic stages([27]) but where these birth-

dated circuits cannot be tracked by MRI, hubs are related to neurogenesis. A recent

computational study of neocortical circuits reported that targeted injury to layer 5 hub

neurons produced the greatest damage to the structural-functional integrity of

neocortical circuits([28]). Significantly, these layer 5 neurons are amongst the earliest

neurons to be generated in the neocortex. Neuronal types generated earlier in the

hippocampus, cortex, brainstem and mesencephalon might also adopt a hub role due

to the timing of their neurogenesis. For example, stimulation of Purkinje cells in the

cerebellum (an early born GABAergic neuronal type in a circuit that develops early) has

been shown to inhibit spontaneous hippocampal seizures in a mouse model of temporal

lobe epilepsy([29]).

In terms of the topology of brain networks, we observe a decreasing gradient in the

probability and strength of connections as a function of the difference in time of

neurogenesis, both in structural and functional networks in the high and low degree of

spatial resolution. Therefore, nodes generated at similar times showed a higher

probability and strength of connection. This latter observation supports the hypothesis

that brain networks follow a "preferential age attachment rule", whereby nodes are

more likely to connect if the time of their neurogenesis differs little. From a biological

point of view, this might explain why neurons establish their main connections in a short

temporal window of plasticity, named the critical period([30,31]), before becoming

functional and maturing. Therefore, neurons with an equivalent birthdate tend to share

these permissive temporal windows, which allows their mutual/reciprocal connection.

In this context, a recent study has shown in the hippocampal network that sequential

neurogenesis affects connectivity and coactivity of neurons so that same birthdate

neurons joined into cell assemblies [32].

The rules of "preferential age attachment" and "older gets richer" extrapolate to the

context of brain networks, which will therefore adhere to the pioneering model of

Barabasi-Albert on the construction of scale-free networks and the genesis of network

hubs.

Brain connectivity, like other biological features, is ruled and limited by natural

selection. Due to the intimate relationship between development and evolution, it is

likely that the rules that relate neurogenic birth to connectivity are conserved across

species. Indeed, previous studies on the development of brain networks in *C. elegans*

highlight the early evolutionary appearance of neuronal hubs([33]). Moreover, neurons

linked by long-range connections tend to be generated at around the same neurogenic

time and early in development([33]), reinforcing the concept of a temporal windows of

plasticity around neuronal birthdates. Ultimately, all connections are shaped

evolutionarily to optimize axon length and the speed of connections([34,35]), and other

vertebrate brains are likely governed by the same rules of development and

connectivity.

Indeed, the brain regions shown here to have the most hubness are highly conserved in

the vertebrate brain([36]), suggesting their crucial importance in all vertebrates. These

regions include the posterior regions of the brain, such as the brainstem and

mesencephalon, and the cerebellum, which are known to be relatively similar across the

vertebrate taxa([37]). In part, the rule of the older gets richer suggests that ancient circuits

-such as those in brainstem that are related to autonomic animal functions and directly

involved in the animal's survival- are more influential in the network and more stable

across evolution. For example, the highly conserved neurogenic formation of the

cerebellum was recently demonstrated in several vertebrate species([38]) and its

remarkable orchestrating hub role has an impact on epileptic brain dynamics([29]). At the

other end of the spectrum, circuits related to associative tasks, more strongly linked to

cognition and human-specific behaviour, are not necessarily conserved and they appear

later in neurogenesis. In high-resolution networks we show here how their structural

hubness (low) is inversely related to their functional one (high).

In relation to brain transcriptomics, we studied how spatial maps of gene expression in

adult brains correlate with FirsT and structural-functional centrality maps. We found

that genes whose expression is positively correlated with the centrality of structural

circuits tend to show stronger expression in the circuits generated first. By contrast,

genes whose expression is anti-correlated with the centrality of structural circuits show

stronger expression in circuits in which neurogenesis occurs later. The opposite was

observed for functional networks and thus, the transcriptomics of structural and

functional networks appears to be driven distinctly by genes whose expression is

developmentally regulated. Previous studies into the genetic basis of human brain

network structure and function presented converging evidence that anatomical

connectivity is more strongly shaped by genetics than functional connectivity (see([39]) for

a review). Here, we add that the developmental sequence (specifically the first time

point of neurogenesis) is an additional key variable to interpret the genetic influence on

the connectome, providing a key feature linking embryogenesis to the topology of adult

brain networks. It also sheds light on the longitudinal reconstruction of developing brain

networks, a phenomenon that presents key technical challenges([5]).

Since enrichment analysis in relation to human diseases suggested a role in mental and

neurological disorders, next we explored if genes with a known role in neurological

diseases are linked closely to network centrality and neurogenesis with the potential to

exert an influence on key network hubs. Interestingly, in a recent similar study on the

UK Biobank database focused on the genetic architecture (heritability) underlying the

white matter connectome (i.e. solely focused on structural connectivity) [40], enrichment

analyses highlighted the key role of neurodevelopmental processes including

neurogenesis, neural differentiation, neural migration, neural projection guidance, and

axon development. In addition, structural connectivity displayed significant associations

also to polygenic scores for psychiatric, neurological, and behavioral traits.

In our research, we studied the correlation between FirsT, structural and functional

network centrality maps, and the expression of genes associated with epilepsy, Autistic

Spectrum Disorder (ASD), Parkinson's disease (PD) and Alzheimer's disease (AD), models

of neurodevelopmental and neurodegenerative diseases. First, we observed that a

significant number of genes linked to ASD and epilepsy show high positive correlations

to neurogenesis and functional centrality maps in agreement with the evidence that ASD

and several forms of epilepsy (such as Dravet syndrome and infant epilepsy) is linked to

late brain developmental processes ([41]) and supporting the idea that ASD and infant

genetic forms of developmental epilepsy can lead to functional brain maps' lesion rather

than structural ones ([42]). Distinctly, in our model a significant population of genes

related to PD express similarly to structural centrality maps, so highly expressed in early

generated structures (i.e. structural brain hubs) and with lower expression in later

developed structures. This support the idea that PD is a disease linked to targeted

structural lesions such as those ones occurring in the dopaminergic neuronal population

in the basal midbrain (which is the third in the rank of neurogenesis). In the case of the

AD, we identified a significant high number of genes with anti-correlated expression to

functional centrality, so potentially poorly expressed in functional hubs (late structures)

and viceversa. Therefore, we conclude that dysfunctional properties of the proteins

coded by these genes, maybe also associated to alterations in their expression levels,

can have potential high impact in brain hubs at developmental, functional and structural

levels and can play a key role in the aetiology and associated dysfunctions of brain

pathologies. Highlighting some well-known genes with top correlation ranks (see

supplementary tables) in relation to the studied brain pathologies, we found a

contribution of four genes known to drive PD: *SNCA(*[43,44]) that encodes the

parkinsonism-associated Lewy body protein alpha synuclein, *MAPT(*[45]) encoding the

microtubule-associated protein tau, GBA [46] encoding the lysosomal enzyme

glucocerebrosidase and SLC41A1 [47] encoding a magnesium transporter (a candidate for

the causative gene in the PARK16). Together our results indicate that *SNCA*, *MAPT* and

GBA are expressed more weakly in nodes with higher structural centrality under

physiological conditions and viceversa for SLC41A1 (Fig. 5A). We also found associations

for the gene encoding apolipoprotein E, *APOE*, which is directly related to AD, cerebral

amyloid angiopathy, PD and other neurological diseases([48]), and for *TREM2*, a receptor

expressed in myeloid cells 2 that is related to microglial biology, a high risk of developing

AD([49]) and prion diseases([50]). In addition, the *SORL1* gene that encodes the sortilin-

related receptor, and that has been associated with early and late-onset AD([51]), was also

found to have a strong positive correlation with functional centrality. In the case of ASD,

we highlight the gene MEF2C[52] which regulates cortical inhibitory and excitatory

synaptic balance and CACNA1C coding for a calcium channel linked to selective

autophagy and altered axon targeting and behavior[53]. For epilepsy, we highlight the

presence of the *SCN1A* gene, which encodes the Alpha 1 Subunit of the Sodium Voltage-

Activated Channel, occupying position 144 in correlation with the functional centrality

of 20,787 genes considered, and with a known causal relationship to Dravet Syndrome,

a rare genetic disease causing one of the most devastating forms of childhood

generalized epilepsy([54]).

Notably, many genes known to underlie neurodevelopmental disorders (as the ones for

example highlighted in [55]) lie in the extreme percentiles of the FirsT or centrality

distributions (such as CACNA1A, DEPDC5, SCN8A, and more).

As a result of both human disease enrichment and deeper analyses of four neurological

conditions, our results provide support for the hypothesis that neurological diseases

may be linked to key topological alterations to brain networks in relation to node

hubness([56]). Thus, altered transcription of genes whose expression correlates closely

with network centrality could lead to profound neurological lesions and disease. In this

context, stroke and other circuit damage to early-generated brain structures (such as

those in the brainstem region) are known to have life-threatening consequences relative

to later-generated cortical damage.

MAIN REFERENCES

1.　　Betzel, R. F. & Bassett, D. S. Multi-scale brain networks. *NeuroImage* **160**, 73–83
(2017).

2.　　Buzsáki, G. *Rhythms of the brain*. (Oxford University Press, 2006).

3.　　Fox, M. D. Mapping Symptoms to Brain Networks with the Human Connectome. *N Engl*
J Med **379**, 2237–2245 (2018).

4.　　Cisek, P. & Hayden, B. Y. Neuroscience needs evolution. *Phil. Trans. R. Soc. B* **377**,
20200518 (2022).

5.　　Oldham, S. & Fornito, A. The development of brain network hubs. *Developmental*
Cognitive Neuroscience **36**, 100607 (2019).

6.　　Cimini, G. *et al.* The statistical physics of real-world networks. *Nat Rev Phys* **1**, 58–71
(2019).

7.　　Broido, A. D. & Clauset, A. Scale-free networks are rare. *Nat Commun* **10**, 1017 (2019).

8.　　Bassett, D. S., Khambhati, A. N. & Grafton, S. T. Emerging Frontiers of
Neuroengineering: A Network Science of Brain Connectivity. *Annu. Rev. Biomed. Eng.* **19**, 327–
352 (2017).

9.　　Barabási, A.-L. & Albert, R. Emergence of Scaling in Random Networks. *Science* **286**,
509–512 (1999).

10.　　van den Heuvel, M. P. & Sporns, O. Network hubs in the human brain. *Trends in*
Cognitive Sciences **17**, 683–696 (2013).

11.　　Bonifazi, P. *et al.* GABAergic Hub Neurons Orchestrate Synchrony in Developing
Hippocampal Networks. *Science* **326**, 1419–1424 (2009).

12.　　Picardo, M. A. *et al.* Pioneer GABA Cells Comprise a Subpopulation of Hub Neurons in
the Developing Hippocampus. *Neuron* **71**, 695–709 (2011).

13.　　Bocchio, M. *et al.* Hippocampal hub neurons maintain distinct connectivity throughout
their lifetime. *Nat Commun* **11**, 4559 (2020).

14.　　Martínez, G. F. *et al.* Extracellular matrix stiffness negatively affects axon elongation,
growth cone area and F-actin levels in a collagen type I 3D culture. *J Tissue Eng Regen Med* **16**,
151–162 (2022).

15.　　Stiles, J. & Jernigan, T. L. The Basics of Brain Development. *Neuropsychol Rev* **20**, 327–
348 (2010).

16.　　Altman, J. & Bayer, S. A. Development of the brain stem in the rat. I. Thymidine-
radiographic study of the time of origin of neurons of the lower medulla. *J. Comp. Neurol.* **194**,
1–35 (1980).

17.　　Altman, J. & Bayer, S. A. Development of the brain stem in the rat. III. Thymidine-
radiographic study of the time of origin of neurons of the vestibular and auditory nuclei of the
upper medulla. *J. Comp. Neurol.* **194**, 877–904 (1980).

18. Altman, J. & Bayer, S. A. Development of the brain stem in the rat. V. Thymidine-radiographic study of the time of origin of neurons in the midbrain tegmentum. *J. Comp. Neurol.* **198**, 677–716 (1981).

19. Steindler, D. A. & Trosko, B. K. Two types of locus coeruleus neurons born on different embryonic days in the mouse. *Anat Embryol* **179**, 423–434 (1989).

20. Altman, J. & Bayer, S. A. Development of the brain Stem in the rat. II. Thymidine-radiographic study of the time of origin of neurons of the upper medulla, excluding the vestibular and auditory nuclei. *J. Comp. Neurol.* **194**, 37–56 (1980).

21. Kebschull, J. M. *et al.* High-Throughput Mapping of Single-Neuron Projections by Sequencing of Barcoded RNA. *Neuron* **91**, 975–987 (2016).

22. Goto, T. *et al.* Neurodegeneration of Trigeminal Mesencephalic Neurons by the Tooth Loss Triggers the Progression of Alzheimer's Disease in 3×Tg-AD Model Mice. *JAD* **76**, 1443–1459 (2020).

23. Beardmore, R., Hou, R., Darekar, A., Holmes, C. & Boche, D. The Locus Coeruleus in Aging and Alzheimer's Disease: A Postmortem and Brain Imaging Review. *JAD* **83**, 5–22 (2021).

24. Jacobs, H. I. L. *et al.* In vivo and neuropathology data support locus coeruleus integrity as indicator of Alzheimer's disease pathology and cognitive decline. *Sci. Transl. Med.* **13**, eabj2511 (2021).

25. Cossart, R. Operational hub cells: a morpho-physiologically diverse class of GABAergic neurons united by a common function. *Current Opinion in Neurobiology* **26**, 51–56 (2014).

26. Mòdol, L. *et al.* Spatial Embryonic Origin Delineates GABAergic Hub Neurons Driving Network Dynamics in the Developing Entorhinal Cortex. *Cerebral Cortex* **27**, 4649–4661 (2017).

27. Molnár, Z. *et al.* New insights into the development of the human cerebral cortex. *J. Anat.* **235**, 432–451 (2019).

28. Gal, E. *et al.* The Role of Hub Neurons in Modulating Cortical Dynamics. *Front. Neural Circuits* **15**, 718270 (2021).

29. Krook-Magnuson, E., Szabo, G. G., Armstrong, C., Oijala, M. & Soltesz, I. Cerebellar Directed Optogenetic Intervention Inhibits Spontaneous Hippocampal Seizures in a Mouse Model of Temporal Lobe Epilepsy. *eneuro* **1**, ENEURO.0005-14.2014 (2014).

30. Dehorter, N. & Del Pino, I. Shifting Developmental Trajectories During Critical Periods of Brain Formation. *Front. Cell. Neurosci.* **14**, 283 (2020).

31. Wong-Riley, M. T. T. The critical period: neurochemical and synaptic mechanisms shared by the visual cortex and the brain stem respiratory system. *Proc. R. Soc. B.* **288**, 20211025 (2021).

32. Huszár, R., Zhang, Y., Blockus, H. & Buzsáki, G. Preconfigured dynamics in the hippocampus are guided by embryonic birthdate and rate of neurogenesis. *Nat Neurosci* **25**, 1201–1212 (2022).

33. Varier, S. & Kaiser, M. Neural Development Features: Spatio-Temporal Development of the Caenorhabditis elegans Neuronal Network. *PLoS Comput Biol* **7**, e1001044 (2011).

34.   Pavlovic, D. M., Vértes, P. E., Bullmore, E. T., Schafer, W. R. & Nichols, T. E. Stochastic
Blockmodeling of the Modules and Core of the Caenorhabditis elegans Connectome. *PLoS ONE*
9, e97584 (2014).

35.   Bassett, D. S. *et al.* Efficient Physical Embedding of Topologically Complex Information
Processing Networks in Brains and Computer Circuits. *PLoS Comput Biol* **6**, e1000748 (2010).

36.   Nieuwenhuys, R. *et al. The Central Nervous System of Vertebrates*. (Springer Berlin /
Heidelberg, 1997).

37.   Suárez, L. E. *et al. A connectomics-based taxonomy of mammals*.
http://biorxiv.org/lookup/doi/10.1101/2022.03.11.483995 (2022)
doi:10.1101/2022.03.11.483995.

38.   Rueda-Alaña, E. & García-Moreno, F. Time in Neurogenesis: Conservation of the
Developmental Formation of the Cerebellar Circuitry. *Brain Behav Evol* 1–15 (2021)
doi:10.1159/000519068.

39.   Arnatkeviciute, A., Fulcher, B. D., Bellgrove, M. A. & Fornito, A. Where the genome
meets the connectome: Understanding how genes shape human brain connectivity.
NeuroImage **244**, 118570 (2021).

40.   Sha, Z., Schijven, D., Fisher, S. E. & Francks, C. Genetic architecture of the white matter
connectome of the human brain. *Sci. Adv.* **9**, eadd2870 (2023).

41.   Lee, B. H., Smith, T. & Paciorkowski, A. R. Autism spectrum disorder and epilepsy:
Disorders with a shared biology. *Epilepsy & Behavior* **47**, 191–201 (2015).

42.   Ghacibeh, G. A. & Fields, C. Interictal epileptiform activity and autism. *Epilepsy &*
Behavior **47**, 158–162 (2015).

43.   Stefanis, L. -Synuclein in Parkinson's Disease. *Cold Spring Harbor Perspectives in*
Medicine **2**, a009399–a009399 (2012).

44.   Zarranz, J. J. *et al.* The new mutation, E46K, of α-synuclein causes parkinson and Lewy
body dementia: New α-Synuclein Gene Mutation. *Ann Neurol.* **55**, 164–173 (2004).

45.   Paul, K. C. *et al.* APOE, MAPT, and COMT and Parkinson's Disease Susceptibility and
Cognitive Symptom Progression. *JPD* **6**, 349–359 (2016).

46.   Sidransky, E. *et al.* Multicenter Analysis of Glucocerebrosidase Mutations in
Parkinson's Disease. *N Engl J Med* **361**, 1651–1661 (2009).

47.   Cornell, R. SLC41A1 and TRPM7 in magnesium homeostasis and genetic risk for
Parkinson?s disease. *J Neurol Neuromedicine* **1**, 23–28 (2016).

48.   Verghese, P. B., Castellano, J. M. & Holtzman, D. M. Apolipoprotein E in Alzheimer's
disease and other neurological disorders. *The Lancet Neurology* **10**, 241–252 (2011).

49.   Ulland, T. K. & Colonna, M. TREM2 — a key player in microglial biology and Alzheimer
disease. *Nat Rev Neurol* **14**, 667–675 (2018).

50.   Diaz-Lucena, D. *et al.* TREM2 expression in the brain and biological fluids in prion
diseases. *Acta Neuropathol* **141**, 841–859 (2021).

51.　　Cuccaro, M. L. *et al. SORL1* mutations in early- and late-onset Alzheimer disease.
Neurol Genet **2**, e116 (2016).

52.　　Harrington, A. J. *et al.* MEF2C regulates cortical inhibitory and excitatory synapses and
behaviors relevant to neurodevelopmental disorders. *eLife* **5**, e20059 (2016).

53.　　Buddell, T., Friedman, V., Drozd, C. J. & Quinn, C. C. An autism-causing calcium channel
variant functions with selective autophagy to alter axon targeting and behavior. *PLoS Genet* **15**,
e1008488 (2019).

54.　　Claes, L. *et al.* De Novo Mutations in the Sodium-Channel Gene SCN1A Cause Severe
Myoclonic Epilepsy of Infancy. *The American Journal of Human Genetics* **68**, 1327–1332 (2001).

55.　　Parenti, I., Rabaneda, L. G., Schoen, H. & Novarino, G. Neurodevelopmental Disorders:
From Genetics to Functional Pathways. *Trends in Neurosciences* **43**, 608–621 (2020).

56.　　Bassett, D. S. & Bullmore, E. T. Human brain networks in health and disease. *Current
Opinion in Neurology* **22**, 340–347 (2009).

57.　　Workman, A. D., Charvet, C. J., Clancy, B., Darlington, R. B. & Finlay, B. L. Modeling
Transformations of Neurodevelopmental Sequences across Mammalian Species. *Journal of
Neuroscience* **33**, 7368–7383 (2013).

58.　　Glasser, M. F. *et al.* The minimal preprocessing pipelines for the Human Connectome
Project. *NeuroImage* **80**, 105–124 (2013).

59.　　Oldham, S. *et al.* Consistency and differences between centrality measures across
distinct classes of networks. *PLoS ONE* **14**, e0220061 (2019).

60.　　Arnatkevičiūtė, A., Fulcher, B. D. & Fornito, A. A practical guide to linking brain-wide
gene expression and neuroimaging data. *NeuroImage* **189**, 353–367 (2019).

61.　　Diez, I. & Sepulcre, J. Unveiling the neuroimaging-genetic intersections in the human
brain. *Current Opinion in Neurology* **Publish Ahead of Print**, (2021).

62.　　Zhou, Y. *et al.* Metascape provides a biologist-oriented resource for the analysis of
systems-level datasets. *Nat Commun* **10**, 1523 (2019).

63.　　Piñero, J. *et al.* The DisGeNET knowledge platform for disease genomics: 2019 update.
Nucleic Acids Research gkz1021 (2019) doi:10.1093/nar/gkz1021.

64.　　Fulcher, B. D., Arnatkeviciute, A. & Fornito, A. Overcoming false-positive gene-category
enrichment in the analysis of spatially resolved transcriptomic brain atlas data. *Nat Commun*
12, 2669 (2021).

65.　　Burt, J. B., Helmer, M., Shinn, M., Anticevic, A. & Murray, J. D. Generative modeling of
brain maps with spatial autocorrelation. *NeuroImage* **220**, 117038 (2020).

TABLES

Table 1

	Major developmental brain area	Circuit	fMRI Comprised structures	Translated earliest stage in humans (embryonic days)	Earliest stage in other species	source references	Observations	ROIs #
1	Rhombencephalon	Locus Coeruleus	Locus Coeruleus	28	Rat - Soon before E12; Mouse - E9	T1–4	Likely one of the earliest generated nucleus. Noradrenergic neurons are the earliest-born within the LC	2
2	Rhombencephalon	Medulla Oblongata	Pons and Medulla oblongata	31	Rat - Soon after 12	T1	Brainstem neurogenesis starts much earlier than in the rest of the brain. However, we could not segment its many different compartments by fMRI and so it appears fused in one single circuit.	40
3	Mesencephalon	Midbrain	Superior colliculus, inferior colliculus, tegmental midbrain	32	Mouse - E10.5	T5,6	Substantia nigra pars reticulata is the earliest mesencephalic nucleus to start neurogenesis	23
4	Secondary Prosencephalon	Hypothalamus	Hypothalamus, mammilary bodies	34	Rat - E12; Macaque - E33; Cat - E21	T7–9; translatingtime.org	Lateral hypothalamic nuclei (lateral hypothalamic nucleus, lateral preoptic area and lateral mammilary bodies) start hypothalamic neurogenesis at	18
	Telencephalon	Olfactory	Olfactory Cortices	34	Mouse E11	T10; translatingtime.org	Unique segmentation and ROI identification with Hypothalamus	
5	Rhombencephalon	Raphe Nuclei	Raphe nuclei	35	Rat E12	T1	Neurogenesis in the Raphe nuclei starts in the Raphe Magnus nucleus	3
6	Telencephalon	Basal Ganglia	Caudate, Putamen, Accumbens, Septum	36	Mouse - E11; Macaque - E38	T11	Accumbens starts neurogenesis at E38 in macaque; Striatum and globus pallidus start soon after.	77
7	Rhombencephalon	Cerebellum	Cerebellar cortex, deep cerebellar nuclei	36	Mouse E11; Rat - E13	T12–14	Earliest neurons are projection neruons of the deep cerebellar nuclei, followed by Purkinje neurons.	275
8	Telencephalon	Entorhinal	Entorhinal cortex	37	Rat - soon after E14	T15	Neurogenesis starts earlier to other parahippocampal cortices	11
9	Telencephalon	Amygdala	Pallial amygdala, subpallial amygdala	38	Mouse E11	T16,17	Neurogenesis starts with GABAergic populations of central and medial nuclei, followed by the glutamatergic neurons of the baso-lateral complex	8
10	Telencephalon	Hippocampus	Amon horn, Dentate gyrus	40	Rat E14.5	T15	Earliest neurons occupy the superficial and deep layers of the Ammon's horn (CA1 to CA3), and the hilus	21
11	Telencephalon	Posterior cingulum	Posterior cingular cortex	42	Rat E14.5	translatingtime.org	Inferred from translatingtime.org Earlier than the anterior cingulate cortex	22
12	Telencephalon	Parahippocampal cortices	Parahippocampal cortex	44	Rat E15	T15	Includes subiculum and parasubiculum. Unique segmentation and ROI identification with Insula	107
	Telencephalon	Insula	Insular cortex	44	Rat E15	T18	First insular neurons occupy the dep layers of the ventral agranular insular cortex.	
13	Telencephalon	Anterior Cingulate cortex	Caudal Anterior Cingulate, Rostral Anterior Cingulate	45	Macaque - E40	T19	First Layer VI neurons in the anterior cingulate cortex were revealed with [3H]-thymidine injection at E40 in the macaque brain	37
14	Diencephalon	Thalamus	Thalamic nuclei: LP, VL, VPL, LGN, MGN, Pulvinar,	46	Rat - E14	T20	Several thalamic nuclei (VB, VE, LGN, PO, MGN and LP) start neurogenesis at E14 in the rat brain.	28
15	Telencephalon	Frontal cortex	Caudal middle frontal, lateral orbito frontal, medial orbito frontal, paracentral, pars opercularis, pars orbitalis, pars triangularis, precentral, rostral middle frontal, frontal pole and superior frontal cortices	48	Macaque - E40; Ferret - E30; Mouse - E11.5	T21–23 translatingtime.org	Neurogenesis in the cortex starts synchronously throughout all cortical areas. The earliest-generated cortical populations are the Cajal-Retzius cells of the marginal zone and the subplate neurons of layer 6b. Most of these neurons die early after birth, which is why we considered the early generation of deep cortical layers as the onset of a functional cortical neurogenesis. In these layers, there is a gradient of neurogenesis starting first at the rostral pole of the cortex, and finishing last at the caudal regions. In macaque, frontal cortex starts at E45, and occipital cortex starts at E54. In human, we estimate ths gradient as a 2-day difference between cortical lobes.	719
16	Telencephalon	Parietal cortex	Inferior parietal, postcentral, precuneus, superior parietal and supramarginal cortices	50				502
17	Telencephalon	Temporal cortex	Banks of the Superior Temporal Sulcus, fusiform, inferior temporal, superior temporal, and transverse temporal cortices	52				384
18	Telencephalon	Occipital cortex	Cuneus, Lateraloccipital cortex, lingual cortex, pericalcarine cortex	54				289

Table 1 Developmental information of the human brain macrocircuits studied. The

table includes the list of 18 MACs, the developmental brain region from which each MAC

is derived, the main anatomical brain structures contained in each MAC, information

about the earliest stage at which neurogenesis (FirsT) is observed or predicted in the

human brain, and in other mammalian species, and the number of ROIs in each MAC.

References in Supplementary Material.

FIGURE LEGENDS

Figure 1. From the circuits' embryogenic age to brain networks and transcriptomics.

(A) Sagittal schemes of the early embryonic human brain. The upper scheme represents

the neuromeres and fundamental regions of development, the lower scheme displays

the location of the 18 MACs within the early embryonic human brain, colour-coded

according to their FirsT. (B) Scheme of the neuroimage analysis pipeline. Brain networks

were reconstructed from healthy adult subjects scanned by MRI at 7T within the HCP

initiative. For each subject, the structural and functional networks were reconstructed

using probabilistic tractography and resting-state activity, respectively. For every pair of

ROIs (2,566 ROIs), the structural connectivity matrix represents the putative number of

fibres connecting them, while the functional connectivity matrix reports the correlation

in their activity as revealed by the BOLD time series. The averages across subjects were

used as final representative brain networks (see Methods). From the 2566 ROIs'

matrices, the 18 MACs' networks were reconstructed and represents for each pair of

MACs the average links between the ROIs forming them (see Methods). To identify

patterns between networks' topology and neurogenesis, the correlation between the

nodes' metrics (centrality or segregation) and embryonic age was calculated. (C) Brain

transcriptome data from AHBA dataset (see methods) was used to search for protein

coding genes with a high similarity between its spatial brain expression and embryonic

age. Functional annotations of the obtained genes were further computed using

overrepresentation analysis to find significantly associated biological processes and

cellular components associated with embryonic age.

Figure 2. Scatter plots of the nodes' centrality and the links' probability/weight

relative to the FirsT, in the high and low resolution structural-functional networks. (A)

Scatter plots of the nodes' centrality as a function of embryonic age. Red and blue

indicates the structural and functional cases, respectively. The first and second columns

from left represent the results for the 2,566 ROIs as violin plots, where the mean of each

group (i.e. ROIs within a MAC) is plotted in black, and the third and fourth columns

represents the 18 MACs. All plots have a logarithmic scale in the y-axis. (B) Scatter plots

of the link weights (left y-axis) as a function of the differences in the first neurogenic

birthdate. The colours and columns are as indicated in panel A. The violin plots for the

2,566 ROIs show the link weight distributions while the solid blue/red lines plot the

average values. Black solid lines show the exponential (linear) fit on the average values

for the structural (functional) case. The link probability is represented by the right y-axes

and displayed as dots, and the broken black line showing the exponential fit. All plots

have a logarithmic scale in the y-axis.

Figure 3. Visualization of brain connectivity and the first neurogenic birthdate. (A) The

colour coded glass brain maps of the eighteen MACs represents the FirsT (A1; see colour

bar on the left) and the z-score (see colour bar on the right) of the eigenvector centrality

for the SC^{LR} (A2) and the FC^{LR} (A3). (B) Linear temporal representation with arches of the

MACs structural connectivity for the earliest (B1) generated MAC (Locus Coeruleus, FirsT

28), for a representative mid-term (B2) generated MAC (Hippocampus, FirsT 40) and for

the latest (B3) generated MAC (Occipital Cortex, FirsT 54). The x-axis represents the

developmental time of several FirsT, where the different MACs are located according to

their FirsT. Note that the Basal Ganglia and Cerebellum overlap since their neurogenesis

is initiated on day 36. The thickness of top black arches is proportional to the link weight

between the MACs extracted from the 18 node adjacency matrix. Broken black lines

mark the presence of connections but with little weight (i.e. below the lowest threshold

for the visualization of solid lines).

Figure 4. Transcriptome of brain circuits according to embryogenic age. (A) Mean

spatial similarity between the expression of 20,737 protein-coding genes with the time

of birth and structural and functional centrality. Structural and functional centrality of

the 2,566 brain regions was downsampled to 90 region (with gene expression data) and

for each of the 90 regions one of the 18 FirsT was assigned. (B) 1,256 genes from the

positive tail (6% of protein coding genes) – genes whose spatial expression is most

similar to FirsT and functional and structural centrality – were used to compute the

overrepresentation analysis for different annotations: biological processes, cell

components, cell types, and human diseases.

Figure 5. Correlations between the maps of the gene transcriptome, neurogenesis and

hubness in relation to Autism Spectrum Disorder, epilepsy, Parkinson's and

Alzheimer's disease. For each gene the spatial expression in the brain was correlated to

the spatial maps of eigenvector centrality and neurogenesis (note that from the sub-

sampled brain transcriptomes, 90 brain regions were considered: see Methods). **A**

Probability distribution of the correlations (light gray histogram, left axis) between

transcriptomics (20,787 genes) and neurogenesis (top plot), functional centrality

(middle plot) and structural centrality (bottom plot). The normalized relevance score of

the GWAS genes listed for each disease (obtained from the GeneCards databases; Note

that normalization is computed separately for each disease; see Methods) has been

plotted with dots (right axis). The colors used to display the results for the different

diseases in pnale A-C, are shown in the inset legend of the top plot of panel B. . Vertical

broken lines highlight the lowest and top five percentile interval. Coloured labels and

arrows mark the most relevant representative GWAS genes with correlations in the

highlighted extreme percentiles. **B** For each disease, the normalized count (percentage

of gene per interval) of GWAS genes in five percentile intervals of the correlation

distribution (light gray histogram of panel A) is plotted for the different diseases (legend

of colours is in the inset of the top plot)). **C** P-value of the normalized gene count in each

percentile. The threshold for p<0.01 calculated from the null model (based on a

thousand reshuffled replications) which keeps into account spatial dependencies of

both centrality maps and gene-expression is plotted as a broken horizontal black line.

The arrows mark the extremities of the correlation distributions with a significantly

higher number of GWAS-genes than what expected by chance.

METHODS

The codes to reproduce the presented analyses will be made available to interested

researchers upon a motivated request.

Neurogenesis timeline of brain circuits – Selection of circuits.

To select the circuits to study, we needed to reach a compromise between the

conflicting resolution of MRI and the developmental structural units. Certainly, there are

hundreds of brain structures and circuits that can be segmented by MRI (cf. the list of

areas that were identified in the AAL classification (49) and others (50)), although many

of these regions do not develop independently (for example, the many neocortical

areas). By contrast, there are brain regions that are developmentally relevant but that

cannot be precisely segmented by MRI (e.g. thalamic nuclei). Therefore, the two criteria

followed to select the list of brain circuits that ultimately studied were developmental

independence and MRI segmentability.

Developmental independence: Based on how the brain is generated during embryonic

development, we chose brain regions that were derived from independent units of the

early differentiated brain, called fundamental morphogenic units (FMUs (51)). Early in

development, all vertebrate brains divide into different vesicles: secondary

prosencephalon, diencephalon, midbrain, and hindbrain. Each of these vesicles can be

segmented into several neuromeres, which are developmentally independent rings of

the neural tube destined to form the major brain regions. Each neuromere contains

several FMUs along its dorsal-ventral axis (figure 1A) and the neurons within a given

FMU are formed mostly by neurogenesis. For example, the m1 neuromere in the

mesencephalon gives rise to most of the neurons in the superior colliculus, so we

consider the superior colliculus to be a developmental unit. When a FMU generates

through development several brain regions that we can resolve through MRI (e.g. the

occipital neocortex), we have merged these structures into a single MAC, as those

regions are not developmentally independent and their FirsTs are therefore the same.

The literature available shows the specific date of birth of the different populations in

each developmental unit in several mammalian species (see below).

MRI segmentability: The human brain evolved following an unprecedented expansion

of the neocortex, which develops from a single developmental unit (52). But this

expansion came at the expense of shrinking many brain regions, such that some are so

small that they cannot be segmented at MRI resolution. For example, the brainstem and

pons are generated from the rhombencephalon, and there are up to 11 rhombomeres,

each of which hosts a number of FMUs but none of which can be segmented by MRI.

Thus, we identified consistently segmentable brain structures from MRIs. Consequently,

no circuit was assigned to two different birth dates. Most brain structures were

generated during a neurogenic period but the first neurogenic time (FirsT) was

considered to be the first day after which neurons began to appear.

Neurogenesis timeline of brain circuits - Estimation of FirsT.

We hypothesize that the strength of adult connectivity in a given region depends on the

developmental stage at which it forms. ([57]) . The same method to estimate human FirsT

was applied for all circuits, this is a summary of the pipeline:

1.   Experimental birthdating of the neurons of a given brain area through DNA-

synthesis labelling (via BrdU or [3]H-Thymidine administration) in non-human species

(references in Table 1).

2.   Determination of the developmental stage at which the first neurons of that

brain region were generated in that species (stages on Table 1).

3.   Translation of this non-human stage into human developmental stage via

translatingtime.org ([57]) which provides the FirsT value of the given brain region. This is

possible and reliable thanks to the great conservation of the developmental sequence

found across mammals ([57]).

Each FirsT data point is based on at least one article, all articles are referenced in Table

1. When several independent articles displayed research on the same structure (i.e., the

locus coeruleus), these led to the same FirsT measurement. Therefore, our FirsT data

points represent a single embryonic day as FirsT, not a range/interval.

No single database contains birthdate information for all brain regions examined.

MRI DATASET AND PROCESSING

Participants

Neuroimaging data was acquired from the Human Connectome Project (WU-Minn

Consortium - PIs David Van Essen and Kamil Ugurbil: 1U54MH091657), funded by the 16

National Institutes of Health (NIH) that support the NIH Blueprint for Neuroscience

Research and by the McDonnell Center for Systems Neuroscience (Washington

University). For this study, we took data from N = 184 healthy subjects acquired at 7

Tesla (72 males, 112 females; 24 participants between 22-25 years, 84 between 26-30

years, 73 between 31-35 and 2 over 36). High-resolution structural T1 images, functional

magnetic resonance images at rest (fMRI) and diffusion tensor images (DTI) were used

(for more information on the acquisition parameters see Supplementary Methods and

the   Human   Connectome   Project   documentation   -

http://www.humanconnectome.org/).

Image preprocessing

Minimal pre-processed data was downloaded from the Human Connectome([58]3). T1

structural images were aligned to the anterior and posterior commissures, skull

stripped, corrected for gradient distortion and bias fields, and the non-linear

transformation to the MNI152 standard space was computed. Using a one-step

resampling approach, resting state functional data was corrected for gradient-non-

linearity-induced distortion and movement within runs using a rigid body

transformation (six parameter linear transformation) and EPI distortion. Both the

transformation from the reference image to the structural T1 image, and from T1 to the

standard space were also performed through a one-step resampling approach. A subject

specific functional data projection to MNI152 was obtained. Diffusion images were

normalized for b0 intensity and EPI distortion, movement and eddy currents were

corrected, as were gradient non-linearities. The B0 image was co-registered to the

subject's anatomical T1 images. For more details on the minimal pre-processing of the

human connectome refer to the original paper([58]3). To further reduce the noise in the

functional data, a general linear model was used to remove linear and quadratic trends,

as well as the contribution of movement, the cerebrospinal fluid and white matter

signals. Additionally, band-pass filtering (0.01–0.08 Hz) and spatial smoothing with an

isotropic Gaussian kernel of 6 mm FWHM was applied.

Brain macrocircuits (MACs) at low resolution

Several regions from different brain MRI atlases were combined to generate 18 different

macrocircuits. The atlases used included the Automated Anatomical Labeling (AAL([49])),

Freesurfers Desikan-killiany atlas([54]), CIT168 subcortical atlas([55]) and locus coeruleus

atlas([56]). The LC and raphe nuclei were chosen as the functional ROIs that englobed

them (for further details see Table 1).

Brain regions of interest (ROIs) at high resolution

High resolution parcellation was performed after clustering the functional data to study

network properties with about 2,500 ROIs. In particular (and similar to([57])), clustering

was performed based on temporal correlations between pairwise voxel time series that

impose a constraint to ensure spatially contiguous ROIs. This was performed in two

stages: 1. clustering at the single subject level; and 2. a second clustering applied to

individual subject data. To avoid ROIs with voxels belonging to several circuits rather

than performing clustering on the whole brain, this strategy was applied separately for

each circuit. Small circuits were neglected and those voxels were inserted into the

nearest circuit. After parcellation, we took all ROIs overlapping some of the existing

circuits, resulting in 2,566 regions covering the whole brain.

STRUCTURAL AND FUNCTIONAL BRAIN NETWORKS

High-resolution

After image processing, the connectivity matrices were obtained at high-resolution, representing two modalities composed of 2,566 ROIs: structural and functional networks. Individual subject functional connectivity matrices were obtained by averaging all the time series of the voxels belonging to each ROI generated, and by calculating the Pearson correlation value between the time series corresponding to each pair of ROIs. The average of the functional matrices across all subjects was then calculated. Only positive links with an average minimal correlation of 0.19 were considered. This value corresponds to a p-value of 0.7×10^{-9} when calculated on time series composed by 900 points. Such p-value threshold was chosen in order to apply a strict Bonferroni correction on a 0.01 p-value of multiple comparisons, i.e. divided by the number of links in the adjacency matrix (which correspond to half of the elements in matrix excluding the diagonal considering that adjacency matrices are symmetric by definition in our methodology, i.e. (2566*2566)-2566/2). This final matrix defined the average high-resolution functional connectivity matrix: FC^{HR}, and it had a connectivity density of 6.5%.

FSL functions were used to quantify the individual subject structural connectivity matrices. First FSL BEDPOSTX was used to model crossing fibres and subsequently, probabilistic tractography was performed with PROBTRACKX to generate a #ROI × #ROI matrix per subject. Matrices were normalized between 0 and 1 dividing each element by their maximum. The average across subject matrices were finally obtained and only

the top 5% of higher weights were considered to create the high-resolution structural

connectivity matrix, SC^{HR}, consistent with previous studies (58). ROIs' hubness

quantified with eigenvector centrality in the SC^{HR} in the dense (when all links were

considered) or sparse (when only top 5% links were considered) adjacency matrix,

correlated with a 0.99 value ($p<10^{-10}$) so they did not show significant differences. For

both the functional and structural matrices, the principal diagonal elements were set to

zero to avoid ROI self-connectivity interactions.

Low-resolution

Each individual subject functional connectivity matrix, i.e. an adjacency matrix of

2566x2566 elements, was threshold at 0.19 in order to keep only positive significant

links as described for the calculation of the average high-resolution functional

connectivity matrix FC^{HR}. We call the individual high resolution functional matrix as

$subFC^{HR}$. Next for each subject we calculated the individual low-resolution functional

connectivity matrix ($subFC^{LR}$) as described in the following. Given a pair of macro-

circuits, MAC_i and MAC_j *(*with *i≠j)*, and the two sets of ROIs belonging to them $R_i=\{ROI_{i1},$

… , ROI_{iN}} and $R_j=\{ROI_{j1}, … , ROI_{iM}\}$, where *N* and *M* are the respective number of ROIs,

the average of all functional links in $subFC^{HR}$ connecting the two sets defined the

functional link between MAC_i and MAC_j, i.e. the low-resolution matrix element $subFC_{ij}^{LR}$

according to:

$$subFC_{ij}^{LR} = \frac{1}{N * M} \sum_{a \in R_i} \sum_{b \in R_j} subFC_{ab}^{HR}$$

803   Note that for FC^{LR} the principal diagonal has non-zero elements, representing the

804   internal connectivity within a given MAC, i.e.:

805
$$subFC_{ii}^{LR} = \frac{1}{N*(N-1)} \sum_{a \in R_i} \sum_{b \in R_i} subFC_{ab}^{HR}$$

In relation to supplementary figure 3, given a MAC, we call internal links (*int-links*) the

subset of links connecting the ROIs belonging to the MAC. Similarly, we define external

links (*ext-links*) all the links connecting the same ROIs to the rest of the ROIs in the brain.

In relation to the structural connectivity, we considered the average adjacency high-

resolution matrix SC^{HR}, and we built the low-resolution structural connectivity matrix,

SC^{LR}, where the MACs' volumes (V_i and V_j) were now used for normalization, i.e.:

812
$$SC_{ij}^{LR} = \frac{1}{V_i \times V_j} \sum_{a \in R_i} \sum_{b \in R_j} SC_{ab}^{HR}$$

813 and for the diagonal elements:

814
$$SC_{ij}^{LR} = \frac{1}{V_i^2} \sum_{a \in R_i} \sum_{b \in R_i} SC_{ab}^{HR}$$

To calculate the *MAC$_i$* volume, *V*, we summed the volumes of the corresponding ROIs

(set *R$_i$*) expressed as the number of voxels.

COMPLEX NETWORKS ANALYSIS

Through their construction, all the networks analysed were symmetric and weighted,

with non-negative values. We used MATLAB (Mathworks) to run different metrics

implemented   in   the   Brain   Connectivity   Toolbox

(*https://sites.google.com/site/bctnet/measures/list*). In particular, we initially chose

Node Strength (NS), Eigenvector Centrality (EC), Subgraph Centrality (SC), Betweenness

Centrality (BC) and Clustering Coefficient (CC) and we discarded the node degree (as this

is not meaningful in fully connected conditions), the flow coefficient (which is not

applicable in fully connected conditions), the local efficiency (diverging computational

time high number of nodes) and Pagerank Centrality (as this is a close variant of the EC).

Finally, being centrality measures very highly correlated in most networks including the

brain[59] we chose eigenvector centrality as main metric in our study being the one

showing less variability when considering all link weights or when applying thresholds

to adjacency matrices (as described previously in the Methods at the "high resolution"

section).   To remove the potential bias introduced by ROIs with a stronger influence of

white or grey matter, each individual partial volume of white and grey matter was

estimated using the FSL FAST tool(59). For each functional ROI the average white and

grey matter partial volume estimates were computed from the group average. We used

a Spearman partial correlation to calculate the correlation between structural

(functional) centrality and FirsT, while regressing out white (grey) matter partial volume

estimates per ROI or per MAC (considering the average across the ROIs within each

MAC) in the case of low- and high-resolution networks respectively.

The difference in neurogenesis time between two ROIs or MACs was defined as ΔFirsT.

With regards the average connection probability and weight between nodes, as a

function of the difference in neurogenesis time ΔFirsT (see fig. 2), the matrix

representing ΔFirsT for pairs of ROIs (ΔFirsT^{HR}) was first computed according to:

$$\Delta FirsT_{ab}^{HR} = FirsT_a - FirsT_b$$

where *(a,b)* is a pair of ROIs.

Similarly, the matrix was calculated for the low-resolution network case for pairs of

MACs (ΔFirsT^{LR}). Note that by construction, ΔFirsT^{HR} and ΔFirsT^{LR} are antisymmetric. In

the case of the high-resolution networks, the Spearman correlation of connection

probability and average link weight as a function of ΔFirsT was only calculated

considering non-negative elements of ΔFirsT^{HR} and the corresponding elements in the

SC^{HR} (FC^{HR}: note that the same results would be obtained for the non-positive case),

When F is the set of elements in the matrix satisfying the condition ΔFirsT^{HR}=D (with

0<=D<=max(ΔFirsT)=26 days), the structural (functional) connection probability for

ΔFirsT=D was calculated as the fraction of existing links out of the set F in the adjacency

matrix SC^{HR} (FC^{HR}). Similarly, for the same ΔFirsT=D and set F, the average link weight

was calculated from the adjacency matrix SC^{HR} (FC^{HR}) for the existing links. In the low-

resolution case, since SC^{LR} and FC^{LR} are not a sparse matrix and they have a connectivity

density above 95%, only the link weight was calculated as a function of ⵧFirsT. As for the

high-resolution networks, the Spearman correlation between the link weights and ⵧFirsT

was calculated on the non-negative links of the SC^{LR} (FC^{LR}) and the corresponding

elements in (ΔFirsT^{LR}).

GENETIC FINGERPRINT OF BIRTHDATE CIRCUITS

To investigate the genetic fingerprints of the circuits' birthdays, we used the

transcriptome dataset from the AHBA(60,61). The AHBA provides whole-brain genome-

wide expression values for 20,737 protein encoding genes extracted from 3,702 brain

samples and distributed spatially over six human post-mortem brains. Using the brain

sample information, brain maps representing the spatial distribution of each gene in the

18 circuits was generated based on recent recommendations([60,61]63): i) expression

values from multiple probes were averaged; ii) each sample was mapped to one of the

18 circuit atlases, with samples falling outside mapped to the nearest circuit if this was

closer than 3 mm; iii) for each individual the median expression values were calculated

across all samples that mapped to the same circuit; iv) for each donor a z-score was

applied to normalize each gene spatial distribution and minimize inter-subject variation;

v) a group expression map was computed by calculating the mean expression values of

the six individual donors. While AHBA dataset has not enough spatial coverage for the

computation of genetic expression values at each of the 2,566 regions, we generated a

90 regions parcellation to gain more detailed expression maps compared to the 18

regions. The expression atlas containing 90 brain regions was based on the 68 cortical

regions of the Desikan-Killiany atlas, the 16 subcortical regions from freesurfer, the

cerebellum, brainstem, LC and the dorsal raphe nucleus (see Supplementary figure 6).

This 90 regions' atlas was used to assess the genetic associations of the structural and

functional organization of the brain. The centrality of the 2,566 values falling into each

90 regions was averaged to transform 2,566 functional and structural centrality maps to

90 regions.

To search the underlying genetic fingerprints of both the microcircuit FirsT, and the brain

functional and structural centrality, a combined score for each of the 20,737 protein-

coding genes was computed.

889
$$r(g) = \frac{r_{FirsT,90}(g) - r_{SCC,90}(g) + r_{FCC,90}(g)}{3}$$

The score r for a given gene g is the average Spearman correlation between the

expression of gene g with the FirsT projected to 90 regions $r_{FirsT,90}(g)$, with structural

connectivity centrality in 90 regions $r_{SCC,90}(g)$ and with functional connectivity

centrality in 90 regions $r_{FCC,90}(g)$. The sign of the Spearman correlation of structural

connectivity was inverted as it follows an inverse pattern relative to the FirsT and

functional connectivity. Genes with a score >1.64 standard deviations from the

distribution obtained were used for further analyses (r>0.43; 1,256 genes).

An overrepresentation analysis was used to find common functional annotations in the

obtained gene list. Gene set enrichment was computed using Metascape[62]

(http://www.metascape.org/) for biological processes and cell component annotations

from Gene Ontology, human diseases annotations from DisGeNET[63], and cell type

signatures [62]. Only terms surviving to multiple comparisons are reported; FDR q<0.05.

We used Monte Carlo simulation to assess significance of each of the obtained

significant annotations, to overcome false-positives bias due to gene co-expression

spatial autocorrelation[64]. First, we used BrainSMASH[65] to generate 1,000 surrogate

maps with matching spatial autocorrelation with 90 regions FirsT, and structural and

functional centrality maps. Then, for each surrogate map we computed the spatial

similarity (Spearman correlation) with 20,737 protein coding genes and we used the list

of genes with a similarity >1.64 standard deviations from the obtained distribution to

perform an over-representation analysis. For each of the annotations we quantified how

many times the enrichment value was higher in the original data compared to the

surrogate maps and we generated a p-value of the significance. All the annotations that

were not significant (FDR-corrected) were removed.

Regarding genetic and neurological diseases, our analysis was based on the full gene

lists related to Epilepsy, ASD, PD and AD as reported by the Genome-Wide Association

Study (GWAS) and downloaded from https://www.ebi.ac.uk/gwas/ (GWAS). For these

lists of genes, we used the corresponding relevance score for each disease from

Genecards database (www.genecards.org). . Given a pathology, we normalized the

relevance scores dividing by its maximum, in this way we could use a common index

wchi could be plot across pathologies (see normalized relevance score in fig. 5) For a

detailed definition of the relevance score relative to the disease, see

https://www.genecards.org/Guide/Search. The distribution of the genes from the

GWAS Catalogue (figure 5) was analysed in intervals of five percentiles. The enrichment

for GWAS Catalogue genes in the outermost percentiles vs the rest of the distribution

was tested using the previously generated thousand randomizations that preserved

both spatial dependencies in the variable representing neurogenesis, functional or

structural centrality. In regard to gene expression dependency, the correlation in the

expression of the genes in the brain parcellation was preserved in each random trial by

reshuffling in the same order the values of the expression across the brain parcellation.

Given one thousand randomizations, p-values reported in figure 5 C were calculated

with a 0.001 resolution.

The lists of genes from the GWAS list in the outermost percentiles (see vertical black

broken lines in Fig. 5A) are reported as supplementary tables, with information on the

structural and functional correlations (value, rank and percentile), and the relevance

score (value, rank and percentile).

METHODS REFERENCES

49. Rolls, E. T., Huang, C.-C., Lin, C.-P., Feng, J. & Joliot, M. Automated anatomical labelling

atlas 3. *NeuroImage* **206**, 116189 (2020).

50. Nowinski, W. L. Evolution of Human Brain Atlases in Terms of Content, Applications,

Functionality, and Availability. *Neuroinform* **19**, 1–22 (2021).

51. Nieuwenhuys, R. & Puelles, L. *Towards a New Neuromorphology*. (Springer International

Publishing, 2016). doi:10.1007/978-3-319-25693-1.

52. Nieuwenhuys, R. & Nieuwenhuys, R. Principles of Current Vertebrate Neuromorphology.

Brain Behav Evol **90**, 117–130 (2017).

53. Glasser, M. F. *et al.* The minimal preprocessing pipelines for the Human Connectome

Project. *NeuroImage* **80**, 105–124 (2013).

54. Desikan, R. S. *et al.* An automated labeling system for subdividing the human cerebral

cortex on MRI scans into gyral based regions of interest. *NeuroImage* **31**, 968–980

(2006).

55. Pauli, W. M., Nili, A. N. & Tyszka, J. M. A high-resolution probabilistic in vivo atlas of

human subcortical brain nuclei. *Sci Data* **5**, 180063 (2018).

56. Edlow, B. L. *et al.* Neuroanatomic Connectivity of the Human Ascending Arousal System

Critical to Consciousness and Its Disorders. *J Neuropathol Exp Neurol* **71**, 531–546

(2012).

57. Diez, I. *et al.* A novel brain partition highlights the modular skeleton shared by structure

and function. *Sci Rep* **5**, 10532 (2015).

58. Bonifazi, P. *et al.* Structure–function multi-scale connectomics reveals a major role of

the fronto-striato-thalamic circuit in brain aging. *Hum Brain Mapp* **39**, 4663–4677

(2018).

59. Zhang, Y., Brady, M. & Smith, S. Segmentation of brain MR images through a hidden

Markov random field model and the expectation-maximization algorithm. *IEEE Trans.*

Med. Imaging **20**, 45–57 (2001).

60. Hawrylycz, M. J. *et al.* An anatomically comprehensive atlas of the adult human brain

transcriptome. *Nature* **489**, 391–399 (2012).

61. Shen, E. H., Overly, C. C. & Jones, A. R. The Allen Human Brain Atlas. *Trends in*

Neurosciences **35**, 711–714 (2012).

62. Arnatkevičiūtė, A., Fulcher, B. D. & Fornito, A. A practical guide to linking brain-wide

gene expression and neuroimaging data. *NeuroImage* **189**, 353–367 (2019).

63. Diez, I. & Sepulcre, J. Unveiling the neuroimaging-genetic intersections in the human

brain. *Current Opinion in Neurology* **Publish Ahead of Print**, (2021).

64. Zhou, Y. *et al.* Metascape provides a biologist-oriented resource for the analysis of

systems-level datasets. *Nat Commun* **10**, 1523 (2019).

65. Shannon, P. *et al.* Cytoscape: A Software Environment for Integrated Models of

Biomolecular Interaction Networks. *Genome Res.* **13**, 2498–2504 (2003).

ACKNOWLEDGEMENTS

We thank M. De Pittá, D. Papo, D. Marinazzo, A. Mazzoni, Y. Ben-Ari for helpful

suggestions and critical comments.

Funding: ANR was supported by grant MTM2017-86061-C2-2-P funded

by MCIN/AEI/10.13039/501100011033 and "ERDF A way of making Europe"; PB

acknowledge financial support from Ikerbasque (The Basque Foundation for Science)

and from the Ministerio Economia, Industria y Competitividad (Spain) and FEDER (grant

SAF2015-69484-R and AI-2021-039). Authors declare that they have no competing

interests. F.G.-M. is supported by Ikerbasque, and holds Spanish Ministry MICINN

PGC2018-096173-A-I00 and PID2021-125156NB-I00 grants and a Basque Government

PIBA_2022_1_0027 grant.

Author contributions: Conceptualization: FGM, PB; Methodology: ID, FGM, NCS, SS,

ANR, MDA, JMC, PB; Writing – original draft: FGM, ID, MDA, JMC, PB; Funding

acquisition: FGM, JMC, PB.

ADDITIONAL INFORMATION

Competing interests

The authors declare no competing interests.

Supplementary materials

Supplementary Information is available for this paper.

Fig S1 – S6

References (T01 to T18)

Auxiliary supplementary tables 01-13

Correspondence and requests for materials should be addressed to Dr. Paolo Bonifazi.

Reprints and permissions information is available at www.nature.com/reprints.

1000

1001

1002   **FIGURES**

Figure 1

1003

Figure 1. From the circuits' embryogenic age to brain networks and transcriptomics.

(A) Sagittal schemes of the early embryonic human brain. The upper scheme represents

the neuromeres and fundamental regions of development, the lower scheme displays

the location of the 18 MACs within the early embryonic human brain, colour-coded

according to their FirsT. (B) Scheme of the neuroimage analysis pipeline. Brain networks

were reconstructed from healthy adult subjects scanned by MRI at 7T within the HCP

initiative. For each subject, the structural and functional networks were reconstructed

using probabilistic tractography and resting-state activity, respectively. For every pair of

ROIs (2,566 ROIs), the structural connectivity matrix represents the putative number of

fibres connecting them, while the functional connectivity matrix reports the correlation

in their activity as revealed by the BOLD time series. The averages across subjects were

used as final representative brain networks (see Methods). From the 2566 ROIs'

matrices, the 18 MACs' networks were reconstructed and represents for each pair of

MACs the average links between the ROIs forming them (see Methods). To identify

patterns between networks' topology and neurogenesis, the correlation between the

nodes' metrics (centrality or segregation) and embryonic age was calculated. (C) Brain

transcriptome data from AHBA dataset (see methods) was used to search for protein

coding genes with a high similarity between its spatial brain expression and embryonic

age. Functional annotations of the obtained genes were further computed using

overrepresentation analysis to find significantly associated biological processes and

cellular components associated with embryonic age.

Figure 2

1025

Figure 2. Scatter plots of the nodes' centrality and the links' probability/weight

relative to the FirsT, in the high and low resolution structural-functional networks. (A)

Scatter plots of the nodes' centrality as a function of embryonic age. Red and blue

indicates the structural and functional cases, respectively. The first and second columns

from left represent the results for the 2,566 ROIs as violin plots, where the mean of each

group (i.e. ROIs within a MAC) is plotted in black, and the third and fourth columns

represents the 18 MACs. All plots have a logarithmic scale in the y-axis. (B) Scatter plots

of the link weights (left y-axis) as a function of the differences in the first neurogenic

birthdate. The colours and columns are as indicated in panel A. The violin plots for the

2,566 ROIs show the link weight distributions while the solid blue/red lines plot the

average values. Black solid lines show the exponential (linear) fit on the average values

for the structural (functional) case. The link probability is represented by the right y-axes

and displayed as dots, and the broken black line showing the exponential fit. All plots

have a logarithmic scale in the y-axis.

Figure 3

1040

Figure 3. Visualization of brain connectivity and the first neurogenic birthdate. (A) The

colour coded glass brain maps of the eighteen MACs represents the FirsT (A1; see colour

bar on the left) and the z-score (see colour bar on the right) of the eigenvector centrality

for the SC^{LR} (A2) and the FC^{LR} (A3). (B) Linear temporal representation with arches of the

MACs structural connectivity for the earliest (B1) generated MAC (Locus Coeruleus, FirsT

28), for a representative mid-term (B2) generated MAC (Hippocampus, FirsT 40) and for

the latest (B3) generated MAC (Occipital Cortex, FirsT 54). The x-axis represents the

developmental time of several FirsT, where the different MACs are located according to

their FirsT. Note that the Basal Ganglia and Cerebellum overlap since their neurogenesis

is initiated on day 36. The thickness of top black arches is proportional to the link weight

between the MACs extracted from the 18 node adjacency matrix. Broken black lines

mark the presence of connections but with little weight (i.e. below the lowest threshold

for the visualization of solid lines).

Figure 4

1054

Figure 4. Transcriptome of brain circuits according to embryogenic age. (A) Mean

spatial similarity between the expression of 20,737 protein-coding genes with the time

of birth and structural and functional centrality. Structural and functional centrality of

the 2,566 brain regions was downsampled to 90 region (with gene expression data) and

for each of the 90 regions one of the 18 FirsT was assigned. (B) 1,256 genes from the

positive tail (6% of protein coding genes) – genes whose spatial expression is most

similar to FirsT and functional and structural centrality – were used to compute the

overrepresentation analysis for different annotations: biological processes, cell

components, cell types, and human diseases.

Figure 5

1064

Figure 5. Correlations between the maps of the gene transcriptome, neurogenesis and hubness in relation to Autism Spectrum Disorder, epilepsy, Parkinson's and Alzheimer's disease. For each gene the spatial expression in the brain was correlated to the spatial maps of eigenvector centrality and neurogenesis (note that from the sub-sampled brain transcriptomes, 90 brain regions were considered: see Methods). **A** Probability distribution of the correlations (light gray histogram, left axis) between transcriptomics (20,787 genes) and neurogenesis (top plot), functional centrality (middle plot) and structural centrality (bottom plot). The normalized relevance score of the GWAS genes listed for each disease (obtained from the GeneCards databases; Note that normalization is computed separately for each disease; see Methods) has been plotted with dots (right axis). The colors used to display the results for the different diseases in pnale A-C, are shown in the inset legend of the top plot of panel B. . Vertical broken lines highlight the lowest and top five percentile interval. Coloured labels and

arrows mark the most relevant representative GWAS genes with correlations in the

highlighted extreme percentiles.**B** For each disease, the normalized count (percentage

of gene per interval) of GWAS genes in five percentile intervals of the correlation

distribution (light gray histogram of panel A) is plotted for the different diseases (legend

of colours is in the inset of the top plot)). **C** P-value of the normalized gene count in each

percentile. The threshold for p<0.01 calculated from the null model (based on a

thousand reshuffled replications) which keeps into account spatial dependencies of

both centrality maps and gene-expression is plotted as a broken horizontal black line.

The arrows mark the extremities of the correlation distributions with a significantly

higher number of GWAS-genes than what expected by chance.