Explaining flexible continuous speech comprehension from individual motor rhythms

Christina Lubinus[1], Anne Keitel[2], Jonas Obleser[3,4], David Poeppel[5,6,7,] and Johanna M. Rimmele[1,6]

[1]Department of Neuroscience and Department of Cognitive Neuropsychology
Max-Planck-Institute for Empirical Aesthetics
60322 Frankfurt am Main, Germany,

[2]Psychology
University of Dundee,
Dundee DD1 4HN, UK,

[3]Department of Psychology
University of Lübeck,
Lübeck, Germany,

[4]Center for Brain, Behavior, and Metabolism
University of Lübeck,
Lübeck, Germany,

[5]Department of Psychology
New York University,
New York, NY, USA

[6]Max Planck NYU Center for Language, Music, and Emotion
Frankfurt am Main, Germany, New York, NY, USA

[7]Ernst Strüngmann Institute for Neuroscience (in Cooperation with Max Planck Society)
Frankfurt am Main, Germany

Corresponding author:

Christina Lubinus
Email: christina.lubinus@ae.mpg.de
Phone: +49 69 8300 479-348

Abstract

When speech is too fast, the tracking of the acoustic signal along the auditory pathway deteriorates, leading to suboptimal speech segmentation and decoding of speech information. Thus, speech comprehension is limited by the temporal constraints of the auditory system. Here we ask whether individual differences in auditory-motor coupling strength in part shape these temporal constraints. In two behavioral experiments, we characterize individual differences in the comprehension of naturalistic speech as function of the individual synchronization between the auditory and motor systems and the preferred frequencies of the systems. Obviously, speech comprehension declined at higher speech rates. Importantly, however, both higher auditory-motor synchronization and higher spontaneous speech motor production rates were predictive of better speech-comprehension performance. Furthermore, performance increased with higher working memory capacity (Digit Span) and higher linguistic, model-based sentence predictability – particularly so at higher speech rates and for individuals with high auditory-motor synchronization. These findings support the notion of an individual preferred auditory–motor regime that allows for optimal speech processing. The data provide evidence for a model that assigns a central role to motor-system-dependent individual flexibility in continuous speech comprehension.

Keywords: speech perception; speech production; auditory-motor synchronization; oscillations; audiomotor.

1. Introduction

Speech comprehension relies on temporal processing, as speech and other naturalistic signals

have a complex temporal structure with information at different timescales(1). The temporal

constraints of the auditory system limit our ability to understand speech at fast rates(2,3). In-

terestingly, the motor system can under certain conditions provide temporal predictions that

aid auditory perception(4,5). Accordingly, current oscillatory models of speech comprehension

propose that properties of the auditory but also the motor system affect the quality of auditory

processing(6,7). In two behavioral experiments, we investigate how the auditory, the motor

system, and their synchronization shape individual flexibility of comprehending fast continuous

speech.

Auditory temporal constraints have been observed as preferred rates of auditory speech(8,9)

processing (but also of tones(10,11), and amplitude modulated sounds(11–14)) and explained

in the context of neurocognitive models of speech perception. According to such proposals,

humans capitalize on temporal information by dynamically aligning ongoing brain activity in

auditory cortex to the temporal patterns inherent to the acoustic speech signal(15–18). By hy-

pothesis, endogenous theta brain rhythms in auditory cortex partition the continuous auditory

stream into smaller chunks at roughly the syllabic scale by tracking quasi-rhythmic temporal

fluctuations in the speech envelope. This chunking mechanism allows for the decoding of seg-

mental phonology – and ultimately linguistic meaning(15,18–20). The decoding of the speech

signal is accomplished seemingly effortlessly within an optimal range centered in the traditional

theta band(18), whereas comprehension deteriorates strongly for speech presented beyond

~9 Hz(2,3). While much research has focused on the apparent *stability* of the average acoustic

modulation rate at the syllabic scale(8,9), the *flexibility* in speech comprehension(9,21), that

is, what constitutes individual differences in understanding fast speech rates, is poorly under-

stood.

The motor system, and neural auditory-motor coupling in particular, is a plausible candidate to

facilitate individual differences in auditory speech processing abilities. Two arguments sup-

porting this notion are the motor systems' modulatory effect on auditory perception(22–24) and

its susceptibility to training(25–27). While there is evidence suggesting that the auditory and

speech motor brain areas are intertwined during speech comprehension(28–32), the extent to

which speech motor processing modulates auditory processing is debated(5,33,34). Specifi-

cally, endogenous brain rhythms in both auditory(20,35) and motor(35,36) cortex have been

observed to track the acoustic speech signal, and are characterized by preferred frequencies(19,37,38). In contrast to neural measures of preferred frequencies(37–39), here we used a behavioral estimate termed "preferred" or "spontaneous" rate. Furthermore, neural coupling between auditory and motor brain areas during speech processing(35,36,40,41) has been hypothesized to provide temporal predictions about upcoming sensory events to the auditory cortex(4,41–43). The precision of these predictions may be proportional to the strength of auditory-motor cortex coupling.

Auditory-motor cortex coupling strength varies across the population, as shown by recent work(6,10,40,44,45). Assaneo et al.(40) developed a behavioral protocol (spontaneous speech synchronization test; SSS-test) which quantifies the strength of auditory-to-motor synchronization during speech production in individuals. The authors reported that auditory-motor synchronization is characterized by a bimodal distribution in the population, classifying individuals into high versus low synchronizers. (The rejection of unimodality has been previously shown with large sample sizes(40, see also: ,46).) Importantly, in addition to superior behavioral synchronization, high synchronizers have stronger structural and functional connectivity between auditory and speech motor cortices (see 40, Figure 3A and B). Thus, the SSS-test provides not only a behavioral measure but also approximates individual differences in neuronal auditory-motor coupling strength. We propose that the individual variability in auditory-motor synchronization, previously observed to predict differences in word learning(40), syllable detection(6), and rate discrimination(10), as well as the individual variability in preferred auditory and motor rate, predicts differences in an individuals' ability to comprehend continuous speech at fast syllabic rates.

The influence of individual auditory-motor coupling strength on behavioral performance has so far been established for behavioral paradigms using rather basic auditory and speech stimuli, e.g. tones or syllables(6,10,40). The current study assesses its importance in a more naturalistic context: during the comprehension of continuous speech. This adds several layers of complexity. First, as speech unfolds over time, processing of continuous, i.e. longer and more complex, speech naturally demands more working memory capacity for maintenance and access to linguistic and context information(47). Second, rich linguistic context is used to derive linguistic predictions about upcoming words and sentences(48–51). When linguistic predictability of a sentence is high(52), speech comprehension is improved, even in adverse listening situations(53,54). Thus, similar to auditory-motor synchronization, linguistic predictability offers a compensatory mechanism when comprehension is difficult.

In summary, we investigate the role of auditory-motor synchronization with the SSS-test and

the role of preferred rhythms of the auditory and motor systems for the individual flexibility of

the comprehension of continuous speech. First, based on an established literature(3,18,55–

57), we expected a decline in comprehension performance at syllabic rates beyond the theta

range. Second, as a faciliatory effect of auditory-motor coupling on auditory processing has

been observed(6,10,40), we hypothesized that individual differences in comprehension perfor-

mance could be predicted by individual auditory-motor synchronization, with superior speech

comprehension for high synchronizers. Such a faciliatory effect might be strongest in demand-

ing listening situations, such as at fast syllabic rates(5,10). Third, while the consequences of

potential individual variation in the preferred rates of the motor and auditory systems are not

clearly understood, based on previous findings(35) we expected a systematic relation of both

preferred auditory and motor rates with individual speech comprehension performance. Fi-

nally, we hypothesized that linguistic predictability and working memory span should positively

affect speech comprehension. Similar to auditory-motor synchronization, we expected linguis-

tic predictability to interact with syllabic rate, such that both systems would become stronger

predictors for speech comprehension as syllabic rate increases.

2. Methods

Two behavioral experiments and a control experiment were conducted: Experiment 1 was per-

formed in the laboratory and investigated the influence of the spontaneous speech motor pro-

duction rate on speech comprehension performance. In Experiment 2 we aimed to understand

the complex interplay of multiple variables during speech comprehension beyond the sponta-

neous speech motor production rate. To this end, we additionally measured participants' pre-

ferred auditory rate, auditory-motor synchronization, and working memory capacity. Experi-

ment 2 and the control experiment were online studies. All studies were approved by the local

ethics committees (Experiment 1: committee of the School of Social Sciences, University of

Dundee, UK (No. UoD-SoSS-PSY-UG-2019-88), Experiment 2 and control experiment: pro-

cedures were approved by the Ethics Council of the Max Planck Society (2017_12)).

      2.1. Participants

Participants were English native speakers with normal hearing and no neurological or psycho-

logical disorders (Exp 1: N = 34, Exp 2: N = 82, Control: N = 39). Participation was voluntary.

For a detailed description of participants, stimuli, exclusion criteria, and tasks please refer to

Supplementary Methods, Figures 1-2, and Tables 1-2.

2.2. Design and materials

Speech comprehension task

In two speech comprehension tasks, we measured participants ability to comprehend sen-

tences at various syllabic rates. Sentences were presented at 7 (Exp 1: [8.2, 9.0, 9.8, 11.0,

12.1, 14.0, 16.4]) or 6 (Exp 2: [5.00, 10.69, 12.48, 13.58, 14.38, 15.00]) rates. In Experiment

1, participants performed a classic intelligibility task, i.e. also termed "word identification task"

(58,59, for review: ,60). On each trial (N = 70), a sentence was presented through headphones

and participants verbally repeated the sentence as accurately as possible (Fig. 1A). Re-

sponses were recorded.

In Experiment 2, speech comprehension was measured by a word-order task. Participants

listened to one sentence per trial (N = 240), followed by the presentation of two words from the

sentence on screen. Participants indicated via button press which word they heard first (Fig.

2A).

Speech production task

In the speech production tasks we estimated participants individual spontaneous speech motor

production rate. In Experiment 1, the speech production task was operationalized by partici-

pants reading a text excerpt (216 words) from a printout. Participants were instructed to read

the text excerpt out loud at a comfortable and natural pace while their speech was recorded

(Fig. 1B).

In Experiment 2, participants were asked to produce continuous, "natural" speech. To facilitate

fluent production, they were prompted by a question/statement belonging to six thematic cat-

egories (6 trials; own life, preferences, people, culture/traditions, society/politics, general

knowledge, see Supplementary Table 2). Each response period lasted 30 seconds and trials

were separated by self-paced breaks (Fig. 2C). While speaking, participants simultaneously

listened to white noise. The white noise was introduced to measure the preferred rate of the

motor system, without potential interference from auditory feedback. A second reason was to
be consistent with the protocol from the SSS-test(40,61; also see below). Note that this proce-
dure was not applied in Experiment 1.

Auditory rate task (only Exp 2)

To measure participants preferred auditory rate, we implemented a two-interval forced choice
(2IFC) task, presenting a reference and a comparison stimulus at each trial. Participants indi-
cated via button press which stimulus they preferred (Fig. 2B). Stimuli were presented at syl-
labic rates from 3.00 to 8.50 syllables/s (3.00, 3.92, 4.83, 5.75, 6.67, 7.58, 8.50). A reference
rate, e.g. 3.00 syllables/s, was compared to all syllabic rates, including itself. For each refer-
ence/comparison pair the same sentence was presented – the stimuli only differed in their
syllabic rates. Additionally, the task included catch trials to measure participant's engagement
(see Supplementary Methods for details).

Spontaneous speech synchronization (SSS) test (only Exp 2)

We measured participant's auditory-motor synchronization using the SSS-test (for details: ,40).
In the main task, participants listened to a random syllable train and whispered along for a
duration of 80s. They were instructed to synchronize their own syllable production to the stim-
ulus presented through their headphones (Fig. 2D). The syllable rate in the auditory stimulus
progressively increased in frequency from 4.3 to 4.7 syllables/s in increments of 0.1 syllables/s,
every 60 syllables. Participants completed two trials, while the whispering was recorded.

Participants' syllable production was masked by the simultaneously presented auditory sylla-
ble train. The masking procedure suppresses auditory feedback, allowing us better to isolate
the synchronization of motor production to the auditory input, without interference of auditory
feedback(44).

Digit span test (only Exp 2)

Working memory capacity was quantified using the forward and backward(62) digit span test.
As for the backward test data is missing for N = 21 participants, only the forward span is re-
ported. Digit spans were presented auditorily and participants typed in their responses(63).

Control Experiment

We designed a control experiment to test if the correct word order from the word order task of

Experiment 2 could be guessed from the target words alone, that is, without understanding the

sentence. The task consisted in judging which of two words would be more likely to occur first

in a *hypothetical sentence*. On each trial, two words were presented on screen and participants

indicated their choice via button press. Importantly, 1) participants did not listen to a full sen-

tence at any time and 2) the target words were taken from the stimulus materials actually

presented in Experiment 2.

2.3. Analysis

Spontaneous speech motor production rate (Exp 1 + 2)

The individual *spontaneous speech motor production rate*, i.e. articulation rate(64), was com-

puted using Praat software(65) by automatically detecting syllable nuclei. The number of syl-

lable nuclei was divided by the duration of the utterance, disregarding silent pauses. For Ex-

periment 1, the production rate was computed across the entire reading paragraph. For Ex-

periment 2, it was first calculated for each trial (30 s) separately. The motor rate was then

averaged across all trials.

Preferred auditory rate (Exp 2)

First, participants with low performance in the catch trials of the preferred auditory rate task

(below 75% correct) were excluded; amongst the remaining participants ($N = 82$) catch trial

performance was very high ($M = 98.48\%$, $SD = 3.71$). To compute the preferred auditory rate,

a distribution of preferred frequencies was derived from all trials -except catch trials- by aggre-

gating the frequency of each trials' preferred item. Then a gaussian function was fitted to each

participants' distribution and two parameters were extracted: the peak as index for the pre-

ferred frequency and the full-width-at-half-maximum (FWHM) as index for the specificity of the

response (lower FWHM equals stronger preference for one frequency).

Auditory-motor synchronization (Exp 2)

From the SSS-test(40) we derived the participant's auditory-motor synchronization by calcu-

lating the phase-locking value (PLV)(66) between the (cochlea) envelopes of the auditory and

the speech signals.

180
$$PLV = \frac{1}{T} \left| \sum_{t=1}^{T} e^{i(\theta_1(t) - \theta_2(t))} \right| \tag{1}$$

where T is the total number of time points, t denotes the discretized time, and θ_1 and θ_2 are
the phase of the first and the second signals, respectively.

To obtain the cochlear envelope of the syllable train (auditory channels: 180–7,246 Hz), we
used the Chimera Software toolbox(67). For the recorded speech signal the amplitude enve-
lope was quantified as the absolute value of the Hilbert transform. Both envelopes were
downsampled to 100 Hz and bandpass filtered (3.5-5.5 Hz) before their phase was extracted
by means of the Hilbert transform. The PLV was first estimated for each trial of the SSS-test
(time windows 5s, overlap 2s) and then averaged across runs, resulting in a mean PLV. The
distribution of mean PLV values was subjected to a k-means algorithm(68) ($k = 2$) to split
participants into a high- and a low-synchronizer group. Speech auditory-motor synchronization
(PLV) was treated as bimodal variable based on previous research that rejected unimodality
based on larger samples(40, see also: ,46).

Linguistic predictability – Recurrent neural network (Exp 2)
Linguistic predictability of all stimulus sentences was measured by deriving single-sentence
perplexity from a recurrent neural network language model. A language model, such as a re-
current neural network, assigns probabilities to all words in a sequence of words. From the
single-word probabilities, we derived one value per sentence, quantifying its predictabil-
ity(69,70). This so-called perplexity is the most common intrinsic evaluation metric of language
models(71–73). It is computed as the inverse of the mean probability of a sentence weighted
by sentence length(69), i.e. lower perplexity values equal higher sentence predictability (see
Supplementary Methods for full details on RNN and perplexity).

Mixed effects models
For both experiments we performed mixed effects analyses to quantify how speech compre-
hension was affected by all variables of interest. Mixed models were computed using the R
packages *lme4* (v1.1-29) and *mgcv* (v1.8-39), as set up in Rstudio (version 2022.2.1.461).
Mixed-effects, rather than fixed-effects models were chosen to account for idiosyncratic varia-
tion within variables, i.e. repeated measures and therefrom resulting interdependencies

between data points(74,75). Thus, both models included random intercepts for *participant* and

items.

In Experiment 1, we computed a generalized additive mixed-effects model (GAMM) using the

mgcv:gam function. For the dependent variable *speech comprehension*, we calculated the

percentage of correctly repeated words for each sentence and subject from the speech com-

prehension task. The number of correct words was counted manually and transformed into a

percentage. Then the dependent variable (single-trial data) was modelled as a function of the

fixed effects *syllabic rate* and *spontaneous speech motor production rate*. A random slope for

syllabic rate could not be included because the model failed to converge, thus the model in-

cluded only random intercepts. Overall, the model explained ~77% of the variance.

In Experiment 2, the dependent variable *speech comprehension* was binary (*correct* vs *incor-*

rect word order judgment). Thus, we employed a generalized linear mixed-effects model

(GLMM; *lme4:glmer* function) with a binomial logit link function. In terms of fixed effects, the

model included all variables of interest: *syllabic rate*, *preferred motor rate*, *preferred auditory*

rate, *auditory-motor synchronization*, *working memory, sentence predictability*. Additionally,

we introduced several linguistic and other covariates for nuisance control(76): *predictability*

target 1, *predictability target 2, sentence length* (# of words), *target distance* (i.e., distance in

words between the target words), *compression/dilation* of audio file. In addition to random in-

tercepts, the model contained a by-participant random slope for *syllabic rate*, allowing the

strength of the effect of the rate manipulation on the dependent variable to vary between par-

ticipants(74,75). Continuous predictor variables were z-transformed to facilitate the interpreta-

tion and comparison of the strength of the different predictors(77). Thus, the coefficients of all

continuous predictors reflect log changes in comprehension for each unit (*SD*) increase in a

given predictor. We observed no problems with (multi-)collinearity, all variance inflation factors

were < 1.2 (package car version 3.0-10(78)). Overall, the model explained ~38% of the vari-

ance.

Control experiment

For each trial, we computed how many participants correctly guessed the word order (in per-

cent, "*word order index*"). In a new GLMM analysis, this *word order index* was added as co-

variate into the model from the main analysis while all other parameters remained the same.

3. Results

3.1. Experiment 1

In Experiment 1, we asked the question: to what extent is speech comprehension affected by one's spontaneous speech motor production rate? *Speech comprehension* was measured as the percentage of correctly repeated words in an intelligibility task (2.75% to 93.70% on average across participants). We observed a mean *spontaneous speech motor production rate* of 4.11 syllables per second (*SD* = 0.35, min = 3.35, max = 4.85) across participants (Fig. 1C).

As expected, the GAMM revealed a main effect of *syllabic rate*: slower speech stimuli were associated with better speech comprehension (edf = 4.61, *F* = 1260.90, *p* < .001, Fig. 1D, see Supplementary Table 3). Importantly, we observed that the *spontaneous speech motor production rate* influenced speech comprehension: the higher the individual *spontaneous speech motor production rate*, the better the speech comprehension performance (edf = 1.00, *F* = 4.37, *p* = .036, Fig. 1E).

Figure 1. A. Example trial for the speech comprehension task. Participants fixated on a green fixation dot while presented auditorily with a sentence. On stimulus offset the fixation dot turned red, indicating to commence recall, i.e., reporting the sentence back. **B.** Spontaneous speech motor production rate task. Participants read a stimulus paragraph from a paper. **C.** Spontaneous speech motor production rate. We observed spontaneous speech motor production rates between 3.35 and 4.85 syllables/s (*M* = 4.11 syllables/s, left). The violin and boxplot show summary statistics and density: the median center line, 25th to 75th percentile hinges, whiskers indicate minimum and maximum within 1.5 × interquartile range. Grey dots represent participants individual speech motor productions rates, averaged across 6 trials. **D.** Main effect of syllabic rate. Plot shows the predicted main effect of syllabic rate from the generalized additive mixed model (GAMM). Black line indicates the predicted effect with 95% confidence interval in grey. Black dots show trial-level speech comprehension performance per subject and rate

condition. **E.** Main effect of spontaneous speech motor production rate. Plot shows the predicted main
effect of spontaneous speech motor production rate from the GAMM. Colored lines indicate the pre-
dicted effect with 95% confidence interval in the corresponding color. Colored dots show trial-level
speech comprehension performance per subject and rate condition.

3.2. Experiment 2

First, in line with the first experiment, we observed a mean *spontaneous speech motor pro-*

duction rate of 4.32 syllables per second across participants (SD = 0.45, Min = 3.36, Max =

5.38 syllables per second, Fig. 2G). Within-subject variance was low (Supplementary Fig. 3),

suggesting that participants' articulation rate was stable across trials. Second, participants

showed a *preferred auditory rate* of ~5.57 syllables per second (peak: M = 5.57, SD = 0.86,

Min = 4.16, Max = 7.92; FWHM, M = 4.89, SD = 0.50, Min = 3.23, Max = 5.50; Fig. 2F). Single-

subject raw data can be inspected in Supplementary Fig. 4. Third, *auditory-to-motor speech*

synchronization was quantified using the SSS-test(40), classifying participants as HIGH or

LOW synchronizers (mean PLV HIGHs = 0.73, SD = 0.09, mean PLV LOWs = 0.36, SD = 0.09,

Fig. 2E). Fourth, *working memory* was measured by means of the digit span test (62) which

revealed a mean forward digit score of M = 8.46 (SD = 2.12, Min = 5.00, Max = 13.00, Fig.

2H).

A   Speech comprehension task

fixation   stimulus   garden   house   WO-task
                        targets

B   Preferred auditory rate task

fixation   reference   fixation   comparison   2IFC
           stimulus   stimulus

C   Spontaneous speech motor production rate task

fixation   Describe a traditional   free speech
           Christmas Holiday.   for 30s
           prompt

D   SSS-test

foo beh lah...

te te te ...

E   Auditory-motor synchronization

Figure 2. Panels **A, B,** and **C** visualize example trials for the speech comprehension task, the preferred auditory rate task, and the spontaneous speech motor production task, respectively. **D.** Schematic representation of the SSS-test, used to measure auditory-motor synchronization. Participants whisper a syllable (here /te/). **E.** Histogram of auditory-motor synchronization strength, obtained with the SSS-test. Participants were classified into high and low synchronizers (highs, lows) based on their PLV using k-means clustering. Group affiliation is overlaid by colored lines representing fitted normal distributions. **F.** Participants showed a mean preferred auditory rate of 5.57 syllables/s (*SD* = 0.86), with no differences between high and low synchronizers (*U* = 897.5, *p* = .48). **G.** We observed a spontaneous speech motor production rate between 3.36 and 5.38 syllables/s (*M* = 4.32, *SD* = 0.45) and no group difference between high and low synchronizers (*U* = 767.0, *p* = .60). **H.** Working memory capacity was indicated by a mean digit-span forward score of 8.46 (*SD* = 2.12) and the score did not differ between high and low synchronizers (*U* = 666.5, *p* = .14).

The GLMM revealed that *syllabic rate* significantly influenced participants' comprehension ac-
curacy: for each increase of syllabic rate by one syllable/s, the odds of a correct word order
judgment decreased (*odds ratio (OR)* = 0.65, *std. error (SE)* = 0.04, *p* < .001, Fig. 3A). This
main effect of syllabic rate is consistent with a decline of speech comprehension performance
at higher syllabic rates (3). In line with our hypothesis, we observed main effects for *spontane-*
ous speech motor production rate and *auditory-motor synchronization*. The higher a partici-
pant's *spontaneous speech motor production rate*, the better the performance in the word order
task (*OR* = 1.19, *SE* = 0.09, *p* = .014, Fig. 3C), replicating our finding from the first experiment.
For *auditory-motor synchronization*, being a dichotomous variable (i.e., HIGH vs. LOW)(40),
performance in the word order judgment task was higher for high compared to low synchroniz-
ers (*OR* = 1.34, *SE* = 0.20, *p* = .048, Fig. 3B). That is, across all trials, high synchronizers were
more likely to correctly perform the task. Additionally, the model revealed a positive effect for
working memory score (*OR* = 1.20, *SE* = 0.09, *p* = .012, Fig. 3D). This main effect suggests
that better working memory performance enabled participants to better perform on the speech
comprehension task. We did not observe a reliable effect of *preferred auditory rate* on speech
comprehension (*OR* =1.14, *SE* = 0.08, *p* = .072). In contrast to our hypothesis, we observed
no interaction effect of *syllabic rate* and *auditory-motor synchronization* on speech compre-
hension (*OR* = 0.97, *SE* = 0.07, *p* = .602).

Figure 3. Significant main effects predicting speech comprehension performance. The generalized lin-
ear mixed effects model revealed a negative main effect of syllabic rate (A) and positive main effects of
auditory-motor synchronization (B), spontaneous speech motor production rate (C), and working
memory score (D). For stimulus perplexity we observed a negative main effect (E). In all panels, error
shades indicate 95% confidence intervals. Note that the predictors are shown as a function of syllabic
rate for visualization purposes only.

Linguistic predictability and further linguistic variables

To account for the effect of linguistic attributes, we expanded the GLMM by adding several

(information-theoretic) linguistic variables: *perplexity*, *probability of target words*, *target dis-*

tance, and *stimulus length*. Adding these variables (with linguistic variables, AIC: 12675) im-

proved model fit (without linguistic variables, AIC: 12848), as measured by a likelihood ratio

test ($\chi 2(6) = 184.24$, p < .001, see Supplementary Table 4).

The full GLMM revealed that *perplexity* had a statistically reliable, negative effect on speech

comprehension (*OR* = 0.84, *SE* = 0.04, *p* = .001, Fig. 3E) such that sentences with lower

perplexity (which is equal to higher sentence predictability) lead to better speech comprehen-

sion performance. Additionally, we observed significant negative effects for *probability of target*

word 1 (*OR* = 0.93, *SE* = 0.03, *p* = .026) and *target word 2* (*OR* = 0.92, *SE*: 0.03, *p* = .021).

Contrary to the perplexity effect, this suggests that task performance in the comprehension

task was increased for unexpected target words.

Furthermore, the model revealed a positive effect for *target distance* (*OR* = 1.48, *SE*: 0.05, *p*

< .001), suggesting that larger distance between targets was associated with better speech

comprehension performance. In contrast, suggesting the opposite relation, for *stimulus length*

we observed a negative effect (*OR* = 0.61, *SE*: 0.03, *p* < .001), i.e., shorter sentences resulted

in higher comprehension performance. Due to the large number of variables introduced for

nuisance control, we applied a control for multiple comparisons (i.e. false discovery rate; for

full results see Supplementary Table 5). All effects remained robust after FDR correction: syl-

labic rate: *p* < .001; spontaneous speech motor production rate: *p* = .023; preferred auditory

rate: *p* = .078; working memory score: *p* = .022; perplexity: *p* = .003; probability target 1: *p* =

.034; probability target 2: *p* = .030; compression: *p* < .001; sentence length: *p* < .001; target

distance: *p* < .001. Only auditory-motor synchronization changed from a significant effect to a

trend (*p* = 0.057) (Note that this was a planned comparison and therefore is discussed).

Finally, we explored interaction effects between *syllabic rate*, *auditory-motor synchronization*,

and *perplexity*. Adding the interaction term improved model fit ($\chi^2 (3) = 13.84$, *p* = .004 (AIC

without interaction term: 12675, AIC with interaction term: 12668)). The model revealed two

significant 2-way interaction effects: *syllabic rate × perplexity* (OR = 0.88, SE = 0.05, p = .015)

and *auditory-motor synchronization × perplexity* (OR = 0.86, SE = 0.04, p = .003; see Supple-

mentary Fig. 5 and Supplementary Table 6). The interaction effect between *syllabic rate* and

perplexity indicates that particularly comprehension of sentences at fast syllabic rates im-

proves when *perplexity* is low. Furthermore, the *auditory-motor synchronization × perplexity*

interaction effect suggests that while having better overall speech comprehension, high syn-

chronizers show a stronger effect of *perplexity* compared to low synchronizers, with even better

speech comprehension for more predictable sentences. The *syllabic rate × auditory-motor*

synchronization effect (OR = 0.94, SE = 0.07, p = .392), as tested before, and the three-way

interaction effect of *syllabic rate × auditory-motor interaction × perplexity* (OR = 1.09, SE =

0.06, p = .106) did not show a statistically reliable effect on speech comprehension.

Control experiment

In Experiment 2, speech comprehension performance was exceptionally good, even at high

syllabic rates. To ensure the high performance was not an artifact of the task or stimuli, we

conducted a control experiment. The analysis revealed that *word order index* did not influence

speech comprehension in a statistically meaningful way (OR = 0.96, SE = 0.07, p = .219, see

Supplementary Table 7).

4. Discussion

In two behavioral experiments, we show clear effects of *syllabic rate* on the comprehension of

continuous speech. This finding is in line with proposals of speech comprehension being tem-

porally constrained such that it is optimal for speech at lower syllabic rates. Crucially, in both

protocols we observed that speech comprehension across a wide range of frequencies (5-15

syllables/s) was affected by participants' *spontaneous speech motor production rate*, with

higher rates predicting better speech comprehension. In the second experiment we showed

that, beyond the spontaneous rate of the speech motor system, the individual strength of

speech auditory-motor synchronization also affected comprehension. In contrast, the preferred

speech perception rate was not related to speech comprehension performance. Together,

these findings suggest that while speech comprehension is limited by general processing char-

acteristics of the auditory system, interindividual differences in comprehension flexibility are

moderated by the motor system and interactions between the auditory and motor systems (Fig.

4). Our findings furthermore allow us to generalize the effects of individual differences in the

motor system on auditory perception, which have been previously shown for simpler stim-

uli(6,10,40,79), to more natural continuous speech.

Figure 4. Schematic illustrating the relationship between speech comprehension performance and au-
ditory-motor synchronization (A), the preferred rates of the motor (B) and of the auditory systems (C).
All three predictor variables are represented by the corresponding distribution generated from our ex-
perimental data. We propose that better speech comprehension at demanding rates - and by hypothesis
auditory behavior more generally - is accompanied by a higher preferred rate of the motor system as
well as stronger auditory-motor synchronization. In contrast, the preferred rate of the auditory system
seems not to determine auditory behavior. Circled A and M illustrate the auditory and motor systems.
The arrows connecting them express the relevance of synchronization between the systems for the
variable in question.

As expected(2,18,55–57), we observed that speech comprehension accuracy declined as syl-

labic rate increased. Although speech comprehension dropped at higher rates in both para-

digms, the overall level of comprehension accuracy was much higher in Experiment 2, with

accuracy remaining very high (~85%), even for speech as fast as 15 syllables/s. In contrast,

in Experiment 1 the increase in syllabic rate resulted in a dramatic drop of comprehension

performance. This is in line with our expectations, as the nature of the word-order task is likely

to yield overall better performance than the classic intelligibility task. Additionally, our control

experiment rules out a potential confound by demonstrating that the high performance in

397 Experiment 2 is not due to simple guessing of the correct word order (see Results section and
398 Supplementary Table 7). Interestingly, however, in both experiments performance decreased
399 later than previously observed, that is, beyond rates of 9 syllables/s(56,80). However, in line
400 with our findings, several other studies, also observed shallower decreases in speech compre-
401 hension, with relatively high comprehension at higher syllable rates (~12 sylla-
402 bles/s)(3,56,81,82). We consider several possible explanations for these discrepancies. One
403 explanation for the different and higher speech-rate decline in comprehension performance is
404 that naturally produced fast speech (with matched degrees of compression across syllabic
405 rates, as used in Experiment 2), in contrast to linearly compressed speech, results in more
406 variance of the speech rate and thus allows for part of the sentences to be understood. How-
407 ever, this explanation does not account for Experiment 1, in which all stimuli were synthesized
408 at the same rate (varying in degrees of compression). Furthermore, the high performance level
409 might be related to different complexity between more naturalistic sentences, providing
410 stronger context information to compensate loss of information, as compared to the words(18),
411 digits(83), or simple sentences(55) used in previous work. Finally, it is notable that while some
412 studies conceptualized the syllabic rate based on the 'theta-syllable' (an information unit defi-
413 ned by cortical function(84)), we define syllabic rate as linguistically defined syllables per sec-
414 ond, following other studies(36).

415 Auditory-motor speech synchronization, a behavioral estimate of auditory-motor cortex cou-
416 pling strength(40), had a modulatory -albeit small- effect on speech comprehension. We ob-
417 served that high compared to low synchronizers exhibited better speech comprehension per-
418 formance. These results expand on findings which showed superior statistical word learn-
419 ing(40) or syllable discrimination(6) for individuals with stronger auditory-motor coupling by
420 showing a similar effect for comprehending more naturalistic, continuous speech. Note that
421 this effect requires further validation as it did not survive control for multiple comparisons (Sup-
422 plementary Table 5). Additionally, we expected an interaction of syllabic rate and auditory-
423 motor synchronization, as reported for rate discrimination in tone sequences(10). However,
424 the modulation observed here occurred across all syllabic rates, suggesting that an interaction
425 effect may be masked and compensated for by context and linguistic information in continuous
426 speech comprehension. Alternatively, it is possible -although unlikely- that the interaction of
427 syllabic rate and auditory-motor synchronization was not observed here due to the different
428 frequency resolution at low frequencies. The difference between HIGHs and LOWS in Kern et

al.(10) manifested between 7.14 and 10.29 Hz. In contrast, in the present experiment, there

was no frequency condition between 5 and 10.69 syllables/s.

Importantly, the spontaneous motor production rate affected speech comprehension, suggest-

ing that individuals with higher spontaneous motor production rate have increased speech

comprehension abilities (at the higher range). We replicated this finding in the second experi-

ment. The finding might reflect a complex interplay of auditory and motor cortex during speech

comprehension wherein not only the coupling strength, but also the preferred rates of the motor

cortex affect speech perception. A possible role of the preferred speech motor rate for speech

processing has been previously discussed(35). Furthermore, our findings are in line with an

oscillatory model of speech comprehension(6). An alternative interpretation of our findings

might be that general processes such as vigilance and fatigue are equally reflected in the

spontaneous speech motor production rate and the speech comprehension performance. This

could be, because speech comprehension naturally is tightly intertwined with production, and

vigilance effects on production for example might similarly affect comprehension. The behav-

ioral protocol does not allow to completely discard such an alternative interpretation. However,

given that no correlation of a demanding cognitive task (Digit span) with the spontaneous

speech motor production rates was observed (see Supplementary Material), we consider this

unlikely. Furthermore, for the effects of speech auditory-motor synchronization on syllable dis-

crimination others have ruled out such an interpretation(6).

Interestingly, the preferred auditory rate (~5.55 syllables/s) had no effect on speech compre-

hension in our study. A possible explanation is that preferred rates in auditory cortex are less

flexible compared to preferred rates in motor cortex and thus less prone to individual difference

related improvements of speech comprehension. However, comparing the variances of the

distribution of preferred auditory ($s^2 = 0.74$) and motor ($s^2 = 0.20$) rates revealed bigger vari-

ance in the auditory rate ($F(1,162) = 22.39$, $p < .001$). Another possibility is that the behavioral

estimation of preferred auditory cortex rates were not optimally operationalized. This might

also explain the lack of correlation between preferred auditory and spontaneous speech pro-

duction rates (see Supplementary Material), which we expected to be correlated. Generally,

our behavioral protocol only allows for an indirect assessment of preferred neural rates. Nev-

ertheless, behavioral measures have been regarded as proxy for underlying intrinsic brain

rhythms(45). Finally, the rates at which speech comprehension decreases are much higher

than the preferred auditory and spontaneous speech motor production rates. While the pre-

ferred rates were well within the expected range(7,8), the mismatch between maximal

comprehension rates and preferred rates was due to the high speech comprehension ability
of participants even at high rates.

We show that continuous speech comprehension is additionally affected by other higher cog-
nitive and linguistic factors. The relevance of linguistic predictability and working memory ca-
pacity have been shown in multiple studies(53,54). In agreement with these studies, such cog-
nitive variables explained a large amount of variance in speech comprehension. Interestingly,
our findings suggest that the faciliatory effect of linguistic predictability is particularly effective
at fast rates. Second, we tentatively interpret that facilitation due to linguistic predictability may
be used more efficiently from individuals with stronger auditory-motor synchronization. A rele-
vant question arising from this is: under what conditions is the impact of the motor system on
speech comprehension the strongest? Previous work observed an impact of the motor system
on speech comprehension in demanding listening conditions, such as listening to speech in
noise (5,33). Our data suggests that this view might extend towards conditions of fast speech
(which requires more experiments) or might interact with linguistic predictability.

Speech comprehension is a highly predictive process which is affected by different sources of
predictions. Here we show that, while speech comprehension is optimal in a preferred auditory
temporal regime, the motor-system provides a role for individual flexibility in continuous speech
comprehension. Additionally, we report that the well-known faciliatory effects of linguistic pre-
dictability on speech comprehension interact with individual differences in the motor system.
This sets the stage for future assessments of how predictions from these systems interact and
under what circumstances the human brain relies more on one over the other.

Ethics

Experiment 1 was approved by the ethics committee of the School of Social Sciences, Univer-

sity of Dundee, UK (No. UoD-SoSS-PSY-UG-2019-88). Procedures for Experiment 2 and the

Control Experiment were approved by the Max Planck Society (No. 2017_12).

Data accessibility

Preprocessed data and analysis scripts will be available via OSF upon publication.

Competing interests

We declare we have no competing interests.

Funding

This work was supported by the Max Planck Society.

Acknowledgments

We thank Anna Broggi and Harry Watt for help with data acquisition (Exp. 1), Dr. Klaus Frieler

for methodological advice, and Dr. Gregory Hickok for valuable comments on a previous ver-

sion of the manuscript.

References

1.   Rosen S. Temporal information in speech: acoustic, auditory and linguistic aspects.
Philosophical Transactions of the Royal Society of London Series B: Biological Sciences. 1992
Jun 29;336(1278):367–73.

2.   Nourski KV, Reale RA, Oya H, Kawasaki H, Kovach CK, Chen H, et al. Temporal En-
velope of Time-Compressed Speech Represented in the Human Auditory Cortex. Journal of
Neuroscience. 2009 Dec 9;29(49):15564–74.

3.   Brungart DS, van Wassenhove V, Brandewie E, Romigh G. The effects of temporal
acceleration and deceleration on AV speech perception. AVSP. 2007;27–34.

4.   Morillon B, Schroeder CE, Wyart V. Motor contributions to the temporal precision of
auditory attention. Nature Communications. 2014 Dec;5(1).

5.   Stokes RC, Venezia JH, Hickok G. The motor system's [modest] contribution to speech
perception. Psychon Bull Rev. 2019 Aug;26(4):1354–66.

6.   Assaneo MF, Rimmele JM, Sanz Perl Y, Poeppel D. Speaking rhythmically can shape
hearing. Nat Hum Behav. 2021 Jan;5(1):71–82.

7.   Poeppel D, Assaneo MF. Speech rhythms and their neural foundations. Nat Rev Neu-
rosci. 2020 May 6;

8.   Ding N, Patel AD, Chen L, Butler H, Luo C, Poeppel D. Temporal modulations in speech
and music. Neuroscience & Biobehavioral Reviews. 2017 Oct;81:181–7.

9.   Pellegrino F, Coupé C, Marsico E. Across-Language Perspective on Speech Infor-
mation Rate. Language. 2011;87(3):539–58.

10.   Kern P, Assaneo MF, Endres D, Poeppel D, Rimmele JM. Preferred auditory temporal
processing regimes and auditory-motor synchronization. Psychon Bull Rev. 2021 Jun 7;

11.   Teng X, Tian X, Rowland J, Poeppel D. Concurrent temporal channels for auditory pro-
cessing: Oscillatory neural entrainment reveals segregation of function at different scales. Jen-
sen O, editor. PLOS Biology. 2017 Nov 2;15(11):e2000812.

12.   Drake C, Botte MC. Tempo sensitivity in auditory sequences: Evidence for a multiple-
look model. Perception & Psychophysics. 1993 May;54(3):277–86.

13.   Teng X, Poeppel D. Theta and Gamma Bands Encode Acoustic Dynamics over Wide-
Ranging Timescales. Cerebral Cortex. 2020 Apr 14;30(4):2600–14.

14.   Viemeister NF. Temporal modulation transfer functions based upon modulation thresh-
olds. The Journal of the Acoustical Society of America. 1979 Nov;66(5):1364–80.

15.   Giraud AL, Poeppel D. Cortical oscillations and speech processing: emerging compu-
tational principles and operations. Nat Neurosci. 2012 Apr;15(4):511–7.

16.   Gross J, Hoogenboom N, Thut G, Schyns P, Panzeri S, Belin P, et al. Speech rhythms
and multiplexed oscillatory sensory coding in the human brain. Poeppel D, editor. PLoS Biol.
2013 Dec;11(12):e1001752.

17.   Peelle JE, Davis MH. Neural Oscillations Carry Speech Rhythm through to Compre-
hension. Frontiers in Psychology. 2012;3.

18.   Ghitza O, Greenberg S. On the possible role of brain rhythms in speech perception:
Intelligibility of time-compressed speech with periodic and aperiodic insertions of silence.

Phonetica. 2009;66(1–2):113–26.

19.   Giraud AL, Kleinschmidt A, Poeppel D, Lund TE, Frackowiak RSJ, Laufs H. Endoge-
nous cortical rhythms determine cerebral specialization for speech perception and production.
Neuron. 2007 Dec;56(6):1127–34.

20.   Luo H, Poeppel D. Phase patterns of neuronal responses reliably discriminate speech
in human auditory cortex. Neuron. 2007 Jun;54(6):1001–10.

21.   Greenberg S, Carvey H, Hitchcock L, Chang S. Temporal properties of spontaneous
speech—a syllable-centric perspective. Journal of Phonetics. 2003 Jul;31(3–4):465–85.

22.   Morillon B, Baillet S. Motor origin of temporal predictions in auditory attention. Proceed-
ings of the National Academy of Sciences. 2017 Oct 17;114(42):E8913–21.

23.   Schubotz RI. Prediction of external events with our motor system: towards a new frame-
work. Trends in Cognitive Sciences. 2007 May;11(5):211–8.

24.   Grahn JA, Brett M. Rhythm and Beat Perception in Motor Areas of the Brain. Journal
of Cognitive Neuroscience. 2007 May 1;19(5):893–906.

25.   Cason N, Astésano C, Schön D. Bridging music and speech rhythm: Rhythmic priming
and audio–motor training affect speech perception. Acta Psychologica. 2015 Feb;155:43–50.

26.   Du Y, Zatorre RJ. Musical training sharpens and bonds ears and tongue to hear speech
better. Proc Natl Acad Sci USA. 2017 Dec 19;114(51):13579–84.

27.   Grahn JA, Rowe JB. Feeling the Beat: Premotor and Striatal Interactions in Musicians
and   Nonmusicians   during   Beat   Perception.   Journal   of   Neuroscience.   2009   Jun
10;29(23):7540–8.

28.   Cheung C, Hamilton LS, Johnson K, Chang EF. The auditory representation of speech
sounds in human motor cortex. eLife. 2016 Mar 4;5:e12577.

29.   Evans S, Davis MH. Hierarchical Organization of Auditory and Motor Representations
in Speech Perception: Evidence from Searchlight Similarity Analysis. Cereb Cortex. 2015
Dec;25(12):4772–88.

30.   Hickok G, Poeppel D. The cortical organization of speech processing. Nat Rev Neuro-
sci. 2007 May;8(5):393–402.

31.   Morillon B, Arnal LH, Schroeder CE, Keitel A. Prominence of delta oscillatory rhythms
in the motor cortex and their relevance for auditory and speech perception. Neuroscience &
Biobehavioral Reviews. 2019 Dec;107:136–42.

32.   Scott SK, McGettigan C, Eisner F. A little more conversation, a little less action — can-
didate roles for the motor cortex in speech perception. Nat Rev Neurosci. 2009 Apr;10(4):295–
302.

33.   Wu ZM, Chen ML, Wu XH, Li L. Interaction between auditory and motor systems in
speech perception. Neurosci Bull. 2014 Jun;30(3):490–6.

34.   Rogalsky C, Basilakos A, Rorden C, Pillay S, LaCroix AN, Keator L, et al. The Neuro-
anatomy of Speech Processing: A Large-scale Lesion Study. Journal of Cognitive Neurosci-
ence. 2022 Jun 10;1–21.

35.   Assaneo MF, Poeppel D. The coupling between auditory and motor cortices is rate-
restricted: Evidence for an intrinsic speech-motor rhythm. Sci Adv. 2018 Feb;4(2):eaao3842.

36.   Keitel A, Gross J, Kayser C. Perceptually relevant speech tracking in auditory and mo-
tor cortex reflects distinct linguistic features. Bizley J, editor. PLOS Biology. 2018
Mar;16(3):e2004473.

37.   Keitel A, Gross J. Individual human brain areas can be identified from their character-
istic spectral activation fingerprints. Engel AK, editor. PLOS Biology. 2016
Jun;14(6):e1002498.

38.   Lubinus C, Orpella J, Keitel A, Gudi-Mindermann H, Engel AK, Roeder B, et al. Data-
Driven Classification of Spectral Profiles Reveals Brain Region-Specific Plasticity in Blindness.
Cerebral Cortex. 2021 Mar 31;31(5):2505–22.

39.   Rosanova M, Casali A, Bellina V, Resta F, Mariotti M, Massimini M. Natural Frequen-
cies of Human Corticothalamic Circuits. Journal of Neuroscience. 2009 Jun 17;29(24):7679–
85.

40.   Assaneo MF, Ripollés P, Orpella J, Lin WM, de Diego-Balaguer R, Poeppel D. Spon-
taneous synchronization to speech reveals neural mechanisms facilitating language learning.
Nat Neurosci. 2019 Apr;22(4):627–32.

41.   Park H, Ince RAA, Schyns PG, Thut G, Gross J. Frontal Top-Down Signals Increase
Coupling of Auditory Low-Frequency Oscillations to Continuous Speech in Human Listeners.
Current Biology. 2015 Jun;25(12):1649–53.

42.   Haegens S, Zion Golumbic E. Rhythmic facilitation of sensory processing: A critical
review. Neuroscience & Biobehavioral Reviews. 2018 Mar;86:150–65.

43.   Rimmele JM, Morillon B, Poeppel D, Arnal LH. Proactive sensing of periodic and ape-
riodic auditory patterns. Trends in Cognitive Sciences. 2018 Oct;22(10):870–82.

44.   Assaneo MF, Rimmele JM, Orpella J, Ripollés P, de Diego-Balaguer R, Poeppel D.
The lateralization of speech-brain coupling Is differentially modulated by intrinsic auditory and
top-down mechanisms. Front Integr Neurosci. 2019 Jul 17;13:28.

45.   McPherson T, Berger D, Alagapan S, Fröhlich F. Intrinsic Rhythmicity Predicts Syn-
chronization-Continuation Entrainment Performance. Scientific Reports. 2018 Dec;8(1).

46.   Rimmele JM, Kern P, Lubinus C, Frieler K, Poeppel D, Assaneo MF. Musical Sophisti-
cation and Speech Auditory-Motor Coupling: Easy Tests for Quick Answers. Front Neurosci.
2022 Jan 4;15:764342.

47.   Emmorey K, Giezen MR, Petrich JAF, Spurgeon E, O'Grady Farnady L. The relation
between working memory and language comprehension in signers and speakers. Acta Psy-
chologica. 2017 Jun;177:69–77.

48.   Arnal LH, Wyart V, Giraud AL. Transitions in neural oscillations reflect prediction errors
generated in audiovisual speech. Nat Neurosci. 2011 Jun;14(6):797–801.

49.   Zhang Y, Frassinelli D, Tuomainen J, Skipper JI, Vigliocco G. More than words: word
predictability, prosody, gesture and mouth movements in natural language comprehension.
Proc R Soc B. 2021 Jul 28;288(1955):20210500.

50.   Lewis AG, Bastiaansen M. A predictive coding framework for rapid neural dynamics
during sentence-level language comprehension. Cortex. 2015 Jul;68:155–68.

51.   Kuperberg GR, Jaeger TF. What do we mean by prediction in language comprehen-
sion? Language, Cognition and Neuroscience. 2016 Jan 2;31(1):32–59.

52.   Grant KW, Seitz PF. The recognition of isolated words and words in sentences: Indi-
vidual variability in the use of sentence context. The Journal of the Acoustical Society of Amer-
ica. 107.(2):1000–11.

53.   Obleser J, Wise RJS, Alex Dresner M, Scott SK. Functional Integration across Brain
Regions Improves Speech Perception under Adverse Listening Conditions. Journal of Neuro-
science. 2007 Feb 28;27(9):2283–9.

54.   Obleser J, Kotz SA. Expectancy Constraints in Degraded Speech Modulate the Lan-
guage Comprehension Network. Cerebral Cortex. 2010 Mar 1;20(3):633–40.

55.   Ahissar E, Nagarajan S, Ahissar M, Protopapas A, Mahncke H, Merzenich MM. Speech
comprehension is correlated with temporal response patterns recorded from auditory cortex.
Proceedings of the National Academy of Sciences. 2001 Nov;98(23):6.

56.   Dupoux E, Green K. Perceptual adjustment to highly compressed speech. Journal of
Experimental Psychology: Human Perception and Performance. 1997;23(3):914–27.

57.   Pefkou M, Arnal LH, Fontolan L, Giraud AL. θ-Band and β-Band Neural Activity Re-
flects Independent Syllable Tracking and Comprehension of Time-Compressed Speech. The
Journal of Neuroscience. 2017 Aug 16;37(33):7930–8.

58.   Beukelman DR, Yorkston KM. The relationship between information transfer and
speech intelligibility of dysarthric speakers. Journal of Communication Disorders. 1979
Jan;12(3):189–96.

59.   Schiavetti N, Sitler RW, Metz DE, Houde RA. Prediction of Contextual Speech Intelligi-
bility from Isolated Word Intelligibility Measures. J Speech Lang Hear Res. 1984
Dec;27(4):623–6.

60.   Schiavetti N. 1. Scaling procedures for the measurement of speech intelligibility. In:
Kent RD, editor. Studies in Speech Pathology and Clinical Linguistics. Amsterdam: John Ben-
jamins Publishing Company; 1992. p. 11.

61.   Lizcano-Cortés F, Gómez-Varela I, Mares C, Wallisch P, Orpella J, Poeppel D, et al.
Speech-to-Speech Synchronization protocol to classify human participants as high or low au-
ditory-motor synchronizers. STAR Protocols. 2022 Jun;3(2):101248.

62.   Richardson JTE. Measures of Short-Term Memory: A Historical Review. Cortex. 2007
Jan;43(5):635–50.

63.   Olsthoorn NM, Andringa S, Hulstijn JH. Visual and auditory digit-span performance in
native and non-native speakers. International Journal of Bilingualism. 2014 Dec;18(6):663–73.

64.   de Jong NH, Wempe T. Praat script to detect syllable nuclei and measure speech rate
automatically. Behavior Research Methods. 2009 May;41(2):385–90.

65.   Boersma P, Weenik D. Praat: doing phonetics by computer [Computer program]. 2020
Aug;Version 6.0.40(retrieved from http://www.praat.org/).

66.   Lachaux JP, Rodriguez E, Martinerie J, Varela FJ. Measuring phase synchrony in brain
signals. Human brain mapping. 1999;8(4):194–208.

67.   Smith ZM, Delgutte B, Oxenham AJ. Chimaeric sounds reveal dichotomies in auditory
perception. Nature. 2002 Mar;416(6876):87–90.

68.   MacQueen J. Some methods for classification and analysis of multivariate observa-
tions. Proceedings of the fifth Berkeley symposium on mathematical statistics and probability.

1967 Jun;1(14):281–97.

69.   Jurafsky D, Martin JH. Speech and language processing: an introduction to natural
language processing, computational linguistics, and speech recognition. 2. ed. [Nachdr.]. Up-
per Saddle River, NJ: Prentice Hall; 2009. 988 p.

70.   Chien JT, Ku YC. Bayesian Recurrent Neural Network for Language Modeling. IEEE
Trans Neural Netw Learning Syst. 2016 Feb;27(2):361–74.

71.   Mikolov T. Recurrent neural network based language model. Eleventh annual confer-
ence of the international speech communication association. 2010;24.

72.   Merity S, Keskar NS, Socher R. Regularizing and Optimizing LSTM Language Models
[Internet]. arXiv; 2017 [cited 2022 Oct 25]. Available from: http://arxiv.org/abs/1708.02182

73.   Fernandez J, Downey D. Sampling Informative Training Data for RNN Language Mod-
els. In: Proceedings of ACL 2018, Student Research Workshop [Internet]. Melbourne, Aus-
tralia: Association for Computational Linguistics; 2018 [cited 2022 Oct 25]. p. 9–13. Available
from: http://aclweb.org/anthology/P18-3002

74.   Baayen RH, Davidson DJ, Bates DM. Mixed-effects modeling with crossed random ef-
fects for subjects and items. Journal of Memory and Language. 2008 Nov;59(4):390–412.

75.   Barr DJ, Levy R, Scheepers C, Tily HJ. Random effects structure for confirmatory hy-
pothesis testing: Keep it maximal. Journal of Memory and Language. 2013 Apr;68(3):255–78.

76.   Sassenhagen J, Alday PM. A common misapplication of statistical inference: Nuisance
control with null-hypothesis significance tests. Brain and Language. 2016 Nov;162:42–5.

77.   Schielzeth H. Simple means to improve the interpretability of regression coefficients:
Interpretation of regression coefficients. Methods in Ecology and Evolution. 2010
Jun;1(2):103–13.

78.   Fox J, Weisberg S. An R companion to applied regression. 3rd ed. Thousand Oaks CA:
Sage Publications; 2019.

79.   Assaneo MF, Orpella J, Ripollés P, Noejovich L, López-Barroso D, de Diego-Balaguer
R, et al. Population-level differences in the neural substrates supporting Statistical Learning.
bioRxiv; 2020 Jul.

80.   Ghitza O. Behavioral evidence for the role of cortical theta oscillations in determining
auditory channel capacity for speech. Front Psychol. 2014 Jul 4;5.

81.   Giroud J, Lerousseau JP, Pellegrino F, Morillon B. The channel capacity of multilevel
linguistic features constrains speech comprehension [Internet]. bioRxiv; 2021 Dec. Available
from: http://biorxiv.org/lookup/doi/10.1101/2021.12.08.471750

82.   Verschueren E, Gillis M, Decruy L, Vanthornhout J, Francart T. Speech Understanding
Oppositely Affects Acoustic and Linguistic Neural Tracking in a Speech Rate Manipulation
Paradigm. J Neurosci. 2022 Sep 28;42(39):7442–53.

83.   Doelling KB, Arnal LH, Ghitza O, Poeppel D. Acoustic landmarks drive delta–theta os-
cillations to enable speech comprehension by facilitating perceptual parsing. NeuroImage.
2014 Jan;85:761–8.

84.   Ghitza O. The theta-syllable: a unit of speech information defined by cortical function.
Front Psychol. 2013;4.