Mmp14-controlled extracellular matrix homeostasis is required for circadian rhythm

Richa Garva[1][ψ], Ching-Yan Chloé Yeung[1][ψ], Adam Pickard[1][ψ], Yinhui Lu[1], Venkatesh Mallikarjun[1], Joe Swift[1], Susan H. Taylor[1], Jyoti Rai[2], David R. Eyre[2], Mayank Chaturvedi[3], Yoshifumi Itoh[3], Qing-Jun Meng[1], Cornelia Mauch[4], Paola Zigrino[4], and Karl E. Kadler[1]*

[1] Wellcome Centre for Cell-Matrix Research, Faculty of Biology, Medicine & Health, University of Manchester, Manchester Academic Health Science Centre, Manchester, M13 9PT UK

[2] Department of Orthopedics, University of Washington, Seattle, WA, USA

[3] Kennedy Institute of Rheumatology, University of Oxford, Oxford, UK

[4] Department of Dermatology and Venereology, University of Cologne, Faculty of Medicine and University Hospital Cologne, 50937 Cologne, Germany

C-YCY current address: Institute of Sports Medicine Copenhagen, Department of Orthopedic Surgery, Copenhagen University Hospital – Bispebjerg and Frederiksberg, Copenhagen, Denmark, and Center for Healthy Aging, Department of Clinical Medicine, University of Copenhagen, Denmark

[ψ] Equal first authors.

*Corresponding author: K.E.K. (email: karl.kadler@manchester.ac.uk)

Keywords: MT1-MMP; chronomatrix; circadian clock; tendon, actin, focal adhesions

SUMMARY

The circadian clock regulates extracellular matrix (ECM) homeostasis but whether altered ECM homeostasis regulates the clock was unknown. Thus, we generated mice with a Col1a2-Cre-ERT2 conditional deletion of matrix metalloproteinase-14 (*Mmp14*) to disrupt matrix homeostasis in postnatal mice. We show here that tamoxifen-treated Col1a2-Cre-ERT2::Mmp14[F/F] mice developed a severe tendon matrix phenotype with dorsiflexion of hind limbs, tissue thickening, and accumulation of hydroxylysine aldehyde (pyridinoline) crosslinked narrow-diameter collagen fibrils typically seen accumulating during embryonic development. Proteomic analysis of tamoxifen-treated postnatal mice showed ~16% of detected proteins lost (176 proteins) or reversed (19 proteins) time-dependent rhythmicity. Ontological analysis identified 'cytoskeleton' as an affected system. Subsequent studies in CRISPR-Cas9 *Mmp14* null cells showed decreased F-/G-actin ratio, decreased cell spreading, elevated levels of pyridinoline crosslink factor *Plod2,* nuclear translocation of transcription factor *Pitx2*, and loss of circadian rhythm. Thus, our study shows a bidirectional dependence of collagen-ECM homeostasis and the circadian clock.

INTRODUCTION

MT1-MMP (gene name *Mmp14*) is a transmembrane matrix metalloproteinase first identified for its ability to activate progelatinase A (proMMP2) and to enhance cellular invasion of reconstituted basement membrane [1]. Extensive studies have showed that it can cleave a wide range of cellular and extracellular substrates including protease inhibitors, chemokines, cytokines, cell receptors, proMMP13, fibronectin and collagens (see [2]). Therefore, disrupted *Mmp14* expression has a major disruptive effect on matrix homeostasis. In humans, *Mmp14* has been implicated in cancer cell invasion [3,4], which has been attributed to its ability to degrade extracellular matrix (ECM) macromolecules, especially type I collagen (collagen-I) [5-7]. Like other vertebrate collagenases, it cleaves fibrillar collagens at the characteristic ¾-¼ site between residues 775 and 776 in each collagen α-chain [8]. MT1-MMP is distinct from other MMPs in that its deletion in mice results in early postnatal death [9,10]. The mice fail to thrive and exhibit a wide range of pathologies including defective osteogenesis and angiogenesis [9-11], altered white adipose tissue [12] and tendon [13]

development, modulation of inflammatory responses [14], activation of TGFβ1 [15], and regulation of α5β1 integrin endocytosis [16].

We noted that the organ systems affected in the *Mmp14* null mouse overlap with organ systems that are affected in mice that lack a functional circadian clock. Circadian clocks are cell-autonomous time keeping mechanisms that optimize cellular activities in anticipation of varying time-of-day demands on the cell [17, 18]. Whilst the suprachiasmatic nucleus is the master circadian pacemaker, almost all tissues have self-sustaining circadian pacemakers that synchronize rhythmic tissue-specific gene expression in anticipation of environmental light and dark cycles [19, 20]. Of particular interest, bone formation [19], tendon homeostasis [21, 22], adipogenesis [23], inflammatory immune responses [24], and angiogenesis [25] are processes that are regulated by the circadian clock and are affected in *Mmp14*-deficient mice. A search of public databases showed that transcripts for *Mmp14* are under circadian clock control in mouse suprachiasmatic nucleus (SCN), lung, heart, and adipose tissue[26]. Time-series transcriptomic profiling of mouse tendon (ArrayExpress accession no. E-MTAB-7743 [21]) showed circadian expression of the *Mmp14* transcript. However, direct mechanistic support for the importance of *Mmp14* rhythmicity was lacking.

Here we show in mouse tendon that *Mmp14* (RNA) and MT1-MMP (protein) are under robust circadian clock control and that fibroblasts lacking MT1-MMP are arrhythmic. We generated a type I collagen-specific conditional *Mmp14* knockout (CKO) mouse (*Col1a2^CreERT2^::Mmp14^F/F^*). Loss of circadian rhythm in the mice resulted in a severe tendon phenotype manifested as tissue thickening and dysregulation of cytoskeletal and matrisome proteins. Embryonic tendon contains exclusively 50-nm diameter collagen fibrils attached to cells via fibripositors. At birth, the fibrils are released from cells and grow in diameter to generate a multimodal diameter distribution, which is a developmental process that requires *Mmp14* [13]. Adult tendon contains a small population of narrow collagen fibrils (D1, ~50 nm diameter) that are under circadian clock control [22]. We show here that D1 fibrils accumulate in *Mmp14* CKO tendons. We also showed that recombinant MT1-MMP was effective at cleaving soluble collagen but not fibrillar collagen; thus the accumulation of collagen fibrils in the absence of MT1-MMP was the result of elevated fibrillogenesis and not the lack of turnover of collagen fibrils. One of the major changes in *Mmp14* null tendon was upregulation of proteins whose expression is controlled by the transcription factor *Pitx2* as well as *Plod2, Plod3,* and *LOX* that render the collagen network insoluble. We subsequently showed that collagen in *Mmp14* null tendons is stabilized by hydroxylysine aldehyde (pyridinoline) crosslinks. Therefore, the complex phenotype and progressive tissue fibrosis in *Mmp14*-deficient mice is most likely the consequence of disruption of circadian rhythm, loss of cytoskeleton in fibroblasts, accumulation of proteins that would normally be substrates for MT1-MMP, and constitutive expression of the lysyl hydroxylase/lysyl oxidase crosslinking program to generate insoluble collagen fibrils.

RESULTS

***Mmp14* (RNA) and MT1-MMP (protein) show circadian rhythms in expression in tendon**

The expression of *Mmp14*, but not other MMPs, is 24-hour rhythmic in tendon [21]. The readouts from 3 independent microarray probe sets showed the highest expression of *Mmp14* at the night-day transition (**Figure 1a**). Time-series western blot analysis showed peak levels of MT1-MMP at CT3 (**Figure 1b**), 4 hours after the peak in transcript. Next, we showed that time-of-day dependent expression of *Mmp14* (in mouse Achilles and tail tendon) was abolished in Scx-Cre::Bmal1^F/F^ and CLOCKΔ19 mice (**Figure 1c-e**), which lack circadian rhythms in their tendons [22]. As such, the rhythmic expression of *Mmp14* depends on an intact local circadian clock in tendon tissues.

We isolated tail tendon fibroblasts from the PER2::Luc mouse (iTTF-PER2::Luc or abbreviated to 'iTTF'), which enabled us to monitor the circadian clock in cultured cells using bioluminescence analysis using previously described methods [22]. In initial experiments we performed siRNA knockdown of *Mmp14.* Q-PCR showed that

knockdown was highly effective at reducing the expression of *Mmp14* (**Extended data figure 1a**) and
bioluminescence showed that the circadian clock rhythm was dampened in si-Mmp14 treated cells
(**Extended data figure 1b**). In a separate experiment, we incubated iTTF cells with GM6001 (a broad
spectrum MMP inhibitor) (**Extended data figure 1c**), which was also highly effective at dampening circadian
rhythm in treated cells compared to untreated controls (**Extended data figure 1d**). Next, we performed
clustered regularly interspaced short palindromic repeats (CRISPR)-associated protein 9 (Cas9) mediated
knockout of *Mmp14* in PER2::Luc fibroblasts (**Figure 1f, g**). Western blot analysis using an anti-MT1-MMP
antibody showed absence of MT1-MMP in 7 different clones (**Figure 1h**). Clones 1 and 2 were unable to
sustain a circadian rhythm in response to dexamethasone (**Figure 1i**), forskolin or 50% serum (**Extended data
figure 1e, f**), compared to matched cells, and restoring Mmp14 levels by lentiviral transduction rescued this
defect enabling robust oscillations (**Figure 1j**).

Accumulation of narrow collagen fibrils in tendons of the *Mmp14* conditional knockout mouse

Because global knockout of *Mmp14* results in perinatal death, we generated the tamoxifen-inducible
collagen-I-specific *Mmp14* knockout mouse (*Col1a2^{CreERT2}::Mmp14$^{F/F}$*) [27] to study the consequences of
postnatal *Mmp14* KO on the circadian clock. Thus, we treated five-week-old *Col1a2^{CreERT2}::Mmp14$^{F/F}$* mice
with tamoxifen for 4 weeks and collected mice at 10-, 14-, 18- and 23-week of age (see schematic in **Figure
2a**). We noticed that tamoxifen-treated mice had severe dorsiflexion of the hind limbs and difficulties
walking (**Figure 2b**). Therefore, under Home Office guidance, only small numbers of mice were available for
study. Examination of the limbs showed thickening of tendon bundles (**Figure 2c**). Transmission EM analysis
showed a progressive increase in the abundance of D1 (~ 50 nm diameter) collagen fibrils during T0 to T4 in
mice treated with tamoxifen compared to wild-type littermates (**Figure 2d, e** and **Extended data figure 2**).
We also noted a gradual increase in irregular fibril profiles from T0 to T4 (**Figure 2f**). Serial block face-
scanning electron microscopy, using established methods [28], showed gross disruption of the collagen fibril
network and accumulation of narrow fibrils in CKO tendons compared to tendons of littermates not treated
with tamoxifen (**Figure 2g, Movie 1** and **Movie 2**).

Recombinant MT1-MMP more effective at cleaving soluble collagen-I than fibrillar collagen-I

The accumulation of D1 collagen fibrils in the CKO tendons could have been explained by the absence of fibril
turnover in the absence of MT1-MMP. To explore this possibility, we released collagen fibrils from 6-week-
old mouse Achilles tendon and examined the diameters after incubation with recombinant soluble Fc-fusion
MT1-MMP *in vitro* [29]. The fibril diameters were derived from mass per length measurements using scanning
transmission electron microscopy [30]. The results showed no differences between rMT1-MMP treated and
untreated samples (**Figure 3a**). We then incubated the suspension of fibrils and soluble collagen-I with rMT1-
MMP at 30 °C and 37 °C. Gel electrophoresis confirmed efficient cleavage of soluble collagen-I by rMT1-
MMP at 30 °C, which was inhibited by EDTA (**Figure 3b**, red box). Liquid chromatography with tandem mass
spectrometry (LC-MS/MS) analysis of the cleavage products showed the presence of a semi-tryptic peptide
from the ¾ α1(I) fragment corresponding to the expected ¾-¼ cleave site, which confirmed that rMT1-MMP
had cleaved soluble collagen-I molecules at the expected Gly775 and Ile776 site (**Figure 3c** and **d**). However,
the results obtained when the suspension of fibrils was incubated with rMT1-MMP were different from those
obtained with soluble collagen-I. Here, only a small proportion of the collagen was cleaved when the reaction
was carried out at 30 °C (**Figure 3b**, white box); as expected, the cleavage was inhibited by EDTA. When the
reaction was carried out at 37 °C, no cleavage of collagen α-chains was observed. We did notice a downward
shift in mobility of β-chains (crosslinked α-chains), which was not inhibited by EDTA. It is likely that this was
the result of non-MMP activity in the collagen fibril preparation (**Figure 3b**). We noted that rMT1-MMP
cleaved a small proportion of the collagen-I from the fibril suspension when the reaction was performed at
30 °C. The most likely explanation was partial depolymerization of the fibrils at this temperature, resulting

in release of collagen molecules that were then susceptible to cleavage by rMT1-MMP. These results showed
that collagen is largely protected from cleavage by rMT1-MMP when the collagen is assembled into fibrils.

Tendons of the *Col1a2*^{CreERT2}*:Mmp14*^{F/F} null mice lose time-dependent features

Next, using mice kept in 12-hr light/12-hr dark conditions, we performed proteomics analysis of tendon
harvested at two time points during 24 h (zeitgeber time, ZT) ZT3 (09:00-hrs; 3 hours into light) and ZT15
(21:00; 3 hours into dark) from CKO *Mmp14* mice (n = 3) and control littermates (n = 3) at T2 (14 weeks of
age; 5 weeks after tamoxifen induction) (see schematic, **Figure 4a**). T2 (5 weeks post tamoxifen treatment)
was chosen because there were sufficient numbers of healthy mice in the control and CKO groups (whilst
allowing other mice to age to T4 for the electron microscopy study) and T2 was the time where a more
prominent fibrotic phenotype was detected in skin and tendons and deletion of *Mmp14* was complete [27]. We
extracted proteins using methods previously described [22]. Analysis of samples isolated at ZT3 and ZT15
identified 1195 proteins with ≥ 2 peptides per protein of which 212 showed time-dependent abundance, as
shown by MetaCycle analysis (**Figure 4a**), with good reproducibility between biological replicates as indicated
by Pearson analysis across consecutive 24-hour periods (**Extended data figure 3a**). Comparing logged
ZT15/ZT3 fold changes for 1195 proteins between CKO and wild-type tendons showed that the absence of
Mmp14 disrupted time-dependent protein abundances (**Extended data figure 3b**). Of the time-dependent
proteins detected in WT tendons, 17 have a statistically significant (FDR adjusted p-value <0.05) difference
in expression between the MMP14KO_CT3 and MMP14KO_CT15 but remained time-dependent, 19 had
reversed time-dependence (**Figure 4b**), and 176 had lost time-dependent expression in CKO tissues (**Figure
4c**). Functional annotation enrichment of the proteomic data using DAVID [31] showed that the proteins that
had remained time dependent were predominantly involved in intermediate filament assembly, metabolism,
cell-matrix adhesion and microtubule transport (**Figure 4d**). Those that had flipped time dependence (i.e.
those normally expressed at night and were expressed in the day in KO cells, and vice versa) and those that
had lost time-dependent expression were associated with actin cytoskeleton remodeling (**Figure 4e**) and
extracellular matrix including fibronectin, members of the small proteoglycan family, and enzymes involved
in collagen crosslinking. (**Supplementary data 1**).

***CRISPR/Cas9 knockout of Mmp14* knocks out the circadian rhythm and disrupts F-actin**

In the next series of experiments, we used CRISPR-Cas9 to knockout *Mmp14* transcripts in fibroblasts and in
MC3T3 osteoblasts (which also express *Mmp14*) and performed cell adhesion assays. The first observation
was that *Mmp14* KO fibroblasts and osteoblasts were defective at spreading on fibronectin compared to
control (**Figure 5a-c**). Secondly, the number (but not the size) of focal adhesions per cell was reduced in
Mmp14 KD fibroblasts (**Figure 5d, e**). Thirdly, staining with phalloidin and immunofluorescence using an anti-
vinculin antibody showed absence of an F-actin cytoskeleton in the *Mmp14* KO cells (**Figure 5f**). Western
blot analysis showed that the CKO cells had a decreased F-/G-actin ratio (**Figure 5g**), which was consistent
with the ontological analysis performed on the proteomics datasets (**Figure 4e**).

Circadian actin dynamics have been shown to drive rhythmic fibroblast mobilization during wound healing
[32], therefore we incubated fibroblasts with cytochalasin B and measured the circadian rhythm by
bioluminescence. The results show that disruption of the cytoskeleton dampened, but did not eradicate, the
circadian rhythm in tendon fibroblasts (**Figure 6a**). The organization of focal adhesions is complex and
involves cytoskeletal proteins such as vinculin and protein tyrosine kinases such as FAK (see [33]). We showed
that shRNA knockdown of vinculin dampened but did not stop the circadian rhythm (**Figure 6b, Extended
data figure 1g-h**). Furthermore, treatment with the FAK inhibitor had a pronounced effect on the amplitude
of the bioluminescence signal in the iTTFs and was able to rescue the circadian rhythm in *Mmp14* knockout
cells (**Figure 6c-e**).

Elevated collagen crosslinking in the absence of *Mmp14*

There was clear discrepancy between the increased tendon thickening and increased collagen fibril number
in the CKO mice (see **Figure 2**) compared to the low levels of collagen-I detected in the time-dependent
proteomics of CKO tissues (**Supplementary data 1**). Therefore, we collected tendons from T2 wild-type and
CKO mice at ZT3 and ZT15 and performed hydroxyproline analysis on whole tendons. We also subjected
tendons to extraction in SL-DOC (sodium laurate-sodium deoxycholate) buffer to isolate soluble proteins
followed by hydroxyproline quantification. As shown in **Table 1**, analysis of whole tendons showed that CKO
samples contained a higher content of collagen (as measured by hydroxyproline) than WT samples, which
was consistent with the increased tendon thickening and increased numbers of collagen fibrils in the CKO
mice. However, the amount of hydroxyproline in the SL-DOC samples showed less collagen in the CKO
samples than in the WT controls. We then subjected CKO and control samples to cross-linking analyses. The
results showed ~ 8-fold increase in hydroxylysine aldehyde (pyridinoline) in the CKO tendons compared to
control tendons. Mass spectrometry ruled out significant amounts of collagen-III as the source of pyridinoline
in the CKO samples. Therefore, the discrepancy in the number of collagen-I peptides extracted into SL-DOC
buffer, and the marked accumulation of collagen fibrils in the CKO tendons, was the result of decreased
extractability caused by increased hydroxylysine aldehyde cross-linking of collagen-I [34] in the CKO tissues. In
further experiments we cultured WT and CRISPR-Cas9 *Mmp14* knockout fibroblasts and determined
hydroxyproline and hydroxylysine aldehyde crosslink content, as above. The results showed an increase in
collagen and hydroxylysine aldehyde crosslink content in *Mmp14* KO cultures (**Table 2**).

Dysregulation of the matrisome in the absence of *Mmp14*
Comparison of over-represented proteins in the CKO tendons showed a preponderance of proteins
characteristic of fibrous tissues (see **Supplementary Data 1** and selected plots in **Extended data figure 4**).
These included fibronectin (up 6-fold), tenascin C (up 4-fold), COL5A1 (up 10-fold), and TIMP2 (3-fold
increase). Of particular interest, enzymes that catalyze posttranslational modifications in collagen leading to
crosslink formation were also elevated, and included PLOD2 (up 13-fold), and LOXL1 (up 20-fold). Also,
decorin and fibromodulin were decreased (~ 2-fold) whilst lumican was increased (~ 2-fold). These small
proteoglycans are regulators of collagen fibrillogenesis [35] and C-telopeptide cross-linking of collagen in
tendon [36]. Inflammasome markers, including NOD-like receptor family proteins, were either not detected or
unchanged.

Elevated Pitx2 and PITX2 nuclear localization in *Mmp14* knockout cells
Analysis of gene ontology terms associated with differentially expressed proteins in the CKO tendons showed
that PLOD1, 2, 3 and TPT1 are associated with the transcription factor *Pitx2*. The increased pyridinoline
crosslinking in the CKO tendons and the over-representation of the lysyl hydroxylases motivated a study of
Pitx2, which has been reported to regulate PLOD2 [37, 38], as well as expression of collagen I and V [39]. Firstly,
RT-qPCR showed that levels of *Pitx2* mRNA are low in murine tail tendon at both ZT3 and ZT15 but were
markedly elevated in the CKO tendons 5 weeks post tamoxifen treatment (**Figure 7a)**. Likewise, PLOD2 levels
were low in WT tendon but markedly elevated in the CKO tendons (**Figure 7b**). To facilitate further studies,
we expressed PITX2-GFP in iTTFs. Over-expression of PITX2-GFP was confirmed by RT-qPCR (**Figure 7c**) and
resulted in a 2-fold increase in expression of *Plod2* (**Figure 7d**). PITX2-GFP localized to cell nuclei but also
occurred in the cell body (**Figure 7e,** left column). The translocation of PITX2 to the nucleus has been
reported to be influenced by mechanosensors and reactive oxygen species [40]. Co-expression of mCherry-
MT1-MMP and PITX2-GFP showed that MT1-MMP and PITX2 localize to distinct puncta in the cell body
(**Figure 7e,** second column). However, in CRISPR-Cas9 KO *Mmp14* cells, PITX2-GFP was predominantly in the
nucleus (**Figure 7e,** third column). In a rescue experiment, we over-expressed mCherry-MT1-MMP in PITX2-
GFP expressing cells in which *Mmp14* was either CRISPR-Cas9 knocked out or not (**Figure 7e**, right hand
column). The rescue *Mmp14* carried a silent mutation (AGG to AAA) in the PAM site which did not alter the

amino acid sequence of MT1-MMP. The results show that nuclear localization of PITX2-GFP is evident in cells
that have not been rescued by mCherry-MT1-MMP.

Over-expression of mCherry-MT1-MMP (**Figure 7f**) suppressed both *Pitx2* and *Plod2* expression (**Figure 7g,h**,
whilst also resulting in a marked upregulation of *Col1a1* (**Figure 7i**). These results were in-line with a direct
relationship between *Pitx2* and *Plod2* as previously reported and the inverse relationship between *Pitx2* and
Mmp14 in the CKO tendons, compared to control. The relationship between *Mmp14* and *Pitx2* in terms of
expression levels and cell localization prompted us to determine if Pitx2 levels could impact on circadian
rhythm. As shown in **Figure 8**, constitutive expression of PITX2 or PITX2-GFP increased the bioluminescence
amplitude in Per2::Luc iTTF cells.

DISCUSSION

In this study we showed that *Mmp14* is under robust circadian clock control and its absence, either by
conditional deletion in mice or CRISPR-Cas9 gene editing of cultured cells, causes major matrisome disruption
and loss of circadian rhythm. In mice, the consequence was a severe tendon phenotype marked by
dorsiflexion of hind limbs, tissue thickening and mistiming and altered levels of 195 proteins. Previous studies
showing that the matrix is under circadian clock control and that disruption of the circadian clock leads to
tissue fibrosis, and, the results shown here that disruption of the matrix (by deletion of *Mmp14*) results in
loss of time-dependent protein expression, identifies a bi-directional dependence of matrix homeostasis and
the circadian clock.

Circadian clocks are maintained at the molecular level by BMAL1, CLOCK, CRYs and PERs, which are core
elements of a transcriptional/translational negative feedback loop that controls the rhythmic expression of
genes in a cell-specific manner [41]. Circadian clock responsive genes contain one or more transcriptional
elements, RORE, DBPE and E-box [42]. Noteworthy, a CACGTT (E-box) sequence is located 1.6 kb upstream of
the translational start site of murine and human *Mmp14*. *Mmp14* gene expression has been shown to be
rhythmically expressed in murine lung, heart, kidney and adipose as well as the suprachiasmatic nucleus [43].
These studies showed that limiting the expression of *Mmp14* to a specific time of day is likely to be conserved
across species and is likely coincident with *Bmal1* expression. We showed that the peak of *Mmp14*
expression in murine tendon is at CT23, which is in good agreement with the peak of *Bmal1* expression [21].
Therefore, it is likely that *Mmp14* is a Bmal1 responsive gene. The peak of expression of MT1-MMP was
particularly narrow (~ 4 hours after the peak of *Mmp14*) (**Figure 1**), which is consistent with observations
that MT1-MMP has a fast turnover of 4.8 hours, which puts it in the top 1% of rapidly turned-over proteins
in the proteome [44, 45].

Loss of *Mmp14* and circadian rhythmicity in the CKO cells affected ~ 16% of soluble proteins including
intracellular and soluble ECM proteins (**Figure 4**). The actin cytoskeleton and F-/G-actin ratio was disrupted
in the absence of *Mmp14*, and cells were less able to spread on a fibronectin matrix *in vitro*. In the absence
of *Mmp14 in vivo*, circadian clock-controlled D1 fibrils accumulate (**Figure 2**) leading to major disruption of
the ECM. Of note, accumulation of D1 fibrils also occurs in clockless fibroblasts [22]. The loss of circadian clock
rhythmicity in the absence of *Mmp14* could partially be explained by the depolymerization of the
cytoskeleton. However, cytochalasin-D (which binds cofilin [46] and disrupts the F/G-actin dynamics under
normal clock control [32]) and removal of vinculin only dampened the rhythm (**Figure 6**). Therefore, we propose
that the altered matrisome containing large numbers of crosslinked collagen fibrils, decreased decorin,
decreased fibromodulin, and elevated lumican, contribute to the loss of circadian rhythm in *Mmp14* deficient
cells.

A previous study showed that levels of soluble collagen and the formation of D1 collagen fibrils in healthy
murine tendon are under circadian clock control [22]. The peak of soluble collagen occurs in the early part of
the day (light cycle), which was attributed to the circadian regulation of the secretory pathway and the

rhythmic synthesis and removal of a sacrificial pool of collagen to maintain tissue health. MT1-MMP is a potent collagenase and, as we show, is under strict circadian control. Therefore, if MT1-MMP had a role in degrading collagen fibrils, its expression might be expected to peak at the day/night transition when fibril numbers decrease. However, as shown in **Figure 1**, both *Mmp14* and MT1-MMP peak at the night/day transition, which coincides with when collagen fibrils are assembling. This suggests a role for MT1-MMP in regulating the availability of soluble collagen for fibril assembly. This conclusion is supported by the finding shown in **Figure 3** that MT1-MMP is more effective in cleaving soluble collagen than fibrillar collagen. In further support of this conclusion, mapping functional domains of collagen-I onto the proposed structure of a collagen-I-containing fibril suggests that the fibril surface partially obscures accessibility of MMPs to the 775-776 cleavage site [47].

Hydroxyproline analysis showed a higher content of collagen in CKO compared to WT samples, which was consistent with the observed accumulation of D1 collagen fibrils in CKO tendons. However, the amount of collagen in the SL-DOC protein extracts was lower in the CKO than in WT. The discrepancy was the result of elevated hydroxylysine aldehyde crosslinking in the CKO tendons that prevented collagen-I from being extracted into SL-DOC buffer. To seek an explanation for elevated crosslinking in the CKO samples, we noted in the LC-MS/MS data elevated PLOD2, PLOD3 and LOX enzymes. Furthermore, there was a 50-fold increase in mRNA levels of *Pitx2* in CKO samples compared to control (**Figure 7**). Previous studies had shown that PITX2 regulates PLOD gene expression [37, 38]. Therefore, a possible explanation for the increased crosslinking in CKO samples was abnormally high activity of PITX2 in the absence to *Mmp14*. Fluorescent tagging of PITX2 and MT1-MMP followed by fluorescence microscopy analysis showed that MT1-MMP regulates the intracellular localization of PITX2 protein; in the absence of MT1-MMP, PITX2 is constitutively nuclear located, which according to previous studies would upregulate PLOD2 and the pyridinoline crosslink program. The overall outcome was the accumulation of crosslinked collagen chains in insoluble collagen fibrils.

We noticed high peptide counts in the mass spectrometry data for fibronectin, MMP2 and stromelysin 1, 2 and 3 in CKO compared to WT littermates. MMP3 and 10 can cleave a range of structural matrix proteins including collagens, proteoglycans, fibronectin and laminin whilst MMP11 cleaves α1-antitrypsin inhibitor and weakly degrades matrix proteins including collagen-IV [48]. MMPs can be activated intracellularly or extracellularly by proteolytic removal of a pro-domain by serine proteinases (e.g. furin[49]). Therefore, whilst these MMPs were elevated in the absence of *Mmp14*, it is possible that they existed in an inactive form. However, removal of the prodomain is not the only mechanism of activation; proMMPs can be activated by conformational changes induced either chemically [50] or by post-translational modifications of the cysteine switch [51]. MMPs can also have intracellular substrates and can reenter the cell (e.g. MT1-MMP [52]), be inefficiently secreted, or have organelle localization signals (reviewed by [53]). Therefore, the changes in the intracellular proteome that we detected by LC-MS/MS might also be expected to affect intracellular processes. Whilst fibronectin and collagen levels were elevated in the absence of MT1-MMP, the levels of small proteoglycans such as decorin, fibromodulin keratocan, and lubricin decreased (**Supplementary data 1**). The notable exception was lumican, which was elevated ~2-fold in CKO tendons. Lumican is a collagen-binding proteoglycan [54] and was proposed to facilitate interchain collagen crosslinking by interacting with collagen and lysyl oxidase (reviewed by [55]). Lumican has a critical role in the organization of collagen fibrils in tendons and cornea and is profibrotic in heart [56]. Furthermore, lumican knockout leads to reduced collagen crosslinking [57]. Therefore, elevated levels of lumican in the absence of *Mmp14* might also contribute to altered matrix homeostasis and elevated collagen crosslinks in the CKO mice.

In conclusion, the inter-dependence between the circadian clock and matrix homeostasis, especially collagen homeostasis, is regulated by the transmembrane metalloproteinase *Mmp14*/MT1-MMP. Lack of collagen homeostasis is a hallmark of fibrosis, which is associated with 45% of all deaths including those caused by

cancer [58]. Thus, we propose that targeting the 'circadian clock-matrix homeostasis axis' may provide opportunities for the treatment of fibrosis.

Acknowledgements

The research was funded by Wellcome (110126/Z/15/Z and 203128/Z/16/Z) and the BBSRC (BB/T001984/1) to KEK. CM and PZ were supported by Deutsche Forschungsgemeinschaft through SFB 829 (Project-ID 73111208-SFB829 (B4)). VM was supported by a studentship from the Sir Richard Stapley Educational Trust. JS was funded by a Biotechnology and Biological Sciences Research Council (BBSRC) David Phillips Fellowship (BB/L024551/1). BC is supported by a Wellcome PhD studentship (210062/Z/17/Z). MC and YI was funded by Polish Ministry of Science & Higher Education (1291/MOB/IV/2015/0). Q-JM was supported by Medical Research Council project grants MR/T016744/1 and MR/P010709/1 and Versus Arthritis Senior Research Fellowship Award 20875. National Institutes of Health (NIH) Grant: (NIAMS) AR037318 to DRE. The authors would like to thank Ronen Schweitzer (Shriners Hospital for Children, Portland, OR, USA) for *Scx-Cre* mice, Jan Zamek, Alison Hallworth and Raymond Hodgkiss, and the University of Manchester Biological Support Facility for assistance in animal welfare and husbandry. The proteomics was performed at the Biological Mass Spectrometry Facility in the Faculty of Biology, Medicine and Health (University of Manchester) with the assistance of Stacey Warwood and Ronan O'Cualain, and electron microscopy was performed in the Electron Microscopy Facility, Faculty of Biology, Medicine and Health (University of Manchester).

Author contributions:

Conceived the project and provided mechanistic insight: AP, C-YCY, KEK, RG.

Interpretation of results regarding the circadian clock: Q-JM.

Wrote the manuscript: AP, C-YCY, KEK, RG.

Designed experiments and interpreted data: AP, C-YCY, DE, JR, MC, PZ, RG, ST.

Provided reagents: CM, PZ, YI.

Performed the electron microscopy: YL.

Performed bioinformatics analysis: JS, VM.

Competing Interests:

The authors declare no competing interests.

Table 1: Collagen content in wild type and CKO tendons determined by hydroxyproline assay.

	Control		CKO	
	ZT3	ZT15	ZT3	ZT15
Hydroxyproline content of whole tissue (%)	70.97 ± 4.27	62.30 ± 5.25	71.43 ± 1.74	70.40 ± 6.35
Hydroxyproline content of SL-DOC extract (%)	1.50 ± 0.10	0.98 ± 0.31*	1.39 ± 0.12	0.81 ± 0.05*
Hydroxylysine aldehyde (pyridinoline) content (residue/residue hydroxyproline)	0.009 ± 0.003	0.090 ± 0.033	0.095 ± 0.033**	0.095 ± 0.007**

Two-way ANOVA: *p=0.0146 for time difference (total collagen), **p=0.0014 for genotype difference.

Table 2: Comparison of hydroxyproline and hydroxylysine aldehyde-pyridinoline content in wild type and *Mmp14* KO fibroblasts.

Sample	lyophilized wt (in mg)	Total collagen	% collagen	HP res/collagen
WT1	2.09	1.609	77.0	0.0055
WT2	2.07	1.298	62.7	0.0141
WT3	2.18	1.597	73.2	0.0084
WT4	1.17	0.760	65.0	0.0168
WT5	1.13	0.590	52.2	0.0102
WT6	2.3	1.605	69.8	0.0108
			66.6	**0.0110**
KO1	1.75	1.256	71.8	0.0234
KO2	1.89	1.290	68.3	0.1249
KO3	2.19	1.626	74.3	0.1205
KO4	1.09	0.900	82.6	0.1046
KO5	1.06	0.715	67.5	0.1002
KO6	1.32	0.807	61.2	0.0805
			70.9	**0.0923**

 FIGURE LEGENDS

Figure 1. *Mmp14* expression and MT1-MMP translation in tendon are under circadian clock control.

A, Expression of *Mmp14* mRNA in mouse tail tendons analysed by microarray every 4 h during 48 h, relative
to the first time point. Three coloured lines indicate the results from 3 probe sets on the microarray chip,
n=2; CT, circadian time (free-running time, in constant darkness). **B**, Expression of MT1-MMP in mouse tail
tendons analysed by western blotting. Levels of GAPDH protein served as a loading control. **C**, Time-
dependent expression of *Mmp14*, measured at zeitgeber time 3 (ZT3; 3 hours into the light phase) and ZT15
in mouse Achilles and tail tendons of wild type mice and **D**, tendon-specific *Bmal1* knockout (*Scx-
Cre::Bmal1$^{lox/lox}$*) mice, n=3; bars show SEM; **p=0.0043 for WT Achilles tendon, **p=0.0093 for WT tail
tendon, unpaired t-tests. **E**, relative Mmp14 expression in wild-type and CLOCKΔ19 Achilles and tail tendons,
n=3; **F**, Sanger sequencing of *Mmp14* in immortalised tail tendon fibroblasts (iTTFs) derived from the
Per2::Luc mouse (iTTFs, upper) and a single cell clone with *Mmp14* CRISPR-Cas9 knockout. Sequences
proximal to the gRNA binding site are shown. **G**, Western blot detection in control Per2::luc iTTFs and single
cell clones of MMP14 CRISPR-Cas9 knockouts. **H**, bioluminescence recordings from Per2::Luc iTTFs (black)
and single cell clones of CRISPR-Cas9 induced knockout of *Mmp14* Per2::Luc fibroblasts (blue) after
synchronisation with dexamethasone. **I,** bioluminescence recordings from Per2::Luc iTTFs (grey) or *Mmp14*
KO fibroblasts (blue) that have been transduced with a lentivirus expressing *Mmp14* (carrying a silent
mutation in the CRISPR-Cas9 PAM site) from a CMV promoter (black and red). Restoration of *Mmp14* levels
in *Mmp14* KO fibroblasts reinstated dexamethasone induced circadian rhythms (n=2 independent repeats
are shown, error bar represents standard deviation, and represent data from n=4 independent experiments.
J). Quantification of Per2::Luc amplitudes assessed over 3 days post dexamethasone treatment (n=4).

**Figure 2. Accumulation of collagen fibrils in tendons from tamoxifen-induced conditional deletion of
Mmp14 of *Col1a2CreERT2:Mmp14*$^{F/F}$ mice.**

A, Schematic of *Mmp14* deletion in fibroblasts of *Col1a2CreERT2:Mmp14*$^{F/F}$ (*Mmp14* CKO) mice *in vivo* by
tamoxifen-chow feeding at 5 weeks old for 4 weeks. Tissue was collected at T0 (before tamoxifen induction),
T1, 2, 3 and 4 (1, 5, 9 and 14 weeks after tamoxifen induction). **B-C**, Photographs showing thickening of skin
of *Mmp14* CKO at site-matched locations of the limb. **C**, thickening of tendons in the tail compared to control
litter mates at T=4. **D**, Transmission electron microscopy analysis showed *Mmp14* CKO tendons accumulated
narrow diameter fibrils both in the matrix and inside cells (black arrows, inset). increase in ~50 nm ⊘ fibrils
in *Mmp14* CKO tendons. **E**, Fibril diameter distributions of fibrils measured in control and *Mmp14* CKO
tendons at T4 show an increase in narrow-diameter (~50 nm) fibrils in *Mmp14* CKO tendons. **F**, Circularity of
collagen fibril cross sections of *Mmp14* CKO tendons showed significant irregularity compared to Ctrl fibrils
at all time points after tamoxifen induction, and increasing irregularity over time (270-470 fibrils measured
per time point; ***p<0.0001 between Ctrl and CKO, two way ANOVA, Bonferroni post tests; ###p<0.0001
comparison within CKO samples, Kruskal-Wallis one way ANOVA and Dunn's Multiple Comparison tests). **G**,
Single images selected from step-through Movie 1 and Movie 2 generated from volume-EM analysis of WT
and CKO tendons. Scale bar, 1 μm.

Figure 3. Recombinant MT1-MMP cleaves soluble but not fibrillar collagen-I.

A, Fibril diameter distribution of collagen fibrils dispersed from mouse Achilles tendons before treatment
and 24 h after treatment with and without rhMT1-MMP at 30 °C. *Effective hydrated fibril diameter (>500
fibrils measured). **B** Coomassie blue-stained gel showing incubation of monomeric and fibrillar collagen-I
with recombinant human MT1-MMP at 30 °C and 37 °C, in the presence or absence of 50 mM EDTA. M,
monomeric; F, fibrillar collagen dispersed from Achilles tendon. rhMT1-MMP generated ¾α1(I) and ¾α2(I)
fragments from monomeric collagen I (red box) and from fibrillar collagen (white box). Asterisk shows the
β-crosslinked chains of collagen-I. A prominent faster migrating β-dimer is seen in the MT1-MMP treated
samples which occurs in the presence and absence of 50 mM EDTA. **C**, Mass spectrum of rhMT1-MMP

generated fragments from monomeric collagen I shows the mass to charge ratio of individual ions resulting from MS/MS fragmentation pattern between Gly$_{775}$ and Ile$_{776}$. **D,** Precise fragmentation pattern that contributes to the generation of b and y ions.

Figure 4. Proteomic analysis of tendons from conditionally-deleted *Mmp14* mice.

A, Experimental strategy and results summary of proteomics analysis of protein abundance in tendons of *Col1a2CreERT2:Mmp14*$^{F/F}$ (CKO) compared to control littermates at ZT3, when *Mmp14* levels are highest, and at 12 h apart (ZT15) (n=3). 1195 proteins were identified, 212 were identified as significantly time-dependent in control tendons (p<0.05), of which 17 remained time-dependent, 19 showed a reversed abundance pattern, and 176 were no longer time-dependent in *Mmp14* CKO tendons. **B**, Heat map representation of proteins that remained time-dependent in *Mmp14* CKO tendons, including proteins with abundance in anti-phase to wild type tendons. **C**, Heat map representation of the abundance of 176 proteins that are time-dependent in control tendons but are no longer time-dependent in *Mmp14* CKO tendons. **D**, Functional analysis of proteins, of which the abundance remains time-dependent in *Mmp14* CKO tendons. The top five significantly (p<0.05) enriched gene ontology terms in biological function are shown. Numbers in bars show p values. **E**, Functional analysis of proteins, of which the time-dependent abundance have been deregulated *Mmp14* CKO tendons. The top significantly (p<0.05) enriched gene ontology terms in biological function and molecular function are shown. Numbers in bars show p values.

Figure 5. Focal adhesion site formation requires *Mmp14*.

Representative images of control and *Mmp14* KO Per2::Luc iTTFs (**A**) and *Mmp14* KO MC3T3 cells (**B**) stained with phalloidin (magenta) and vinculin (green). Bars represent 10 μm. **C**, percentage of cell spreading on fibronectin after 10 min. Quantification from n=3 independent experiments, a minimum of 2000 cells per replicate were scored. **D**, Numbers of vinculin-positive focal adhesions were counted in control Per2::Luc iTTFs or CRISPR-Cas9 *Mmp14* KO cells, a minimum of 2000 cells were counted per replicate, n=4, the numbers of focal adhesions per cell are reported relative to control iTTFs. **E**) The average focal adhesion size was unaltered in *Mmp14* KO cells although large focal adhesions were less frequently observed. **F**, Confocal microscopy images of control and *Mmp14* KO iTTFs immunolabelled for vinculin and collagen and stained with phalloidin after 72 hrs in culture. Bar, 10 μm. **G**, representative western blot of filamentous actin and soluble globular actin separated through ultra-centrifugation. **H**, Quantification of the F/G actin ratio in *Mmp14* KO cells, n=3. * p<0.05, ** p<0.01, one-way ANOVA.

Figure 6. Cells with *Mmp14* knockdown have fewer actin filaments and reduced ability to spread.

A, Bioluminescence recordings from Per2::Luc iTTFs (black) treated with cytochalasin B for 1 hour prior to dexamethasone synchronisation. **B,** bioluminescence recordings from Per2::Luc iTTFs transduced with control (black) or shRNA targeting vinculin (shVinc, blue) after dexamethasone synchronisation. **C,** Immunofluorescent detection of vinculin and collagen along with actin fibres detected with phalloidin in *Mmp14* knockout cells following addition of 10 μM FAK Inhibitor, PF-573228. **D,** Bioluminescence recordings from Per2::Luc iTTFs without synchronisation (red), after dexamethasone treatment (blue) or after pre-treatment with 10 μM PF-573228 for 5 hours prior to dexamethasone treatment. **E,** FAK inhibition strengthened and prolonged dexamethasone induced circadian rhythms and was also observed in *Mmp14* KO Per2::luc iTTFs

Figure 7. Pitx2 Is elevated in *Mmp14* CKO tendons.

A, Real-time PCR detection of Pitx2 mRNA levels in WT and *Mmp14* CKO tail tendons 5 weeks post tamoxifen treatment, at ZT3 (09:00 hr) and ZT15 (21:00 hr). **B,** Real-time PCR of PLOD2 mRNA expression in WT and *Mmp14* CKO tail tendons (n=3 animals per time point) 5 weeks post tamoxifen treatment, at ZT3 (09:00 hr) and ZT15 (21:00 hr). **C,** Per2::Luc iTTFs stably transduced with control or Pitx2-GFP lentivirus, demonstrating enhanced expression of Pitx2 and a concomitant increase in Plod2 mRNA expression **D,** (n=3 and n=5

respectively). **E,** Live cell Airyscan imaging of Per2::Luc iTTFs and MMP14 CRISPR-Cas9 KO ITTFs stably transduced with Pitx2-GFP lentivirus. Cells were transiently transduced with mCherry tagged MMP14 carrying a silent mutation in the CRISPR-Cas9 PAM site. Pitx2-GFP was readily detected within the nucleus but also with punctate localisation within the cell body in control cell, these correlated with regions also positive for MT1-MMP. Nuclear Pitx2-GFP signals were stronger within *Mmp14* KO cells and also displayed reduced localisation within the cell body. Re-expression of *Mmp14* within knockout cells reduced nuclear Pitx-GFP signals. **F-I,** Real-time PCR demonstrating expression of *Mmp14* in Pitx2-GFP expressing iTTFs, which resulted in reduced Pitx2 and Plod2 mRNA expression, but enhanced Col1a1 mRNA (n=3).

Figure 8. Higher amplitude of circadian clock rhythm in Per2::Luc fibroblasts overexpressing Pitx2.
A, Bioluminescence recordings from Per2::Luc iTTFs and the same cells stably overexpressing Pitx2 and Pitx2-GFP after synchronisation with dexamethasone. **B,** bioluminescence recordings from Per2::Luc iTTFs (n=3) and Pitx2 overexpressing fibroblasts that have been transduced with a lentivirus expressing Pitx2.

Extended data figure 1: Depletion or inactivation of *Mmp14* weakens circadian entrainment.
A, left, Mmp14 mRNA expression detected by real-time PCR 72 hours after transfection of Per2::Luc iTTFs with scrambled (si-scram) or *Mmp14* targeting siRNA (siMmp14), (n=3 biological replicates), **B,** representative bioluminescence recordings from si-scram and si-MMP14 transfected Per2::Luc fibroblasts, after synchronisation with dexamethasone (n=3 biological replicates). **C,** Dose response curves showing the effects of GM6001 on the viability of Per2::luc iTTFs assessed by Alamar blue. **D,** representative bioluminescence recordings from DMSO or GM6001 treated Per2::Luc fibroblasts, with and without synchronisation with dexamethasone (n=3 biological replicates). **E,** bioluminescence recordings from Per2::Luc iTTFs (black) and CRISPR-Cas9 edited *Mmp14* Per2::Luc fibroblasts (red) after synchronisation with forskolin, **F,** bioluminescence recordings from Per2::Luc iTTFs (black) and CRISPR-Cas9 edited *Mmp14* Per2::Luc fibroblasts (red) after synchronisation with 50% serum shock. **G,** After flow cytometry sorting of Per2::Luc iTTFs transduced with retrovirus expressing scrambled or vinculin targeting shRNA and GFP. GFP positive cells were sorted and mRNA levels of vinculin were assessed by real-time PCR, vinculin mRNA levels relative to control iTTFs are shown (n=3 biological repeats). **H,** Representative images showing the morphology of Per2::Luc iTTFs transduced with scrambled and vinculin targeting shRNA, captured 24 hours after seeding cells onto plastic tissue culture dishes.

Extended data figure 2: *Mmp14*-deficient tendons accumulate D1 collagen fibrils.
Transmission electron microscopy of wildtype (A) and Mmp14 null (B) mouse tail tendons. Scale bar, 0.5 μm.

Extended data figure 3: Depletion or inactivation of *Mmp14* weakens circadian entrainment
A, Correlation matrix showing Pearson's R-squared values between datasets resulting from mass spectrometry analysis of tail tendon tissue from *Col1a2CreERT2:Mmp14*^{F/F} (CKO) and control (Ctrl) at ZT3 and ZT15 timepoints within the circadian cycle (n=3 biological repeats; see Figure 4). Greater similarity was observed between samples at different timepoints than between CKO vs. Ctrl. **B,** Plot showing fold-changes in protein abundance, correlating changes observed in *Mmp14* CKO tendon tissues between ZT15 vs. ZT3 time points, to changes observed in control tissue between corresponding time points. The data showed significant positive correlation, with a Pearson's R-squared value of 0.2421 ($p < 0.0001$). However, the difference between the ZT15 vs. ZT3 timepoints was compressed in the *Mmp14* cKO tissues, evidenced by linear fit to the data (shown in red, with gradient lower than y=x), and consistent with a dampened circadian response (n=3 biological repeats).

Extended data figure 4: Different protein expression in *Mmp14* CKO.
Plots of proteins that remained time dependent; ECM proteins (COL1A1, COL1A2, COL5A1, FN1), Metalloproteinases (MMP14, TIMP2), Lysyl oxidase (PLOD2, LOXL2), Proteoglycans (DCN, LUM, FMOD, TNC);

482 in tendons of Col1a2CreERT2:Mmp14^{F/F} (CKO) compared to control littermates at ZT3, when *Mmp14* levels
483 are highest, and at 12 hrs apart (ZT15) at T2 (14 weeks old, 5 weeks after tamoxifen chow feeding) (n=3).

MATERIALS AND METHODS

Mice

The care and use of all mice in this study was carried out in accordance with the UK Home Office regulations, UK Animals (Scientific Procedures) Act of 1986 under the Home Office licence (70/8858) and with German legal guidelines. Ethical approval was granted by the University of Manchester and by the Regierungspräsidium Köln, Germany (NRW authorization 50.203.2- K 37a, 20/05 and AZ2010.A342). The permission included the generation of CKO out animals. All mice were housed in 12-h light/12-h dark cycle (LD) and are indicated where zeitgeber time (ZT) is used, with the exception of wild type C57BL/6 mice housed in 12-h dark/12-h dark cycle (DD) that were used for circadian time (CT) time-series microarray and protein extraction as previously described [21]. To generate inducible CKO mice in which *Mmp14* is ablated in fibroblasts upon induction with tamoxifen, we crossed mice expressing CreER recombinase under the control of the *Col1a2* promoter (*Col1a2-CreERT2*; C57BL/6) [59] with C57BL/6 mice carrying the floxed exons (exons 2 to 4) of the *Mmp14* gene [60] to produce *Col1a2-CreERT2::Mmp14^{lox/lox}* CKO mice. To induce *Mmp14* ablation, mice were fed tamoxifen-formulated chow at 5-weeks old for 4 weeks. To generate mice in which *Bmal1* is ablated in tendon-lineage cells, we crossed mice expressing Cre recombinase under the control of the *Scleraxis* promoter (*Scx-Cre*; C57BL/6) [61] with C57BL/6 mice carrying loxP sites flanking the exon encoding the BMAL1 basic helix-loop-helix domain [62] bred on a background with *PER2::Luciferase* knock-in [18] to produce *Scx-Cre::Bmal1^{lox/lox}* KO mice. *Scx-Cre::Bmal1^{lox/lox}* KO mice were validated previously [22]. *Col1a1^{r/r}* mice were imported from Jackson Laboratory (*B6;129S4-Col1a1^{tm1Jae/J}*) [63]. Mice that showed signs of distress or had problems walking were humanely sacrificed using Home Office approved cervical dislocation.

RNA isolation and qPCR analysis

RNA was isolated from Achilles and tail tendons as described previously [64]. In brief, tissues were washed with ice-cold PBS 3 times and snap frozen in liquid nitrogen with 500 μL TRIzol Reagent (Thermo Fisher Scientific) using a Mikro Dismembrator S (B. Braun Biotech International) at 2000 Hz for 90 seconds two times. After, 500 μL TRIzol was added to the homogenized samples and RNA was isolated according to the manufacturer's protocol. DNase treatment of RNA was performed using RQ1 RNase-free DNase (Promega) according to manufacturer's protocol. RNA integrity was assessed by gel electrophoresis and RNA concentration was measured using a NanoDrop 2000 (Thermo Fisher Scientific). TaqMan Reverse Transcription Reagents (Thermo Fisher Scientific) were used for complementary DNA synthesis and SensiFAST SYBR kit reagents (Bioline) were used for qPCR. Primers sequences used were: *Mmp14* AGT GAC AGG CAA GGC TGA TT and GCC CAC CTT AGG GGT GTA AT, and *Gapdh* AAC TTT GGC ATT GTG GAA GG and ACA CAT TGG GGG TAG GAA CA. The $2^{-\Delta\Delta Ct}$ method [65] was used to analyse relative fold changes in gene expression compared to the first time point. Microarray data for *Mmp14* expression during 48 h was collected previously [21] and the data is available on ArrayExpress (accession number E-MTAB-7743).

Transmission electron microscopy and morphometric analyses

Tendons from mice were prepared for transmission electron microscopy (TEM) as described previously [28]. Fibril diameter, perimeter and area were measured using ImageJ software to calculate circularity and fibril volume fraction as described previously [28]. Fibril diameter distributions were plotted and fitted to 3-Gaussian distribution using Sigma Plot 12.0 software (Systat Software) as previously described [28].

Western blotting

Protein purification and western blot analyses were performed as described previously [22]. Wild type mice were kept in DD for 36 h to allow endogenous circadian rhythm of gene expression. Protein was purified

from tail tendons at 4-h intervals. Primary antibodies used were mouse mAb to MT1-MMP (1:500; clone LEM-2/15.8, Millipore) and mAb GAPDH (1:10000; clone Gapdh71.1; Merck).

Sample preparation for proteomic analysis

Tail tendons from wild type and *Mmp14* CKO mice were collected at ZT3 and ZT15. Tissues were homogenized with a bullet blender (with 1.6 mm steel beads; Next Advance) at maximum speed at 4 °C for 5 min in 200 μL of SL-DOC buffer (1.1% sodium laurate, 0.3% sodium deoxycholate, 0.5 mM dithiothreitol (DTT) in 25 mM ammonium bicarbonate), supplemented with protease and phosphatase inhibitor cocktails (Roche). Samples were incubated at 4 °C for 5 min, alkylated with 12 μL of 30 mM of iodoacetamide for 20 min at RT, followed by quenching with 12 μL of 30 mM DTT. Samples were centrifuged at maximum speed for 10 min. Supernatants were transferred into LoBind tubes (Eppendorf) and protein concentrations measured using a Millipore Direct Detect spectrometer. A total of 2 μg protein per sample was digested using trypsin beads in accordance with the manufacturer's protocol (SMART digestion kit, Thermo Scientific). Supernatants were extracted from the beads and 10% formic acid added to adjust to pH 3. Samples were cleaned using an organic phase extraction method: 400 μL ethylacetate (Sigma Aldrich) was added, the resulting solution thoroughly mixed, centrifuged, and the organic phase extracted and discarded; this process was then repeated. Samples were then desalted, in accordance with the manufacturer's protocol, using POROS R3 beads (Thermo Fisher) and lyophilized in a speed-vac centrifuge (Heto).

Liquid chromatography-coupled tandem mass spectrometry (LC-MS/MS)

Samples were dissolved in 15 μl of 5% acetonitrile with 0.1% formic acid. Digested samples were analyzed using an UltiMate® 3000 Rapid Separation liquid chromatography system (RSLC, Dionex Corporation) coupled to either Q Exactive HF (Thermo Fisher Scientific) mass spectrometer. Peptides were separated in a gradient of 95% A and 5% B to 7% B at 1 min, 18% B at 58 min, 27% B in 72 min and 60% B at 74 min at 300 nL/min using a 75 mm x 250 μm inner diameter 1.7 μM CSH C18, analytical column (Waters). Peptides were selected for fragmentation automatically by data dependent analysis.

Proteomics data processing

Spectra from multiple samples were aligned using Progenesis QI (Nonlinear Dynamics) Peak-picking sensitivity was set to 4/5 and all other parameters were left as defaults. Only peptides with charge between +1 to +4, with 2 or more isotopes were taken for further analysis. Filtered peptides were identified using Mascot (Matrix Science UK), by searching against the SwissProt and TREMBL mouse databases. The peptide database was modified to search for alkylated cysteine residues (monoisotopic mass change, 57.021 Da), oxidized methionine (15.995 Da), hydroxylation of asparagine, aspartic acid, proline or lysine (15.995 Da) and phosphorylation of serine, tyrosine, threonine, histidine or aspartate (79.966 Da). A maximum of 2 missed cleavages was allowed. Peptide detection intensities were exported from Progenesis QI as Excel (Microsoft) spread sheets for further processing using code written in-house in MATLAB with the bioinformatics toolbox (R2015a, The MathWorks, USA). Raw proteomic data was deposited to the ProteomeXchange Consortium via the PRIDE partner repository with the data set identifier PXD031692. Fold-change differences in the quantity of proteins detected in different samples were calculated by fitting a linear regression model to take into account inter-sample variation [66, 67]. Briefly, peptide intensities were logged and normalised by the median intensity; protein fold-changes were calculated using a mixed-effects linear regression model, considering random (peptides, biological replicates) and fixed (controlled variables) effects. Pathway enrichment analysis was performed using the DAVID bioinformatics resource [31]. Peptide identifications were filtered via Mascot scores so that only those with a Benjamini-Hochberg false discovery rate (FDR) $\geq$ 0.05 remained. FDR filtering was performed on combined phospho-enriched and non-enriched datasets as described in [68]. Raw ion intensities from peptides belonging to proteins with fewer than 2 unique peptides (by sequence) per protein in the dataset were excluded from quantification. Remaining intensities were logged and normalized by the median logged peptide intensity. Peptides assigned to different isoforms were

grouped into a single "protein group" by gene name. Only peptides observed in at least 2 samples were used in quantification. Missing values were assumed as missing due to low abundance, an assumption others have shown is justified [69]. Imputation was performed at the peptide level following normalization using a method similar to that employed by Perseus [70] whereby missing values were imputed randomly from a normal distribution centered on the apparent limit of detection for this experiment. The limit of detection in this instance was determined by taking the mean of all minimum logged peptide intensities and down-shifting it by 1.6σ, where σ is the standard deviation of minimum logged peptide intensities. The width of this normal distribution was set to 0.3σ as described in [70]. Fold-change differences in the quantity of proteins detected in different time-points were calculated by fitting a mixed-effects linear regression model for each protein with Huber weighting of residuals (doi: 10.1214/aoms/1177703732) as described in [69] using the fitglme Matlab (The MathWorks, USA) function with the formula:

$$y_{iptb} = \beta_t + \beta_p + \beta_b$$

Where y_{ipt} represents the $\log_2$ (intensity) of peptide p belonging to protein i, at time-point t, in batch b. βs represent effect sizes for the indicated coefficients. Peptide was entered as a random effect whereas time-point was entered as a fixed effect. Standard error estimates were adjusted with Empirical Bayes variance correction according to [71, 72]. Conditions were compared with Student's t-tests with Benjamini-Hochberg correction for false positives (doi:10.1111/j.2517-6161.1995.tb02031.x).

Bioluminescence imaging and recordings

Bioluminescence was recorded as described previously [22]. Briefly, tendon fibroblasts were cultured on 35 mm culture dish in recording media containing 0.1 mM luciferin (Merck) in the presence of 100 nM dexamethasone (Merck) and real-time quantitative bioluminescence was recorded using lumiCycle apparatus (Actimetrics). Baseline subtraction was carried out using a 24-hr moving average. Amplitude was calculated as peak-trough difference in bioluminescence of the second peak using base-line subtracted data.

Fusion protein constructs

Lentivirus expression of *Mmp14* was achieved using vectors generated by Vectorbuilder. Murine *Mmp14* coding sequences were expressed under a CMV promoter, with mutation of the *Mmp14* targeting gRNA, AGATCAAGGCCAATGTTCGGAG, PAM site (AGG to AAA). For mCherry tagging the mCherry coding sequence was introduced after amino acid 536 of murine *Mmp14* following the design of Dr. Philippe Chavrier [73], these vectors also carried blasticidin resistance gene. Pitx2GFP expressing cells were cloned into pLenti PGK Neo DEST (Addgene plasmid # 19067 ref[74]), EGFP was cloned at the C terminus of the coding sequence of Pitx2 isoform 1 (Origene; MR227617). Lentivirus particles were generated as previously described [75]. Stable cells were created by neomycin selection (250 μg/mL) or 5 μg/mL blasticidin.

Light microscopy

Live cell super resolution microscopy for **Figure 7** was performed using a Zeiss LSM 880 Axio Observer microscope with Airyscan detection system in SR mode using a Plan-Apochromat 63×/1.4 Oil DIC M27 objective (Carl Zeiss, Jena, Germany). Excitation was achieved using the 488 nm line from an argon laser and 594 nm HeNe laser and emitted light was collected through appropriate filters to eliminate any spill-over. Raw images were Airyscan processed using the default settings and analyzed using ZEN 2.1 software and the image brightness and contrast adjusted using BestFit.

CRISPR/Cas9

To generate *Mmp14* knockout fibroblasts, immortalized tail tendon fibroblasts from the Per2::Luc mouse, generated at previously described [76], were transduced with lentivirus particles created from vectors co-expressing Cas9 and gRNA (ABMgood 302361140595). *Mmp14* knockout were achieved using vectors encoding the gRNA AGATCAAGGCCAATGTTCGGAG. After puromycin selection (2.5 μg/mL) we isolated single cell clones by FACS and confirmed knockout of *Mmp14* by real-time PCR and western blotting.

Collagen content and crosslink analyses

The hydroxylysine aldehyde (pyridinoline) cross-link content of tendon samples was determined by HPLC after acid hydrolysis in 6 N HCl for 24 hrs at 108 °C as previously described [34]. Results were expressed in residues per mole of collagen based on the hydroxyproline content assayed on the same hydrolysate as a measure of tissue collagen content.

Statistical analysis

Data were evaluated using Two-tailed Student's t-test, two-way ANOVA or non-parametric, two-tailed Mann-Whitney test. Results were presented as mean ± SEM from at least three independent experiments. Differences were considered significant at the values of $*P < 0.05$, $**P < 0.01$ and $***P < 0.001$.

REFERENCES

1. Sato, H. *et al.* A matrix metalloproteinase expressed on the surface of invasive tumour cells. *Nature* **370**, 61-65 (1994).
2. Tam, E.M., Morrison, C.J., Wu, Y.I., Stack, M.S. & Overall, C.M. Membrane protease proteomics: Isotope-coded affinity tag MS identification of undescribed MT1-matrix metalloproteinase substrates. *Proc Natl Acad Sci U S A* **101**, 6917-6922 (2004).
3. Hotary, K.B. *et al.* Membrane type I matrix metalloproteinase usurps tumor growth control imposed by the three-dimensional extracellular matrix. *Cell* **114**, 33-45 (2003).
4. Sodek, K.L., Ringuette, M.J. & Brown, T.J. MT1-MMP is the critical determinant of matrix degradation and invasion by ovarian cancer cells. *Br J Cancer* **97**, 358-367 (2007).
5. Ellerbroek, S.M., Wu, Y.I., Overall, C.M. & Stack, M.S. Functional interplay between type I collagen and cell surface matrix metalloproteinase activity. *J Biol Chem* **276**, 24833-24842 (2001).
6. Holmbeck, K., Bianco, P., Yamada, S. & Birkedal-Hansen, H. MT1-MMP: a tethered collagenase. *J Cell Physiol* **200**, 11-19 (2004).
7. Itoh, Y. Membrane-type matrix metalloproteinases: Their functions and regulations. *Matrix Biol* **44-46**, 207-223 (2015).
8. Tam, E.M., Wu, Y.I., Butler, G.S., Stack, M.S. & Overall, C.M. Collagen binding properties of the membrane type-1 matrix metalloproteinase (MT1-MMP) hemopexin C domain. The ectodomain of the 44-kDa autocatalytic product of MT1-MMP inhibits cell invasion by disrupting native type I collagen cleavage. *J Biol Chem* **277**, 39005-39014 (2002).
9. Holmbeck, K. *et al.* MT1-MMP-deficient mice develop dwarfism, osteopenia, arthritis, and connective tissue disease due to inadequate collagen turnover. *Cell* **99**, 81-92 (1999).
10. Zhou, Z. *et al.* Impaired endochondral ossification and angiogenesis in mice deficient in membrane-type matrix metalloproteinase I. *Proc Natl Acad Sci U S A* **97**, 4052-4057 (2000).
11. Hiraoka, N., Allen, E., Apel, I.J., Gyetko, M.R. & Weiss, S.J. Matrix metalloproteinases regulate neovascularization by acting as pericellular fibrinolysins. *Cell* **95**, 365-377 (1998).
12. Chun, T.H. *et al.* A pericellular collagenase directs the 3-dimensional development of white adipose tissue. *Cell* **125**, 577-591 (2006).
13. Taylor, S.H. *et al.* Matrix metalloproteinase 14 is required for fibrous tissue expansion. *Elife* **4**, e09345 (2015).
14. Shimizu-Hirota, R. *et al.* MT1-MMP regulates the PI3Kdelta.Mi-2/NuRD-dependent control of macrophage immune function. *Genes Dev* **26**, 395-413 (2012).
15. Mu, D. *et al.* The integrin alpha(v)beta8 mediates epithelial homeostasis through MT1-MMP-dependent activation of TGF-beta1. *J Cell Biol* **157**, 493-507 (2002).
16. Shi, F. & Sottile, J. MT1-MMP regulates the turnover and endocytosis of extracellular matrix fibronectin. *J Cell Sci* **124**, 4039-4050 (2011).
17. Yagita, K., Tamanini, F., van Der Horst, G.T. & Okamura, H. Molecular mechanisms of the biological clock in cultured fibroblasts. *Science* **292**, 278-281 (2001).
18. Yoo, S.H. *et al.* PERIOD2::LUCIFERASE real-time reporting of circadian dynamics reveals persistent circadian oscillations in mouse peripheral tissues. *Proc Natl Acad Sci U S A* **101**, 5339-5346 (2004).

19.   Roenneberg, T. & Merrow, M. Circadian clocks - the fall and rise of physiology. *Nat Rev Mol Cell Biol*
6, 965-971 (2005).
20.   Cermakian, N. & Sassone-Corsi, P. Multilevel regulation of the circadian clock. *Nat Rev Mol Cell Biol*
1, 59-67 (2000).
21.   Yeung, C.Y. *et al.* Gremlin-2 is a BMP antagonist that is regulated by the circadian clock. *Sci Rep* **4**,
5183 (2014).
22.   Chang, J. *et al.* Circadian control of the secretory pathway maintains collagen homeostasis. *Nat Cell*
Biol **22**, 74-86 (2020).
23.   Aggarwal, A. *et al.* The Circadian Clock Regulates Adipogenesis by a Per3 Crosstalk Pathway to Klf15.
Cell Rep **21**, 2367-2375 (2017).
24.   Keller, M. *et al.* A circadian clock in macrophages controls inflammatory immune responses. *Proc*
Natl Acad Sci U S A **106**, 21407-21412 (2009).
25.   Jensen, L.D. *et al.* Opposing effects of circadian clock genes bmal1 and period2 in regulation of VEGF-
dependent angiogenesis in developing zebrafish. *Cell Rep* **2**, 231-241 (2012).
26.   Pizarro, A., Hayer, K., Lahens, N.F. & Hogenesch, J.B. CircaDB: a database of mammalian circadian
gene expression profiles. *Nucleic Acids Res* **41**, D1009-1013 (2013).
27.   Zigrino, P. *et al.* Fibroblast-Derived MMP-14 Regulates Collagen Homeostasis in Adult Skin. *J Invest*
Dermatol **136**, 1575-1583 (2016).
28.   Starborg, T. *et al.* Using transmission electron microscopy and 3View to determine collagen fibril size
and three-dimensional organization. *Nat Protoc* **8**, 1433-1448 (2013).
29.   Itoh, Y. *et al.* A common SNP risk variant MT1-MMP causative for Dupuytren's disease has a specific
defect in collagenolytic activity. *Matrix Biol* **97**, 20-39 (2021).
30.   Sherratt, M.J., Graham, H.K., Kielty, C.M. & Holmes, D.F. Mass-mapping of ECM macromolecules by
scanning transmission electron microscopy. *Methods Mol Biol* **522**, 151-161 (2009).
31.   Huang da, W., Sherman, B.T. & Lempicki, R.A. Systematic and integrative analysis of large gene lists
using DAVID bioinformatics resources. *Nat Protoc* **4**, 44-57 (2009).
32.   Hoyle, N.P. *et al.* Circadian actin dynamics drive rhythmic fibroblast mobilization during wound
healing. *Sci Transl Med* **9** (2017).
33.   Fife, C.M., McCarroll, J.A. & Kavallaris, M. Movers and shakers: cell cytoskeleton in cancer
metastasis. *Br J Pharmacol* **171**, 5507-5523 (2014).
34.   Hudson, D.M. *et al.* Age-related type I collagen modifications reveal tissue-defining differences
between ligament and tendon. *Matrix Biol Plus* **12**, 100070 (2021).
35.   Chakravarti, S. *et al.* Lumican regulates collagen fibril assembly: skin fragility and corneal opacity in
the absence of lumican. *J Cell Biol* **141**, 1277-1286 (1998).
36.   Kalamajski, S. *et al.* Increased C-telopeptide cross-linking of tendon type I collagen in fibromodulin-
deficient mice. *J Biol Chem* **289**, 18873-18879 (2014).
37.   Ghosh, P., Saha, S.K., Nandi, S.S., Bhattacharya, S. & Roy, S.S. Involvement of Pitx2, a homeodomain
transcription factor, in hypothyroidism associated reproductive disorders. *Cell Physiol Biochem* **20**,
887-898 (2007).
38.   Hjalt, T.A., Amendt, B.A. & Murray, J.C. PITX2 regulates procollagen lysyl hydroxylase (PLOD) gene
expression: implications for the pathology of Rieger syndrome. *J Cell Biol* **152**, 545-552 (2001).
39.   Hendee, K.E. *et al.* PITX2 deficiency and associated human disease: insights from the zebrafish
model. *Hum Mol Genet* **27**, 1675-1695 (2018).
40.   Tao, G. *et al.* Pitx2 promotes heart repair by activating the antioxidant response after cardiac injury.
Nature **534**, 119-123 (2016).
41.   Takahashi, J.S. Transcriptional architecture of the mammalian circadian clock. *Nat Rev Genet* **18**,
164-179 (2017).
42.   Yamamoto, T. *et al.* Transcriptional oscillation of canonical clock genes in mouse peripheral tissues.
BMC Mol Biol **5**, 18 (2004).
43.   Li, X. *et al.* CirGRDB: a database for the genome-wide deciphering circadian genes and regulators.
Nucleic Acids Res **46**, D64-D70 (2018).
44.   Schwanhausser, B. *et al.* Corrigendum: Global quantification of mammalian gene expression control.
Nature **495**, 126-127 (2013).

45.   Schwanhausser, B. *et al.* Global quantification of mammalian gene expression control. *Nature* **473**,
337-342 (2011).
46.   Shoji, K., Ohashi, K., Sampei, K., Oikawa, M. & Mizuno, K. Cytochalasin D acts as an inhibitor of the
actin-cofilin interaction. *Biochem Biophys Res Commun* **424**, 52-57 (2012).
47.   Orgel, J.P. *et al.* Collagen fibril surface displays a constellation of sites capable of promoting fibril
assembly, stability, and hemostasis. *Connect Tissue Res* **52**, 18-24 (2011).
48.   Motrescu, E.R. *et al.* Matrix metalloproteinase-11/stromelysin-3 exhibits collagenolytic function
against collagen VI under normal and malignant conditions. *Oncogene* **27**, 6347-6355 (2008).
49.   Pei, D. & Weiss, S.J. Furin-dependent intracellular activation of the human stromelysin-3 zymogen.
Nature **375**, 244-247 (1995).
50.   Nagase, H., Enghild, J.J., Suzuki, K. & Salvesen, G. Stepwise activation mechanisms of the precursor of
matrix metalloproteinase 3 (stromelysin) by proteinases and (4-aminophenyl)mercuric acetate.
Biochemistry **29**, 5783-5789 (1990).
51.   Okamoto, T. *et al.* Activation of matrix metalloproteinases by peroxynitrite-induced protein S-
glutathiolation via disulfide S-oxide formation. *J Biol Chem* **276**, 29596-29602 (2001).
52.   Galvez, B.G. *et al.* Caveolae are a novel pathway for membrane-type 1 matrix metalloproteinase
traffic in human endothelial cells. *Mol Biol Cell* **15**, 678-687 (2004).
53.   Bassiouni, W., Ali, M.A.M. & Schulz, R. Multifunctional intracellular matrix metalloproteinases:
implications in disease. *FEBS J* (2021).
54.   Kalamajski, S. & Oldberg, A. Homologous sequence in lumican and fibromodulin leucine-rich repeat
5-7 competes for collagen binding. *J Biol Chem* **284**, 534-539 (2009).
55.   Kalamajski, S. & Oldberg, A. The role of small leucine-rich proteoglycans in collagen fibrillogenesis.
Matrix Biol **29**, 248-253 (2010).
56.   Engebretsen, K.V. *et al.* Lumican is increased in experimental and clinical heart failure, and its
production by cardiac fibroblasts is induced by mechanical and proinflammatory stimuli. *FEBS J* **280**,
2382-2398 (2013).
57.   Mohammadzadeh, N. *et al.* The extracellular matrix proteoglycan lumican improves survival and
counteracts cardiac dilatation and failure in mice subjected to pressure overload. *Sci Rep* **9**, 9206
(2019).
58.   Wynn, T.A. Common and unique mechanisms regulate fibrosis in various fibroproliferative diseases. *J
Clin Invest* **117**, 524-529 (2007).
59.   Zheng, B., Zhang, Z., Black, C.M., de Crombrugghe, B. & Denton, C.P. Ligand-dependent genetic
recombination in fibroblasts : a potentially powerful technique for investigating gene function in
fibrosis. *Am J Pathol* **160**, 1609-1617 (2002).
60.   Zigrino, P. *et al.* Loss of epidermal MMP-14 expression interferes with angiogenesis but not with re-
epithelialization. *Eur J Cell Biol* **91**, 748-756 (2012).
61.   Blitz, E., Sharir, A., Akiyama, H. & Zelzer, E. Tendon-bone attachment unit is formed modularly by a
distinct pool of Scx- and Sox9-positive progenitors. *Development* **140**, 2680-2690 (2013).
62.   Storch, K.F. *et al.* Intrinsic circadian clock of the mammalian retina: importance for retinal processing
of visual information. *Cell* **130**, 730-741 (2007).
63.   Liu, X. *et al.* A targeted mutation at the known collagenase cleavage site in mouse type I collagen
impairs tissue remodeling. *J Cell Biol* **130**, 227-237 (1995).
64.   Yeung, C.Y. *et al.* Chick tendon fibroblast transcriptome and shape depend on whether the cell has
made its own collagen matrix. *Sci Rep* **5**, 13555 (2015).
65.   Livak, K.J. & Schmittgen, T.D. Analysis of relative gene expression data using real-time quantitative
PCR and the 2(-Delta Delta C(T)) Method. *Methods* **25**, 402-408 (2001).
66.   Mallikarjun, V., Richardson, S.M. & Swift, J. BayesENproteomics: Bayesian Elastic Nets for
Quantification of Peptidoforms in Complex Samples. *J Proteome Res* **19**, 2167-2184 (2020).
67.   Gilbert, H.T.J. *et al.* Nuclear decoupling is part of a rapid protein-level cellular response to high-
intensity mechanical loading. *Nat Commun* **10**, 4149 (2019).
68.   Bekker-Jensen, D.B. *et al.* An Optimized Shotgun Strategy for the Rapid Generation of
Comprehensive Human Proteomes. *Cell Syst* **4**, 587-599 e584 (2017).

69. Goeminne, L.J., Gevaert, K. & Clement, L. Peptide-level Robust Ridge Regression Improves
Estimation, Sensitivity, and Specificity in Data-dependent Quantitative Label-free Shotgun
Proteomics. *Mol Cell Proteomics* **15**, 657-668 (2016).
70. Tyanova, S., Temu, T. & Cox, J. The MaxQuant computational platform for mass spectrometry-based
shotgun proteomics. *Nat Protoc* **11**, 2301-2319 (2016).
71. Smyth, G.K. Linear models and empirical bayes methods for assessing differential expression in
microarray experiments. *Stat Appl Genet Mol Biol* **3**, Article3 (2004).
72. Kammers, K., Cole, R.N., Tiengwe, C. & Ruczinski, I. Detecting Significant Changes in Protein
Abundance. *EuPA Open Proteom* **7**, 11-19 (2015).
73. Rosse, C. *et al.* Control of MT1-MMP transport by atypical PKC during breast-cancer progression.
Proc Natl Acad Sci U S A **111**, E1872-1879 (2014).
74. Campeau, E. *et al.* A versatile viral system for expression and depletion of proteins in mammalian
cells. *PLoS One* **4**, e6529 (2009).
75. Calverley, B.C., Kadler, K.E. & Pickard, A. Dynamic High-Sensitivity Quantitation of Procollagen-I by
Endogenous CRISPR-Cas9 NanoLuciferase Tagging. *Cells* **9** (2020).
76. Pickard, A. *et al.* Preservation of circadian rhythms by the protein folding chaperone, BiP. *FASEB J* **33**,
7479-7489 (2019).
789

Figure 1

Figure 2

a ± Tamoxifen-chow for 4 weeks / Normal chow

Age (weeks) 0 5 9 10 14 18 23
T0 T1 T2 T3 T4

b Control / *Mmp14* cKO

c Control / *Mmp14* cKO

d T1 T2 T3 T4

Control

Mmp14 cKO

e Control / *Mmp14* cKO

Number / Fibril diameter (mm)

f Circularity

Ctrl cKO Ctrl cKO Ctrl cKO Ctrl KO WT KO
T=0 T=1 T=2 T=3 T=4

###
*** *** *** ***

g T4 control / T4 CKO

Figure 3

a

0 h untreated

24 h 30°C buffer control

24 h 30°C rhMMP14

Number

Fibril diameter* (nm)

b

Collagen-I			30°C					37°C		
Collagen-I	M	−	M	M	F	F	F	F	F
rMMP14	−	+	+	+	+	+	−	+	+
EDTA	−	−	−	+	−	+	−	−	+

α1(I)
α2(I)
¾ α1(I)
¾ α2(I)
rhMMP14

*

c

% of base peak

y3 y4 b5 b6 b7 y6 y7 b9 b10 y9 y10 y5 y8 b11 b12 b13 y12 y13 b14 y14 y15

m/z

d

G S P GADGP A G S P G T P G P Q G
Ox   Ox   Ox

y18 y17   y15 y14 y13 y12   y10 y9 y8 y7 y6 y5 y4 y3

b5 b6 b7 b8 b9   b11 b12 b13 b14   b16   b18

Figure 4

Figure 5

Figure 6

Figure 7

Figure 8

a

b

Extended data figure 1

Extended data figure 2

Extended data figure 3

a

b

Extended data figure 4

Unprocessed blots for Figure 1

WT mouse tail tendons

Control lysates:
*Scx-Cre::Mmp14*fl/fl

b, bone
m, muscle
t, tendon

CT 3 7 11 15 19 23   b   m   t

WB: MT1-MMP

WB: GAPDH

WB: Calnexin