Western spotted skunks provide important food web linkages in forest of the Pacific Northwest

Running head: Western spotted skunk diet

Authors:

Marie I. Tosa[1], Damon B. Lesmeister[1,2], Jennifer M. Allen[1], Taal Levi[1]

[1]Department of Fisheries, Wildlife, and Conservation Sciences, Oregon State University, OR,

USA 97331

[2]Pacific Northwest Research Station, U.S. Department of Agriculture Forest Service, Corvallis,

OR, USA 97331

Corresponding author: Marie Tosa, tosa.marie@gmail.com

Abstract

There are increasing concerns about the decreasing population trends of small mammalian carnivores around the world. With limited knowledge about their ecology and natural history, small mammal conservation and management remains difficult. To address one of these deficiencies for western spotted skunks (*Spilogale gracilis*), we investigated their diet in the Oregon Cascades of the Pacific Northwest during 2017 – 2019. We collected 130 spotted skunk scats opportunistically and with detection dog teams and identified prey items using DNA metabarcoding and mechanical sorting. Western spotted skunk diet consisted of invertebrates such as wasps, millipedes, and gastropods, vertebrates such as small mammals, amphibians, and birds, and plants such as Gaultheria, Rubus, and Vaccinium. Diet also consisted of items such as black-tailed deer that were likely scavenged. Comparison in diet by season revealed that spotted skunks consumed more insects during the dry season (June - August) and marginally more mammals during the wet season (September – May). We observed similar diet in areas with no record of human disturbance and areas with a history of logging. Western spotted skunks provide important food web linkages between aquatic, terrestrial, and arboreal systems by facilitating energy and nutrient transfer, and serve important functional roles of seed dispersal and scavenging. Through prey-switching, western spotted skunks may dampen the effects of irruptions of prey, such as wasps during dry springs and summers, which could then provide ecosystem resilience to environmental change.

Keywords:

diet, DNA metabarcoding, generalist predator, small carnivore, land-use change, *Spilogale gracilis*

Introduction

Globally, many small mammalian carnivores (< 16 kg) face decreasing population trends due to multiple threats including land use change, disease, and overhunting (Belant, Schipper, and Conroy 2009; Marneweck et al. 2021). Even small carnivore species that were previously widely distributed and considered "least concern" by the IUCN, such as weasels (*Neogale* spp.), have shown signs of significant population decline (Jachowski et al. 2021). These declines are problematic because small carnivores can play important roles in ecosystem function, predator-prey dynamics, and disease transmission dynamics (Roemer, Gompper, and Van Valkenburgh 2009). Despite their potentially important roles, many small carnivores remain neglected in ecological research, data deficient, or understudied (Marneweck et al. 2021; Proulx 2010). Limited knowledge about small carnivore ecology and natural history continues to hinder management and conservation of declining populations.

In the Pacific Northwest, the western spotted skunk (*Spilogale gracilis*) is putatively a common forest carnivore, but little is known about their ecology due to their nocturnal nature (Verts, Carraway, and Kinlaw 2001) (Neiswenter and Dowler 2007; Doty and Dowler 2006; Neiswenter, Dowler, and Young 2010). Most spotted skunk literature is derived from the island spotted skunk subspecies (*S. g. amphialus*) (K. Crooks 1994b; 1994a), the congeneric eastern spotted skunk (*S. putorius*) (Lesmeister, Gompper, and Millspaugh 2009; Lesmeister et al. 2010; Kinlaw 1995) and plains spotted skunk subspecies (*S. p. interrupta*) (Crabb 1948), or other skunk species such as the pygmy skunk (*S. pygmaea*) (Cantú-Salazar et al. 2005). These spotted skunks, however, inhabit markedly different ecosystems such as on islands, prairie, or desert where there is limited forest vegetation. The Pacific Northwest, in comparison, is a temperate rainforest system that is dominated by large coniferous trees, and the functional role of western

spotted skunks in this system is largely unknown. Due to their dietary plasticity, spotted skunks

could vary from omnivorous generalist (e.g., eastern and plains spotted skunk) (Baker and Baker

1975; Cheeseman, Tanis, and Finck 2021; Selko 1937; Crabb 1941), insectivorous specialist

(e.g., pygmy skunk) (Cantú-Salazar et al. 2005), or key carnivorous predator of small vertebrates

(e.g., island spotted skunk) (K. R. Crooks and Van Vuren 1995) in Pacific Northwest forests.

Eastern and plains spotted skunk populations are in severe decline and as a result, the

eastern spotted skunk is now listed as Vulnerable by the IUCN (Gompper and Jachowski 2016),

and the plains spotted skunk subspecies had been petitioned for listing under the US Endangered

Species Act (US Fish and Wildlife Service 2012). The mechanism for these declines are poorly

understood, but multiple mechanisms have been proposed, including land-use change, disease

outbreaks, or changes in predator communities (Gompper and Hackett 2005; Gompper 2017; D.

Blake Sasse 2021). Although western spotted skunks are still relatively common and considered

a species of Least Concern by the IUCN (Cuarón, Helgen, and Reid 2016), western spotted

skunks may be prone to future, rapid declines similar to those of eastern and plains spotted

skunks. Studying western spotted skunks provides an opportunity to understand the functional

roles of the species and amass basic ecological knowledge that may inform conservation and

land management decisions.

Land use change is a potent disturbance in the Pacific Northwest given that it is an

internationally important center of timber production (Simmons et al. 2016). Forest management

can influence the structure and composition of forests with unknown consequences on the

ecology of western spotted skunks. One way that land use change could cause declines in spotted

skunk population densities is by causing declines in prey populations. For example, potential

small mammal prey such as Trowbridge's shrews (*Sorex trowbridgii*), shrew moles

(*Neurotrichus gibbsii*), red tree voles (*Arborimus longicaudus*), and flying squirrels (*Glaucomys*

oregonensis), are less abundant in young forest stands than in mature and old-growth forest

stands (Gilbert and Allwine 1991; Carey 1995; 1989). Disturbances such as commercial thinning

can reduce small mammal prey density (Manning, Hagar, and McComb 2012). In contrast, forest

management can increase the abundance of flowers and fruits of understory plants through

increased light penetration onto the forest floor (Wender, Harrington, and Tappeiner 2004). Thus,

it remains important to investigate how western spotted skunk diets are impacted by forest

management.

Characterizing the diet of small carnivores, however, has been difficult. New techniques

such as detection dogs and DNA metabarcoding have improved our ability to find scat and

identify prey items, respectively. Previously, diets of spotted skunks were difficult to study

because scats were often deposited in rest sites (Selko 1937; Lesmeister, Gompper, and

Millspaugh 2008), not on trails, and because spotted skunks exhibit an omnivorous diet

consisting of insects, small vertebrates, and fruit (Baker and Baker 1975; K. R. Crooks and Van

Vuren 1995; Crabb 1941; Howell 1906). These scats were typically collected opportunistically,

mechanically sorted, and morphologically identified (Ewins et al. 1994; Sándor and Ionescu

2009), but these processes had biases related to digestion that may render prey items

unrecognizable (Symondson 2002; Galan, Pagès, and Cosson 2012) and misidentification of rare

species (Massey et al. 2021). This can be particularly problematic for small omnivorous

predators that consume a wide breadth of prey items including plants, animals, and invertebrates

because identifiers must have taxonomic expertise. Moreover, misidentification of carnivores

from scat morphology has been problematic and has potentially led to biased results (Morin et al.

2016). Newer genetic approaches such as DNA metabarcoding (Ji et al. 2013; Monterroso et al.

2019) can increase confidence in correctly identifying the carnivore (Morin et al. 2016), increase

the number of prey items that can be correctly identified (Massey et al. 2021), and increase

efficiency of identifying diet for a high volume of samples (Kartzinel et al. 2015), especially for

omnivorous species (De Barba et al. 2014).

Here we use DNA metabarcoding and mechanical sorting to provide the first

comprehensive analysis of western spotted skunk diet in the Pacific Northwest, and quantified

seasonal variability in diet as a function of land use change. This improved understanding of

western spotted skunk foraging ecology can elucidate their functional role as small vertebrate

predators, insectivores, and frugivores.

Methods

Study Area

This study was centered around the H. J. Andrews Experimental Forest (HJA), which is

located on the western slope of the Cascade Mountain Range near Blue River, Oregon (Figure 1).

The is surrounded by the McKenzie River Ranger District of the Willamette National Forest.

Elevations range from 410 m to 1,630 m. The maritime climate consists of warm, dry summers

and mild, wet winters. Mean monthly temperatures range from 1°C in January to 18°C in July.

Precipitation falls primarily as rain, is concentrated from November through March, and

averages 230 cm at lower elevations and 355 cm at higher elevations (Greenland 1993; Swanson

and Jones 2002). During 2018 – 2019, western Oregon experienced an extreme drought (USDM

2022). In Lane County, drought severity was greatest during August 2018 – February 2019, but

abnormally dry conditions began as early as January 2018 and moderate drought conditions

began as early as June 2018 (Figure S1).

Lower elevation forests are dominated by Douglas-fir (*Pseudotsuga menziesii*), western

hemlock (*Tsuga hetemphylla*), and western red cedar (*Thuja plicata*). Upper elevation forests are

dominated by noble fir (*Abies procera*), Pacific silver fir (*Abies amabilis*), Douglas-fir, and

western hemlock. The understory is variable and ranged from open to dense shrubs. Common

shrubs included Oregon grape (*Mahonia aquifolium*), salal (*Gaultheria shallon*), sword fern

(*Polystichum munitum*), vine maple (*Acer circinatum*), Pacific rhododendron (*Rhododendron*

macrophyllum), huckleberry (*Vaccinium* spp.), and blackberry and salmonberry (*Rubus* spp.).

Before timber cutting in 1950, 65% of the HJA was covered in old-growth forest.

Approximately 30% of the HJA was clear cut or shelterwood cut to create plantation forests

varying in tree composition, stocking level, and age. In 1980, the HJA became a charter member

of the Long Term Ecological Research network and no logging has occurred since 1985. The

Willamette National Forest immediately surrounding the HJA has a similar logging history, but

logging continues to occur. Currently, the HJA consists of a higher percentage of old-growth

forest than the surrounding Willamette National Forest (approximately 58% in the HJA vs. 37%

in the study area) (Davis et al. *In Press*). Wildfires are the primary disturbance type, followed by

windthrow, landslides, root rot infections, and lateral stream channel erosion. Mean fire return

interval of partial or complete stand-replacing fires for this area is 166 years and ranges from 20

years to 400 years (Teensma 1987; Morrison and Swanson 1990).

Field methods

Our western spotted diet study was part of a larger study on their spatial ecology in the

temperate rainforest ecosystem of western Oregon that was conducted between April 2017 –

September 2019. During this study, we set and maintained 112 baited trail cameras and captured

and tracked western spotted skunks using Tomahawk traps (Model 102 and 103, Tomahawk Live

Trap Co., Hazelhurst, WI) and VHF radio-collars (M1545, 16 g; Advanced Telemetry Systems,

Isanti, MN). Cameras placed in the HJA were paired with previously established long-term

songbird monitoring (Frey, Hadley, and Betts 2016) and small mammal monitoring sites (Weldy

et al. 2019). Cameras placed outside of the HJA were stratified based on elevation and old-

growth structural index (Spies and Franklin 1988) and chosen randomly within logistical

constraints. Both cameras and live traps were baited with a frozen house mouse (*Mus musculus*),

a can of sardines (*Culpidae*), and/or various carnivore scent lures. We located skunks using

radio-telemetry triangulation and homing techniques daily, weather permitting. Homing

techniques were mainly used to locate rest site locations during the day whereas triangulation

was used to locate skunks during the night when skunks were most active. All animal capture

and handling were conducted in accordance with the guidelines set by the American Society of

Mammalogists and were approved by the USDA Forest Service Institutional Animal Care and

Use Committee (IACUC #2016-015) and the Oregon Department of Fish and Wildlife.

166   We collected western spotted skunk scat in multiple ways: 1) during western spotted

skunk capture, 2) opportunistically while tracking western spotted skunks with radio-collars and

checking trail cameras, and 3) using detection dog teams (summer and fall of 2018). Detection

dog teams either surveyed 3 x 3 km grids within the study area for a minimum of 6 hours near

camera trap locations where we detected western spotted skunk or focused their surveys around

known spotted skunk rest sites. Focused surveys were necessary to increase scat sample sizes

and increase spotted skunk scat detection rates. Moreover, western spotted skunk scats were

difficult to locate opportunistically because typically, they were deposited after we tracked

skunks to their rest sites, were in hard to search locations such as in hollow logs or a short

distance from the rest site. We froze all scat samples until we processed them in the laboratory,

and processed scats were dried for long-term storage.

Laboratory methods

In the lab, we identified the diet of western spotted skunks using DNA metabarcoding

(Massey et al. 2021; Eriksson et al. 2019) and mechanical sorting. For DNA metabarcoding, we

extracted DNA in a laboratory dedicated to processing degraded DNA using the DNeasy Blood

and Tissue kit (Qiagen, Germantown, Maryland) or the QIAamp Fast DNA Stool Mini Kit

(Qiagen, Germantown, Maryland). We included an extraction blank with every batch of

extractions as a negative control, where we used the same protocol but without a fecal sample

(hereafter called extraction blanks). We kept extraction blanks throughout the DNA

metabarcoding process.

Following DNA extraction, we amplified 3 regions of the mitochondria and chloroplast

DNA. First, we amplified a ~100 base-pair DNA segment of the ribosomal mitochondrial 12S

gene using universal vertebrate primers (12S-V5-F': YAGAACAGGCTCCTCTAG and 12S-V5-

R: TTAGATACCCCACTATGC) (Kocher et al. 2017; Riaz et al. 2011) and the trnL gene using

universal plant primers (g-F: GGGCAATCCTGAGCCAA and h-R:

CCATYGAGTCTCTGCACCTATC) (Taberlet et al. 2007) in a multiplex polymerase chain

reaction (PCR). In a separate singleplex PCR reaction, we amplified the COI gene using ANML

universal arthropod primers (LCO1490-F: GGTCAACAAATCATAAAGATATTGG and CO1-

CFMRa-R: GGWACTAATCAATTTCCAAATCC) (Jusino et al. 2019). We performed 3 PCR

replicates per sample using the QIAGEN Multiplex PCR kit (Qiagen, Germantown, Maryland).

To aid in identifying contamination, we performed PCR on a negative control on each plate

(hereafter called PCR blanks) in addition to the extraction blanks. Each reaction was amplified

with identical 8 base pair tags on the 5' end of the forward and reverse primer that were unique

to each sample to identify individual sample after pooling and to prevent misidentification of

prey samples due to tag jumping (Schnell, Bohmann, and Gilbert 2015). 12S and trnL PCR

reactions were carried out in a total volume of 20 μL using the following reagent mixtures: 10 μL

QIAGEN Multiplex PCR Master Mix, 4 μL of forward and reverse primer mix for a final primer

concentration of 200 nM, 0.2 μL of bovine serum albumin (BSA), 0.8 μL of water, and 1 μL of

DNA template. After 15 minutes of initial denaturation at 95°C, we conducted 40 cycles of 94°C

for 30 seconds, 58°C for 90 seconds, 72°C for 90 seconds, and a final extension at 72°C for 10

minutes. COI PCR reactions were carried out in a total volume of 20 μL using the following

reagent mixtures: 4 μL of GoTaq Flexi Buffer, 1.2 μL of MgCl$_2$, 0.132 μL of GoTaq Polymerase,

0.4 μL of dNTPs, 0.064 μL of BSA, 6.204 μL of water, 4 μL of each primer for a final

concentration of 200 nM, and 4 μL of final DNA extract elution. After 2 minutes of initial

denaturation at 95°C, we conducted 5 cycles of 94°C for 60 seconds, 45°C for 90 seconds, 72°C

for 90 seconds, followed by 40 cycles of 94°C for 60 seconds, 50°C for 90 seconds, 72°C for 60

seconds, and a final extension at 72°C for 7 minutes. We normalized and pooled the PCR

products and used NEBNext Ultra II Library Prep Kit (New England BioLabs, Ipswich,

Massachusetts) to adapt the library pools into Illumina sequencing libraries (Illumina Inc., San

Diego, California). We purified libraries using the Solid Phase Reversible Immobilization beads

and sent libraries to the Center for Genome Research and Biocomputing at Oregon State

University for 150 base pair paired-end sequencing on the Illumina HiSeq 3000.

We paired raw sequence reads using PEAR (Zhang et al. 2014) and demultiplexed

samples based on the 8-base pair-index sequences using a custom shell script (Text S1). We

counted unique reads from each sample replicate and assigned taxonomy using BLAST against

the 12S, COI, and trnL sequences in a local database and GenBank

(www.ncbi.nlm.nih.gov/blast). Scat amplification was considered successful if DNA sequencing

produced over 100 total reads per replicate, and we limited the effects of contamination by

retaining only species that consisted of more than 1% of the total reads. Furthermore, we used

extraction and PCR negative controls to set additional filtering thresholds for species read counts.

Species were only retained in the final species list if it was present in at least 2 of the 3 replicates

and if their species distribution maps included our study area or were included on the species

lists of the study area (https://andrewsforest.oregonstate.edu/about/species). We identified plants

to genus since congeners are difficult to differentiate using these primers.

To mechanically sort scats, we identified remains to the lowest taxonomic order possible.

We used mechanical sorting to augment results from DNA metabarcoding because of known

biases introduced by mismatches in the universal invertebrate ANML primers we used, which is

attributed to a lack of conserved regions across all invertebrates (Deagle et al. 2014).

We confirmed scats as defecated by western spotted skunks using the metabarcoding data

following criteria: 1) western spotted skunk was the only carnivore (order: Carnivora) identified

in the scat, or 2) western spotted skunk was one of the carnivores identified in the scat and the

other carnivores consisted of less than 10% of the read count.

Data analysis

We conducted analyses and produced figures using the Program R (R Core Team 2019).

We quantified the importance of each taxonomic group by first calculating the frequency of

occurrence of broader taxonomic group of prey (i.e., vertebrate, invertebrate, or plant). We

calculated frequency of occurrence as a proportion of the number of scats in which each

taxonomic group was present divided by the total number of scats. Due to the broad breadth of

diet, we then calculated conditional frequency of occurrence of each species as the number of

scats that contained the prey species divided by the total number of scats containing the broader

taxonomic group. We also calculated the importance of each prey item by calculating the relative

read abundance (RRA) for items identified through DNA metabarcoding by:

$$RRA_i = \frac{1}{S} \sum_{k=1}^{S} \frac{n_{i,k}}{\sum_{i=1}^{T} n_{i,k}}$$

where $n_{i,k}$ is the number of sequences of prey species i in sample k, S is the number of scat

samples, and T is the number of species. We produced figures relating taxonomy of prey items

using the *metacoder* package (Foster, Sharpton, and Grünwald 2017), and we produced

rarefaction curves for each taxonomic group (species for vertebrates, genus for invertebrates and

plants) using the *iNEXT* package (Hsieh, Ma, and Chao 2016) to estimate completeness and

expected taxonomic richness of diet based on sample size.

To investigate the effect of season and logging on western spotted skunk diet, we split

data according to season (dry: June – September, wet: October – May) and location where the

scat was collected (previously logged area vs. no record of logging in area). We summarized the

presence or absences of each taxonomic class (e.g., Mammalia, Insecta, Gastropoda) per sample

and fit a generalized linear model for multivariate data using the manyglm function in the

mvabund package (Wang et al. 2012) to compare diet composition across season and past

disturbance to the area.

Results

During October 2017 – August 2019, we collected and genetically confirmed 130

western spotted skunk scats ($n_{summer} = 47$, $n_{fall} = 62$, $n_{opportunistic} = 21$). 58 scats ($n_{dry} = 32$, $n_{wet} =$

26) were collected from previously logged areas and 72 (n_{dry} = 25, n_{wet} = 47) scats were collected

from areas with no record of timber harvest.

We identified 27 vertebrate species, 43 plant genera, 15 arthropod species, and 3 mollusk

species as prey using DNA metabarcoding (Figure 2). The most frequent prey item (n = 15) was

Atlantic herring (*Clupea harengus*), which we used to bait skunks to trail cameras and traps.

Atlantic herring was the only prey item in 2 scats, so we removed these samples from the

following analyses. After removing Atlantic herring, invertebrates were the most common prey

items identified through metabarcoding and mechanical sorting. Invertebrates occurred in 85.2%

of all scats (n = 109). Vertebrates were the next most common prey item (58.6%, n = 75), and we

detected mammals in 46.9% (n = 60), birds in 14.1% (n = 18), and amphibians in 13.3% (n = 17)

of all scats. Finally, we detected plants in 28.9% of all scats (n = 37).

Wasps (*Vespula* spp.) and millipedes (Diplopoda) were the top invertebrate prey items

comprising 67.0% (n = 73) and 40.4% (n = 44) of scats containing invertebrates, respectively

(Figure 3D). The most frequent vertebrate naturally occurring prey items were shrew mole

(*Neurotrichus gibbsii*), Pacific tree frog (*Pseudacris regilla*), Townsend's chipmunk (*Neotamias*

townsendii), Swainson's thrush (*Catharus ustulatus*), clouded salamander (*Aneides ferreus*),

and Humboldt's flying squirrel (*Glaucomys oregonensis*) (Figure 3E). The most frequent plant

items were Douglas fir (*Pseudotsuga*), maple (*Acer*), Hemlock (*Tsuga*), Gaultheria (*Gaultheria*),

Alder (*Alnus*), and Rhododendron (*Rhododendron*) (Figure 3F).

Of the 130 scats, 51 scats (39.2%) only amplified western spotted skunk DNA and 67

scats (51.5%) did not contain any vertebrate DNA other than Atlantic herring or western spotted

skunk. Manual inspection of all scat samples revealed that the majority consisted of invertebrate

body parts such as wasps (Vespidae) (n = 71), millipedes (Diplopoda) (n = 43), and snail shells

(Gastropoda) (n = 6) that failed to amplify with ANML primers. We also found feathers (n = 3),

snake skin (n = 4), and plant material such as Douglas fir needles and bark (likely consumed

incidentally along with other food items), indicating that DNA quality in some scats did not

allow for detection of all diet items.

Diet composition of western spotted skunks differed based on season (LRT_{season} = 32.0, p

= 0.001), but not based on the collection site's logging history (LRT_{logged} = 10.7, p = 0.35).

During the dry season, diet was composed primarily of insects, mainly wasps (LRT_{insect} = 25.4, p

= 0.001) (Figure 4). Although plant material consumed was similar across season and the

collection site's logging history (LRT_{season} = 0.36, p = 0.996; LRT_{logged} = 0.007, p = 0.98), plants

from genera that produce fruit such as Rubus (n = 6), Vaccinium (n = 1), Gaultheria (n = 10)

were higher during the dry season. During the wet season, western spotted skunks consumed

Gaultheria (n = 3) and Vaccinium (n = 3), but no Rubus. Similarly, although mammals and

amphibians consumed were similar across season and collection site's logging history, wet

season scats consisted of more amphibians such as salamanders (*Rhyacotriton*, *Plethodon*,

Ambystoma, and *Aneides*) and small mammals (*Neotamias townsendii*, *Neurotrichus gibbsii*,

Myodes spp., *Sorex* spp.) (Figure 4).

Discussion

This study provides the first data on the diet of western spotted skunks in the Pacific

Northwest and represents the first use of DNA metabarcoding for high resolution spotted skunk

diet analysis. In the coniferous forests of the Oregon Cascades, the western spotted skunk diet

was highly diverse and included mammals, birds, amphibians, reptiles, insects, gastropods, and

plants. The combined methods of DNA metabarcoding and mechanical sorting revealed that

invertebrates were the primary diet items and mammals were the secondary diet item for western

spotted skunks, which is consistent among other food habit studies conducted on skunks (Selko

1937; Crabb 1941; Baker and Baker 1975; Cantú-Salazar et al. 2005). The importance of these

diet items shifted by season, where skunks consumed more insects during the dry season.

We detected substantially higher frequency of Vespidae than in diets reported on the

island spotted skunk (K. R. Crooks and Van Vuren 1995), the pygmy skunk (Cantú-Salazar et al.

2005), and the plains spotted skunk (Howell 1906; Selko 1937; Crabb 1941). In these studies,

grasshoppers and crickets (Orthoptera) and beetles (Coleoptera) were consumed more frequently

(Howell 1906; Crabb 1941; Baker and Baker 1975; Cantú-Salazar et al. 2005). The scats

collected for this study were collected during a year when the Oregon Cascades were unusually

dry during the spring and summer (Figure S1). These conditions have been shown to be

correlated with irruptions in wasp populations (Akre and Reed 1981; Dejean et al. 2011), and

wasps were observed to be more abundant on the landscape (W. Gerth, personal communication).

Thus, higher consumption of wasps during these irruptions suggests that western spotted skunks

can switch from one prey item to another and capitalize on abundant resources as generalist

predators.

As expected, the vertebrate prey base of the western spotted skunk in Oregon was much

more diverse than that of the island spotted skunk that only consumed one mammal species

(*Peromyscus maniculatus*) (K. R. Crooks and Van Vuren 1995) given that mammal diversity is

greater in the Oregon Cascades. Unlike many of the other studies on skunks (except Howell

1906; Sprayberry and Edelman 2016), the western spotted skunk in the Oregon Cascades

consumed a variety of amphibian species. DNA metabarcoding also revealed that western

spotted skunks consumed 11 avian species that would have been difficult to identify with

mechanical sorting. Given their generalist diet, the western spotted skunk may provide important

linkages between terrestrial, aquatic, and arboreal systems in the Pacific Northwest by

facilitating energy and nutrient transfer.

In addition to linking these disparate systems, western spotted skunks may perform

important roles in ecosystem as seed dispersers, scavengers, and disease reservoirs. We

identified plants in western spotted skunk scats with fruiting bodies including berries (e.g.,

Vaccinium, Rubus, Gaultheria) and mast (e.g., Acer, Quercus, Corylus). Western spotted skunks

may provide key movements that allow for long-distance dispersal of seeds and may influence

plant communities similar to martens and foxes (Jordano et al. 2007; González-Varo, López-Bao,

and Guitián 2013). We also identified black-tailed deer in 3 scats that we collected

opportunistically. We observed radio-collared western spotted skunks scavenging on kills made

by mountain lions (*Puma concolor*) on trail camera videos on multiple occasions and tracked

western spotted skunks to rest sites adjacent to mountain lion kill sites. This behavior has been

observed in other systems (e.g., California scrub oak forest), where the western spotted skunk

has the ability to displace gray fox (*Urocyon cinereoargenteus*) from carcasses and mountain

lions from their kill sites (Allen, Elbroch, and Wittmer 2013; Allen et al. 2016). Furthermore,

this signifies the importance of these kill sites and carrion as food resources that are worth the

risks associated with being near or directly encountering larger predators (Briffa and Sneddon

2007; Allen et al. 2016; Ruprecht et al. 2021). Finally, we identified possible pathways for

transmission of nematode parasites such as *Skrjabingylus* spp. which require spotted skunks as

definitive hosts to complete their life cycle (Kirkland Jr. and Kirkland 1983; Higdon and

Gompper 2020; LaRose, Lesmeister, and Gompper 2021; Lesmeister et al. 2008). Western

spotted skunks have been shown to exhibit high prevalence and high severity of *Skrjabingylus*

spp. infection (Higdon and Gompper 2020). Direct transmission of this nematode may occur

through consumption of gastropods, which are the obligate intermediate host (Lankester and

Anderson 1971; Kirkland Jr. and Kirkland 1983), and indirect transmission may occur through

consumption of gastropod-consuming vertebrates that serve as paratenic hosts such as

chipmunks *(Neotamias townsendii),* shrew moles (*Neurotrichus* gibsii), shrews (*Sorex*

trowbridgii), voles *(Myodes spp.),* and amphibians (Gamble and Riewe 1982). Given that

Skrjabingylus spp. can cause significant osteologic damage to the cranium, it is possible that this

parasite could have significant impacts on individual fitness and population dynamics (Hughes et

al. 2018; Lankester and Anderson 1971).

Seasonal changes in diet for western spotted skunks were similar to other skunk species,

where skunks switched from consuming more insects during the dry season to more vertebrate

prey during the wet season (K. R. Crooks and Van Vuren 1995; Crabb 1941). Island spotted

skunks increased their consumption of crickets during the dry season and mice during the wet

season (K. R. Crooks and Van Vuren 1995). Prairie spotted skunks primarily consumed rabbits

and mice during the winter and spring and insects during the summer and fall (Crabb 1941).

These seasonal changes in diet likely reflect changes in availability and abundance of

invertebrate resources and plasticity in spotted skunk diet (Cantú-Salazar et al. 2005). In addition

to the changes in availability of invertebrate resources, western spotted skunks may switch their

diet to one that includes more vertebrates because they need more caloric and protein-rich input

to thermoregulate and survive the harsher, colder weather (Moors 1977). Moreover, western

spotted skunks breed during the fall during the wet season (Mead 1968) and, if like eastern

spotted skunks (Lesmeister, Gompper, and Millspaugh 2009), males have larger home ranges

when questing for mates, this could increase their energetic requirements, and therefore require

an increase in caloric input.

This was the first study to examine western spotted skunk diet across scat collected from

areas with different logging histories. The similarities in diet across logging history, however, is

not surprising. The western spotted skunk appears to be a generalist species, and many prey

species such as chipmunks may be distributed equally across these forest types. Although some

of the prey species are associated with old-growth forest such as flying squirrels (*Glaucomys*

oregonensis), these species still occur in logged forest, but at lower densities (Carey 1989). A

relatively large portion of our study area (41.5%) is still composed of old-growth forest (Davis et

al. *In Press*), and this amount of old-growth in the larger landscape may help support old-growth

associated species within logged areas. Results may differ in landscapes with few remaining old-

growth stands or in landscapes with more intensive logging operations. In addition, spotted

skunks are highly mobile (Lesmeister, Gompper, and Millspaugh 2009) and can easily move

between logged and unlogged areas in this area. Another possible explanation for similarities in

diet across logged and unlogged areas is that during irruptions, species may spill over into areas

that they are not typically found given lower population densities (Bock and Lepthien 1976).

Although we detected a wide variety of items in western spotted skunk scats,

discrepancies between DNA metabarcoding and mechanical sorting indicated that the ANML

invertebrate primers that we used poorly amplified Vespidae even though we detected Vespula in

some of our samples. The potential for mismatch in the universal invertebrate primers and the

biases introduced by the ANML primers we used is well-known because of a lack of conserved

regions across all invertebrates (Deagle et al. 2014). Still, we used these primers because of the

extensive COI reference library (Jusino et al. 2019; Elbrecht et al. 2019). This highlights the

need for genera-specific primers, better universal invertebrate primers, a panel of invertebrate

primers, or shotgun sequencing that does not rely on PCR to amplify target sequences so that key

prey items are not missed in the future.

We detected a wide variety of plants in the western spotted skunk diet through DNA

metabarcoding, but many of the genera identified may not have been consumed by western

spotted skunks as a food source. When mechanically sorting scats, we discovered many intact

Douglas fir needles and bark imbedded in the scat that were likely consumed incidentally,

environmentally contaminated following defecation, or contaminated during scat collection.

Therefore, care should be taken when interpreting some DNA metabarcoding results using plant

primers.

Noticeably missing from our analysis are the fungal components of the western spotted

skunk diet. Fungi are likely to contribute important nutrients (e.g., vitamins) to the western

spotted skunk diet (Maser, Claridge, and Trappe 2008) and fungal dispersal by mammals is

essential to plants, fungal diversity, and ecosystem function (Nuske et al. 2017). Eastern spotted

skunks have been documented bringing fungal sporocarps to den sites (Sprayberry and Edelman

2016), and we have also recorded western spotted skunks carrying fungal sporocarps on trail

cameras during this study. The importance of fungal diet items remains unknown for western

spotted skunks and an important area of future research.

Together, understanding the diet of western spotted skunks revealed that they may be an

important component of the Pacific Northwest food web. In the face of climate change, western

spotted skunks may possess greater ability to withstand environmental change due to their

dietary flexibility (Reed and Tosh 2019; Ducatez et al. 2020), and through their broad diet and

omnivory, they may help stabilize Pacific Northwest forest food webs (Fagan 1997; Kratina et al.

2012). Through prey-switching, western spotted skunks may capitalize on, help control, and

dampen the effects of irruptions of prey, such as wasps during dry springs and summers, which

could then provide ecosystem resilience to environmental change. Similarly, western spotted

skunks may allow species to persist and recover from low abundances through prey switching.

Acknowledgments

We thank field technicians A. Coombs, B. Murley, K. Van Neste, and R. Rich and

Rogue Detection Dogs (formerly with Conservation Canines) – H. Smith, J. Hartman,

M. Poisson, C. Yee, Chester, Jack, and Scooby – for field support. We thank Levi Lab

technicians for laboratory support. We also thank J. Hedges, S. Speir, and H. Thomas

for mechanically sorting scats. Facilities were provided by the H. J. Andrews

Experimental Forest and the Long Term Ecological Research program, administered

cooperatively by the USDA Forest Service Pacific Northwest Research Station,

Oregon State University, and the Willamette National Forest. Funding for this project

was provided by Oregon State University, the USDA Forest Service, the National

Science Foundation (LTER7 DEB-1440409), the Northwest Ecological Research

Institute, and the ARCS Oregon Chapter. This publication represents the views of the

authors, and any use of trade, firm, or product names is for descriptive purposes only

and does not imply endorsement by the United States Government.

References

Akre, Roger D., and Hal C. Reed. 1981. "Population Cycles of Yellowjackets
(Hymenoptera:Vespinae) in the Pacific Northwest 1." *Environmental Entomology* 10 (3):
267–74. https://doi.org/10.1093/ee/10.3.267.

Allen, Maximilian L., L. Mark Elbroch, and Heiko U. Wittmer. 2013. "Encounter Competition
between a Cougar, *Puma Concolor*, and a Western Spotted Skunk, *Spilogale Gracilis*."
The Canadian Field-Naturalist 127 (1): 64. https://doi.org/10.22621/cfn.v127i1.1410.

Allen, Maximilian L., Christopher C. Wilmers, L. Mark Elbroch, Julie M. Golla, and Heiko U.
Wittmer. 2016. "The Importance of Motivation, Weapons, and Foul Odors in Driving
Encounter Competition in Carnivores." *Ecology* 97 (8): 1905–12.
https://doi.org/10.1002/ecy.1462.

Baker, Rollin H., and Mary W. Baker. 1975. "Montane Habitat Used by the Spotted Skunk
(*Spilogale Putorius*) in Mexico." *Journal of Mammalogy* 56 (3): 671–73.
https://doi.org/10.2307/1379480.

Belant, Jerrold L, Jan Schipper, and James Conroy. 2009. "The Conservation Status of Small
Carnivores in the Americas." *Small Carnivore Conservation* 41: 3–8.

Bock, Carl E., and Larry W. Lepthien. 1976. "Synchronous Eruptions of Boreal Seed-Eating
Birds." *The American Naturalist* 110 (974): 559–71. https://doi.org/10.1086/283091.

Briffa, M., and L. U. Sneddon. 2007. "Physiological Constraints on Contest Behaviour."
Functional Ecology 21 (4): 627–37. https://doi.org/10.1111/j.1365-2435.2006.01188.x.

Cantú-Salazar, Lisette, Mircea G. Hidalgo-Mihart, Carlos A. López-González, and Alberto
González-Romero. 2005. "Diet and Food Resource Use by the Pygmy Skunk (*Spilogale*

Pygmaea) in the Tropical Dry Forest of Chamela, Mexico." *Journal of Zoology* 267 (3):

283–89. https://doi.org/10.1017/S0952836905007417.

Carey, Andrew B. 1989. "Wildlife Associated with Old-Growth Forests in the Pacific Northwest."

Natural Areas Journal 9 (3): 151–62.

Carey, Andrew B. 1995. "Sciurids in Pacific Northwest Managed and Old-Growth Forests."

Ecological Applications 5 (3): 648–61. https://doi.org/10.2307/1941974.

Cheeseman, Amanda E., Brian P. Tanis, and Elmer J. Finck. 2021. "Quantifying Temporal

Variation in Dietary Niche to Reveal Drivers of Past Population Declines." *Functional

Ecology* n/a (n/a). https://doi.org/10.1111/1365-2435.13765.

Crabb, Wilfred D. 1941. "Food Habits of the Prairie Spotted Skunk in Southeastern Iowa."

Journal of Mammalogy 22 (4): 349–64. https://doi.org/10.2307/1374928.

———. 1948. "The Ecology and Management of the Prairie Spotted Skunk in Iowa." *Ecological

Monographs* 18 (2): 201–32. https://doi.org/10.2307/1948639.

Crooks, Kevin. 1994a. "Demography and Status of the Island Fox and the Island Spotted Skunk

on Santa Cruz Island, California." *The Southwestern Naturalist* 39 (3): 257–62.

https://doi.org/10.2307/3671590.

———. 1994b. "Den-Site Selection in the Island Spotted Skunk of Santa Cruz Island,

California." *The Southwestern Naturalist* 39 (4): 354–57.

Crooks, Kevin R., and Dirk Van Vuren. 1995. "Resource Utilization by Two Insular Endemic

Mammalian Carnivores, the Island Fox and Island Spotted Skunk." *Oecologia* 104 (3):

301–7. https://doi.org/10.1007/BF00328365.

Cuarón, A.D., K. Helgen, and F. Reid. 2016. "*Spilogale Gracilis*. The IUCN Red List of

Threatened Species 2016: E.T136797A45221721." International Union for Conservation

of Nature. https://doi.org/10.2305/IUCN.UK.2016-1.RLTS.T136797A45221721.en.

D. Blake Sasse. 2021. "Reexamination of the Purported Rapid Population Decline of Plains

Spotted Skunks in the Mid-Twentieth Century." *Southeastern Naturalist* 20 (sp11): 83–

94. https://doi.org/10.1656/058.020.0sp1109.

Davis, Raymond J., David M. Bell, Matthew J. Gregory, Zhiqiang Yang, Andrew N. Gray, Sean

P. Healey, and Andrew E. Stratton. In Press. "Northwest Forest Plan - the First 25 Years:

Status and Trends of Late Successional and Old-Growth Forests." General Technical

Report PNW-GTR. Portland, Oregon: U.S. Department of Agriculture, Forest Service,

Pacific Northwest Research Station.

De Barba, M., C. Miquel, F. Boyer, C. Mercier, D. Rioux, E. Coissac, and P. Taberlet. 2014.

"DNA Metabarcoding Multiplexing and Validation of Data Accuracy for Diet

Assessment: Application to Omnivorous Diet." *Molecular Ecology Resources* 14 (2):

306–23. https://doi.org/10.1111/1755-0998.12188.

Deagle, Bruce E., Simon N. Jarman, Eric Coissac, François Pompanon, and Pierre Taberlet. 2014.

"DNA Metabarcoding and the Cytochrome c Oxidase Subunit I Marker: Not a Perfect

Match." *Biology Letters* 10 (9): 20140562. https://doi.org/10.1098/rsbl.2014.0562.

Dejean, Alain, Régis Céréghino, James M. Carpenter, Bruno Corbara, Bruno Hérault, Vivien

Rossi, Maurice Leponce, Jérome Orivel, and Damien Bonal. 2011. "Climate Change

Impact on Neotropical Social Wasps." *PLOS ONE* 6 (11): e27004.

https://doi.org/10.1371/journal.pone.0027004.

Doty, Jeffrey B., and Robert C. Dowler. 2006. "Denning Ecology in Sympatric Populations of

Skunks (*Spilogale Gracilis* and *Mephitis Mephitis*) in West-Central Texas." *Journal of*

Mammalogy 87 (1): 131–38. https://doi.org/10.1644/04-MAMM-A-172R2.1.

Ducatez, Simon, Daniel Sol, Ferran Sayol, and Louis Lefebvre. 2020. "Behavioural Plasticity Is

Associated with Reduced Extinction Risk in Birds." *Nature Ecology & Evolution* 4 (6):

788–93. https://doi.org/10.1038/s41559-020-1168-8.

Elbrecht, Vasco, Thomas W A Braukmann, Natalia V Ivanova, Sean W J Prosser, Mehrdad

Hajibabaei, Michael Wright, Evgeny V Zakharov, Paul D N Hebert, and Dirk Steinke.

2019. "Validation of COI Metabarcoding Primers for Terrestrial Arthropods." *PeerJ* 7

(October): e7745–e7745. https://doi.org/10.7717/peerj.7745.

Eriksson, Charlotte E., Katie M. Moriarty, Mark A. Linnell, and Taal Levi. 2019. "Biotic Factors

Influencing the Unexpected Distribution of a Humboldt Marten (*Martes Caurina*

Humboldtensis) Population in a Young Coastal Forest." *PLOS ONE* 14 (5): e0214653.

https://doi.org/10.1371/journal.pone.0214653.

Ewins, P. J., D. V. Weseloh, J. H. Groom, R. Z. Dobos, and P. Mineau. 1994. "The Diet of

Herring Gulls (*Larus Argentatus*) during Winter and Early Spring on the Lower Great

Lakes." *Hydrobiologia* 279 (1): 39–55. https://doi.org/10.1007/BF00027839.

Fagan, William F. 1997. "Omnivory as a Stabilizing Feature of Natural Communities." *The*

American Naturalist 150 (5): 554–67. https://doi.org/10.1086/286081.

Foster, Zachary S. L., Thomas J. Sharpton, and Niklaus J. Grünwald. 2017. "Metacoder: An R

Package for Visualization and Manipulation of Community Taxonomic Diversity Data."

PLOS Computational Biology 13 (2): e1005404.

https://doi.org/10.1371/journal.pcbi.1005404.

Frey, Sarah J. K., Adam S. Hadley, and Matthew G. Betts. 2016. "Microclimate Predicts Within-

Season Distribution Dynamics of Montane Forest Birds." *Diversity and Distributions* 22

(9): 944–59. https://doi.org/10.1111/ddi.12456.

Galan, Maxime, Marie Pagès, and Jean-François Cosson. 2012. "Next-Generation Sequencing

for Rodent Barcoding: Species Identification from Fresh, Degraded and Environmental

Samples." *PLOS ONE* 7 (11): e48374. https://doi.org/10.1371/journal.pone.0048374.

Gamble, R. Lloyd, and R. R. Riewe. 1982. "Infestations of the Nematode Skrjabingylus Nasicola

(Leukart 1842) in Mustela Frenata (Lichtenstein) and M. Erminea (L.) and Some

Evidence of a Paratenic Host in the Life Cycle of This Nematode." *Canadian Journal of*

Zoology 60 (1): 45–52. https://doi.org/10.1139/z82-006.

Gilbert, Frederick F, and Rochelle Allwine. 1991. "Small Mammal Communities in the Oregon

Cascade Range." *USDA Forest Service General Technical Report PNW-GTR-Pacific*

Northwest Research Station (USA).

Gompper, Matthew E. 2017. "Range Decline and Landscape Ecology of the Eastern Spotted

Skunk." In *Biology and Conservation of Musteloids*, 478–92. Oxford University Press.

https://doi.org/10.1093/oso/9780198759805.003.0025.

Gompper, Matthew E., and H. Mundy Hackett. 2005. "The Long-Term, Range-Wide Decline of

a Once Common Carnivore: The Eastern Spotted Skunk (*Spilogale Putorius*)." *Animal*

Conservation 8 (2): 195–201. https://doi.org/10.1017/S1367943005001964.

Gompper, Matthew E., and David S Jachowski. 2016. "*Spilogale Putorius*. The IUCN Red List

of Threatened Species 2016: E.T41636A45211474." International Union for

Conservation of Nature. https://doi.org/10.2305/IUCN.UK.2016-

1.RLTS.T41636A45211474.en.

González-Varo, Juan P., José V. López-Bao, and José Guitián. 2013. "Functional Diversity

among Seed Dispersal Kernels Generated by Carnivorous Mammals." *Journal of Animal*

Ecology 82 (3): 562–71. https://doi.org/10.1111/1365-2656.12024.

Greenland, David. 1993. "The Climate of the HJ Andrews Experimental Forest, Oregon, and Its

Regional Synthesis. USDA Forest Service." *Pacific Northwest Research Station*, 92–

0221.

Higdon, Summer D., and Matthew E. Gompper. 2020. "Prevalence and Severity of Skunk

Cranial Worm (Skrjabingylus Chitwoodorum) Damage Increase with Precipitation in

Spotted Skunks (Spilogale Spp.)." *Journal of Wildlife Diseases* 56 (3): 597–608.

https://doi.org/10.7589/2019-07-182.

Howell, Arthur Holmes. 1906. *Revision of the Skunks of the Genus Spilogale*. US Government

Printing Office.

Hsieh, T. C., K. H. Ma, and Anne Chao. 2016. "INEXT: An R Package for Rarefaction and

Extrapolation of Species Diversity (Hill Numbers)." *Methods in Ecology and Evolution* 7

(12): 1451–56. https://doi.org/10.1111/2041-210X.12613.

Hughes, Malorri R., Nicholas J. Negovetich, Bonny C. Mayes, and Robert C. Dowler. 2018.

"Prevalence and Intensity of the Sinus Roundworm (*Skrjabingylus Chitwoodorum*) in

Rabies-Negative Skunks of Texas, USA." *Journal of Wildlife Diseases* 54 (1): 85–94.

https://doi.org/10.7589/2017-02-023.

Jachowski, David, Roland Kays, Andrew Butler, Anne M. Hoylman, and Matthew E. Gompper.

2021. "Tracking the Decline of Weasels in North America." *PLOS ONE* 16 (7):

e0254387. https://doi.org/10.1371/journal.pone.0254387.

Ji, Yinqiu, Louise Ashton, Scott M. Pedley, David P. Edwards, Yong Tang, Akihiro Nakamura,

Roger Kitching, et al. 2013. "Reliable, Verifiable and Efficient Monitoring of

Biodiversity via Metabarcoding." Edited by Marcel Holyoak. *Ecology Letters* 16 (10):

1245–57. https://doi.org/10.1111/ele.12162.

Jordano, P., C. García, J. A. Godoy, and J. L. García-Castaño. 2007. "Differential Contribution

of Frugivores to Complex Seed Dispersal Patterns." *Proceedings of the National*

Academy of Sciences 104 (9): 3278. https://doi.org/10.1073/pnas.0606793104.

Jusino, Michelle A., Mark T. Banik, Jonathan M. Palmer, Amy K. Wray, Lei Xiao, Emma Pelton,

Jesse R. Barber, et al. 2019. "An Improved Method for Utilizing High-Throughput

Amplicon Sequencing to Determine the Diets of Insectivorous Animals." *Molecular*

Ecology Resources 19 (1): 176–90. https://doi.org/10.1111/1755-0998.12951.

Kartzinel, Tyler R., Patricia A. Chen, Tyler C. Coverdale, David L. Erickson, W. John Kress,

Maria L. Kuzmina, Daniel I. Rubenstein, Wei Wang, and Robert M. Pringle. 2015.

"DNA Metabarcoding Illuminates Dietary Niche Partitioning by African Large

Herbivores." *Proceedings of the National Academy of Sciences* 112 (26): 8019.

https://doi.org/10.1073/pnas.1503283112.

Kinlaw, Al. 1995. "Spilogale Putorius." *Mammalian Species*, no. 511 (October): 1–7.

https://doi.org/10.2307/0.511.1.

Kirkland Jr., Gordon L., and Carol J. Kirkland. 1983. "Patterns of Variation in Cranial Damage

in Skunks (Mustelidae: Mephitinae) Presumably Caused by Nematodes of the Genus

Skrjabingylus Petrov 1927 (Metastrongyloidea)." *Canadian Journal of Zoology* 61 (12):

2913–20. https://doi.org/10.1139/z83-379.

Kocher, Arthur, Benoit de Thoisy, François Catzeflis, Mailis Huguin, Sophie Valière, Lucie

Zinger, Anne-Laure Bañuls, and Jérôme Murienne. 2017. "Evaluation of Short

Mitochondrial Metabarcodes for the Identification of Amazonian Mammals." *Methods in*

Ecology and Evolution 8 (10): 1276–83. https://doi.org/10.1111/2041-210X.12729.

Kratina, Pavel, Robin M. LeCraw, Travis Ingram, and Bradley R. Anholt. 2012. "Stability and

Persistence of Food Webs with Omnivory: Is There a General Pattern?" *Ecosphere* 3 (6):

art50. https://doi.org/10.1890/ES12-00121.1.

Lankester, Murray W., and Roy C. Anderson. 1971. "The Route of Migration and Pathogenesis

of Skrjabingylus Spp. (Nematoda: Metastrongyloidea) in Mustelids." *Canadian Journal*

of Zoology 49 (9): 1283–93. https://doi.org/10.1139/z71-193.

LaRose, Summer H., Damon B. Lesmeister, and Matthew E. Gompper. 2021. "Addressing a

Knowledge Gap in Spilogale Disease Ecology: Skunk Cranial Worm, Skrjabingylus

Chitwoodorum, in Spilogale Putorius Interrupta." *Southeastern Naturalist* 20 (sp11):

173–80. https://doi.org/10.1656/058.020.0sp1114.

Lesmeister, Damon B., Matthew E. Gompper, and Joshua J. Millspaugh. 2008. "Summer Resting

and Den Site Selection by Eastern Spotted Skunks (*Spilogale Putorius*) in Arkansas."

Journal of Mammalogy 89 (6): 1512–20. https://doi.org/10.1644/07-MAMM-A-361.1.

———. 2009. "Habitat Selection and Home Range Dynamics of Eastern Spotted Skunks in the

Ouachita Mountains, Arkansas, USA." *The Journal of Wildlife Management* 73 (1): 18–

25.

Lesmeister, Damon B., Joshua J. Millspaugh, Matthew E. Gompper, and Tony W. Mong. 2010.

"Eastern Spotted Skunk (*Spilogale Putorius*) Survival and Cause-Specific Mortality in

618     the Ouachita Mountains, Arkansas." *American Midland Naturalist* 164 (1): 52–60.

619     https://doi.org/10.1674/0003-0031-164.1.52.

620 Lesmeister, Damon B., Joshua J. Millspaugh, Susan E. Wade, and Matthew E. Gompper. 2008.

621     "A Survey of Parasites Identified in the Feces of Eastern Spotted Skunks (*Spilogale*

622     *Putorius*) in Western Arkansas." *Journal of Wildlife Diseases* 44 (4): 1041–44.

623     https://doi.org/10.7589/0090-3558-44.4.1041.

624 Manning, Tom, Joan C. Hagar, and Brenda C. McComb. 2012. "Thinning of Young Douglas-Fir

625     Forests Decreases Density of Northern Flying Squirrels in the Oregon Cascades." *Forest*

626     *Ecology and Management* 264 (January): 115–24.

627     https://doi.org/10.1016/j.foreco.2011.09.043.

628 Marneweck, C., A.R. Butler, L.C. Gigliotti, S.N. Harris, A.J. Jensen, M. Muthersbaugh, B.A.

629     Newman, et al. 2021. "Shining the Spotlight on Small Mammalian Carnivores: Global

630     Status and Threats." *Biological Conservation* 255 (March): 109005.

631     https://doi.org/10.1016/j.biocon.2021.109005.

632 Maser, Chris, Andrew W Claridge, and James M Trappe. 2008. *Trees, Truffles, and Beasts*.

633     Rutgers University Press.

634 Massey, Aimee L., Gretchen H. Roffler, Tessa Vermeul, Jennifer M. Allen, and Taal Levi. 2021.

635     "Comparison of Mechanical Sorting and DNA Metabarcoding for Diet Analysis with

636     Fresh and Degraded Wolf Scats." *Ecosphere* 12 (6): e03557.

637     https://doi.org/10.1002/ecs2.3557.

638 Mead, Rodney A. 1968. "Reproduction in Western Forms of the Spotted Skunk (Genus

639     *Spilogale*)." *Journal of Mammalogy* 49 (3): 373–90. https://doi.org/10.2307/1378196.

Monterroso, Pedro, Raquel Godinho, Teresa Oliveira, Pablo Ferreras, Marcella J. Kelly, Dana J.

Morin, Lisette P. Waits, Paulo C. Alves, and L. Scott Mills. 2019. "Feeding Ecological

Knowledge: The Underutilised Power of Faecal DNA Approaches for Carnivore Diet

Analysis." *Mammal Review* 49 (2): 97–112. https://doi.org/10.1111/mam.12144.

Moors, P. J. 1977. "Studies of the Metabolism, Food Consumption and Assimilation Efficiency

of a Small Carnivore, the Weasel (Mustela Nivalis L.)." *Oecologia* 27 (3): 185–202.

https://doi.org/10.1007/BF00347466.

Morin, Dana J., Summer D. Higdon, Jennifer L. Holub, David M. Montague, Michael L. Fies,

Lisette P. Waits, and Marcella J. Kelly. 2016. "Bias in Carnivore Diet Analysis Resulting

from Misclassification of Predator Scats Based on Field Identification." *Wildlife Society*

Bulletin 40 (4): 669–77. https://doi.org/10.1002/wsb.723.

Morrison, Peter H, and Frederick J Swanson. 1990. "Fire History and Pattern in a Cascade Range

Landscape." General Technical Report PNW-GTR-254. Portland, Oregon: U.S.

Department of Agriculture, Forest Service, Pacific Northwest Research Station.

Neiswenter, Sean A., and Robert C. Dowler. 2007. "Habitat Use of Western Spotted Skunks and

Striped Skunks in Texas." *Journal of Wildlife Management* 71 (2): 583–86.

https://doi.org/10.2193/2005-623.

Neiswenter, Sean A., Robert C. Dowler, and John H. Young. 2010. "Activity Patterns of Two

Sympatric Species of Skunks (*Mephitis Mephitis* and *Spilogale Gracilis*) in Texas."

Southwestern Naturalist 55 (1): 16–21. https://doi.org/10.1894/PS-51.1.

Nuske, S.J., K. Vernes, T.W. May, A.W. Claridge, B.C. Congdon, A. Krockenberger, and S.E.

Abell. 2017. "Redundancy among Mammalian Fungal Dispersers and the Importance of

Declining Specialists." *Fungal Ecology* 27 (June): 1–13.

https://doi.org/10.1016/j.funeco.2017.02.005.

Proulx, Gilbert. 2010. "We Need More Research and Conservation Programs on

Mesocarnivores." *Canadian Wildlife Biology and Management* 9: 1–2.

R Core Team. 2019. *R: A Language and Environment for Statistical Computing* (version 3.6.1).

Vienna, Austria. https://www.R-project.org/.

Reed, Daniel T., and Colin R. Tosh. 2019. "Diversity Loss Is Predicted to Increase Extinction

Risk of Specialist Animals by Constraining Their Ability to Expand Niche." *Journal of*

Theoretical Biology 476 (September): 44–50. https://doi.org/10.1016/j.jtbi.2019.05.016.

Riaz, Tiayyba, Wasim Shehzad, Alain Viari, François Pompanon, Pierre Taberlet, and Eric

Coissac. 2011. "EcoPrimers: Inference of New DNA Barcode Markers from Whole

Genome Sequence Analysis." *Nucleic Acids Research* 39 (21): e145–e145.

https://doi.org/10.1093/nar/gkr732.

Roemer, Gary W., Matthew E. Gompper, and Blaire Van Valkenburgh. 2009. "The Ecological

Role of the Mammalian Mesocarnivore." *BioScience* 59 (2): 165–73.

https://doi.org/10.1525/bio.2009.59.2.9.

Ruprecht, Joel, Charlotte E. Eriksson, Tavis D. Forrester, Derek B. Spitz, Darren A. Clark,

Michael J. Wisdom, Marcus Bianco, et al. 2021. "Variable Strategies to Solve Risk–

Reward Tradeoffs in Carnivore Communities." *Proceedings of the National Academy of*

Sciences 118 (35): e2101614118. https://doi.org/10.1073/pnas.2101614118.

Sándor, Attila D, and Dan Traian Ionescu. 2009. "Diet of the Eagle Owl (*Bubo Bubo*) in Braşov,

Romania." *North-Western Journal of Zoology* 5 (1).

Schnell, Ida Bærholm, Kristine Bohmann, and M. Thomas P. Gilbert. 2015. "Tag Jumps

Illuminated – Reducing Sequence-to-Sample Misidentifications in Metabarcoding

Studies." *Molecular Ecology Resources* 15 (6): 1289–1303. https://doi.org/10.1111/1755-

0998.12402.

Selko, Lyle F. 1937. "Food Habits of Iowa Skunks in the Fall of 1936." *The Journal of Wildlife*

Management 1 (3/4): 70–76. https://doi.org/10.2307/3795763.

Simmons, Eric A, Micah G Scudder, Todd A Morgan, Erik C Berg, and Glenn A Christensen.

2016. "Oregon's Forest Products Industry and Timber Harvest 2013 with Trends through

2014." *Gen. Tech. Rep. PNW-GTR-942. Portland, OR: US Department of Agriculture,*

Forest Service, Pacific Northwest Research Station. 58 p. 942.

Spies, Thomas A, and Jerry F Franklin. 1988. "Old Growth and Forest Dynamics in the Douglas-

Fir Region of Western Oregon and Washington." *Natural Areas Journal* 8 (3): 190–201.

Sprayberry, Tyler R., and Andrew J. Edelman. 2016. "Food Provisioning of Kits by a Female

Eastern Spotted Skunk." *Southeastern Naturalist* 15 (4).

https://doi.org/10.1656/058.015.0417.

Swanson, Frederick J, and Julia A Jones. 2002. "Geomorphology and Hydrology of the HJ

Andrews Experimental Forest, Blue River, Oregon." *Field Guide to Geologic Processes*

in Cascadia 36: 289–313.

Symondson, W. O. C. 2002. "Molecular Identification of Prey in Predator Diets." *Molecular*

Ecology 11 (4): 627–41. https://doi.org/10.1046/j.1365-294X.2002.01471.x.

Taberlet, Pierre, Eric Coissac, François Pompanon, Ludovic Gielly, Christian Miquel, Alice

Valentini, Thierry Vermat, Gérard Corthier, Christian Brochmann, and Eske Willerslev.

706   2007. "Power and Limitations of the Chloroplast TrnL (UAA) Intron for Plant DNA

707   Barcoding." *Nucleic Acids Research* 35 (3): e14–e14. https://doi.org/10.1093/nar/gkl938.

708   Teensma, Peter Dominic Adrian. 1987. "Fire History and Fire Regimes of the Central Western

709   Cascades of Oregon." Eugene, Oregon: University of Oregon.

710   US Fish and Wildlife Service. 2012. "Endangered and Threatened Wildlife and Plants; 90-Day

711   Finding on a Petition to List the Prairie Gray Fox, the Plains Spotted Skunk, and a

712   Distinct Population Segment of the Mearn's Eastern Cottontail in East-Central Illinois

713   and Western Indiana as Endangered or Threatened Species." *Federal Register* 77: 71159–

714   771.

715   USDM. 2022. "U.S. Drought Monitor." Lincoln, Nebraska, USA: National Drought Mitigation

716   Center at the University of Nebraska-Lincoln, U.S. Department of Agriculture, and

717   National Oceanic and Atmospheric Administration. https://droughtmonitor.unl.edu/.

718   Verts, B. J., Leslie N. Carraway, and Al Kinlaw. 2001. "Spilogale Gracilis." *Mammalian Species*,

719   June, 1–10. https://doi.org/10.1644/1545-1410(2001)674<0001:SG>2.0.CO;2.

720   Wang, Yi, Ulrike Naumann, Stephen T. Wright, and David I. Warton. 2012. "Mvabund– an R

721   Package for Model-Based Analysis of Multivariate Abundance Data." *Methods in*

722   *Ecology and Evolution* 3 (3): 471–74. https://doi.org/10.1111/j.2041-210X.2012.00190.x.

723   Weldy, Matthew J., Clinton W. Epps, Damon B. Lesmeister, Tom Manning, Mark A. Linnell,

724   and Eric D. Forsman. 2019. "Abundance and Ecological Associations of Small

725   Mammals." *The Journal of Wildlife Management* 83 (4): 902–15.

726   https://doi.org/10.1002/jwmg.21641.

Wender, Bryan W, Constance A Harrington, and John C Tappeiner. 2004. "Flower and Fruit

Production of Understory Shrubs in Western Washington and Oregon." *Northwest*

Science 78 (2): 124–40.

Zhang, Jiajie, Kassian Kobert, Tomáš Flouri, and Alexandros Stamatakis. 2014. "PEAR: A Fast

and Accurate Illumina Paired-End ReAd MergeR." *Bioinformatics* 30 (5): 614–20.

https://doi.org/10.1093/bioinformatics/btt593.

Figures

Figure 1. Study area within the Willamette National Forest in the Cascade Range of Oregon,

USA, and locations of western spotted skunk (*Spilogale gracilis*) scats (black crosses). Detection

dog tracks during the summer and fall of 2018 shown in orange, 3 x 3 km survey grids shown in

grey dashed lines, and locations of camera traps shown in maroon circles. Previously logged

areas shown in white. Roads shown in dark grey lines, and outlines of watersheds shown in thick

light grey lines.

Figure 2. Estimation of prey taxonomic richness for western spotted skunks (*Spilogale gracilis*)

in the Willamette National Forest. Vertebrate taxonomic richness (blue line) represents species

richness. Invertebrate (red line) and plant (green line) richness represents genus richness.

Figure 3. Contents of western spotted skunk (*Spilogale gracilis*) scats (n = 130) collected from 2017-2019 in the Willamette National

Forest. Contents determined through DNA metabarcoding and mechanical sorting and organized by (A and D) invertebrates (n = 109),

(B and E) vertebrates (n = 75), and (C and F) plants (n = 37). Taxonomic relationships of diet items shown in top row; color and size

of nodes represent number of occurrences. Frequency of occurrence (FOO) conditional on presence of taxonomic group show in

bottom row.

Note only plants detected in more than one scat were shown.

Figure 4. Frequency of occurrence of taxonomic groups in western spotted skunk (*Spilogale gracilis*) scats (n = 128) collected from

2017-2019 in the Willamette National Forest. Contents determined by DNA metabarcoding and mechanical sorting. Left panel shows

frequency of occurrence by season (dry in red, wet in blue), right panel shows frequency of occurrence by amount of disturbance in

the location we collected the scat (previously logged in tan, no record of logging in green). * represents significant differences in

frequency of occurrence by taxonomic group.

Supplemental Material

Western spotted skunks provide important food web linkages in forest of the Pacific Northwest

Marie I. Tosa, Damon B. Lesmeister, Jennifer M. Allen, Taal Levi

Ecosphere

This material includes:

Text S1. Bioinformatics pipeline

Figure S1**Figure S1.** Weekly climate values for Lane County, Oregon during 2017 – 2019. (A)

Values indicating the percentage of county in drought categories of abnormally dry (D0),

moderate drought (D1), severe drought (D2), extreme drought (D3) and exceptional drought

(D4).

Figure S2. Relative read abundances of (A) invertebrates, (B) vertebrates, and (C) plants in

western spotted skunks (*Spilogale gracilis*) diets during 2017-2019 in the Willamette National

Forest near Blue River, Oregon. Note figure only represents diet items identified through DNA

metabarcoding.

Figure S3. Plant diet of western spotted skunks (*Spilogale gracilis*) identified through DNA

metabarcoding. (A) Plant identified in all scats collected from 2017-2019 (n = 37), (B) plants

identified in scats collected during the dry season (n = 18), and (C) plants identified in scats

collected during the wet season (n = 19) in the Willamette National Forest.

Text S1. Bioinformatics pipeline

1. Pair reads from HiSeq 3000 using PEAR

pearrun -f lane8-s001-index-ATCACG-LibA_S1_L008_R1_001.fastq -r lane8-s001-index-

ATCACG-LibA_S1_L008_R2_001.fastq -n 80 -j 12 -o ./PAIRED/SetA.pear.fastq

2. demultiplex paired reads and BLAST against NCBI database

#!/bin/bash

while read sample lib_index lib f_barcode r_barcode f_primer r_primer locus

do

SampleID=$(echo $sample"_LIB_"$lib"_FBAR_"$f_barcode"_RBAR_"$r_barcode"_LOCUS_"$locus)

793   DIR=$(echo "LIB_"$lib)

794   [-d $DIR] || mkdir $DIR

795   #Make reverse complements of our primer sets

796   f_primer_adj=$(echo $f_primer|sed 's/Y/[CT]/g'|sed 's/W/[AT]/g')

797   r_primer_adj=$(echo $r_primer|sed 's/Y/[CT]/g'|sed 's/W/[AT]/g')

799   f_search=$f_barcode$f_primer_adj

800   rcf_search=$(echo $f_barcode$f_primer_adj|rev|tr ACGT[] TGCA][)

801   r_search=$r_barcode$r_primer_adj

802   rcr_search=$(echo $r_barcode$r_primer_adj|rev|tr ACGT[] TGCA][)

804   for i in $2

805   do

806   grep -oP '(?<='$f_search').*(?='$rcr_search')' $i| sed 's/^/>\n/g' >>

807         $DIR/$SampleID.fasta

808   grep -oP '(?<='$r_search').*(?='$rcf_search')' $i|tr ACGT TGCA| rev | sed

809   's/^/>\n/g' >> $DIR/$SampleID.fasta

```
810
811                            cat $DIR/$SampleID.fasta | fastx_collapser > $DIR/$SampleID.clust.fasta
812        if [ "$locus" == "12s" ]
813        then
814                    MIDORIDB="/nfs1/FW_HMSC/Levi_Lab/Databases/MIDORIpluslocal_UNIQUE_202006
815        18_srRNA_SINTAX.fasta"
816                            VERTMINPROB=0.8
817                            usearch -threads 1 -sintax $DIR/$SampleID.clust.fasta \
818                              -db ${MIDORIDB} -strand plus -sintax_cutoff ${VERTMINPROB} \
819                                -tabbedout $DIR/$SampleID.clust.fasta.usearch
820
821                            blastn -db /nfs1/FW_HMSC/Levi_Lab/Databases/Marten.Nov2019.Blast.fasta \
822                            -query $DIR/$SampleID.clust.fasta \
823                            -outfmt "6 qseqid sseqid sscinames staxids pident qcovs evalue bitscore qseq
824        sseq" \
825                            -max_target_seqs 1 -evalue 1e-5 \
826                            >> $DIR/$SampleID.clust.fasta.assigned
827
828                            blastn -db /nfs1/FW_HMSC/Levi_Lab/Databases/nt_12s_eukaryotes \
829                            -query $DIR/$SampleID.clust.fasta \
830                            -outfmt "6 qseqid sseqid sscinames staxids pident qcovs evalue bitscore qseq
831        sseq" \
832                            -max_target_seqs 1 -evalue 1e-5  \
833                            >> $DIR/$SampleID.clust.fasta.assigned
834
835                            awk -F "\t" '$8 > maxvals[$1] {lines[$1]=$0 ; maxvals[$1]=$8}END { for (tag
836        in lines) print lines[tag] }' $DIR/$SampleID.clust.fasta.assigned | sort -nk1 >
837        $DIR/$SampleID.clust.fasta.assigned.bestmatch
```

```
838    elif [ "$locus" == "COI" ]
839    then
840    MIDORIDBCOI="/nfs1/FW_HMSC/Levi_Lab/Databases/MIDORI_UNIQUE_20180221_COI_SINTA
841    X.fasta"
842            INVERTMINPROB=0.8
843            usearch -threads 1 -sintax $DIR/$SampleID.clust.fasta \
844                -db ${MIDORIDBCOI} -strand plus -sintax_cutoff ${INVERTMINPROB} \
845                    -tabbedout $DIR/$SampleID.clust.fasta.usearch
846
847                        blastn -db /nfs1/FW_HMSC/Levi_Lab/Databases/nt_COI_eukaryotes \
848                            -query $DIR/$SampleID.clust.fasta \
849                            -outfmt "6 qseqid sseqid sscinames staxids pident qcovs evalue bitscore qseq
850    sseq" \
851                            -max_target_seqs 1 -evalue 1e-5 \
852                            >> $DIR/$SampleID.clust.fasta.assigned
853
854                            cat $DIR/$SampleID.clust.fasta.assigned | sort -nk1 >
855    $DIR/$SampleID.clust.fasta.assigned.bestmatch
856
857    elif [ "$locus" == "ITS" ]
858    then
859    ITS="/nfs1/FW_HMSC/Levi_Lab/Databases/sh_general_release_dynamic_s_02.02.2019.fasta"
860            ITSMINPROB=0.8
861            usearch -threads 1 -sintax $DIR/$SampleID.clust.fasta \
862                -db ${ITS} -strand plus -sintax_cutoff ${ITSMINPROB} \
863                    -tabbedout $DIR/$SampleID.clust.fasta.usearch
864
865
```

```
866                              blastn -db

867     /nfs1/FW_HMSC/Levi_Lab/Databases/sh_general_release_dynamic_s_02.02.2019.fasta \

868                         -query $DIR/$SampleID.clust.fasta \

869                         -outfmt "6 qseqid sseqid sscinames staxids pident qcovs evalue bitscore qseq sseq" \

870                         -max_target_seqs 1 -evalue 1e-5 \

871                         >> $DIR/$SampleID.clust.fasta.assigned

872

873              cat $DIR/$SampleID.clust.fasta.assigned | sort -nk1 >

874     $DIR/$SampleID.clust.fasta.assigned.bestmatch

875

876        elif [ "$locus" == "trnL" ]

877        then

878                         blastn -db /nfs1/FW_HMSC/Levi_Lab/Databases/trnL_Kwhite.fasta -task

879     blastn-short \

880                         -query $DIR/$SampleID.clust.fasta \

881                         -outfmt "6 qseqid sseqid sscinames staxids pident qcovs evalue bitscore qseq

882     sseq" \

883                         -max_target_seqs 1 -evalue 1e-5 \

884                         >> $DIR/$SampleID.clust.fasta.ALASKA.assigned

885            cat $DIR/$SampleID.clust.fasta.ALASKA.assigned | sort -nk1 >

886     $DIR/$SampleID.clust.fasta.ALASKA.assigned.bestmatch

887

888

889                         blastn -db /nfs1/FW_HMSC/Levi_Lab/Databases/trnLsequences_nospace.fasta -

890     task blastn-short \

891                         -query $DIR/$SampleID.clust.fasta \

892                         -outfmt "6 qseqid sseqid sscinames staxids pident qcovs evalue bitscore qseq

893     sseq" \
```

```
894                              -max_target_seqs 1 -evalue 1e-5 \

895                              >> $DIR/$SampleID.clust.fasta.trnLnew.assigned

896                    cat $DIR/$SampleID.clust.fasta.trnLnew.assigned | sort -nk1 >

897          $DIR/$SampleID.clust.fasta.trnLnew.assigned.bestmatch

898

899          else

900                              blastn -db nt \

901                              -query $DIR/$SampleID.clust.fasta \

902                              -task blastn -outfmt "6 qseqid sseqid sscinames staxids pident qcovs evalue

903          bitscore qseq sseq" \

904                              -max_target_seqs 1 -evalue 1e-5 \

905                              >> $DIR/$SampleID.clust.fasta.assigned

906

907                    cat $DIR/$SampleID.clust.fasta.assigned | sort -nk1 >

908          $DIR/$SampleID.clust.fasta.assigned.bestmatch

909

910          fi

911                    done

912          done<$1

913
```

Figure S1. Weekly climate values for Lane County, Oregon during 2017 – 2019. (A) Values indicating the percentage of county in

drought categories of abnormally dry (D0), moderate drought (D1), severe drought (D2), extreme drought (D3) and exceptional

drought (D4). (B) Mean temperature difference (°C) compared to mean monthly temperature calculated from data from 1901 – 2000.

Figure S2. Relative read abundances of (A) invertebrates, (B) vertebrates, and (C) plants in western spotted skunks (*Spilogale*

gracilis) diets during 2017-2019 in the Willamette National Forest near Blue River, Oregon. Note figure only represents diet items

identified through DNA metabarcoding.

Figure S3. Plant diet of western spotted skunks (*Spilogale gracilis*) identified through DNA metabarcoding. (A) Plant identified in all

scats collected from 2017-2019 (n = 37), (B) plants identified in scats collected during the dry season (n = 18), and (C) plants

identified in scats collected during the wet season (n = 19) in the Willamette National Forest.