Data integration across conditions improves turnover number estimates and metabolic predictions

Philipp Wendering[1†], Marius Arend[1,2†], Zahra Razaghi-Moghadamkashani[2], Zoran Nikoloski[1,2*]

[1] Bioinformatics, Institute of Biochemistry and Biology, University of Potsdam, Potsdam, Germany

[2] Systems Biology and Mathematical Modelling, Max Planck Institute of Molecular Plant Physiology, Potsdam, Germany

* Zoran Nikoloski

† authors contributed equally

Email: nikoloski@mpimp-golm.mpg.de

Competing Interest Statement: The authors declare no competing interest.

Abstract

Turnover numbers characterize a key property of enzymes, and their usage in constraint-based

metabolic modeling is expected to increase prediction accuracy of diverse cellular phenotypes. *In*

vivo turnover numbers can be obtained by ranking of estimates obtained by integrating reaction

rate and enzyme abundance measurements from individual experiments; yet, their contribution to

improving predictions of condition-specific cellular phenotypes remains elusive. Here we show that

available *in vitro* and *in vivo* turnover numbers lead to poor prediction of condition-specific growth

rates with protein-constrained models of *Escherichia coli* and *Saccharomyces cerevisiae,*

particularly in the ultimate test scenario when protein abundances are integrated in the model. We

demonstrate that *in vivo* estimation of turnover number by simultaneous consideration of

heterogeneous physiological data leads to improved prediction of condition-specific growth rates.

Moreover, the obtained estimates are more precise than the available *in vivo* turnover numbers.

Therefore, our approach provides the means to decrease the bias of *in vivo* turnover numbers and

paves the way towards cataloguing *in vivo* kcatomes of other organisms.

Main Text
30
31   **Introduction**
32
Genome-scale metabolic models (GEMs) together with advances in constrained-based modeling
have led to improved understanding of how cellular resources are used to fulfill different cellular
tasks (Goelzer *et al*, 2015; Lerman *et al*, 2012; O'Brien *et al*, 2013). Recent advances are largely
propelled by the development of protein-constrained GEMs (pcGEMs) in which the catalytic
capacities of individual enzymes are linked to the allocation of enzyme abundances (Chen &
Nielsen, 2021a). Such models have led to more accurate predictions of maximum specific growth
rates on different carbon sources (Adadi *et al*, 2012; Beg *et al*, 2007; Sánchez *et al*, 2017), flux
distributions (Sánchez *et al*, 2017), and other complex phenotypes (Malina *et al*, 2021) in
Escherichia coli and *Saccharomyces cerevisiae*. However, the development of pcGEMs critically
depends on integration of organism-specific enzyme turnover numbers, k_{cat}, comprising the
kcatome of an organism (Nilsson *et al*, 2017).
Measuring the kcatome of an organism based on *in vitro* characterization is limited due to
impossibility to purify specific enzymes, lack of availability of substrates, and knowledge of required
cofactors, such that their relevance for studies of *in vivo* phenotypes remains questionable (van
Eunen & Bakker, 2014; Labhsetwar *et al*, 2017). Proxies for turnover numbers, referred to as
maximal *in vivo* catalytic rates, can be obtained by combining constrained-based approaches for
flux prediction with measurements of protein abundance under different growth conditions or
genetic modifications (Davidi *et al*, 2016; Heckmann *et al*, 2020; Xu *et al*, 2021). The results from
this approach, that entails ranking of condition-specific estimates that use individual data sets, have
shown that the proxies for *in vivo* turnover numbers generally concur with *in vitro* k_{cat} values in *E.*
coli (Davidi *et al*, 2016). However, applications with data from *S. cerevisiae* (Chen & Nielsen,
2021b; Heckmann *et al*, 2020) and *A. thaliana* (Küken *et al*, 2020) indicated that these proxies for
in vivo turnover numbers do not reflect *in vitro* measurements. Another approach to estimate the
kcatome relies exclusively on machine and deep learning methods that use variety of features of
enzymes (e.g. network-based, structure-based, and biochemical) (Zikmanis & Kampenusa, 2012;
Preprint: Li et al, 2021; Heckmann *et al*, 2018), resulting in predictive models that can explain up
to 70% of the variance in turnover numbers obtained *in vitro*.
The estimates of turnover numbers are integrated into metabolic models by different constraint-
based approaches that have been grouped into coarse-grained (e.g. MOMENT (Adadi *et al*, 2012),
sMOMENT (Bekiaris & Klamt, 2020), eMOMENT (Wendering & Nikoloski, 2022), and GECKO
(Sánchez *et al*, 2017)) and fine-grained (e.g. resource balance analysis (Goelzer *et al*, 2015;
Lerman *et al*, 2012) and ME-models (O'Brien *et al*, 2013)). Of these, GECKO (Sánchez *et al*, 2017)
has been adopted in several recent studies due to the elegantly structured formulation of the protein
constraints. In addition, GECKO allows for integration of protein contents and correction factors

that account for mass fraction of enzymes (f) included in the model as well as average *in vivo*
saturation (σ) of all enzymes, facilitating the development of condition-specific models.

While data-driven estimation of *in vivo* turnover numbers improves the coverage of k_{cat} values
in pcGEMs, the available estimates usually lead to over-constrained models when using the
allocation of total protein mass, not considered in flux balance analysis (FBA). In addition, the bias
in the estimates of k_{cat} values can be determined in the ultimate test, in which they are used
alongside constraints from enzyme abundances to predict growth--a scenario that has not yet been
examined. Here, we propose PRESTO (for protein-abundance-based correction of turnover
numbers), a scalable constraint-based approach to correct turnover numbers by matching
predictions from pcGEMs with measurements of cellular phenotypes – simultaneously – over
multiple conditions. As a constraint-based approach, PRESTO facilitates the investigation of
variability of the proposed corrections. We show that predictions of growth by pcGEMs of *S.*
cerevisiae with turnover numbers corrected by PRESTO are more accurate than those based on
the models that include k_{cat} values corrected based on a contending heuristic that relies on enzyme
control coefficients (Preprint: Domenzain et al, 2021). We also demonstrate that same conclusions
hold when enzyme abundances are integrated into the *E. coli* model using PRESTO. Therefore,
PRESTO paves the way to broaden the applicability of pcGEMs for organisms with biotechnological
applications and to arrive at genotype-specific estimates of the kcatome.

85
86 **Results and Discussion**
87
PRESTO – protein-abundance-based correction of turnover numbers

For a given data set of protein abundances over a set of conditions, the enzymes with turnover
numbers in a pcGEM can be partitioned into three groups. For instance, a data set of protein
abundances that was recently used to estimate *in vivo* turnover numbers in *S. cerevisiae* (Chen &
Nielsen, 2021b) includes 45%, 41%, and 14% measured over all, at least one (but not all), and
none of the 27 used conditions, respectively. Therefore, there is then a different data support for
correcting the k_{cat} values of these classes of proteins. PRESTO relies on solving a linear program
that minimizes a weighted linear combination of the average relative error for predicted specific
growth rates and the correction of the initial turnover numbers integrated in the pcGEM model (Fig.
1, Methods). It further employs K-fold cross validation (here, K = 3) with 10 repetitions while
ensuring steady state and integrating protein constraints for proteins measured over all conditions
(Fig 1, Methods). The training set of conditions is used to generate a single set of corrected k_{cat}
values, that provide estimates of *in vivo* turnover numbers, by using the respective protein
abundances. The resulting corrected k_{cat} values are in turn used to determine the relative error of
predicted specific growth rate for each condition in the test set using flux balance analysis with the
pcGEM, while only applying a constraint on total protein content and measured uptake rates. The

relative error of predicted specific growth rate along with the sum of introduced corrections are
lastly used to select the value for the tuning parameter λ in the objective function of PRESTO, as
done in machine learning approaches that rely on regularization.

PRESTO outperforms a contending heuristic for correction of turnover numbers in *S. cerevisiae*

To determine the performance of PRESTO and compare it to that of contending heuristics, we used
a data set comprising protein abundances and exchange fluxes from 27 diverse conditions, as
supported by the principal component analysis (Fig S1). Application of PRESTO with a pcGEM of
S. cerevisiae with initial *in vitro* turnover numbers obtained from BRENDA resulted in a mean
relative error of 0.68 from the cross-validation procedure, yielding a correction of on average 213
turnover numbers (Fig S2a). For the *S. cerevisiae* pcGEM, we found a value of 10^{-7} for the
parameter λ in the PRESTO objective provides the optimal trade-off between both the relative error
and the sum of introduced corrections (see Methods). Moreover, we observed a high overlap
between the sets of proteins with corrected turnover numbers in the cross-validation (average
Jaccard distance of 0.07 (Fig S2b, c), suggesting that the integrated data from different conditions
point at a specific subset of enzymes that need to be corrected to improve performance of growth
prediction.

Unlike PRESTO, GECKO implements a heuristic for correction of turnover that is based on the
control coefficient of a protein in each condition (Fig S3) (Preprint: Domenzain et al, 2021). The
control coefficient of a protein is determined by increasing the turnover number by 1000-fold and
scoring the effect on the predicted specific growth rate. The proteins are then ranked in a
decreasing order of their control coefficients, and the turnover number of the first enzyme in the list
is changed to the maximum value found in BRENDA for this enzyme across all organisms. This
procedure is repeated with the remaining enzymes until the pcGEM predicts a growth rate that is
at most 10% smaller than the measured specific growth rate for that condition or no additional k_{cat}
value that strongly constraints the solution can be found. In contrast to this procedure, PRESTO
corrects at once the turnover numbers of multiple enzymes that are measured in all investigated
conditions by simultaneously leveraging the data from the different conditions, considerably
reducing runtime and the number of solved problems. As a result, rather than deriving condition-
specific corrected k_{cat} values, which are difficult to use in making predictions for unseen scenarios,
PRESTO results in a single set of corrected k_{cat} values.

Next, we compared the performance of PRESTO with the heuristic implemented in GECKO in
three modeling scenarios that consider: (i) only condition-specific total protein content, (ii) both total
protein content and uptake constraints, and (iii) additional constraints from abundances of enzymes

measured in all conditions. For corrections of turnover number from PRESTO, we observed that
the relative error spans the range from 0.15 to 0.88 in the least constrained scenario (i) (Fig 2a)
and from 0.69 to 0.98 in the most constrained scenario (iii) (Fig 2c). In contrast, the relative error
with the corrections of turnover numbers from the GECKO heuristic is in the range from 0.96 to
1.00 in scenario (iii) (Fig 2c). In addition, in scenario (iii), the relative error in the case of the GECKO
heuristic for each condition is larger than the relative error of the PRESTO predicted specific growth
rate (Fig 2c). We observed that predictions from FBA, considering enzyme abundances, without a
constraint on the total protein content, led to an average relative error of 0.70 with k_{cat} values
corrected according to PRESTO and 0.99 with k_{cat} values corrected according to GECKO (Table
S1A).

We also performed sensitivity analysis by investigating a smaller value, of 10^{-10}, for the
weighting factor λ used in the PRESTO objective. We found that when the weighting factor is 10^{-10}
(at which the total corrections of the initial k_{cat} values plateaus), the relative errors from PRESTO
can be further decreased to 0.69 with the constraint on the total protein content, with no effects on
the other findings (Fig S2a). We also note that the relative error lies in the range from 0.35 to 0.80
over the considered weighting factors in the range from 10^{-14} to 10^{-1}. Together, these results
demonstrated that k_{cat} values corrected according to PRESTO provide better model performance
than the values obtained by the contending heuristic in the case of *S. cerevisiae*.

PRESTO provides precise corrections and estimates of *in vivo* turnover numbers

In the following, we investigated the precision of the corrected k_{cat} values from the application of
PRESTO to data and pcGEM model of *S. cerevisiae*. To this end, we determined the range that
the correction of the k_{cat} value of each enzyme can take, while fixing the relative error in specific
growth rate and total corrections from the optimum of PRESTO (Methods). Moreover, we
complemented this analysis by sampling corrected k_{cat} values that achieve the optimum of
PRESTO with two values of the weighting factor λ of 10^{-7} and 10^{-10}.

In the case of the corrected k_{cat} values for *S. cerevisiae* with a weighting factor of 10[-7], we found
that the k_{cat} values with the largest corrections are more precisely determined (Fig S4). In addition,
the sampled corrections per enzyme show an average Euclidean distance to the respective mean
of 4.88 s[-1], indicating that the corrected values are more precise than the values in BRENDA,
exhibiting an average Euclidean distance of 27.54 s[-1] to the mean per EC number (Fig S5).
Importantly, while k_{cat} values with smaller correction showed larger variability, the 25 and 75
percentiles of the sampled corrections for 42 enzymes are concentrated around those resulting
from PRESTO. Repeating the analysis with a weighting factor of 10[-10] showed that the larger total
corrections of the initial k_{cat} values resulted in also larger variability for the corrections over all k_{cat}

(Fig S6). Here, too, for 62 enzymes the 25 and 75 percentiles of the sampled corrections are
concentrated around those resulting from PRESTO. Therefore, we concluded that the corrections
from PRESTO are precise and can be used in downstream analyses.

**175   Pathways of carbon metabolism are enriched in enzymes with corrected turnover numbers
in pcGEM model of *S. cerevisiae***

In pcGEMs generated by the GECKO toolbox (Sánchez *et al*, 2017), turnover numbers are
assigned to each enzymes in the GEM using a fuzzy matching algorithm. It takes into account the
organism, substrate, and EC number of a BRENDA entry. When we investigated the magnitude of
the turnover number correction dependent on the quality of the match between BRENDA entry and
the corresponding enzyme, we found that k_{cat} values measured in *S. cerevisae* were associated
with smaller corrections then those from other organisms (Fig S7a).

To check which metabolic processes are more likely to require correction of *in vitro* k_{cat} values,
we next conducted an enrichment analysis based on the KEGG pathway terms linked to corrected
k_{cat} values (Methods). The most prominent pathway in this analysis was the synthesis of secondary
metabolites, particularly the synthesis of cofactors and terpenoids (Fig 3a). However, several terms
linked to central carbon metabolism, such as: the tricarboxylic acid cycle and oxidative
phosphorylation, were also significantly enriched. Interestingly, amino acid synthesis was the only
term linked to nitrogen metabolism that came up in this analysis, although many pathways of
nitrogen metabolism were among the tested terms. This analysis suggested that particularly *in vitro*
turnover numbers in carbon metabolism need to be corrected, due to underestimation in *in vitro*
assays.

Next, we aimed to identify the extent to which the corrected k_{cat} values differ between PRESTO
and the GECKO approach. To this end, we determined the intersection of enzymes with k_{cat} values
corrected manually (Sánchez *et al*, 2017), by PRESTO, and by the GECKO heuristic. For this
comparison, we considered the union of all condition-specific corrected k_{cat} values from the
GECKO approach. With the weighing factor $\lambda = 10^{-7}$, PRESTO adapted the k_{cat} values of 48% of
enzymes corrected by the GECKO heuristic (Fig 3b, Table S1B). We did not find a significant
Spearman correlation ($\rho_S = 0.17$, $P = 0.45$) between the log-transformed k_{cat} values in this
intersection (Fig 3c), owing to the different principles employed in the two procedures. To determine
the pathways that comprise enzymes whose turnover number are corrected by GECKO and
PRESTO, we next repeated the pathway enrichment analysis for the enzymes in the overlap.
Among the significant terms, like in PRESTO, we again found 2-Oxocarboxylic acid, amino acid,
and secondary metabolism were enriched (Fig 3a, S8). However, the more specific pathway terms
were associated with pathways that are part of carbohydrate metabolism and aromatic amino acid

metabolism corrected by both approaches (Fig S8). In addition, the intersection between enzymes
with manually corrected values and those corrected by the GECKO heuristic is higher than with
PRESTO. This is expected since the manual curation partly aimed at correcting the most
constraining turnover numbers (Sánchez *et al*, 2017).

We also compared the k_{cat} values adjusted by GECKO against values for the maximum
apparent catalytic rate obtained by parsimonious FBA (pFBA) using the same proteomics data
(Chen & Nielsen, 2021b) (Fig S9 a, b). We confirmed the low correspondence ($\rho_S = 0.23$) between
the k_{cat} values obtained from BRENDA, included in the GECKO model without manual
modifications, and the maximum apparent catalytic rates. As expected, the correspondence of the
maximum apparent catalytic rates to the turnover numbers corrected based on PRESTO was
higher ($\rho_S = 0.34$). To investigate how turnover numbers obtained from pFBA perform as model
parameters, we also generated a pcGEM in which BRENDA values were substituted by the
maximum apparent catalytic rates from (Chen & Nielsen, 2021b), whenever available. In scenarios
without enzyme abundance values, this model performed worse than that including the k_{cat} values
corrected by PRESTO as well as the model combining the maximum of all condition-specific
GECKO corrections (Fig S10 a, b). In the enzyme abundance constrained scenario, the model with
turnover numbers obtained from pFBA performed slightly better than GECKO but still only achieved
a minimum relative error of 0.93, which is larger than 0.69 resulting from PRESTO (Fig S10c).
These results demonstrated the value of the PRESTO in combining the genome-scale coverage of
BRENDA with *in vivo* proteomics chemostat measurements to obtain less biased estimates of *in*
vivo k_{cat} values.

Application of PRESTO with protein-constrained model of *E. coli* metabolism

To demonstrate the applicability of PRESTO across species, we deployed it with a pcGEM of *E.*
coli (eciML1515) (Preprint: Domenzain et al, 2021; Monk *et al*, 2017). To this end, we used a large
dataset comprising 31 different growth conditions (Valgepea *et al*, 2013; Peebo *et al*, 2015; Schmidt
et al, 2016; Davidi *et al*, 2016). Due to the lack of data on nutrient exchange rates, the same GAM
value (i.e., 75.55 $\frac{mmol}{gDWh}$) was used across all conditions. Similarly, we used the same value for total
protein content since condition-specific measurements were not available (see Methods).

By applying a three-fold cross-validation, we found the optimal value for the λ parameter to be
10^{-5} (Fig S11a). This value was associated with an average relative error of 1.95 (overall average:
3.32) and 73 corrected turnover numbers, while on average 156 k_{cat} values were corrected across
all explored values for λ. On average, the Jaccard distance between cross-validation folds was
0.13 (Fig S11b), while the average Jaccard distance between unique sets of enzymes with

corrected turnover numbers for each λ parameter was three-fold larger (0.4, Fig S11c). Thus, the

corrected k_{cat} values among cross-validation folds for each λ are more similar (maximum Jaccard

distance of 0.29). Moreover, the union of the set of enzymes with corrected k_{cat} values can remain

similar over a range of chosen λ parameters up to four orders of magnitude (Fig S11c),

demonstrating the robustness of the method.

The performance of PRESTO was assessed and compared to GECKO using scenarios (i) and

(iii) since no condition-specific uptake rates were available. With default uptake rates, the relative

error for predicted growth ranged between 0.01 and 8.56 in the less constrained scenario (i) (Fig

4a). Further, we obtained relative errors between 0.01 and 0.88 for the more constrained scenario

(iii), when using the k_{cat} values corrected by PRESTO (Fig 4b). In contrast, when using the k_{cat}

values from the GECKO approach, the relative error was in the range between 0.01 and the 4.89

for scenario (i) and between 0.89 and 0.99 for scenario (iii). In this scenario, too, we observed that

the relative error using k_{cat} values corrected by GECKO was consistently larger than the relative

error resulting from the single set of corrected k_{cat} values obtained by PRESTO (Fig 4a, b).

The sum of introduced k_{cat} corrections reached a plateau at 10^{-11} for the weighting factor λ in

the PRESTO objective. We found that the relative cross-validation error at this value was 5.26,

which is 2.7-fold larger than the relative error obtained using the optimal λ. Hence, allowing for

more and larger corrections in PRESTO leads to a decrease of the overall relative error within the

PRESTO program at the cost of highly biased parameters. The predictions with the highly biased

parameters are worse in the test conditions and result in larger specific growth rate when no

enzyme abundance constraints are enforced. This observation is in line with the small number of

corrections introduced by the GECKO approach, where only the pool constraint is considered. We

conclude that the prediction performance of the eciML1515 model was improved by using turnover

numbers corrected by PRESTO only when enzyme abundances are integrated.

To assess the precision of the introduced k_{cat} corrections, we performed variability analysis and

sampling (see Methods) of the introduced corrections to the initial k_{cat} values for two values of the

weighting factor λ, namely 10^{-5} and 10^{-11}. We observed that the 25 and 75 percentiles enclose a

narrow interval around the values resulting from PRESTO (Fig S12) and are thus not evenly

distributed across the respective interval determined by the variability analysis. We further noted

that here, the predictions of smaller δ are generally more precise than the large corrections ($\delta \geq$

p_{50}), which span 2.12 orders of magnitude (small δ (< p_{50}): 1.83, Fig S12). However, we also

observed that the precision decreased when more corrections were allowed in PRESTO. This

further justified our choice for the optimal parameter λ, which results in a lower number of 73

corrections compared to 204 at $\lambda = 10^{-11}$, and moreover guarantees more precise estimates (Fig

S13). In conclusion, the application of PRESTO is not limited to a single species but presents a
versatile tool for the correction of turnover numbers across species.

In contrast to the observations made in *S. cerevisiae* we found that a model parameterized with
the turnover numbers estimated by pFBA (Davidi *et al*, 2016) outperformed both PRESTO and
GECKO in the modeling scenario where no enzyme abundance constraints are taken into account
(Fig S10d). This is due to the fact, that pFBA, in contrast to PRESTO and GECKO, allows for the
decrease of *in vitro* k_{cat} values, in turn leading to more accurate predictions. However, in the
scenario with enzyme abundance constraints PRESTO predicts specific growth rates closer to the
experimental observation in 87% of the conditions (Fig S10e). Thus, in this scenario the integration
of information from different modeling conditions achieved in PRESTO serves to obtain *in vivo* k_{cat}
value that perform better than the k_{app}^{max} approach applied by (Davidi *et al*, 2016).

Interestingly, in contrast *S. cerevisiae*, we did not observe larger corrections by PRESTO for
organism-unspecific k_{cat} values (Fig S7b). The low number of corrections introduced by GECKO
leads to an overlap of only three (75%) enzymes whose k_{cat} values were also corrected by
PRESTO (Table S1C, Fig S14a). These three enzymes catalyze reactions in three distinct
metabolic pathways: Phosphoribosylformylglycinamidine synthase acts in the synthesis of purines,
while serine acetyltransferase and NADP dependent Ketol-acid reductoisomerase are involved in
the synthesis of sulfur aminoacids and hydrophic amino acids, respectively. The pathway
enrichment analysis for all PRESTO corrections at $\lambda = 10^{-5}$, indeed also identified amino acid and
secondary metabolite synthesis as significantly enriched terms among the enzymes with corrected
turnover numbers (Fig S14b). These results argue for a systematic underestimation of *in vivo*
turnover numbers in these pathways compared to *in vitro* data, irrespective of the investigated
organism. However, the lower order KEGG pathway terms enriched in *E. coli* do not overlap with
the ones found in *S. cerevisiae*. Here, fatty acid metabolism and the synthesis of hydrophobic
amino acids are among the pathways requiring correction of turnover numbers.

Robustness of turnover number corrections
All of the approaches for estimation of turnover numbers rely on predicted (or estimated) fluxes and
protein abundances from multiple conditions (Davidi *et al*, 2016; Chen & Nielsen, 2021b; Xu *et al*,
2021), but have not investigated the robustness of the estimates to the number of conditions used.
Therefore, next, we investigate the difference in the sets of enzymes with corrected turnover
numbers and the concordance of their corrections when ten randomly sampled subcollections of M
experimental conditions (M = 3, 5, 10, 15) was used instead of all experiments. The differences
and concordance were quantified with respect to the estimates obtained by considering data from
all available experiments using the Jaccard index and the Pearson correlation coefficient,

respectively. In the case of *S. cerevisiae,* we found that the smallest Jaccard difference over 200
scenarios was 0.36, while for *E. coli* this was 0.41 (Figs S15 and S16). In addition, the Pearson
correlation coefficient between the (log-transformed) corrected turnover numbers with
consideration of all versus a subcollection of M experiments in *S. cerevisiae* ranged from 0.99 to
1.00 (for M = 15) to 0.11 and 1.00 (for M = 3) (Fig S15). Repeating the analysis in the case of *E.*
coli, we found that that the Pearson correlation coefficient ranged from 0.15 to 1.00 (for M = 15) to
0.14 and 1.00 (for M = 3) (Fig S16). This is in line with the expectation that the corrections stabilize
with increasing number of experiments. Altogether, these findings pointed out the robustness of
turnover number corrections derived from PRESTO with the number of available experiments.
316
317 **Conclusion**
318
Characterization of enzyme parameters that can inform models of reaction rates is key to
expanding and further propelling the usage of metabolic models in diverse biotechnological
applications. While the generation of pcGEMs has facilitated the integration of more biophysically
relevant constraints, it necessitates access to estimates of turnover numbers as key enzyme
parameters. The bias in the available *in vitro* and *in vivo* turnover numbers can readily be assessed
by considering the accuracy of growth predictions based on the integration of protein abundances.
Indeed, we showed that condition-specific growth rates cannot be reliably predicted with pcGEMs
of *S. cerevisiae* and *E. coli* when available *in vitro* and *in vivo* estimates of turnover numbers are
used.
We use the modeling scenario that considers measured protein abundances as the ultimate test
scenario not only for the prediction of metabolic fluxes, but also for the prediction of specific growth
rates as it contains considerably more biochemically relevant constraints. The usage of protein
abundances for estimation of proxies for *in vivo* turnover numbers also warrants their integration in
pcGEMs to predict specific growth rates. However, we found that the integration of protein
abundances resulted in poor predictions of specific growth rates. GECKO resolves this issue by
flexibilizing measured protein abundances without considering physiological information during the
procedure. To overcome this limitation, we developed PRESTO, which corrects turnover numbers
and facilitates the integration of enzyme abundance constraints.
In contrast to PRESTO, GECKO uses measured total protein content from a single condition to
achieve specific growth rates in the process of correcting the turnover numbers. As a result, the
corrected turnover numbers vary between different experiments. Like in all existing approaches for
estimation of turnover numbers based on GEMs, we integrated protein abundance data directly to
correct turnover numbers. Following this strategy in PRESTO is further justified by the observation
that the turnover numbers included in pcGEMs are often neither from the same enzyme (i.e., EC
number), substrate, nor organism. While *in vivo* turnover number estimates can be adjusted by

considering recently proposed Bayesian statistical learning {Formatting Citation}, this approach has
not considered protein abundance information from proteomics measurements.

To resolve this issue, we proposed PRESTO, a constraint-based approach that simultaneously
considers heterogeneous physiological read-outs and enzyme abundance measurements to
correct *in vitro* turnover numbers. Through a series of comparative analyses, we demonstrated that
the *in vivo* estimates of turnover numbers from PRESTO ultimately increase the prediction accuracy
of condition-specific growth for the two organisms when enzyme abundance data are integrated in
the corresponding pcGEMs. We also showed that the maximal *in vivo* catalytic rates, obtained by
ranking of condition-specific estimates that use proteomics and fluxomics data, are more highly
(but modestly) correlated to estimates from PRESTO than to *in vitro* turnover numbers. Owing to
the constraint-based formulation of PRESTO, we also determined the precision of the *in vivo*
estimates of turnover numbers. Previous studies have shown that even for the well-studied model
organism *Sacharomyces cerevisiae*, only 52% of enzyme turnover numbers in the pcGEM can be
obtained from organism-specific *in vitro* measurements (Preprint: Domenzain et al, 2021). Using
organism unspecific k_{cat} values for parameterization and correction of pcGEMs, as done in the
GECKO pipeline, assumes that enzyme kinetic properties are comparable within one EC number
class (Bar-Even *et al*, 2011; Davidi *et al*, 2018). However, we did not identify clear differences
between EC classes, down to the second digit, when considering the distribution of k_{cat} similarities
within EC classes (Fig S17). Indeed, it has been reported that EC class plays only a minor role in
the prediction of turnover numbers (Heckmann *et al*, 2018) and show stronger similarity with
concordant GO categories (Mao & Ma, 2019). Interestingly, our findings show that the *in vivo*
turnover numbers obtained from PRESTO are more centered around the means of the of the EC
classes for both studied organisms (Fig S5). Together, these findings demonstrated PRESTO can
be readily used to decrease the bias of *in vitro* and *in vivo* estimates of turnover numbers. This
paves the way for employing the outcome of PRESTO and future extensions towards effectively
predicting the kcatome from available protein sequences.

370
Materials and Methods

Experimental data

S. cerevisiae We made use of a dataset gathered by (Chen & Nielsen, 2021b) from four different
studies (Lahtvee *et al*, 2017; Yu *et al*, 2020, 2021; Di Bartolomeo *et al*, 2020), which included
protein abundance data as well as measured growth or dilution rates and nutrient exchange fluxes.
Exchange fluxes missing in certain conditions were set to 1000 mmol/gDW/h if the nutrient was
present in the used culture media. We further augmented this data set by adding total protein
content measurements from the original studies. For subsequent analyses, we used the maximum
abundance of each protein over all replicates per experimental condition. Similarly, we used the

average value for specific growth rates and nutrient exchange rates. Since no measurement of total
protein content was available for the two conditions evaluated in the Di Bartolomeo study (Di
Bartolomeo *et al*, 2020), we used the maximum protein content measured across the remaining
conditions for these conditions (i.e., $0.67\ g/gDW$). Moreover, we excluded three temperature
stress conditions (i.e., Lahtvee2017_Temp33, Lahtvee2017_Temp36, Lahtvee2017_Temp38)
from the analysis since temperature can have a large effect on the catalytic activity of an enzyme.
Gene names in the proteomics dataset were translated to UniProt identifiers using the batch
retrieval service of the UniProt REST API (Bateman, 2019).

E. coli We used a dataset comprising 31 experimental conditions, which was gathered by Davidi
and colleagues and augmented by Xu et al. (Davidi *et al*, 2016; Xu *et al*, 2021) from three
publications (Valgepea *et al*, 2013; Schmidt *et al*, 2016; Peebo *et al*, 2015). Here, too, we used the
maximum protein abundance over all replicates. Due to the absence of total protein content
measurements in two of the original studies, we relied on the maximum protein content measured
in the Valgepea study (i.e., $0.61\ g/gDW$) to be used for all conditions. Since precise data on
nutrient uptake rates were only given for a few conditions, we assigned a default upper bound of
1000 mmol/gDW/h to all nutrients contained in the minimal medium (Table S2) with additional
carbon sources as specified. Gene identifiers were translated to UniProt similar as for *S. cerevisiae*.

Model preparation

The proposed approach aims at parsimonious correction of turnover values in genome-scale
enzyme-constraint metabolic models using measured protein abundances. Therefore, it is
important to consider differential association between enzymes and reactions, i.e., isozymes,
enzyme complexes, and promiscuous enzymes. We decided to use the GECKO formalism
(Sánchez *et al*, 2017), which deals with these problems elegantly by directly encoding the required
information into the stoichiometric matrix. The genome-scale metabolic models for *S. cerevisiae*
(YeastGEM v.8.5.0) and *E. coli* (iML1515) were obtained from the yeast-GEM and ecModels
GitHub repository, respectively [(Preprint: Domenzain et al, 2021; Lu *et al*, 2019),
https://github.com/SysBioChalmers ; accessed on 22.08.2021]. For subsequent steps, functions of
the COBRA v3.0 toolbox (Heirendt *et al*, 2019) and GECKO2.0 toolbox (Preprint: Domenzain et al,
2021) were employed, of which several functions were adapted for our purposes.

To arrive at raw enzyme-constraint models for both organisms, the GECKO2.0 model enhancement
pipeline was adapted to allow the k_{cat} correction procedure to be omitted. Moreover, any manual
corrections of turnover numbers were excluded from model generation. In the process of adapting
the raw pcGEM to the respective experimental conditions for both organisms, the GAM value per
condition was fitted using the *scaleBioMass* function of GECKO2.0, based solely on the condition-

specific nutrient exchange rates, and returning the minimum $9 \frac{mmol}{gDWh}$) or maximum ($161 \frac{mmol}{gDWh}$)

interval boundary if reached (only *S. cerevisiae*). Furthermore, we omitted enzyme abundances,

which were not measured across all experiments as the approach proposed here only works for

enzymes with measured abundances ($E_{measured}$).

PRESTO approach

In the design of PRESTO, we modified the enzyme mass-balance constraints of the augmented

stoichiometric matrix, created by GECKO, from

421
$$-\frac{1}{k_{cat}^{ij}} v_i + e_i = 0 \tag{1}$$

422 to inequality constraints that use the measured protein abundance directly and further assume a

423 single turnover number per enzyme i over all catalyzed reactions $\left(k_{cat_i}^{min} = \min k_{cat_{i.}}\right)$:

424
$$\sum_{r \in GPR(E_i)} v_r \le k_{cat_i}^{min} \cdot [E_i] . \tag{2}$$

GPR stands for gene-protein-reaction rule that associates reactions with underlying genes and

proteins. We justify making this assumption based on our observation that most enzymes in the *S.*

cerevisiae model are associated with no more than four reactions. Further, the vast majority of

enzymes are assigned a single unique turnover number even though they catalyze multiple

reactions (Fig S16).

We then introduced a correction factor δ, which is added to each k_{cat} if the protein abundances for

the underlying enzyme were available:

432
$$\sum_{r \in GPR(E_i)} v_r \le \left(k_{cat_i}^{min} + \delta_i\right)E_i, \ \forall i \in E_{measured} . \tag{3}$$

433 The value for each δ was constrained by the fold change ε with respect to $k_{cat_i}^{min}$ and a cut-off value

434 K^{max}, which denotes the maximum allowed k_{cat} value.

435 To find a biologically relevant minimal set of adaptations with respect to the sum of δ, we minimized

436 the weighted sum of the average relative error, ω, between measured (μ^{exp}) and predicted specific

437 growth rates (v_{bio}) over all experimental conditions C, and the average δ:

438
$$\min_{v,\delta,\omega} \frac{1}{|C|} \sum_{j \in conditions} |\omega_j| + \frac{\lambda}{|E_{measured}|} \sum_{i \in E_{measured}} \delta_i . \tag{4}$$

439 Finally, the linear programming formulation of the k_{cat} correction in PRESTO is the following:

$$440 \qquad \min_{v,\delta,\omega} \frac{1}{|C|} \sum_{j\in conditions} |\omega_j| + \frac{\lambda}{|E_{measured}|} \sum_{i\in E_{measured}} \delta_i$$

441   subject to

$$442 \qquad \mathbf{Nv_j = 0},\ 1 \le j \le |C| \tag{5}$$

$$443 \qquad \sum_{r\in GPR(E_i)} v_r \le \left(k_{cat,i}^{min} + \delta_i\right)E_{ij},\ \forall i \in E_{measured}$$

$$444 \qquad \mathbf{v_j^{min} \le v_j \le v_j^{max}} \tag{6}$$

$$445 \qquad \delta_i \le (\varepsilon - 1)\cdot k_{cat,i}^{min} \tag{7}$$

$$446 \qquad k_{cat,i}^{min} + \delta_i \le K^{max} \tag{8}$$

$$447 \qquad v_{bio,j}\cdot \omega_j \ge \mu_j^{exp} - v_{bio,j} \tag{9}$$

$$448 \qquad v_{bio,j}\cdot \omega_j \ge v_{bio,j} - \mu_j^{exp} \tag{10}$$

$$449 \qquad \omega \le \theta,\ \delta \ge 0\,.$$

The value for ω was bound from above by a value θ, which was set to 0.6. Further, ε was set to 10^5
since lower values did not yield solutions. The value for K^{max} was set to 57,500,000 s^{-1}(5.3.1.1,
Pyrococcus furiosus (Sharma & Guptasarma, 2015)).

The parameter λ controls the trade-off between both minimization objectives (see Eq. 4). As λ is
unknown and may also be condition- and model-specific, it was fitted using a 3-fold cross-validation
scheme, which was repeated for 10 iterations. To this end, we scanned a log-scale interval between
10^{-14} and 10^{-1}. In each iteration, we performed k_{cat} corrections on two folds of condition-specific
models and validated the obtained corrections on the remaining fold of condition-specific models.
The relative errors (e_r) and the sum of δ (i.e., Δ) were then used to calculate the scores s_λ, which
helped us choose the optimal value for λ:

$$460 \qquad s_\lambda = \frac{1}{10}\sum_{\tau=1}^{10} \frac{e_{r,\lambda,\tau} - e_{r,\lambda,\tau}^{min}}{e_{r,\lambda,\tau}^{max} - e_{r,\lambda,\tau}^{min}} \cdot \frac{\log_{10}\frac{\Delta_{\lambda,\tau}}{\Delta_{\lambda,\tau}^{min}}}{\log_{10}\frac{\Delta_{\lambda,\tau}^{max}}{\Delta_{\lambda,\tau}^{min}}}\cdot \tag{11}$$

More precisely, the optimal value was then determined by finding the first sign change in the second

numerical gradient over s_λ, starting from the maximum value for λ. In addition to the optimal λ, we

also compared our results to a second λ, where the sum of δ reached a plateau ($\lambda = 10^{-10}$ for *S.*

cerevisiae and $\lambda = 10^{-11}$ for *E. coli*, Figs S2a and S10a).

Variability analysis for δ

While PRESTO considers multiple experimental conditions to find a set of universal corrections for

k_{cat} values, it does not provide an exhaustive view over all possible solutions to this problem. To

assess the precision of the corrections, we first performed a variability analysis for δ to find the

minimum and maximum possible values. To guarantee that a solution of equal quality is found with

respect to the previously determined sum of δ and the relative errors to experimentally measured

specific growth rates (i.e., ω^{opt}), corresponding constraints were added to arrive at the following

linear programming problem:

473
$$\min/\max_{v,\delta,\omega} \delta_i$$

474
$$\text{s.t.}$$

475
$$\mathbf{N v_j} = \mathbf{0}, \ 1 \leq j \leq |C|$$

476
$$\sum_{r \in GPR(E_i)} v_r \leq \left(k_{cat,i}^{min} + \delta_i\right) E_{ij}, \ \forall i \in E_{measured}$$

477
$$\mathbf{v_j^{min}} \leq \mathbf{v_j} \leq \mathbf{v_j^{max}}$$

478
$$\delta_i \leq (\varepsilon - 1) \cdot k_{cat,i}^{min}$$

479
$$k_{cat,i}^{min} + \delta_i \leq K^{max}$$

480
$$v_{bio,j} \cdot \omega_j \geq \mu_j^{exp} - v_{bio,j}$$

481
$$v_{bio,j} \cdot \omega_j \geq v_{bio,j} - \mu_j^{exp}$$

482
$$0.99 \cdot \omega_j^{opt} \leq \omega_j \leq 1.01 \cdot \omega^{opt} \tag{12}$$

483
$$\Delta - 10^{-3} \leq \sum_{i \in E_{measured}} \delta_i \leq \Delta + 10^{-3} \tag{13}$$

484
$$\omega \leq \theta, \ \delta \geq 0 \,.$$

485 As the distribution within the obtained min/max intervals can be skewed, we sampled 10,000 points
486 within the obtained intervals. For uniform random sampling, we created random vectors of
487 corrections δ^* within the determined intervals and projected them onto the solution space by
488 minimizing the distance of δ to the respective random vector. Therefore, we updated the objective
489 of the program above:

490
$$\min_{v,\delta,\omega} \sum_{i \in E_{measured}} |\delta_i - \delta_i^*|. \tag{14}$$

To ensure reproducibility and compatibility with the COBRA toolbox (Heirendt *et al*, 2019), we
solved all optimization problems using the *optimizeCbModel* of the COBRA toolbox. Within this
environment, we used the Gurobi solver v9.1.1 (Gurobi Optimization, 2021) but we note that any
other supported solver can also be used. As we observed numerical instability of the problems in
some cases, we decreased the feasibility tolerance (i.e., *feasTol* parameter) for the COBRA solver
to 10^{-9} for all predictions.

Validation of corrected models

We used the adapted GECKO pipeline (fitting a condition specific GAM; excluding manual k_{cat}
adaptions) to obtain models with k_{cat} values adapted according to the objective control coefficient
heuristic. We note that, when no manual modifications were introduced to the *Saccharomyces*
cerevisiae models, the k_{cat} adaption of the GECKO pipeline would stop because no objective
control coefficient above the threshold of 0.001 could be found and corrected models would still be
below the predicted growth error tolerance of 10%. To compare the predictive performance of
PRESTO and GECKO corrected models, the models were adapted with the same condition-
specific GAM, biomass reaction and total protein content, $P_{tot.}$ Additionally, PRESTO models where
constrained using the same condition-specific saturation rate, σ and enzyme mass fraction, f, as
obtained from the GECKO pipeline. In contrast to the GECKO formulation, we did not subtract the
mass of measured enzymes from the total protein pool constraint but instead introduced the
measured protein concentration as upper bound on the enzyme usage reaction, E_i, in the
respective scenario. This formulation still guarantees that the mass of all used enzymes is lower or
equal to the approximated cellular protein pool according to

512
$$\sum_i E_{i,j} \cdot M_i \leq P_{tot,j} \cdot f \cdot \sigma_j, \tag{15}$$

where M is the respective molecular weight of the protein. By considering measured and
unmeasured enzymes in (15) we do not have to change *f* and use the same factor as for the
scenario where no protein abundance measures are used (Sánchez *et al*, 2017). Maximum growth
was predicted in three different constraint scenarios: (i) using only the protein pool constrain and

default uptake rates (1000 mmol/gDW/h), (ii) using the pool constrain and experimentally measured
uptake rates, (iii) using the previous constraints plus the absolute enzyme abundance.

The two studies which generated *in vivo* k_{cat} estimates from pFBA (Davidi *et al*, 2016; Chen &
Nielsen, 2021b)calculated a single value per reaction irrespective of the presence of isoenzymes.
Thus, to parameterize the raw pcGEM (containing only uncorrected BRENDA values) we
substituted the k_{cat} values of all isoenzyme reactions with the same estimate provided in the study.
Reactions catalyzed by complexes were not corrected. Since PRESTO and the pFBA studies
provide a single condition independent model, we generated a condition independent GECKO
model by following the maximum over all conditions approach: For the comparisons in Fig S9 the
condition wise GECKO models were aggregated into a single union model in which for each
reaction the maximum k_{cat} value was used.

Pathway enrichment analysis

The KEGG pathway terms (Kanehisa & Goto, 2000), associated with each enzyme that was
measured in all conditions, were acquired using the KEGG REST API. The one-sided p-value, p,
for significant enrichment of a pathway term among the enzymes with corrected k_{cat} values was
calculated using the hypergeometric density distribution:

$$533 \qquad p(x) = 1 - \sum_{i=1}^{x-1} \frac{\binom{K}{i}\binom{M-K}{N-i}}{\binom{M}{N}} \ . \tag{16}$$

Only KEGG pathway terms associated to at least two corrected enzymes were taken into
consideration. The p-values associated with all tested pathway terms were corrected for a false
discovery rate of 0.05 using the Benjamini-Hochberg correction (Benjamini & Hochberg, 1995).

Funding

P.W. and Z.N. would like to thank the Research Focus Group "Evolutionary Systems Biology" of
University of Potsdam for funding. Z.N., M.A., and Z.R. would like to thank the Max Planck Society
for funding. Z.R. was supported by the European Union's Horizon 2020 research and innovation
programme grant 862201 (to Z.N.) (this publication reflects only the author's view and the
Commission is not responsible for any use that may be made of the information it contains).

Author Contributions: PW, MA performed research and analyzed data, PW contributed code for
PRESTO approach, MA assessed model performance and performed statistical analysis, PW,
MA, ZR, ZN designed research, PW, MA, ZN wrote the paper.

549
Code Availability

All code that was used to generate the results of this study, including the PRESTO method, are available at https://github.com/pwendering/PRESTO.

553
References

Adadi R, Volkmer B, Milo R, Heinemann M & Shlomi T (2012) Prediction of microbial growth rate versus biomass yield by a metabolic network with kinetic parameters. *PLoS Comput Biol* 8: e1002575

Bar-Even A, Noor E, Savir Y, Liebermeister W, Davidi D, Tawfik DS & Milo R (2011) The Moderately Efficient Enzyme: Evolutionary and Physicochemical Trends Shaping Enzyme Parameters. *Biochemistry* 50: 4402–4410

Di Bartolomeo F, Malina C, Campbell K, Mormino M, Fuchs J, Vorontsov E, Gustafsson CM & Nielsen J (2020) Absolute yeast mitochondrial proteome quantification reveals trade-off between biosynthesis and energy generation during diauxic shift. *Proc Natl Acad Sci U S A* 117: 7524–7535

Bateman A (2019) UniProt: A worldwide hub of protein knowledge. *Nucleic Acids Res* 47: D506–D515

Beg QK, Vazquez A, Ernst J, de Menezes MA, Bar-Joseph Z, Barabási A-L & Oltvai ZN (2007) Intracellular crowding defines the mode and sequence of substrate uptake by Escherichia coli and constrains its metabolic activity. *Proc Natl Acad Sci* 104: 12663–12668

Bekiaris PS & Klamt S (2020) Automatic construction of metabolic models with enzyme constraints. *BMC Bioinformatics* 21: 19

Benjamini Y & Hochberg Y (1995) Controlling the False Discovery Rate: A Practical and Powerful Approach to Multiple Testing. *J R Stat Soc Ser B* 57: 289–300

Chen Y & Nielsen J (2021a) Mathematical modeling of proteome constraints within metabolism. *Curr Opin Syst Biol* 25: 50–56

Chen Y & Nielsen J (2021b) In vitro turnover numbers do not reflect in vivo activities of yeast enzymes. *Proc Natl Acad Sci U S A* 118: 2108391118

Davidi D, Longo LM, Jabłońska J, Milo R & Tawfik DS (2018) A Bird's-Eye View of Enzyme Evolution: Chemical, Physicochemical, and Physiological Considerations. *Chem Rev* 118: 8786–8797

Davidi D, Noor E, Liebermeister W, Bar-Even A, Flamholz A, Tummler K, Barenholz U, Goldenfeld M, Shlomi T & Milo R (2016) Global characterization of in vivo enzyme catalytic rates and their correspondence to in vitro kcat measurements. *Proc Natl Acad Sci* 113: 3401–3406

Domenzain I, Sánchez B, Anton M, Kerkhoven EJ, Millán-Oropeza A, Henry C, Siewers V, Morrissey JP, Sonnenschein N & Nielsen J (2021) Reconstruction of a catalogue of genome-scale metabolic models with enzymatic constraints using GECKO 2.0. *bioRxiv doi:*

10.1101/2021.03.05.433259 [Preprint]

van Eunen K & Bakker BM (2014) The importance and challenges of *in vivo*-like enzyme kinetics.
Perspect Sci 1: 126–130

Goelzer A, Muntel J, Chubukov V, Jules M, Prestel E, Nölker R, Mariadassou M, Aymerich S,
Hecker M, Noirot P, *et al* (2015) Quantitative prediction of genome-wide resource allocation
in bacteria. *Metab Eng* 32: 232–243

Gurobi Optimization L (2021) Gurobi Optimizer Reference Manual (https://www.gurobi.com)

Heckmann D, Campeau A, Lloyd CJ, Phaneuf P V., Hefner Y, Carrillo-Terrazas M, Feist AM,
Gonzalez DJ & Palsson BO (2020) Kinetic profiling of metabolic specialists demonstrates
stability and consistency of *in vivo* enzyme turnover numbers. *Proc Natl Acad Sci U S A*
117: 23182–23190

Heckmann D, Lloyd CJ, Mih N, Ha Y, Zielinski DC, Haiman ZB, Desouki AA, Lercher MJ &
Palsson BO (2018) Machine learning applied to enzyme turnover numbers reveals protein
structural correlates and improves metabolic models. *Nat Commun* 9: 1–10

Heirendt L, Arreckx S, Pfau T, Mendoza SN, Richelle A, Heinken A, Haraldsdóttir HS, Wachowiak
J, Keating SM, Vlasov V, *et al* (2019) Creation and analysis of biochemical constraint-based
models using the COBRA Toolbox v.3.0. *Nat Protoc* 14: 639–702

Jeske L, Placzek S, Schomburg I, Chang A & Schomburg D (2019) BRENDA in 2019: a
European ELEXIR core data resource. *Nucleic Acids Res* 47: D542–D549

Kanehisa M & Goto S (2000) KEGG: kyoto encyclopedia of genes and genomes. *Nucleic Acids*
Res 28: 27–30

Küken A, Gennermann K & Nikoloski Z (2020) Characterization of maximal enzyme catalytic
rates in central metabolism of *Arabidopsis thaliana*. *Plant J* 103: 2168–2177

Labhsetwar P, Melo MCR, Cole JA & Luthey-Schulten Z (2017) Population FBA predicts
metabolic phenotypes in yeast. *PLoS Comput Biol* 13: e1005728

Lahtvee PJ, Sánchez BJ, Smialowska A, Kasvandik S, Elsemman IE, Gatto F & Nielsen J (2017)
Absolute Quantification of Protein and mRNA Abundances Demonstrate Variability in Gene-
Specific Translation Efficiency in Yeast. *Cell Syst* 4: 495-504.e5

Lerman JA, Hyduke DR, Latif H, Portnoy VA, Lewis NE, Orth JD, Schrimpe-Rutledge AC, Smith
RD, Adkins JN, Zengler K, *et al* (2012) In silico method for modelling metabolism and gene
product expression at genome scale. *Nat Commun* 3: 929

Li F, Yuan L, Lu H, Li G, Chen Y, Engqvist MKM, Kerkhoven EJ & Nielsen J (2021) Deep learning
based kcat prediction enables improved enzyme constrained model reconstruction. *bioRxiv*
doi: 10.1101/2021.08.06.455417 [Preprint]

Lu H, Li F, Sánchez BJ, Zhu Z, Li G, Domenzain I, Marcišauskas S, Anton PM, Lappa D, Lieven
C, *et al* (2019) A consensus *S. cerevisiae* metabolic model Yeast8 and its ecosystem for
comprehensively probing cellular metabolism. *Nat Commun* 10: 3586

Malina C, Yu R, Bjorkeroth J, Kerkhoven EJ & Nielsen J (2021) Adaptations in metabolism and

protein translation give rise to the Crabtree effect in yeast. *Proc Natl Acad Sci U S A* 118:
e2112836118

Mao Z & Ma H (2019) iMTBGO: An Algorithm for Integrating Metabolic Networks with
Transcriptomes Based on Gene Ontology Analysis. *Curr Genomics* 20: 252–259

Monk JM, Lloyd CJ, Brunk E, Mih N, Sastry A, King Z, Takeuchi R, Nomura W, Zhang Z, Mori H,
et al (2017) iML1515, a knowledgebase that computes *Escherichia coli* traits. *Nat*
Biotechnol 35: 904–908

Nilsson A, Nielsen J & Palsson BO (2017) Metabolic Models of Protein Allocation Call for the
Kinetome. *Cell Syst* 5: 538–541

O'Brien EJ, Lerman JA, Chang RL, Hyduke DR & Palsson BØ (2013) Genome-scale models of
metabolism and gene expression extend and refine growth phenotype prediction. *Mol Syst*
Biol 9: 693

Peebo K, Valgepea K, Maser A, Nahku R, Adamberg K & Vilu R (2015) Proteome reallocation in
Escherichia coli with increasing specific growth rate. *Mol Biosyst* 11: 1184–1193

Sánchez BJ, Zhang C, Nilsson A, Lahtvee P, Kerkhoven EJ & Nielsen J (2017) Improving the
phenotype predictions of a yeast genome-scale metabolic model by incorporating enzymatic
constraints. *Mol Syst Biol* 13: 935

Schmidt A, Kochanowski K, Vedelaar S, Ahrné E, Volkmer B, Callipo L, Knoops K, Bauer M,
Aebersold R & Heinemann M (2016) The quantitative and condition-dependent *Escherichia*
coli proteome. *Nat Biotechnol* 34: 104–110

Sharma P & Guptasarma P (2015) 'Super-perfect' enzymes: Structural stabilities and activities of
recombinant triose phosphate isomerases from *Pyrococcus furiosus* and *Thermococcus*
onnurineus produced in *Escherichia coli*. *Biochem Biophys Res Commun* 460: 753–758

Valgepea K, Adamberg K, Seiman A & Vilu R (2013) *Escherichia coli* achieves faster growth by
increasing catalytic and translation rates of proteins. *Mol Biosyst* 9: 2344–2358

Wendering P & Nikoloski Z (2022) Genome-Scale Modeling Specifies the Metabolic Capabilities
of *Rhizophagus irregularis*. *mSystems* 7: e01216-21

Xu R, Razaghi-Moghadam Z & Nikoloski Z (2021) Maximization of non-idle enzymes improves
the coverage of the estimated maximal in vivo enzyme catalytic rates in Escherichia coli.
Bioinformatics 37: 3848–3855

Yu R, Campbell K, Pereira R, Björkeroth J, Qi Q, Vorontsov E, Sihlbom C & Nielsen J (2020)
Nitrogen limitation reveals large reserves in metabolic and translational capacities of yeast.
Nat Commun 11: 1881

Yu R, Vorontsov E, Sihlbom C & Nielsen J (2021) Quantifying absolute gene expression profiles
reveals distinct regulation of central carbon metabolism genes in yeast. *Elife* 10: e65722

Zikmanis P & Kampenusa I (2012) Relationships between kinetic constants and the amino acid
composition of enzymes from the yeast *Saccharomyces cerevisiae* glycolysis pathway.
Eurasip J Bioinforma Syst Biol 2012
665

Fig 1. Schematic overview of the PRESTO approach for k_{cat} correction. The approach uses a

GECKO-formatted pcGEM (Sánchez *et al*, 2017) for an organism containing turnover numbers

from BRENDA (Jeske *et al*, 2019). Using available data from n experimental conditions, n

condition-specific models are generated using nutrient uptake rates and protein contents. PRESTO

then uses data on abundances for the enzymes measured across the n investigated conditions

and solves a linear program that minimizes a weighted sum of two objectives, the relative error to

measured specific growth rates and the sum of positive k_{cat} corrections, δ. The optimal weighting

factor, λ, which modulates the trade-off between the two objectives, is then determined by cross-

validation (Tr: training set; Ts: test set) , choosing the parameter, which is associated with the

lowest average relative error. Using the optimal value for λ, PRESTO combines all models for the

experimental conditions to find a k_{cat} correction for each enzyme with measured abundance. Last,

the precision of δ values are assessed by variability analysis as well as by sampling and corrected

k_{cat} values are validated by comparing them to values obtained from other approaches.

Fig 2. Comparison of predicted growth of *S. cerevisiae* from pcGEMs with k_{cat} corrections

from GECKO and PRESTO. Condition-specific pcGEMs with corrected k_{cat} values generated in

GECKO were used to predict the specific growth rate in each condition. The boxplots indicate the

distribution of the relative error resulting from each set of condition-specific corrected k_{cat} values

obtained from the GECKO heuristic. Relative prediction error from each set is indicated by a circle.

The red rhombus shows the relative error of predicted specific growth rate from the PRESTO model

($\lambda = 10^{-7}$) by using the single set of corrected k_{cat} values in the respective pcGEM. **(a)** Only the

measured total protein pool was used to constrain the solution and condition-specific uptake rates

were bounded by 1000 $\frac{mmol}{h\ gDW}$; **(b)** measured uptake rates were also considered; **(c)** abundances of

enzymes measured in all conditions were used as additional constraints. The compared pcGEMs

in each condition used the same respective biomass coefficients, GAM, σ, and $Ptot$ values

(Methods). L: Lahtvee et al. 2017, D: Di Bartolomeo et al. 2020, Y20: Yu et al. 2020, Y21: Yu et al.

2021.

Fig 3. Comparison of enzymes with corrected k_{cat} values by both GECKO and PRESTO. (a) KEGG Pathway terms significantly enriched in the set of enzymes corrected by PRESTO ($\lambda = 10^{-7}$) in the *S. cerevisiae* model. The x-axis gives the number of corrected enzymes linked to the given term. The one-sided p-value was calculated using the hypergeometric density distribution and was corrected using the Benjamini-Hochberg procedure (Benjamini & Hochberg, 1995). **(b)** Venn diagram showing the overlap of enzymes whose k_{cat} values were manually corrected (Sánchez *et al*, 2017) ("Manual"), automatically corrected by the GECKO heuristic in any of the conditions ("GECKO"), or corrected by PRESTO ("PRESTO"). **(c)** Log-transformed k_{cat} values corrected using both the GECKO heuristic and PRESTO are not associated (Spearman correlation coefficient of 0.186, p-value = 0.385).

Fig 4. Comparison of predicted growth of *E. coli* from pcGEMs with k_{cat} corrections from GECKO and PRESTO. Condition-specific pcGEMs with corrected k_{cat} values generated in GECKO were used to predict the specific growth rate in each condition. Grey dots mark the relative error resulting from each set of condition-specific corrected k_{cat} values obtained from the GECKO heuristic. Relative prediction error from each set is indicated by a circle. The red rhombus shows the relative error of predicted specific growth rate from the PRESTO model ($\lambda = 10^{-5}$) by using the single set of corrected k_{cat} values in the respective pcGEM. **(a)** Only the measured total protein pool was used to constrain the solution and condition-specific uptake rates were bounded by 1000 $\frac{mmol}{h\,gDW}$; **(b)** abundances of enzymes measured in all conditions were used as additional constraints. Missing data points originate from infeasibility of the respective models. The compared pcGEMs in each condition used the same respective biomass coefficients, GAM, $\sigma,$ and P_{tot} values (Methods). P: Peebo et al. 2015 , V: Valgepea et al. 2013, S: Schmidt et al. 2015.