Title: CCR4 and CCR7 differentially regulate thymocyte localization with distinct outcomes for central

tolerance

Authors: Yu Li[1†], Pablo Guaman Tipan[1†], Hilary J. Selden[1], Jayashree Srinivasan[1], Laura P. Hale[2], Lauren

I.R. Ehrlich[1,3 *]

[1] Department of Molecular Biosciences, The University of Texas at Austin, Austin, TX 78712, USA

[2] Department of Pathology, Duke University School of Medicine, Durham, NC 27710

[3] Department of Oncology, LIVESTRONG Cancer Institutes, The University of Texas at Austin Dell Medical

School, Austin, TX 78712, USA

[†] These authors contributed equally

[*] Correspondence: lehrlich@austin.utexas.edu (L.I.R. Ehrlich)

Abstract

Central tolerance ensures autoreactive T cells are eliminated or diverted to the regulatory T cell lineage,

thus preventing autoimmunity. To undergo central tolerance, thymocytes must enter the medulla to

test their TCRs for autoreactivity against the diverse self-antigens displayed by antigen presenting cells

(APCs). While CCR7 is known to promote thymocyte medullary entry and negative selection, our

previous studies implicate CCR4 in these processes, raising the question of whether CCR4 and CCR7 play

distinct or redundant roles in central tolerance. Here, synchronized positive selection assays, 2-photon

timelapse microscopy, and quantification of TCR-signaled apoptotic thymocytes, demonstrate that CCR4

and CCR7 promote medullary accumulation and central tolerance of distinct post-positive selection

thymocyte subsets. CCR4 is upregulated within hours of positive selection signaling and promotes

medullary entry and clonal deletion of immature post-positive selection thymocytes. In contrast, CCR7 is

expressed several days later and is required for medullary localization and negative selection of mature

thymocytes. In addition, CCR4 and CCR7 differentially enforce self-tolerance, with CCR4 enforcing

tolerance to self-antigens presented by activated APCs, which express CCR4 ligands. Our findings show

that CCR7 expression is not synonymous with medullary localization and support a revised model of

central tolerance in which CCR4 and CCR7 promote early and late stages of negative selection,

respectively, via interactions with distinct APC subsets.

Keywords: Thymocytes, Chemokine receptors, Negative selection, Central tolerance, *Ccr7*, *Ccr4*

Introduction

Self-tolerance of the T-cell repertoire is established in thymus through the process of central tolerance,

which encompasses both negative selection and regulatory T cell (Treg) induction (Klein et al., 2014). To

avoid autoimmunity, developing T cells must be broadly tolerized not only to ubiquitously expressed

self-antigens, but also to proteins expressed by distinct differentiated cell types. The thymic medulla is a

specialized environment in which central tolerance to such diverse self-antigens is enforced. After

positive selection in the cortex, thymocytes migrate into the medulla, where they interact with

medullary antigen presenting cells (APCs), including thymic dendritic cells (DCs) and medullary thymic

epithelial cells (mTECs). Collectively, these APCs enforce self-tolerance by presenting self-peptides from

the majority of the proteome on MHC complexes, such that thymocytes expressing autoreactive T cell

receptors undergo clonal deletion or diversion to the Treg lineage (Ehrlich, 2016; Klein et al., 2014; Klein

et al., 2019; Lancaster et al., 2019). AIRE[+] mTECs express >80% of the proteome, including *Aire-*

dependent tissue restricted antigens (TRAs) that are otherwise expressed in only a few peripheral

tissues (Abramson & Anderson, 2017; Anderson & Su, 2016; Mathis & Benoist, 2007; Meredith et al.,

2015). DCs also present numerous self-antigens, including those acquired from mTECs, from circulation,

or from peripheral tissues and trafficked into the thymus (Atibalentja et al., 2009; Bonasio et al., 2006;

Koble & Kyewski, 2009). The importance of inducing thymocyte tolerance to medullary self-antigens is

evidenced by multi-organ autoimmunity that ensues in *Aire*-deficient mice and APECED patients

(Anderson et al., 2002; Finnish-German, 1997; Nagamine et al., 1997), in whom thymic expression of

TRAs is greatly diminished. Failure to express even a single medullary TRA can result in impaired

tolerance and subsequent T-cell mediated autoimmune pathology (DeVoss et al., 2006). Notably,

individual TRAs are expressed by only 1-3% of *Aire*[+] mTECs (Brennecke et al., 2015; Meredith et al.,

2015; Sansom et al., 2014), resulting in a sparse mosaic of self-antigen display throughout the medulla.

Because thymocytes reside in the medulla for only 4-5 days (McCaughtry et al., 2007), tight

spatiotemporal regulation is required to ensure that post-positive selection thymocytes encounter the

full spectrum of medullary self-antigens required to enforce self-tolerance.

To facilitate medullary entry after positive selection, thymocytes upregulate chemokine receptors that

promote their directional migration towards medullary-biased chemokine gradients (Bleul & Boehm,

2000; Hu, Lancaster, & Ehrlich, 2015; Lancaster et al., 2018; Petrie & Zuniga-Pflucker, 2007). Notably,

CCR7 has been shown to play a critical role in directing chemotaxis of post-positive selection thymocytes

towards the medulla, where CCR7 ligands are expressed, enhancing thymocyte accumulation within the

medulla, enforcing negative selection to TRAs, and averting autoimmunity (Ehrlich et al., 2009; Kozai et

al., 2017; Kurobe et al., 2006; Nitta et al., 2009; Ueno et al., 2004). Given the significance of CCR7 in

promoting thymocyte medullary entry, CCR7 expression is widely considered to be synonymous with

thymocyte medullary localization. Based largely on this definition, recent studies indicate that despite

the well-established role of the medulla in inducing central tolerance, ~75% of negative selection occurs

in CCR7$^-$ 'cortical' thymocytes, while only 25% occurs in CCR7$^+$ 'medullary' cells (Breed et al., 2019; Daley

et al., 2013; Hu et al., 2016). It has also been suggested that cortical negative selection may eliminate

thymocytes reactive to ubiquitous self-antigens while medullary deletion tolerizes TRA-responsive cells.

Our previous research implicates chemokine receptors other than CCR7 in promoting thymocyte

medullary localization (Ehrlich et al., 2009). We found that CCR4, which is upregulated by post-positive

selection thymocytes, also promotes medullary entry and negative selection (Hu, Lancaster,

Sasiponganan, et al., 2015). These findings raise the question of whether CCR4 and CCR7 play distinct or

redundant roles in promoting thymocyte medullary entry and central tolerance.

In this paper, we use a combination of approaches, including chemotaxis assays, 2-photon live-cell

microscopy, and synchronized positive selection assays, to distinguish the contributions of CCR4 versus

CCR7 to thymocyte medullary entry and negative selection. We find that CCR4 is upregulated by

thymocytes as early as a few hours after positive selection, while CCR7 is expressed days later by mature

post-positive selection cells. CCR4 expression promotes chemotaxis and medullary entry of early post-

positive selection thymocytes, which do not yet express CCR7. Initial CCR7 expression results in only

moderate thymocyte chemotaxis towards CCR7 ligands and modest accumulation in the medulla, such

that these thymocytes migrate in both the cortex and medulla. These findings indicate that CCR7

expression is not a definitive marker of medullary versus cortical localization. CCR7 is, however, required

for robust accumulation of mature (CD4SP) thymocytes in the medulla. Notably, consistent with their

differential activity in distinct thymocyte subsets, CCR4 and CCR7 are required in early and late phases of

polyclonal negative selection, respectively. While CCR7 is known to promote tolerance to TRAs (Kozai et

al., 2017; Nitta et al., 2009), we present evidence that CCR4 promotes tolerance to self-antigens

presented by activated APCs. Collectively, our study establishes non-redundant roles for CCR4 and CCR7

in governing localization of post-positive selection thymocyte subsets and central tolerance.

Results

CCR4 is expressed by immature post-positive selection thymocyte subsets, while CCR7 expression is

restricted to more mature thymocyte subsets

We first investigated expression of CCR4 and CCR7 by distinct thymocyte subsets using cell surface

markers that delineate developmental stages after positive selection (Figure 1A) (Sinclair et al., 2013;

Xing et al., 2016). To validate the developmental trajectory of these thymocyte subsets, we analyzed

their GFP expression levels in Rag2p-GFP mice, in which GFP expression is driven by the *Rag2* promoter.

In this model, expression of GFP distinguishes newly generated thymocytes (GFP$^+$) from recirculating T

cells (GFP$^-$), and declining GFP expression reflects time elapsed after positive selection, when the *Rag2*

promoter becomes inactive (Boursalian et al., 2004). In pre-positive selection CD4$^+$CD8$^+$ double-positive

(DP) CD3$^-$CD69$^-$ thymocytes, the *Rag2* promoter is active, resulting in maximal GFP expression (Figure

1B). After receiving a TCR signal, thymocytes upregulate CD3 and CD69 (Fu et al., 2009), generating early

post-positive selection DP CD3^{lo}CD69$^+$ cells; the brief time elapsed after positive selection is indicated by

continued high GFP expression by this subset (Figure 1B). To further evaluate if DP CD3^{lo}CD69$^+$

thymocytes represent post-positive selection thymocytes, as opposed to cells undergoing strong TCR

signaling driving clonal deletion, we evaluated thymocyte subsets from *Nur77^{GFP}* mice, in which GFP

levels reflect the strength of TCR signaling (Figure 1—figure supplement 1) (Moran et al., 2011).

Elevation of GFP expression by DP CD3^{lo}CD69$^+$ cells, relative to DP CD3$^-$CD69$^-$ cells, confirmed this subset

was TCR-signaled. Thymocytes rescued from negative selection express Nur77^{GFP} at levels comparable to

thymic Tregs, which express self-reactive TCRs (Stritesky et al., 2013). GFP levels were substantially

lower in DP CD3^{lo}CD69$^+$ cells than in CD25$^+$ CD4SPs, which are mostly Tregs (Figure 1—figure supplement

1A). Together, these data are consistent with DP CD3^{lo}CD69$^+$ cells representing mainly post-positive

selection DPs, and not strongly self-reactive thymocytes undergoing negative selection. Sequential

maturation through by CD4$^+$single-positive semi-mature cells (CD4SP SM) and then CD4SP or CD8SP

mature 1 (M1) and mature 2 (M2) subsets was confirmed by progressive reductions in GFP levels for

both the CD4 and CD8 lineages (Figure 1B) (Xing et al., 2016). A distinct population of DP CD3$^+$CD69$^+$

cells expressed lower levels of GFP than CD4SP SM cells (Figure 1B), placing it temporally between

CD4SP SM and CD8SP M1 cells (Figure 1B). Thus, we considered whether DP CD3$^+$CD69$^+$ cells represent

MHCI-restricted thymocytes in the process of differentiating into CD8SP cells through co-receptor

reversal downstream of the CD4SP SM stage (Brugnera et al., 2000). Consistent with this possibility, DP

CD3$^+$CD69$^+$ cells were significantly reduced in β2m$^{-/-}$ mice, in which positive selection of CD8SP

thymocytes is abrogated, as is selection of innate iNKT cells and CD8αα+ IELps (Bendelac et al., 1994;

Ruscher et al., 2017). Furthermore, DP CD3$^+$CD69$^+$ cells were enriched in MHCII$^{-/-}$ mice in which only

MHCI-restricted thymocytes are positively selected (Figure 1—figure supplement 1B-C)). Together, these

data demonstrate that DP CD3$^+$CD69$^+$ cells are MHC-I restricted and have persisted for an intermediate

time between CD4SP SM and CD8SP M1 cells after positive selection, consistent with a co-receptor

reversing DP subset.

Having delineated the temporal sequence of thymocyte development post-positive selection, we

quantified expression of CCR4 and CCR7 by each subset. CCR4 is upregulated by the majority of early

post-positive selection DP CD3^{lo}CD69$^+$ thymocytes, which do not yet express CCR7 (Figure 1C-D, Figure

1—figure supplement 2A-B). CCR7 is subsequently expressed at high levels by ~40% CD4SP SM cells, and

CCR4 expression persists on this subset (Figure 1C-D). In comparison to *Ccr7*[-/-] cells, low-level expression

of CCR7 can be detected on additional *Ccr7*[+/+] CD4SP SM cells, consistent with initial upregulation of

CCR7 by this subset (Figure 1—figure supplement 2A-B). Expression of CCR4 and CCR7 by MHCI-

restricted DP CD3[+]CD69[+] cells is similar to that of CD4SP SM cells, with slightly more CCR7 and less CCR4

(Figure 1C-D, Figure 1—figure supplement 2A-B). As thymocytes mature through CD4SP and CD8SP M1

and M2 stages, CCR4 is progressively downregulated. CCR7 expression peaks in both CD4SP and CD8SP

M1 subsets and is diminished in M2 cells (Figure 1C-D, Figure 1—figure supplement 2A-B). Because cell-

surface expression of CCR7 on mature CD4[+] T cells can be downregulated due to ligand-induced

internalization of the chemokine receptor (Britschgi et al., 2008), we tested whether receptor

internalization could have obscured detection of CCR7 or CCR4 on ex vivo thymocyte subsets.

Thymocytes were incubated at 37°C for up to 2 hours, to allow re-expression of internalized receptors,

before immunostaining with anti-CCR7 and anti-CCR4 antibodies. Increased expression of cell-surface

CCR7 or CCR4 was not detected with increasing incubation time (Figure 1—figure supplement 2C-D).

These findings confirm that CCR4 expression is an early indicator of positive selection at the DP stage,

while CCR7 expression is upregulated later by more mature CD4SP and CD8SP subsets.

Early post-positive selection thymocytes undergo chemotaxis towards CCR4 ligands, while more

mature subsets respond to CCR7 ligands

We next determined whether expression of CCR4 and CCR7 corresponds to thymocyte chemotactic

responses to their respective ligands. DP CD3[lo]CD69[+] cells, which express CCR4 but not CCR7, underwent

chemotaxis in response to the CCR4 ligand CCL22, but not the CCR7 ligands CCL19 or CCL21 (Figure 2A-

D, Figure 2—figure supplement 1A). CCL22 induced chemotaxis of all subsequent post-positive selection

thymocytes, except the most mature M2 thymocyte subsets (Figure 2B, Figure 2—figure supplement

1A), largely in keeping with CCR4 expression patterns. Although CD4SP SM and DP CD3+CD69+ subsets

expressed CCR7 (Figure 1C), CCR7-mediated chemotaxis of these subsets did not reach statistical

significance when compared to all thymocyte subsets (Figure 2C-D). However, when considered

separately, moderate CCR7-mediated chemotaxis of these subsets was revealed, although DP

CD3^{lo}CD69$^+$ cells still did not undergo CCR7-mediated chemotaxis (Figure 2—figure supplement 1A).

Consistent with expression of CCR7 by more mature thymocyte subsets, the CCR7 ligands CCL19 and

CCL21 induced highly efficient chemotaxis of mature CD4SP and CD8SP M1 and M2 subsets (Figure 2C-

D).

A recent study indicated that expression of the chemokine receptor CXCR4 must be extinguished before

post-positive selection thymocytes can leave the cortex, where CXCL12 is expressed by cortical thymic

epithelial cells, to migrate into the medulla (Kadakia et al., 2019). Thus, we sought to place CXCR4

expression and function into context with CCR4 and CCR7. CXCR4 expression decreased after positive

selection, but all post-positive selection subsets continued to express CXCR4 (Figure 3A-B), and all

except for CD4SP SM and DP CD3$^+$CD69$^+$ subsets underwent chemotaxis in response to CXCL12 (Figure

3C). Although the earliest post-positive selection thymocyte subsets exhibited reduced CXCR4

responsiveness, perhaps allowing them to exit the cortex, the ability to respond to CXCR4 signals does

not preclude medullary localization, as evidenced by CXCR4-mediated chemotaxis of medullary CD4SP

and CD8SP M2 cells. Taken together, chemotaxis assays show that early post-positive selection DP

CD3^{lo}CD69$^+$ thymocytes are responsive to CCR4 but not CCR7 ligands, with low responsiveness to CXCR4

ligands. At the next stage, CD4SP SM cells also undergo chemotaxis towards CCR4 ligands, and begin to

migrate at a low level towards CCR7 ligands, but not to CXCR4 ligands. As thymocytes mature to the

CD4SP and CD8SP M1 and M2 stages, they progressively gain the ability to respond robustly to CCR7

ligands, regain CXCR4 responsiveness, and lose responsiveness to CCR4 ligands (Figure 3D).

Initial expression of CCR7 by CD4SP SM thymocytes does not result in strong chemotaxis to CCR7 ligands

(Figure 2D), and expression of CXCR4 by this subset does not enable detectable chemotaxis towards

CXCL12 (Figure 3C), highlighting a temporal disconnect between chemokine receptor expression and

function on maturing post-positive selection thymocytes. We considered the possibility that CD4SP SM

cells could be intrinsically unable to respond to chemokine receptor signaling. However, these cells

responded efficiently to CCR4 ligands (Figure 2B), and they underwent chemotaxis to the CCR9 ligand

CCL25 at levels comparable to DP CD3^{lo}CD69$^+$ cells (Figure 2—figure supplement 2A), consistent with a

previous study showing CCL25 induces migration of most thymocyte subsets (Campbell et al., 1999).

Thus, regulation of thymocyte responsiveness to chemokine receptors is more complex than altered

cell-surface expression levels.

CCL21a establishes a chemokine gradient that drives accumulation of CD4SP thymocytes in the medulla

(Kozai et al., 2017; Ueno et al., 2002; Ueno et al., 2004). Because we found that functional CCR4 is

expressed by early post-positive selection DP thymocytes, we examined whether CCL22 could establish a

similar chemokine gradient to recruit these CCR4-responsive cells into the medulla.

Immunofluorescence analysis revealed that CCL22 is expressed predominantly in the medulla (Figure

2—figure supplement 2B), consistent with a potential role in inducing medullary entry of early post-

positive selection thymocytes.

The timing of CCR4 upregulation following positive selection correlates with thymocyte medullary

entry

Given that the majority of early post-positive selection DP thymocytes respond to CCR4 but not CCR7

ligands, we sought to determine how rapidly CCR4 versus CCR7 are upregulated following initiation of

positive selection and to assess whether expression of either receptor correlates with the timing of

medullary entry. To accomplish these goals, we used a synchronized positive selection thymic slice

assay, coupled with flow cytometry or 2-photon microscopy (Figure 4A). OT-II$^+$ *Rag2$^{-/-}$ MHCII$^{-/-}$* mice

served as a source of pre-positive selection DP thymocytes that did not yet express CCR4 or CCR7 due to

lack of positive selection in the absence of MHCII expression (Figure 4B). OT-II$^+$ *Rag2$^{-/-}$ MHCII$^{-/-}$*

thymocytes were fluorescently labeled prior to culturing on MHC-II-sufficient, positively selecting (wild-

type (WT or pCX-EGFP) or non-positively selecting ($MHCII^{-/-}$) live thymic slices. At various time points,

expression of CCR4 and CCR7 by OT-II thymocytes in the thymic slices was assayed by flow cytometry, or

medullary entry was quantified by 2-photon live-cell microscopy (Figure 4A).

OT-II$^+$ $Rag2^{-/-}$ $MHCII^{-/-}$ DP cells upregulated CCR4 as early as 3 hours (3h) after being introduced onto

positively selecting thymic slices (Figure 4C). The frequency of CCR4$^+$ cells continued to increase over 24

hours on WT slices. Positive selection was required for initiation of CCR4 expression, as demonstrated by

the failure to upregulate CCR4 on non-selecting $MHCII^{-/-}$ slices. CCR7 was not upregulated within 24h of

initiating positive selection (Figure 4C). When labeled OT-II$^+$ $Rag2^{-/-}$ $MHCII^{-/-}$ thymocytes were introduced

into positively selecting pCX-EGFP thymic slices and imaged by 2-photon microscopy, they were initially

localized to the cortex, consistent with our previous findings that pre-positive selection thymocytes

cannot access the medulla (Video S1) (Ehrlich et al., 2009). By 9-12 hours after positive selection, OT-II$^+$

thymocytes entered the medulla, as evidenced by a significant increase in medullary density (Figure 4D-

E; Video S2). As CCR7 is not yet expressed by these cells (Figure 4C), CCR4 likely drives medullary entry

of early post-positive selection thymocytes.

To test if the timing of CCR4 and CCR7 upregulation and medullary entry are comparable for MHCI-

restricted thymocytes, we carried out similar experiments using pre-positive selection OT-I$^+$ $Rag2^{-/-}$

thymocytes that matured in the non-selecting thymuses of $\beta 2m^{-/-}$ hosts. Although OT-I positive selection

was largely inhibited in these bone marrow chimeras, about 10% of the DP cells expressed CCR4 at

baseline (Figure 4—figure supplement 1A-B). Nonetheless, CCR4 was upregulated as early as 3h after

initiating positive selection in WT thymic slices, but not in non-selecting $\beta 2m^{-/-}$ slices and increased over

24 hours (Figure 4—figure supplement 1A-B). CCR7 was not expressed during the first 24h after

initiation of positive selection and only became detectable as CD8SP cells differentiated at the 72h

timepoint (Figure 4—figure supplement 1 B-C), consistent with prior observations (Lutes et al., 2021).

Furthermore, 2-photon imaging of pre-selection OT-I$^+$ $Rag2^{-/-}$ thymocytes within the cortex and medulla

of thymic slices showed that despite medullary entry of some thymocytes at 1.5h (Video S3), likely due

to the observed baseline CCR4 expression in this model, the medullary density of OT-I thymocytes

increased significantly by 12h after the initiation of positive selection (Figure 4—figure supplement 1 D-

E; Video S4).

Together, these findings show that both MHCI- and MHCII-restricted thymocytes upregulate CCR4

within a few hours of the initiation of positive selection and accumulate significantly in the medulla by 9-

12h. Thymocyte medullary entry precedes CCR7 upregulation, which does not occur until 48h-72h after

positive selection. The kinetics of CCR4 upregulation are consistent with a role for CCR4 in promoting

early medullary entry of positively selected thymocytes. However, CCR7 is known to be required for

efficient medullary entry of SP thymocytes (Ehrlich et al., 2009; Kurobe et al., 2006), raising the

possibility that these two chemokine receptors promote medullary localization of different thymocyte

subsets.

CCR4 and CCR7 are required for medullary accumulation of distinct post-positive selection thymocyte

subsets

To test our hypothesis that CCR4 and CCR7 are required for medullary accumulation of distinct post-

positive selection thymocytes, we FACS purified thymocyte subsets from WT, $Ccr4^{-/-}$, $Ccr7^{-/-}$ and $Ccr4^{-/-}$

$Ccr7^{-/-}$ (DKO) mice, labeled them with red or blue fluorescent dyes, and allowed them to migrate in pCX-

EGFP live thymic slices. 2-photon time-lapse microscopy was used to image the sorted cells, so we could

determine their migratory properties and densities in the medulla versus cortex (Figure 5—figure

supplement 1A). EGFP is expressed ubiquitously in pCX-EGFP mice, such that the cortex and medulla of

thymic slices can be distinguished by differences in stromal cell morphology and EGFP intensity (Figure

5A) (Ehrlich et al., 2009).

Purified WT DP CD3^{lo}CD69$^+$ cells entered and migrated within the medulla, accumulating at a 2-fold

higher density than in the cortex (Figure 5A-B; Video S5). As expected, the majority of the sorted early

post-positive selection cells expressed CCR4, but not CCR7 (Figure 5—figure supplement 1B-C).

Medullary enrichment of this subset was abolished when the cells were purified from *Ccr4$^{-/-}$* or DKO

mice, but *Ccr7* deficiency did not diminish their medullary accumulation (Figure 5B). Paired analyses of

WT versus *Ccr4$^{-/-}$* and *Ccr7$^{-/-}$* versus DKO DP CD3^{lo}CD69$^+$ cells imaged together in the same slices

confirmed that *Ccr4* deficiency significantly impaired medullary accumulation of this early post-positive

selection DP subset irrespective of *Ccr7* genotype, but genetic deficiency of *Ccr7* at this stage did not

alter medullary accumulation (Figure 5C). Altogether, CCR4 is required for the 2-fold accumulation of DP

CD3^{lo}CD69$^+$ cells in the medulla relative to the cortex.

Despite the fact that roughly half of the purified CD4SP SM cells expressed CCR7 (Figure 5—figure

supplement 1B-C), which is considered to be a hallmark of thymocyte medullary entry, accumulation of

these cells within the medulla did not further increase, but instead remained at a 2-fold medullary:

cortical density ratio, comparable to that of DPCD3^{lo}CD69$^+$ cells (Figure 5A-B; Video S6). Medullary

accumulation of this subset was significantly reduced by CCR4 deficiency, regardless of the *Ccr7*

genotype (Figure 5B-C), indicating an important role for CCR4 in promoting medullary localization of

CD4SP SM cells. In addition, paired analysis of *Ccr7$^{-/-}$* versus WT CD4SP SM cells imaged together in the

same slices showed that CCR7 also contributes to medullary accumulation of CD4SP SM cells (Figure 5C).

Together, these data indicate that CCR4 and CCR7 cooperate to promote medullary accumulation of

CD4SP SM thymocytes.

CD4SP M1+M2 thymocytes, the majority of which express only CCR7 (Figure 5—figure supplement 1C),

accumulated in the medulla to a greater extent than either of the previous subsets, with a ~4-5-fold

medullary:cortical density ratio (Figure 5B-C; Video S7, S8), consistent with the high migration index of

these cells to CCR7 ligands (Figure 2C-D). CCR4 deficiency did not significantly impact medullary

accumulation of this subset (Figure 5B-C), consistent with the lack of responsiveness of M2 cells to CCR4

ligands (Figure 2A). However, the trend towards decreased medullary enrichment of *Ccr4* deficient cells

may reflect the activity of CCR4 in CD4SP M1 cells (Figure 2A). Medullary accumulation of M1+M2

CD4SP thymocytes was highly dependent on CCR7, as demonstrated by the significant decrease in

medullary accumulation of both *Ccr7*$^{-/-}$ and DKO cells relative to WT or *Ccr4*$^{-/-}$ cells (Figure 5B, Video S7,

S8).

We also considered whether CCR4 and CCR7 alter thymocyte migratory properties. Deficiency in *Ccr4*

and/or *Ccr7* did not impact the speed of DP CD3^{lo}CD69^{+} or CD4SP SM cells. However, CCR7 deficiency

resulted in a significant decline in the speed of CD4SP M1+M2 cells, consistent with our previous finding

that CCR7 promotes rapid motility of CD4SP thymocytes (Ehrlich et al., 2009). Interestingly, double

deficiency of *Ccr4* and *Ccr7* resulted in a significant increase in the average path straightness of all

imaged thymocyte subsets (Figure 5E), suggesting that thymocytes responding to CCR4 and CCR7 ligands

migrate in a more tortuous path.

Overall, CCR4 and CCR7 were required for medullary accumulation of distinct post-positive selection

thymocyte subsets, largely consistent with their expression patterns and chemotactic function (Figure

3D). Notably, our results show that CCR4 directs early post-positive selection thymocytes into the

medulla, demonstrating that lack of CCR7 expression by post-positive selection thymocytes does not

preclude medullary localization. Conversely, expression of CCR7 by CD4SP SM cells does not result in

enhanced medullary accumulation over the previous subset. Thus, CCR7 expression on its own is not

indicative of robust medullary accumulation. Interestingly, only mature CD4SP M1+M2 cells accumulate

robustly in the medulla and migrate rapidly in a CCR7-dependent manner.

CCR4 and CCR7 contribute to early versus late phases of negative selection, respectively

Given the differential impact of CCR4 versus CCR7 on migration and medullary accumulation of post-

positive selection thymocyte subsets, we hypothesized that these chemokine receptors are required for

negative selection of early versus late post-positive selection cells, respectively. Consistent with our

prior observations (Hu, Lancaster, Sasiponganan, et al., 2015), the frequencies of post-positive selection

thymocyte subsets were comparable between littermate control WT and $Ccr4^{-/-}$ mice (Figure 6—figure

supplement 1A). CD5 expression levels, which are a proxy for self-reactivity (Azzam et al., 1998; Hawiger

et al., 2004; Persaud et al., 2014), were also comparable in the absence of $Ccr4$ (Figure 6—figure

supplement 1). To test the impact of CCR4 and CCR7 on polyclonal negative selection, we quantified the

frequency of cleaved-caspase 3$^+$ thymocytes that had undergone TCR signaling in WT, $Ccr4^{-/-,}$ $Ccr7^{-/-}$, and

DKO mice. Intracellular cleaved-caspase 3 in TCR-signaled thymocytes has been shown to mainly reflect

negative selection (Breed et al., 2019; Hu et al., 2016), although other apoptotic stimuli, such as

glucocorticoid hormones, could also induce cleaved-caspase 3 in thymocytes (Alam et al., 1997;

Marchetti et al., 2003). Previous studies showed that negative selection occurs in 2 main phases,

generally defined to occur at the CCR7$^-$ DP stage in the cortex versus the CCR7$^+$ CD4SP stage in the

medulla (Breed et al., 2019; Daley et al., 2013; Hu et al., 2016; Stritesky et al., 2013). Thus, we

quantified the impact of $Ccr4$ versus $Ccr7$ deficiency on "early-phase" and "late-phase" negative

selection (Figure 6A-B). Interestingly, the majority of early-phase DP cells undergoing negative selection

expressed only CCR4, as did ~60% of late-phase CD4SP cells; thus, the majority of thymocytes

undergoing clonal deletion express CCR4, but not CCR7. CCR7 was expressed by ~40% CD4SP and 60% of

CD8SP cells undergoing negative selection (Figure 6B). The frequency of total thymocytes undergoing

negative selection declined significantly only in DKO mice (Figure 6C). However, CCR4 deficiency resulted

in a significant decrease in the frequency of thymocytes undergoing early-phase, but not late-phase

negative selection (Figure 6C). In contrast, CCR7 deficiency resulted in a significantly lower frequency of

thymocytes undergoing late-phase negative selection, with no impact on early-phase negative selection.

Double-deficiency for CCR4 and CCR7 significantly diminished both early-phase and late-phase negative

selection (Figure 6C). It was reported that most negative selection occurs at the earlier phase (Breed et

al., 2019; Stritesky et al., 2013), which is consistent with the 2:1 ratio of WT thymocytes undergoing

early- to late-phase negative selection (Figure 6D). This ratio falls significantly in $Ccr4^{-/-}$ mice and

increases significantly in $Ccr7^{-/-}$ and DKO mice, highlighting the respective contributions of CCR4 and

CCR7 to early versus late phases of negative selection, respectively (Figure 6D). Altogether, these data

indicate that, consistent with its role in promoting medullary entry of early-post positive selection

thymocytes, CCR4 is required for efficient early-phase negative selection of polyclonal DP cells. In

contrast, CCR7, which promotes medullary accumulation of more mature CD4SP and CD8SP cells, is

required for efficient negative selection of late-phase CD4SP and CD8SP cells.

CCR4 and CCR7 suppress autoinflammation in distinct tissues, with CCR4 promoting tolerance to

activated antigen-presenting cells

Given that CCR4 and CCR7 enforced central tolerance at distinct stages of T-cell development when

thymocytes might interact with different APCs, these chemokine receptors could promote tolerance to

distinct self-antigens. To evaluate this possibility, we assessed the presence of spontaneous

autoimmunity/autoinflammation in multiple organs of WT, $Ccr4^{-/-}$, $Ccr7^{-/-}$ and $Ccr4^{-/-}Ccr7^{-/-}$ mice. We

previously found that $Ccr4^{-/-}$ and $Ccr7^{-/-}$ mice had anti-nuclear autoantibodies by ~12-months of age (Hu

et al., 20115). Here we detected anti-nuclear autoantibodies in the serum of 67% of $Ccr4^{-/-}$,100% of

$Ccr7^{-/-}$, and 75% of $Ccr4^{-/-}Ccr7^{-/-}$ mice between 5 to 6.5 months of age, relative to 22% of WT mice (Figure

7—figure supplement 1A). Thus, we evaluated other organs from mice in this age range for lymphocytic

infiltrates or inflammation. Colon inflammation was not observed in $Ccr7^{-/-}$ or WT mice. Colons from

Ccr4[-/-] mice showed mild epithelial hyperplasia, with small foci of acute inflammation. In contrast, Ccr4[-/-]

Ccr7[-/-] mice demonstrated moderate to severe colonic mucosal hyperplasia, with abundant acute and

chronic inflammation extending into the submucosa. Active ulcerations were observed in 40% of *Ccr4*[-/-]

Ccr7[-/-] mice, with crypt abscesses and regenerative changes consistent with prior ulceration in most

Ccr4[-/-] *Ccr7*[-/-] mice (Figure 7A, left). Histologic scores were markedly and significantly higher in *Ccr4*[-/-]

Ccr7[-/-] mice compared to WT, with a trend toward higher scores in *Ccr4*[-/-] mice that is absent in *Ccr7*[-/-]

mice (Figure 7A, right), suggesting that CCR4 and CCR7 collaboratively maintain self-tolerance in the

colon (Figure 7A). Immune infiltrates were elevated in the liver, lacrimal glands, and submandibular

glands of *Ccr7*[-/-] and *Ccr4*[-/-]*Ccr7*[-/-] mice, but not *Ccr4*[-/-] mice at 6-months of age (Figure 7B, Figure 7- figure

supplement 1B-D), indicating that CCR7 plays a dominant role in promoting tolerance to endocrine

organs, consistent with the established requirement for CCR7 to mediate negative selection against

mTEC-expressed TRAs, but not ubiquitous self-antigens (Nitta et al., 2009). Interestingly, 70-75% of *Ccr4*[-/-]

mice, compared to 17-20% of WT and 0% of *Ccr7*[-/-] mice displayed moderate lymphoid hyperplasia in

the spleen and mesenteric lymph nodes of 12-month old mice (Figure 7C, right). Immunofluorescent

staining of lymph nodes from 6-month old *Ccr4*[-/-] mice revealed abnormal CD4 and CD8

distributions/expansions in T-cell zones relative to WT mice (Figure 7C, left). As expected, T cell zones

from *Ccr7*[-/-] mice were hypoplastic (not shown), reflecting the role of CCR7 in mediating T-cell entry into

LNs. These results indicate that CCR4 maintains T cell homeostasis in secondary lymphoid organs.

Given the distinct contributions of CCR4 and CCR7 to establishment of central tolerance and

maintenance of self-tolerance across different organs, we considered the possibility that CCR4 and CCR7

might promote thymocyte selection against antigens presented by different types of thymic APCs.

Thymic APC subsets were FACS purified (Figure 7—figure supplement 2A), and expression of CCR4 and

CCR7 ligands was quantified by qRT-PCR. Consistent with our previous study (Hu, Lancaster,

Sasiponganan, et al., 2015), the CCR4 ligands CCL17 and CCL22 were expressed by Sirpα[+] cDC2 cells.

Activated thymic CCR7[+] cDCs (Ardouin et al., 2016; Hu et al., 2017; Oh et al., 2018) also expressed CCR4

ligands, with the highest levels of CCL22 expressed by activated DCs (Figure 7D). Thymic B cells also

expressed CCL22, consistent with a recent report that CCL22 expressed by activated B cells promotes

interactions with T cells in the germinal center (Liu et al., 2021). The CCR7 ligands CCL19 and CCL21 were

expressed by mTEC subsets, with the highest levels of CCL21 expression by mTEC[lo] cells (Figure 7E),

consistent with previous reports (Bornstein et al., 2018; Kozai et al., 2017). These results suggest that

CCR4 may promote thymocyte interactions with B cells and cDCs, including activated CCR7[+] cDCs that

express elevated levels of MHCII and co-stimulatory molecules (Ardouin et al., 2016; Hu et al., 2017),

while CCR7 may promote thymocyte interactions primarily with mTECs.

Because a large proportion of thymic B cells (Cepeda et al., 2018) and cDCs (Ardouin et al., 2016; Hu et

al., 2017) have activated phenotypes, and T-cell zones in the lymph nodes of *Ccr4[-/-]* mice are

hyperplastic (Figure 7C), we considered the possibility that CCR4-directed negative selection could

facilitate establishment of central tolerance to antigens expressed by activated APCs. To explore this

hypothesis, CD45.1[+] congenic splenocytes were stimulated overnight with toll-like receptor (TLR)

ligands. We confirmed that stimulation with TLR ligands induced activation of the splenic cDCs and B

cells based on upregulation of MHCII, CD80, and CD86 (Figure 7—figure supplement 2B). The activated

splenocytes were then co-cultured with CD45.2[+] CD4[+] conventional T cells isolated from WT or *Ccr4[-/-]*

spleens, and the frequency of T cells induced to proliferate was quantified after 3-5 days (Figure 7F).

Notably, LPS-stimulated splenocytes induced more proliferation of *Ccr4[-/-]* versus WT CD4[+] T cells at days

3 and 5 (Figure 7G). Splenocytes that were unstimulated or activated by other TLR ligands did not

preferentially induce *Ccr4[-/-]* T cell proliferation (Figure 7E, Figure 7—figure supplement 2C). Taken

together, these data suggest that CCR4 promotes thymocyte central tolerance to self-antigens

expressed by LPS-activated APCs, possibly by promoting interactions with such APCs in the thymus.

Discussion

Our data demonstrate that CCR4 and CCR7 play distinct roles in thymocyte localization and the

induction of T-cell central tolerance. We find that CCR4 is upregulated by thymocytes within a few hours

of the initiation of positive selection, largely coincident with the timing of medullary entry. CCR4 is

expressed by almost all early post-positive selection DP CD3^{lo}CD69^{+} cells and is required for their 2-fold

accumulation in the medulla and efficient negative selection. Notably, the majority of thymocytes

undergoing negative selection are DP CD3^{lo}CD69^{+} cells, which do not express CCR7. Thus, while the role

of CCR4 in negative selection has been questioned (Cowan et al., 2014), our analyses reveal an

important role for this chemokine receptor in localization and central tolerance of early post-positive

selection polyclonal thymocytes, in keeping with medullary localization of the CCR4 ligand CCL22. In

contrast, our study and a previous report (Lutes et al., 2021) show that CCR7 is not upregulated until 48-

72 hours after positive selection. CCR7 is expressed by about half of the CD4SP SM cells, which continue

to express CCR4 and respond to CCR4 ligands. In keeping with dual expression, CCR4 and CCR7

cooperate to induce the 2-fold accumulation of CD4SP SM cells in the medulla. As thymocytes mature

further to the CD4SP and CD8SP M1 and M2 stages, CCR7 is upregulated, CCR4 is downregulated, and

the cells become increasingly chemotactic to CCR7 ligands. We find that CCR7 is required for the robust

accumulation of CD4SP M1+M2 cells in the medulla and for late-phase negative selection of CD4SP and

CD8SP thymocytes. The finding that the overall frequency of negative selection in the thymus is reduced

significantly only in the absence of both CCR4 and CCR7 further indicates that these chemokine

receptors both contribute to the induction of central tolerance. Altogether, our findings show a

temporally correlated shift after positive selection from a requirement for CCR4 to a requirement for

CCR7 for thymocyte medullary localization and negative selection.

Several of our findings raise questions about the current model of negative selection, which assumes

that CCR7 expression is a surrogate indicator of thymocyte medullary versus cortical localization (Breed

et al., 2019; Hu et al., 2016). This assumption is based on studies, including our own, showing that CCR7

is expressed by CD4SP and CD8SP thymocytes, which are localized mainly within the thymic medulla at

steady-state, as evidenced by immunostaining (Ehrlich, 2016). In addition, prior studies showed that

accumulation of CD4SP cells in the medulla is CCR7-dependent (Ehrlich et al., 2009; Ueno et al., 2004);

consistent with these findings, we confirm that robust medullary accumulation of the more mature

CD4SP M1 and M2 cells is CCR7-dependent. Equating CCR7 expression with medullary localization has

led to a model in which the early phase of negative selection is thought to impact CCR7$^-$ DP cells in the

cortex, while the later phase impacts more mature CCR7$^+$ cells in the medulla (Breed et al., 2019; Hu et

al., 2016). In support of cortical negative selection, apoptotic TCR transgenic cells were located in the

cortex of the H-Y^{CD4} model of negative selection (McCaughtry et al., 2008). Also, TCR-signaled

thymocytes are present at the cortical side of the CMJ in *Bim*$^{-/-}$ mice, in which apoptosis following strong

TCR signaling is impaired (Stritesky et al., 2013), suggesting thymocytes encounter negatively selecting

self-antigens in the cortex. Furthermore, CCR7 deficiency impairs negative selection to TRAs, but not to

ubiquitous self-antigens (Kurobe et al., 2006; Nitta et al., 2009), indicating that CCR7$^+$ mature

thymocytes undergo negative selection in the medulla, where TRAs are expressed by mTECs.

While our data confirm that CCR7 is critical for medullary accumulation of more mature thymocyte

subsets, we find that CCR7 expression is not a reliable indicator of medullary localization in earlier post-

positive selection thymocytes. CCR7$^-$ DP CD3^{lo}CD69$^+$ enter the medulla, where they accumulate in a

CCR4-dependent manner with a 2-fold higher density than in the cortex, so CCR7 is not required for

medullary entry. Conversely, although almost half of CD4SP SM cells express CCR7, their medullary:

cortical density is also 2-fold. CCR7 deficiency only modestly diminished medullary accumulation of

CD4SP SM cells, consistent with their moderate chemotaxis towards CCR7 ligands; instead, CCR4 played

a greater role in medullary accumulation of this subset. Thus, the two earliest post-positive selection

subsets accumulate in the medulla at similar densities, despite CCR7 upregulation by one of these

subsets. DP CD3^{lo}CD69$^+$ and CD4SP SM cells both migrate in the cortex and in the medulla and could

thus encounter APCs presenting cognate self-antigens that drive negative selection in either region.

Based on the 2-fold enrichment of DP CD3^{lo}CD69^{+} cells in the medulla, a substantial proportion of CCR7^{-}

early phase negative selection may occur in the medulla. Moreover, some late phase negative selection,

defined by either CCR7 expression or maturation to the CD4SP SM stage, may occur in the cortex. These

data indicate that CCR7 expression cannot be used as a reliable surrogate biomarker of medullary

localization. The location in which thymocytes undergo negative selection impacts which APC subsets

will be encountered, altering the spectrum of self-antigens to which thymocytes are tolerized. Thus, it

will be important to determine where distinct thymocyte subsets undergo negative selection and which

APCs promote tolerance of these different subsets.

It has recently been suggested that CXCR4 expression must be extinguished after positive selection to

release thymocytes from binding to cTECs in the cortex, thus enabling medullary entry (Kadakia et al.,

2019). However, we found that while CXCR4 is partially downregulated on the cell surface of post-

positive selection thymocytes, all subsets except CD4SP SM and DP CD3^{+}CD69^{+} cells undergo chemotaxis

to CXCL12, despite their accumulation in the medulla. These findings indicate that CXCR4 expression and

activity are compatible with medullary localization. This seeming discrepancy may reflect the

overexpression system used to reveal that persistent CXCR4 expression prevented thymocyte medullary

entry, perhaps resulting in superphysiological tethering of cells to the cortex (Kadakia et al., 2019). In

keeping with the ability of CXCR4^{+} thymocytes to enter the medulla, higher CXCR4 surface expression

and responsiveness are observed in mature CD4SP and CD8SP M1 and M2 subsets, which accumulate in

the medulla to the greatest extent. It is notable that chemotactic activity of CXCR4 is completely

extinguished at the CD4SP SM stage, despite continued CXCR4 expression, but not at the more mature

CD4SP and CD8SP stages, suggesting a thymocyte-intrinsic mechanism of diminished responsiveness to

CXCR4, similar to the lack of responsiveness to CCR7 ligands by this subset. It is possible that the lack of

CXCR4 activity in CD4SP SM cells may permit medullary entry, consistent with the need to release post-

positive selection thymocytes from binding to cTECs (Kadakia et al., 2019); however, DP CD3[lo]CD69[+] cells

enter the medulla just as efficiently as CD4SP SM cells despite continued CXCR4 responsiveness,

suggesting factors other than reduced CXCR4 activity likely enable medullary entry. The role for

increased CXCR4 activity in more mature SP thymocytes remains to be determined but could impact

thymocyte egress. Identifying intrinsic regulators of chemokine receptor responsiveness in different

thymocyte subsets will further our understanding of the complex mechanisms underlying the

orchestration of thymocyte movement and selection in the thymus.

A major outstanding question raised by our findings is whether the combinatorial and temporally

regulated changes in chemokine receptor expression on post-positive selection thymocytes drive

interactions with distinct APC subsets. Here we show that CCR7 ligands are expressed by mTECs, while

CCR4 ligands are expressed mainly by cDC2s and activated cDCs, with some expression by thymic B cells.

Thus, CCR7 activity may drive interactions with mTECs, possibly explaining why CCR7 is particularly

required for negative selection to TRAs, consistent with the finding that CCR7 deficiency results in

autoimmune infiltrates in endocrine organs, as occurs in Aire-deficient mice (Anderson et al., 2002;

Kurobe et al., 2006; Nitta et al., 2009). In contrast, CCR4 activity may promote interactions with and

tolerance to cDC2s, B cells, and activated cDCs, consistent with our finding that *Ccr4*[-/-] mice showed T

cell hyperplasia in secondary lymphoid organs and that *Ccr4*[-/-] T cells undergo increased proliferation to

LPS-activated splenic APCs. Recently, a subset of cDC2s has been shown to present circulating antigens

to induce negative selection, and another subset has been shown to traffic microbial antigens to the

thymus, impacting thymocyte selection (Atibalentja et al., 2009; Vollmann et al., 2021; Zegarra-Ruiz et

al., 2021). B cells also present self-antigens to induce thymocyte negative selection (Perera et al., 2016;

Yamano et al., 2015). While we find CCL22 is expressed predominantly in the medulla, and CCR4

promotes medullary accumulation of early- post positive selection DPs, our study does not rule out the

possibility that CCR4 could also enforce interactions between CCR4-responsive thymocytes and cortical

DCs expressing CCL22, which could contribute to cortical negative selection. This possibility is

particularly intriguing in light of the fact that DP CD3^{lo}CD69^{+} and CD4SP SM cells respond to CCR4 ligands

and migrate both in the cortex and medulla. The finding that CCR4 and CCR7 cooperate to prevent

inflammation in the colon is unexpected and warrants further exploration. Whether the inflammatory

phenotypes observed reflect impaired central tolerance due to diminished interactions between

thymocytes and distinct thymic APCs and/or impaired peripheral tolerance remains to be determined.

Altogether, our findings suggest a layered process of central-tolerance induction in which CCR4 first

promotes interactions with DCs and B cells, driving self-tolerance to activated APCs, which may be

present in both the cortex and/or medulla. As thymocytes mature further, CCR7 is expressed and

becomes active, drawing the cells robustly into the medulla, where the partially tolerized repertoire can

be focused on tolerance induction to the sparse TRAs. One limitation of our current system is that ex

vivo live thymic slices are not equivalent to in situ thymi, as they lack circulation, for example. We note,

however, that short-term live thymic slice cultures have been widely used to investigate the

development, localization, migration, and positive and negative selection of thymocytes, as they have

been shown to faithfully reflect these in vivo processes, including confirming that CCR7 signaling induces

chemotaxis of mature thymocytes from the cortex into the medulla (Au-Yeung et al., 2014; Dzhagalov et

al., 2013; Ehrlich et al., 2009; Lancaster et al., 2019; Melichar et al., 2013; Ross et al., 2014). Because our

study measured differences in medullary accumulation of thymocytes that differed genetically only in

expression of CCR4 and/or CCR7, we attribute altered localization to the impact of these chemokine

receptors on thymocyte migration. However, we note the possibility that CCR4 and CCR7 could signal in

conjunction with survival cues, like cytokines, localized to different thymic microenvironments, which

could also impact survival of thymocytes in different regions. As thymocytes were imaged within a few

hours of entering slices, differential survival in the cortex and medulla due to cytokine cues is somewhat

less likely than differential migration. Multiple aspects of our revised model of thymocyte migration and

central tolerance, including the impact of CCR4 and CCR7 on interactions with distinct APCs during

tolerance induction and on selection of distinct TCR clones will be tested in future studies.

Materials and Methods

Mice

C57BL/6J (wild-type), B6.SJL-Ptprc^a PepC^b (CD45.1), C57BL/6-Tg(TcraTcrb)1100Mjb/J (OT-I), B6.Cg-

Tg(TcraTcrb)425Cbn/J (OT-II), B6(Cg)-Rag2$^{tm1.1Cgn}$/J ($Rag2^{-/-}$), B6.129P2(C)Ccr7^{tm1Rfo}r/J ($Ccr7^{-/-}$), B6.129S2-

H2$^{dlAb1-Ea}$/J ($MHCII^{-/-}$), B6.129P2-B2m^{tm1Unc}/DcrJ ($β2m^{-/-}$), C57BL/6-Tg(Nr4a1-EGFP/cre)820Khog/J

(Nur77^{GFP}) were purchased from the Jackson Laboratory. $Ccr4^{-/-}$ (Chvatchko et al., 2000), pCX-EGFP

(Wright et al., 2001), and Rag2p-GFP (Boursalian et al., 2004) strains were generously provided by A.D.

Luster (Massachusetts General Hospital, Boston, MA), Irving L. Weissman (Stanford University, Stanford,

CA), and Ellen R. Richie (the University of Texas MD Anderson Center, Houston, Texas, United States),

respectively. $Ccr4^{-/-}$ $Ccr7^{-/-}$, OT-I $Rag2^{-/-}$, and OT-II $Rag2^{-/-}$ $MHCII^{-/-}$ strains were bred in house. Experiments

were performed using mice 4-8 weeks old, except for for autoimmune/inflammatory studies which were

carried out in older mice, as specified. All strains were bred and maintained under specific pathogen-

free conditions at the Animal Resources Center, the University of Texas at Austin, with procedure

approval from the Institutional Animal Care and Use Committee, the University of Texas at Austin.

Flow cytometry

$5×10^6$ thymocytes were stained with fluorescently conjugated antibodies and a viability dye, and

incubated on ice for 30 minutes in the dark. Unless specified, for stains including anti-CCR7, cells were

instead incubated in a 37°C water bath for 45 minutes in the dark. Stained cells were washed and

resuspended in FACS Wash Buffer (PBS + 2% FBS (GemCell™, Gemini, CA, USA)) + 1μg/mL propidium

iodine (PI) if a fixable viability dye was not used. For cleaved caspase 3 stains, surface-stained cells were

fixed and permeabilized with the BD Cytofix/Cytoperm kit (BD, NJ, USA) according to manufacturer's

instructions prior to staining for anti-cleaved caspase 3 (Cell Signaling Technology, MA, USA). All flow

cytometry data were acquired on an LSR Fortessa flow cytometer (BD) and analyzed with FlowJo

ver.10.8.0 (BD).

Transwell Chemotaxis assays

Thymocyte chemotaxis assays were performed as previously described (Campbell et al., 1999). Briefly,

5×10^5 thymocytes were resuspended in 100µL RPMI 1640 (Gibco, MA, USA) + 10% Fetal Bovine Serum

(Gemini) and added to the top chamber of 5µm pore transwell tissue culture inserts (Corning, NY, USA)

in 24-well plates. The bottom chamber of the transwells contained recombinant mouse chemokines

CCL17, CCL22, CCL19, CCL21 (Peprotech, NJ, USA) or CCL25 (R&D Systems, MN, USA) at 100nM, 10nM

and 1nM, diluted in 500µL RPMI 1640 + 10% FBS. The plate was cultured at 37°C, 5% CO_2 for 2 hours.

Cells that migrated to the bottom wells, and 5×10^5 input cells, were analyzed and quantified by flow

cytometry. A standard number of 15µm polyester beads were added to each well to calculate the

absolute number of thymocytes that migrated. Migration percentages of each subset were calculated by

dividing the number of migrated cells of each subset by the number of cells of the same subset in the

input cell sample, and the migration index was calculated as the ratio of the migration percentage in

wells containing chemokine to the average migration percentage in wells without chemokine (Blank

wells).

Ex vivo **thymic slice preparation**

Ex vivo thymic slices were prepared as previous described (Lancaster & Ehrlich, 2017). Briefly, surgically

removed thymi were cleaned of residual connective tissue, and the two lobes were separated and

embedded in 4% low melting point agarose (Lonza, NJ, USA). 400µm thymic slices were generated by

slicing the trimmed agarose blocks on a VT 1000S vibratome (Leica, Germany) and kept on ice,

submerged in complete RPMI until addition of thymocytes. Slices were placed on 0.4µm cell culture

inserts (Millipore, MA, USA) in 35mm dishes containing 1mL complete RPMI (RPMI 1640 supplemented

with 1X GlutaMAX, 1mM Sodium Pyruvate, 1X Penicillin-Streptomycin-glutamine, 1X MEM Non-Essential

Amino Acid, and 50µM β-mercaptoethanol (All from Gibco), and 10% FBS (Gemini)).

Synchronized positive selection thymic slice assays

Synchronized positive selection assays were set up as previously described (Ross et al., 2014). Briefly, to

generate a source of pre-positive selection OT-I thymocytes, bone marrow from OT-I *Rag2$^{-/-}$* hosts was

magnetically depleted of CD3$^+$, CD8$^+$, CD25$^+$, B220$^+$, Gr-1$^+$, CD11b$^+$, Ter119$^+$ cells using Rat anti-mouse

antibodies (BioXcell, NH, USA) and Dynabeads™ sheep anti-rat IgG beads (Invitrogen, MA, USA)

according to manufacturer's recommendations, then 10^7 cells were injected into lethally irradiated *β2m$^-$*

$^{/-}$ hosts. The bone marrow chimera recipients were maintained for 4-6 weeks before collecting OT-I pre-

selection thymocytes. For pre-positive selection OT-II thymocytes, OT-II *Rag2$^{-/-}$ MHCII$^{-/-}$* mice were bred.

Thymic single cell suspensions from the OT-I *Rag2$^{-/-}$* -> *β2m$^{-/-}$* bone marrow chimeras or OT-II *Rag2$^{-/-}$*

MHCII$^{-/-}$ mice were isolated, stained in 5mL RPMI 1640 + 5µM CellTrace™ Violet (Invitrogen) for 30

minutes in the dark at 37°C, washed once with warm complete RPMI, resuspended in fresh, warm

complete RPMI, and kept in the dark at 37°C prior to slice addition. Prepared cells were overlaid on WT,

β2m$^{-/-}$, *MHCII$^{-/-}$* or pCX-EGFP slices and incubated at 5% CO_2 37°C. Slices were gently washed at 3h with

warm complete RPMI in a dish to prevent further entry of new cells. At the indicated time points,

thymocytes were harvested by mechanical disruption of the slices for flow cytometry or intact slices

were imaged by 2-photon microscopy.

2-photon imaging and analysis of thymic slices

For imaging of purified thymocyte subsets, sorted cells from each host were centrifuged for collection,

then separately stained in RPMI 1640 with 5µM CellTracker™ Red CMTPX (Thermo Fisher Scientific, MA,

USA) or Indo-1 AM (eBioscience, CA, USA) for 30 minutes in 37°C water bath, away from light. Dye used

for each host was swapped between experiments to account for potential effect of dyes on cell viability

and mobility. For imaging of pre-positive selection thymocytes, whole thymus single cells were stained

in RPMI 1640 with 5µM CellTracker™ Red CMTPX in the same way. Stained cells were collected by

centrifugation and resuspended in fresh warm cRPMI, incubated for 30 minutes in 37°C water bath,

away from light. Prepared cells were collected by centrifugation, then washed 2 times with fresh warm

complete RPMI to remove dye residues, then laid onto pCX-EGFP slices, and incubated in 5% CO_2

incubator at 37°C for a minimum of 1 hour before 2-photon imaging.

For image and Video acquisition, thymic slices were secured to an imaging chamber (Harvard Apparatus,

MA, USA) perfused with heated DRPMI (Corning) + 2g/L sodium bicarbonate, 5mM HEPES (Sigma-

Aldrich, MA, USA) and 1.25mM $CaCl_2$, pH = 7.4, aerated with 95% O_2 + 5% CO_2 at flow rate of 100mL/h.

The imaging chamber is secured on a heated stage, and a temperature probe was inserted to monitor

and maintain the chamber temperature at 37°C. Images were acquired using an Ultima IV microscope

controlled by PrairieView software (V5.4, Bruker, MA, USA), with a 20X water immersion objective, NA =

1.0 (Olympus, Japan). Time-lapse Videos were acquired by a t-series of 60 rounds of 15s acquisition,

through 40µm depth at 5µm intervals. Samples were illuminated with a Mai Tai Ti:sapphire laser

(Spectra Physics) tuned to 865nm for simultaneous excitation of EGFP and CMTPX, and an additional

InSight Ti:sapphire laser (Spectra Physics, CA, USA) tuned to 730nm for excitation of Indo-1 AM, if

necessary. Emitted light was passed through 473/24, 525/50 and 605/70 band pass filters (Chroma, VT,

USA) to separate PMTs for detecting Indo-1 AM (low calcium), EGFP and CMTPX signals, respectively.

Captured images were analyzed using Imaris v9.7.2 (Bitplane, Switzerland). For medullary enrichment

analysis, medullary and cortical volumes were established by manually generating surface objects by

morphological distinction. Fluorescently labelled cells were identified using Spot tools from random

frames of each Video, and medullary or cortical cellular localization was determined by calculating the

spot's distance to the medullary surface object using the Spot distance to Surface function in ImarisXT.

Densities of cells within each region were calculated by dividing the number of cells in the respective

region by the volume of the surface object, and medullary-to-cortical ratio was calculated by dividing

the medullary density to the cortical density within the same Video, using Excel (Microsoft). For

thymocyte migration parameters, trajectory of thymocytes were established with the automated Spot

tracking feature within Imaris. Only trajectories with time span ≥ 3 minutes were included in the

analysis. Average track speed and straightness were generated by Imaris for analysis.

cDNA preparation and qPCR

FACS purified, frozen thymic APC subsets were thawed and lysed in TRIzol (Thermo Fisher), RNA was

extracted, and cDNA waw synthesized using the qScript cDNA Synthesis Kit (QuantaBio, MA, USA), per

manufacture's recommendations. qRT-PCR was performed using SYBR Green PCR Master Mix (Applied

Biosystems, MA, USA), on a ViiA 7 Real-Time PCR System (Applied Biosystems), using the following

primers: *Ccl17* Forward: 5'-AGT GGA GTG TTC CAG GGA TG-3', *Ccl17* Reverse: 5'-CCA ATC TGA TGG CCT

TCT TC-3', *Ccl19* Forward: 5'-GCT AAT GAT GCG GAA GAC TG-3', *Ccl19* Reverse: 5'-ACT CAC ATC GAC TCT

CTA GG-3', *Ccl21* Forward: 5'-GCA GTG ATG GAG GGG GTC AG-3', *Ccl21* Reverse: 5'-CGG GGT GAG AAC

AGG ATT GC-3', *Ccl22* Forward: 5'-AGG TCC CTA TGG TGC CAA TGT-3', *Ccl22* Reverse: 5'- CGG CAG GAT

TTT GAG GTC CA-3', β-actin Forward: 5'- CAC TGT CGA GTC GCG TCC A-3', β-actin Reverse: 5'- CAT CCA

TGG CGA ACT GGT GG-3'. ddCT Relative expression levels of target genes in each subset were quantified

by first normalizing to actin expression within each subset, then normalizing between subsets for

expression of target genes relative to a subset previously shown to express high levels of that gene. For

CCL17 and CCL22, expression levels were normalized to cDC2; For CCL19 and CCL21, expression levels

were normalized to mTEC^{lo}.

Congenic TLR stimulation and co-culture experiment

To activate APCs, congenic CD45.1 mouse spleen was isolated and red blood cells were lysed with RBC

lysis buffer (BioLegend, CA, USA). Leukocytes were then resuspended in complete RPMI with 1µg/mL

high molecular weight Poly(I:C), 1µg/mL low molecular weight Poly(I:C), 100ng/mL LPS, 500nM

ODN1826 (Invivogen, CA, USA), or without any TLR ligands, and incubated in 5% CO_2 incubator at 37°C

overnight. The next day, CD45.2 WT and $Ccr4^{-/-}$ CD4$^+$ T cells were isolated from spleen by magnetically

depleting CD8$^+$, CD25$^+$, B220$^+$, CD11b$^+$, Gr-1$^+$ and Ter119$^+$ cells with corresponding rat anti-mouse

antibodies (BioXCell) and Dynabeads® sheep anti-rat IgG (Invitrogen), and resuspended in complete

RPMI. Stimulated CD45.1 splenocytes were washed three times with fresh complete RPMI. $2.5×10^4$ WT

or $Ccr4^{-/-}$ CD4$^+$ purified T cells were plated with or without $1.25×10^5$ unstimulated or stimulated

splenocytes per well in complete RPMI in a 96-well, U-bottom plate. At Day 3 and Day 5, each well was

harvested and subjected to flow cytometry analysis. 15µm polyester beads of known quantities were

added to quantify cells in each sample.

Immunofluorescent analyses of thymic and LNs cryosections and detection of autoantibodies

To detect anti-nuclear autoantibodies in mouse serum, 7-µm cryosections were prepared from $Rag2^{-/-}$

kidneys (C57BL/6J background). Cryosections were fixed in acetone (-20°C for 20 minutes), rinsed 3X (2x

PBS rinses and 1x PBS with 0.1% Tween 20), blocked with 9% donkey serum (Jackson ImmunoResearch)

in PBS for 20 minutes and incubated with undiluted mouse serum from mice of the indicated genotypes

for 2 hours at room temperature. After washing, auto-antibodies were detected by staining with anti-

mouse IgG AF594 in PBS for 1 hour at room temperature. After washing, slides were incubated with

DAPI, washed, and coverslips were mounted in ProLong™ Gold Antifade reagent (ThermoFisher). For

CCL22 immunofluorescent staining and inguinal lymph node immunostaining, 7-µm cryosections were

prepared from the thymus of 1 month old C57BL/6J mice and from lymph nodes of 5–6-month-old mice,

respectively. All slides were fixed and stained as above. Thymus sections were stained with anti-CCL22

AF439, anti-CD11c-biotin and anti-CD31-APC overnight at room temperature. Inguinal lymph nodes

were stained with anti-CCL22, anti-CD11c -biotin, and anti-CD31 - APC overnight at room temperature.

For both, after washing, anti-goat-Dylight 488 and streptavidin Alexa Fluor 594 secondary antibodies

were incubated for 1 hour at room temperature. The DAPI nuclear stain was added and slides mounted

as above. Immunofluorescent images were acquired on a DMi8 microscope (Leica), using a 10x/0.4 NA

objective or a 20X/0.7NA objective. Stitched images were generated with LasX software (Leica). All

images were uniformly processed and converted into Tiffs using Fiji software (ImageJ),

H&E staining and quantification

Lacrimal glands, submandibular glands, colon, and liver were extracted from 6–8-month-old mice and

fixed in formalin (Fisherbrand) for 48 hours before storing in 70% ethanol. Organs were sent to the

Histology and Immunohistochemistry Laboratory at the University of Texas San Antonio for embedding,

sectioning, and staining. Spleen and mLNs were extracted and processed by the Histology and Tissue

Processing core at the University of Texas MD Anderson Cancer Center Science Park (Smithville, TX).

Histological analysis of spleen, inguinal lymph nodes, liver and colon were done by a veterinary

pathologist, while infiltrates in lacrimal and submandibular glands stains were analyzed in-house, as in

previously described (Lieberman et al., 2015). Briefly, we scored foci composed of at least 50

mononuclear cells. In the cases where multiple foci coalesced, we assigned those foci a score value of 3

for statistical analysis. The number of inflammatory foci per 10 mm^2 was calculated by counting the

total number of foci by standard light microscopy using a 10 × objective and dividing that by the surface

area of sections measured by Fiji software. Colon sections were scored for mucosal changes such as

hyperplasia and architectural distortion, inflammation severity, and the percentage of the tissue

affected as described previously (Hale et al., 2005), but with a single score that reflected the entire

segment examined (total possible scores ranged from 0 – 15).

Statistical analysis

All statistical analysis was conducted using Graphpad PRISM v9.8 (Graphpad Software, CA, USA), with

the corresponding statistical tests and multiple comparison corrections listed in the figure legends.

Acknowledgements: The authors thank the Animal Resource Center staff at the University of Texas at

Austin for assistance with mouse maintenance, Dr. Jessica Lancaster for assistance with 2-photon

microscopy training, Richard Salinas at the Center for Biomedical Research Support, and Dr. Ellen Richie

for providing advice. Graphical illustrations were created with Biorender.com. This research was

supported by a grant from the National Institutes of Health R01AI104870 to L.I.R.E.

Author contributions: Y.L. and P.G.T. designed and conducted experiments, analyzed data, and wrote

and edited the manuscript. J.S. and H.J.S. conducted experiments. L.P.H carried out histologic analyses

and edited the manuscript. L.I.R.E. designed experiments, analyzed data, and edited the manuscript.

Competing Interests statement: The authors have no conflicts of interest to declare.

References

Abramson, J., & Anderson, G. (2017). Thymic Epithelial Cells. *Annu Rev Immunol*, *35*(1), 85-118.
https://doi.org/10.1146/annurev-immunol-051116-052320
Alam, A., Braun, M. Y., Hartgers, F., Lesage, S., Cohen, L., Hugo, P., Denis, F., & Sékaly, R.-P. (1997).
Specific Activation of the Cysteine Protease CPP32 during the Negative Selection of T Cells in the
Thymus. *Journal of Experimental Medicine*, *186*(9), 1503-1512.
https://doi.org/10.1084/jem.186.9.1503
Anderson, M. S., & Su, M. A. (2016). AIRE expands: new roles in immune tolerance and beyond. *Nat Rev*
Immunol, *16*(4), 247-258. https://doi.org/10.1038/nri.2016.9
Anderson, M. S., Venanzi, E. S., Klein, L., Chen, Z., Berzins, S. P., Turley, S. J., von Boehmer, H., Bronson,
R., Dierich, A., Benoist, C., & Mathis, D. (2002). Projection of an immunological self shadow
within the thymus by the aire protein. *Science*, *298*(5597), 1395-1401.
https://doi.org/10.1126/science.1075958
Ardouin, L., Luche, H., Chelbi, R., Carpentier, S., Shawket, A., Montanana Sanchis, F., Santa Maria, C.,
Grenot, P., Alexandre, Y., Gregoire, C., Fries, A., Vu Manh, T. P., Tamoutounour, S., Crozat, K.,
Tomasello, E., Jorquera, A., Fossum, E., Bogen, B., Azukizawa, H., . . . Malissen, B. (2016). Broad
and Largely Concordant Molecular Changes Characterize Tolerogenic and Immunogenic
Dendritic Cell Maturation in Thymus and Periphery. *Immunity*, *45*(2), 305-318.
https://doi.org/10.1016/j.immuni.2016.07.019
Atibalentja, D. F., Byersdorfer, C. A., & Unanue, E. R. (2009). Thymus-blood protein interactions are
highly effective in negative selection and regulatory T cell induction. *J Immunol*, *183*(12), 7909-
7918. https://doi.org/10.4049/jimmunol.0902632
Au-Yeung, B. B., Melichar, H. J., Ross, J. O., Cheng, D. A., Zikherman, J., Shokat, K. M., Robey, E. A., &
Weiss, A. (2014). Quantitative and temporal requirements revealed for Zap70 catalytic activity
during T cell development. *Nat Immunol*, *15*(7), 687-694. https://doi.org/10.1038/ni.2918
Azzam, H. S., Grinberg, A., Lui, K., Shen, H., Shores, E. W., & Love, P. E. (1998). CD5 expression is
developmentally regulated by T cell receptor (TCR) signals and TCR avidity. *J Exp Med*, *188*(12),
2301-2311. https://doi.org/10.1084/jem.188.12.2301
Bendelac, A., Killeen, N., Littman, D. R., & Schwartz, R. H. (1994). A subset of CD4+ thymocytes selected
by MHC class I molecules. *Science*, *263*(5154), 1774-1778.
https://doi.org/10.1126/science.7907820
Bleul, C. C., & Boehm, T. (2000). Chemokines define distinct microenvironments in the developing
thymus. *Eur J Immunol*, *30*(12), 3371-3379. https://doi.org/10.1002/1521-
4141(2000012)30:12<3371::AID-IMMU3371>3.0.CO;2-L
Bonasio, R., Scimone, M. L., Schaerli, P., Grabie, N., Lichtman, A. H., & von Andrian, U. H. (2006). Clonal
deletion of thymocytes by circulating dendritic cells homing to the thymus. *Nat Immunol*, *7*(10),
1092-1100. https://doi.org/10.1038/ni1385
Bornstein, C., Nevo, S., Giladi, A., Kadouri, N., Pouzolles, M., Gerbe, F., David, E., Machado, A., Chuprin,
A., Toth, B., Goldberg, O., Itzkovitz, S., Taylor, N., Jay, P., Zimmermann, V. S., Abramson, J., &
Amit, I. (2018). Single-cell mapping of the thymic stroma identifies IL-25-producing tuft epithelial
cells. *Nature*, *559*(7715), 622-626. https://doi.org/10.1038/s41586-018-0346-1
Boursalian, T. E., Golob, J., Soper, D. M., Cooper, C. J., & Fink, P. J. (2004). Continued maturation of
thymic emigrants in the periphery. *Nat Immunol*, *5*(4), 418-425. https://doi.org/10.1038/ni1049
Breed, E. R., Watanabe, M., & Hogquist, K. A. (2019). Measuring Thymic Clonal Deletion at the
Population Level. *J Immunol*, *202*(11), 3226-3233. https://doi.org/10.4049/jimmunol.1900191
Brennecke, P., Reyes, A., Pinto, S., Rattay, K., Nguyen, M., Kuchler, R., Huber, W., Kyewski, B., &
Steinmetz, L. M. (2015). Single-cell transcriptome analysis reveals coordinated ectopic gene-

expression patterns in medullary thymic epithelial cells. *Nat Immunol*, *16*(9), 933-941. https://doi.org/10.1038/ni.3246

Britschgi, M. R., Link, A., Lissandrin, T. K. A., & Luther, S. A. (2008). Dynamic Modulation of CCR7 Expression and Function on Naive T Lymphocytes In Vivo1. *The Journal of Immunology*, *181*(11), 7681-7688. https://doi.org/10.4049/jimmunol.181.11.7681

Brugnera, E., Bhandoola, A., Cibotti, R., Yu, Q., Guinter, T. I., Yamashita, Y., Sharrow, S. O., & Singer, A. (2000). Coreceptor reversal in the thymus: signaled CD4+8+ thymocytes initially terminate CD8 transcription even when differentiating into CD8+ T cells. *Immunity*, *13*(1), 59-71. https://doi.org/10.1016/s1074-7613(00)00008-x

Campbell, J. J., Pan, J., & Butcher, E. C. (1999). Cutting edge: developmental switches in chemokine responses during T cell maturation. *J Immunol*, *163*(5), 2353-2357. https://doi.org/ji_v163n5p2353 [pii]

Cepeda, S., Cantu, C., Orozco, S., Xiao, Y., Brown, Z., Semwal, M. K., Venables, T., Anderson, M. S., & Griffith, A. V. (2018). Age-Associated Decline in Thymic B Cell Expression of Aire and Aire-Dependent Self-Antigens. *Cell Rep*, *22*(5), 1276-1287. https://doi.org/10.1016/j.celrep.2018.01.015

Chvatchko, Y., Hoogewerf, A. J., Meyer, A., Alouani, S., Juillard, P., Buser, R., Conquet, F., Proudfoot, A. E. I., Wells, T. N. C., & Power, C. A. (2000). A Key Role for Cc Chemokine Receptor 4 in Lipopolysaccharide-Induced Endotoxic Shock. *Journal of Experimental Medicine*, *191*(10), 1755-1764. https://doi.org/10.1084/jem.191.10.1755

Cowan, J. E., McCarthy, N. I., Parnell, S. M., White, A. J., Bacon, A., Serge, A., Irla, M., Lane, P. J., Jenkinson, E. J., Jenkinson, W. E., & Anderson, G. (2014). Differential requirement for CCR4 and CCR7 during the development of innate and adaptive alphabetaT cells in the adult thymus. *J Immunol*, *193*(3), 1204-1212. https://doi.org/10.4049/jimmunol.1400993

Daley, S. R., Hu, D. Y., & Goodnow, C. C. (2013). Helios marks strongly autoreactive CD4+ T cells in two major waves of thymic deletion distinguished by induction of PD-1 or NF-kappaB. *J Exp Med*, *210*(2), 269-285. https://doi.org/10.1084/jem.20121458

DeVoss, J., Hou, Y., Johannes, K., Lu, W., Liou, G. I., Rinn, J., Chang, H., Caspi, R. R., Fong, L., & Anderson, M. S. (2006). Spontaneous autoimmunity prevented by thymic expression of a single self-antigen. *J Exp Med*, *203*(12), 2727-2735. https://doi.org/10.1084/jem.20061864

Dzhagalov, I. L., Chen, K. G., Herzmark, P., & Robey, E. A. (2013). Elimination of self-reactive T cells in the thymus: a timeline for negative selection. *PLoS Biol*, *11*(5), e1001566. https://doi.org/10.1371/journal.pbio.1001566

Ehrlich, L. I., Oh, D. Y., Weissman, I. L., & Lewis, R. S. (2009). Differential contribution of chemotaxis and substrate restriction to segregation of immature and mature thymocytes. *Immunity*, *31*(6), 986-998. https://doi.org/10.1016/j.immuni.2009.09.020

Control of Migration during Intrathymic T Cell Development, 1 Encyclopedia of Immunobiology 249-262 (Elsevier 2016).

Finnish-German, A. C. (1997). An autoimmune disease, APECED, caused by mutations in a novel gene featuring two PHD-type zinc-finger domains. *Nat Genet*, *17*(4), 399-403. https://doi.org/10.1038/ng1297-399

Fu, G., Vallee, S., Rybakin, V., McGuire, M. V., Ampudia, J., Brockmeyer, C., Salek, M., Fallen, P. R., Hoerter, J. A., Munshi, A., Huang, Y. H., Hu, J., Fox, H. S., Sauer, K., Acuto, O., & Gascoigne, N. R. (2009). Themis controls thymocyte selection through regulation of T cell antigen receptor-mediated signaling. *Nat Immunol*, *10*(8), 848-856. https://doi.org/10.1038/ni.1766

Hale, L. P., Greer, P. K., Trinh, C. T., & Gottfried, M. R. (2005). Treatment with oral bromelain decreases colonic inflammation in the IL-10-deficient murine model of inflammatory bowel disease.

Clinical Immunology, *116*(2), 135-142.
https://doi.org/https://doi.org/10.1016/j.clim.2005.04.011
Hawiger, D., Masilamani, R. F., Bettelli, E., Kuchroo, V. K., & Nussenzweig, M. C. (2004). Immunological
unresponsiveness characterized by increased expression of CD5 on peripheral T cells induced by
dendritic cells in vivo. *Immunity*, *20*(6), 695-705. https://doi.org/10.1016/j.immuni.2004.05.002
Hu, D. Y., Yap, J. Y., Wirasinha, R. C., Howard, D. R., Goodnow, C. C., & Daley, S. R. (2016). A timeline
demarcating two waves of clonal deletion and Foxp3 upregulation during thymocyte
development. *Immunol Cell Biol*, *94*(4), 357-366. https://doi.org/10.1038/icb.2015.95
Hu, Z., Lancaster, J. N., & Ehrlich, L. I. (2015). The Contribution of Chemokines and Migration to the
Induction of Central Tolerance in the Thymus. *Front Immunol*, *6*, 398.
https://doi.org/10.3389/fimmu.2015.00398
Hu, Z., Lancaster, J. N., Sasiponganan, C., & Ehrlich, L. I. (2015). CCR4 promotes medullary entry and
thymocyte-dendritic cell interactions required for central tolerance. *J Exp Med*, *212*(11), 1947-
1965. https://doi.org/10.1084/jem.20150178
Hu, Z., Li, Y., Van Nieuwenhuijze, A., Selden, H. J., Jarrett, A. M., Sorace, A. G., Yankeelov, T. E., Liston, A.,
& Ehrlich, L. I. R. (2017). CCR7 Modulates the Generation of Thymic Regulatory T Cells by
Altering the Composition of the Thymic Dendritic Cell Compartment. *Cell Rep*, *21*(1), 168-180.
https://doi.org/10.1016/j.celrep.2017.09.016
Kadakia, T., Tai, X., Kruhlak, M., Wisniewski, J., Hwang, I. Y., Roy, S., Guinter, T. I., Alag, A., Kehrl, J. H.,
Zhuang, Y., & Singer, A. (2019). E-protein-regulated expression of CXCR4 adheres preselection
thymocytes to the thymic cortex. *J Exp Med*, *216*(8), 1749-1761.
https://doi.org/10.1084/jem.20182285
Ki, S., Park, D., Selden, H. J., Seita, J., Chung, H., Kim, J., Iyer, V. R., & Ehrlich, L. I. R. (2014). Global
transcriptional profiling reveals distinct functions of thymic stromal subsets and age-related
changes during thymic involution. *Cell Rep*, *9*(1), 402-415.
https://doi.org/10.1016/j.celrep.2014.08.070
Klein, L., Kyewski, B., Allen, P. M., & Hogquist, K. A. (2014). Positive and negative selection of the T cell
repertoire: what thymocytes see (and don't see). *Nat Rev Immunol*, *14*(6), 377-391.
https://doi.org/10.1038/nri3667
Klein, L., Robey, E. A., & Hsieh, C. S. (2019). Central CD4(+) T cell tolerance: deletion versus regulatory T
cell differentiation. *Nat Rev Immunol*, *19*(1), 7-18. https://doi.org/10.1038/s41577-018-0083-6
Koble, C., & Kyewski, B. (2009). The thymic medulla: a unique microenvironment for intercellular self-
antigen transfer. *J Exp Med*, *206*(7), 1505-1513. https://doi.org/10.1084/jem.20082449
Kozai, M., Kubo, Y., Katakai, T., Kondo, H., Kiyonari, H., Schaeuble, K., Luther, S. A., Ishimaru, N.,
Ohigashi, I., & Takahama, Y. (2017). Essential role of CCL21 in establishment of central self-
tolerance in T cells. *J Exp Med*, *214*(7), 1925-1935. https://doi.org/10.1084/jem.20161864
Kurobe, H., Liu, C., Ueno, T., Saito, F., Ohigashi, I., Seach, N., Arakaki, R., Hayashi, Y., Kitagawa, T., Lipp,
M., Boyd, R. L., & Takahama, Y. (2006). CCR7-dependent cortex-to-medulla migration of
positively selected thymocytes is essential for establishing central tolerance. *Immunity*, *24*(2),
165-177. https://doi.org/10.1016/j.immuni.2005.12.011
Lancaster, J. N., & Ehrlich, L. I. R. (2017). Analysis of Thymocyte Migration, Cellular Interactions, and
Activation by Multiphoton Fluorescence Microscopy of Live Thymic Slices. In G. E. Rainger & H.
M. McGettrick (Eds.), *T-Cell Trafficking: Methods and Protocols* (pp. 9-25). Springer New York.
https://doi.org/10.1007/978-1-4939-6931-9_2
Lancaster, J. N., Li, Y., & Ehrlich, L. I. R. (2018). Chemokine-Mediated Choreography of Thymocyte
Development and Selection. *Trends Immunol*, *39*(2), 86-98.
https://doi.org/10.1016/j.it.2017.10.007

Lancaster, J. N., Thyagarajan, H. M., Srinivasan, J., Li, Y., Hu, Z., & Ehrlich, L. I. R. (2019). Live-cell imaging
reveals the relative contributions of antigen-presenting cell subsets to thymic central tolerance.
Nat Commun, *10*(1), 2220. https://doi.org/10.1038/s41467-019-09727-4

Liu, B., Lin, Y., Yan, J., Yao, J., Liu, D., Ma, W., Wang, J., Liu, W., Wang, C., Zhang, L., & Qi, H. (2021).
Affinity-coupled CCL22 promotes positive selection in germinal centres. *Nature*, *592*(7852), 133-
137. https://doi.org/10.1038/s41586-021-03239-2

Lutes, L. K., Steier, Z., McIntyre, L. L., Pandey, S., Kaminski, J., Hoover, A. R., Ariotti, S., Streets, A., Yosef,
N., & Robey, E. A. (2021). T cell self-reactivity during thymic development dictates the timing of
positive selection. *eLife*, *10*, 1-28. https://doi.org/10.7554/eLife.65435

Marchetti, M. C., Di Marco, B., Cifone, G., Migliorati, G., & Riccardi, C. (2003). Dexamethasone-induced
apoptosis of thymocytes: role of glucocorticoid receptor-associated Src kinase and caspase-8
activation. *Blood*, *101*(2), 585-593. https://doi.org/10.1182/blood-2002-06-1779

Mathis, D., & Benoist, C. (2007). A decade of AIRE. *Nat Rev Immunol*, *7*(8), 645-650.
https://doi.org/10.1038/nri2136

McCaughtry, T. M., Baldwin, T. A., Wilken, M. S., & Hogquist, K. A. (2008). Clonal deletion of thymocytes
can occur in the cortex with no involvement of the medulla. *J Exp Med*, *205*(11), 2575-2584.
https://doi.org/10.1084/jem.20080866

McCaughtry, T. M., Wilken, M. S., & Hogquist, K. A. (2007). Thymic emigration revisited. *J Exp Med*,
204(11), 2513-2520. https://doi.org/10.1084/jem.20070601

Melichar, H. J., Ross, J. O., Herzmark, P., Hogquist, K. A., & Robey, E. A. (2013). Distinct temporal
patterns of T cell receptor signaling during positive versus negative selection in situ. *Sci Signal*,
6(297), ra92. https://doi.org/10.1126/scisignal.2004400

Meredith, M., Zemmour, D., Mathis, D., & Benoist, C. (2015). Aire controls gene expression in the thymic
epithelium with ordered stochasticity. *Nat Immunol*, *16*(9), 942-949.
https://doi.org/10.1038/ni.3247

Misslitz, A., Pabst, O., Hintzen, G., Ohl, L., Kremmer, E., Petrie, H. T., & Forster, R. (2004). Thymic T cell
development and progenitor localization depend on CCR7. *J Exp Med*, *200*(4), 481-491.
https://doi.org/10.1084/jem.20040383

Nagamine, K., Peterson, P., Scott, H. S., Kudoh, J., Minoshima, S., Heino, M., Krohn, K. J., Lalioti, M. D.,
Mullis, P. E., Antonarakis, S. E., Kawasaki, K., Asakawa, S., Ito, F., & Shimizu, N. (1997). Positional
cloning of the APECED gene. *Nat Genet*, *17*(4), 393-398. https://doi.org/10.1038/ng1297-393

Nitta, T., Nitta, S., Lei, Y., Lipp, M., & Takahama, Y. (2009). CCR7-mediated migration of developing
thymocytes to the medulla is essential for negative selection to tissue-restricted antigens. *Proc*
Natl Acad Sci U S A, *106*(40), 17129-17133. https://doi.org/10.1073/pnas.0906956106

Oh, J., Wu, N., Barczak, A. J., Barbeau, R., Erle, D. J., & Shin, J. S. (2018). CD40 Mediates Maturation of
Thymic Dendritic Cells Driven by Self-Reactive CD4(+) Thymocytes and Supports Development of
Natural Regulatory T Cells. *J Immunol*, *200*(4), 1399-1412.
https://doi.org/10.4049/jimmunol.1700768

Perera, J., Zheng, Z., Li, S., Gudjonson, H., Kalinina, O., Benichou, J. I. C., Block, K. E., Louzoun, Y., Yin, D.,
Chong, A. S., Dinner, A. R., Weigert, M., & Huang, H. (2016). Self-Antigen-Driven Thymic B Cell
Class Switching Promotes T Cell Central Tolerance. *Cell Rep*, *17*(2), 387-398.
https://doi.org/10.1016/j.celrep.2016.09.011

Persaud, S. P., Parker, C. R., Lo, W. L., Weber, K. S., & Allen, P. M. (2014). Intrinsic CD4+ T cell sensitivity
and response to a pathogen are set and sustained by avidity for thymic and peripheral
complexes of self peptide and MHC. *Nat Immunol*, *15*(3), 266-274.
https://doi.org/10.1038/ni.2822

Petrie, H. T., & Zuniga-Pflucker, J. C. (2007). Zoned out: functional mapping of stromal signaling
microenvironments in the thymus. *Annu Rev Immunol*, *25*, 649-679.
https://doi.org/10.1146/annurev.immunol.23.021704.115715

Ross, J. O., Melichar, H. J., Au-Yeung, B. B., Herzmark, P., Weiss, A., & Robey, E. A. (2014). Distinct phases
in the positive selection of CD8+ T cells distinguished by intrathymic migration and T-cell
receptor signaling patterns. *Proc Natl Acad Sci U S A*, *111*(25), E2550-2558.
https://doi.org/10.1073/pnas.1408482111

Ruscher, R., Kummer, R. L., Lee, Y. J., Jameson, S. C., & Hogquist, K. A. (2017). CD8alphaalpha
intraepithelial lymphocytes arise from two main thymic precursors. *Nat Immunol*, *18*(7), 771-
779. https://doi.org/10.1038/ni.3751

Sansom, S. N., Shikama-Dorn, N., Zhanybekova, S., Nusspaumer, G., Macaulay, I. C., Deadman, M. E.,
Heger, A., Ponting, C. P., & Hollander, G. A. (2014). Population and single-cell genomics reveal
the Aire dependency, relief from Polycomb silencing, and distribution of self-antigen expression
in thymic epithelia. *Genome Res*, *24*(12), 1918-1931. https://doi.org/10.1101/gr.171645.113

Sinclair, C., Bains, I., Yates, A. J., & Seddon, B. (2013). Asymmetric thymocyte death underlies the
CD4:CD8 T-cell ratio in the adaptive immune system. *Proc Natl Acad Sci U S A*, *110*(31), E2905-
2914. https://doi.org/10.1073/pnas.1304859110

Stritesky, G. L., Xing, Y., Erickson, J. R., Kalekar, L. A., Wang, X., Mueller, D. L., Jameson, S. C., & Hogquist,
K. A. (2013). Murine thymic selection quantified using a unique method to capture deleted T
cells. *Proc Natl Acad Sci U S A*, *110*(12), 4679-4684. https://doi.org/10.1073/pnas.1217532110

Ueno, T., Hara, K., Willis, M. S., Malin, M. A., Hopken, U. E., Gray, D. H., Matsushima, K., Lipp, M.,
Springer, T. A., Boyd, R. L., Yoshie, O., & Takahama, Y. (2002). Role for CCR7 ligands in the
emigration of newly generated T lymphocytes from the neonatal thymus. *Immunity*, *16*(2), 205-
218. https://doi.org/10.1016/s1074-7613(02)00267-4

Ueno, T., Saito, F., Gray, D. H., Kuse, S., Hieshima, K., Nakano, H., Kakiuchi, T., Lipp, M., Boyd, R. L., &
Takahama, Y. (2004). CCR7 signals are essential for cortex-medulla migration of developing
thymocytes. *J Exp Med*, *200*(4), 493-505. https://doi.org/10.1084/jem.20040643

Vollmann, E. H., Rattay, K., Barreiro, O., Thiriot, A., Fuhlbrigge, R. A., Vrbanac, V., Kim, K. W., Jung, S.,
Tager, A. M., & von Andrian, U. H. (2021). Specialized transendothelial dendritic cells mediate
thymic T-cell selection against blood-borne macromolecules. *Nat Commun*, *12*(1), 6230.
https://doi.org/10.1038/s41467-021-26446-x

Wright, D. E., Cheshier, S. H., Wagers, A. J., Randall, T. D., Christensen, J. L., & Weissman, I. L. (2001).
Cyclophosphamide/granulocyte colony-stimulating factor causes selective mobilization of bone
marrow hematopoietic stem cells into the blood after M phase of the cell cycle. *Blood*, *97*(8),
2278-2285. https://doi.org/10.1182/blood.v97.8.2278

Xing, Y., Wang, X., Jameson, S. C., & Hogquist, K. A. (2016). Late stages of T cell maturation in the thymus
involve NF-kappaB and tonic type I interferon signaling. *Nat Immunol*, *17*(5), 565-573.
https://doi.org/10.1038/ni.3419

Yamano, T., Nedjic, J., Hinterberger, M., Steinert, M., Koser, S., Pinto, S., Gerdes, N., Lutgens, E.,
Ishimaru, N., Busslinger, M., Brors, B., Kyewski, B., & Klein, L. (2015). Thymic B Cells Are Licensed
to Present Self Antigens for Central T Cell Tolerance Induction. *Immunity*, *42*(6), 1048-1061.
https://doi.org/10.1016/j.immuni.2015.05.013

Zegarra-Ruiz, D. F., Kim, D. V., Norwood, K., Kim, M., Wu, W. H., Saldana-Morales, F. B., Hill, A. A.,
Majumdar, S., Orozco, S., Bell, R., Round, J. L., Longman, R. S., Egawa, T., Bettini, M. L., & Diehl,
G. E. (2021). Thymic development of gut-microbiota-specific T cells. *Nature*, *594*(7863), 413-417.
https://doi.org/10.1038/s41586-021-03531-1

Figures and Legends

Figure 1. CCR4 and CCR7 are expressed by immature and mature subsets of post-positive selection
thymocytes, respectively.

A) Flow cytometry gating scheme to delineate thymocyte subsets in Rag2p-GFP mice. Post-positive
selection DP (CD3^{lo}CD69$^+$ and CD3$^+$CD69$^+$), CD4SP (SM, M1, and M2), and CD8SP (M1 and M2) subsets
are annotated. **B)** Relative Rag2p-GFP Geometric Mean Fluorescent Intensity (GMFI) of each subset
normalized to pre-positive selection DP CD3$^-$CD69$^-$ cells. Individual data points represent average GFP
GMFI of the indicated subset from each mouse. **C)** Representative flow cytometry plots showing CCR4
and CCR7 expression by post-positive selection thymocyte subsets. **D)** Quantification of the percentage
of cells of the indicated subset that express CCR4, CCR7, or both chemokine receptors. For B) and D),
data were compiled from 4 independent experiments (Mean ± SEM; N=10 mice).

Figure 2. **CCR4 and CCR7 promote chemotaxis of immature and mature post-positive selection thymocyte subsets, respectively.**

A-D) Transwell assays were used to quantify chemotaxis of thymocyte subsets to the indicated concentrations of the CCR4 ligands CCL17 (A) and CCL22 (B) and the CCR7 ligands CCL19 (C) and CCL21 (D). Migration indexes were calculated as the frequencies of input cells of each subset that migrated towards the chemokine relative to the frequencies of input cells that migrated in the absence of chemokine (Blank). Data were compiled from 3 independent experiments with triplicate wells per assay. 2-way ANOVA with Dunnett's multiple comparison correction were used for statistical analysis. (Mean ± SEM; *: $p < 0.05$, **: $p < 0.01$, ***: $p < 0.001$, ****: $p < 0.0001$)

Figure 3. Persistent CXCR4 expression was detected on post-positive selection thymocytes, and CXCR4 activity declines only in intermediate subsets.

A) Representative flow cytometry plots showing CXCR4 surface expression by the indicated thymocyte subsets, compared to isotype control stains. **B)** Quantification of relative GMFI of CXCR4 for the indicated thymocyte subsets. (Mean ± SEM; N=10) **C)** Transwell assays were used to quantify chemotaxis of thymocyte subsets to the indicated concentrations of CXCL12. Migration indexes were calculated as the frequencies of input cells of each subset that migrated towards the chemokine relative to the frequencies of input cells that migrated in the absence of chemokine (Blank). Data were compiled from 3 independent experiments with triplicate wells per assay. 2-way ANOVA with Dunnett's multiple comparison correction were used for statistical analysis. (Mean ± SEM; *: p<0.05, **: p<0.01, ***: p<0.001, ****: p<0.0001) **D)** Graphical summary showing relative expression and chemotactic activity of CXCR4, CCR4 and CCR7 on post-positive selection thymocyte subsets based on data in Figures 1-3.

Figure 4. Rapid upregulation of CCR4 following positive selection of OT-II thymocytes correlates with medullary entry.

A) Experimental schematic of synchronized thymic slice positive selection assay to determine timing of CCR4 and CCR7 expression by flow cytometry and medullary entry by 2-photon microscopy. **B)** Representative flow cytometry plots showing CCR4 and CCR7 expression of OT-II$^+$ *Rag2$^{-/-}$ MHCII$^{-/-}$* thymocytes added to thymic slices (left) and analyzed 24h after incubation in WT or *MHCII$^{-/-}$* slices, as indicated (right). **C)** Percentages of CCR4$^+$ and CCR4$^+$CCR7$^+$ OT-II$^+$ thymocytes in thymic slices of the indicated genotypes at the indicated time points. Data were compiled from 3 experiments with triplicate slices each. Mixed-effect analysis with Šídák's multiple comparison correction was used for statistical analysis. (Mean ± SEM; *: p<0.05, **: p<0.01) **D)** Representative maximum intensity projections of 2-photon imaging data showing a pCX-EGFP thymic slice (green) containing CMTPX-labeled OT-II$^+$ *Rag2$^{-/-}$ MHCII$^{-/-}$* thymocytes (red) at 1.5h and 12h after thymocytes were added to the slices. Medullary and cortical volumes were demarcated as indicated by the masked regions. **E)** Quantification of medullary density of OT-II$^+$ *Rag2$^{-/-}$ MHCII$^{-/-}$* thymocytes at indicated time points after incubation on pCX-EGFP thymic slices. Individual data points represent relative medullary input cell densities from each Video compared to the average of medullary densities from all 1.5h Videos. Data were compiled from 4 time-course experiments. One-way ANOVA with Tukey's multiple comparison correction were used for statistical analysis. (Mean ± SEM; *: p<0.05, **: p<0.01, ***: p<0.001, ****: p<0.0001)

Figure 5. CCR4 and CCR7 are required for medullary accumulation of distinct post-positive selection thymocyte subsets.

A) Representative maximum intensity projections of 2-photon imaging data showing pCX-EGFP thymic slices (green) containing CMTPX (red)- and Indo-1 AM (blue)-labeled FACS sorted DP CD3^{lo}CD69$^+$, CD4SP SM or CD4SP M1+M2 cells from WT or *Ccr4$^{-/-}$* mice, as indicated, imaged 1-4 hours after slice entry. The cortex and medulla are demarcated by white dotted lines, and cell tracks are color encoded for elapsed imaging time. **B)** Medullary-to-cortical thymocyte density ratios were quantified from 2-photon imaging data as in A), with individual data points representing ratios from individual Videos. **C)** Paired analysis of medullary-to-cortical density ratios of WT versus *Ccr4$^{-/-}$*, WT versus *Ccr7$^{-/-}$*, or *Ccr7$^{-/-}$* versus *Ccr4$^{-/-}$Ccr7$^{-/-}$* thymocyte subsets imaged simultaneously within the same slices, with individual data points representing ratios from single videos. Paired datasets are taken from the same data shown in B), and paired t-tests were used for statistical analysis. (**: p<0.01, ***: p<0.001) **D-E)** Quantification of D) average thymocyte track speeds and E) average track straightness from imaging data reported in B). Individual data points represent average track speeds or straightness of thymocytes of the indicated genotype in individual Videos. For B), D) and E), data were compiled from 19 independent experiments. One-way ANOVA with Tukey's multiple comparison correction were used for statistical analysis. (Mean ± SEM; *: p<0.05, **: p<0.01, ***: p<0.001, ****: p<0.0001)

Figure 6. CCR4 and CCR7 contribute to early versus late phases of negative selection, respectively.

A) Flow cytometric gating scheme to quantify thymocytes undergoing early-phase (DP) or late-phase (CD4SP+CD8SP) negative selection. Representative data from a WT mouse is shown. **B)** Representative flow cytometry plots showing CCR4 and CCR7 expression by thymocytes undergoing early- and late-phase negative selection, as gated in C). **C)** Frequencies of total, early-phase (DP), and late-phase (CD4SP+CD8SP) negative selection among all live thymocytes in WT, *Ccr4*[-/-], *Ccr7*[-/-], and *Ccr4*[-/-]*Ccr7*[-/-] mice. **D)** Ratio of thymocytes undergoing early- to late-phase negative selection in WT, *Ccr4*[-/-], *Ccr7*[-/-], and *Ccr4*[-/-]*Ccr7*[-/-] mice. For C) and D), One-way ANOVA with Dunnett's multiple comparison correction were used for statistical analysis. (n=22 for WT, n=20 for *Ccr4*[-/-], n=14 for *Ccr7*[-/-], n=24 for *Ccr4*[-/-] *Ccr7*[-/-]. Mean ± SEM; *: p<0.05, **: p<0.01, ***: p<0.001, ****: p<0.0001)

1005

1006

Figure 7. CCR4 and CCR7 promote tolerance to distinct organs, with CCR4 promoting tolerance to activated antigen-presenting cells.

A) Representative images of H&E stains from the colon of 5–6-month-old WT, *Ccr4-/-*, *Ccr7-/-* or *Ccr4-/-Ccr7-/-* mice (left). Bars, 200 μm. Histological lesion score was quantified by a pathologist blinded to treatment group (right). One-way ANOVA was used to determine significance between groups (Mean ± SEM; ***: p<0.001).**B)** Representative images of H&E stains from extraorbital lacrimal glands of 5–6-month-old WT, *Ccr4-/-*, *Ccr7-/-* or *Ccr4-/-Ccr7-/-* mice (left). Bars, 500 μm. Arrows indicate lymphocytic infiltrate foci. The number of infiltrate foci per 10mm^2 was quantified (right), and one-way ANOVA was used to test significance between groups (Mean ± SEM; *: p<0.05, ***: p>0.001): C) Representative confocal images of CD4 (green), CD8 (red), and B220 (blue) on immunostained cryosections of inguinal lymph nodes from 5-6 month-old WT and *Ccr4-/-* mice(left). Frequency of moderate lymphoid hyperplasia in mLNs and spleens from 12-13 month-old WT, *Ccr4-/-*and *Ccr7-/-* mice, as determined by a veterinary pathologist (right). Fisher's exact test was used to determine significance. **D-E)** Expression of CCR4 ligands (D) and CCR7 ligands (E) by distinct thymic antigen presenting cell subsets, assessed by qRT-PCR. Expression levels were normalized to subsets previously reported to expresses the respective ligands: CCL17 and CCL22 expression were normalized to cDC2 (Hu, Lancaster, Sasiponganan, et al., 2015), and CCL19 and CCL21 expression to mTEClo (Ki et al., 2014; Misslitz et al., 2004). Data were compiled from 3 independent experiments with triplicates. One-way ANOVA with Dunnett's multiple comparison correction were used for statistical analysis. (Mean ± SEM; ns: p>0.05, **: p<0.01) **F)** The percentage of WT or *Ccr4-/-* T cells that proliferated at day 3 and day 5 when cultured without

1027   splenocytes, with unstimulated splenocytes, or with LPS-stimulated splenocytes, as indicated. Data were
1028   compiled from 4 independent experiments with triplicate wells, and unpaired t-tests with Holm-Šídák's
1029   multiple comparison correction was used for statistical analysis. (Mean ± SEM; *: $p < 0.05$)

1030

1032 **Supplemental Figures and Legends**

1033

Figure 1—figure supplement 1. DP CD3^{lo}CD69^{+} cells are consistent with post-positive selection DPs
and DP CD3^{+}CD69^{+} cells represent co-receptor reversing MHCI-restricted thymocytes.

A) Quantification of the GFP MFI of the indicated thymocyte subsets from Nur77^{GFP} mice. Data were
compiled from 3 independent experiments (Nur77^{GFP+} N=4 mice, Nur77^{GFP-} N=3 mice). **B)** Flow cytometry
plots showing the absence of DP CD3^{+}CD69^{+} thymocytes from *β2m$^{-/-}$*, but not *MHCII$^{-/-}$* thymus. **C)**
Quantification of the frequency of DP CD3^{+}CD69^{+} in all DP thymocytes from wild-type, *β2m$^{-/-}$*, and *MHCII^{-}*
$^{/-}$ mice. One-way ANOVA with Dunnett's multiple comparison correction was used for statistical analysis.
(n=3 for wild-type, n=4 for *β2m$^{-/-}$*, n=3 for *MHCII$^{-/-}$*. Mean ± SEM; *: p<0.05, ***: p<0.001).

1042

1043

Figure 1—figure supplement 2. CCR7 expression on developing thymocytes begin at the CD4SP SM stage.

A) Representative flow cytometry overlays showing CCR4 and CCR7 expression by post-positive selection thymocyte subsets from WT and *Ccr7*^-/- mice, respectively. **B)** Quantification of the CCR7 MFI within each subset from data in A (N = 4 mice of each genotype). **C-D)** Quantification of CCR7 and CCR4 MFIs for each thymocyte subset after incubation at 37°C for the indicated times, prior to antibody staining at 4°C. Data were compiled from 2 independent experiments (n=4 biologic replicates).

1051

A

1052

Figure 2—figure supplement 1. Chemokine signaling is not diminished in CD4SP SM thymocytes.

A) Migration indices from data in Fig. 2 for DP CD3[lo]CD69[+], CD4SP SM and DP CD3[lo]CD69[+] were analyzed separately to compare the relative efficiency of CCR4 ligands and CCR7 ligands at inducing chemotaxis of each subset. Ordinary one-way ANOVA with Dunnett's multiple comparison corrections were used for statistical analysis. (Mean ± SEM; *: p<0.05, **: p<0.01, ***: p<0.001, ****: p<0.0001)

1058

1059

Figure 2—figure supplement 2. CD4SP SM thymocytes migrate towards CCL25 and CCL22 is localized in the medulla.

 A) Transwell assays were used to quantify chemotaxis of thymocyte subsets to the indicated concentrations of the CCR9 ligand, CCL25. Migration indexes were calculated as the frequencies of input cells of each subset that migrated towards the chemokine relative to the frequencies of input cells that migrated in the absence of chemokine (Blank). Data were compiled from 3 independent experiments with triplicate wells per assay. 2-way ANOVA with Dunnett's multiple comparison correction were used for statistical analysis. (Mean ± SEM; *: $p<0.05$, **: $p<0.01$, ****: $p<0.0001$). **B)** Representative confocal images of CCL22 (green), CD11c (red), and CD31 (purple) on immunostained cryosections from WT thymus. The cortex (C) and medulla (M) are demarcated by white dotted lines.

1070

1071

Figure 4— figure supplement 1. Rapid upregulation of CCR4 following positive selection of OT-I thymocytes correlates with medullary entry.

A) Representative flow cytometry plots of CCR4 and CCR7 expression by pre-positive selection OT-I$^+$ *Rag2$^{-/-}$* thymocytes from *β2m$^{-/-}$* hosts that were added to thymic slices (left) or analyzed 24h after incubation in WT or *β2m$^{-/-}$* slices, as indicated (right). **B)** Percentages of CCR4$^+$, CCR4$^+$CCR7$^+$, CCR7$^+$ or CD8SP OT-I$^+$ *Rag2$^{-/-}$* thymocyte safter incubation on thymic slices of the indicated genotypes for the indicated time period up to 24h. **C)** Percentages of CCR7$^+$ or CD8SP OT-I$^+$ *Rag2$^{-/-}$* thymocytes at 48h and 72h time points after initiation of thymic slices assays. Data for B-C were compiled from 3 OT-I$^+$ *Rag2$^{-/-}$* BM -> *β2m$^{-/-}$* experiments, with triplicate slices per experiment. Mixed-effect analysis with Šídák's multiple comparison correction were used for statistical analysis. (Mean ± SEM; *: p<0.05, **: p<0.01, ****: p<0.0001) **D)** Representative maximum intensity projections of 2-photon imaging data showing a pCX-EGFP thymic slice (green) containing CMTPX-labeled OT-I$^+$ *Rag2$^{-/-}$* BM -> *β2m$^{-/-}$* thymocytes (red) at 1.5h and 12h after thymocytes were added to the slices. Medullary and cortical volumes were demarcated as indicated by the masked regions. **E)** Quantification of medullary density of OT-I$^+$ *Rag2$^{-/-}$* cells at the indicated time points. Individual data points represent medullary density calculated from each Video, relative to the average medullary density of all Videos taken at 1.5 hour. Data were compiled from 3 time-course experiments. One-way ANOVA with Tucky's multiple comparison correction were used for statistical analysis. (Mean ± SEM; *: p<0.05)

1090

1091

Figure 5— figure supplement 1. Experimental approach to test if CCR4 and CCR7 are required for medullary entry of distinct FACS- purified post-positive selection thymocyte subsets that differentially express CCR4 and CCR7.

A) Experimental schematic of 2-photon imaging of sorted, fluorescently labeled post-positive selection thymocyte subsets. WT cells were imaged in the same slices with *Ccr4*[-/-] or *Ccr7*[-/-] cells, and *Ccr7*[-/-] cells were imaged in the same slices with *Ccr4*[-/-] *Ccr7*[-/-] cells. Dyes were swapped between different experiments, and we did not find that use of CMTPX versus Indo-1 altered the results. **B-C)** Representative flow cytometry plots showing B) the gating scheme for sorting DP CD3[lo]CD69[+], CD4SP SM and CD4SP M1+M2 subsets for 2-photon imaging, and C) CCR4 and CCR7 expression of the FACS purified thymocyte subsets from a WT mouse.

1102

1103

Figure 6— figure supplement 1. Thymocyte composition and CD5 levels are not altered in *Ccr4*^{-/-} mice.

A) Thymocyte subset frequencies in littermate WT and *Ccr4*^{-/-} mice. **B)** CD5 expression levels on the indicated thymocyte subsets from littermate WT and *Ccr4*^{-/-} mice, normalized to the DP CD3⁻CD69⁻ subset of each mouse. For A) and B), data were compiled from 6 independent experiments and significance determined using unpaired t-tests with Holm-Šídák's multiple comparison correction (Mean ± SEM; n=13 for WT, n=18 for *Ccr4*^{-/-}).

1110

1111

Figure 7— figure supplement 1. CCR4 and CCR7 support self-tolerance in different organs.

A) Representative images of Rag2$^{-/-}$ kidney sections immunostained with serum from 5–6.5-month-old *WT, Ccr4$^{-/-}$, Ccr7$^{-/-}$ or Ccr4$^{-/-}$Ccr7$^{-/-}$* mice (left). Anti-nuclear autoantibodies were detected as an overlay of anti–mouse IgG (red) and the nuclear DAPI stain (blue). Proportion of mice containing anti-nuclear serum autoantibodies was quantified (right). p < 0.05 **: p<0.01 (Fisher's exact test)**. B)** Representative images of H&E stains from the liver of 5–6-month-old *WT, Ccr4$^{-/-}$, Ccr7$^{-/-}$* and *Ccr4$^{-/-}$Ccr7$^{-/-}$* mice (left). Bars, 500 μm. Histological lesion score was quantified by a veterinary pathologist (right). One-way ANOVA was used to test significance between groups (Mean ± SEM; **: p>0.01). **C-D)** Representative images of H&E stains from intraorbital lacrimal glands (C) and submandibular salivary glands (D) of 5–6-month-old *WT, Ccr4$^{-/-}$, Ccr7$^{-/-}$ or Ccr4$^{-/-}$Ccr7$^{-/-}$* mice (left). Bars, 500 μm. Arrows indicate lymphocytic infiltrate foci. The number of infiltrate foci per 10mm^2 was quantified (right). One-way ANOVA was used to determine the significant differences between groups (Mean ± SEM; *: p<0.05, **: p<0.01, ***: p<0.001).

1125

1126

Figure 7— figure supplement 2. CCR4 promotes tolerance to activated antigen presenting cells. A)
Gating scheme for FACS sorting of thymic APC subsets analyzed by qRT-PCR for expression of CCR4 and
CCR7 ligands. **B)** Surface expression of MHCII, CD80 and CD86 on APC subsets from unstimulated or TLR
ligand-stimulated splenocytes used in the proliferation assays in Figure 7F . Individual data points
represent the GMFI of the indicated markers on each respective APC subset. Data were compiled from 5
individual experiments. One-way ANOVA with Dunnett's multiple comparison correction was used for
statistical analysis. (Mean ± SEM; **: p<0.01, ***: p<0.001 ****: p<0.0001) **C)** The percentage of WT
and *Ccr4*$^{-/-}$ T cells that proliferated at days 3 and 5 when cultured with High Molecular Weight (HMW)
poly(I:C), Low Molecular Weight (LMW) poly(I:C), or ODN1826 (CpG)-stimulated splenocytes. Data were
compiled from 3-4 independent experiments with triplicate wells, and unpaired t-tests with Holm-
Šídák's multiple comparison correction was used for statistical analysis. (Mean ± SEM)

1138

Supplemental Video Legends

Video S1. Two-photon live imaging of OT-II+ *Rag2$^{-/-}$ MHCII$^{-/-}$* input thymocytes (red) migrating in a pCX-EGFP thymic slice (green) 1.5 hours after addition to positively selecting thymic slices, as illustrated in Figure 3D. Images were acquired for 15 minutes at 15 second time intervals through a depth of 40mm, and maximum intensity projections of input cells shown side by side.

Video S2. Two-photon live imaging of OT-II+ *Rag2$^{-/-}$ MHCII$^{-/-}$* input thymocytes (red) migrating in a pCX-EGFP thymic slice (green) 12 hours after addition to positively selecting thymic slices, as illustrated in Figure 3D. Images were acquired for 15 minutes at 15 second time intervals through a depth of 40mm, and maximum intensity projections of input cells shown side by side.

Video S3. Two-photon live imaging of OT-I+ *Rag2$^{-/-}$* input thymocytes from a *β2m$^{-/-}$* bone marrow chimera host (red) migrating in a pCX-EGFP thymic slice (green) 1.5 hours after addition to positively selecting thymic slices, as illustrated in Supplemental Figure 3D, respectively. Images were acquired for 15 minutes at 15 second time intervals through a depth of 40mm, and maximum intensity projections of input cells shown side by side.

Video S4. Two-photon live imaging of OT-I+ *Rag2$^{-/-}$* input thymocytes from a *β2m$^{-/-}$* bone marrow chimera host (red) migrating in a pCX-EGFP thymic slice (green) 12 hours after addition to positively selecting thymic slices, as illustrated in Supplemental Figure 3D, respectively. Images were acquired for 15 minutes at 15 second time intervals through a depth of 40mm, and maximum intensity projections of input cells shown side by side.

Video S5. Two-photon live imaging of WT (red) or *Ccr4$^{-/-}$* (blue) FACS sorted DP CD3^{lo}CD69^{+} thymocytes migrating in pCX-EGFP thymic slices (green), as illustrated in Figure 4A. Images were acquired for 15 minutes at 15 second time intervals through a depth of 40mm, and maximum intensity projections of input cells with and without cell tracks, color-coded for elapsed imaging time, are shown side by side.

Video S6. Two-photon live imaging of WT (red) or *Ccr4$^{-/-}$* (blue) FACS sorted CD4SP SM thymocytes migrating in pCX-EGFP thymic slices (green), as illustrated in Figure 4A. Images were acquired for 15 minutes at 15 second time intervals through a depth of 40mm, and maximum intensity projections of input cells with and without cell tracks, color-coded for elapsed imaging time, are shown side by side.

Video S7. Two-photon live imaging of WT (red) or *Ccr4$^{-/-}$* (blue) FACS sorted CD4SP M1+M2 thymocyte subsets, migrating in pCX-EGFP thymic slices (green), as illustrated in Figure 4A. Images were acquired for 15 minutes at 15 second time intervals through a depth of 40mm, and maximum intensity projections of input cells with and without cell tracks, color-coded for elapsed imaging time, are shown side by side.

1178   **Video S8.** Two-photon live imaging of $Ccr4^{-/-}Ccr7^{-/-}$ (red) or $Ccr7^{-/-}$ (blue) FACS sorted CD4SP M1+M2
1179   subsets, migrating in pCX-EGFP thymic slices (green). Images were acquired for 15 minutes at 15 second
1180   time intervals through a depth of 40mm, and maximum intensity projections of input cells with and
1181   without cell tracks, color-coded for elapsed imaging time, are shown side by side.