Overactivation of a sleep-active neuron decouples survival from the need to sleep

Inka Busack[1], Henrik Bringmann[1]*

1 BIOTEC, Technical University Dresden, Germany

*Corresponding author: henrik.bringmann@tu-dresden.de

Abstract

Sleep is a state of behavioral quiescence that is closely associated with survival. Sleep-

active neurons promote sleep and survival. It is not known, however, whether sleep-active

neurons need to cause sleep to promote survival. Here, we tested how depolarization of the

sleep-active RIS neuron in *Caenorhabditis elegans* controls sleep and survival during

larval starvation. Survival always increased with raised RIS depolarization. RIS

depolarization promoted sleep over a long range. Unexpectedly, however, high levels of

RIS depolarization caused a nearly complete loss of sleep. Similarly, overexpression of

sleep-inducing FLP-11 neuropeptides in RIS inhibited sleep, indicating that overactive

transmission from RIS inhibits sleep. Despite the loss of sleep, survival was normal

following FLP-11 overexpression. Thus, RIS overactivation abolishes sleep yet supports

survival. These results indicate that regulation of sleep and survival are separable functions

of a sleep-active neuron that are normally coupled but can be uncoupled by sleep-neuron

over activation.

Author Summary

Sleep behavior and survival benefits are independently regulated functions of the sleep
neuron RIS in *Caenorhabditis elegans.*

Introduction

Sleep is defined by behavioral inactivity and reduced responsiveness [1]. It is a physiological state that conserves and allocates energy, promotes restorative processes, and is important for brain and cognitive functions [2-4]. Sleep disorders and deviations from the normal sleep amount are widespread in modern societies and pose an unresolved medical and economical problem [5-7].

Central to the induction of sleep in all species are so-called sleep-active neurons that depolarize to inhibit wakefulness behavioral circuits and thus cause sleep [8, 9]. Sleep-active neurons are also important for survival, for example during starvation or after wounding [10, 11]. Little is known about how sleep-active neurons promote survival. Different hypotheses are conceivable. For example, sleep-active neurons might promote survival by reducing behavioral activities during sleep, thus conserving and allocating energy or optimizing behavior [12, 13]. The close association of sleep and physiological benefits has suggested a causal relationship. For example, memory and information processing are thought to be improved in the human brain during sleep when information inputs are blocked [14, 15], and energy is conserved by reduced behavioral activity [12]. A less explored additional hypothesis is that sleep-active neurons provide benefits during sleep, but independently of sleep behavior. According to this hypothesis, sleep-active neurons have two parallel functions. The first is to induce sleep, and the second is to promote survival. This hypothesis implies that physiological benefits of sleep-active neuron functions can be exerted also in the absence of sleep, providing a novel potential starting point for treating health problems associated with sleep loss.

Sleep is found in most animals ranging from cnidarians to humans. *C. elegans* has a small and invariant nervous system and sleep is controlled by relatively few neurons. RIS is a GABAergic and peptidergic interneuron that activates during sleep, and its ablation virtually abolishes sleep behavior [16-19]. Optogenetic manipulations showed that acute depolarization of RIS inhibits wakefulness activity such as feeding and locomotion [16, 19], whereas hyperpolarization inhibits sleep and shortens survival [10, 20-22]. RIS is

required for sleep during various stages and conditions [10, 11, 16, 17, 19, 23-27]. Larval

starvation increases RIS activity and RIS is essential for starvation survival as it counteracts

aging phenotype progression [10]. RIS promotes survival by controlling the transcriptional

response that underlies the arrest, and promotes the activity of the FOXO transcription

factor DAF-16, which is required for survival (our unpublished data) [28, 29]. Thus, RIS

is a sleep-active neuron that is crucial for *C. elegans* sleep and survival.

In order to find out how RIS links sleep and survival, we determined the dose-response

relationship of RIS polarization, sleep and survival. Surprisingly, we found that RIS can

promote survival independently of sleep, indicating that promotion of sleep and survival

are separable functions of RIS that are normally coupled during sleep. Uncoupling of the

two processes provides a strategy to promote survival while gaining waking time.

Results

Generation of a strain set for controlling RIS depolarization levels

Sleep can be controlled by steering the activity of sleep-active neurons [30, 31]. We set out

to control RIS activity to control sleep and survival during L1 arrest. We manipulated RIS

to generate a quantitative dose-response relationship between the polarization state of RIS

and sleep. For this analysis, we generated strains that express constitutively active ion

channels of different strength and permeability in RIS.

To constitutively depolarize RIS, we expressed a K2P channel mutant that is permeable to

sodium (*unc-58(L428F)*, Fig S1A) [32] (Thomas Boulin, personal communication) tagged

with mKate2 at either a lower or a higher level in RIS. The different expression levels were

achieved by generating two different expression loci for the channel in the genome. To

achieve a strong expression of the *unc-58(L428F)* channel (*RIS::unc-58gf(strong)*), we

turned the endogenous *flp-11* locus into an operon that expressed both *flp-11* and the

channel. To achieve a weaker expression of the channel (*RIS::unc-58gf(weak)*), we

expressed mKate2-tagged *unc-58(L428F)* from a single-copy knock-in locus on

chromosome III [33], which contained only a minimal *flp-11* promoter and 3' UTR (Fig 1

A-D). Fluorescent imaging after two days of L1 arrest showed that the stronger strain

expressed at about double the level of the weaker strain and in both genetic systems, the

channel was expressed prominently in RIS and was targeted to the cell membrane (Fig 1A-

D, S1). Even after 21 days of arrest, mKate2-tagged *unc-58(L428F)* expression could be

detected in the strong strain indicating that RIS as well as the sodium channel are intact

across the entire arrest (FigS1E).

To constitutively hyperpolarize RIS, we expressed potassium channels that carry mutations

that cause potassium leak currents. We used different channel genes and mutations to

achieve different strengths of hyperpolarization (*RIS::egl-23gf(weak)* expresses *egl-*

23(L229N) in RIS, *RIS::egl-23gf(strong)* expresses *egl-23(A383V)* in RIS, and *RIS::twk-*

18gf expresses *twk-18(e1913)* in RIS) [32]. In all three strains, the channel genes were

knocked into the endogenous *flp-11* locus behind the *flp-11* gene to generate operons as

above. All protein fusions localized to the cell membrane of the sleep neuron RIS (Fig 1E-

H).

We measured RIS depolarization activity using calcium imaging as a proxy in these

transgenic strains [16]. We were able to measure RIS activity up until the first 12 days of

starvation and found that the effects of the channel transgenes were consistent across the

entire time of starvation that we analyzed (Fig. 1I for two-day starvation, S2C for all time

points). The mean RIS baseline calcium activity levels were reduced by the strong

hyperpolarization transgenes *twk-18gf* and *egl-23gf(strong)*. Depolarization levels were

modestly increased in *unc-58gf(weak)* and strongly increased by the *unc-58gf(strong)*

transgenes. The magnitude of the baseline calcium signal alterations of the different

transgenes were thus in accordance with the expression levels and the expected strengths

of the leak currents (Fig 1I). Hence, we created a set of strains that allows probing dose-

response relationships of depolarization levels and survival as well as sleep.

Survival is a function of RIS depolarization

RIS impairment has been shown to blunt survival during larval arrest and wounding [10,
11, 31]. We thus investigated how RIS depolarization levels affect survival in arrested first-
stage larval worms. For this experiment, we measured survival and the ability to recover
and develop to the adult stage after refeeding [10, 34]. *RIS::twk-18gf* larvae, in which RIS
was strongly hyperpolarized, showed strongly reduced survival by about half compared to
the wild type. This result is in accordance with previous reports that showed that RIS
ablation shortens L1 arrest survival by half, suggesting that strongly hyperpolarized RIS
causes similar effects on lifespan as RIS ablation [10]. *egl-23(strong)* resulted in a more
moderate yet significant lifespan reduction of approximately 21%. Weak hyperpolarization
by *egl-23(weak)* or depolarization by *unc-58gf(weak)* of RIS activity did not significantly
change survival. Strong RIS depolarization by *unc-58gf(strong)* had an increased survival
of around 19% compared to wild type (Fig 1J). To test whether this survival-extending
effect was caused by *unc-58gf* expression in RIS*, we measured survival of *RIS::unc-*
58gf(strong) in the *aptf-1(-)* background, in which expression in RIS is abolished (Fig
S1G). In the combined *RIS::unc-58gf(strong); aptf-1(-)* strain the lifespan prolongation
was extremely shortened compared with wild type (FigS2D) and was similar to *aptf-1(-)*
[10] or *RIS::twk-18gf* (Fig1J), indicating that the lifespan extension indeed stemmed from
unc-58gf expression in RIS. Correlation analysis of survival time versus RIS baseline
activity showed that survival precipitates drastically with decreased RIS baseline activity,
yet increases only mildly with strong depolarization of RIS (Fig 1K). The recovery on food
matched the survival curves (Fig S2E). These results indicate that survival is a function of
RIS depolarization. The more RIS is depolarized, the more it promotes survival.
144
145 **RIS depolarization promotes survival independently of sleep behavior**
146
We next tracked mobility quiescence behavior to estimate sleep during L1 arrest and
quantified RIS activation transients [16]. Consistent with previous reports [10, 16, 18],
wild-type RIS showed activation transients over a constant activity baseline, and calcium
transients were accompanied by phases of behavioral quiescence (Fig 2A-G). Calcium
transients were completely abolished in both the strongest hyperpolarization strains (Fig
2A, B) as well as in the strong depolarization strain (2F). In the mild hyperpolarization

(Fig. 2C) or depolarization strains (Fig. 2E), calcium transients were still visible and correlated with behavioral quiescence, but were decreased in magnitude (Fig. 2G, S2A). The results suggest that RIS activation transients are required for sleep induction and that strong hyperpolarization prevents these transients. Mobility quiescence decreased with hyperpolarization and was increased with modest depolarization as expected. We anticipated that strong RIS depolarization would even further increase sleep. Surprisingly, however, mobility quiescence was completely abolished by strong RIS depolarization in most of the animals (Fig. 2H). These findings were additionally confirmed for a longer period of starvation up to day 12 (S2F). The loss of quiescence of the strong RIS depolarization strain was similar to, but not as strong as in *aptf-1(-)* (Fig. S2G). Mobility quiescence can thus be described as a parabolic function of RIS baseline calcium activity, with the wild type located in the rising flank of the parabolic curve. This indicates that there is a sensitive regulation of RIS-dependent sleep by modulation of RIS excitability (Fig. 2I). This parabolic relationship also suggests that there is an upper limit for RIS-induced sleep that might be near 50% of the time, which is consistent with the observation that RIS-dependent sleep typically occurs as bouts of quiescence that are regularly intermitted by bouts of wakefulness activity [10, 20, 35]. In summary, RIS polarization correlates with behavioral quiescence over a wide range of conditions, except for a very high level of RIS depolarization, which causes an apparent loss of sleep. This result is astonishing, as strong RIS depolarization promotes survival, which could be expected to depend on increased sleep. Surprisingly, strongly depolarized RIS leads to a combination of increased survival with a nearly complete loss of sleep behavior. Hence, the positive correlation between sleep behavior and survival that holds true for large ranges of RIS polarization inverses at high levels of RIS depolarization. These data thus indicate that RIS promotes sleep behavior and survival by different, separable mechanisms.

Constant and strong RIS depolarization prevents the shutdown of wake-active circuits

C. elegans sleep is characterized as a state of globally reduced nervous system activity [10, 18, 27]. To test how the RIS polarization status affects neuronal activity, we quantified

neuronal calcium activity in the presence of different RIS polarization levels. First, we
measured the overall neuronal activity by utilizing a strain that expresses GCaMP6s in all
neurons [36]. We measured both motion behavior and overall neuronal activity by
extracting the calcium activity signal of all neurons as one combined signal. As expected,
periods of behavioral quiescence were accompanied by strongly reduced global neuronal
activity (Fig 3A, B, S3A-F) [10, 18, 27]. This experimental setup allowed extracting phases
of neuronal inactivity as well as phases of behavioral inactivity. In all strains, neuronal
inactivity directly correlated with immobility, suggesting that behavioral quiescence and
neuronal quiescence are correlated in all strains irrespective of the level of RIS polarization
(Fig 3C). In addition, behavioral quiescence correlated in a parabolic manner with RIS
activity (Fig 3D). Thus, both behavioral inactivity and brain inactivity can serve as
equivalent readouts for detecting sleep in this system. Strong RIS hyperpolarization led to
an increased mean overall neuronal activity of 50.7% compared to wild type. Intermediate
strength of hyperpolarization did not affect mean overall neuronal activity. Strong RIS
depolarization caused a significant increase of mean overall neuronal activity of 35.7%
(Fig 3E). These results support the notion that worms with strongly depolarized RIS are in
a state of wakefulness for most of the time and cannot sleep.
201
To further characterize the effects of our RIS-manipulating transgenes on the neuronal
state, we looked at the activity of specific neurons. We focused on the RIM neurons, which
are second layer interneurons that integrate information from sensory and internal states to
control motion behaviors [37, 38]. Again, we used GCaMP imaging in microfluidic
chambers to correlate neuronal and behavioral activity. RIM showed activation transients
during phases of behavioral activity. During phases of behavioral quiescence, RIM activity
was strongly reduced and activation transients were lacking (Fig 3F, G, S3G-L). Similar to
the pan-neuronal measurements, the time fraction during which RIM was inactive
correlated with the fraction of motion quiescence (Fig 3H, I) and followed again a parabolic
distribution of RIS baseline activity (Fig 3I-J). Thus, sleep can be detected consistently in
the RIS polarization variants by both behavioral quiescence as well as neuronal activity
measurements, and strong RIS depolarization inhibits neuronal signatures that are typical
for sleep.

Bouts of behavioral quiescence occur during transients of RIS activation, and increasing RIS activity correlates with decreasing behavioral activity [10, 16]. The lack of RIS activation transients caused by the strong RIS depolarization transgene might thus cause a lack of behavioral quiescence. To test whether RIS depolarization transients can be triggered in the strongly RIS-depolarized transgenic strain, we stimulated RIS optogenetically. Optogenetic activation of RIS in the presence of the strongly depolarized RIS strain did not increase RIS activation or behavioral quiescence. This result is consistent with the view that strong levels of RIS depolarization prevent activation transients and thus suppress sleep bouts (Fig S3M-N). In summary, these results demonstrate that sleep amount is determined by the levels of RIS depolarization with both hyperpolarization and strong RIS depolarization causing sleep loss.

Both deletion and overexpression of *flp-11* inhibit sleep behavior

How can strong and constant RIS depolarization inhibit sleep? Constant RIS depolarization might cause an increased secretion of neurotransmitters from RIS, which in turn could lead to desensitization or habituation effects of the sleep-inducing system, ultimately leading to reduced sleep. FLP-11 is a neuropeptide of RIS that contributes to sleep induction [17]. Expression from the *flp-11* promoter depends on neuronal activity levels [39], and is increased in *RIS::unc-58gf(strong)* (Fig S2B). To directly test the role of FLP-11 in quiescence during L1 arrest, we measured the effects of overexpressing *flp-11* in RIS (*flp-11OE*) by using a multicopy transgenic extrachromosomal array that expresses FLP-11 from its own promoter, and compared the effects of *flp-11OE* with the effects of *flp-11* deletion (*flp-11(-)*) (Fig S4A-B) [17]. Both *flp-11OE* and *flp-11(-)* strongly reduced quiescence behavior, and where thus similar to *RIS::unc-58gf(strong)* (Fig 4A). *RIS::unc-58gf(strong)* still displayed a small fraction of quiescence behavior during L1 arrest. To test whether this residual quiescence was caused by *flp-11*, we combined *RIS::unc-58gf(strong)* with either *flp-11(-)* or *flp-11OE,* and quantified quiescence behavior. For this experiment, we deleted the entire *flp-11* gene in the *RIS::unc-58gf(strong)* operon locus. The sleep loss of *RIS::unc-58gf(strong)* was reduced further by both *flp-11(-)* and also by

flp-11OE, suggesting that the residual quiescence behavior in *RIS::unc-58gf* indeed

depends on *flp-11* (Fig 4A).

How does *flp-11OE* affect RIS activity? To answer this question, we first investigated

calcium activity of RIS in *flp-11OE* animals. RIS calcium activity stayed at a constant level

and did not show any activation transients in most of the animals that were tested (Fig 4B-

D). We next tested whether RIS can be optogenetically excited in *flp-11OE*. Unlike in wild-

type animals, optogenetic activation of RIS in *flp-11OE* animals did not increase calcium

activity (Fig 4E). In summary these experiments indicate that excess FLP-11 inhibits the

activation dynamics of RIS and the behavioral quiescence of sleep, suggesting that similar

effects might underly inhibition of sleep in *RIS::unc-58gf(strong)*.

FLP-11 neuropeptides support survival

flp-11 is required for most of the sleep during L1 arrest. We hence tested whether FLP-11

neuropeptides are also required for survival. Deletion of *flp-11* reduced survival by 24.7%,

suggesting that *flp-11* contributes to survival (Fig. 4F). To test whether *flp-11* is required

for increased survival caused by *RIS::unc-58gf(strong)*, we measured survival in the

RIS::unc-58gf(strong), flp-11(-) strain. *flp-11(-)* suppressed almost the entire survival

extension caused by *RIS::unc-58gf(strong)* (Fig. 4F). We next tested the effects of *flp-*

11OE on survival. *flp-11OE* did not significantly change survival, and it efficiently

suppressed the extended survival of *RIS::unc-58gf(strong)* (Fig. 4G). Consistent with the

survival assays, the recovery of larvae from the arrested first larval stage was significantly

reduced by *flp-11(-)*, yet unaffected by *flp-11OE* (Fig S4C-D). Together these experiments

indicate that *flp-11* plays an important role in survival and recovery of L1 arrest both in the

wild type and in *RIS::unc-58gf(strong)*.

These results suggest that *flp-11* plays two roles, one role is to promote sleep and the other

is to promote survival. *flp-11OE* uncouples those two functions of *flp-11*. Hence, *flp-11OE*

presents a condition in which, similar to *RIS::unc-58gf(strong)*, the animals survive well

without much sleep. Deletion of *flp-11* blunts survival less than strong RIS
hyperpolarization, suggesting that additional neurosecretory factors might also contribute
to survival. As RIS expresses multiple neuropeptides and also GABA [17, 19, 23], it may
use multiple transmitters to control survival. In summary these data suggest that FLP-11
neuropeptides are the major transmitters released by RIS to promote both sleep as well
survival during L1 arrest.

Discussion

Sleep-active neurons inhibit wakefulness to cause sleep and promote survival [30, 31]. A
parsimonious explanation linking these two observations would be that sleep promotes
survival, for example through energy conservation or behavioral optimization [13]. Our
data, however, suggest that the sleep-active RIS neuron promotes survival rather
independently of sleep. This suggests that RIS has two parallel functions. The first function
is the induction of sleep, a state of reduced nervous system activity and behavioral
quiescence. The reduction of brain and muscle activity likely contributes to survival by
conserving energy. While energy consumption appears to be playing a role in survival,
behavioral activity and thus energy consumption does not correlate with survival when
transmission from RIS is strongly increased. In strongly depolarized *RIS::unc-58gf(strong)*
animals, neuronal and behavioral activity are high and consume energy, yet survival is
increased. Similarly, in *flp-11OE*, behavioral activity is as high as in RIS loss-of-function
mutations, which should cause high energy expenditure, yet the survival is normal. Hence
RIS appears to be able to promote survival rather independently of sleep behavior and
associated energy conservation. As both sleep and survival are functions of RIS
depolarization over a wide range of conditions, the two processes are normally coupled
during sleep. During each sleep bout, RIS depolarizes and thus promotes both sleep as well
as survival. This coupling of sleep and survival processes might ensure that phases of brain
and behavioral inactivity coincide with phases during which survival-promoting processes
are active. This tight coupling might serve to optimize energy expenditure, as energy
conserved by sleep could be directly allocated to survival-promoting processes [12].

Both, the induction of sleep and the promotion of survival by RIS depend on FLP-11

neuropeptides, which are released upon depolarization. Hence the role of FLP-11 on

survival could include the conservation of energy during sleep plus an additional survival

component that is independent of sleep. The effects of FLP-11 on sleep and survival can

be separated by their different response to increased FLP-11 activity. The two functions of

FLP-11 might separate mechanistically at the level of divergent receptors, signaling

pathways or cell types downstream of FLP-11 that mediate either the sleep or the survival

response.

By controlling the polarization state of RIS, we showed that the survival function and the

sleep function of this sleep neuron can be decoupled. By strongly depolarizing RIS, we

generated a condition in which wakefulness neural activity and sleep-active neuron

depolarization coexist. The coupling of sleep and survival functions by sleep-active

neurons also could provide a starting point for treating the consequences of insomnia in

humans. Monitoring and managing the activity of sleep-active neurons might be key to not

only restore sleep but also to promote essential physiological functions.

Figures

Fig. 1 – RIS depolarization promotes survival

A) Schematic representation of the genetic design of the different RIS depolarization

transgenes.

B) Quantification of the expression levels of UNC-58gf::mKate2 in the two RIS depolarization strains. ***p<0.001, Welch test.

C) Localization and expression of the strong depolarization strain (*RIS::unc-58gf(strong)*).

D) Localization and expression of the weak depolarization strain (*RIS::unc-58gf(weak)*).

E) Scheme of the genetic design of the different RIS hyperpolarization transgenes.

F-H) Localization and expression of mKate2, which is translationally fused to the ion channels, in the different hyperpolarization strains.

I) Ion channel expression in RIS results in a dose-response strain set with varying levels of RIS baseline activity. n.s. p>0.05, **p<0.001, ***p<0.001, Welch test with FDR correction for multiple testing.

J) Strong hyperpolarization significantly decreases survival whereas strong depolarization prolongs survival (three-parameter logistic fit, see also supplementary table S1, Fisher's Exact Test was conducted when 50% of wild-type worms were alive (day 22)). The p-values were FDR corrected with Benjamini-Hochberg procedure with a 5% false discovery rate. The plot includes data from three replicates. ***p<0.001.

K) Survival time increases with mean RIS baseline activity from a minimum median of about 10 days to a maximum median of about 25 days (Belehradek fit, timepoints when 50% of worms of a strain were still alive are plotted for all three replicates).

Fig. 2 – Both, strong hyperpolarization as well as strong depolarization of RIS

prevent sleep behavior

A-F) Ion channel expression in RIS controls calcium activity and sleep behavior.

Mobility quiescence bouts (grey shade) correlate with RIS transients A) RIS-

hyperpolarized strain *RIS::twk-18gf* and B) RIS-hyperpolarized strain *RIS::egl-*

23gf(strong) larvae do not have detectable mobility quiescence bouts or RIS transients.

C) The weakly RIS hyperpolarized strain *RIS::egl-23gf(weak)* has reduced mobility

quiescence bouts. D) Wild-type RIS activation transients and immobility bouts. E) The

weakly RIS-depolarized strain *RIS::unc-58gf(weak)* spends more time in mobility

quiescence bouts. F) The strongly RIS depolarized strain *RIS::unc-58gf(strong)* has no

detectable mobility quiescence bouts and no RIS transients but a high constant activity.

G) RIS activation and immobility are correlated. Alignment of RIS activity and speed to

mobility quiescence bout onset. **p<0.01, ***p<0.001, Wilcoxon signed rank test.

H-I) Both, strong hyperpolarization as well as strong depolarization of RIS prevent sleep

behavior. I) mobility quiescence is a function of RIS activity levels n.s. p>0.05,

**p<0.001, ***p<0.001, Welch test with FDR correction for multiple testing. J) second-

degree polynomial fit for mobility quiescence fraction against mean RIS baseline activity.

Most of the channel transgenes are expressed as an operon with *flp-11*, which should not

grossly impair expression of the *flp-11* neuropeptide gene that is crucial for quiescence

induction [17, 19]. The correlation of quiescence with RIS polarization levels and the

preservation of most sleep bouts in the weak *egl-23* knock-in transgene support this

notion.

Fig. 3 – Constant strong RIS depolarization fails to inhibit wake-active circuits

A-B) Reduced global neuronal activity (grey shading) correlates with motion quiescence

(supplementary movies S1-S6). A) sample trace, B) alignment of overall neuronal

activity to mobility quiescence bout onset. ***p<0.001, Wilcoxon signed rank test.

C) The mean neuronal inactivity fraction correlates with the mean mobility quiescence

fraction (linear fit) for all strains.

D) Both, strongly hyperpolarized as well as strongly depolarized strains spend only a

small fraction of time in states of neuronal inactivity (parabolic fit).

E) Strong RIS hyper- as well as depolarization caused an overall increase of neuronal

activity. n.s. $p > 0.05$, *$p<0.05$, **$p<0.01$, ***$p < 0.001$, Welch test with FDR correction

for multiple testing.

F-G) RIM does not activate during mobility quiescence bouts. F) sample trace (grey

shade is RIM inactivity bout). G) Alignment of RIM activity and speed to mobility

quiescence bout onset. *$p<0.05$, **$p<0.01$, ***$p<0.001$, Wilcoxon signed rank test.

H) The mean RIM inactivity fraction correlates with the mean mobility quiescence

fraction (linear fit) for all strains.

I) Both, strongly hyperpolarized as well as strongly depolarized strains spend only a

small fraction of time in states of RIM inactivity (parabolic fit).

J) Strong RIS hyper- as well as depolarization caused an overall increase of RIM activity.

**$p < 0.01$, ***$p < 0.001$, Welch test with FDR correction for multiple testing.

Fig 4 – FLP-11 controls sleep and survival during L1 arrest

A) Deletion as well as overexpression of the neuropeptide gene *flp-11* causes loss of

sleep. The strongly RIS depolarized worms (*RIS::unc-58gf(strong)*) sleep even less when

flp-11 is deleted or overexpressed. ***p<0.001, Welch test with FDR correction for

multiple testing.

B) Sample trace of RIS activity and speed of a wild type worm.

C) Sample trace of RIS activity and speed of a worm overexpressing *flp-11*.

D) *flp-11OE* worms do not show many RIS peaks. ***p<0.001, Welch test.

E) Optogenetic activation of RIS causes a strong RIS activity increase and reduction of

speed in the wild type, but there is no response in the *flp-11OE* strain. n.s. p>0.05,

***p<0.001, Wilcoxon signed rank test.

F) Loss of the neuropeptide FLP-11 leads to strongly reduced survival (three-parameter

logistic fit, see also supplementary table S1, Fisher's Exact Test was conducted on day

18) for comparisons when *flp-11(-)* was the shortest lived condition, day 20) when

RIS::unc-58gf(strong), flp-11(-) was the shortest lived condition or day 24) when wild

type was the shortest lived condition. The p-values were FDR corrected with Benjamini-

Hochberg procedure with a 5% false discovery rate. The plot includes data from three

replicates. ***p<0.001.

G) Overexpression of FLP-11 does not reduce survival (three-parameter logistic fit, see

also supplementary table S1. Fisher's Exact Test was conducted on day 23). The p-values

were FDR corrected with Benjamini-Hochberg procedure with a 5% false discovery rate.

The plot includes data from three replicates. *p<0.05, ***p<0.001.

Materials and Methods

Resource and Materials Availability

Key *C. elegans* lines that were used in this project are available at the Caenorhabditis

elegans Genetics Center. Other *C. elegans* strains, plasmids, or additional reagents

generated for this study are available upon reasonable request.

Data and Code Availability

Raw and processed data as well as MATLAB scripts will be made publicly available.

Experimental model and subject details

C. elegans hermaphrodites were maintained at 20°C. They were kept on Nematode Growth

Medium Plates (NGM). The *Escherichia coli* strain OP50 served as a food source during

maintenance [40].

Method Details

Creation of transgenic animals

Crossing *C. elegans* strains

Males were generated via heat shock or maintained by a continuation of crossings of

worms of the same genotype [41]. Single worms of the F2 generation were genotyped

using duplex PCR genotyping [42]. For crossing fluorescent transgenes, fluorescent

markers were used. All strains and primers of this study can be found in the

supplementary tables S2 and S3.

Creating mutants with the CRISPR/Cas9 system

We designed the sequences of the following transgenes in silico. Transgenes were

synthesized and inserted into the genome by using CRISPR/Cas9 by SunyBiotech.

Transgene sequences were verified through Sanger sequencing.

PHX1433   *flp-11(syb1433 [flp-11-SL2-egl-23cDNA(A383V)-linker-mKate2]) X*

PHX1445   *aptf-1(gk794) II; flp-11(syb1445[flp-11-SL2-unc-58(L428F)-linker-*
456   *mKate2]) X*

457 PHX1464   *flp-11(syb1464[flp-11-SL2-egl-23cDNA(L229N)-linker-mKate2]) X*

458 PHX2193   *flp-11(syb2193[flp-11b-SL2(gpd-2)-mKate2-linker-twk-18(e1913)]) X*

459 PHX2493   *lgc-38(syb2346[flp-11p::dpy-10 site::flp-11 3'UTR], syb2493[ReaChR-*
460   *linker-mKate2]) III.*

461 PHX3190   *lgc-38(syb2346[flp-11p::dpy-10 site::flp-11 3'UTR] ,syb3190[unc-*
462   *58(e665)-linker(GSGSGSGSG)-mKate2]) III*

463 PHX4110   *lgc-38(syb2346[flp-11p::dpy-10 site::flp-11 3'UTR], syb4110[unc-58gf-*
464   *CAI-1.0-linker(GSGSGSGSG)-mKate2]) III*
465 PHX4416   *aptf-1(gk794) II; flp-11(syb1445 syb4416) X*
466

The coding sequences of the ion channel transgenes were synthesized and consisted each
of the spliced coding sequence of the ion channel including the mutation that caused the
leak current [32], a C-terminal linker sequence followed by a C-terminal mKate2 gene.
The coding sequence of mKate2 was codon optimized and intronized [43]. The coding
sequences of the ion channels were not codon optimized to allow for proper folding of
the membrane proteins. We tried codon-optimization [43] of the *unc-58(L428F)* channel
gene and mKate2 to achieve high expression, but this also caused aggregation of the
fluorescent fusion protein (Fig S1A,D). Hence, we used the endogenous gene sequence
for the channel and only codon-optimized mKate2.

Analysis of the RIS transcriptome showed that *flp-11* has the strongest and most specific
promoter of all RIS-expressed genes, with 157-fold higher expression in RIS than in other
neurons [23]. *flp-11* is also expressed in other neurons, albeit at low levels [17, 23, 44].
Using imaging settings that cause oversaturation of the mKate2 signal in RIS revealed that
there is a weak expression of *unc-58gf::mKate2* in a few neurons in the ventral cord in
RIS::unc-58gf(strong) (Fig S1F). Expression from the *flp-11* promoter in RIS but not in
other neurons depends on APTF-1 [17]. We hence imaged mKate2 expression in *aptf-1(-*
). Consistent with previous results [17], mKate2 expression was abolished in RIS but was
still present in the ventral cord (Fig S1G). Thus, expression from the *flp-11* promoter is
strong, has a high level of RIS specificity, and RIS expression depends on *aptf-1*.
486

The sequences of the ion channels of the strains PHX1433, PHX1445, PHX1464 and
PHX2193 were inserted after the stop codon of the *flp-11* endogenous coding sequence
together with an SL2 site. Thus, an operon was created that expressed both *flp-11* and the ion
channel gene from the *flp-11* promoter. The transgenes of PHX2493, PHX3190 and
PHX4110 were generated via a single-copy knock-in loci for defined gene expression (SKI
LODGE) system [33]. For the SKI LODGE expression in RIS we first inserted the minimal

flp-11 3' and 5'UTRs separated by a *dpy-10* sgRNA cutting site. The expression from these

minimal UTR sequences allowed specific expression in RIS, but the expression levels were

lower compared with the endogenous *flp-11* locus (Fig 1A-D, S1A-D). The final sequences

of all transgenes can be found in the supplementary Text S1.

498   Imaging

500   Microfluidic devices

Worms were filmed in microfluidic devices made from agarose hydrogel as previously

described [45, 46]. In brief, a PDMS mold was used to cast box-shaped indentations from

melted 5% agarose diluted in M9 buffer. After agarose hydrogel solidification the stamp was

removed resulting in hundreds of microchambers sized 110x110x10μm per agarose slab.

Depending on the experiment, approximately 9-20 pretzel-stage eggs per strain were then

transferred without food into the indentations in the hydrogel and the chambers were closed

with a coverslip and kept in a temperature-controlled dish. Hatching of the larvae in the absence

of food caused L1 arrest. Chambers were timed so that the worms were incubated for 48h at

25°C before imaging. The number of biological replicates (n) imaged in microfluidic devices

is stated in each figure. There were two or more technical replicates (different microfluidic

devices) imaged for each experiment.

514   Software

516   All image acquisition was performed with Nikon NIS Elements 5 software (Nikon, Tokyo).

518   Dark Differential Contrast (DIC) Imaging

519   Long term dark differential contrast (DIC) images were acquired with a standard 100W

520   halogen lamp and an infrared filter (Semrock Brightline HC 785/62) (Idex Health and Science,

521   New York). The images were taken 48h upon starvation for 12h with a frame rate of 0.1Hz.

522   A 20x objective for imaging was combined with a 0.7 lens.

524   Functional Ca^{2+}-imaging

For calcium imaging we used either a back-illuminated sCMOS camera (Photometrics Prime
95B, 1200x1200 pixels) (Figure 1I, 2, S2A-C) or an Andor iXon EMCCD camera (512x512
pixels, Andor Technology Ltd., Belfast). (Figure 3, 4B-D, S3A-L). The gain of the iXon
camera was set to 200. The iXon camera's read out mode was 10MHz at 14-bit. The images
of the Photometrics camera were 2 x 2 binned.

A 20x objective was used to simultaneously image 3 x 3 adjacent microchambers containing
one worm each (This configuration was used to obtain data for Figures 3, 4B-D, S3A-L). A
10x objective combined with a 1.5x lens was used to simultaneously image 6 x 6 adjacent
microchambers containing one worm each (this configuration was used to obtain data for
Figures 1I, 2, S2A-C).

For calcium imaging we utilized a 480nm LED and a standard GFP filter set (EGFP,
Chroma). The EMCCD camera TTL signal triggered the LED. For data in Figures 1I, 2 and
S2A-B the 490nm intensities were set to either 0.23mW/mm^2 with an exposure time of 50ms
(Figure 1I, 2, S2A-B), 0.33mW/mm^2 (Figure 3A) or 0.51 mW/mm^2 (Figure 3B), both with an
exposure time of 5ms. Additional mKate2 fluorescence images were acquired for the RIS and
RIM activity experiments (Figure 1I, 2, 3B S2A, S3G-L). For this measurement we used a
585nm LED and a set of standard Texas Red filters (Chroma). The 585nm LED intensity was
set to 0.11mW/mm^2 while imaging with a 50ms exposure time (Figure 1I, 2, S2A, B) and to
0.15 mW/mm^2 for acquiring images with a 100ms exposure time (Figure 3F-J). Calcium and
mKate2 imaging were carried out with a frame rate of 0.1Hz for 3h. As the ion channels were
additionally tagged with an mKate2, the parental strains without the GCaMP/mKate2
construct were imaged with the same settings with the 585nm LED. Here, no signal could be
observed in the parental strains.
For the long-term calcium imaging from day 1 to day 12 we utilized slightly different LED
light intensities. Here, the 490nm intensity was set to 0.34mW/mm^2 and the 585nm light had
an intensity of 0.21mW/mm^2.
RIS in the *flp-11OE* strain could only be imaged for GCaMP as the overexpression construct
carried mKate2, which caused an extremely bright signal that did not allow for normalization
of the GCaMP signal. A 490nm light of 0.15mW/mm^2 intensity was utilized.

Optogenetic activation

For optogenetic RIS activation experiments, the agarose containing the microchambers was
supplemented with 10µl of 10mM all-trans-retinal (ATR, Sigma Aldrich) 1h before the start of
image acquisition. Control worms and worms before the stimulation did not receive ATR
treatment.

In experiment in Fig S3M-N, A baseline of GCaMP images was recorded for 1min with a
frame rate of 0.33Hz. This was followed by a stimulation phase of 1min, in which again
GCaMP images were acquired with a frame rate of 0.33Hz. In between the camera exposures,
the worms were illuminated with 585nm LED light with an intensity of $0.17mW/mm^2$ for
800ms triggered by the software. After the stimulation period, a recovery period of 1min was
imaged, in which GCaMP images were acquired with a frame rate of 0.33Hz. The 490nm LED
intensities for imaging were set to $0.39mW/mm^2$ and images were acquired with an exposure
time of 5ms.

In the optogenetic experiment in Fig 4E the worms were first imaged without ATR for 5min.
With a 490nm LED of $0.36mW/mm^2$ intensity, GCaMP was imaged with a frame rate of 0.2Hz
and exposure time of 50ms. mKate2 images were additionally acquired with a 585nm LED
with an intensity of $0.25mW/mm^2$ for 50ms all 5s. After the pre stimulation phase, 10µl of
10mM all-trans-retinal (ATR, Sigma Aldrich) was added to the worms and the imaging
protocol was then repeated after 1h to allow for sufficient time for the worms to ingest the
ATR.

Expression of mKate2 of the RIS activity strains

Larvae were synchronized via bleaching [23] and hatched in M9 buffer at 25°C. Imaging of
most strains was performed after 48h in the L1 arrest stage by transferring worms with a 10
µl liquid drop onto an agarose pad (5% agarose solved in M9 buffer). The *flp-11OE* and
RIS::unc-58gf, flp-11(-) worms were imaged 24h after bleaching. Larvae were then fixed
through the addition of 10 µl of 25mM levamisole (Sigma Aldrich).

To image the expression levels of the different RIS activity transgenes, mKate2 was imaged

with an Andor Revolution disc system (Andor Technology Ltd.) and a CSU-X1 spinning disc

head (Yokogawa). The 561nm laser light intensity was set to 0.49mW/mm^2. The worms were

imaged through a 100x oil objective and by an Andor iXon Ultra EMCCD camera

(1024x1024 pixels) (Andor Technology Ltd., Belfast). An EM gain of 200 was included and

the camera read-out time was set to 30MHz. A z-stack of 21planes, each 0.5μm apart (10μm

in total) was acquired. For comparison, a maximum z-stack projection was calculated.

Image analysis

DIC images analysis

601 Mobility of worms was analyzed from DIC images though image subtraction [47]. The matrix

602 of intensity values of the preceding image was subtracted from the matrix of intensity of the

603 following image. A mean of all values of the difference matrix was then calculated to attain a

604 mean smoothed image subtraction value, which represents the movement of the worm.

Analyzing functional Ca^{2+}-images

Neuronal intensities were automatically analyzed via custom-written MATLAB scripts.

These codes extracted the regions of interest through an intensity threshold and acquired

the intensities and position data. The extracted regions were set to be slightly bigger than

the actual size of the neuron to allow for background intensity detection and background

subtraction. The pan-neuronal activity was measured treating the head neurons as a

combined signal. The RIS GCaMP activity in Figure 2 was determined by extracting the

position data of the corresponding mKate2 signal. A larger region of interest was extracted

and a fitting percentage of that was calculated as the signal. The extracted position data of

the signal served to calculate the speed values.

RIS and RIM activities were normalized by dividing GCaMP intensities through

corresponding mKate2 intensities. In the optogenetic experiment, the baseline

measurements were used for normalization.

Mobility quiescence detection

First, speeds or image subtraction mean intensities were smoothed over 30 time points
through the *smooth* function in MATLAB, which is a first-degree polynomial local
regression model. Smoothed speeds and image subtraction values were normalized
between 0 and 1 for each individual animal with 0 being the lowest speed value and 1 the
highest speed value the animal possessed. A mobility quiescence bout was counted when
the normalized speed was below a specific percentage for a certain amount of time. For
calcium imaging experiments with the 20x objective (Figure 3), the normalized speed had
to be below 10% for at least 3min. The RIS activity experiment, which was filmed with a
10x objective, required a percentage threshold of 5% for a minimum of 4min. For DIC
images filmed with the 20x objective, the normalized mean image subtraction values had
to be below 15% for at least 3min to be counted as sleep time points.

Mobility quiescence bout onset alignment

Speeds and GCaMP intensities were aligned to the start of mobility quiescence bouts, which
had been detected by the mobility quiescence bout analysis. Mobility quiescence bouts were
only counted for the alignment when the worm was mobile throughout 3min prior to the
mobility quiescence bout onset. This alignment was carried out with a custom-written
MATLAB script.

Neuronal inactivity detection

The neuronal activities were treated similarly as speed values for the mobility quiescence
detection. However, the nervous system was counted as inactive when it was below 20%
of the smoothed and normalized activity for at least 3min.

RIM neuron inactivity detection

RIM peaks were detected through custom-written MATLAB scripts. First, the activity values
were smoothed using the inbuilt *smooth* function over 5 time points. Then, peaks of a
minimum prominence of 0.2 were extracted via the inbuilt function *islocalmax*. RIM was
counted as inactive when there were no detected peaks for at least 5min.

Wide RIS peak extraction

To quantify RIS transients, a custom-written MATLAB script was utilized. The in-build

function *findpeaks* detected peaks with a minimum peak width of 9 frames, which

corresponds to peak time of 1.5min.

Survival assay

Worms were synchronized by bleaching [23]. 3-4 days before bleaching, 10-12 L4 stage

worms were transferred onto NGM plates with a diameter of 6cm. 1-2 plates were bleached

per strain and the day of bleaching was defined as day 0. The following day was counted

as day 1 of the lifespan. During the survival assay, the worms were kept in liquid culture

in 1ml M9 buffer in 2ml Eppendorf tubes. This resulted in a population density of

approximately 100 worms per 10μl. The tubes were placed in a rotator at 20°C, which

rotated at 20rpm. For counting, 2x5μl drops of the liquid culture were pipetted onto fresh

NGM plates for each strain respectively. The worms were allowed to recover for at least 2h

after pipetting before counting. Counting was performed manually with the assistance of a

clicker. The recovery number, i.e., the number of worms that developed into at least the L4

stage, was counted 3-5 days after pipetting. Around 50-100 worms were counted for each strain

of each replicate on each time point except for the *flp-11OE* strains. As the *flp-11OE* strains

were extra chromosomal with medium transmission rate only 20-60 worms could be counted

for each strain on one day in each replicate.

Fitting

Origin software was used for the regression analyses. A second-degree polynomial was

fitted to the data in Figure 2I, 3D and 3I. A linear regression was fitted to the data in Figure

3C and 3H. For data in Figure 1K a Belehradek fit was conducted. R^2 values can be found

in the corresponding figures. The exact functions are in Table 1.

Figure panel	fit	function
1K	Belehradek fit	$y = 28.693 * (x - 0.97774)^{0.14966}$
2I	2nd degree polynomial	$y = -5.7210 + 9.2995x - 3.5059x^2$
3C	linear	$y = 0.116 + 0.5494x$

3D	2nd degree polynomial	$y = -3.067 + 5.163x - 1.95x^2$
3H	linear	$y = 0.21203 + 1.10949x$
3I	2nd degree polynomial	$y = -6.38841 + 10.8x - 4.1x^2$

680   Table 1. Calculated functions for the regressions.

Quantification and Statistical Analysis

The experimenter was blind to the genotype information during the lifespan experiments.
All other experiments relied on an automated analysis and hence were not blinded. The
Wilcoxon signed rank test was utilized for the comparison of speeds and GCaMP intensities
within the same strain. The Welch test was conducted to compare data from different strains.
The survival assay experiments were statistically analyzed by utilizing the Fisher's exact test
at 50% survival of the wild type. Experiments with four or more genotypes were FDR corrected
by the Benjamini-Hochberg procedure with a 5% false discovery rate. The graphs depict the
mean and standard error of the mean. Box plots display individual data points, the interquartile
range and the median. The whiskers represent the 10-90th percentile.

Acknowledgements

We thank the *Caenorhabditis* Genetics Center supported by the National Institutes of
Health Office of Research Infrastructure Programs (P40 OD010440) for strains. The
CRISPR strains were generated by SunyBiotech based on our design. We thank Thomas
Boulin for communicating the sodium permeability of *unc-58gf* and suggesting its use as a
tool for neuronal depolarization. This work was supported by a European Research Council
Starting Grant (ID: 637860, SLEEPCONTROL).

Author contributions

702   I.B. designed, executed and analyzed all experiments. H.B. conceived the overall structure
703   of the study, designed the transgenes, acquired funding and supervised the project. All
704   authors wrote the paper.

Declaration of Interest

707   The authors declare no competing interests.

References

1.   Campbell SS, Tobler I. Animal sleep: a review of sleep duration across phylogeny. Neurosci Biobehav Rev. 1984;8(3):269-300. Epub 1984/01/01. PubMed PMID: 6504414.
2.   Krueger JM, Frank MG, Wisor JP, Roy S. Sleep function: Toward elucidating an enigma. Sleep medicine reviews. 2016;28:46-54. Epub 2015/10/09. doi: 10.1016/j.smrv.2015.08.005. PubMed PMID: 26447948; PubMed Central PMCID: PMC4769986.
3.   Anafi RC, Kayser MS, Raizen DM. Exploring phylogeny to find the function of sleep. Nature reviews Neuroscience. 2019;20(2):109-16. Epub 2018/12/24. doi: 10.1038/s41583-018-0098-9. PubMed PMID: 30573905.
4.   Diekelmann S, Born J. The memory function of sleep. Nature reviews Neuroscience. 2010;11(2):114-26. Epub 2010/01/05. doi: 10.1038/nrn2762. PubMed PMID: 20046194.
5.   Van Cauter E, Spiegel K, Tasali E, Leproult R. Metabolic consequences of sleep and sleep loss. Sleep Med. 2008;9 Suppl 1:S23-8. Epub 2008/12/17. doi: 10.1016/S1389-9457(08)70013-3. PubMed PMID: 18929315; PubMed Central PMCID: PMCPMC4444051.
6.   Irwin MR. Why sleep is important for health: a psychoneuroimmunology perspective. Annu Rev Psychol. 2015;66:143-72. Epub 2014/07/26. doi: 10.1146/annurev-psych-010213-115205. PubMed PMID: 25061767; PubMed Central PMCID: PMCPMC4961463.
7.   Hublin C, Partinen M, Koskenvuo M, Kaprio J. Sleep and mortality: a population-based 22-year follow-up study. Sleep. 2007;30(10):1245-53. Epub 2007/11/01. doi: 10.1093/sleep/30.10.1245. PubMed PMID: 17969458; PubMed Central PMCID: PMCPMC2266277.
8.   Saper CB, Fuller PM, Pedersen NP, Lu J, Scammell TE. Sleep state switching. Neuron. 2010;68(6):1023-42. Epub 2010/12/22. doi: 10.1016/j.neuron.2010.11.032. PubMed PMID: 21172606; PubMed Central PMCID: PMC3026325.
9.   Bringmann H. Sleep-Active Neurons: Conserved Motors of Sleep. Genetics. 2018;208(4):1279-89. Epub 2018/04/06. doi: 10.1534/genetics.117.300521. PubMed PMID: 29618588; PubMed Central PMCID: PMC5887131.
10.   Wu Y, Masurat F, Preis J, Bringmann H. Sleep Counteracts Aging Phenotypes to Survive Starvation-Induced Developmental Arrest in C. elegans. Curr Biol. 2018;28(22):3610-24 e8. Epub 2018/11/13. doi: 10.1016/j.cub.2018.10.009. PubMed PMID: 30416057; PubMed Central PMCID: PMCPMC6264389.
11.   Sinner M, Masurat F, Ewbank J, Pujol N, Bringmann H. Antimicrobial peptides signal need for sleep from peripheral wounds to the nervous system. bioRxiv. 2020:2020.07.02.183806. doi: 10.1101/2020.07.02.183806.
12.   Schmidt MH. The energy allocation function of sleep: a unifying theory of sleep, torpor, and continuous wakefulness. Neurosci Biobehav Rev. 2014;47:122-53. Epub 2014/08/15. doi: 10.1016/j.neubiorev.2014.08.001. PubMed PMID: 25117535.

13.   Siegel JM. Sleep viewed as a state of adaptive inactivity. Nature reviews
Neuroscience. 2009;10(10):747-53. Epub 2009/08/06. doi: 10.1038/nrn2697. PubMed
PMID: 19654581.
14.   Krause AJ, Simon EB, Mander BA, Greer SM, Saletin JM, Goldstein-Piekarski
AN, et al. The sleep-deprived human brain. Nature reviews Neuroscience.
2017;18(7):404-18. Epub 2017/05/19. doi: 10.1038/nrn.2017.55. PubMed PMID:
28515433.
15.   Klinzing JG, Niethard N, Born J. Mechanisms of systems memory consolidation
during sleep. Nat Neurosci. 2019;22(10):1598-610. Epub 2019/08/28. doi:
10.1038/s41593-019-0467-3. PubMed PMID: 31451802.
16.   Turek M, Lewandrowski I, Bringmann H. An AP2 transcription factor is required
for a sleep-active neuron to induce sleep-like quiescence in C. elegans. Curr Biol.
2013;23(22):2215-23. Epub 2013/11/05. doi: 10.1016/j.cub.2013.09.028. PubMed PMID:
24184105.
17.   Turek M, Besseling J, Spies JP, Konig S, Bringmann H. Sleep-active neuron
specification and sleep induction require FLP-11 neuropeptides to systemically induce
sleep. eLife. 2016;5. Epub 2016/03/08. doi: 10.7554/eLife.12499. PubMed PMID:
26949257; PubMed Central PMCID: PMC4805538.
18.   Nichols ALA, Eichler T, Latham R, Zimmer M. A global brain state underlies C.
elegans sleep behavior. Science. 2017;356(6344). Epub 2017/06/24. doi:
10.1126/science.aam6851. PubMed PMID: 28642382.
19.   Steuer Costa W, Van der Auwera P, Glock C, Liewald JF, Bach M, Schuler C, et
al. A GABAergic and peptidergic sleep neuron as a locomotion stop neuron with
compartmentalized Ca2+ dynamics. Nat Commun. 2019;10(1):4095. Epub 2019/09/12.
doi: 10.1038/s41467-019-12098-5. PubMed PMID: 31506439.
20.   Maluck E, Busack I, Besseling J, Masurat F, Turek M, Busch KE, et al. A wake-
active locomotion circuit depolarizes a sleep-active neuron to switch on sleep. PLoS
biology. 2020;18(2):e3000361. Epub 2020/02/23. doi: 10.1371/journal.pbio.3000361.
PubMed PMID: 32078631.
21.   Busack I, Jordan F, Sapir P, Bringmann H. The OptoGenBox - a device for long-
term optogenetics in C. elegans. J Neurogenet. 2020:1-9. Epub 2020/06/17. doi:
10.1080/01677063.2020.1776709. PubMed PMID: 32543249.
22.   Sinner MP, Masurat F, Ewbank JJ, Pujol N, Bringmann H. Innate Immunity
Promotes Sleep through Epidermal Antimicrobial Peptides. Curr Biol. 2021;31(3):564-77
e12. Epub 2020/12/02. doi: 10.1016/j.cub.2020.10.076. PubMed PMID: 33259791.
23.   Konietzka J, Fritz M, Spiri S, McWhirter R, Leha A, Palumbos S, et al. Epidermal
Growth Factor Signaling Promotes Sleep through a Combined Series and Parallel Neural
Circuit. Curr Biol. 2020;30(1):1-16 e3. Epub 2019/12/17. doi:
10.1016/j.cub.2019.10.048. PubMed PMID: 31839447.
24.   Gonzales DL, Zhou J, Fan B, Robinson JT. A microfluidic-induced C. elegans
sleep state. Nat Commun. 2019;10(1):5035. Epub 2019/11/07. doi: 10.1038/s41467-019-
13008-5. PubMed PMID: 31695031.
25.   Makino M, Ulzii E, Shirasaki R, Kim J, You YJ. Regulation of Satiety
Quiescence by Neuropeptide Signaling in Caenorhabditis elegans. Front Neurosci.
2021;15:678590. Epub 2021/08/03. doi: 10.3389/fnins.2021.678590. PubMed PMID:
34335159; PubMed Central PMCID: PMCPMC8319666.

26.   Xu Y, Zhang L, Liu Y, Topalidou I, Hassinan C, Ailion M, et al. Dopamine
receptor DOP-1 engages a sleep pathway to modulate swimming in C. elegans. iScience.
2021;24(4):102247. Epub 2021/04/03. doi: 10.1016/j.isci.2021.102247. PubMed PMID:
33796839; PubMed Central PMCID: PMCPMC7995527.
27.   Skora S, Mende F, Zimmer M. Energy Scarcity Promotes a Brain-wide Sleep
State Modulated by Insulin Signaling in C. elegans. Cell reports. 2018;22(4):953-66.
Epub 2018/02/02. doi: 10.1016/j.celrep.2017.12.091. PubMed PMID: 29386137.
28.   Kenyon CJ. The genetics of ageing. Nature. 2010;464(7288):504-12. Epub
2010/03/26. doi: 10.1038/nature08980. PubMed PMID: 20336132.
29.   Baugh LR, Hu PJ. Starvation Responses Throughout the Caenorhabditis elegans
Life Cycle. Genetics. 2020;216(4):837-78. Epub 2020/12/04. doi:
10.1534/genetics.120.303565. PubMed PMID: 33268389; PubMed Central PMCID:
PMCPMC7768255.
30.   Weber F, Dan Y. Circuit-based interrogation of sleep control. Nature.
2016;538(7623):51-9. Epub 2016/10/07. doi: 10.1038/nature19773. PubMed PMID:
27708309.
31.   Bringmann H. Genetic sleep deprivation: using sleep mutants to study sleep
functions. EMBO Rep. 2019;20(3). Epub 2019/02/26. doi: 10.15252/embr.201846807.
PubMed PMID: 30804011; PubMed Central PMCID: PMCPMC6399599.
32.   Ben Soussia I, El Mouridi S, Kang D, Leclercq-Blondel A, Khoubza L, Tardy P,
et al. Mutation of a single residue promotes gating of vertebrate and invertebrate two-
pore domain potassium channels. Nat Commun. 2019;10(1):787. Epub 2019/02/17. doi:
10.1038/s41467-019-08710-3. PubMed PMID: 30770809; PubMed Central PMCID:
PMCPMC6377628.
33.   Silva-Garcia CG, Lanjuin A, Heintz C, Dutta S, Clark NM, Mair WB. Single-
Copy Knock-In Loci for Defined Gene Expression in Caenorhabditis elegans. G3
(Bethesda). 2019;9(7):2195-8. Epub 2019/05/09. doi: 10.1534/g3.119.400314. PubMed
PMID: 31064766; PubMed Central PMCID: PMCPMC6643881.
34.   Roux AE, Langhans K, Huynh W, Kenyon C. Reversible Age-Related
Phenotypes Induced during Larval Quiescence in C. elegans. Cell Metab.
2016;23(6):1113-26. Epub 2016/06/16. doi: 10.1016/j.cmet.2016.05.024. PubMed PMID:
27304510.
35.   Iwanir S, Tramm N, Nagy S, Wright C, Ish D, Biron D. The microarchitecture of
C. elegans behavior during lethargus: homeostatic bout dynamics, a typical body posture,
and regulation by a central neuron. sleep. 2013;36(3):385-95. Epub 2013/03/02. doi:
10.5665/sleep.2456. PubMed PMID: 23449971; PubMed Central PMCID: PMC3571756.
36.   Yemini E, Lin A, Nejatbakhsh A, Varol E, Sun R, Mena GE, et al. NeuroPAL: A
Multicolor Atlas for Whole-Brain Neuronal Identification in C. elegans. Cell.
2021;184(1):272-88 e11. Epub 2020/12/31. doi: 10.1016/j.cell.2020.12.012. PubMed
PMID: 33378642.
37.   Piggott BJ, Liu J, Feng Z, Wescott SA, Xu XZ. The neural circuits and synaptic
mechanisms underlying motor initiation in C. elegans. Cell. 2011;147(4):922-33. Epub
2011/11/15. doi: 10.1016/j.cell.2011.08.053. PubMed PMID: 22078887; PubMed Central
PMCID: PMC3233480.
38.   Li W, Kang L, Piggott BJ, Feng Z, Xu XZ. The neural circuits and sensory
channels mediating harsh touch sensation in Caenorhabditis elegans. Nat Commun.

2011;2:315. Epub 2011/05/19. doi: 10.1038/ncomms1308. PubMed PMID: 21587232;
PubMed Central PMCID: PMC3098610.
39.   Laurent P, Soltesz Z, Nelson GM, Chen C, Arellano-Carbajal F, Levy E, et al.
Decoding a neural circuit controlling global animal state in C. elegans. eLife. 2015;4.
Epub 20150311. doi: 10.7554/eLife.04241. PubMed PMID: 25760081; PubMed Central
PMCID: PMCPMC4440410.
40.   Brenner S. The genetics of Caenorhabditis elegans. Genetics. 1974;77(1):71-94.
Epub 1974/05/01. PubMed PMID: 4366476.
41.   Stiernagle T. Maintenance of C. elegans. WormBook. 2006:1-11. Epub
2007/12/01. doi: 10.1895/wormbook.1.101.1. PubMed PMID: 18050451.
42.   Ahringer A. Reverse genetics. WormBook, ed The C elegans Research
Community. 2006. doi: doi/10.1895/wormbook.1.47.1.
43.   Redemann S, Schloissnig S, Ernst S, Pozniakowsky A, Ayloo S, Hyman AA, et
al. Codon adaptation-based control of protein expression in C. elegans. Nat Methods.
2011;8(3):250-2. Epub 2011/02/01. doi: nmeth.1565 [pii]10.1038/nmeth.1565. PubMed
PMID: 21278743.
44.   Taylor SR, Santpere G, Weinreb A, Barrett A, Reilly MB, Xu C, et al. Molecular
topography of an entire nervous system. Cell. 2021. Epub 2021/07/09. doi:
10.1016/j.cell.2021.06.023. PubMed PMID: 34237253.
45.   Turek M, Besseling J, Bringmann H. Agarose Microchambers for Long-term
Calcium Imaging of Caenorhabditis elegans. Journal of visualized experiments : JoVE.
2015;(100):e52742. Epub 2015/07/02. doi: 10.3791/52742. PubMed PMID: 26132740;
PubMed Central PMCID: PMC4544933.
46.   Bringmann H. Agarose hydrogel microcompartments for imaging sleep- and
wake-like behavior and nervous system development in Caenorhabditis elegans larvae. J
Neurosci Methods. 2011;201(1):78-88. Epub 2011/08/02. doi: S0165-0270(11)00418-3
[pii]10.1016/j.jneumeth.2011.07.013. PubMed PMID: 21801751.
47.   Nagy S, Raizen DM, Biron D. Measurements of behavioral quiescence in
Caenorhabditis elegans. Methods. 2014. Epub 2014/03/20. doi:
10.1016/j.ymeth.2014.03.009. PubMed PMID: 24642199.
874
875