A sleep-active neuron can promote survival while sleep behavior is disturbed

Inka Busack[1], Henrik Bringmann[1]*

[1] BIOTEC, Technical University Dresden, Germany

*Corresponding author: henrik.bringmann@tu-dresden.de

Abstract

Sleep is controlled by neurons that induce behavioral quiescence and physiological

restoration. It is not known, however, how sleep neurons link sleep behavior and survival.

In *Caenorhabditis elegans*, the sleep-active RIS neuron induces sleep behavior and is

required for survival of starvation and wounding. Sleep-active neurons such as RIS might

hypothetically promote survival primarily by causing sleep behavior and associated

conservation of energy. Alternatively, RIS might provide a survival benefit that does not

depend on behavioral sleep. To probe these hypotheses, we tested how activity of the sleep-

active RIS neuron in *Caenorhabditis elegans* controls sleep behavior and survival during

larval starvation. To manipulate the activity of RIS, we expressed constitutively active

potassium channel (*twk-18gf* and *egl-23gf*) or sodium channel (*unc-58gf*) mutant alleles in

this neuron. Low levels of *unc-58gf* expression in RIS increased RIS calcium transients

and sleep. High levels of *unc-58gf* expression in RIS elevated baseline calcium activity and

inhibited calcium activation transients, thus locking RIS activity at a high but constant

level. This manipulation caused a nearly complete loss of sleep behavior but increased

survival. Long-term optogenetic activation also caused constantly elevated RIS activity

and a small trend towards increased survival. Disturbing sleep by lethal blue-light

stimulation also overactivated RIS, which again increased survival. FLP-11 neuropeptides

were important for both, induction of sleep behavior and starvation survival, suggesting

that FLP-11 might have divergent roles downstream of RIS. These results indicate that

promotion of sleep behavior and survival are separable functions of RIS. These two

functions may normally be coupled but can be uncoupled during conditions of strong RIS

activation or when sleep behavior is impaired. Through this uncoupling, RIS can provide

survival benefits under conditions when behavioral sleep is disturbed. Promoting survival

in the face of impaired sleep might be a general function of sleep neurons.

Author Summary

The sleep neuron RIS in *Caenorhabditis elegans* is required to induce sleep behavior and

to promote survival upon stress. Here we show that RIS can provide survival benefits

even when sleep is disturbed.

Introduction

Sleep is defined by behavioral inactivity and reduced responsiveness [1]. It is a physiological state that conserves and allocates energy, promotes restorative processes, and is important for brain and cognitive functions [2-4]. Sleep disorders and deviations from the normal sleep amount are widespread in modern societies and pose an unresolved medical and economical problem [5-7].

Sleep is found in most animals ranging from cnidarians to humans. Central to the induction of sleep in all species are so-called sleep-active neurons that activate to inhibit wakefulness behavioral circuits and thus cause sleep behavior [8, 9]. *C. elegans* has a small and invariant nervous system and sleep is controlled by relatively few neurons. RIS is a GABAergic and peptidergic interneuron that activates during sleep, and its ablation virtually abolishes sleep behavior during molting of developing larvae, in arrested larvae, and in adults following cellular stress [10-21]. Optogenetic manipulations showed that acute activation of RIS inhibits wakefulness activity such as feeding and locomotion [10, 15], whereas inactivation of RIS inhibits sleep [12, 18, 22, 23]. RIS is important for survival during starvation and after wounding [12, 18, 24]. This neuron counteracts aging phenotype progression during starvation-induced arrest [12]. It supports the protective transcriptional response that underlies the arrest, and promotes the activity of the FOXO transcription factor DAF-16, which is required for survival [24-26]. Thus, RIS is a sleep-active neuron that is crucial for *C. elegans* sleep and stress survival.

Little is known about how sleep-active neurons such as RIS link sleep behavior and survival. Different hypotheses are conceivable. For example, sleep-active neurons might promote survival by reducing behavioral activities during sleep, thus conserving and allocating energy or optimizing behavior [27, 28]. The close association of sleep and physiological benefits has suggested a causal relationship. For example, memory and information processing are thought to be improved in the human brain during sleep when information inputs are blocked [29, 30], and energy is conserved by reduced behavioral activity [28]. A less explored additional hypothesis is that sleep-active neurons provide benefits during sleep, but independently of sleep behavior. According to this hypothesis,

69 sleep-active neurons have two parallel functions. The first is to induce sleep behavior, and
70 the second is to promote survival. This hypothesis implies that physiological benefits of
71 sleep-active neuron functions can be exerted also when sleep behavior is impaired.
72
In order to find out how RIS links sleep and survival, we manipulated RIS activity through
transgenic expression of ion channel mutants in RIS. By strongly expressing *unc-58gf* in
RIS, we achieved an overall increase of RIS activity that was caused by constantly elevated
baseline calcium activity in RIS. RIS appeared to be locked at this higher activity level, as
it was not able to activate further upon stimulation. Survival was increased in this strain,
but sleep behavior was almost completely lacking. The sleep loss phenotype included a
lack of dampening of the activity of interneurons, sensory neurons and muscles that is
typical for normal sleep. Thus, strong expression of RIS led to an apparent uncoupling of
sleep behavior and survival. Long-term optogenetic activation also generated constantly
increased calcium activity and a small trend towards increased survival. Stimulation with
blue light disturbed sleep and increased RIS activation, which in turn extended survival.
Finally, we find that RIS promotes sleep behavior and promotes survival through FLP-11
neuropeptides.
86
RIS hence appears to be able to promote survival independently of sleep behavior,
indicating that promotion of sleep behavior and survival are separable functions of RIS that
are normally coupled during sleep but that can be uncoupled. Uncoupling appears to occur
during conditions of stimulation that disturb sleep and increase RIS activity. This suggests
that an important function of RIS overactivation is to promote survival not only during
normal sleep, but also and in particular when sleep behavior is disturbed. Sleep-active
neurons might generally function to promote survival both during normal sleep and even
more so when sleep behavior is disturbed.
95
96
97 **Results**
98
Generation of a strain set for controlling RIS activity

We set out to control RIS activity to study sleep and survival during L1 arrest. For this

study, we generated strains that express constitutively active ion channels with different

ion permeability and expression levels in RIS. To constitutively inactivate RIS, we

expressed potassium channels that carry mutations that cause potassium leak currents using

the *flp-11* promotor that is strong and highly specific for RIS [11, 17, 31]. We used two

different *egl-23* channel mutations to achieve different strengths of inactivation of RIS

(*RIS::egl-23gf(weak)* expresses *egl-23(L229N)* in RIS, *RIS::egl-23gf(strong)* expresses

egl-23(A383V) in RIS) [32]. To create a strong tool for RIS inhibition, we expressed a *twk-*

18 mutant allele that is known to cause a very strong potassium current (*RIS::twk-18gf*

expresses *twk-18(e1913)* in RIS) [33]. In all three strains, the channel genes were knocked

into the endogenous *flp-11* locus behind the *flp-11* gene to generate operons that express

both *flp-11* and the mutant ion channel. All fusion proteins localized to the cell membrane

of RIS (Fig 1A-D).

To constitutively over activate RIS, we used an unusual K2P channel mutant that is

permeable to sodium [32] (Thomas Boulin, personal communication). We expressed the

ion channel mutant in RIS (*unc-58(L428F),* Fig S1A) tagged with mKate2 at either a lower

or a higher level in RIS. The different expression levels were achieved by generating two

different expression loci for the channel in the genome. To achieve a strong expression of

the *unc-58(L428F)* channel (*RIS::unc-58gf(strong)*), we turned the endogenous *flp-11*

locus into an operon as above. To achieve a weaker expression of the channel (*RIS::unc-*

58gf(weak)), we expressed mKate2-tagged *unc-58(L428F)* from a single-copy knock-in

locus on chromosome III [34], which contained only a minimal *flp-11* promoter and 3'

UTR [11] (Fig 1 E-H). Fluorescent imaging after two days of L1 arrest showed that the

stronger strain expressed at about double the level of the weaker strain and in both

transgenes, the channel was expressed prominently in RIS and was targeted to the cell

membrane (Fig 1E-H, S1). Even after 21 days of arrest, mKate2-tagged *unc-58(L428F)*

expression could be detected in the strong strain indicating that the sodium channel mutant

is properly folded and targeted to the plasma membrane across the entire arrest (FigS1E).

RIS::unc-58gf(strong) promotes survival of L1 arrest

Previous studies have shown that RIS is required for survival during larval arrest and following wounding of the adult [12, 18, 35]. RIS activity controls the gene expression changes that underly the physiological adaptation to starvation [24]. *RIS::unc-58gf(strong)* and *RIS::twk-18gf* displayed opposing gene expression changes compared with the wild type. *RIS::twk-18gf* is more similar to *aptf-1(-)*, which causes loss of RIS function [10, 24]. To identify additional genes whose expression depend on RIS activity, we directly compared the gene expression profiles of *RIS::unc-58gf(strong)* and *RIS::twk-18gf* during L1. This analysis identified 4676 differentially expressed genes including 1910 novel genes that were not previously reported to be controlled by RIS (Fig 2A, Table S5). This comparison confirms the notion that activation and inactivation of RIS have opposing effects on gene expression [24]. These data support the idea that the expression of *twk-18gf* and *unc-58gf* in RIS cause RIS inactivation and RIS activation gene expression phenotypes, respectively [24].

To get an understanding of the role of RIS activation on survival, we investigated how the different RIS-expressed ion channels affect survival in arrested L1 larval worms. For this experiment, we kept arrested L1 larvae as liquid culture, as this allowed for a higher number of animals to be processed. We regularly took a sample of the larvae from the liquid culture and measured two different readouts. We first measured starvation survival by scoring the fraction of animals that had survived and secondly, we scored the ability of starved animals to recover and develop to the adult stage after refeeding [12, 23, 24, 36]. *RIS::twk-18gf* larvae showed strongly reduced starvation survival by about half compared to the wild type. *egl-23(strong)* resulted in a more moderate yet significant starvation survival reduction of approximately 21%. Weak inactivation by *egl-23(weak)* or overactivation by *unc-58gf(weak)* of RIS activity did not significantly change starvation survival. Strong RIS overactivation by *unc-58gf(strong)* had an increased starvation survival of around 19% compared to wild type (Fig 2B). These results are in accordance with previous reports and the expected changes caused by the ion channel transgenes [12,

24, 32, 33]. The recovery upon refeeding matched the survival curves (Fig 2C). To test

whether this survival-extending effect was caused by *unc-58gf* expression in RIS, we

measured survival of *RIS::unc-58gf(strong)* in an *aptf-1(-)* background, in which

expression of *unc-58gf* in RIS is abolished (Fig S1G). In the combined *RIS::unc-*

58gf(strong); aptf-1(-) strain starvation survival was extremely shortened compared with

wild type (Fig S2) and was similar to *aptf-1(-)* [12] or *RIS::twk-18gf* (Fig 2B), indicating

that the starvation survival extension indeed stemmed from *unc-58gf* expression in RIS.

Ion channel mutant expression in RIS controls baseline RIS activity

How are RIS activity, sleep behavior and survival linked? To shed light on this question,

we imaged RIS calcium activity in these mutants. [10]. We used microfluidic chambers to

keep and image the larvae. This method allowed for a quantification of RIS baseline

activity, RIS calcium activation transients, and sleep behavior [37, 38]. We were able to

measure RIS activity for the first 12 days of starvation, and found that the effects of the

channel transgenes were consistent across the entire time of starvation that we analyzed

(Fig 3A for two-day starvation, 3B for all time points). We first analyzed the baseline

activity of RIS in the absence of calcium transients. The mean RIS baseline calcium activity

levels were reduced by the strongly inactivating transgenes *twk-18gf* and *egl-23gf(strong)*.

Baseline activity levels were modestly increased by *unc-58gf(weak)* and strongly increased

by *unc-58gf(strong)*. The magnitude of the baseline calcium signal alterations of the

different transgenes were thus in accordance with the expression levels and the expected

strengths of the leak currents (Fig 1F). These data demonstrate that transgenic *unc-58gf*

can be used as a tool for increasing calcium activity in a neuron that does not normally

express *unc-58* [17, 31, 32]. We used these tools to create a set of strains that allows for

probing associations of baseline activity levels and survival as well as sleep.

RIS::unc-58gf(strong) **inhibits sleep behavior during L1 arrest**

We next quantified motion quiescence as a readout for sleep and tracked mobility behavior

during L1 arrest inside microfluidic chambers. We used GCaMP fluorescence to quantify

RIS activation transients across the entire measurement [10]. Consistent with previous

reports [10, 12, 20], wild-type RIS showed activation transients over a constant baseline,

and calcium transients were accompanied by phases of behavioral quiescence (Fig 4A-G).

Calcium transients were completely abolished in both the strongest deactivation strains

(Fig 4A, B) as well as in the strong activation strain (4F). In the mild deactivation strain

calcium transients were still visible and correlated with behavioral quiescence, but were

decreased in magnitude (Fig 4C/G, S3A). In *RIS::unc-58gf(weak)*, RIS calcium transient

activity and bouts of behavioral activity were increased (Fig 4E). These results suggest that

RIS activation transients are required for sleep induction and that strong inactivation

prevents these transients. Mobility quiescence decreased with inactivation and was

increased with modest activation as expected. We anticipated that the elevated RIS calcium

baseline of *RIS::unc-58gf(strong)* might be associated with a further increase in sleep

behavior compared with the weak activation. Mobility quiescence was, however,

completely abolished by strong RIS activation in most of the animals (Fig 4H). These

findings were additionally confirmed for a longer period of starvation up to day 12 (Fig

4I). The loss of quiescence of the strong RIS activation strain was similar to, but not as

strong as in *aptf-1(-)* (Fig S2B). The sleep amounts for each strain are consistent for several

days in L1 arrest (Fig 4I). In summary, baseline RIS activity correlates with behavioral

quiescence over a wide range of conditions, except for the *RIS::unc-58gf(strong)*

transgene, which appears to cause a lack of RIS calcium transients and sleep behavior.

RIS::unc-58gf impairs sleep behavior during larval development and following

cellular stress in the adult

C. elegans sleeps during various stages and conditions. Previous studies showed that

cellular stress increases adult sleep [17, 39-42]. During development, sleep is coupled to

the molting cycle and occurs during a period called lethargus, which coincides with

formation of a new cuticle [10, 43, 44]. To test whether constant RIS baseline activation is

also associated with a loss of behavioral quiescence during stress-induced sleep and during

lethargus sleep, we tested whether *RIS::unc-58gf* affects sleep behavior during L1

development as well as following heat stress. For testing for stress-induced sleep, we kept

young adult worms inside microfluidic devices and first measured baseline behavioral

activity at 22°C, and then exposed the animals to heat stress at 37°C for 20 minutes. We

then followed the behavior again after the end of the heat shock [17]. The wild type

immobilized during the heat shock and displayed an increase in behavioral quiescence

following the heat shock. The quiescence response was attenuated in all strains in which

RIS was manipulated. Both *unc-58gf* expressing strains showed decreased behavioral

quiescence. Thus, a reduction of behavioral sleep can also be caused by *RIS::unc-*

58gf(weak) (Fig 5A-G).

To test for sleep that occurs during development, we quantified behavioral quiescence of

RIS::unc-58gf(strong) during lethargus in fed L1 larvae. Behavioral quiescence of

RIS::unc-58gf(strong) was significantly reduced compared with the wild type. The loss of

sleep behavior in lethargus was, however, not as strong as during L1 arrest or following

stress-induced sleep (Fig 5H). Hence, in all the stages and conditions that we tested, sleep

behavior was reduced by *RIS::unc-58gf(strong)*.

We also tested whether the lack of sleep behavior in *RIS::unc-58gf(strong)* during L1 arrest

causes a compensatory increase in sleep when the animals resume development. For this

experiment, we starved larvae for 12 days and then provided them with food, which

allowed for a resumption of development. We then imaged sleep behavior during L1

lethargus in these recovered worms. There was no increase of sleep behavior in recovered

RIS::unc-58gf(strong) larvae compared with wild type (Fig S4). Thus there was no

indication for rebound sleep occurring at the subsequent lethargus stage in *RIS::unc-*

58gf(strong).

RIS::unc-58gf(strong) impairs the dampening of wake-active neuronal circuits

during L1 arrest

RIS::unc-58gf(strong) caused strongly reduced sleep behavior, but the nature of the residual sleep behavior in this mutant was unclear. Did the residual sleep behavior present a generally weakened sleep state or a compensatory deep sleep state? To answer this question, we characterized neuronal activity during in *RIS::unc-58gf(strong)* mutants. For this set of experiments, we imaged larvae in microfluidic chambers and imaged neuronal activity with GCaMP during L1 starvation-induced arrest.

C. elegans sleep is characterized as a state of globally reduced nervous system activity [12, 13, 20]. We measured the overall neuronal activity by utilizing a strain that expresses GCaMP6s in all neurons [45]. We measured both motion behavior and overall neuronal activity by extracting the calcium activity signal of all neurons as one combined signal [12, 22]. Consistent with previous reports, periods of behavioral quiescence were accompanied by strongly reduced global neuronal activity (Fig 6A, B, S5) [12, 13, 20, 22]. We extracted phases of neuronal inactivity as well as phases of behavioral inactivity from the data set. In all strains, neuronal inactivity directly correlated with immobility, suggesting that behavioral quiescence and neuronal quiescence are correlated in all strains irrespective of the level of RIS baseline activity (Fig 6C). Thus, both behavioral inactivity and brain inactivity can serve as equivalent readouts for detecting sleep in this system. Strong RIS inactivation led to an increased mean overall neuronal activity of 50.7% compared to wild type. Intermediate strength of inactivation did not affect mean overall neuronal activity. *RIS::unc-58gf(strong)* caused a significant increase of mean overall neuronal activity of 35.7% (Fig 6D). These results support the notion that *RIS::unc-58gf(strong)* worms are in a state of wakefulness for most of the time and lack phases of strong nervous system activity dampening.

To characterize the effects of *RIS::unc-58gf(strong)* at the level of specific individual neurons, we imaged activity of the RIM neurons, which are second layer interneurons that integrate information from sensory and internal states to control motion behaviors [46, 47]. Again, we used GCaMP imaging in microfluidic chambers to correlate neuronal and

behavioral activity. RIM showed activation transients during phases of behavioral activity.

During phases of behavioral quiescence, RIM activity was strongly reduced and activation

transients were lacking (Fig 6E, F, S6). Similar to the pan-neuronal measurements, the time

fraction during which RIM was inactive correlated with phases of motion quiescence (Fig

6G). Consistent with the pan-neuronal imaging data, *RIS::unc-58gf(strong)* lacked phases

of RIM inactivity (Fig 6H).

An increased arousal threshold and attenuated sensory information processing are key

characteristics of sleep [1, 48]. In *C. elegans*, sensory information processing is inhibited

during sleep in developing larvae and the activity of sensory neurons is reduced [49-51].

For example, the mechanosensory neurons ALM and PLM display dampened activation

transients during sleep in developing L1 larvae [49, 50]. We measured PLM activity in

wild-type, *RIS::unc-58gf(strong)* and *RIS::twk-18gf* larvae. In wild-type animals, PLM

showed activation transients only during phases of wakefulness, whereas PLM appeared

to be rather inactive during sleep. In both *RIS::unc-58gf(strong)* and *RIS::twk-18gf*

mutants, PLM showed activation transients almost continuously throughout the time, with

only small episodes where PLM seemed to be not active (Fig 7). Thus, extended periods

of PLM neuron inactivity were absent in both *RIS::unc-58gf(strong)* and *RIS::twk-18gf*

mutants. Thus, *RIS::unc-58gf(strong)* inhibits the dampening of neuronal activity that is

associated with wakefulness behavior and thus lacks the neuronal signature of deep sleep.

***RIS::unc-58gf(strong)* promotes muscle activity**

An absence of voluntary movement, reduced muscle contraction and a relaxed body

posture are hallmarks of sleep across species including *C. elegans* [44, 52-54]. If *RIS::unc-*

58gf(strong) lacks deep sleep bouts then this should be reflected by the absence of extended

periods of movement quiescence that coincide with reduced muscle activity. During

wakefulness, *C. elegans* shows locomotion behavior and feeds by contractions of the

pharynx muscle [55, 56]. Locomotion and pharyngeal pumping stops during sleep and the

animals assume a relaxed body posture [10, 44, 52, 54]. Underlying these effects, RIS

inhibits both locomotion and pharyngeal pumping [10, 12, 15, 57, 58]. To test the hypothesis that *RIS::unc-58gf(strong)* lacks long phases of muscle relaxation that are typical for sleep, we first imaged arrested larvae inside microfluidic chambers and quantified more precisely movement activity of the body and the pharynx.

To quantify mobility behavior of the head and tail regions, we tracked the positions of neurons that are located in these body parts. To calculate movement speed of the head, we tracked the position of the RIM neurons. For tracking the tail we used the position of the PLM neurons [59]. The tracking data revealed that in *RIS::twk-18gf*, both head and tail movement was similar to the wild type during wakefulness. During the few quiescence phases of *RIS::twk-18gf*, movement speed was reduced less than in the wild type, indicating that the quiescence phases that remain in *RIS::twk-18gf* are less pronounces compared with the wild type. In *RIS::unc-58gf(strong)*, head movement was decreased compared to wild-type worms, whereas there was no difference in tail movement (Fig 8A-B). *RIS::unc-58gf(strong)* worms were uncoordinated and the body was more curved (Fig S7, Movies S7 and S8), which might explain the difference of head and tail movement speeds. In the few locomotion quiescence phases of *RIS::unc-58gf(strong)*, there appeared to be less of a reduction of movement speed.

We next imaged our transgenic strains in the arrested first larval stage for 1 min with a framerate of 10Hz and manually scored pumping rate. This analysis revealed no significant difference in pharyngeal contraction rates for all strains compared to wild type (Fig S8).

To test how *RIS::unc-58gf(strong)* and *RIS::twk-18gf* affect muscle activity, we measured and quantified muscle GCaMP [52]. In the wild type, muscle calcium activity was high during wakefulness, and strongly reduced during sleep. In *RIS::twk-18gf*, muscle calcium was almost constantly elevated and resembled the activity levels of the wild type. *RIS::unc-58gf(strong)* worms showed a general increase in muscle calcium activity (Fig 8C-F). The increased muscle activity is consistent with and might contribute to the slightly uncoordinated movement of this mutant. Phases of behavioral and muscle activity

dampening occurred less often, and if they occurred, then these dampened states appeared

to be less deep. (Fig 8G, H).

In *RIS::unc-58gf(strong)*, RIS calcium activity cannot be easily elevated further by

optogenetic stimulation

Calcium imaging suggested that the RIS baseline in *RIS::unc-58gf(strong)* is substantially

elevated and that calcium transients are lacking. We aimed to better understand the nature

of these physiological changes in RIS that are caused by expression of *RIS::unc-*

58gf(strong). Based on the known characteristics of the *unc-58gf* mutant channel, the

expression of *unc-58gf* might plausibly cause a constant depolarization of the membrane

of RIS [32]. Depolarization in turn should lead to the activation of voltage-gated calcium

channels, and RIS expresses the voltage-gated calcium channels *egl-19* and *unc-2* [31].

The conductivity of voltage-gated calcium channels is increased by depolarization with an

optimum in the depolarized range. Further depolarization causes inhibition and long-term

stimulation reduces conductivity by desensitization [60-65]. RIS depolarization in

RIS::unc-58gf(strong) could thus perhaps be increased to allow for elevated RIS baseline

calcium activity, but might also inhibit further activation of calcium channel activity. We

tested the ability of RIS to activate by optogenetically stimulating RIS in *RIS::unc-*

58gf(strong) inside microfluidic chambers and monitored the calcium activity of this

neuron as well as the motion behavior of the larvae. For this experiment, we used the red-

shifted channelrhodopsin variant ReaChR in combination with GCaMP3 [12, 22, 57]. In

the wild type, optogenetic activation caused a strong increase in intracellular calcium and

a strong induction of motion quiescence, which is consistent with previous results [12, 15,

22, 57]. By contrast, optogenetic activation of RIS in *RIS::unc-58gf(strong)* did not

increase RIS activity further and did not induce behavioral quiescence (Fig 9A-B). This

result is consistent with the view that *RIS::unc-58gf(strong)* causes a heightened level of

RIS activity that cannot be easily increased further with optogenetic activation. Hence, in

RIS::unc-58gf(strong), RIS appears to be locked in a state of heightened activity while

further activation is inhibited.

Long-term optogenetic activation of RIS increases RIS baseline activity and inhibits

activation transients

RIS::unc-58gf(strong) appears to cause increased RIS activity that cannot easily be

increased further. This could hypothetically be explained by a high level of depolarization

in *RIS::unc-58gf(strong)* that causes both RIS baseline elevation as well as inhibition of

RIS calcium transients. To test the idea that constant depolarization elevates baseline RIS

activity while inhibiting activation transients, we used long-term optogenetic activation.

We used again ReaChR to depolarize the membrane and followed the activity of RIS.

Optogenetic tools like ReaChR typically display a brief photocurrent peak activity within

the first second of light exposure and then plateau at a lower photocurrent level for an

extended period of time [66]. We previously used ReaChR to establish long-term

optogenetic manipulations in *C. elegans* [23]. To test for the activation of RIS during

prolonged optogenetic activation, we activated ReaChR in RIS with orange light for 11h

and followed the activity of RIS using GCaMP and measured mobility of the larvae that

were kept inside microfluidic chambers. During the first hour of optogenetic stimulation

of RIS, there was a strong increase of calcium activity as well as of behavioral quiescence

(Fig 10A-C). After this first peak period, the ongoing optogenetic stimulation led to lower

but more constant increase of RIS baseline activation (which was, however, weaker than

the activation observed in *RIS::unc-58gf(strong)* average calcium baseline activity increase

in *RIS::unc-58gf(strong)* compared with wild type was 1.55, and average calcium activity

baseline increase from 2-11h of activation in *RIS::ReaChR* compared with wild type was

1.25) (Fig 10D, E). Long-term optogenetic activation caused reduced calcium transients,

and the residual calcium transients did not correlate with phases of immobility. This

indicates that long-term optogenetic activation has an inhibitory effect on RIS calcium

transients (Fig 10F). The optogenetic long-term RIS activation did not cause a measurable

decrease in behavioral quiescence (Fig 10C). Thus, long-term optogenetic stimulation

leads to a constantly increased RIS baseline activity and an inhibition of RIS calcium

transients that is similar to, but weaker than, the activation observed in *RIS::unc-*

58gf(strong). This difference in effect strength might be explained by the difference in ion

conductivity of these channels [32, 66]. These data indicate that long-term optogenetic

stimulation can lead to a constant increase in baseline RIS calcium activity at a level that

is above the baseline activity without optogenetic stimulation, yet lower than the maximum

possible optogenetic RIS activation peak intensity. These optogenetic data are consistent

with the idea that increased depolarization of RIS leads to a constant elevation of baseline

calcium activity and impairs calcium transients.

Long-term optogenetic activation of RIS causes a small increase in survival

Our data using *RIS::unc-58gf(strong)* indicated that extended survival of L1 arrest is

associated with increased overall RIS activity. To confirm this result with an independent

approach, we used *RIS::ReaChR* for long-term activation of RIS using the OptoGenBox

system [23]. We kept L1 larvae inside microchambers and constantly stimulated the entire

chambers with orange light. The optogenetic stimulation was only paused briefly once

every day to monitor the survival of the larvae. The stimulation protocol was applied until

all larvae had died. The functionality of ReaChR in *C. elegans* requires the addition of all-

trans retinal (ATR). Addition of ATR shortened the survival (Fig 10G). We hence

compared the survival of N2 exposed to ATR to worms expressing ReaChR that were also

exposed to ATR. By this comparison, optogenetic long-term RIS activation caused a small

trend for increased survival that was statistically significant according to Fisher's Exact

Test but not significant according to the Logrank test (p=0.02 Fisher's Exact Test, p=0.09

Logrank, Fig 10G). While this result borders statistical significance, it is consistent with

the view that activation of RIS promotes survival. The small effect size is consistent with

the small RIS activity increase caused by extended optogenetic stimulation. Thus, this

experiment supports the view that RIS activation promotes survival of L1 arrest.

Lethal blue-light stimulation activates RIS, and RIS supports survival of lethal blue

light

Disturbing sleep by sensory stimulation leads to arousal that forces wakefulness and prevents the dampening of underlying neuronal circuits [12, 20, 22, 44, 67, 68]. Sleep disturbance or deprivation can cause increased RIS activation [22, 24, 69]. The high level of RIS activity in turn could be protective under such conditions and promote survival [24]. Thus, *RIS::unc-58gf(strong)* might cause a state that is similar to a state caused by prolonged sleep deprivation or disturbed sleep. In both, disturbed sleep and *RIS::unc-58gf(strong)*, the activity of wakefulness circuits is not dampened yet coincides with elevated RIS activity.

To further illustrate the idea that sensory stimulation activates RIS while impairing sleep, we stimulated worms with a lethal intensity of blue light. We used arrested L1 larvae after 48h in the absence of food, a condition during which the animals normally show regular RIS activation transients and sleeping behavior [12]. We kept L1 larvae in microchambers and repeatedly applied pulses of blue light while monitoring RIS activity and motion behavior. For comparison, we used control animals that were exposed to shorter pulses of blue light of only one tenth the length, thus still allowing for calcium imaging. Control larvae displayed sleeping behavior detected by motion quiescence and RIS calcium activity during about a quarter of the time (Fig 11A, B). Strong blue-light stimulation caused an increase in RIS calcium activity compared with control animals (after 1h of strong blue-light stimulation, the activity of RIS was approximately 5.6-fold higher compared with worms that were stimulated with a lower dose). Many of the calcium transients were very short, however, and did not coincide with increased motion quiescence (Fig 11B-C). Hence, the stronger blue-light stimulus increased RIS activity but apparently did not increase sleep behavior.

To probe for the role of RIS in survival of lethal blue-light stimulation, we compared wild-type and *RIS::twk-18gf* animals, in which RIS activated less under blue-light stimulation (Fig 11C). We scored the time point at which the animals became terminally immobile, which presents an established proxy to score for death [70]. Control animals with functional RIS died around 7-8h after strong blue-light stimulus onset. RIS-impaired

animals already died at around 6h after stimulus onset. The presence of functional RIS

hence extended survival by 15% (Fig 11D).

Thus, stimulation by a lethal blue-light stimulus can increase RIS calcium activity without

increasing RIS-dependent sleep behavior. While all animals were eventually killed by the

blue-light stimulus, RIS activity slightly extended survival. The blue-light experiment thus

supports the idea that RIS is a protective neuron that activates upon stressful stimulation.

As in *RIS::ReaChR* and during long-term optogenetic stimulation of RIS, increased RIS

activity was not associated with increased sleep behavior.

***flp-11* is a key neuropeptide gene required for quiescence behavior in L1 arrest**

What are the key transmitters that RIS uses to control sleep during L1 arrest? GABA is

expressed in RIS but has not been shown to play a substantial role in promoting sleep [10,

15]. FLP-11 is a group of neuropeptides of RIS that present the major transmitters of RIS

required for sleep induction during development [11]. Expression from the *flp-11* promoter

depends on neuronal activity levels [71], and is increased in *RIS::unc-58gf(strong)* (Fig

S3B), suggesting that FLP-11 might also be a major transmitter for sleep induction during

L1 arrest. To directly test the role of FLP-11 in quiescence during L1 arrest, we measured

the effects of *flp-11* deletion (*flp-11(-)*) using microfluidic chamber imaging [11]. *flp-11(-*

) strongly reduced quiescence behavior (Fig 12A). *RIS::unc-58gf(strong)* still displayed a

small fraction of quiescence behavior during L1 arrest. To test whether this residual

quiescence was caused by *flp-11*, we deleted the entire *flp-11* gene in the *RIS::unc-*

58gf(strong) operon locus (Fig S9A) and quantified quiescence behavior. The sleep loss of

RIS::unc-58gf(strong) was reduced further by *flp-11(-),* suggesting that the residual

quiescence behavior in *RIS::unc-58gf* indeed depends on *flp-11* (Fig 12A).

FLP-11 neuropeptides support survival

What is the key transmitter of RIS that promotes survival? RIS is GABAergic, but GABA

has been suggested to be a negative regulator of lifespan [72]. We showed above that *flp-*

11 is required for most of the immobility during L1 arrest sleep. We hence tested whether

FLP-11 neuropeptides are also required for survival. Deletion of *flp-11* reduced survival

by 24.7%, suggesting that *flp-11* contributes to survival (Fig 12B). To test whether *flp-11*

is required for increased survival caused by *RIS::unc-58gf(strong)*, we measured survival

in the *RIS::unc-58gf(strong), flp-11(-)* strain. *flp-11(-)* suppressed almost the entire

survival extension caused by *RIS::unc-58gf(strong)* (Fig 12B). Similarly, the recovery

upon feeding was suppressed by *flp-11(-)* in *RIS::unc-58gf(strong)* (Fig S9B). Together,

these experiments indicate that *flp-11* plays an important role in survival and recovery of

L1 arrest both in the wild type and in *RIS::unc-58gf(strong)*.

These results suggest that *flp-11* plays two roles, one role is to promote sleep and the other

is to promote survival. Deletion of *flp-11* blunts survival less than strong RIS inactivation,

indicating that additional neurosecretory factors might also contribute to survival. As RIS

expresses multiple neuropeptides and also GABA [11, 15, 17], it may use multiple

transmitters to control sleep behavior and survival. In summary these data suggest that

FLP-11 neuropeptides are the major transmitters released by RIS to promote both sleep as

well as survival during L1 arrest.

Discussion

Tools to manipulate neuronal activity are instrumental for studying neuronal circuits and

physiology. Here we establish the constitutively active sodium channel gene *unc-58gf* as a

tool for activation of a neuron that normally does not express *unc-58* [31, 32]. Modest

expression of *unc-58gf* in RIS during L1 arrest caused an increased baseline activity of RIS

and increased the probability of RIS to experience calcium activation transients. It thus

appears that *unc-58gf* can activate RIS when expressed at lower level. Strong expression

of *unc-58gf* in RIS caused constant high levels of baseline RIS activity yet prevented

further activation transients. When expressed highly, *unc-58gf* thus locks RIS activity at a

level that is higher than the normal baseline and at the same time also inhibits RIS calcium
transients. Thus, *unc-58gf* can be used as a tool to create different and complex patterns of
neuronal activity, but the level of expression is crucial to achieve different effects. Together
with constitutively active potassium ion channels that cause various levels of inhibition,
activity of a neuron can be modulated from strong inhibition to constant elevation of
baseline activity.

RIS inhibits wakefulness neurons to cause sleep behavior and it promotes survival under
stressful conditions such as larval starvation or wounding [10, 12, 18, 24, 35]. A
parsimonious explanation linking these two observations would be that sleep promotes
survival, for example through energy conservation or behavioral optimization [27]. Energy
conservation and behavioral optimization certainly are important functions of sleep
behavior [28]. In addition, sleep neurons could promote physiological benefits by
mechanisms that do not necessarily depend on sleep behavior.

How can the manipulation of neuronal activity be employed to solve how neurons control
behavioral and physiological function? Here we used transgenic expression of ion channel
mutants in RIS to shed light on the relationship between RIS activity, sleep behavior and
survival. *RIS::unc-58gf(strong)* promoted survival, but also inhibited sleep behavior. Thus,
RIS::unc-58gf(strong) allowed for an uncoupling of sleep behavior and survival functions
of RIS. Our data thus suggest that the sleep-active RIS neuron can promote survival
independently of sleep.

What are hypothetical mechanisms that allow for the uncoupling of sleep behavior and
survival benefits of RIS? The loss of sleep behavior in *RIS::unc-58gf(strong)* could
hypothetically be explained by the absence of calcium transients in RIS. This could, for
example, be caused by desensitization of wakefulness circuits that are downstream of RIS.
According to this hypothesis, wakefulness circuits would decrease their activity with
increasing RIS activity, and would start getting more active again only after a period of
stagnating or decreasing RIS activity. Both, the induction of sleep behavior and the
promotion of survival by RIS depend on FLP-11 neuropeptides, which are thought to be

released upon RIS activation [11]. The two functions of FLP-11 might mechanistically
separate at the level of divergent receptors, signaling pathways or cell types downstream
of FLP-11 that mediate either the sleep or the survival response. For example, the receptors
and downstream cell types that mediate the effects of FLP-11 on dampening wakefulness
circuits might desensitize quickly, thus limiting the amount of behavioral sleep that can be
induced by FLP-11 released from RIS. Along these lines, RIS activity might exert a
survival-promoting effect through a mechanism with different desensitization properties.
Survival appears to be a function of RIS activity, depends on *flp-11*, but does not seem to
depend on activation transients. Thus, the underlying signaling components do not seem to
desensitize quickly but might respond to lower levels of RIS activity or might integrate
RIS activity over time. This could hypothetically be achieved by FLP-11 activating
downstream receptors and cell types that react sensitively to FLP-11 concentration, that do
not easily desensitize and that promote survival upon sensing FLP-11. The key FLP-11
receptors required for promoting sleep and survival and the cells in which these receptors
function have not yet been identified [11]. The identification of FLP-11 receptors will
allow for further studies of the mechanisms by which FLP-11 causes sleep behavior and
survival. Another hypothetical explanation for changes in sleep behavior upon expression
of ion channel mutants might be changes in the underlying neuronal connectivity. Further
experimentation is required to test these speculative hypotheses.

What is the mechanism and role of homeostatic compensation during sleep deprivation?
Homeostatic regulation can compensate for lost or disturbed sleep across species and plays
an important role in the regulation of sleep in *C. elegans* [22, 53, 67, 69, 73]. The typical
effects of disturbing sleep or depriving an animal of sleep are a reduction of sleep behavior
caused by continued activity of wakefulness circuits that steer behavioral activity.
Increased behavioral activity is often accompanied by a subsequent overactivation of sleep-
active neurons that promote a rebound phase of increased sleep duration or intensity.
Rebound sleep in turn is thought to compensate for lost or disturbed sleep, at least partially
[8, 22, 24, 67, 69, 73, 74].

How is *RIS::unc-58gf(strong)* related to sleep deprivation and sleep homeostasis? Does
RIS::unc-58gf(strong) trigger homeostatic compensation of the lost sleep? Does the
transgene cause a state that is similar to the effects of sleep-deprivation? To answer these
questions, we first studied behavioral sleep across stages. Across the life of arrested larvae,
behavioral quiescence was reduced in *RIS::unc-58gf(strong)*. Furthermore, sleep was
reduced by *RIS::unc-58gf(strong)* during L1 lethargus and after stress in the adult. We also
studied the activity of excitable cells in *RIS::unc-58gf(strong)* in arrested L1 larvae. We
studied the activity of sensory neurons, interneurons and body wall muscle. All of these
excitable cells are known to be rather inactive during sleep and thus the study of the activity
of these excitable cells allows for a quantitative description of sleep depth. Our data showed
that the sleep bouts in *RIS::unc-58gf(strong)* are less pronounced and hence appear to be
less deep. Hence we could not find any signs of deeper or compensatory sleep in *RIS::unc-*
58gf(strong). By contrast, the analysis of excitable cells suggested that the residual bouts
of behavioral quiescence in *RIS::unc-58gf(strong)* were generally weakened. *RIS::unc-*
58gf(strong) thus displays the neuronal signature of sleep deprivation or disturbed sleep,
i.e. the combination of impaired wakefulness circuit dampening and over activation of RIS.
Thus, *RIS::unc-58gf(strong)* might be understood and employed as a model of disturbed
sleep or of sleep deprivation.
612
Does RIS activate when the animals are disturbed to promote survival [12, 18, 22, 24, 69]?
To further illustrate the idea of RIS acting as a promoter of survival, we disturbed animals
during sleep via a lethal blue-light stimulus. Lethal blue light increased RIS activity and
functional RIS was required for extending the survival of the animals during blue-light
stimulation yet did not increase sleep behavior. This supports the idea that RIS activates
upon sensory stimulation to promote survival independently of increasing sleep behavior.
619
620 What is the mechanism by which RIS promotes survival? Is there an important role of RIS
621 to conserve energy by shutting down behavior? The reduction of brain and muscle activity
622 likely contributes to survival by conserving energy. While energy consumption appears to
623 be playing a role in survival, behavioral activity and thus energy consumption does not
624 correlate with survival when transmission from RIS is strongly increased. In *RIS::unc-*

58gf(strong) animals, neuronal, muscle and behavioral activity are high and consume energy, yet survival is increased. Hence RIS appears to be able to promote survival independently of sleep behavior and despite the energy loss that can be expected to be caused by the increased activity of excitable cells [28]. As both sleep and survival are functions of RIS activity over a wide range of conditions, the two processes should be coupled during normal sleep. During each sleep bout, RIS depolarizes and thus should promote both, sleep and survival. Such coupling of sleep and survival during normal sleep might ensure that phases of brain and behavioral inactivity coincide with phases during which survival-promoting processes are active. Tight coupling might serve to optimize energy expenditure, as energy conserved by sleep could be directly allocated to survival-promoting processes [28].

How does RIS cause physiological protection and increased survival independently of conserving energy? RIS promotes survival of stressful conditions and underlying changes in protective gene expression [12, 18, 24]. Together with the apparent strong reduction of sleep behavior in *RIS::unc-58gf(strong)*, this might suggest that the role of RIS overactivation upon disturbance or sleep deprivation is not only to promote rebound sleep behavior, but also to promote protective physiological benefits that support survival. An important function of RIS might thus be to promote survival when the animal is disturbed by sensory stimulation. Disturbance, e.g. by blue light, might directly be stressful and thus require an increased survival response. Disturbed sleep is often associated with stress, for example when mechanical stimuli are experienced. Hence RIS might hypothetically also act as part of a sensory system that activates during stimulation to signal an increased demand of physiological protection.

The FOXO transcription factor DAF-16 has been shown to be linked to the control and functions of sleep. It is required for sleep under various conditions including during L1 arrest. Also, FOXO has been shown to be activated upon sleep deprivation during both lethargus as well as during L1 arrest and to be required for increased rebound sleep following sleep disturbances [24, 58, 67, 73] (Fig S10). A plausible model hence suggests that RIS activation promotes DAF-16-dependent gene expression changes in order to

support survival [24]. Our data indicate that RIS strongly promotes protective gene

expression despite the loss of sleep behavior. Thus, *RIS::unc-58gf(strong)* intensifies the

survival function of RIS while inhibiting the sleep function of RIS.

By controlling the activity of RIS, we showed that the survival function and the sleep

behavior function of this neuron can be decoupled. By constantly elevating RIS baseline

activity, we generated a condition in which wakefulness neural activity and sleep-active

neuron activity coexist. A similar condition often underlies disturbed sleep. Our data

suggest that when sleep is disturbed, RIS promotes survival despite the loss of behavioral

sleep. It would be interesting to study whether sleep neurons in other species also can

promote physiological benefits in the face of lost sleep.

Materials and Methods

Resource and Materials Availability

Key *C. elegans* lines that were used in this project are available at the Caenorhabditis elegans Genetics Center. Other *C. elegans* strains, plasmids, or additional reagents generated for this study are available upon reasonable request.

Data Availability

All relevant data are within the manuscript and its supporting Information files. Matlab scripts can be found at Github (https://github.com/ibusack/A-sleep-active-neuron-can-promote-survival-while-sleep-behavior-is-disturbed).

Experimental model and subject details

C. elegans hermaphrodites were maintained at 20°C. They were kept on Nematode Growth Medium Plates (NGM). The *Escherichia coli* strain OP50 served as a food source during maintenance [75].

Method Details

Crossing *C. elegans* strains

Males were generated via heat shock or maintained by a continuation of crossings of worms of the same genotype [76]. Single worms of the F2 generation were genotyped using duplex PCR genotyping [77]. For crossing fluorescent transgenes, fluorescent markers were used. All strains and primers of this study can be found in the supplementary tables S2 and S3.

Creating mutants with the CRISPR/Cas9 system

We designed the sequences of the following transgenes in silico. Transgenes were synthesized and inserted into the genome by using CRISPR/Cas9 by SunyBiotech. Transgene sequences were verified by Sanger sequencing.

PHX1433   *flp-11(syb1433 [flp-11-SL2-egl-23cDNA(A383V)-linker-mKate2]) X*

PHX1445   *aptf-1(gk794) II; flp-11(syb1445[flp-11-SL2-unc-58(L428F)-linker-*
           mKate2]) X

PHX1464   *flp-11(syb1464[flp-11-SL2-egl-23cDNA(L229N)-linker-mKate2]) X*

PHX2193   *flp-11(syb2193[flp-11b-SL2(gpd-2)-mKate2-linker-twk-18(e1913)]) X*

PHX2493   *lgc-38(syb2346[flp-11p::dpy-10 site::flp-11 3'UTR], syb2493[ReaChR-*
           *linker-*mKate2]) III.

PHX3190   *lgc-38(syb2346[flp-11p::dpy-10 site::flp-11 3'UTR] ,syb3190[unc-*
           58(e665)-linker(GSGSGSGSG)-mKate2]) III

PHX4110   *lgc-38(syb2346[flp-11p::dpy-10 site::flp-11 3'UTR], syb4110[unc-58gf-*
           CAI-1.0-linker(GSGSGSGSG)-mKate2]) III

PHX4416   *aptf-1(gk794) II; flp-11(syb1445 syb4416) X*

The coding sequences of the ion channel transgenes were synthesized and consisted each of the spliced coding sequence of the ion channel including the mutation that caused the leak current [32], a nine amino acid long C-terminal linker sequence (GSGSGSGSG) followed by a C-terminal mKate2 gene. The coding sequence of mKate2 was codon optimized and two introns were inserted [78]. The coding sequences of the ion channels were not codon optimized to allow for proper folding of the membrane proteins. We tried codon-optimization [78] of the *unc-58(L428F)* channel gene and mKate2 to achieve high expression, but this also caused aggregation of the fluorescent fusion protein (Fig S1A, D). Hence, we used the endogenous gene sequence for the channel and only codon-optimized mKate2. The sequences of the ion channel transgenes can be found in the supplemental information.

Analysis of the RIS transcriptome showed that *flp-11* has the strongest and most specific promoter of all RIS-expressed genes, with 157-fold higher expression in RIS than in other neurons [17]. *flp-11* is also expressed in other neurons, albeit at low levels [11, 17, 31]. Using imaging settings that cause oversaturation of the mKate2 signal in RIS revealed that there is a weak expression of *unc-58gf::mKate2* in a few neurons in the ventral cord in

RIS::unc-58gf(strong) (Fig S1F). Expression from the *flp-11* promoter in RIS but not in

other neurons depends on *aptf-1* [11]. We hence imaged mKate2 expression in *aptf-1(-).*

Consistent with previous results [11], mKate2 expression was abolished in RIS but was

still present in the ventral cord (Fig S1G). Thus, expression from the *flp-11* promoter is

strong, has a high level of RIS specificity, and RIS expression depends on *aptf-1*.

The sequences of the ion channels of the strains PHX1433, PHX1445, PHX1464 and

PHX2193 were inserted after the stop codon of the *flp-11* endogenous coding sequence

together with an SL2 site. Thus, an operon was created that expressed both *flp-11* and the

ion channel gene from the *flp-11* promoter. The transgenes of PHX2493, PHX3190 and

PHX4110 were generated via a single-copy knock-in loci for defined gene expression (SKI

LODGE) system [34]. For the SKI LODGE expression in RIS we first inserted the minimal

flp-11 3' and 5'UTRs separated by a *dpy-10* sgRNA cutting site. The expression from these

minimal UTR sequences allowed specific expression in RIS, but the expression levels were

lower compared with the endogenous *flp-11* locus (Fig 1A-D, S1A-D). The final sequences

of all transgenes can be found in the supplementary Text S1.

Imaging using microfluidic devices

Worms were filmed in microfluidic devices made from agarose hydrogel as previously

described [37, 38]. In brief, a PDMS mold was used to cast box-shaped indentations from

melted 5% agarose diluted in M9 buffer. After agarose hydrogel solidification the stamp

was removed resulting in hundreds of microchambers sized 110x110x10μm for L1 arrest

chambers or 190x190x15μm for L1 lethargus chambers per agarose slab. Depending on

the experiment, approximately 9-36 pretzel-stage eggs per strain were then transferred

without food for L1 arrest or with food for lethargus into the indentations in the hydrogel

and the chambers were closed with a coverslip and kept in a temperature-controlled dish.

Hatching of the larvae in the absence of food caused L1 arrest. Chambers were timed so

that the worms were incubated for 48h at 25°C before imaging. The number of biological

replicates (n) imaged in microfluidic devices is stated in each figure. One or several

technical replicates (different microfluidic devices) were imaged for each experiment.

Software for imaging

All image acquisition was performed with Nikon NIS Elements 5 software (Nikon,

Tokyo).

Dark Differential Contrast (DIC) Imaging

Long-term dark differential contrast (DIC) images were acquired with a standard 100W

halogen lamp and an infrared filter (Semrock Brightline HC 785/62) (Idex Health and

Science, New York). The images for L1 arrest imaging were taken 48h upon starvation for

12h with a frame rate of 0.1Hz. A 20x objective for imaging was combined with a 0.7 lens.

Lethargus data was acquired with similar settings except the frame rate was 0.2Hz.

Functional Ca^{2+}-imaging

For calcium imaging we used either a back-illuminated sCMOS camera (Photometrics

Prime 95B, 1200x1200 pixels) (Fig 3, 4, 8C-H, S3) or an Andor iXon EMCCD camera

(512x512 pixels, Andor Technology Ltd., Belfast). (Fig 6, 7, 8A-B, S5, S6). The gain of

the iXon camera was set to 200. The iXon camera's read out mode was 10MHz at 14-bit.

The images of the Photometrics camera were binned 2 x 2.

A 20x objective was used to simultaneously image 3 x 3 adjacent microchambers

containing one worm each (This configuration was used to obtain data for Fig 6, 7, 8A-B,

S5, S6). A 10x objective combined with a 1.5x lens was used to simultaneously image 6 x

6 adjacent microchambers containing one worm each (this configuration was used to obtain

data for Fig 3, 4, 8C-H, S3).

For calcium imaging we utilized a 480nm LED and a standard GFP filter set (EGFP,

Chroma). The EMCCD camera TTL signal triggered the LED. The 490nm intensities were

set to either 0.23mW/mm^2 with an exposure time of 50ms (Fig 3A, 4A-H, 8C-H),

0.33mW/mm^2 (Fig 6A-D) or 0.51 mW/mm^2 (Fig 6E-H), both with an exposure time of

5ms or 0.33mW/mm^2 with an exposure time of 10ms (PLM). Additional mKate2

fluorescence images were acquired for the RIS, RIM and PLM activity experiments. For

this measurement we used a 585nm LED and a set of standard Texas Red filters (Chroma).

The 585nm LED intensity was set to 0.11mW/mm^2 while imaging with a 50ms exposure

time (RIS) and to 0.15 mW/mm^2 for acquiring images with a 100ms exposure time (RIM,

PLM). Calcium and mKate2 imaging were carried out with a frame rate of 0.1Hz for 3h.

As the ion channels were additionally tagged with an mKate2, the parental strains without

the GCaMP/mKate2 construct were imaged with the same settings with the 585nm LED.

Here, no signal could be observed in the parental strains.

For the long-term calcium imaging from day 1 to day 12 we utilized slightly different LED

light intensities. Here, the 490nm intensity was set to 0.34mW/mm^2 and the 585nm light

had an intensity of 0.21mW/mm^2.

Optogenetic activation

For optogenetic RIS activation experiments, the agarose containing the microchambers

was supplemented with 10μl of 10mM all-trans-retinal (ATR, Sigma Aldrich) dissolved in

ethanol 1h before the start of image acquisition. In experiment in Fig 9, A baseline of

GCaMP images was recorded for 1min with a frame rate of 0.33Hz. This was followed by

a stimulation phase of 1min, in which again GCaMP images were acquired with a frame

rate of 0.33Hz. In between the camera exposures, the worms were illuminated with 585nm

LED light with an intensity of 0.17mW/mm^2 for 800ms triggered by the software. After

the stimulation period, a recovery period of 1min was imaged, in which GCaMP images

were acquired with a frame rate of 0.33Hz. The 490nm LED intensities for imaging were

set to 0.39mW/mm^2 and images were acquired with an exposure time of 5ms.

For the long-term optogenetic imaging worms (Fig 10), a stimulation phase of 11h was

recorded in which GCaMP images were acquired with a frame rate of 0.33Hz. In between

the camera exposures, the worms were illuminated with 585nm LED light with an intensity

of 0.17mW/mm^2 for 800ms triggered by the software.

Quantification of expression of mKate2 of the RIS activity strains

Larvae were synchronized via bleaching [17] and hatched in M9 buffer at 25°C. Imaging

of most strains was performed after 48h in the L1 arrest stage by transferring worms with

a 10 µl liquid drop onto an agarose pad (5% agarose solved in M9 buffer). The *RIS::unc-*

58gf and flp-11(-) worms were imaged 24h after bleaching. Larvae were then fixed through

the addition of 10 µl of 25mM levamisole (Sigma Aldrich).

To image the expression levels of the different RIS activity transgenes, mKate2 was

imaged with an Andor Revolution disc system (Andor Technology Ltd.) and a CSU-X1

spinning disc head (Yokogawa). The 561nm laser light intensity was set to 0.49mW/mm^2.

The worms were imaged through a 100x oil objective and by an Andor iXon Ultra EMCCD

camera (1024x1024 pixels) (Andor Technology Ltd., Belfast). An EM gain of 200 was

included and the camera read-out time was set to 30MHz. A z-stack of 21planes, each

0.5µm apart (10µm in total) was acquired. For comparison, a maximum z-stack projection

was calculated.

DIC image analysis

Mobility of worms was analyzed from DIC images though image subtraction as previously

described [17, 79]. The matrix of intensity values of the preceding image was subtracted

from the matrix of intensity of the following image. A mean of all values of the difference

matrix was then calculated to attain a mean smoothed image subtraction value, which

represents the movement of the worm.

To identify lethargus in fed worms, the non-pumping time prior to molting was identified.

For the sleep analysis in recovery lethargus after 12 days of starvation followed by

refeeding, 3h prior to molt were analyzed for mobility quiescence.

Analyzing functional Ca^{2+}-images

Neuronal intensities were automatically analyzed via custom-written MATLAB scripts

that are available at Github (https://github.com/ibusack/A-sleep-active-neuron-can-

promote-survival-while-sleep-behavior-is-

disturbed/blob/main/analysisfluorescentsignal.m). These codes extracted the regions of

interest through an intensity threshold and acquired the intensities and position data. The

extracted regions were set to be slightly bigger than the actual size of the neuron to allow

for background intensity detection and background subtraction. The pan-neuronal activity

was measured treating the head neurons as a combined signal. The RIS GCaMP activity in

Figure 6 was determined by extracting the position data of the corresponding mKate2

signal. A larger region of interest was extracted and a fitting percentage of that was

calculated as the signal. The extracted position data of the signal served to calculate the

speed values.

RIS, RIM and PLM activities were normalized by dividing GCaMP intensities through

corresponding mKate2 intensities. In the optogenetic experiment, the baseline

measurements were used for normalization.

Mobility quiescence detection

First, speeds or image subtraction mean intensities were smoothed over 30 time points

through the *smooth* function in MATLAB, which is a first-degree polynomial local

regression model. The code is available at Github (https://github.com/ibusack/A-sleep-

active-neuron-can-promote-survival-while-sleep-behavior-is-

disturbed/blob/main/boutdetection.m). Smoothed speeds and image subtraction values

were normalized between 0 and 1 for each individual animal with 0 being the lowest speed

value and 1 the highest speed value the animal possessed. A mobility quiescence bout was

counted when the normalized speed was below a specific percentage for a certain amount

of time. For calcium imaging experiments with the 20x objective (Fig 6), the normalized

speed had to be below 10% for at least 3min. The RIS activity experiment, which was

filmed with a 10x objective, required a percentage threshold of 5% for a minimum of 4min.

For DIC images filmed with the 20x objective, the normalized mean image subtraction

values had to be below 15% for at least 3min to be counted as sleep time points. The code

for detecting sleep bouts based on RIM tracking can be found at Github

(https://github.com/ibusack/A-sleep-active-neuron-can-promote-survival-while-sleep-

behavior-is-disturbed/blob/main/RIM_sleepboutdetection.m.)

Mobility quiescence bout onset alignment

Speeds and GCaMP intensities were aligned to the start of mobility quiescence bouts, which had been detected by the mobility quiescence bout analysis. Mobility quiescence bouts were only counted for the alignment when the worm was mobile throughout 3min prior to the mobility quiescence bout onset. This alignment was carried out with a custom-written MATLAB script which is published on Github (https://github.com/ibusack/A-sleep-active-neuron-can-promote-survival-while-sleep-behavior-is-disturbed/blob/main/boutdetection.m).

Neuronal inactivity detection

The neuronal activities were treated similarly as speed values for the mobility quiescence detection. Hence the boutdetection MATLAB script was utilized, which can be found on Github (https://github.com/ibusack/A-sleep-active-neuron-can-promote-survival-while-sleep-behavior-is-disturbed/blob/main/boutdetection.m).

The nervous system was counted as inactive when it was below 20% of the smoothed and normalized activity for at least 3min.

RIM neuron inactivity detection

RIM peaks were detected through a custom-written MATLAB script, which is available at Github (https://github.com/ibusack/A-sleep-active-neuron-can-promote-survival-while-sleep-behavior-is-disturbed/blob/main/analyze_RIM.m). First, the activity values were smoothed using the inbuilt *smooth* function over 5 time points. Then, peaks of a minimum prominence of 0.2 were extracted via the inbuilt function *islocalmax*. RIM was counted as inactive when there were no detected peaks for at least 5min.

Induction of cellular stress by heat shock

The heat shock was carried out following a previously described protocol and utilizing a previously described heat shock device [17]. To summarize, young adult worms were imaged in 370x370x45μm microfluidic chambers and for a total of 4h. 5 to 12 worms were transferred from plates grown at 20°C to the microchamber. The worms were starved for

90min at 20°C before filming. During the entire duration of the movie, DIC images were

acquired at a frequency of 0.1Hz. First, a baseline was recorded for 1h with the sample

holder temperature set to 22°C and the heating lid set to 25°C. Then the heat shock was

conducted with the sample holder temperature set to 37°C and the heating lid set to 39°C

for 20 minutes. Finally, a recovery phase was recorded with the sample holder temperature

set to 22°C and the heating lid set to 25°C for 2h 40min.

Pumping assay

Arrested L1 worms were imaged in 110x110x10μm microfluidic chambers. DIC images

were acquired with a frequency of 10 Hz for 1min. Pumps of the grinder were then

counted manually with the help of a clicker.

Survival assay in liquid culture

RIS impairment has been shown to confer a reduction of survival in both liquid culture and

in microfluidic chambers [12, 23, 24, 36]. Worms were synchronized by bleaching [17]. 3-

4 days before bleaching, 10-12 L4 stage worms were transferred onto NGM plates with a

diameter of 6cm. 1-2 plates were bleached per strain and the day of bleaching was defined

as day 0. The worms were kept in 1ml of M9 buffer in 2ml Eppendorf tubes in a rotator at

20°C for most survival assays. The following day was counted as day 1 of the lifespan.

During the survival assay, the worms were kept in liquid culture in 1ml M9 buffer in 2ml

Eppendorf tubes. This resulted in a population density of approximately 100 worms per

10μl. The tubes were placed in a rotator at 20°C, which rotated at 20rpm. For counting,

10μl drops of the liquid culture were pipetted onto fresh NGM plates for each strain

respectively. The worms were allowed to recover for at least 2h after pipetting before

counting. Counting was performed manually with the assistance of a clicker. The recovery

number, i.e., the number of worms that developed into at least the L4 stage, was counted

3-5 days after pipetting. Around 50-100 worms were counted for each strain of each

replicate on each time point.

Survival assay in the OptoGenBox

For the long-term optogenetic survival assay worms were kept in microfluidic chambers

(110x110x15μm) as described above. Between 27 and 43 worms were used for each strain

per agarose slab each containing many microfluidic chambers. A maximum of one worm

was placed per microfluidic chamber. Three replicates were performed per condition. We

then calculated the average of all individuals of all three replicates combined. For the

optogenetic activation 10μl of 10mM all-trans-retinal (ATR, Sigma Aldrich) dissolved in

ethanol was added every 3-4 days. 20μl of 10μg/ml nystatin dissolved in DMSO was added

3-4 times throughout the lifespan to prevent fungal contamination. To prevent drying out

of the agarose, 20μl of sterile water was added every 2 days. In the first week, the worms

were counted every second day, then the survival was counted every day until all worms

had died. A worm was counted as dead if a 2 min blue-light stimulus did not cause a visible

behavioral response.

For long-term optogenetic activation of RIS, worms were placed in the OptoGenBox [23]

and continuously illuminated with 10mW at 20°C.

Blue light survival assay

For the blue-light survival assays worms were imaged in 110x110x10μm microchambers.

A blue light stimulus of $0.74mW/mm^2$ was given for 1s every 5s while simultaneously

taking a GCaMP image. For the control, the same intensity stimulus was given for only

100ms through the camera exposure time.

Imaging of growing worms

The images of growing plates were taken at a Leica M125 stereomicroscope. A Jenoptik

ProgRes MF captured the images utilizing the corresponding software ProgResCapturePro

2.10.0.1.

RNA sequencing data

RNA sequencing data is described in [24].

Fitting

Origin software was used for the regression analyses. A linear regression was fitted to the
data in Figure 6C and 6G. R^2 values can be found in the corresponding figures. The exact
functions are in Table 1.

Figure panel	fit	function
6C	linear	$y = 0.116 + 0.5494x$
6G	linear	$y = 0.21203 + 1.10949x$

Table 1. Calculated functions for the regressions.

Quantification and Statistical Analysis

The experimenter was blind to the genotype information during the lifespan experiments.
All other experiments relied on an automated analysis and hence were not blinded. The
Wilcoxon signed rank test was utilized for the comparison of speeds and GCaMP
intensities within the same strain. The Welch test was conducted to compare data from
different strains. The survival assay experiments were statistically analyzed by utilizing the
Fisher's exact test at 50% survival of the wild type. Experiments with four or more
genotypes were FDR corrected by the Benjamini-Hochberg procedure with a 5% false
discovery rate. The graphs depict the mean and standard error of the mean. Box plots
display individual data points, the interquartile range and the median. The whiskers
represent the 10-90[th] percentile. A table listing all exact p-values can be found in the
supplemental information (Table S4).

Acknowledgements

Some strains were provided by the CGC, which is funded by NIH Office of Research
Infrastructure Programs (P40 OD010440). The CRISPR strains were generated by
SunyBiotech based on our design. We thank Thomas Boulin for communicating the sodium
permeability of *unc-58gf* and suggesting its use as a tool for neuronal activation. This work
was supported by a European Research Council Starting Grant (ID: 637860,
SLEEPCONTROL).

Author contributions

I.B.: Data curation, Formal analysis, Investigation, Methodology, Project administration,

Resources, Software, Visualization, Writing – review & editing.

H.B.: Conceptualization, Funding acquisition, Investigation, Methodology, Project

administration, Resources, Supervision, Writing – original draft.

1005

1006 **Declaration of Interest**

1007 The authors declare no competing interests.

1008   **References**

1009

1.   Campbell SS, Tobler I. Animal sleep: a review of sleep duration across
phylogeny. Neurosci Biobehav Rev. 1984;8(3):269-300. Epub 1984/01/01. PubMed
PMID: 6504414.
2.   Diekelmann S, Born J. The memory function of sleep. Nature reviews
Neuroscience. 2010;11(2):114-26. Epub 2010/01/05. doi: 10.1038/nrn2762. PubMed
PMID: 20046194.
3.   Krueger JM, Frank MG, Wisor JP, Roy S. Sleep function: Toward elucidating an
enigma. Sleep medicine reviews. 2016;28:46-54. Epub 2015/10/09. doi:
10.1016/j.smrv.2015.08.005. PubMed PMID: 26447948; PubMed Central PMCID:
PMC4769986.
4.   Anafi RC, Kayser MS, Raizen DM. Exploring phylogeny to find the function of
sleep. Nature reviews Neuroscience. 2019;20(2):109-16. Epub 2018/12/24. doi:
10.1038/s41583-018-0098-9. PubMed PMID: 30573905.
5.   Hublin C, Partinen M, Koskenvuo M, Kaprio J. Sleep and mortality: a population-
based 22-year follow-up study. Sleep. 2007;30(10):1245-53. Epub 2007/11/01. doi:
10.1093/sleep/30.10.1245. PubMed PMID: 17969458; PubMed Central PMCID:
PMCPMC2266277.
6.   Van Cauter E, Spiegel K, Tasali E, Leproult R. Metabolic consequences of sleep
and sleep loss. Sleep Med. 2008;9 Suppl 1:S23-8. Epub 2008/12/17. doi: 10.1016/S1389-
9457(08)70013-3. PubMed PMID: 18929315; PubMed Central PMCID:
PMCPMC4444051.
7.   Irwin MR. Why sleep is important for health: a psychoneuroimmunology
perspective. Annu Rev Psychol. 2015;66:143-72. Epub 2014/07/26. doi:
10.1146/annurev-psych-010213-115205. PubMed PMID: 25061767; PubMed Central
PMCID: PMCPMC4961463.
8.   Saper CB, Fuller PM, Pedersen NP, Lu J, Scammell TE. Sleep state switching.
Neuron. 2010;68(6):1023-42. Epub 2010/12/22. doi: 10.1016/j.neuron.2010.11.032.
PubMed PMID: 21172606; PubMed Central PMCID: PMC3026325.
9.   Bringmann H. Sleep-Active Neurons: Conserved Motors of Sleep. Genetics.
2018;208(4):1279-89. Epub 2018/04/06. doi: 10.1534/genetics.117.300521. PubMed
PMID: 29618588; PubMed Central PMCID: PMC5887131.
10.   Turek M, Lewandrowski I, Bringmann H. An AP2 transcription factor is required
for a sleep-active neuron to induce sleep-like quiescence in C. elegans. Curr Biol.
2013;23(22):2215-23. Epub 2013/11/05. doi: 10.1016/j.cub.2013.09.028. PubMed PMID:
24184105.
11.   Turek M, Besseling J, Spies JP, Konig S, Bringmann H. Sleep-active neuron
specification and sleep induction require FLP-11 neuropeptides to systemically induce
sleep. eLife. 2016;5. Epub 2016/03/08. doi: 10.7554/eLife.12499. PubMed PMID:
26949257; PubMed Central PMCID: PMC4805538.
12.   Wu Y, Masurat F, Preis J, Bringmann H. Sleep Counteracts Aging Phenotypes to
Survive Starvation-Induced Developmental Arrest in C. elegans. Curr Biol.
2018;28(22):3610-24 e8. Epub 2018/11/13. doi: 10.1016/j.cub.2018.10.009. PubMed
PMID: 30416057; PubMed Central PMCID: PMCPMC6264389.

13.   Skora S, Mende F, Zimmer M. Energy Scarcity Promotes a Brain-wide Sleep
State Modulated by Insulin Signaling in C. elegans. Cell reports. 2018;22(4):953-66.
Epub 2018/02/02. doi: 10.1016/j.celrep.2017.12.091. PubMed PMID: 29386137.
14.   Gonzales DL, Zhou J, Fan B, Robinson JT. A microfluidic-induced C. elegans
sleep state. Nat Commun. 2019;10(1):5035. Epub 2019/11/07. doi: 10.1038/s41467-019-
13008-5. PubMed PMID: 31695031.
15.   Steuer Costa W, Van der Auwera P, Glock C, Liewald JF, Bach M, Schuler C, et
al. A GABAergic and peptidergic sleep neuron as a locomotion stop neuron with
compartmentalized Ca2+ dynamics. Nat Commun. 2019;10(1):4095. Epub 2019/09/12.
doi: 10.1038/s41467-019-12098-5. PubMed PMID: 31506439.
16.   Grubbs JJ, Lopes LE, van der Linden AM, Raizen DM. A salt-induced kinase is
required for the metabolic regulation of sleep. PLoS biology. 2020;18(4):e3000220. Epub
20200421. doi: 10.1371/journal.pbio.3000220. PubMed PMID: 32315298; PubMed
Central PMCID: PMCPMC7173979.
17.   Konietzka J, Fritz M, Spiri S, McWhirter R, Leha A, Palumbos S, et al. Epidermal
Growth Factor Signaling Promotes Sleep through a Combined Series and Parallel Neural
Circuit. Curr Biol. 2020;30(1):1-16 e3. Epub 2019/12/17. doi:
10.1016/j.cub.2019.10.048. PubMed PMID: 31839447.
18.   Sinner MP, Masurat F, Ewbank JJ, Pujol N, Bringmann H. Innate Immunity
Promotes Sleep through Epidermal Antimicrobial Peptides. Curr Biol. 2021;31(3):564-77
e12. Epub 2020/12/02. doi: 10.1016/j.cub.2020.10.076. PubMed PMID: 33259791.
19.   Makino M, Ulzii E, Shirasaki R, Kim J, You YJ. Regulation of Satiety
Quiescence by Neuropeptide Signaling in Caenorhabditis elegans. Front Neurosci.
2021;15:678590. Epub 2021/08/03. doi: 10.3389/fnins.2021.678590. PubMed PMID:
34335159; PubMed Central PMCID: PMCPMC8319666.
20.   Nichols ALA, Eichler T, Latham R, Zimmer M. A global brain state underlies C.
elegans sleep behavior. Science. 2017;356(6344). Epub 2017/06/24. doi:
10.1126/science.aam6851. PubMed PMID: 28642382.
21.   Xu Y, Zhang L, Liu Y, Topalidou I, Hassinan C, Ailion M, et al. Dopamine
receptor DOP-1 engages a sleep pathway to modulate swimming in C. elegans. iScience.
2021;24(4):102247. Epub 2021/04/03. doi: 10.1016/j.isci.2021.102247. PubMed PMID:
33796839; PubMed Central PMCID: PMCPMC7995527.
22.   Maluck E, Busack I, Besseling J, Masurat F, Turek M, Busch KE, et al. A wake-
active locomotion circuit depolarizes a sleep-active neuron to switch on sleep. PLoS
biology. 2020;18(2):e3000361. Epub 2020/02/23. doi: 10.1371/journal.pbio.3000361.
PubMed PMID: 32078631.
23.   Busack I, Jordan F, Sapir P, Bringmann H. The OptoGenBox - a device for long-
term optogenetics in C. elegans. J Neurogenet. 2020:1-9. Epub 2020/06/17. doi:
10.1080/01677063.2020.1776709. PubMed PMID: 32543249.
24.   Koutsoumparis A, Welp LM, Wulf A, Urlaub H, Meierhofer D, Börno S, et al.
Sleep neuron depolarization promotes protective gene expression changes and FOXO
activation. Current Biology. 2022. doi: https://doi.org/10.1016/j.cub.2022.04.012.
25.   Kenyon CJ. The genetics of ageing. Nature. 2010;464(7288):504-12. Epub
2010/03/26. doi: 10.1038/nature08980. PubMed PMID: 20336132.
26.   Baugh LR, Hu PJ. Starvation Responses Throughout the Caenorhabditis elegans
Life Cycle. Genetics. 2020;216(4):837-78. Epub 2020/12/04. doi:

10.1534/genetics.120.303565. PubMed PMID: 33268389; PubMed Central PMCID:
PMCPMC7768255.
27.   Siegel JM. Sleep viewed as a state of adaptive inactivity. Nature reviews
Neuroscience. 2009;10(10):747-53. Epub 2009/08/06. doi: 10.1038/nrn2697. PubMed
PMID: 19654581.
28.   Schmidt MH. The energy allocation function of sleep: a unifying theory of sleep,
torpor, and continuous wakefulness. Neurosci Biobehav Rev. 2014;47:122-53. Epub
2014/08/15. doi: 10.1016/j.neubiorev.2014.08.001. PubMed PMID: 25117535.
29.   Krause AJ, Simon EB, Mander BA, Greer SM, Saletin JM, Goldstein-Piekarski
AN, et al. The sleep-deprived human brain. Nature reviews Neuroscience.
2017;18(7):404-18. Epub 2017/05/19. doi: 10.1038/nrn.2017.55. PubMed PMID:
28515433.
30.   Klinzing JG, Niethard N, Born J. Mechanisms of systems memory consolidation
during sleep. Nat Neurosci. 2019;22(10):1598-610. Epub 2019/08/28. doi:
10.1038/s41593-019-0467-3. PubMed PMID: 31451802.
31.   Taylor SR, Santpere G, Weinreb A, Barrett A, Reilly MB, Xu C, et al. Molecular
topography of an entire nervous system. Cell. 2021. Epub 2021/07/09. doi:
10.1016/j.cell.2021.06.023. PubMed PMID: 34237253.
32.   Ben Soussia I, El Mouridi S, Kang D, Leclercq-Blondel A, Khoubza L, Tardy P,
et al. Mutation of a single residue promotes gating of vertebrate and invertebrate two-
pore domain potassium channels. Nat Commun. 2019;10(1):787. Epub 2019/02/17. doi:
10.1038/s41467-019-08710-3. PubMed PMID: 30770809; PubMed Central PMCID:
PMCPMC6377628.
33.   Kunkel MT, Johnstone DB, Thomas JH, Salkoff L. Mutants of a temperature-
sensitive two-P domain potassium channel. J Neurosci. 2000;20(20):7517-24. doi:
10.1523/JNEUROSCI.20-20-07517.2000. PubMed PMID: 11027209; PubMed Central
PMCID: PMCPMC6772866.
34.   Silva-Garcia CG, Lanjuin A, Heintz C, Dutta S, Clark NM, Mair WB. Single-
Copy Knock-In Loci for Defined Gene Expression in Caenorhabditis elegans. G3
(Bethesda). 2019;9(7):2195-8. Epub 2019/05/09. doi: 10.1534/g3.119.400314. PubMed
PMID: 31064766; PubMed Central PMCID: PMCPMC6643881.
35.   Bringmann H. Genetic sleep deprivation: using sleep mutants to study sleep
functions. EMBO Rep. 2019;20(3). Epub 2019/02/26. doi: 10.15252/embr.201846807.
PubMed PMID: 30804011; PubMed Central PMCID: PMCPMC6399599.
36.   Roux AE, Langhans K, Huynh W, Kenyon C. Reversible Age-Related
Phenotypes Induced during Larval Quiescence in C. elegans. Cell Metab.
2016;23(6):1113-26. Epub 2016/06/16. doi: 10.1016/j.cmet.2016.05.024. PubMed PMID:
27304510.
37.   Bringmann H. Agarose hydrogel microcompartments for imaging sleep- and
wake-like behavior and nervous system development in Caenorhabditis elegans larvae. J
Neurosci Methods. 2011;201(1):78-88. Epub 2011/08/02. doi: S0165-0270(11)00418-3
[pii]10.1016/j.jneumeth.2011.07.013. PubMed PMID: 21801751.
38.   Turek M, Besseling J, Bringmann H. Agarose Microchambers for Long-term
Calcium Imaging of Caenorhabditis elegans. Journal of visualized experiments : JoVE.
2015;(100):e52742. Epub 2015/07/02. doi: 10.3791/52742. PubMed PMID: 26132740;
PubMed Central PMCID: PMC4544933.

39.   Van Buskirk C, Sternberg PW. Epidermal growth factor signaling induces
behavioral quiescence in Caenorhabditis elegans. Nat Neurosci. 2007;10(10):1300-7.
Epub 2007/09/25. doi: 10.1038/nn1981. PubMed PMID: 17891142.
40.   Hill AJ, Mansfield R, Lopez JM, Raizen DM, Van Buskirk C. Cellular stress
induces a protective sleep-like state in C. elegans. Curr Biol. 2014;24(20):2399-405.
Epub 2014/09/30. doi: 10.1016/j.cub.2014.08.040. PubMed PMID: 25264259; PubMed
Central PMCID: PMC4254280.
41.   Nelson MD, Lee KH, Churgin MA, Hill AJ, Van Buskirk C, Fang-Yen C, et al.
FMRFamide-like FLP-13 neuropeptides promote quiescence following heat stress in
Caenorhabditis elegans. Curr Biol. 2014;24(20):2406-10. Epub 2014/09/30. doi:
10.1016/j.cub.2014.08.037. PubMed PMID: 25264253; PubMed Central PMCID:
PMC4254296.
42.   Iannacone MJ, Beets I, Lopes LE, Churgin MA, Fang-Yen C, Nelson MD, et al.
The RFamide receptor DMSR-1 regulates stress-induced sleep in C. elegans. eLife.
2017;6. Epub 2017/01/18. doi: 10.7554/eLife.19837. PubMed PMID: 28094002; PubMed
Central PMCID: PMCPMC5241116.
43.   Cassada RC, Russell RL. The dauerlarva, a post-embryonic developmental variant
of the nematode Caenorhabditis elegans. Dev Biol. 1975;46(2):326-42. Epub 1975/10/01.
doi: 0012-1606(75)90109-8 [pii]. PubMed PMID: 1183723.
44.   Raizen DM, Zimmerman JE, Maycock MH, Ta UD, You YJ, Sundaram MV, et
al. Lethargus is a Caenorhabditis elegans sleep-like state. Nature. 2008;451(7178):569-
72. Epub 2008/01/11. doi: nature06535 [pii]10.1038/nature06535. PubMed PMID:
18185515.
45.   Yemini E, Lin A, Nejatbakhsh A, Varol E, Sun R, Mena GE, et al. NeuroPAL: A
Multicolor Atlas for Whole-Brain Neuronal Identification in C. elegans. Cell.
2021;184(1):272-88 e11. Epub 2020/12/31. doi: 10.1016/j.cell.2020.12.012. PubMed
PMID: 33378642.
46.   Piggott BJ, Liu J, Feng Z, Wescott SA, Xu XZ. The neural circuits and synaptic
mechanisms underlying motor initiation in C. elegans. Cell. 2011;147(4):922-33. Epub
2011/11/15. doi: 10.1016/j.cell.2011.08.053. PubMed PMID: 22078887; PubMed Central
PMCID: PMC3233480.
47.   Li W, Kang L, Piggott BJ, Feng Z, Xu XZ. The neural circuits and sensory
channels mediating harsh touch sensation in Caenorhabditis elegans. Nat Commun.
2011;2:315. Epub 2011/05/19. doi: 10.1038/ncomms1308. PubMed PMID: 21587232;
PubMed Central PMCID: PMC3098610.
48.   McCormick DA, Bal T. Sensory gating mechanisms of the thalamus. Curr Opin
Neurobiol. 1994;4(4):550-6. doi: 10.1016/0959-4388(94)90056-6. PubMed PMID:
7812144.
49.   Schwarz J, Lewandrowski I, Bringmann H. Reduced activity of a sensory neuron
during a sleep-like state in Caenorhabditis elegans. Curr Biol. 2011;21(24):R983-4. Epub
2011/12/24. doi: S0960-9822(11)01207-3[pii]10.1016/j.cub.2011.10.046. PubMed
PMID: 22192827.
50.   Schwarz J, Bringmann H. Reduced sleep-like quiescence in both hyperactive and
hypoactive mutants of the Galphaq Gene egl-30 during lethargus in Caenorhabditis
elegans. PLoS One. 2013;8(9):e75853. Epub 2013/09/28. doi:

10.1371/journal.pone.0075853. PubMed PMID: 24073282; PubMed Central PMCID:
PMC3779211.
51.   Cho JY, Sternberg PW. Multilevel modulation of a sensory motor circuit during
C. elegans sleep and arousal. Cell. 2014;156(1-2):249-60. Epub 2014/01/21. doi:
10.1016/j.cell.2013.11.036. PubMed PMID: 24439380; PubMed Central PMCID:
PMC3962823.
52.   Schwarz J, Spies JP, Bringmann H. Reduced muscle contraction and a relaxed
posture during sleep-like Lethargus. Worm. 2012;1(1):12-4. Epub 2012/01/01. doi:
10.4161/worm.19499. PubMed PMID: 24058817; PubMed Central PMCID:
PMC3670164.
53.   Iwanir S, Tramm N, Nagy S, Wright C, Ish D, Biron D. The microarchitecture of
C. elegans behavior during lethargus: homeostatic bout dynamics, a typical body posture,
and regulation by a central neuron. sleep. 2013;36(3):385-95. Epub 2013/03/02. doi:
10.5665/sleep.2456. PubMed PMID: 23449971; PubMed Central PMCID: PMC3571756.
54.   Tramm N, Oppenheimer N, Nagy S, Efrati E, Biron D. Why do sleeping
nematodes adopt a hockey-stick-like posture? PLoS One. 2014;9(7):e101162. Epub
2014/07/16. doi: 10.1371/journal.pone.0101162. PubMed PMID: 25025212; PubMed
Central PMCID: PMC4099128.
55.   Schwartz HT. A protocol describing pharynx counts and a review of other assays
of apoptotic cell death in the nematode worm Caenorhabditis elegans. Nature protocols.
2007;2(3):705-14. doi: 10.1038/nprot.2007.93. PubMed PMID: 17406633.
56.   Song BM, Avery L. The pharynx of the nematode C. elegans: A model system for
the study of motor control. Worm. 2013;2(1):e21833. doi: 10.4161/worm.21833. PubMed
PMID: 24058858; PubMed Central PMCID: PMCPMC3670459.
57.   Urmersbach B, Besseling J, Spies JP, Bringmann H. Automated analysis of sleep
control via a single neuron active at sleep onset in C. elegans. Genesis. 2016;54(4):212-9.
Epub 2016/02/03. doi: 10.1002/dvg.22924. PubMed PMID: 26833569.
58.   Bennett HL, Khoruzhik Y, Hayden D, Huang H, Sanders J, Walsh MB, et al.
Normal sleep bouts are not essential for C. elegans survival and FoxO is important for
compensatory changes in sleep. BMC neuroscience. 2018;19(1):10. Epub 2018/03/11.
doi: 10.1186/s12868-018-0408-1. PubMed PMID: 29523076.
59.   White JG, Southgate E, Thomson JN, Brenner S. The structure of the nervous
system of the nematode Caenorhabditis elegans. Philos Trans R Soc Lond B Biol Sci.
1986;314(1165):1-340. Epub 1986/11/12. PubMed PMID: 22462104.
60.   Schafer WR, Kenyon CJ. A calcium-channel homologue required for adaptation
to dopamine and serotonin in Caenorhabditis elegans. Nature. 1995;375(6526):73-8.
Epub 1995/05/04. doi: 10.1038/375073a0. PubMed PMID: 7723846.
61.   Lee RY, Lobel L, Hengartner M, Horvitz HR, Avery L. Mutations in the alpha1
subunit of an L-type voltage-activated Ca2+ channel cause myotonia in Caenorhabditis
elegans. EMBO J. 1997;16(20):6066-76. Epub 1997/10/08. doi:
10.1093/emboj/16.20.6066. PubMed PMID: 9321386.
62.   Mathews EA, Garcia E, Santi CM, Mullen GP, Thacker C, Moerman DG, et al.
Critical residues of the Caenorhabditis elegans unc-2 voltage-gated calcium channel that
affect behavioral and physiological properties. J Neurosci. 2003;23(16):6537-45. doi:
10.1523/JNEUROSCI.23-16-06537.2003. PubMed PMID: 12878695; PubMed Central
PMCID: PMCPMC6740628.

63.   Laine V, Frokjaer-Jensen C, Couchoux H, Jospin M. The alpha1 subunit EGL-19,
the alpha2/delta subunit UNC-36, and the beta subunit CCB-1 underlie voltage-
dependent calcium currents in Caenorhabditis elegans striated muscle. The Journal of
biological chemistry. 2011;286(42):36180-7. Epub 2011/09/01. doi:
10.1074/jbc.M111.256149. PubMed PMID: 21878625; PubMed Central PMCID:
PMC3196126.
64.   Schuler C, Fischer E, Shaltiel L, Steuer Costa W, Gottschalk A. Arrhythmogenic
effects of mutated L-type Ca 2+-channels on an optogenetically paced muscular pump in
Caenorhabditis elegans. Sci Rep. 2015;5:14427. Epub 20150924. doi:
10.1038/srep14427. PubMed PMID: 26399900; PubMed Central PMCID:
PMCPMC4585839.
65.   Huang YC, Pirri JK, Rayes D, Gao S, Mulcahy B, Grant J, et al. Gain-of-function
mutations in the UNC-2/CaV2alpha channel lead to excitation-dominant synaptic
transmission in Caenorhabditis elegans. eLife. 2019;8. Epub 20190805. doi:
10.7554/eLife.45905. PubMed PMID: 31364988; PubMed Central PMCID:
PMCPMC6713474.
66.   Lin JY, Knutsen PM, Muller A, Kleinfeld D, Tsien RY. ReaChR: a red-shifted
variant of channelrhodopsin enables deep transcranial optogenetic excitation. Nat
Neurosci. 2013;16(10):1499-508. Epub 2013/09/03. doi: 10.1038/nn.3502. PubMed
PMID: 23995068; PubMed Central PMCID: PMC3793847.
67.   Driver RJ, Lamb AL, Wyner AJ, Raizen DM. DAF-16/FOXO Regulates
Homeostasis of Essential Sleep-like Behavior during Larval Transitions in C. elegans.
Curr Biol. 2013;23(6):501-6. Epub 2013/03/13. doi: 10.1016/j.cub.2013.02.009. PubMed
PMID: 23477722.
68.   Choi S, Chatzigeorgiou M, Taylor KP, Schafer WR, Kaplan JM. Analysis of
NPR-1 reveals a circuit mechanism for behavioral quiescence in C. elegans. Neuron.
2013;78(5):869-80. Epub 2013/06/15. doi: 10.1016/j.neuron.2013.04.002. PubMed
PMID: 23764289; PubMed Central PMCID: PMC3683153.
69.   Spies J, Bringmann H. Automated detection and manipulation of sleep in C.
elegans reveals depolarization of a sleep-active neuron during mechanical stimulation-
induced sleep deprivation. Sci Rep. 2018;8(1):9732. Epub 2018/06/29. doi:
10.1038/s41598-018-28095-5. PubMed PMID: 29950594.
70.   Stroustrup N, Ulmschneider BE, Nash ZM, Lopez-Moyado IF, Apfeld J, Fontana
W. The Caenorhabditis elegans Lifespan Machine. Nat Methods. 2013;10(7):665-70.
Epub 20130512. doi: 10.1038/nmeth.2475. PubMed PMID: 23666410; PubMed Central
PMCID: PMCPMC3865717.
71.   Laurent P, Soltesz Z, Nelson GM, Chen C, Arellano-Carbajal F, Levy E, et al.
Decoding a neural circuit controlling global animal state in C. elegans. eLife. 2015;4.
Epub 20150311. doi: 10.7554/eLife.04241. PubMed PMID: 25760081; PubMed Central
PMCID: PMCPMC4440410.
72.   Chun L, Gong J, Yuan F, Zhang B, Liu H, Zheng T, et al. Metabotropic GABA
signalling modulates longevity in C. elegans. Nat Commun. 2015;6:8828. Epub
20151105. doi: 10.1038/ncomms9828. PubMed PMID: 26537867; PubMed Central
PMCID: PMCPMC4667614.
73.   Nagy S, Tramm N, Sanders J, Iwanir S, Shirley IA, Levine E, et al. Homeostasis
in C. elegans sleep is characterized by two behaviorally and genetically distinct

mechanisms. eLife. 2014;3:e04380. Epub 2014/12/05. doi: 10.7554/eLife.04380.
PubMed PMID: 25474127; PubMed Central PMCID: PMC4273442.
74.   Alam MA, Kumar S, McGinty D, Alam MN, Szymusiak R. Neuronal activity in
the preoptic hypothalamus during sleep deprivation and recovery sleep. J Neurophysiol.
2014;111(2):287-99. Epub 2013/11/01. doi: 10.1152/jn.00504.2013. PubMed PMID:
24174649; PubMed Central PMCID: PMC3921380.
75.   Brenner S. The genetics of Caenorhabditis elegans. Genetics. 1974;77(1):71-94.
Epub 1974/05/01. PubMed PMID: 4366476.
76.   Stiernagle T. Maintenance of C. elegans. WormBook. 2006:1-11. Epub
2007/12/01. doi: 10.1895/wormbook.1.101.1. PubMed PMID: 18050451.
77.   Ahringer A. Reverse genetics. WormBook, ed The C elegans Research
Community. 2006. doi: doi/10.1895/wormbook.1.47.1.
78.   Redemann S, Schloissnig S, Ernst S, Pozniakowsky A, Ayloo S, Hyman AA, et
al. Codon adaptation-based control of protein expression in C. elegans. Nat Methods.
2011;8(3):250-2. Epub 2011/02/01. doi: nmeth.1565 [pii]10.1038/nmeth.1565. PubMed
PMID: 21278743.
79.   Nagy S, Raizen DM, Biron D. Measurements of behavioral quiescence in
Caenorhabditis elegans. Methods. 2014. Epub 2014/03/20. doi:
10.1016/j.ymeth.2014.03.009. PubMed PMID: 24642199.
1301
1302

Figure Legends

Fig 1 – Strains expressing ion channel mutants in RIS

A) Scheme of the genetic design of the different RIS potassium channel mutant transgenes.

B-D) Localization and expression of mKate2, which is translationally fused to the ion channels, in the different inactivation strains.

E) Schematic representation of the genetic design of the RIS sodium channel mutant transgenes.

F) Quantification of the expression levels of UNC-58gf::mKate2 in the two RIS overactivation strains. ***p<0.001, Welch test.

G) Localization and expression of the strong overactivation strain (*RIS::unc-58gf(strong)*).

H) Localization and expression of the weak overactivation strain (*RIS::unc-58gf(weak)*).

Fig 2 – Survival of starvation-induced L1 arrest is a function of RIS activity

A) *RIS::twk-18gf* and *RIS::unc-58gf(strong)* strains cause divergent transcriptional profiles. Shown is the log2 fold change for all differentially expressed genes. A gene was counted as differentially expressed with an FDR<0.05. According to our previously published results, *RIS::twk-18gf* expressed 498 genes differentially compared with the wild type. *RIS::twk-18gf* has been shown to be similar to *aptf-1(gk794)* [24]. *RIS::unc-58gf(strong)* was shown to express 2913 genes differentially compared with the wild type [24]. Comparing *RIS::twk-18gf* and *RIS::unc-58gf(strong)* revealed 4676 differentially expressed genes. This analysis supports the view that these two strains have opposing transcriptional profiles and identifies 1910 additional genes that are potentially under the control of RIS activity (Table S5).

B) Strong inactivation significantly decreases survival in L1 arrest whereas strong activation prolongs survival (three-parameter logistic fit, see also supplementary table S1, Fisher's Exact Test was conducted when 50% of wild-type worms were alive (day 22)). The p-values were FDR corrected with Benjamini-Hochberg procedure with a 5% false discovery rate. The plot includes data from three replicates. ***p<0.001.

C) Recovery from L1 arrest depends on RIS activity with strong inactivation leading to a

reduced recovery and constant activation leading to a prolonged ability to recover. For

statistics, the Fisher's Exact Test was conducted at the time when 50% of wild-type

worms recovered (day 23). The p-values were FDR corrected with Benjamini-Hochberg

procedure with a 5% false discovery rate. The plot includes data from two replicates. N.s.

$p > 0.05$, *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$. Fisher's Exact Test.

1340

Fig 3 – Ion channel mutant expression in RIS controls baseline RIS activity

A) Ion channel expression in RIS results in a dose-response strain set with varying levels

of RIS baseline activity. n.s. $p > 0.05$, **$p < 0.001$, ***$p < 0.001$, Welch test with FDR

correction for multiple testing.

B) Long term RIS baseline activity for RIS activity strains from day 1 to day 12 of

starvation.

1347
1348 **Fig 4 - Both, strong inactivation of RIS as well as *RIS::unc-58gf(strong)* prevent**

1349 **sleep behavior during L1 arrest**

1350

A-F) Ion channel expression in RIS controls calcium activity and sleep behavior.

Mobility quiescence bouts (grey shade) correlate with RIS transients. Movement speed

was calculated by tracking the position of the head neuron RIS. A) *RIS::twk-18gf* and B)

RIS::egl-23gf(strong) larvae do not have detectable mobility quiescence bouts or RIS

transients. C) *RIS::egl-23gf(weak)* has reduced mobility quiescence bouts. D) Wild-type

RIS activation transients and immobility bouts. E) *RIS::unc-58gf(weak)* spends more

time in mobility quiescence bouts. F) *RIS::unc-58gf(strong)* has no detectable mobility

quiescence bouts and no RIS transients but a high constant activity.

G) RIS activation and immobility are correlated only for the strains that also show

sleeping behavior. Alignment of RIS activity and speed to mobility quiescence bout

onset. **$p < 0.01$, ***$p < 0.001$, Wilcoxon signed rank test.

H) Mobility quiescence is a function of RIS activity levels n.s. $p > 0.05$, **$p < 0.001$,

***$p < 0.001$, Welch test with FDR correction for multiple testing.

I) Long-term sleep fraction for RIS activity strains from day 1 to day 12 of starvation.

1365

Fig 5 *RIS::unc-58gf* impairs sleep following cellular stress and during the L1 molt in developing larvae

A-F) Sleep fraction of worms of different RIS activity strains during the heat shock experiment. Animals were recorded continuously throughout the experiment. First, a 1h baseline was recorded at 22°C, then a 20min heat shock period of 37°C (in red shade) was applied, after the heat shock, the temperature was lowered to at 22°C again to monitor the response to the heat shock.

G) Quantification of time spent in sleep 20 min after the heat shock. Wild-type worms sleep on average 38.3%. All other RIS manipulations reduced quiescence. *RIS::unc-58gf(strong)* only sleeps on average 6.5% of the time. **$p<0.001$, ***$p<0.001$, Welch test with FDR correction for multiple testing.

H) *RIS::unc-58gf(strong)* reduces lethargus sleep compared to wild type. ***$p<0.001$, Welch test.

Fig 6 – *RIS::unc-58gf(strong)* fails to inhibit wake-active circuits

A-B) Reduced global neuronal activity (grey shading) correlates with motion quiescence (supplementary movies S1-S6). Head speed was measured. A) sample trace, B) alignment of overall neuronal activity to mobility quiescence bout onset. ***$p<0.001$, Wilcoxon signed rank test.

C) The mean neuronal inactivity fraction correlates with the mean mobility quiescence fraction (linear fit) for all strains.

D) Strong RIS inactivation as well as *RIS::unc-58gf(strong)* caused an overall increase of neuronal activity. n.s. $p > 0.05$, *$p<0.05$, **$p<0.01$, ***$p < 0.001$, Welch test with FDR correction for multiple testing.

E-F) RIM does not activate during mobility quiescence bouts. RIM speed was measured

F) Sample trace (grey shade is RIM inactivity bout). Alignment of RIM activity and speed to mobility quiescence bout onset. *$p<0.05$, **$p<0.01$, ***$p<0.001$, Wilcoxon signed rank test.

G) The mean RIM inactivity fraction correlates with the mean mobility quiescence fraction (linear fit) for all strains.

H) Strong RIS inactivation as well as *RIS::unc-58gf(strong)* caused an overall increase of RIM activity. **p < 0.01, ***p < 0.001, Welch test with FDR correction for multiple testing.

Fig 7 – Phases of sensory PLM neuron inactivity occur during sleep and are strongly reduced in *RIS::unc-58gf(strong)* and *RIS::twk-18gf*

A-C) Sample traces of PLM activity and speed in wild type, *RIS::unc-58gf(strong)* and *RIS::twk-18gf*. PLM speed was measured.

D) *RIS::twk-18gf* and *RIS::unc-58gf(strong)* reduced phases of sensory neuron quiescence. *p<0.05, *** p<0.001, Welch test.

E) *RIS::twk-18gf* and *RIS::unc-58gf(strong)* caused an overall increase of PLM activity. **p<0.01, ***p < 0.001, Welch test.

F) *RIS::twk-18gf* causes increased PLM activity in detected quiescence as well as mobility bouts. *p<0.05, **p<0.01, ***p < 0.001, Welch test for comparison between strains, Wilcoxon signed rank test for comparison between the same strain quiescent and mobility bouts.

Fig 8. *RIS::unc-58gf(strong)* causes increased muscle activity.

A) We calculated the movement speed of the RIM head neuron to measure head movement speed. RIM speed was reduced in *RIS::unc-58gf(strong)*. *p<0.05, **p<0.01, ***p < 0.001, Welch test for comparison between strains, Wilcoxon signed rank test for comparison between the same strain quiescent and mobile bouts.

B) We calculated the movement speed of the PLM tail neuron to measure tail movement speed. PLM movement speed was not significantly changed in mobile bouts in *RIS::unc-58gf(strong)* and increased in detected quiescent bouts. *p<0.05, **p<0.01, ***p < 0.001, Welch test for comparison between strains, Wilcoxon signed rank test for comparison between the same strain quiescent and mobile bouts.

C-E) Sample traces of muscle activity for *RIS::twk-18gf* and *RIS::unc-58gf(strong)*.

F) *RIS::unc-58gf(strong)* causes increased muscle activity in quiescence as well as mobility bouts. *p<0.05, **p<0.01, ***p < 0.001, Welch test for comparison between strains,

Wilcoxon signed rank test for comparison between the same strain quiescent and mobility

bouts.

G) A sleep bout alignment reveals that *RIS::unc-58gf(strong)* has increased muscle

activity. **p<0.01, ***p < 0.001, Wilcoxon signed rank test.

H) *RIS::twk-18gf* and *RIS::unc-58gf(strong)* reduced phases of muscle inactivity.

*p<0.05, Welch test.

1433
1434
Fig 9 *RIS::unc-58gf(strong)* is refractory to optogenetic activation of RIS

A) Optogenetic activation of RIS during the stimulation time (minute 1-2) leads to an

increase of RIS activity and reduced mobility in wild-type worms but it does not affect

the strong RIS activation strain. RIS speed was measured. A statistical comparison was

made between baseline (0-1min) and stimulation (1-2min). n.s. p>0.05, **p<0.01,

***p<0.001, Wilcoxon signed rank test.

B) Control for the optogenetic experiment. Without ATR treatment, neither the wild type

nor the strong RIS activation strain, shows a response to green light. N.s. p>0.05,

Wilcoxon signed rank test.

1444
1445
Fig 10 – Long-term optogenetic activation of RIS increases overall RIS calcium

activity and promotes survival

A) Sample trace of control larvae without retinal in which RIS cannot be activated

optogenetically. RIS speed was measured.

B) Long-term optogenetic RIS activation by orange light in the presence of retinal.

C) Long-term activation of RIS by ReaChR leads to an initial strong increase in calcium

activity in RIS and simultaneous mobility quiescence followed by a mild RIS activity

increase yet no increase in movement speed or movement quiescence.

D) Comparison of RIS activity level changes caused by *RIS::unc-58gf(strong)* and by

optogenetic activation of RIS. The data was normalized to their respective controls.

E) Long-term optogenetic activation of RIS significantly increases RIS baseline. *p<0.05

Welch test.

F) Transients in RIS are significantly reduced in magnitude and don't correlate with quiescence. **$p<0.01$ Welch test.

G) Long-term optogenetic activation of RIS inside microfluidic chambers leads to a small increase in survival ($p=0.09$ Logrank test, $p=0.02$ Fisher's Exact Test, the average of all individual worms of all three technical replicates was averaged, n=92 for *RIS::ReaChR*(+ATR), n=98 *RIS::ReaChR*(-ATR), n=124 wild type (+ATR), n=108 wild type (-ATR))

Fig 11 - RIS activates upon lethal blue-light stimulation and extends survival

A) Sample trace for a control worm stimulated by a short blue light stimulus. RIS speed was measured.

B) Sample traces of wild-type worms upon stimulation with a long blue-light stimulus. The grey shaded area presents detected time in motion quiescence.

C) The strong stimulus increased normalized RIS activity over baseline compared to control worms. ***$p<0.001$, Welch test.

D) Wild-type worms have a survival advantage over RIS-inhibited worms upon exposure to the lethal blue-light stimulus. *$p<0.05$, Welch test.

Fig 12 – FLP-11 is required for sleep during L1 arrest

A) Deletion of the neuropeptide gene *flp-11* causes loss of sleep. Immobility in *RIS::unc-58gf(strong)* is increased when *flp-11* is deleted. ***$p<0.001$, Welch test with FDR correction for multiple testing.

B) Loss of the FLP-11 neuropeptides leads to strongly reduced survival (three-parameter logistic fit, see also supplementary table S1, Fisher's Exact Test was conducted on day 18) for comparisons when *flp-11(-)* was the shortest lived condition, day 20) when *RIS::unc-58gf(strong), flp-11(-)* was the shortest lived condition or day 24) when wild type was the shortest lived condition. The p-values were FDR corrected with Benjamini-Hochberg procedure with a 5% false discovery rate. The plot includes data from three replicates. ***$p<0.001$.

1491

List of Legends

The supplementary file includes:

Supplementary text

CRIPR sequences

Supplementary Figures

Fig S1 to S10

Supplementary Tables

Table S1 fitting parameters

Table S2 strain list

Table S3 primer list

Table S4 p-values

Table S5 DEG genes from the transcriptome data

Movie legends for movies S1-S8

References for the supplementary information

A

RIS manipulation	chromo-some	DNA sequence
RIS::egl-23gf(weak)	X	*flp-11p(endogenous)* | *flp-11b* | SL2 | *egl-23(L229N)* | linker | *mKate2* | *flp-11 3'UTR(endogenous)*
RIS::egl-23gf(strong)	X	*flp-11p(endogenous)* | *flp-11b* | SL2 | *egl-23(A383V)* | linker | *mKate2* | *flp-11 3'UTR(endogenous)*
RIS::twk-18gf	X	*flp-11p(endogenous)* | *flp-11b* | SL2 | *twk-18(e1913)* | linker | *mKate2* | *flp-11 3'UTR(endogenous)*

max z-projection mKate2   single z-plane DIC   z-projection merged   single z-plane mKate2

B *RIS::egl-23gf (weak)*

20µm

C *RIS::egl-23gf (strong)*

20µm

D *RIS::twk-18gf*

20µm

E

RIS manipulation	chromo-some	DNA sequence
RIS::unc-58gf (strong)	X	*flp-11p(endogenous)* | *flp-11b* | SL2 | *unc-58(L428F)* | linker | *mKate2* | *flp-11 3'UTR(endogenous)*
RIS::unc-58gf (weak)	III	*flp-11p(minimal)* | *unc-58(L428F)* | linker | *mKate2* | *flp-11 3'UTR(minimal)*

F

mKate2 intensity (a.u.)

RIS::unc-58gf (weak) n=12   *RIS::unc-58gf (strong)* n=10

***

max z-projection mKate2   z-projection brightfield   z-projection merged   single z-plane mKate2

G *RIS::unc-58gf (strong)*

RIS

20µm

H *RIS::unc-58gf (weak)*

RIS

20µm