The genetic architectures of vine and skin maturity in tetraploid potato

Maria V. Caraza-Harter (ORCID 0000-0002-5934-1526)
Jeffrey B. Endelman* (ORCID 0000-0003-0957-4337)

Department of Horticulture, University of Wisconsin-Madison, Madison, WI 53706, USA

*corresponding author
Email: endelman@wisc.edu

ABSTRACT

As with most crops, maturity is a critical trait for potato production and therefore breeding. In this study, both vine and skin maturity, which refer to foliar senescence and adherence of the tuber periderm, respectively, were measured in breeding populations (N = 586) over a two-year period. It is well established that *S. tuberosum Cyclin DOF Factor 1 (CDF1)* affects vine maturity, but this is the first study to quantify its effect on skin maturity, using linked markers discovered through genome-wide association studies (GWAS). Skin maturity had slightly less genomic heritability than vine maturity (0.41 vs. 0.47), and the importance of *CDF1* was considerably less for skin maturity, accounting for only 39% of the additive variance vs. 70% for vine maturity. Using path analysis, the additive genetic correlation between the two maturity traits was estimated at 0.65, of which 0.51 was attributed to *CDF1*. Within the GWAS panel was a half-diallel population derived from three parents, which was analyzed by joint linkage mapping. Based on the estimated effects of the parental haplotypes and prior knowledge of the allelic series for *CDF1*, the *CDF1* allele for each haplotype was predicted and ultimately confirmed through whole-genome sequencing. The ability to connect statistical alleles from QTL models with biological alleles based on DNA sequencing represents a new milestone in genomics-assisted breeding for tetraploid species.

INTRODUCTION

Maturity is very important for plant breeding and production and can refer to several, related concepts (Bruns 2007). Perhaps the most common meaning is the time interval between planting and harvest, such that varieties with early maturity can be harvested before those with late maturity. The traits that determine when a crop is ready for harvest vary across species. Although potato (*S. tuberosum*) tubers are not true seeds, they still develop toward a condition of maximum physiological dormancy (Struik and Wiersema 1999). Several maturity traits are used to describe the changes that occur during this process (Bussan et al. 2009). Although these traits can refer to a time interval, more commonly they describe a range of phenotypic variation at one time point. Vine maturity refers to senescence of the foliage and has been traditionally measured based on a visual rating; more recently, remote sensing has been used (Sankaran et al. 2015; Colwell et al. 2021). Chemical maturity refers to the concentration of tuber sucrose, which decreases in late development as starch and dry matter levels increase (Sowokinos 1978).

Skin maturity, which is commonly called skin set, refers to adherence of tuber skin to the underlying tissue. Anatomically, skin is the phellem (outer) layer of the potato periderm, which also contains the phellogen (middle) and phelloderm (inner) layers (Reeve et al. 1969). Phellogen is meristematic tissue, adding cells of suberized phellem to the outside and phelloderm to the inside during tuber growth. Skinning (Figure 1) occurs when the phellem is subjected to shear stress and separates from the phelloderm due to fracture of the phellogen (Lulai and Freeman 2001). Skinning reduces the marketability of tubers and their storability, by compromising the barrier to water loss and pathogen entry. Skin maturity can be quantified based on the percent of the tuber surface without skin, after exposure to the stresses of harvest and grading (Caraza-Harter and Endelman 2020). Another approach is to measure the torque at which

skinning occurs, using a handheld torsion meter (Halderson and Henning 1993; Lulai and Orr 1993).

Little is known about the genetics of skin maturity, in part because it is rarely measured in breeding populations. Some progress has been made by studying patterns of gene expression in select varieties (Neubauer et al. 2013; Vulavala et al. 2019), which indicates it is a complex trait. Our understanding of the relationship between skin and vine maturity is primarily based on research geared toward crop management, not breeding. Management actions that hasten foliar senescence, such as chemical desiccation or reduced fertilization, generally lead to better skin maturity at a fixed days after planting (Murphy 1968; Bethke and Busse 2010). As such, the predominant conceptual model is that skin maturation follows vine maturation. In contrast to skin maturity, vine maturity is routinely measured in breeding and has been the subject of many QTL studies (e.g., Collins et al. 1999; Bradshaw et al. 2004; Uitdewilligen et al. 2013).

Most studies of vine maturity identify a major QTL on potato chromosome 5, which Kloosterman et al. (2013) demonstrated was a potato homolog of the *A. thaliana* gene *Cycling DOF Factor 1* (*CDF1*). In potato, CDF1 promotes expression of florigen (SP3D) and tuberigen (SP6A), which are key signals for the initiation of flowering and tuberization, respectively (Navarro et al. 2011; Abelenda et al. 2014). The connection between this pathway and foliar senescence (i.e., vine maturity) is not well understood. In addition to the DOF (DNA-binding One zinc Finger) domain, the ancestral (i.e., wild-type) *S. tuberosum* CDF1 contains C-terminal domains that interact with the proteins GI and FKF1 in a daylength-dependent manner (Kloosterman et al. 2013; Gutaker et al. 2019). Under short days, such as those found in the Andean region where potato was first domesticated, FKF1 expression lags behind GI (Sawa et al.

2007). Under long days, their expression becomes more synchronized, leading to diminished activity of wild-type CDF1.

Three alleles of *CDF1* that code for truncated proteins have been discovered (Kloosterman et al. 2013; Gutaker et al. 2019). These alleles are numbered 2–4 (wild-type alleles are collectively designated type 1), and all contain an insertion in the same location, creating a premature stop codon. *CDF1.3* has an 865 bp insertion from a putative transposon, while *CDF1.2* and *CDF1.4* have different 7 bp insertions. These truncated CDF1 variants are missing the FKF1 binding domain, which leads to enhanced CDF1 activity and early vine maturity under long days. The large size of the insertion in *CDF1.3* also disrupts a long, non-coding RNA named *FLORE*, which is anti-sense to the *CDF1* transcript (Ramírez Gonzales et al. 2021).

The present study was initiated to study the quantitative genetics of skin maturity and its relationship to vine maturity in breeding populations. Since beginning the study, new software has been developed to characterize the function of haplotypes in tetraploid, multiparental populations (Zheng et al. 2021; Amadeu et al. 2021). Sequencing technology has also advanced, reducing the cost and time needed to determine the sequence and dosage of haplotypes. Using these new tools, the original scope of the study was extended to investigate the sequence-function relationship between *CDF1* and maturity.

MATERIALS AND METHODS

Germplasm, genotyping, genomics, and phenotyping

The study population consisted of two consecutive cohorts from the red-skin potato breeding program at the University of Wisconsin-Madison. The 2017 cohort consisted of 127 clones, of which 47 were selected (based on overall merit) for a second year of evaluation with the 2018

cohort of 459 clones, for a total population size of 586. The 2018 cohort was much larger than normal because minimal selection was practiced in the first two field generations, to prepare for the present study. Breeding lines were genotyped with version 3 of the potato SNP array (Felcher et al. 2012; Vos et al. 2015), and genotype calls were made using R package fitPoly (Voorrips et al. 2011; Zych et al. 2019). Each marker data point is the estimated dosage (0–4) of the B allele for the sample. There were 10,185 SNPs after removing markers for which fewer than 5 clones contained the minor allele. Version 4.03 of the DM reference genome was used for physical positions (Potato Genome Sequencing Consortium 2011; Sharma et al. 2013).

An augmented experimental design was used for field trials at the Hancock Agricultural Research Station (Hancock, Wisconsin), with one plot for each breeding line and three (in 2017) or four (in 2018) plots per check variety. Each plot was planted with 15 seed pieces, with 30 cm in-row spacing and 90 cm between rows. The experiments were planted on April 24, 2017, and May 1, 2018, and harvested 120 days after planting (DAP). Diquat bromide was applied at 14 and 7 days before harvest to promote vine desiccation. Tubers were mechanically harvested into 30×45 cm rigid plastic milk crates, run through our standard washing and grading line, and then crated for storage at 12°C with 95% relative humidity.

Vine maturity was measured at 100 days after planting (DAP) using a 1–9 visual rating: 1 indicates full senescence and 9 indicates flowering. Skin maturity was measured from images collected within a few days of harvest, following the protocol of Caraza-Harter and Endelman (2020). A representative sample of 7-8 tubers from each plot was photographed on both sides against a black background, using a Photosimile 200 Lightbox and CanonEOS T5i camera. All photos included a MacBeth color card to adjust for lighting and exposure variation. Using the ImageJ software (Schneider et al. 2012), hue and brightness thresholds were used to delineate the

skinned surface area for measurement, and each photo was visually inspected to ensure accuracy. Skin maturity is reported as the ratio between skinned area and total surface area, with a log-transformation, $y = \log (x + 1)$, to improve normality of the residuals.

Whole-genome sequencing for two parents from the diallel population (W6511-1R, Villetta Rose) utilized the NovaSeq S1 2x150 flow cell (University of Minnesota Genomics Center). Reads were aligned with BWA-MEM (Li 2013) to the *CDF1.1_scaffold1389* and *CDF1.3_scaffold1390* alleles from the Atlantic reference genome (Hoopes et al. 2022). The software SAMtools (Li et al. 2009) was used to fix paired reads and remove PCR duplicates. The software BEDtools (Quinlan and Hall 2010) was used to remove reads with fewer than 10 bp on both sides of the transposon insertion site (Gutaker et al. 2019). The software IGV (Robinson et al. 2011) was used to visualize the alignments and count *CDF1* allele read depth. A multiple sequence alignment of *CDF1* alleles from the potato pan-genome (Hoopes et al. 2022) was performed using MUSCLE v3.8 (Edgar 2004; Madeira et al. 2019), and the percent identity matrix was used for UPGMA hierarchical clustering with *hclust* in R (R Core Team 2021).

Statistical analysis

Data were analyzed in R (R Core Team 2021) with a two-stage approach (Damesa et al. 2017), using ASReml-R v4 (Butler et al. 2018) for REML estimation of variance components. In Stage 1, the best linear unbiased estimate (BLUE) for genotype was computed separately in each year, with a random cofactor for block. The same model, but with genotype as random, was used to estimate the broad-sense heritability on a plot basis as $V_g/(V_g+V_e)$ from the Stage 1 variance components for genetic (V_g) and residual (V_e) variance.

The BLUEs from Stage 1 were used as the response variable for genome-wide association studies with the software GWASpoly (Rosyara et al. 2016). Population structure was controlled using a kinship matrix (Yu et al. 2006), calculated with the leave-one-chromosome-out (LOCO) method (Yang et al. 2014). The discovery threshold was based on a significance level of 0.1, corrected for multiple testing by the effective number of markers (Moskvina and Schmidt 2008). Partial R^2 values were computed based on the change in the likelihood under backward elimination (Sun et al. 2010).

Partitioning of variance in Stage 2 utilized the BLUEs from Stage 1 as the response variable. The Stage 2 model was

$$\text{BLUE}[g_{ij}] = y_{ij} = \mu_j + \sum_k \beta_k w_{ik} + a_i + d_i + s_{ij} + \varepsilon_{ij} \qquad [1]$$

where g_{ij} is the genotypic value for clone i in year j, μ_j is the fixed effect for year j, w_{ik} is the (centered) allele dosage for clone i at marker k, and β_k is the fixed effect for marker k. The vector of additive values, **a**, follows a multivariate normal distribution with covariance proportional to a genomic relationship matrix (**G**) estimated from markers (VanRaden 2008; Endelman et al. 2018). Independent and identically distributed (i.i.d.) residual genetic values, d_i, were included to capture non-additive effects. The s_{ij} random effect follows a multivariate normal distribution with no free variance parameters: the variance-covariance matrix **Ω** is the direct sum of the variance-covariance matrices of the Stage 1 BLUEs (Damesa et al. 2017). The model residuals ε_{ij} are i.i.d. and represent the genotype x year interaction. The variance for each term in Eq. 1 was computed based on Legarra (2016) to estimate the proportion of variance explained.

Joint analysis of vine and skin maturity also followed a two-stage approach. In Stage 1, an additional parameter for the residual covariance was estimated. The Stage 2 model followed

Eq. 1, using separable covariance structures for the additive, residual genetic, and genotype x year effects:

$$\text{var}[\mathbf{a}] = \mathbf{G} \otimes \boldsymbol{\Gamma}_a$$

$$\text{var}[\mathbf{d}] = \mathbf{I}_n \otimes \boldsymbol{\Gamma}_d \qquad\qquad [2]$$

$$\text{var}[\boldsymbol{\varepsilon}] = \mathbf{I}_t \otimes \boldsymbol{\Gamma}_\varepsilon$$

Effect correlations were calculated from the 2x2 covariance matrices $\boldsymbol{\Gamma}$ in the usual way $\left(\Gamma_{12}/\sqrt{\Gamma_{11}\Gamma_{22}}\right)$, and the contribution of each effect to the phenotypic correlation of the Stage 1 BLUEs was calculated using standard methods in path analysis (Wright 1934).

Genetic analysis of the three-parent diallel population relied on the software packages PolyOrigin for parental phasing and haplotype construction (Zheng et al. 2021) and diaQTL (Amadeu et al. 2021) for QTL mapping. The discovery threshold was based on a genome-wide significance level of 0.05, established using simulation (Amadeu et al. 2021). The number of MCMC iterations was determined using the function *set_params* at the recommended settings. PotVar0079081 was used as an additive covariate to represent *CDF1* in the genome scan for skin maturity.

RESULTS

Over a two-year period, 586 clones from the US red-skin market type were phenotyped for vine and skin maturity. We observed the full range of variation for vine maturity at 100 days after planting (DAP), from a score of 1 for complete senescence to 9 for flowering (Fig. 2). Skin maturity at harvest 120 DAP ranged from 0 to 25% skinned area. Within year, the broad-sense heritability (H^2) for skin maturity was 0.7–0.8 on a plot basis; vine maturity was similar in 2018

but lower in 2017 (Table 1). As expected, clones with early vine maturity tended to have early skin maturity (Fig. 2), with a phenotypic correlation of 0.41 in 2017 and 0.45 in 2018.

GWAS identified a QTL on chromosome 5 in the vicinity of *CDF1*, which explained 33% of the variation for vine maturity but only 14% for skin maturity (Table 2, Fig. S1). Figure 3 illustrates the additive relationship between vine maturity and allele dosage at the most significant SNP, PotVar0079081 (49 kb from *CDF1*), with an average maturity decrease of 1.8 per dose of the "B" allele from the SNP array. Minor QTL on chromosomes 6 and 9 for vine and skin maturity, respectively, explained 3-4% of the variation.

Bivariate trait analysis using a genomic G matrix was used to partition the variance for each trait and their correlation (Table 3). For vine maturity, *CDF1* explained 33% of the variance compared to 14% for the polygenic additive effect, for a total genomic heritability of 0.47. For skin maturity, the additive polygenic effect explained more of the variance (25%) than *CDF1* (16%), and non-additive effects explained 9%. Genotype x year effects were important for both traits (24–27% of the variance). *CDF1* explained nearly half of the total phenotypic correlation between the traits (0.22 out of 0.48), and the second most important contributor was the genotype x year effect.

Within the GWAS panel was a half-diallel of 397 clones derived from three parents (W6511-1R, W9914-1R, and Villetta Rose). As expected from the GWAS results, joint linkage mapping identified a major QTL for vine maturity near *CDF1* (Fig. S2). Based on the estimated additive effects (Fig. 4), a nearly binary classification of haplotypes into late (positive effect) vs. early (negative effect) was possible. The only exception was haplotype 3 (numbering is arbitrary) in parent W6511-1R, which had an intermediate effect on vine maturity. The phased genotype

for PotVar0079081 at the bottom of Fig. 4 shows the "B" allele had perfect linkage with early maturity, excluding the W6511-1R.3 haplotype.

Based on previous research summarized in the Introduction, we hypothesized that parental haplotypes with late maturity were wild-type *CDF1.1* alleles, the early haplotypes were *CDF1.3*, and the intermediate haplotype W6511-1R.3 was either *CDF1.2* or *CDF1.4*. Allele counts from the alignment of Illumina short reads against *CDF1* confirmed this hypothesis (Table 4). The tetraploid *CDF1* genotype for Villetta Rose is 1133 and W6511-1R is 1123 (bottom of Fig. 4). Although W9914-1R no longer existed to submit for sequencing, the most likely *CDF1* genotype is 1333.

The minor QTL for skin maturity on chromosome 9 (chr09) discovered by GWAS was also discovered via joint linkage mapping in the diallel population. The most likely position of the QTL from the joint linkage mapping was 8.9 Mb, compared with 4.1 Mb from GWAS (Fig. 5A). From the estimated haplotype effects (Fig. 5B), we can infer that one haplotype in W6511-1R carries a unique allele for early maturity. The low frequency (1/12) of this allele in the diallel population constrains the percent variation explained (PVE), which was 9% compared to 3% in the GWAS panel (Table 2). Although the difference in PVE could be due to differences in population composition (the GWAS panel contained additional genotypes), the lack of a SNP in our dataset unique to the early haplotype limits the marker-QTL LD and therefore PVE in the GWAS panel.

DISCUSSION

This is the second publication in which QTL haplotype effects for vine maturity were connected to *CDF1* alleles in a tetraploid mapping population. Hoopes et al. (2022) analyzed a single F1 population with *CDF1* parental genotypes 1111 x 1223, and we have extended the

method to a multi-parental population. Our determination of the *CDF1* allelic series 3 > 2 > 1 for early maturity agrees with the results of Hoopes et al. (2022). An additional discovery in our study was the dosage effect of truncated *CDF1* alleles on vine maturity, which could not be observed by Hoopes et al. (2022) based on the parental genotypes. Although not explicitly mentioned by Kloosterman et al. (2013), their supplemental results for vine maturity in a diploid F1 population with *CDF1* parental genotypes 12 x 13 also appear to demonstrate an additive effect for truncated alleles.

The most significant SNP for vine maturity in our GWAS, PotVar0079081, has been reported in other recent publications (Willemsen 2018; Klaassen et al. 2019; Ospina Nieto et al. 2021) that used a GWAS panel with mostly European varieties but also some founders of the US red market type, such as Triumph, Katahdin, and Early Rose. From the pedigree analysis of Vos et al. (2015), we know the SNP predates modern breeding ("pre-1945"). With a physical distance of 49 kb and map distance of 1.5 cM from *CDF1*, PotVar0079081 is not particularly close to the causal locus; there were over 20 SNPs in our dataset closer to *CDF1*. This illustrates a well-known limitation of GWAS panels: the fine-map position of an unobserved causal variant cannot be reliably inferred from marker significance.

From the observation that PotVar0079081 had perfect linkage disequilibrium with the *CDF1.2* and *CDF1.3* alleles (as a group) in the parents of our diallel population, a potential explanation for its high significance in association studies can be offered. Genetic distance analysis suggests many different wild-type alleles existed in the potato gene pool when *CDF1.3* was created by a transposon insertion (Fig. 6). Soon thereafter (on the evolutionary time scale for *CDF1*), the transposon was excised, leaving a 7 bp footprint in *CDF1.2*. If PotVar0079081 arose after diversification of the wild-type alleles, in the lineage where *CDF1.3* arose but before the

transposon insertion, then it would be a good proxy for the dosage of late-maturing, wild-type alleles relative to the dosage of truncated alleles in panels with a low frequency of *CDF1.4* (which arose in a separate lineage, see Fig. 6).

Despite its importance, skin maturity has not been routinely measured in potato breeding as a quantitative trait. One justification is the belief that selection for early vine maturity (which is measured) will also lead to earlier skin maturity. Our study confirms the validity of this approach but also reveals its limitations. Using the results from Table 2, path analysis for the additive component of each trait (Fig. S3) leads to an estimated additive genetic correlation of 0.65. The efficiency of indirect selection for skin maturity based on vine maturity, relative to direct selection, is therefore $0.65(h_{\text{vine}}/h_{\text{skin}}) \approx 0.7$. However, indirect selection is expected to become less efficient in the future, as genetic selection against wild-type *CDF1.1* alleles reduces their frequency and the contribution of *CDF1* to genetic variation for maturity. The results in Table 2 imply the efficiency of indirect selection for skin maturity when *CDF1* becomes fixed may be closer to 0.3. Fortunately, as demonstrated in this study, direct selection is already feasible using image-based phenotyping methods.

REFERENCES

Amadeu RR, Muñoz PM, Zheng C, Endelman JB (2021) QTL mapping in outbred tetraploid (and diploid) diallel populations. Genetics 219(3). https://doi.org/10.1093/genetics/iyab124

Abelenda JA, Navarro C, Prat S (2011) From the model to the crop: genes controlling tuber formation in potato. Current Opinion in Biotechnology 22: 287–292.

Bethke PC, Busse JS (2010) Vine-kill treatment and harvest date have persistent effects on tuber physiology. American Journal of Potato Research 87:299–309. https://doi.org/10.1007/s12230-010-9137-4

Bradshaw JE, Pande B, Bryan GJ, Hackett CA, McLean K, Stewart HE, Waugh R (2004) Interval mapping of quantitative trait loci for resistance to late blight [*Phytophthora infestans* (Mont.) de Bary], height and maturity in a tetraploid population of potato (*Solanum tuberosum* subsp. *tuberosum*). Genetics 168:983–995. https://doi.org/10.1534/genetics.104.030056.

Bruns HA (2007) A survey of factors involved in crop maturity. Agronomy Journal 101: 60-66.

Bussan AJ, Sabba RP, Drilias MJ (2009) Tuber maturation and potato storability: Optimizing skin set, sugars and solids. University of Wisconsin-Extension, Cooperative Extension.

Butler D, Cullis B, Gilmour A, Gogel B, Thompson R (2018) ASReml-R Reference Manual Version 4. VSN International Ltd, Hemel Hempstead, HP1 1ES, UK.

Caraza-Harter MV, Endelman JB (2020) Image-based phenotyping and genetic analysis of potato skin set and color. Crop Science 60:202–210. https://doi.org/10.1002/csc2.20093

Collins A, Milbourne D, Ramsay L, Meyer R, Chatot-Balandras C, Oberhagemann P, De Jong W, Gebhardt C, Bonnel E, Waugh R (1999) QTL for field resistance to late blight in potato are strongly correlated with maturity and vigour. Molecular Breeding 5:387-398.

Colwell FJ, Souter J, Bryan GJ, Compton LJ, Boonham N, Prashar A (2021) Development and validation of methodology for estimating potato canopy structure for field crop phenotyping and improved breeding. Frontiers Plant Science 12:612843. https://doi.org/10.3389/fpls.2021.612843.

Damesa TM, Möhring J, Worku M, Piepho H-P (2017) One step at a time: Stage-Wise analysis of a series of experiments. Agronomy Journal 109:845-857. https://doi.org/10.2134/agronj2016.07.0395

Edgar RC (2004) MUSCLE: a multiple sequence alignment method with reduced time and space complexity. BMC Bioinformatics 5:113. https://doi.org/10.1186/1471-2105-5-113

Endelman JB, Schmitz Carley CA, Bethke PC et al. (2018) Genetic variance partitioning and genome-wide prediction with allele dosage information in autotetraploid potato. Genetics 209:77–87. https://doi.org/10.1534/genetics.118.300685

Felcher KJ, Coombs JJ, Massa AN, Hansey CN, Hamilton JP, Veilleux RE, Buell CB, Douches DS (2012) Integration of two diploid potato linkage maps with the potato genome sequence. PLoS ONE 7(4):e36347. https://doi.org/10.1371/journal.pone.0036347

Gutaker RM, Weiss CL, Ellis D, Anglin NL, Knapp S, Fernández-Alonso JL, Prat S, Burbano HA (2019) The origins and adaptation of European potatoes reconstructed from historical genomes. Nature Ecol Evol 3:1093-1101. https://doi.org/10.1038/s41559-019-0921-3

Halderson JL, Henning RC (1993) Measurements for determining potato tuber maturity. American Potato Journal 70:131–141.

Hoopes G, Meng X, Hamilton JP et al. (2022) Phased, chromosome-scale genome assemblies of tetraploid potato reveal a complex genome, transcriptome, and proteome landscape underpinning phenotypic diversity. Molecular Plant 15:520–536. https://doi.org/10.1016/j.molp.2022.01.003

Klaassen MT, Willemsen JH, Vos PG, Visser RG, Van Eck HJ, Maliepaard C, Trindade LM (2019) Genome-wide association analysis in tetraploid potato reveals four QTLs for protein content. Molecular Breeding 39. https://doi.org/10.1007/s11032-019-1070-8

Kloosterman B, Abelenda JA, Carretero Gomez MM, Oortwijn M, de Boer JM, Kowitwanich K, Horvath BM, van Eck HJ, Smaczniak C, Prat S, Visser RGF, Bachem CWB (2013) Naturally occurring allele diversity allows potato cultivation in northern latitudes. Nature 495:246-250. https://doi.org/10.1038/nature11912

Legarra A (2016) Comparing estimates of genetic variance across different relationship models. Theor. Pop. Biol. 107:26–30. https://doi.org/10.1016/j.tpb.2015.08.005

Li H (2013) Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM. arXiv:1303.3997v2 [q-bio.GN]

Li H, Handsaker B, Wysoker A et al. (2009) The sequence alignment/map format and SAMtools. Bioinformatics 25:2078–2079. https://doi.org/10.1093/bioinformatics/btp352

Lulai EC, Orr PH (1993) Determining the feasibility of measuring genotypic differences in skin-set. American Journal of Potato Research 70: 599-609.

Lulai EC, Freeman TP (2001) The importance of phellogen cells and their structural characteristics in susceptibility and resistance to excoriation in immature and mature potato tuber (*Solanum tuberosum* L.) periderm. Annals of Botany 88:555–561. https://doi.org/10.1006/anbo.2001.1497

Madeira F, Park YM, Lee J, Buso N, Gur T, Madhusoodanan N, Basutkar P, Tivey ARN, Potter SC, Finn RD, Lopez R (2019) The EMBL-EBI search and sequence analysis tools APIs in 2019. Nucleic Acids Research 47(W1):W636–W641. https://doi.org/10.1093/nar/gkz268

Moskvina V, Schmidt KM (2008) On multiple-testing correction in genome-wide association studies. Genetic Epidemiology 32:567–573. https://doi.org/10.1002/gepi.20331

Murphy HJ (1968) Potato vine killing. American Potato Journal 45: 472-478.

Navarro C, Abelenda JA, Cruz-Oró E, Cuéllar CA, Tamaki S, Silva J, Shimamoto K, Prat S (2011) Control of flowering and storage organ formation in potato by FLOWERING LOCUS T. Nature 478:119–122. https://doi.org/10.138/nature10431

Neubauer JD, Lulai EC, Thompson AL, Suttle JC, Bolton MD, Campbell LG (2013) Molecular and cytological aspects of native periderm maturation in potato tubers. Journal of Plant Physiology 170:413–423. https://doi.org/10.1016/j.jplph.2012.10.008

Ospina Nieto CA, Lammerts van Bueren ET, Allefs S, Vos PG, van der Linden G, Maliepaard CA, Struik PC (2021) Association mapping of physiological and morphological traits related to crop development under contrasting nitrogen inputs in a diverse set of potato cultivars. Plants 10:1727. https://doi.org/10.3390/plants10081727

Potato Genome Sequencing Consortium (2011) Genome sequence and analysis of the tuber crop potato. Nature 475:189–195. https://doi.org/10.1038/nature10158

Quinlan AR, Hall IM (2010) BEDTools: a flexible suite of utilities for comparing genomic features. Bioinformatics 26:841–842. https://doi.org/10.1093/bioinformatics/btq033

R Core Team (2021) R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria.

Ramírez Gonzales L, Shi L, Bergonzi SB, Oortwijn M, Franco-Zorrilla JM, Solano-Tavira R, Visser RGF, Abelenda JA, Bachem CWB (2021) Potato CYCLING DOF FACTOR 1 and its lncRNA counterpart *StFLORE* link tuber development and drought response. The Plant Journal 105:855-869. https://doi.org/10.1111/tpj.15093

Reeve RM, Hautala E, Weaver ML (1969) Anatomy and compositional variation within potatoes I. Developmental histology of the tuber. American Journal of Potato Research 46:361–373.

Robinson JT, Thorvaldsdóttir H, Winckler W et al. (2011) Integrative Genomics Viewer. Nature Biotech. 29:24–26.

Rosyara UR, De Jong WS, Douches DS, Endelman JB (2016) Software for genome-wide association studies in autopolyploids and its application to potato. Plant Genome 9:2. https://doi.org/10.3835/plantgenome2015.08.0073

Sankaran S, Khot LR, Espinoza CZ, Jarolmasjed S, Sathuvalli VR, Vandemark GJ, Miklas PN, Carter AH, Pumphrey MO, Knowles NR, Pavek MJ (2015) Low-altitude, high resolution aerial imaging systems for row and field crop phenotyping: A review. European Journal Agronomy 70:112-123. https://doi.org/10.1016/j.eja.2015.07.004.

Sawa M, Nusinow DA, Kay SA, Imaizumi T (2007) FKF1 and GIGANTEA complex formation is required for day-length measurement in *Arabidopsis*. Science 318:261-265.

Schneider C, Rasband W, Eliceiri K (2012) NIH Image to ImageJ: 25 years of image analysis. Nature Methods 9.

Searle SR, Casella G, McCulloch CE (1992) Variance components. John Wiley & Sons, Hoboken, NJ.

Sharma SK, Bolser D, de Boer J et al (2013) Construction of reference chromosome-scale pseudomolecules for potato: integrating the potato genome with genetic and physical maps. G3 3:2031-2047. https://doi.org/10.1534/g3.113.007153

Sowokinos JR (1978) Relationship of harvest sucrose content to processing maturity and storage life of potatoes. American Potato Journal 55:333–344.

Struik PC and Wiersema SG (1999) Seed potato technology. Wageningen Academic Publishers, Wageningen.

Sun G, Zhu C, Kramer MH, Yang S-S, Piepho H-P, Yu J (2010) Variation explained in mixed-model association mapping. Heredity 105:333–340.

Uitdewilligen JGAML, Wolters AMA, D'hoop BB, Borm TJA, Visser RGF, van Eck HJ (2013) A next-generation sequencing method for genotyping-by-sequencing of highly heterozygous autotetraploid potato. PLoS ONE 8:e62355.

VanRaden PM (2008) Efficient methods to compute genomic predictions. J. Dairy Sci. 91:4414–4423.

Voorrips RE, Gort G, Vosman B (2011) Genotype calling in tetraploid species from bi-allelic marker data using mixture models. BMC Bioinformatics 12:172. https://doi.org/10.1186/1471-2105-12-172

Vos PG, Uitdewilligen JGAML, Voorrips RE, Visser RGF, van Eck HJ (2015) Development and analysis of a 20K SNP array for potato (*Solanum tuberosum*): an insight into the breeding history. Theor Appl Genet 128:2387-2401. https://doi.org/10.1007/s00122-015-2593-y

Vulavala VKR, Fogelman E, Faigenboim A, Shoseyov O, Ginzberg I (2019) The transcriptome of potato tuber phellogen reveals cellular functions of cork cambium and genes involved in periderm formation and maturation. Scientific Reports 9:10216. https://doi.org/10.1038/s41598-019-46681-z

Willemsen JH (2018) The identification of allelic variation in potato. Wageningen University, Wageningen, the Netherlands.

Wright S (1934) The method of path coefficients. Ann. Math. Statist. 5:161-215. https://doi.org/10.1214/aoms/1177732676

Yang J, Zaitlen NA, Goddard ME, Visscher PM, Price AL (2014) Advantages and pitfalls in the application of mixed-model association methods. Nature Genetics 467:100–106. https://doi.org/10.1038/ng.2876

Yu J, Pressoir G, Briggs WH, Vroh Bi I, Yamasaki M, Doebley JF, McMullen MD, Gaut BS, Nielsen D, Holland JB, Kresovich S, Buckler ES (2006) A unified mixed-model method for association mapping that accounts for multiple levels of relatedness. Nature Genetics 38:203–208. https://doi.org/10.1038/ng1702

Zheng C, Amadeu RR, Muñoz PM, Endelman JB (2021) Haplotype reconstruction in connected tetraploid F1 populations. Genetics 219(2). https://doi.org/10.1093/genetics/iyab106

Zych K, Gort G, Maliepaard CA, Jansen RC, Voorrips RE (2019) FitTetra 2.0: improved genotype calling for tetraploids with multiple population and parental data support. BMC Bioinformatics 20:148. https://doi.org/10.1186/s12859-019-2703-y

Declarations

Acknowledgments

We thank Peyton Sorensen for assistance with data collection, as well as the staff of the UW Rhinelander and Hancock Agricultural Research Stations.

Funding

Support for data collection was provided by a Wisconsin Dept. of Agriculture Specialty Crop Block Grant (16-02), the Wisconsin Potato and Vegetable Growers Association, and the UW-Madison Office of the Vice Chancellor for Research and Graduate Education. Support for data analysis was provided by the USDA National Institute of Food and Agriculture, Award 2019-67013-29166.

Competing Interests

The authors have no relevant financial or non-financial interests to disclose.

Author Contributions

Study design: JBE. Data collection: MCH. Data analysis: MCH, JBE. Manuscript preparation: MCH, JBE.

Data Availability

The FASTQ files, SNP array marker genotypes, and phenotype data are available from the Dryad depository (https://doi.org/10.5061/dryad.3n5tb2rk0). The genotype input file for joint linkage mapping is distributed with the diaQTL package (https://github.com/jendelman/diaQTL).

FIGURE CAPTIONS

Figure 1. Skinning injury on a red-skin potato tuber. The clone pictured is W10209-7R, one of the checks in the 2017 field trial. Based on image analysis, 18% of the tuber surface was skinned.

Figure 2. Scatterplot of skin maturity vs. vine maturity from a two-year field trial. The points are best linear unbiased estimates (BLUEs) for clone, with a linear trendline in blue. The repeated checks are in red, and the other clones are black. Vine maturity used a 1 (early) to 9 (late) visual rating. Skin maturity is reported as $\log(x+1)$, where x is the skinned area percentage.

Figure 3. Illustrating the additive relationship between vine maturity and allele dosage at SNP PotVar0079081, which is linked to *CDF1*.

Figure 4. Additive haplotype effects for vine maturity (higher = later) at *CDF1* on chromosome 5. Haplotypes are arbitrarily numbered for the parents of the diallel population (Villetta Rose, W6511-1R, W9914-1R). Error bars represent the 90% CI.

Figure 5. (A) Comparison of marker significance scores for skin maturity under GWAS (-$\log_{10}$p, black points) vs. joint linkage mapping (–ΔDIC, red line). (B) Haplotype effects for skin maturity (higher = later) at the QTL on chromosome 9. Haplotypes are arbitrarily numbered for the three parents of the diallel population (Villetta Rose, W6511-1R, W9914-1R). Error bars represent the 90% CI.

Figure 6. *CDF1* dendrogram based on percent identity from a multiple sequence alignment of 12 potato alleles and one tomato allele. The potato alleles are labeled based on the variety and scaffold identifier from Hoopes et al. (2022). Truncated alleles are labeled *CDF1.2*, *CDF1.3*, *CDF1.4*, and the other nine potato alleles are wild-type *CDF1.1*.

TABLES

Table 1. Broad-sense heritability, on a plot basis, within year.

Year	Vine Maturity	Skin Maturity
2017	0.54	0.77
2018	0.78	0.71

Table 2. Genome-wide association study results.

Trait	Marker	Chrom	Position (bp)	% Variance Explained
Vine Maturity	PotVar0079081	5	4,489,481	33
Vine Maturity	PotVar0090539	6	48,596,917	4
Skin Maturity	PotVar0079652	5	4,548,540	14
Skin Maturity	solcap_snp_c2_4158	9	4,091,032	3

Table 3. Partitioning of variance and correlation.

	Vine Maturity[b]	Skin Maturity[b]	Effect Correlation	Phenotypic Correlation[c]
CDF1[a]	0.33	0.16	0.97	0.22
Additive Polygenic	0.14	0.25	0.33	0.06
Genetic Residual	0.00	0.09	NA[d]	0.00
Genotype x Year	0.27	0.24	0.63	0.16
Stage 1 Error	0.25	0.27	0.15	0.04
Total	1.00	1.00		0.48

[a] *CDF1* modeled using fixed effects for PotVar0079081 and PotVar0079652 (see Table 2).

[b] Proportion of variance, excluding the main effect for year.

[c] Contribution toward the phenotypic correlation of the Stage 1 BLUEs.

[d] Not estimable because there was no variance for vine maturity

Table 4. *CDF1* allele read counts from whole-genome sequencing of two parents from the diallel population.

CDF1 Allele	Villetta Rose	W6511-1R
1	36	40
2	0	26
3	45	15

Figure 1. Skinning injury on a red-skin potato tuber. The clone pictured is W10209-7R, one of the checks in the 2017 field trial. Based on image analysis, 18% of the tuber surface was skinned.

Figure 2. Scatterplot of skin maturity vs. vine maturity from a two-year field trial. The points are best linear unbiased estimates (BLUEs) for clone, with a linear trendline in blue. The repeated checks are in red, and the other clones are black. Vine maturity used a 1 (early) to 9 (late) visual rating. Skin maturity is reported as $\log(x+1)$, where x is the skinned area percentage.

Figure 3. Illustrating the additive relationship between vine maturity and allele dosage at SNP PotVar0079081, which is linked to *CDF1*.

Figure 4. Additive haplotype effects for vine maturity (higher = later) at *CDF1* on chromosome 5. Haplotypes are arbitrarily numbered for the parents of the diallel population (Villetta Rose, W6511-1R, W9914-1R). Error bars represent the 90% CI.

Figure 5. (A) Comparison of marker significance scores for skin maturity under GWAS (-log$_{10}$p, black points) vs. joint linkage mapping ($-\Delta$DIC, red line). (B) Haplotype effects for skin maturity (higher = later) at the QTL on chromosome 9. Haplotypes are arbitrarily numbered for the three parents of the diallel population (Villetta Rose, W6511-1R, W9914-1R). Error bars represent the 90% CI.

Figure 6. *CDF1* dendrogram based on percent identity from a multiple sequence alignment of 12 potato alleles and one tomato allele. The potato alleles are labeled based on the variety and scaffold identifier from Hoopes et al. (2022). Truncated alleles are labeled *CDF1.2*, *CDF1.3*, *CDF1.4*, and the other nine potato alleles are wild-type *CDF1.1*.

Supplemental Figures

The genetic architectures of vine and skin maturity in tetraploid potato

Maria V. Caraza-Harter, Jeffrey B. Endelman
Department of Horticulture, University of Wisconsin-Madison, WI 53706, USA

Fig. S1. GWAS results for skin maturity (left) and vine maturity (right).

Fig S2. Joint linkage mapping results on potato chromosome 5 for vine maturity, in the half-diallel with three parents. The y-axis is the change in the deviance information criterion (DIC). The dashed vertical line indicates the position of *CDF1* at 4.54 Mb.

Fig S3. Path analysis for the additive genetic components of vine and skin maturity. The correlation and path coefficients are derived from Table 2.