1

Title
Dark accumulation of downstream glycolytic intermediates confers robust initiation of photosynthesis in cyanobacteria

Authors
Kenya Tanaka[1,2,3], Tomokazu Shirai[2,4], Christopher J. Vavricka[2], Mami Matsuda[2], Akihiko Kondo[1,2,4,5], Tomohisa Hasunuma[1,2,4,*]

Affiliations
1. Engineering Biology Research Center, Kobe University, 1-1 Rokkodai, Nada, Kobe 657-8501, Japan
2. Graduate School of Science, Innovation and Technology, Kobe University, 1-1 Rokkodai, Nada, Kobe 657-8501, Japan
3. Research Center for Solar Energy Chemistry, Graduate School of Engineering Science, Osaka University, Toyonaka, Osaka 560-8531, Japan
4. RIKEN Center for Sustainable Resource Science, 1-7-22 Suehiro, Tsurumi, Yokohama, Kanagawa 230-0045, Japan
5. Department of Chemical Science and Engineering, Graduate School of Engineering, Kobe University, 1-1 Rokkodai, Nada, Kobe 657-8501, Japan

Abstract
Photosynthesis must maintain stability and robustness throughout fluctuating natural environments. In cyanobacteria, dark-to-light transition leads to drastic metabolic changes from dark respiratory metabolism to CO_2 fixation through the Calvin-Benson-Bassham (CBB) cycle using energy and redox equivalents provided by photosynthetic electron transfer. Previous studies showed that catabolic metabolism supports the smooth transition into CBB cycle metabolism. However, metabolic mechanisms for robust initiation of photosynthesis are poorly understood due to lack of dynamic metabolic characterizations of dark-to-light transitions. Here, we show rapid (on a time scale of seconds) dynamic changes in absolute metabolite concentrations and ^{13}C tracer incorporation after strong or weak light irradiation in the cyanobacterium *Synechocystis* sp. PCC 6803. Integration of this data enables estimation of time-resolved nonstationary metabolic flux underlying CBB cycle activation. This dynamic metabolic analysis indicates that downstream glycolytic intermediates including phosphoglycerate and phosphoenolpyruvate accumulate under dark conditions as major substrates for initial CO_2 fixation. Compared with wild-type *Synechocystis*, significant delays in the initiation of oxygen evolution are observed in 12 h dark preincubated mutants deficient in glycogen degradation or oxidative pentose phosphate pathway (*Δzwf*, *Δgnd*, and *ΔglgP*). Accordingly, the degree of delay in the oxygen evolution initiation is proportional to the accumulated pool size of the glycolytic intermediates. These observations indicate that the accumulation of glycolytic intermediates is essential for efficient metabolism switching under fluctuating light environments.

Main text

Photosynthetic organisms have evolved various mechanisms to maintain robust photosynthesis processes throughout natural environmental perturbations. Ancestral oxygenic and phototrophic cyanobacteria contribute to global carbon cycles by photosynthetic CO_2 fixation through the Calvin-Benson-Bassham (CBB) cycle[1–3]. The driving forces of the CBB cycle are ATP and NADPH produced by light energy conversion in photosynthetic electron transfer (PET) reactions; and cyanobacteria have evolved metabolic control mechanisms to coordinate CBB cycle activity with environmental light-dark conditions. One mechanism is enzymatic activity regulation that responds to light-dependent cellular environmental changes. For example, glyceraldehyde 3-phosphate dehydrogenase (GAPDH) and phosphoribulokinase (PRK) are regulated by ternary complex formation with the regulatory protein CP12, depending on redox states of specific cysteine residues and cellular NAD/NADP ratio [4–6].

Maintaining optimal metabolite concentrations is another important mechanism of CBB cycle regulation. In general, concentration robustness is necessary for maintenance of cellular functionality[7]. Recent analyses performed using a theoretical photosynthetic model have predicted that maintaining the appropriate concentrations of CBB cycle intermediates is crucial for the initiation of CO_2 fixation[8]. Experimental evidence in cyanobacteria has confirmed that the transfer of the carbon supply from a storage polymer (glycogen) to the CBB cycle affects photosynthetic efficiency[9,10]. In addition, while enzymatic regulation systems control metabolism, metabolite concentration affects metabolism itself by thermodynamic constraints and effects on enzymatic kinetics as represented by the Michaelis-Menten equation[11,12]. Thus proper regulation of metabolite distribution is considered to be important for activation of the CBB cycle.

However, the metabolic distribution underlying robust CBB cycle operation under light-dark perturbations is still unclear. Ascertaining which metabolite is important for the CBB cycle kickstart requires understanding metabolic dynamics during CBB cycle activation immediately after a dark-to-light transition. For this, we aim to derive dynamic metabolic flux changes during photosynthetic activation. Although metabolic flux analysis (MFA) using an isotope tracer enables the mapping of intracellular carbon fluxes[13–15], the approach cannot be applied to a metabolically non-stationary state with variable metabolic fluxes and pool sizes. On the other hand, previously developed dynamic MFA, which incorporates specialized statistical approaches, has permitted the estimation of long-term flux changes over the culture interval[16,17]. There has been (to our knowledge) no appropriate approach for the quantitative determination of metabolic flux changes throughout short, metabolically transient states, including that of CBB cycle activation.

In this study, we successfully quantify absolute metabolite concentration changes at second-scale intervals after dark-to-light transitions in the cyanobacterium *Synechocystis* sp. PCC 6803 (hereafter *Synechocystis*). Additionally, to identify metabolic pathways that are activated during CBB cycle initiation, dynamic ^{13}C labeling is performed immediately following light irradiation. Accurate concentration and labeling dynamics allow us to conduct short-time dynamic metabolic flux analysis (STD-MFA) to estimate non-stationary flux changes during activation of the CBB cycle. The STD-MFA results show that highly accumulated glycolytic intermediates including 3-phosphoglycerate (3PG), 2-phosphoglycerate (2PG), and phosphoenolpyruvate (PEP) are main substrates for initial CO_2 fixation. This metabolic phenomenon is further characterized by the delay of oxygen evolution in several mutant strains, in which the accumulated pool sizes of the glycolytic intermediates are found to be smaller than that of wild-type cells.

Results
Strategy for estimating metabolic flux dynamics

To estimate non-stationary metabolic flux changes, the following relationship is employed. The accumulation rate of a metabolite in a metabolic network is defined as the sum of all fluxes producing the metabolite minus the sum of all fluxes consuming the metabolite, as represented by following equation:

$$r_i(t) = \sum_j \alpha_j x_j(t) - \sum_k \alpha_k x_k(t) \qquad (1)$$

where, $x_j(t)$ is the flux through reaction j, α_j is a stoichiometric coefficient, and $r_i(t)$ is the accumulation rate of metabolite i[18]. An example with three metabolic fluxes is illustrated in Fig. 1a. To obtain accumulation rates for estimation of the flux values, it is necessary to examine the absolute concentration changes of metabolites with high time resolution. Namely, the metabolic flux during CBB cycle activation can be calculated by quantifying the changes in metabolite concentrations over highly time-resolved intervals immediately following light irradiation.

Absolute concentration changes in central carbon metabolism
We sought to determine the absolute concentration dynamics of central energy metabolites in *Synechocystis*. As light irradiation rapidly causes metabolic variations driven by increases in ATP and NADPH via the PET reaction, rapid sampling and quenching of cellular metabolic activity are necessary to accurately quantify metabolic changes during CBB cycle activation. A standardized procedure for intracellular metabolome quantification requires multiple analytical steps, including quenching of microbial metabolic activity, separation of microbial cells from growth medium, extraction of intracellular metabolites, and finally mass spectrometric metabolite analysis. These multistep processes largely impede accurate estimation of the absolute '*in vivo*' cellular metabolic state[19–21].

To overcome the above problems of the present study, a combination of two strategies were employed. First, to analyze intracellular metabolites, cells are separated from the growth medium and suspended in 1 mM $NaHCO_3$ prior to metabolite sampling to remove extracellular metabolites and medium components. Note that the exchange of medium for $NaHCO_3$ solution is a typical pretreatment for measuring photosynthetic activity of fresh-water cyanobacteria[22]. Carefully dropping the cell suspension into the extraction solution during light irradiation enables the rapid quenching and extraction of intracellular metabolites and avoids metabolic state variations during sampling. A phenol-chloroform-isoamyl alcohol (PCI) solution (pH 7.9) is selected as the extraction solution because PCI extraction enables rapid and efficient extraction of NADPH from *Synechocystis*[23]. Second, stable isotopomers of the analyte compounds are used as internal standards (IS). Experimental protocols using isotope-labeled IS are available for absolute quantification of metabolite concentrations in various microorganisms including cyanobacteria[24–26].

Figure 1b shows the experimental process for the exact determination of absolute metabolite concentration dynamics. A ^{13}C-labeled IS mixture is prepared by extracting metabolites from *Synechocystis* cells cultivated with $NaH^{13}CO_3$ as the sole carbon source. Absolute concentrations of uniformly ^{13}C-labeled metabolites are determined using capillary electrophoresis-mass spectrometry (CE-MS) based on the peak area ratios of unlabeled metabolites (with known concentration) and uniformly ^{13}C-labeled metabolites. Then, to quantify absolute concentration changes during CBB cycle activation, dark-adapted *Synechocystis* cells (grown in darkness for 12 h) are illuminated at a light intensity of 30 μmol m^{-2} s^{-1} (same as the growth light intensity) or 200 μmol m^{-2} s^{-1}. Metabolites of these cells are rapidly extracted using PCI solvent containing the ^{13}C-labeled extract as labeled IS with concentrations determined from another aliquot of the same extraction batch (Fig. 1b). Because of high physicochemical similarities between labeled and unlabeled central metabolites, both isotopomers undergo similar processes of degradation and ion suppression, and therefore accurate concentrations can be calculated[24–26].

Given that absolute metabolite concentrations are determined based on the ratio between unlabeled and uniformly labeled peak areas (Fig. 1b), the process is expected to prevent contamination of unlabeled metabolites in the ^{13}C-labeled IS. The proportions of uniformly ^{13}C-labeled metabolites (compared to the sum of unlabeled and uniformly labeled metabolites) are found to exceed 95% for all metabolites except for citrate (Supplementary Fig. 1). Figure 2 shows temporal changes in absolute metabolite concentrations after the initiation of light irradiation at 30 μmol m^{-2} s^{-1} or 200 μmol m^{-2}

s^{-1}. It should be noted that a small amount of some metabolites are extracellular (Supplementary Fig.
2). A previous study that measured *in vivo* NADPH fluorescence revealed that the amount of NADPH
in *Synechocystis* increases within 0.5 s of illumination[27]. The rapid increase of NADPH observed
within 5 s in the present work is consistent with that of the earlier *in vivo* study[27], confirming that the
current metabolomics strategy can successfully detect the rapid start of photosynthesis. As with
NADPH, the level of the energy cofactor ATP also increases rapidly. Overall, the levels of 3PG, 2PG,
and PEP decrease; the levels of other CBB cycle intermediates (except 3PG, ribulose 5-phosphate
(Ru5P) and xylulose 5-phosphate (Xu5P)) increase, while the levels of tricarboxylic acid (TCA) cycle
intermediates do not change appreciably. Considering these metabolite variations together, we inferred
that accumulated glycolysis intermediates including 3PG, 2PG, and PEP likely were converted to CBB
cycle intermediates at the photosynthetic start.
161
¹³C labeling kinetics
A direct way to confirm this hypothesis is to determine the metabolic flux distribution. However,
precise metabolic flux values cannot be determined when the number of unknown flux parameters is
larger than the number of known metabolite concentration changes in the metabolic flux model. To
accurately perform STD-MFA, we sought to refine our metabolic models by excluding metabolic
pathways that are inactive during photosynthetic activation. The metabolite concentration data shown
in Fig. 2 cannot identify inactive metabolic pathways. Even if the metabolic concentration change is
zero, metabolic flux is not necessarily zero, because such an observation instead may indicate that
metabolic inflow and outflow are balanced. In the case where the concentration of a metabolite is
stable and ¹³C is not incorporated into the metabolite during photosynthetic ¹³CO$_2$ fixation, the
metabolic fluxes around the metabolite are estimated to be very small, and therefore can be neglected.
Accordingly, we next measured the dynamic ¹³C isotope incorporation after the transition from dark
to light conditions, by starting photosynthesis while the cells are suspended in a solution of NaH¹³CO$_3$
(Fig. 1c). As a result, a transient increase in ¹³C fractions of CBB cycle intermediates is observed (Fig.
3a, Supplementary Fig. 3). On the other hand, labeling of 6-phosphogluconate (6PG) and TCA cycle
intermediates (except for malate and fumarate) is not observed. In addition, a low ¹³C fraction of the
glycogen precursor ADP-glucose[28], is observed. When considered together with the stable
concentrations of TCA and oxidative pentose phosphate (OPP) pathway intermediates (Fig. 2), these
results suggest extremely low metabolic fluxes of the TCA cycle, OPP pathway, Entner-Doudoroff
(ED) pathway, and glycogen synthesis pathway, immediately after light irradiation.
182
In addition to ribulose bisphosphate carboxylase/oxygenase (RuBisCO), the primary metabolic
enzymes that incorporate CO$_2$ are phosphoenolpyruvate carboxylase (PepC) and malic enzyme (ME),
which are both related to malate metabolism. Since malate is not labeled until 60 s after the initiation
of light irradiation (at both of the tested light intensities), ¹³C incorporation before 60 s is inferred to
reflect RuBisCO activity. The amount of incorporated ¹³C is calculated by multiplying the measured
¹³C fraction (Fig. 3a, Supplementary Fig. 3) by the absolute carbon concentration of each metabolite
(shown in Fig. 2); these analyses are presented in Supplementary Fig. 4. The total amount of
incorporated ¹³C is derived by summing the number of ¹³C atoms of all measured metabolites (Fig.
3b). Notably, ¹³C incorporation does not start until 12 s (200 μmol m^{-2} s^{-1}) or 20 s (30 μmol m^{-2} s^{-1})
after the initiation of light irradiation. On the other hand, ¹³C incorporation is observed at least 10 s
after the addition of NaH¹³CO$_3$ to cells with activated photosynthesis in response to 5 min of
illumination (Fig. 3c). Together, these results suggest that the observed delay in the start of ¹³C
incorporation is not due to cellular uptake of NaH¹³CO$_3$. Instead, these data indicate that RuBisCO-
mediated CO$_2$ fixation remains inactive during the initial stage of photosynthetic metabolism,
immediately following the start of illumination. This delayed onset of CO$_2$ fixation is consistent with
the photosynthetic induction phenomena observed in other photosynthetic organisms[29].
199
It is interesting to note that labeled carbon is incorporated into malate and fumarate after 120 s,
particularly in cells illuminated at 200 μmol m^{-2} s^{-1}, at a time when other TCA cycle intermediates
remained unlabeled (Supplementary Fig. 3). This result suggests that malate and fumarate are
generated by the reductive TCA pathway, which is known as a fermentation pathway dependent on

NADH oxidation[30]. Thus, the accumulation of NADH may drive the activity of this reductive TCA
pathway (Supplementary Fig. 5).
206
Short-term dynamic metabolic flux analysis (STD-MFA)
Temporal changes in net metabolic fluxes are calculated based on the obtained data. As mentioned
above, during the first 60 s of illumination, the observed net metabolic fluxes in pathways other than
the CBB cycle and glycolysis are very small. In fact, the concentrations of newly synthesized amino
acids do not change significantly within 60 s, even in cells illuminated at 200 μmol m^{-2} s^{-1}, although
the labeled levels of some amino acids (serine, alanine, phenylalanine, tyrosine and ornithine)
increased significantly after the first minute of illumination (Supplementary Fig. 6). Furthermore,
photorespiratory intermediates such as glycolate, glyoxylate and glycerate do not accumulate during
the first 60 s of illumination (Supplementary Fig. 7). Therefore, the metabolic pathways shown in
Supplementary Fig. 8a are used for subsequent calculations, focusing on metabolic dynamics during
the first 60 s following the initiation of illumination.
218
To solve equation (1) and conduct STD-MFA, accumulation rates for metabolites in the metabolic
model shown in Supplementary Fig. 8a are derived from the slopes obtained between two consecutive
concentration data points in Fig. 2. In addition to our use of equation (1), the CO_2 fixation rate is
calculated as the increase in the rate of total ^{13}C incorporation (Fig. 3b). Specifically, the latter
parameter is interpreted as the metabolic flux from ribulose 1,5-bisphosphate (RuBP) to 3PG, given
that the ^{13}C incorporation experiment mentioned above indicates that all carbon atoms fixed during
the first 60 s of illumination are attributable to RuBisCO activity. The net metabolic fluxes of each
time interval are estimated by solving these equations simultaneously (Fig. 4, Supplementary Fig. 8).
The estimated flux values are validated and judged to be reliable, given that almost all of the estimated
metabolite accumulation rate values (which were derived from the estimated fluxes) yielded values
that are within the mean ± SD range of the measured accumulated rate (Supplementary Fig. 9). For
more details of the calculation process and equations, see the Methods section.
231
The time course of metabolic flux and metabolite concentration distribution in Fig. 4 clearly indicates
that photosynthetic metabolism in *Synechocystis* starts by converting dark accumulated 3PG as an
initial substrate into RuBP under both two light intensities. On the other hand, the reduction fluxes of
3PG differ greatly between cells illuminated at 30 and 200 μmol m^{-2} s^{-1}, even before the initiation of
CO_2 fixation (i.e. during the first 5-20 s). In steady-state photosynthesis, the increased rate of electron
donor (NADPH) production from the PET reaction at the higher light intensities is balanced by the
rate of electron acceptor (3PG) production that is increased by the elevated CO_2 fixation rate. On the
other hand, before the onset of CO_2 fixation that follows the initiation of light irradiation, excess
reducing power can be accumulated because 3PG cannot be produced while NADPH is produced by
the PET reaction. Nevertheless, the increase in the reducing fluxes of 3PG with the higher light
intensity (200 μmol m^{-2} s^{-1}) before CO_2 fixation suggests that, even before the onset of CO_2 fixation,
the rate of 3PG consumption is balanced by the rate of NADPH production from the PET reaction,
which varies with light intensity. In addition, it can clearly be seen that 2PG and PEP are converted to
3PG under both light intensities (Fig. 4). Therefore, we propose that the accumulation of these
glycolytic intermediates provides a "stand-by" state which precedes robust initiation of
photosynthesis; thus, these intermediates are expected to play an important role in balancing cofactor
regeneration and consumption during activation of the CBB cycle.
249
Correlation between dark-accumulation of glycolytic intermediates and oxygen evolution
The effects of glycolytic intermediate accumulation on the initiation of photosynthesis are further
examined using mutants lacking enzymes involved in dark metabolism. Under dark conditions,
cyanobacteria obtain NADPH mainly via the OPP pathway[31−33]. As an initial OPP pathway substrate,
glucose 6-phosphate (G6P) is supplied by glycogen degradation through glycogen phosphorylase. In
a glycogen phosphorylase-deficient mutant of *Synechocystis*, decreased accumulation of CBB cycle
metabolites has been suggested as a cause of delayed oxygen evolution after dark incubation[9].
However, the key metabolites that determine CBB cycle activation efficiency have not been reported.

258
To better understand CBB cycle activation, the relationship between absolute metabolite concentration
and oxygen evolution activity, following 12 h dark preincubation periods, is characterized in mutants
lacking glycogen phosphorylase, glucose 6-phosphate dehydrogenase or 6-phosphogluconate
dehydrogenase (*ΔglgP*, *Δzwf*, *Δgnd*). In all strains, the increase of oxygen evolution rate (OER) is
significantly delayed when following a dark preincubation period (Fig. 5a). Especially, the OER of
dark preincubated *Δgnd* does not increase during 5 min of light irradiation. To quantify OER activation
efficiency, ratios of OER values at 60 s after light irradiation of dark preincubated cells to that of non-
dark incubated cells are calculated (Fig. 5b). The resulting OER ratio is highest in wild-type (WT)
Synechocystis, followed by *Δzwf*, *ΔglgP*, *Δgnd*, in descending order.
268
On the other hand, 9 substrates are found to be used for initial CO_2 fixation (measured metabolites
converted into RuBP during the first 60 s of light irradiation in Fig. 4): 3PG, 2PG, PEP,
dihydroxyacetone phosphate (DHAP), fructose 1,6-bisphosphate (FBP), fructose 6-phosphate (F6P),
sedoheptulose 7-phosphate (S7P), Xu5P and Ru5P. The total amounts of these initial substrates in the
12 h dark preincubated cells follows the same trend as that of the OER ratios (Fig. 5c). The three
glycolytic intermediates (3PG, 2PG, and PEP) account for 98% of the initial substrates in WT, 90%
in *Δzwf*, 99% in *ΔglgP*, and 68% in *Δgnd*. Together these results confirm that photosynthesis activation
efficiency is clearly correlated with the accumulated amounts of glycolytic intermediates. These
results demonstrate that the accumulation of glycolytic intermediates under dark conditions is essential
for rapid activation of photosynthetic metabolism.
279
280
Discussion
A new method for estimating metabolic flux kinetics
The STD-MFA method developed in this study is able to quantify rapid changes in metabolic flux
over short intervals (within a few minutes), in response to a dynamic phenomenon. This method
requires the measurement of time-resolved absolute concentration changes, uptake or evolution rate
of one or some metabolites (in present case, carbon uptake rate) and metabolic model refinement based
on isotope incorporation data before flux calculation. Previously developed dynamic metabolic flux
analysis methods typically are intended to estimate flux change over long periods of culture processes,
and require technically specialized mathematical approaches[16,17]. In contrast, STD-MFA permits the
direct determination of metabolic flux changes in exchange, albeit while focusing on short time
intervals. We expect that this method also may be of use for understanding other non-stationary rapid
metabolic changes that have been the recent focus of attention, such as the end of photosynthesis and
adaptation against oxidative stress[34,35].
294
Regulation mechanisms of the CBB cycle
Metabolic regulation of the CBB cycle includes control of enzyme activity and metabolite
accumulation. The present study reveals that accumulation of glycolytic intermediates streamlines
activation of photosynthetic metabolism. On the other hand, the presence of plant-like enzymatic
regulation in cyanobacteria is debatable while plant CBB cycle enzymes are strictly regulated
depending upon light-dark conditions[36–40]. The present *in vivo* metabolomic data in Fig. 2 is consistent
with the following enzymatic activations in *Synechocystis*.
302
Metabolic flux of conversion of 3PG remains small until 5 s (200 µmol m^{-1} s^{-1}) or 12 s (30 µmol m^{-1}
s^{-1}), indicating the presence of regulation in phosphoglycerate kinase (PGK) and/or GAPDH.
Cyanobacterial PGK and GAPDH are activated by reductive dissociation of cysteine pairs[5,41].
Transient accumulation of FBP indicates the existence of an activation period for the fructose-1,6-/
sedohepturose-1,7-bisphosphatase (FBP/SBPase) enzyme. FBP/SBPase in *Synechocystis* is activated
by the reduction of a thiol group and binding of Mg^{2+}, and this enzyme is inhibited allosterically by
the binding of AMP[42,43]. Indeed, a rapid decrease in AMP concentration was confirmed after light
irradiation (Supplementary Fig. 5). In addition, temporal accumulation of DHAP upon irradiation with
200 µmol m^{-2} s^{-1} light may be attributable to the transient rate-limitation of FBP/SBPase. Similar to

the behavior of FBP, the accumulation and subsequent decrease of RuBP suggests a phase shift from
inactivated to activated RuBisCO. RuBisCO activity is activated by various factors: (a) carbamylation
to the activator lysine and binding of a magnesium ion to the carbamate anion as essential
regulations[44,45], (b) effectors such as orthophosphate[46], several phosphorylated sugars and NADPH[47],
(c) cysteine redox state[48,49], (d) RuBisCO activase[37]. Among these activation processes, *Synechocystis*
lacks RuBisCO activase. In addition, cyanobacteria encapsulate RuBisCO enzymes into protein-
encased microcompartments called the carboxysome[50]. Initiation of CO_2 fixation requires entrance of
RuBisCO substrate, RuBP into the carboxysome. Although kinetics of this RuBP uptake is unknown,
this process might affect activation kinetics of CBB cycle. Note that similar dynamics of the RuBP
pool size has been observed after a dark-to-light transition in plant leaves, and is associated with
RuBisCO activation[51,52].
323
324 Free energy analysis provides further insight into the regulation in the CBB cycle. Thermodynamic
325 feasibility is estimated by the reaction Gibbs energy change ($\Delta_r G$), calculated based on the following
326 equation:

$$\Delta_r G = \Delta_r G° + RT ln\, Q \qquad (2)$$

where, $\Delta_r G°$ is the standard reaction Gibbs energy change, R is the ideal gas constant, T is the
temperature, and Q is the reaction quotient, which includes concentration terms. The availability of
absolute concentration data enables assessment of the thermodynamically feasible flux directions.
Corresponding $\Delta_r G°$ values are obtained from the eQuilibrator database[53] (Supplementary Table 1).
Interestingly, conversion of 3PG to DHAP is thermodynamically allowed even under dark conditions
as the $\Delta_r G$ for the reaction from 3PG forming DHAP is a large negative value (Supplementary Fig.
10). This non-equilibrium state supports the presence of dark inactivation of PGK and GAPDH as
mentioned above. The $\Delta_r G$ value of conversion from PEP to pyruvate catalyzed by pyruvate kinase
(PK) is found to be a very negative value compared with that of heterotrophic organisms[12]
(Supplementary Fig. 10). Along with GAPDH and PGK, PK activity regulation may contribute to the
observed accumulation of 3PG, 2PG, and PEP.
339

Essential metabolic pathways for dark accumulation of glycolytic intermediates
Accumulation of glycolytic intermediates decreases in *Δzwf*, *ΔglgP*, and *Δgnd*, suggesting that the
accumulated glycolytic intermediates are derived from glycogen degradation and the OPP pathway.
The first step of the OPP pathway is the irreversible conversion of G6P into 6PG by glucose-6-
phosphate dehydrogenase (G6PDH). Then, 6PG is decarboxylated into Ru5P by 6-phosphogluconate
dehydrogenase (6PGDH). Accumulation of initial CO_2 fixation substrates in *Δgnd* is much lower than
in *Δzwf* (Fig. 5c). This may be explained by the G6PDH-mediated conversion of glycogen-derived
G6P into 6PG, with will then accumulate if it is not readily converted to other metabolites. Indeed,
6PG accumulation in *Δgnd* (45.0 nmol mg^{-1} DCW) is about 4000-fold higher than in the wild-type
(0.011 nmol mg^{-1} DCW) after dark incubation. On the other hand, G6P in *Δzwf* can be metabolized
by other pathways such as the Embden-Meyerhof-Parnas (EMP) pathway, which might prevent the
depletion of glycolytic intermediates.
352

Significance and conservation of intermediate pool size
Although concentration robustness is generally considered to be a fundamental factor for maintaining
cellular functions[7], the precise role of metabolic robustness in photosynthetic organisms has not been
clarified. Balancing photosynthetic supply and metabolic demand of NADPH is important for
operating stable CO_2 fixation and avoiding oxidative stress by redox imbalance state[8,54]. In this study,
it is suggested that accumulation of glycolytic intermediates acts as a buffering system that
compensates for the gap between supply and demand of NADPH, and ensures rapid photosynthesis
start. Indeed, lower accumulation of the glycolytic intermediates resulted in inefficient photosynthetic
activity (Fig. 5). Previous study showed that 3PG is the most abundant in CBB cycle intermediates in
three cyanobacterial species including *Synechocystis*[25]. This metabolite distribution is characteristic
compared with that of heterotrophic organisms[12] (Supplementary Fig. 11). Moreover, rapid
consumption of phosphoglycerate upon illumination like Fig. 2 was also observed in algae, the
chloroplast of spinach protoplasts, and leaves of wheat[55–57]. These facts support functional

significance and conservation of accumulation of the glycolytic intermediates for robust
photosynthesis.
368
Methods
Growth conditions
This work employed a glucose-tolerant wild-type strain of *Synechocystis*[58] that was grown and stored
on solid BG-11 media plates containing 1.5% Bacto agar. Pre-culturing was performed in 200 mL
flasks containing 50 mL each of liquid BG-11 medium buffered with 20 mM 2-[4-(2-hydroxyethyl)-
1-piperazinyl]ethanesulfonic acid (HEPES, pH 7.5); each flask was inoculated with cells harvested
from the agar plates. These flasks then were cultured at 30 ºC in ambient air with shaking at 100 rpm;
illumination at an intensity of 30 µmol m^{-2}s^{-1} was provided by white light-emitting diode (LED) lights.
The main culture was initiated by adding pre-culture grown for 4 days into 70 mL of the BG-11
medium to give an optical density of 0.05 at 730 nm (OD$_{730}$). The growth condition of the main culture
was the same as that of the pre-culture. The dry cell weight (DCW) was measured after harvesting
cells by centrifugation and lyophilization. The obtained relationship between the OD$_{730}$ value and
DCW (DCW = 0.1998 × OD$_{730}$ × V + 0.0974; R^2=0.982; n=9; V is volume of cell suspension) was
used for estimating DCW from the OD$_{730}$ value.
383
Preparation of ^{13}C isotope-labeled internal standard (^{13}C-IS)
Synechocystis cells in pre-culture were inoculated to an OD$_{730}$ of 0.05 in a serum bottle containing 60
mL modified BG-11 medium supplemented with 100 mM HEPES (pH 7.5) and 25 mM NaH^{13}CO$_3$ (in
place of the unlabeled Na$_2$CO$_3$ included in the standard BG-11 medium). The culture serum bottle was
sealed, followed by flushing with N$_2$ gas to remove unlabeled ambient CO$_2$. The growth conditions
used for the labeled culture were the same as those used for the pre-culture and main culture. Cells
grown for 3 days were harvested and re-suspended in fresh modified BG-11 to avoid pH increase
during cultivation with NaH^{13}CO$_3$ as the sole carbon source. The labeled culture then was grown under
the same conditions for another 3 days, followed by extraction of uniformly labeled metabolites, as
described below.
394
The entire volume of the labeled culture was harvested by centrifugation, washed with 1 mM
NaH^{13}CO$_3$ solution, and then re-suspended in 9 mL of 1 mM NaH^{13}CO$_3$. The resulting cell suspension
was illuminated for 90 s with white LED light at an intensity of 200 µmol m^{-2}s^{-1} (to maximize
extraction yield of CBB cycle intermediates), and placed into a mixture of 9 mL PCI (pH 7.9, Nacalai
Tesque, Kyoto, Japan) and 60 µL 0.5 M N-cyclohexyl-3-aminopropanesulfonic acid (CAPS, pH 10.6)
for metabolite extraction. The extraction mixture was centrifuged at 4 ºC at 8,000 ×*g* for 3 min. Twelve
to 13 fractions of the water layer were then aliquoted into microtubes and lyophilized.
402
403 One of the dried aliquots was dissolved in 20 µL of standard mixture containing prepared unlabeled
404 compounds. The sample was analyzed by capillary electrophoresis-mass spectrometry (CE-MS) (CE,
405 Agilent G7100; MS, Agilent G6224AA LC/MSD TOF; Agilent Technologies, Palo Alto, CA, USA)
406 as described previously[59]. Absolute concentrations of the uniformly labeled isotopomer (C_n) were
407 derived as follows:

$$C_n = \frac{S_n}{S_0} C_0 \tag{3}$$

409 where, S_n and S_0 are the peak areas of uniformly ^{13}C-labeled (namely, the number of ^{13}C atoms in a *n*-
410 carbon compound is equal to *n*) and unlabeled standard signals, respectively, and C_0 is the prepared
411 concentration of the unlabeled standard.
412
Metabolite extraction with rapid sampling and analysis
To perform metabolite sampling for analyses of absolute concentration and ^{13}C incorporation after the
transition from dark to light, a main culture grown for 6 days was dark-adapted for 12 h. A calculated
volume (90/OD$_{730}$, in mL) of the dark-adapted cell culture was harvested by centrifugation at 8000×*g*
for 5 min. The cells were washed with 4 mL of 1 mM NaHCO$_3$ (for absolute concentration analysis)
or 1 mM NaH^{13}CO$_3$ (for ^{13}C incorporation analysis), and harvested again by centrifugation. The

washed cells were resuspended in 6.8 mL of 1 mM $NaHCO_3$ (for absolute concentration analysis) or 1 mM $NaH^{13}CO_3$ (for ^{13}C incorporation analysis). Nine aliquots (750 µL each) of the resulting cell suspension were then loaded into clear pipet tips (Supplementary Fig. 12). Up to this point, light irradiation was avoided during the procedure to keep the cells in the dark-adapted state. Irradiation of all the aliquots with white LED light at an intensity of 200 ± 10 µmol $m^{-2}s^{-1}$ or 30 ± 2 µmol $m^{-2}s^{-1}$ was initiated, followed by sampling of the cells at the indicated time points by adding into 750 µL PCI supplemented with ^{13}C-IS dissolved in 20 µL water (for absolute concentration analysis) or with 5 µL of 0.5 M CAPS (pH 10.6) (for ^{13}C incorporation analysis). The cell suspension before illumination was extracted as the nominal 0-s sample. To precisely determine the amount of extracted cells, the OD_{730} of the final cell suspension was measured. After the extraction mixtures were centrifuged at 4 ºC at $12,000\times g$ for 2 min, harvested aliquots of the water layer were lyophilized.

The lyophilized samples were dissolved in 20 µL of 0.4 mM piperazine-1,4-bis(2-ethanesulfonic acid) (PIPES) and analyzed by CE-MS as described above. Absolute concentrations of each metabolite (C') were derived as follows:

$$C' = \frac{S'_0}{WS'_n}C_n V \tag{4}$$

where, S'_0 and S'_n are the peak areas of unlabeled and uniformly ^{13}C-labeled IS signals, respectively, W represents DCW, and V is the sample volume for CE-MS analysis. The intracellular concentration (mM) was calculated by using the values: 5×10^7 cells/mL/OD_{730}[60] and 4.4×10^{-15} L/cell[61]. ^{13}C fractions were calculated as described previously[30].

Short-term dynamic metabolic flux analysis (STD-MFA)
To estimate the temporal changes of 18 fluxes in the metabolic map shown in Supplementary Fig. 8, 18 metabolite accumulation rates $r_i(t)$ in equation (1) were derived from the slope taken between two consecutive concentration data points shown in Fig. 2. Accumulation rates of metabolites under the detection limit (E4P, R5P, GAP <0.001 nmol/mg DCW) and peak-unidentified metabolites (BPG and SBP) were set to zero. In addition, the CO_2 fixation rate derived from the slope taken between two consecutive data points shown in Fig. 3 was regarded as the metabolic flux of RuBisCO (i.e., the conversion of RuBP to 3PG). The set of 19 equations, including the 18 unknown fluxes in the network, was represented in matrix notation as follows:

$$\mathbf{A}\mathbf{x}(t) = \mathbf{r}(t) \tag{5}$$

where, A is the $19 \cdot 18$ matrix of stoichiometric coefficients, $\mathbf{x}(t)$ is the 18-dimensional flux vector, and r(t) is the 19-dimensional metabolite accumulation rate vector. The specific form of equation (4) is shown in Supplementary Fig. 8b. The weighted least squares solution to equation (5) is as follows:

$$\hat{\mathbf{x}}(t) = (\mathbf{A}^T\mathbf{W}\mathbf{A})^{-1}\mathbf{A}^T\mathbf{W}\bar{\mathbf{r}}(t) \tag{6}$$

where, the superscripts ˆ and ⁻ denote estimated and measured quantities (respectively), and $\mathbf{W}$ is a 19-dimensional weighting diagonal matrix, whose diagonal elements are the reciprocal of the variance of the measured metabolic accumulation rate values. Variance values for the accumulation rate of E4P, R5P, GAP, BPG and SBP, and for the CO_2 fixation rate, were set to 10^{-15} (i.e., a value that was effectively zero). To evaluate the reliability of the estimated fluxes, the metabolite accumulation rates were calculated as follows:

$$\mathbf{A}\hat{\mathbf{x}}(t) = \hat{\mathbf{r}}(t) \tag{7}$$

The estimated and measured metabolite accumulation rate values are compared in Supplementary Fig. 9.

References

1. Flombaum, P. et al. Present and future global distributions of the marine Cyanobacteria *Prochlorococcus* and *Synechococcus*. *Proc Natl Acad Sci U S A* **110**, 9824-9829 (2013).
2. Hutchins, D. A. & Fu, F. Microorganisms and ocean global change. *Nat. Microbiol.* **2**, 17058 (2017).

3. Sánchez-Baracaldo, P., Bianchini, G., Wilson, J. D. & Knoll, A. H. Cyanobacteria and biogeochemical cycles through Earth history. *Trends Microbiol.* **30**, 143-157 (2022).

4. Tamoi, M., Miyazaki, T., Fukamizo, T. & Shigeoka, S. The Calvin cycle in cyanobacteria is regulated by CP12 via the NAD(H)/NADP(H) ratio under light/dark conditions. *Plant J.* **42**, 504-513 (2005).

5. McFarlane, C. R. et al. Structural basis of light-induced redox regulation in the Calvin-Benson cycle in cyanobacteria. *Proc. Natl. Acad. Sci. USA.* **116**, 20984-20990 (2019).

6. Gurrieri, L., Fermani, S., Zaffagnini, M., Sparla, F. & Trost, P. Calvin-Benson cycle regulation is getting complex. *Trends Plant Sci.* **26**, 898-912 (2021).

7. Eloundou-Mbebi, J. M. et al. A network property necessary for concentration robustness. *Nat. Commun.* **7**, 13255 (2016).

8. Matuszyńska, A., Saadat, N. P. & Ebenhöh, O. Balancing energy supply during photosynthesis - a theoretical perspective. *Physiol. Plant* **166**, 392-402 (2019).

9. Shimakawa, G. et al. Respiration accumulates Calvin cycle intermediates for the rapid start of photosynthesis in *Synechocystis* sp. PCC 6803. *Biosci. Biotechnol. Biochem.* **78**, 1997-2007 (2014).

10. Shinde, S. et al. Glycogen Metabolism Supports Photosynthesis Start through the Oxidative Pentose Phosphate Pathway in Cyanobacteria. *Plant Physiol.* **182**, 507-517 (2020).

11. Bennett, B. D. et al. Absolute metabolite concentrations and implied enzyme active site occupancy in *Escherichia coli*. *Nat. Chem. Biol.* **5**, 593-599 (2009).

12. Park, J. O. et al. Metabolite concentrations, fluxes and free energies imply efficient enzyme usage. *Nat. Chem. Biol.* **12**, 482-489 (2016).

13. Shimizu, H. & Toya, Y. Recent advances in metabolic engineering-integration of in silico design and experimental analysis of metabolic pathways. *J. Biosci. Bioeng.* **132**, 429-436 (2021).

14. Sauer, U. Metabolic networks in motion: ^{13}C-based flux analysis. *Mol. Syst. Biol.* **2**, 62 (2006).

15. Young, J. D., Shastri, A. A., Stephanopoulos, G. & Morgan, J. A. Mapping photoautotrophic metabolism with isotopically nonstationary ^{13}C flux analysis. *Metab. Eng.* **13**, 656-665 (2011).

16. Leighty, R. W. & Antoniewicz, M. R. Dynamic metabolic flux analysis (DMFA): a framework for determining fluxes at metabolic non-steady state. *Metab. Eng.* **13**, 745-755 (2011).

17. Niklas, J., Schräder, E., Sandig, V., Noll, T. & Heinzle, E. Quantitative characterization of metabolism and metabolic shifts during growth of the new human cell line AGE1.HN using time resolved metabolic flux analysis. *Bioprocess Biosyst. Eng.* **34**, 533-545 (2011).

18. Vallino, J. J. & Stephanopoulos, G. Metabolic flux distributions in *Corynebacterium glutamicum* during growth and lysine overproduction. *Biotechnol. Bioeng.* **41**, 633-646 (1993).

19. Pinu, F. R., Villas-Boas, S. G. & Aggio, R. Analysis of Intracellular Metabolites from Microorganisms: Quenching and Extraction Protocols. *Metabolites.* **7**, E53 (2017).

20. Rabinowitz, J. D. & Kimball, E. Acidic acetonitrile for cellular metabolome extraction from *Escherichia coli. Anal. Chem.* **79**, 6167-6173 (2007).

21. Annesley, T. M. Ion suppression in mass spectrometry. *Clin. Chem.* **49**, 1041-1044 (2003).

22. Shimakawa, G. et al. FLAVODIIRON2 and FLAVODIIRON4 proteins mediate an oxygen-dependent alternative electron flow in *Synechocystis* sp. PCC 6803 under CO_2-limited conditions. *Plant. Physiol.* **167**, 472-480 (2015).

23. Tanaka, K. et al. Quantification of NAD(P)H in cyanobacterial cells by a phenol extraction method. *Photosynth. Res.* **148**, 57-66 (2021).

24. Wu, L. et al. Quantitative analysis of the microbial metabolome by isotope dilution mass spectrometry using uniformly ^{13}C-labeled cell extracts as internal standards. *Anal. Biochem.* **336**, 164-171 (2005).

25. Dempo, Y., Ohta, E., Nakayama, Y., Bamba, T. & Fukusaki, E. Molar-based targeted metabolic profiling of cyanobacterial strains with potential for biological production. *Metabolites.* **4**, 499-516 (2014).

26. Nishino, S., Okahashi, N., Matsuda, F. & Shimizu, H. Absolute quantitation of glycolytic

524    intermediates reveals thermodynamic shifts in *Saccharomyces cerevisiae* strains lacking
525    *PFK1* or *ZWF1* genes. *J. Biosci. Bioeng.* **120**, 280-286 (2015).

526 27. Kauny, J. & Sétif, P. NADPH fluorescence in the cyanobacterium *Synechocystis* sp. PCC
527    6803: a versatile probe for in vivo measurements of rates, yields and pools. *Biochim. Biophys.*
528    *Acta.* **1837**, 792-801 (2014).

529 28. Yoo, S. H., Lee, B. H., Moon, Y., Spalding, M. H. & Jane, J. L. Glycogen synthase isoforms
530    in *Synechocystis* sp. PCC6803: identification of different roles to produce glycogen by
531    targeted mutagenesis. *PLoS One.* **9**, e91524 (2014).

532 29. Pearcy, R. W., Krall, J. P. & Sassenrath-Cole, G. F. Photosynthesis in Fluctuating Light
533    Environments. In: Baker NR, ed. *Photosynthesis and the Environment.* Dordrecht: Kluwer,
534    321–346 (1996).

535 30. Hasunuma, T., Matsuda, M., Kato, Y., Vavricka, C. J. & Kondo, A. Temperature enhanced
536    succinate production concurrent with increased central metabolism turnover in the
537    cyanobacterium *Synechocystis* sp. PCC 6803. *Metab. Eng.* **48**, 109-120 (2018).

538 31. Wan, N. et al. Cyanobacterial carbon metabolism: Fluxome plasticity and oxygen dependence.
539    *Biotechnol. Bioeng.* **114**, 1593-1602 (2017).

540 32. Welkie, D. G. et al. Genome-wide fitness assessment during diurnal growth reveals an
541    expanded role of the cyanobacterial circadian clock protein KaiA. *Proc. Natl. Acad. Sci. U S*
542    *A* **115**, E7174-E7183 (2018).

543 33. Hatano, J. et al. NADPH production in dark stages is critical for cyanobacterial photocurrent
544    generation: a study using mutants deficient in oxidative pentose phosphate pathway.
545    *Photosynth. Res.* 10.1007/s11120-022-00903-0

546 34. Christodoulou, D. et al. Reserve Flux Capacity in the Pentose Phosphate Pathway Enables
547    *Escherichia coli*'s Rapid Response to Oxidative Stress. *Cell Syst.* **6**, 569-578.e7 (2018).

548 35. Maruyama, M. et al. Time-resolved analysis of short term metabolic adaptation at dark
549    transition in *Synechocystis* sp. PCC 6803. *J. Biosci. Bioeng.* **128**, 424-428 (2019).

550 36. Buchanan, B. B. Role of light in the regulation of chloroplast enzymes. *Ann. Rev. Plant*
551    *Physiol.* **31**, 341-374 (1980).

552 37. Portis, Jr. A. R. The regulation of Rubisco by Rubisco activase. *J. Exp. Bot.* **46**, 1285-1291
553    (1995).

554 38. Tamoi, M., Murakami, A., Takeda, T. & Shigeoka, S. Lack of Light/Dark Regulation of
555    Enzymes Involved in the Photosynthetic Carbon Reduction Cycle in Cyanobacteria,
556    Synechococcus PCC 7942 and Synechocystis PCC 6803. *Biosci. Biotechnol. Biochem.* **62**,
557    374-376 (1998).

558 39. Michelet, L. et al. Redox regulation of the Calvin-Benson cycle: something old, something
559    new. *Front Plant Sci* **4**, 470 (2013).

560 40. Gütle, D. D., Roret, T., Hecker, A., Reski, R. & Jacquot, J. P. Dithiol disulphide exchange in
561    redox regulation of chloroplast enzymes in response to evolutionary and structural constraints.
562    *Plant Sci.* **255**, 1-11 (2017).

563 41. Tsukamoto, Y., Fukushima, Y., Hara, S. & Hisabori, T. Redox control of the activity of
564    phosphoglycerate kinase in *Synechocystis* sp. PCC6803. *Plant Cell. Physiol.* **54**, 484-491
565    (2013).

566 42. Feng, L. et al. Structural and biochemical characterization of fructose-1,6/sedoheptulose-1,7-
567    bisphosphatase from the cyanobacterium *Synechocystis* strain 6803. *FEBS J.* **281**, 916-926
568    (2014).

569 43. Mallén-Ponce, M. J., Huertas, M. J., Sánchez-Riego, A. M. & Florencio, F. J. Depletion of
570    m-type thioredoxin impairs photosynthesis, carbon fixation, and oxidative stress in
571    cyanobacteria. *Plant Physiol.* **187**, 1325-1340 (2021).

572 44. Lorimer, G. H., Badger, M. R. & Andrews, T. J. The Activation of Ribulose-1,5,-bisphosphate
573    carboxylase by carbon dioxide and magnesium ions. Equilibria, kinetics, a suggested
574    mechanisms, and physiological implications. *Biochem.* **15**, 529-536 (1976).

575 45. Miziorko, H. M. Ribulose-1,5-biphosphate carboxylase. Evidence in support of the existence
576    of distinct CO_2 activator and CO_2 substrate sites. *J. Biol. Chem.* **254**, 270-272 (1979).

577 46. Marcus, Y. & Gurevitz, M. Activation of cyanobacterial RuBP-carboxylase/oxygenase is

facilitated by inorganic phosphate via two independent mechanisms. *Eur. J. Biochem.* **267**,
5995-6003 (2000).
47. Tabita, F. R. & Colletti, C. Carbon dioxide assimilation in cyanobacteria: regulation of
ribulose, 1,5-bisphosphate carboxylase. *J. Bacteriol.* **140**, 452-458 (1979).
48. Marcus, Y., Altman-Gueta, H., Finkler, A. & Gurevitz, M. Dual role of cysteine 172 in redox
regulation of ribulose 1,5-bisphosphate carboxylase/oxygenase activity and degradation. *J.*
Bacteriol. **185**, 1509-1517 (2003).
49. Moreno, J., García-Murria, M. J. & Marín-Navarro, J. Redox modulation of Rubisco
conformation and activity through its cysteine residues. *J. Exp. Bot.* **59**, 1605-1614 (2008).
50. Rae, B. D., Long, B. M., Badger, M. R. & Price, G. D. Functions, compositions, and evolution
of the two types of carboxysomes: polyhedral microcompartments that facilitate CO_2 fixation
in cyanobacteria and some proteobacteria. *Microbiol. Mol. Biol. Rev.* **77**, 357-379 (2013).
51. Perchorowicz, J. T., Raynes, D. A. & Jensen, R. G. Light limitation of photosynthesis and
activation of ribulose bisphosphate carboxylase in wheat seedlings. *Proc. Natl. Acad. Sci.*
USA. **78**, 2985-2989 (1981).
52. Seemann, J. R., Kirschbaum, M. U., Sharkey, T. D. & Pearcy, R. W. Regulation of Ribulose-
1,5-Bisphosphate Carboxylase Activity in *Alocasia macrorrhiza* in Response to Step Changes
in Irradiance. *Plant Physiol.* **88**, 148-152 (1988).
53. Beber, M. E. et al. eQuilibrator 3.0: a database solution for thermodynamic constant
estimation. *Nucleic Acids Res.* **50**, D603-D609 (2022).
54. Shimakawa, G. & Miyake, C. Oxidation of P700 Ensures Robust Photosynthesis. *Front. Plant*
Sci. **9**, 1617 (2018).
55. Bassham, J. A., Shibata, K., Steenberg, K., Bourdon, J. and Calvin, M. The photosynthetic
cycle and respiration: Light-dark transients. *J. Am. Chem. Soc.* **78**, 4120-4124 (1956).
56. Stitt, M., Wirtz, W. & Heldt, H. W. Metabolite levels during induction in the chloroplast and
extrachloroplast compartments of spinach protoplasts. *Biochim. Biophys. Acta.* **593**, 85-102
(1980).
57. Kobza, J. & Edwards, G. E. The photosynthetic induction response in wheat leaves: net CO_2
uptake, enzyme activation, and leaf metabolites. *Planta* **171**, 549-559 (1987).
58. Williams, J. G. K. Construction of specific mutations in photosystem II photosynthetic
reaction center by genetic engineering methods in *Synechocystis* 6803. *Methods Enzymol.* **167**,
766-778 (1988).
59. Hasunuma, T. et al. Dynamic metabolic profiling of cyanobacterial glycogen biosynthesis
under conditions of nitrate depletion. *J. Exp. Bot.* **64**, 2943-2954 (2013).
60. Kohga, H., Saito, Y., Kanamaru, M., Uchiyama, J. & Ohta, H. The lack of the cell division
protein FtsZ induced generation of giant cells under acidic stress in cyanobacterium
Synechocystis sp. PCC6803. *Photosynth. Res.* **150**, 343-356 (2021).
61. Cameron, J. C. & Pakrasi, H. B. Essential role of glutathione in acclimation to environmental
and redox perturbations in the cyanobacterium *Synechocystis* sp. PCC 6803. *Plant Physiol.*
154, 1672-1685 (2010).

Acknowledgements
We thank Prof. C. Miyake (Kobe University) and Dr. Ginga Shimakawa (Kwansei Gakuin University)
for kindly providing the mutants used in this work. This work was supported by the Mirai Program
[grant number JPMJMI19E4] from the Japan Science and Technology Agency (JST) of the Ministry
of Education, Culture, Sports, Science, and Technology (MEXT), Japan, and JSPS KAKENHI Grant
Number 21J00113 and 22K15142.

625
626   **Author contributions**
627   K.T. and T.H. designed the research. K.T. and M.M. performed research. K.T. and T.S. analyzed data.
628   K.T., T.S., C.J.V., A.K. and T.H. wrote the article.

629
630   **Competing interests**

631   The authors declare no competing interests.
632
633
634

635 **Figures**
636

637
Fig. 1 Process for estimating temporal changes in metabolic flux. a, Relationship between
concentration change and metabolic flux. r_i(t) is the accumulation rate of metabolite M_i. x_n represents
the metabolic flux of the nth reaction. **b,** Experimental procedure for quantifying absolute metabolite
concentration. **c,** Schematic example of isotope-labeled state during $^{13}CO_2$ incorporation. Empty and
red-filled circles represent unlabeled and labeled carbon atoms, respectively. Abbreviations: CBB,
Calvin-Benson-Bassham; CEMS, capillary electrophoresis-mass spectrometry; F6P, fructose 6-
phosphate; IS, internal standard; RuBP, ribulose 1,5-bisphosphate; TCA, tricarboxylic acid; 2OG, 2-
oxoglutarate; 3PG, 3-phosphoglycerate.
646

647
648

649
Fig. 2 Rapid temporal changes in absolute metabolite concentrations during the initiation of
photosynthesis. Intracellular metabolites are extracted at the indicated times after transition from
darkness to light-irradiated condition at light intensities of 200 μmol m^{-2} s^{-1} (red line) or 30 μmol m^{-2}
s^{-1} (blue line). Levels of glyceraldehyde 3-phosphate (GAP), erythrose 4-phosphate (E4P), and ribose
5-phosphate (R5P) are below the detection limit (0.001 nmol/mg DCW). 1,3-Bisphosphoglycerate
(BPG) and sedoheptulose 1,7-bisphosphate (SBP) are not identified due to lack of commercial reagent
for use as a standard. Metabolites are grouped by different colors according to the central schematic
metabolic network. Abbreviations: DCW, dry cell weight; DHAP, dihydroxyacetone phosphate; FBP,
fructose 1,6-bisphosphate; F6P, fructose 6-phosphate; G1P, glucose 1-phosphate; G6P, glucose 6-
phosphate; 2OG, 2-oxoglutarate; PEP, phosphoenolpyruvate; 6PG, 6-phosphogluconate; 2PG, 2-
phosphoglycerate; 3PG, 3-phosphoglycerate; RuBP, ribulose 1,5-bisphosphate; Ru5P, ribulose 5-
phosphate; S7P, sedoheptulose 7-phosphate; Xu5P, Xylulose 5-phosphate. Values are the mean ± SD
(bars) of three independent experiments. Significant differences between the two light intensities were
evaluated by a two-tailed non-paired Student's t test (* $p < 0.05$).
664
665

667
668
Fig. 3 Incorporation of ^{13}C atoms during CBB cycle activation. (a) Time course of ^{13}C fraction of
metabolites following light irradiation at 200 μmol m^{-2} s^{-1} (red line) or 30 μmol m^{-2} s^{-1} (blue line).
Values are the mean ± SD (bars) of three biological replicates. (b) Total absolute amount of ^{13}C-labeled
carbon atoms. The number of incorporated ^{13}C atoms is derived from absolute concentration and ^{13}C
fraction data. Black, green, and orange lines indicate total amount of ^{13}C atoms of all measured
metabolites, CBB cycle and OPP pathway metabolites (G6P, 6PG, and metabolites in green zone in
Fig. 2), and TCA cycle and lower glycolysis metabolites (2PG, PEP, pyruvate, and metabolites in
yellow zone in Fig. 2). Left panel: 200 μmol m^{-2} s^{-1}, right panel: 30 μmol m^{-2} s^{-1}. (c) Effect of pre-
illumination on the start of ^{13}C incorporation. ^{13}C fractions of 3-phosphoglycerate (left panel) and
ribulose 1,5-bisphosphate (right panel) after 12 s of illumination (200 μmol m^{-2} s^{-1}) of the 12-h dark-
adapted cells were lower than those at 10 s after addition of 1 mM NaH^{13}CO$_3$ to cells that had been
pre-illuminated (200 μmol m^{-2} s^{-1}) for 5 min. Presumably, the higher ^{13}C incorporation rate in pre-
illuminated cells reflects the fact that CO$_2$ fixation already had been activated (by pre-illumination)
prior to the addition of NaH^{13}CO$_3$. Thus, lower ^{13}C incorporation by dark-adapted cells can be
attributed to the low activation state of CO$_2$ fixation during photosynthetic induction. Values are the
mean ± SD (bars) of three biological replicates. Significant differences were evaluated by a two-tailed
non-paired Student's t test; p values were as indicated.
686

 Fig. 4 Estimated change in metabolic flux after initiation of light irradiation at 200 μmol m⁻² s⁻¹ (upper row) and 30 μmol m⁻² s⁻¹ (lower row). Metabolic fluxes are indicated by the thickness of arrows with values indicated by the scale bar in the lower right corner. Metabolite concentrations at the beginning and end of each time interval are indicated by the diameters of blue and orange circles, respectively. Overlap between orange and blue circles (namely invariant concentration during a time interval) is shown in gray. Abbreviations are the same as in the Figure 2 legend.

Fig. 5 Effect of dark metabolism on initiation of photosynthesis and accumulation of substrates for initial CO₂ fixation. (a) Oxygen evolution rate (OER) of wild-type (WT) and mutants. Orange and blue lines indicate OER transients of light-adapted and 12 h-dark-adapted cells after light irradiation (200 μmol m^{-2} s^{-1}). (b) Ratio of OER value of dark-adapted cells to that of light-adapted cells at 60 s are compared among mutants. (c) Absolute concentrations of substrates for initial CO₂ fixation in 12 h dark-adapted cells. Values are the mean $\pm$ SD (bars) of three biological replicates. Abbreviations are the same as in the Figure 2 legend.

706
707