Title

Accumulated 3-phosphoglycerate in darkness is a primary starting substrate for stepwise

photosynthetic induction process in cyanobacteria

Authors

Kenya Tanaka[1,2], Tomokazu Shirai[3,4], Mami Matsuda[4], Akihiko Kondo[1,3,4,5], Tomohisa

Hasunuma[1,3,4,*]

Affiliations

1. Engineering Biology Research Center, Kobe University, 1-1 Rokkodai, Nada, Kobe 657-8501,

Japan

2. Research Center for Solar Energy Chemistry, Graduate School of Engineering Science, Osaka

University, Toyonaka, Osaka 560-8531, Japan

3. RIKEN Center for Sustainable Resource Science, 1-7-22 Suehiro-cho, Tsurumi-ku, Yokohama,

Kanagawa 230-0045, Japan

4. Graduate School of Science, Innovation and Technology, Kobe University, 1-1 Rokkodai, Nada,

Kobe 657-8501, Japan

5. Department of Chemical Science and Engineering, Graduate School of Engineering, Kobe

University, 1-1 Rokkodai, Nada, Kobe 657-8501, Japan

*Corresponding author: Tomohisa Hasunuma

Email: hasunuma@port.kobe-u.ac.jp

Abstract

In photosynthetic organisms, a smooth metabolic transition from dark to light conditions is essential

for efficient CO_2 fixation and growth. Dark-to-light transition leads to activation of the enzymes of

the Calvin-Benson-Bassham (CBB) cycle and initiates CO_2 fixation metabolism using energy and

redox equivalents provided by the photosynthetic electron transfer reaction. While many enzymatic

mechanisms of the light activation of the CBB cycle have been clarified to date, the metabolic

distribution and dynamics underlying the smooth start of photosynthetic metabolism remain unclear.

Here, we show rapid (on a time scale of seconds) dynamics of changes in absolute metabolite

concentration and ^{13}C tracer incorporation during photosynthetic induction (PI) in the

cyanobacterium *Synechocystis* sp. PCC 6803. By combining these data, we were able to estimate

time-resolved nonstationary dynamics of metabolic fluxes during PI. This novel metabolic analysis

indicated that the initial step of PI consists of the reduction of 3-phosphoglycerate (3PG), a

metabolite that accumulates in dark conditions; notably, this step occurs before subsequent

generation of ribulose-1,5-bisphosphate (RuBP) and CO_2 fixation by ribulose-1,5-bisphosphate

carboxylase/oxygenase (Rubisco). *In vivo* oxygen evolution measurement in the presence of a CBB

cycle inhibitor further confirmed that the accumulated 3PG is a major initial substrate for RuBP

generation and functions as a key electron sink before the initiation of CO_2 fixation. Our results

indicate a fundamental mechanism supporting the seamless activation of photosynthetic metabolism

after dark-to-light transitions.

Main text

      In leaves and isolated chloroplasts, there is known to be a delay between the onset of illumination and achieving a maximum rate of photosynthetic CO_2 fixation[1]. Since this photosynthetic induction (PI) process was suggested more than a century ago[2], the underlying mechanisms have been investigated intensively. Since the 1960s, the activation processes of many of the enzymes in the Calvin-Benson-Bassham (CBB) cycle have been studied[3], including those of glyceraldehyde 3-phosphate dehydrogenase (GAPDH), fructose-1,6-bisphosphatase (FBPase), sedoheptulose-1,7-bisphosphatase (SBPase), phosphoribulokinase (PRK), and ribulose-1,5-bisphosphate carboxylase/oxygenase (Rubisco). The activities of these enzymes and known to be altered by changes in cellular environmental properties such as redox state, energy charge, and ion concentration; these changes themselves reflect the effects of the photosynthetic electron transfer reaction[4-6]. Eventually, an overview of PI process in the leaves of land plants was developed, and can be summarized as follows: enzymes of RuBP regeneration pathways are activated in the first 1-2 min of illumination, followed by completion of slower Rubisco activation and stomatal opening within 7 to 10 min[7].

      In addition to these enzymatic mechanisms, recent analyses performed using a theoretical photosynthetic model have predicted that maintaining the appropriate concentrations of CBB cycle intermediates is crucial for the initiation of CO_2 fixation[8]. In fact, experimental evidence in cyanobacteria has confirmed that the transfer of the carbon supply from a storage polymer (glycogen) to the CBB cycle affects photosynthetic efficiency[9,10].

      However, details of the metabolic distribution and metabolic flux dynamics essential for efficient PI remain unknown. This shortcoming likely reflects the fact that the metabolic change during PI is poorly understood. Although transient changes in metabolite levels have been reported during PI in algae, in the chloroplasts of spinach protoplasts, and in wheat and maize leaves[11-15], we have yet to obtain systematic understanding of metabolite dynamics due to the lack of comprehensive and quantitative analyses. Instead, the metabolic dynamics and stability of the CBB cycle have been analyzed by computational modeling using random sampling of kinetic parameters instead of using known metabolome and flux distribution as a basis for model parameterization, given the lack of these analyses[16].

      Among oxygenic photosynthetic organisms, cyanobacteria are the only prokaryote; notably, in this class of organisms, most metabolic reactions are performed in a shared space, the cytoplasm. Thus, it is not obvious that the metabolic dynamics of PI in cyanobacteria is similar to that in leaves and chloroplasts. On the other hand, cyanobacteria fix CO_2 via the CBB cycle like that used in higher plants, and these prokaryotes are regarded as the likely ancestor of the plant chloroplast[17]. Despite its importance for further understanding PI processes across photosynthetic organisms, cyanobacterial metabolic dynamics during PI has not (to our knowledge) been studied to date.

The ideal way to systematize complicated metabolic changes during PI is to estimate

metabolic flux changes, which represent the integrated network responses of enzymes and

metabolites. Analysis of metabolic fluxes provides useful information regarding topics such as

pathway bottlenecks and network regulation in biological systems[18]. Although metabolic flux

analysis (MFA) using an isotope tracer enables the mapping of intracellular carbon fluxes[19], the

approach cannot be applied to a metabolically nonstationary state where metabolic fluxes and pool

sizes are variable. On the other hand, previously developed dynamic MFA, which incorporates

specialized statistical approaches, has permitted the estimation of long-term flux changes over the

culture interval[20,21]. There has been (to our knowledge) no appropriate approach for the quantitative

determination of metabolic flux change in short, metabolically transient states, including that of PI.

To estimate nonstationary metabolic flux changes, the following relationship was

employed. The accumulation rate of a metabolite in a metabolic network is defined as the sum of all

fluxes producing that metabolite minus the sum of all fluxes consuming that metabolite, as

represented in following equation:

92
$$r_i(\text{t}) = \sum_j \alpha_j x_j(t) - \sum_k \alpha_k x_k(t) \tag{1}$$

where $x_j(t)$ is the flux through reaction j, α_j is a stoichiometric coefficient, and $r_i(\text{t})$ is the

accumulation rate of metabolite i [22]. An example of three fluxes is illustrated in Fig. 1a. To obtain

accumulation rates for estimation of the flux values, it is necessary to examine the absolute

concentration change of metabolites with high time resolution. Namely, the metabolic flux during PI

can be calculated by quantifying the change in metabolite concentration over highly time-resolved

intervals immediate following light irradiation.

The time-resolved absolute quantification during PI in cyanobacteria has not been

performed so far due to the technical difficulties of absolute quantification and rapid sampling. Here,

by developing the integrated process shown in Fig. 1b, we successfully quantified absolute

metabolite concentration change at second-scale intervals during PI in the cyanobacterium

Synechocystis sp. PCC 6803 (*Synechocystis* hereafter). Additionally, to identify active metabolic

pathways during PI, [13]C labelling dynamics after light irradiation start was analyzed (Fig. 1c). These

accurate concentration and labelling dynamics allowed us to conduct short-time dynamic metabolic

flux analysis (STD-MFA) and to estimate nonstationary flux change during PI. These analyses

demonstrated a stepwise activation behavior that supports the smooth start of photosynthetic

metabolism in *Synechocystis*.

Results

Absolute concentration changes in central carbon metabolic pathways during PI. Firstly, we

sought to determine the absolute concentration dynamics of intermediate metabolites in central

energy metabolism during PI in *Synechocystis*. As light irradiation rapidly causes metabolic variation
driven by increases in ATP and NADPH via the photosynthetic electron transfer reaction, rapid
sampling and quenching of cellular metabolic activity are necessary for detecting metabolic changes
during PI. A standardized procedure for intracellular metabolome quantification requires multiple
analytical steps, including quenching of microbial metabolic activity, separation of microbial cells
from growth medium, and extraction of intracellular metabolites, followed by mass spectrometric
metabolite analysis. These multistep processes largely impede accurate estimation of the '*in vivo*'
cellular metabolic state. Typically, metabolic activities are stopped rapidly by adding a quenching
solution such as a cold methanol solution, followed by harvesting cells. However, metabolite leakage
in the quenching solution is known to interfere with precise quantification of metabolite levels[23].
Although quick filtration usually is used for rapid cell sampling before quenching and extraction
processes, the '*in vivo*' metabolic state may change a great deal due to variation in the light
environment during the filtration process in the case of photosynthetic microorganisms. Furthermore,
extraction yields of water-soluble metabolites depend on the extraction solvent, given that such
metabolites may decompose during extraction processes[24]. The exact determination of metabolite
concentrations also is hampered by ion suppression effects during mass spectrometric analysis[25].

To overcome these problems, in the present study, we employed a combination of two
strategies. Firstly, to analyze intracellular metabolites, cells were separated from growth medium and
suspended in 1 mM $NaHCO_3$ prior to metabolite sampling. This allowed quenching and extraction of
intracellular metabolites merely by directly dropping the cell suspension into the extraction solution
during light irradiation, and avoided metabolic state variation during the sampling process. We chose
phenol-chloroform-isoamylalcohol (PCI) solution (pH 7.9) as an extraction solution because PCI
extraction enabled rapid and efficient extraction of NADPH from *Synechocystis*[26]. Note that the
exchange of medium for $NaHCO_3$ solution is a typical pretreatment for measuring photosynthetic
activity of fresh-water cyanobacteria[27]. Secondly, stable isotopomers of the analyte compounds were
used as an internal standard (IS). Although experimental protocols using the isotope-labeled IS have
been developed for absolute quantification of metabolite concentration in various microorganisms
including cyanobacteria[28-31], few attempts (to our knowledge) have been made to apply this accurate
quantification method to time course analysis.

Figure 1b shows the experimental process for the exact determination of the absolute
metabolite concentration dynamics of PI. We prepared a ^{13}C-labeled IS mixture by extracting
metabolites from *Synechocystis* cells cultivated with $NaH^{13}CO_3$ as the sole carbon source. Absolute
concentrations of uniformly ^{13}C-labeled metabolites were determined using capillary
electrophoresis-mass spectrometry (CE-MS) based on the ratio between peak areas of known
concentrations of unlabeled compounds included as an IS and the uniformly ^{13}C-labeled metabolites.
Then, to quantify absolute concentration change during PI, dark-adapted *Synechocystis* cells (grown

in darkness for 12 h) were illuminated at a light intensity of 30 μmol m^{-2} s^{-1} (the same as the growth

light intensity) or 200 μmol m^{-2} s^{-1}. Metabolites of these cells were extracted rapidly using PCI

solvent containing the ^{13}C-labeled extract as a labeled IS whose concentration was determined using

another aliquot in the same extraction batch. Importantly, because of the high physicochemical

similarities between the labeled IS and the unlabeled analyte, the two mixtures undergo similar

opportunities for degradation and ion suppression effects, permitting compensation for the above

drawbacks[28-31]. Given that the absolute metabolite concentrations were determined based on the ratio

between unlabeled and uniformly labeled peak areas (Fig. 1b), the process was expected to prevent

contamination of unlabeled metabolites in the ^{13}C-labeled IS. The proportions of uniformly

^{13}C-labeled metabolites (compared to sum of unlabeled and uniformly labeled metabolites) were

found to exceed 95% for all metabolites except for citrate (Supplementary Fig. 1). Figure 2 shows

the temporal change of the absolute metabolite concentration after the initiation of light irradiation at

30 μmol m^{-2} s^{-1} or 200 μmol m^{-2} s^{-1}. It should be noted that a small amount of some metabolites

was extracellular (Supplementary Fig. 2). A previous study that measured NADPH fluorescence *in*

vivo revealed that the amount of NADPH in *Synechocystis* increased within 0.5 s of illumination[32].

The rapid increase of NADPH observed within 5 s in the present work was consistent with that

earlier *in vivo* study, confirming that this metabolomics analysis successfully detected the rapid start

of photosynthesis. As with NADPH, the level of the energy cofactor ATP also was observed to

increase rapidly. Overall, the levels of 3-phosphoglycerate (3PG), 2-phosphoglycerate (2PG), and

phosphoenolpyruvate (PEP) decreased; the levels of other CBB cycle intermediates (excepting 3PG,

ribulose 5-phosphate (Ru5P) and xylulose 5-phosphate (Xu5P)) increased, while the levels of

tricarboxylic acid (TCA) cycle intermediates did not change appreciably. Considering these

metabolite variations together, we inferred that accumulated glycolysis intermediates including 3PG,

2PG, and PEP likely were converted to CBB cycle intermediates at the initial step of PI.

^{13}C labeling kinetics during PI. A direct way to confirm this hypothesis is to determine the

metabolic flux distribution at PI. However, the precise metabolic flux values cannot be determined

when the number of unknown flux parameters is larger than that of known metabolite concentration

changes in a metabolic model system used for flux calculation. To perform STD-MFA, we sought to

refine the metabolic model by excluding metabolic pathways that are inactive during PI. However,

the metabolite concentration data in Fig. 2 cannot identify the inactive metabolic pathways. Notably,

even if the metabolic concentration change is zero, metabolic flux is not necessarily zero, because

such an observation instead may indicate that metabolic inflow and outflow are balanced. In the case

where the concentration of a metabolite is stable and ^{13}C is not incorporated into the metabolite

during photosynthetic ^{13}CO$_2$ fixation, the metabolic fluxes around the metabolite are estimated to be

very small, and so can be neglected. Therefore, we next measured the dynamics of ^{13}C isotope

incorporation after the transition from dark to light conditions by starting photosynthesis while the

cells were suspended in a solution of $NaH^{13}CO_3$. Figure 3 shows time courses of transient changes in

the atomic fraction of labeled carbon. In contrast to the high ^{13}C fraction seen for CBB cycle

intermediates, 6-phosphogluconate (6PG) and TCA cycle intermediates (except for malate and

fumarate) were not labeled. In addition, the ^{13}C fraction of ADP-glucose, which is known to be a

precursor of glycogen[33], was low. When these results were considered together with those indicating

that the concentrations of these metabolites did not change appreciably during PI (Fig. 2), we

inferred that the metabolic fluxes in the TCA cycle, the oxidative pentose phosphate pathway

(OPPP), the Entner-Doudoroff pathway (EDP), and the glycogen synthesis pathway are extremely

low immediately after light irradiation.

Other than Rubisco, the primary metabolic enzymes that incorporate CO_2 are

phosphoenolpyruvate carboxylase (PepC) and malic enzyme (ME), which are directly or indirectly

connected to malate (Fig. 3). Since malate was not labelled until 60 s after the initiation of light

irradiation (at both of the tested light intensities), ^{13}C incorporation before 60 s was inferred to

reflect Rubisco activity. The amount of incorporated ^{13}C atom was calculated by multiplying the

measured ^{13}C fraction (shown in Fig. 3) by the absolute carbon concentration of each metabolite

(shown in Fig. 2); these analyses are presented in Extended Data Fig. 1 a and b. The total amount of

incorporated ^{13}C atom was derived by summing the number of ^{13}C atoms of all measured

metabolites (Extended Data Fig. 1c, d). Notably, ^{13}C incorporation did not start until 12 s after the

initiation of light irradiation. On the other hand, ^{13}C incorporation was observed at least 10 s after

the addition of $NaH^{13}CO_3$ to cells whose photosynthetic activity already had been activated by 5 min

of illumination (Extended Data Fig. 2). Together, these results suggested that the observed delay in

the start of ^{13}C incorporation (compared to the start of light irradiation) was not attributable to slow

uptake of $NaH^{13}CO_3$ by the cells. Instead, these data indicated that the fixation of CO_2 through

Rubisco remains inactive at the initial stage of photosynthetic metabolism (i.e., immediately

following the start of illumination). Although the delayed onset of CO_2 fixation is a well-known

phenomenon in plant leaves[1], this observation in the present work is (to our knowledge) the first

direct demonstration of a Rubisco activation effect in the *in vivo* metabolic dynamics in

cyanobacteria.

It is interesting to note that labeled carbon was incorporated into malate and fumarate after

120 s, particularly in cells illuminated at 200 μmol m^{-2} s^{-1}, at a time when other TCA cycle

intermediates remained unlabeled. This result suggested that malate and fumarate are generated by

the reductive TCA pathway, which is known as a fermentation pathway employed for NADH

oxidation[34]. Thus, the accumulation of NADH may drive the activity of this reductive TCA pathway

(Extended Data Fig. 3).

Short-term dynamic metabolic flux analysis (STD-MFA). Finally, we calculated temporal changes in the net metabolic fluxes based on the obtained data. As mentioned above, during the first 60 s of illumination, the net metabolic fluxes in pathways other than the CBB cycle and glycolysis until 60 s appeared to be very small. In fact, the concentrations of newly synthesized (labeled) amino acids before 60 s were hardly changed, even in cells illuminated at 200 μmol m^{-2} s^{-1}, although the labeled levels of some amino acids (serine, alanine, phenylalanine, tyrosine and ornithine) increased significantly after the first minute of illumination (Extended Data Fig. 4). Furthermore, photorespiratory intermediates such as glycolate, glyoxylate, and glycerate did not accumulate during the first 60 s of illumination (Extended Data Fig. 5). Therefore, we assessed the metabolic pathways shown in Extended Data Fig. 6a as part of the subsequent calculations, focusing on metabolic dynamics during the first 60 s following the initiation of illumination.

To solve equation (1) and conduct STD-MFA, accumulation rates for metabolites in the metabolic model shown in Extended Data Fig. 6a were derived from the slopes obtained between two consecutive concentration data points in Fig. 2. In addition to our use of equation (1), we calculated the CO_2 fixation rate as the increase in the rate of total ^{13}C incorporation (Extended Data Fig. 1). Specifically, the latter parameter was interpreted as the metabolic flux from RuBP to 3PG, given that the ^{13}C incorporation experiment mentioned above indicated that all carbon atoms fixed during the first 60 s of illumination were attributable to Rubisco activity. The net metabolic fluxes of each time interval were estimated by solving these equations simultaneously (Fig. 4, Extended Data Fig. 6). The estimated flux values were validated and judged to be reliable, given that almost all of the estimated metabolite accumulation rate values (which were derived from the estimated fluxes) yielded values that were within the mean ± SD range of the measured accumulated rate (Extended Data Fig. 7). For more details of the calculation process and equations, see the Methods section.

The time course of flux distribution clearly indicated that photosynthetic metabolism in *Synechocystis* starts in a stepwise manner: first, activation of the reduction of 3PG to glyceraldehyde 3-phosphate (GAP); second, generation of RuBP; and third, Rubisco activation (Fig. 4, Extended Data Fig. 6c, d). Interestingly, the induction of RuBP generation can be further divided into two sub-steps based on metabolic fluxes, given that the formation fluxes of C6 intermediates (fructose 1,6-bisphosphate (FBP) and fructose 6-phosphate (F6P)) increased faster than that of C7 intermediates (sedoheptulose 1,7-bisphoshate (SBP) and sedoheptulose 7-phosphate (S7P)) (Extended Data Fig. 6c,d). Notably, the C6 and C7 pathways share an enzyme, fructose-1,6-bisphosphatase, in common, which (in *Synechocystis*) is known to be a bi-functional fructose-1,6-/sedoheptulose-1,7-bisphosphatase (FBP/SBPase)[35]. Thus, modification of enzymatic activity is unlikely to be the cause of the activation time gap between the two pathways. In the CBB cycle, the production of erythrose 4-phosphate (E4P), a starting substrate of the C7 pathway, depends on the activity of the C6 pathway. Therefore, we inferred that substrate deficiency in the C7

pathway is attributable to the metabolic pathway topology in the CBB cycle, delaying the C7

pathway relative to the C6 pathway in *Synechocystis*.

The reduction fluxes of 3PG differed greatly between cells illuminated at 30 and 200 μmol

m^{-2} s^{-1}, even before the initiation of CO_2 fixation (i.e., during the first 5-20 s) (Fig. 4). In

steady-state photosynthesis, the increased rate of electron donor (NADPH) production from the

photosynthetic electron transfer reaction at the higher light intensities becomes balanced with the

rate of electron acceptor (3PG) production that also is increased by elevated CO_2 fixation rate. On

the other hand, before the onset of CO_2 fixation immediately following the initiation of light

irradiation, excess reducing power can be accumulated because 3PG cannot be produced while

NADPH is produced by the electron transfer reaction. Nevertheless, the increase in the reducing

fluxes of 3PG with the higher light intensity (200 μmol m^{-2} s^{-1}) before CO_2 fixation suggests that

even before the onset of CO_2 fixation, the rate of 3PG consumption is balanced with the rate of

NADPH production rate from the electron transfer reaction, which varies with light intensities. In

addition, it clearly can be seen that 2PG and PEP were converted to 3PG at the both light intensities

(Fig. 4). Therefore, we proposed that accumulation of these glycolytic intermediates provides a

"stand-by" state and flexibility for the initiation of photosynthetic metabolism before the start of CO_2

fixation; thus, these intermediates are expected to play an important role in balancing cofactor

regeneration and consumption during PI.

Oxygen evolution in the presence of a CBB cycle inhibitor. To confirm the function of 3PG as an

electron acceptor at the start of photosynthesis *in vivo*, we measured the oxygen evolution rate in the

presence of a CBB cycle inhibitor, glycolaldehyde (GA). GA is known to inhibit the conversion of

triose phosphate to RuBP in the CBB cycle, thereby impeding oxygen evolution during steady-state

photosynthesis in cyanobacteria[36]. If 3PG is serving as an electron acceptor for NADPH generated

by the photosynthetic electron transfer reaction, oxygen evolution in light-exposed, GA-treated cells

should proceed only until the accumulated 3PG is exhausted. As expected, GA-treated cells

generated oxygen starting just after the initiation of illumination, but oxygen evolution by these cells

diminished rapidly after 25 s of irradiation with 200 μmol m^{-2} s^{-1} light (Fig. 5a). Based on

theoretical stoichiometry, two molecules of NADPH produced by one molecule of oxygen evolution

reduce two molecules of 3PG. The total decreasing rate of 3PG, 2PG and PEP immediately after 200

μmol m^{-2} s^{-1} light irradiation was about twice that of oxygen evolution rate in GA-treated cells (Fig.

5a, c). This quantitatively supports that the accumulated glycolytic metabolites acted as electron

acceptors for oxygen evolution in GA-treated cells. Moreover, the oxygen evolution rate of

GA-treated cells from the onset of the 200 μmol m^{-2} s^{-1} illumination to 25 s was identical to the

oxygen evolution rate of non-treated cells (Fig. 5a). This suggests that even in a normal PI process of

non-GA-treated cells, the electron acceptors that contributes to the initial oxygen evolution in PI are

the accumulated glycolytic intermediates. In cells irradiated with 30 μmol m^{-2} s^{-1} light irradiation,

no significant difference in oxygen evolution rate dynamics was observed between GA-treated and

non-treated cells; this effect presumably reflected slow consumption and depletion of 3PG in

GA-treated cells (Fig. 5b, d). In further confirmation of our hypothesis, we demonstrated depletion

of 3PG and inhibition of RuBP accumulation in GA-treated cells following irradiation with 200

μmol m^{-2} s^{-1} light (Extended Data Fig. 8). These results clearly indicated that accumulated 3PG

functions as a first electron acceptor for the rapid initiation of photosynthetic metabolism upon

exposure of dark-adapted cyanobacteria to light.

Discussion

In the present study, combined analysis of absolute metabolite concentrations and isotope

incorporation transients of a wide spectrum of metabolites permitted us to visualize changes in

nonstationary metabolic fluxes during PI in *Synechosystis*. The activation order during PI in

Synechosystis was found to be resemble that in land plants, where the activation of RuBP generation

typically precedes the start of CO_2 fixation[7]. However, the detailed order of activation of the various

pathways in plant leaves has not (to our knowledge) been investigated. STD-MFA revealed that the

step of activation of RuBP generation in *Synechosystis* involves two phases, the fast C6 pathway and

the slower C7 pathway. This distinction in timing was inferred to reflect the metabolic pathway

topology inherent to the CBB cycle. Furthermore, 3PG accumulated in the dark period was found to

be depleted rapidly, apparently functioning as a starting substrate for RuBP generation before the

start of CO_2 fixation. This rapid depletion of phosphoglycerate upon illumination also has been

observed in algae, in the chloroplast of spinach protoplasts, and in the leaves of a C3 plant

(wheat)[11,12,15]. In contrast, monotonic accumulation of phosphoglycerate has been reported in the

leaves of a C4 plant (*Zea mays*)[13,14]. We suspect that the difference in phosphoglycerate dynamics

during PI may reflect divergence between C3 and C4 photosynthetic metabolism, but clearly further

investigation is necessary.

A closer look at Fig. 2 permits a more detailed characterization of the details of the

individual metabolic pathways. Although increases in the levels of NADPH and ATP (which occur

within 5 s of the start of illumination) should make the reduction of 3PG to GAP energetically

favorable, a large decrease in the level of 3PG is observed after the 5-s time point. Cyanobacterial

phosphoglycerate kinase (PGK) and GAPDH are activated by reductive dissociation of cysteine

pairs upon light illumination[37,38]. Thus, the delay in 3PG consumption until after 5 s of illumination

likely is due to the need for the redox activation of PGK and GAPDH. Transient accumulation of

FBP indicates the existence of an activation period for the FBP/SBPase enzyme. FBP/SBPase in

Synechocystis is activated by reduction of a thiol group and binding of Mg^{2+}, and this enzyme is

inhibited allosterically by the binding of AMP[39,40]. Indeed, a rapid decrease of AMP concentration

was confirmed after light irradiation (Extended Data Fig. 3). In addition, temporal accumulation of
dihydroxyacetonephosphate (DHAP) upon irradiation with 200 μmol m^{-2} s^{-1} light may be
attributable to the transient rate-limitation of FBP/SBPase. The significantly higher continuous
accumulation of F6P and S7P in cells illuminated at 200 μmol m^{-2} s^{-1} (compared to those
illuminated at 30 μmol m^{-2} s^{-1}) indicates that the reaction catalyzed by the transketolase enzyme is
the major rate-limiting step. This interpretation is consistent with the results of a previous study,
which reported enhanced photosynthetic activity in a *Synechocystis* strain engineered to overproduce
transketolase[41]. Regarding RuBP dynamics, *Synechocystis* (like plants) requires Rubisco activation
processes, although the mechanism of this activation differs slightly between these two classes of
organisms[42]. Thus, the accumulation and subsequent decrease of RuBP in cyanobacteria suggests a
phase shift from inactivated to activated Rubisco. Indeed, similar dynamics of the RuBP pool size
was observed during PI in plant leaves, and is associated with Rubisco activation[43,44]. The levels of
GAP, E4P, and ribose 5-phosphate (R5P) were below the detection limit, suggesting that these
metabolites are consumed rapidly and are not accumulated during PI. The levels of 2PG and PEP
decreased more slowly than those of 3PG. Phosphoglycerate mutase catalyzes the reversible
conversion between 3PG and 2PG, while enolase catalyzes the reversible conversion between 2PG
and PEP. To assess the direction of these processes, the reaction Gibbs energy change ($\Delta_r G$) of these
reactions was calculated based on following equation:

$$\Delta_r G = \Delta_r G° + RT \ln Q \qquad (2)$$

$$Q_{3PG \to 2PG} = \frac{[2PG]}{[3PG]}, Q_{2PG \to PEP} = \frac{[PEP]}{[2PG]}$$

where $\Delta_r G°$ is the standard reaction Gibbs energy change, R is the ideal gas constant, T is the
temperature, and Q is the reaction quotient. As the conversions among 3PG, 2PG, and PEP are
simple reactions, where one reactant species is converted to one product (along with water),
consideration of cell volume is not necessary, allowing relatively precise estimation of $\Delta_r G$.
Therefore, we focused on the $\Delta_r G$ estimation of these reactions. Corresponding $\Delta_r G°$ values were
obtained as the means of the experimentally determined values summarized in Bennett et al.[29]. The
changes in reaction Gibbs energy from 3PG to 2PG and from 2PG to PEP during the first 60 s of
light irradiation were positive values or approximately 0 (Extended Data Fig. 9). These analyses are
consistent with our inference that the direction of net metabolic flux is from PEP to 3PG
immediately following the initiation of light irradiation, as shown in Fig. 4.

Let us consider the mechanisms by which large amounts of 3PG are accumulated in
dark-adapted *Synechocystis*. Cyanobacteria utilize glycogen stored during the day for respiratory
metabolism at night. In mutant strains deficient in the synthesis or degradation of glycogen, the
activation of photosynthetic metabolism is delayed due to lack of CBB cycle intermediates[9,10]. On
the other hand, the enzymatic activities of the TCA cycle in *Synechocystis* are known to be lower

than those in other organisms[45]. Thus, we infer that the generation of glycolytic intermediates by
glycogen degradation and slow catabolic activity via TCA cycle are likely to cause the accumulation
of 3PG along with 2PG and PEP in dark-adapted *Synechocystis*.

The STD-MFA technique developed in this study is a method for quantifying changes over
time of metabolic flux during a dynamic phenomenon that induces rapid metabolic changes that
occur over short intervals (within a few minutes). This method requires measurement of
time-resolved absolute concentration changes and metabolic model refinement based on isotope
incorporation data before flux calculation. Previously developed dynamic metabolic flux analysis
methods typically are intended to estimate flux change over long periods of culture processes, and
require technically specialized mathematical approaches[20,21]. In contrast, STD-MFA permits the
direct determination of metabolic flux changes in exchange, albeit while focusing on short time
intervals. We expect that this method also may be of use for understanding other non-stationary rapid
metabolic changes that have been the recent focus of attention, such as the end of photosynthesis and
adaptation against oxidative stress[46,47].

We have demonstrated that the accumulation of glycolysis intermediates helps smooth
metabolic start of CBB cycle in *Synechocystis*. The rational metabolic engineering based on the
findings in this study may improve CO_2 fixation rate and growth of synthetic autotrophic
microorganisms, which need further metabolic optimization[48,49]. This will lead to construction of
efficient bio-based chemical production from CO_2.

Methods

Growth conditions

This work employed a glucose-tolerant wild strain of *Synechocystis*[50] that was grown and stored on
solid BG-11 media plates (containing 1.5% Bacto agar). Pre-culturing was performed in 200-mL
flasks containing 50 mL each of liquid BG-11 medium buffered with 20 mM
2-[4-(2-hydroxyethyl)-1-piperazinyl]ethanesulfonic acid (HEPES, pH 7.5); each flask was
inoculated with cells from the agar plates. These flasks then were cultured at 30 ºC in ambient air
with shaking at 100 rpm; illumination at an intensity of 30 μmol $m^{-2}s^{-1}$ was provided by white
light-emitting diode (LED) lights. The main culture was initiated by adding pre-culture grown for 4
days into a 70-mL quantity of the BG-11 medium to give an optical density of 0.05 at 730 nm
(OD_{730}). The growth condition of the main culture was the same as that of the pre-culture. The dry
cell weight (DCW) was measured after harvesting cells by centrifugation and lyophilization. The
obtained relationship between OD_{730} value and DCW (DCW = 0.1998 × OD_{730} × V + 0.0974;
R^2=0.982; n=9; V is volume of cell suspension) was used for estimating DCW from the OD_{730} value.

Preparation of ^{13}C isotope-labeled internal standard (^{13}C-IS)

Synechocystis cells in pre-culture were inoculated to an OD_{730} of 0.05 in a serum bottle containing

60 mL modified BG-11 medium supplemented with 100 mM HEPES (pH 7.5) and 25 mM

$NaH^{13}CO_3$ (in place of the unlabeled Na_2CO_3 included in the standard BG-11 medium). The culture

serum bottle was sealed, followed by flushing with N_2 gas to remove unlabeled ambient CO_2. The

growth condition of the labeled culture was the same as that of the pre-culture and main culture.

Cells grown for 3 days were harvested and re-suspended in fresh modified BG-11 to avoid pH

increase during cultivation with $NaH^{13}CO_3$ as the sole carbon source. The labeled culture then was

grown under the same conditions for another 3 days, followed by extraction of uniformly labeled

metabolites as described below.

The entire volume of the labeled culture was harvested by centrifugation, washed with 1 mM

$NaH^{13}CO_3$ solution, and then re-suspended in 9 mL of 1 mM $NaH^{13}CO_3$. The resulting cell

suspension was illuminated for 90 s with white LED light at an intensity of 200 μmol m^{-2}s^{-1} (to

maximize extraction yield of CBB cycle intermediates), and dropped into a mixture of 9 mL PCI (pH

7.9, Nacalai tesque, Kyoto, Japan) and 60 μL 0.5 M N-cyclohexyl-3-aminopropanesulfonic acid

(CAPS, pH 10.6) for metabolite extraction. The extraction mixture was centrifuged at 4 ºC at 8,000

$\times$g for 3 min. Twelve to 13 fractions of the water layer were aliquoted into microtubes and

lyophilized.

One of the dried aliquots was dissolved in 20 μL of standard mixture containing prepared unlabeled

compounds. The sample was analyzed by capillary electrophoresis-mass spectrometry (CE-MS) (CE,

Agilent G7100; MS, Agilent G6224AA LC/MSD TOF; Agilent Technologies, Palo Alto, CA, USA)

as described previously[51]. Absolute concentrations of the uniformly labeled isotopomer (C_n) were

derived as follows:

422
$$C_n = \frac{S_n}{S_0} C_0 \tag{3}$$

where S_n and S_0 are peak areas of uniformly ^{13}C-labeled (namely, the number of ^{13}C atoms in a

n-carbon compound is equal to n) and unlabeled standard signals, respectively, and C_0 is the

prepared concentration of the unlabeled standard.

Metabolite extraction with rapid sampling and analysis

To perform metabolite sampling for analyses of absolute concentration and ^{13}C incorporation after

transition from dark to light, a main culture grown for 6 days was dark-adapted for 12 h. A calculated

volume (90/OD_{730}, in mL) of the dark-adapted cell culture was harvested by centrifugation at

8000×g for 5 min. The cells were washed with 4 mL of 1 mM $NaHCO_3$ (for absolute concentration

analysis) or 1 mM $NaH^{13}CO_3$ (for ^{13}C incorporation analysis), and harvested again by centrifugation.

The washed cells were resuspended in 6.8 mL of 1 mM $NaHCO_3$ (for absolute concentration

analysis) or 1 mM $NaH^{13}CO_3$ (for ^{13}C incorporation analysis). Nine aliquots (750 μL each) of the

resulting cell suspension were loaded into clear pipet tips (Supplementary Fig. 3). Up to this point,

light irradiation was avoided during the procedure to keep the cells in the dark-adapted state.

Irradiation of all the aliquots with white LED light at an intensity of 200 ± 10 μmol m^{-2}s^{-1} or 30 ± 2

μmol m^{-2}s^{-1} was initiated, followed by sampling of the cells at the indicated time points by dropping

into 750 μL PCI supplemented with ^{13}C-IS dissolved in 20 μL water (for absolute concentration

analysis) or with 5 μL of 0.5 M CAPS (pH 10.6) (for ^{13}C incorporation analysis). The cell

suspension before illumination was extracted as the nominal 0-s sample. To precisely determine the

amount of extracted cells, the OD$_{730}$ of the final cell suspension was measured. After the extraction

mixtures were centrifuged at 4 ºC at 12,000×g for 2 min, harvested aliquots of the water layer were

lyophilized.

The lyophilized samples were dissolved in 20 μL of 0.4 mM piperazine-1,4-bis(2-ethanesulfonic

acid) (PIPES) and analyzed by CE-MS as described above. Absolute concentrations of each

metabolite (*C'*) were derived as follows:

448
$$C' = \frac{S'_0}{WS'_n}C_nV \tag{4}$$

where *S'$_0$* and *S'$_n$* are peak areas of unlabeled and uniformly ^{13}C-labeled IS signals, respectively, *W*

represents DCW, and *V* is sample volume for CE-MS analysis. ^{13}C fractions were calculated as

described previously[34].

Short-term dynamic metabolic flux analysis (STD-MFA)

To estimate temporal changes of 18 fluxes in the metabolic map shown in Extended Data Fig. 6,

eighteen metabolite accumulation rates r_i(t) in equation (1) were derived from the slope taken

between two consecutive concentration data points shown in Fig. 2. Accumulation rates of

metabolites under the detection limit (E4P, R5P, GAP < 0.001 nmol/mg DCW) and peak-unidentified

metabolites (BPG and SBP) were set to zero. In addition, the CO$_2$ fixation rate derived from the

slope taken between two consecutive data points shown in Extended Data Fig. 1c and d was

regarded as the metabolic flux of Rubisco (i.e., the conversion of RuBP to 3PG). The set of 19

equations, including the 18 unknown fluxes in the network, was represented in matrix notation as

follows:

463
$$\mathbf{A}\mathbf{x}(t) = \mathbf{r}(t) \tag{5}$$

464   where A was the 19 $\times$ 18 matrix of stoichiometric coefficients, **x**(t) was the 18-dimensional flux

465   vector, and r(t) was the 19-dimensional metabolite accumulation rate vector. The specific form of

466   equation (4) is shown in Extended Data Fig. 6b. The weighted least squares solution to equation (5)

467   is as follows:

468
$$\hat{\mathbf{x}}(t) = (\mathbf{A}^T\mathbf{W}\mathbf{A})^{-1}\mathbf{A}^T\mathbf{W}\bar{\mathbf{r}}(t) \tag{6}$$

where the superscripts ^ and ⁻ denote estimated and measured quantities (respectively), and **W** is

a 19-dimensional weighting diagonal matrix, whose diagonal elements are the reciprocal of the

variance of the measured metabolic accumulation rate values. Variance values for the accumulation

rate of E4P, R5P, GAP, BPG, and SBP, and for the CO_2 fixation rate, were set to 10^{-15} (i.e., a value

that was effectively zero). To evaluate the reliability of the estimated fluxes, metabolite

accumulation rates were calculated as follows:

$$\mathbf{A}\hat{x}(t) = \hat{r}(t) \tag{7}$$

The estimated and measured metabolite accumulation rate values are compared in Extended Data

Fig. 7.

Oxygen evolution measurement

Synechocystis wild-type cells were collected and re-suspended in the same way as the preparation

process in the metabolite sampling experiment described above. Whole-cell oxygen evolution was

measured with light illumination at the intensity of 200 µmol $m^{-2}s^{-1}$ or 30 µmol $m^{-2}s^{-1}$ in a

Clark-type oxygen electrode (Hansatech Instruments) containing 750 µL of the cell suspension.

References

1.   Walker, D. A. Photosynthetic induction phenomena and the light activation of ribulose

diphosphate carboxylase. *New Phytol.* **72**, 209–235 (1973).

2.   Osterhout, W. J. & Haas, A. R. Dynamical Aspects of Photosynthesis. *Proc. Natl. Acad.*

Sci. USA. **4**, 85-91 (1918).

3.   Buchanan, B. B. Role of light in the regulation of chloroplast enzymes. *Ann. Rev. Plant*

Physiol. **31**, 341-374 (1980).

4.   Lorimer, G. H., Badger, M. R. & Andrews, T. J. The Activation of

Ribulose-1,5,-bisphosphate carboxylase by carbon dioxide and magnesium ions. Equilibria,

kinetics, a suggested mechanisms, and physiological implications. *Biochem.* **15**, 529-536

(1976).

5.   Portis, Jr. A. R. The regulation of Rubisco by Rubisco activase. *J. Exp. Bot.* **46**, 1285-1291

(1995).

6.   Michelet, L. et al. Redox regulation of the Calvin-Benson cycle: something old, something

new. *Front Plant Sci* **4**, 470 (2013).

7.   Pearcy, R. W., Krall, J. P. & Sassenrath-Cole, G. F. Photosynthesis in Fluctuating Light

Environments. In: Baker NR, ed. *Photosynthesis and the Environment.* Dordrecht: Kluwer,

321–346 (1996).

8.   Matuszyńska, A., Saadat, N. P. & Ebenhöh, O. Balancing energy supply during

photosynthesis - a theoretical perspective. *Physiol. Plant* **166**, 392-402 (2019).

9.   Shimakawa, G. et al. Respiration accumulates Calvin cycle intermediates for the rapid start

of photosynthesis in Synechocystis sp. PCC 6803. *Biosci. Biotechnol. Biochem.* **78**,
1997-2007 (2014).

10.   Shinde, S. et al. Glycogen Metabolism Supports Photosynthesis Start through the
Oxidative Pentose Phosphate Pathway in Cyanobacteria. *Plant Physiol.* **182**, 507-517
(2020).

11.   Bassham, J. A., Shibata, K., Steenberg, K., Bourdon, J. and Calvin, M. The photosynthetic
cycle and respiration: Light-dark transients. *J. Am. Chem. Soc.* **78**, 4120-4124 (1956).

12.   Stitt, M., Wirtz, W. & Heldt, H. W. Metabolite levels during induction in the chloroplast
and extrachloroplast compartments of spinach protoplasts. *Biochim. Biophys. Acta.* **593**,
85-102 (1980).

13.   Leegood, R. C. & Furbank, R. T. Carbon metabolism and gas exchange in leaves of Zea
mays L.: Changes in CO2 fixation, chlorophyll a fluorescence and metabolite levels during
photosynthetic induction. *Planta.* **162**, 450-456 (1984).

14.   Usuda, H. Changes in Levels of Intermediates of the C(4) Cycle and Reductive Pentose
Phosphate Pathway during Induction of Photosynthesis in Maize Leaves. *Plant Physiol.* **78**,
859-864 (1985).

15.   Kobza, J. & Edwards, G. E. The photosynthetic induction response in wheat leaves: net
CO2 uptake, enzyme activation, and leaf metabolites. *Planta* **171**, 549-559 (1987).

16.   Janasch, M., Asplund-Samuelsson, J., Steuer, R. & Hudson, E. P. Kinetic modeling of the
Calvin cycle identifies flux control and stable metabolomes in *Synechocystis* carbon
fixation. *J. Exp. Bot.* **70**, 973-983 (2019).

17.   Ponce-Toledo, R. I. et al. An Early-Branching Freshwater Cyanobacterium at the Origin of
Plastids. *Curr. Biol.* **27**, 386-391 (2017).

18.   Shimizu, H. & Toya, Y. Recent advances in metabolic engineering-integration of in silico
design and experimental analysis of metabolic pathways. *J. Biosci. Bioeng.* **132**, 429-436
(2021).

19.   Sauer, U. Metabolic networks in motion: 13C-based flux analysis. *Mol. Syst. Biol.* **2**, 62
(2006).

20.   Leighty, R. W. & Antoniewicz, M. R. Dynamic metabolic flux analysis (DMFA): a
framework for determining fluxes at metabolic non-steady state. *Metab. Eng.* **13**, 745-755
(2011).

21.   Niklas, J., Schräder, E., Sandig, V., Noll, T. & Heinzle, E. Quantitative characterization of
metabolism and metabolic shifts during growth of the new human cell line AGE1.HN
using time resolved metabolic flux analysis. *Bioprocess Biosyst. Eng.* **34**, 533-545 (2011).

22.   Vallino, J. J. & Stephanopoulos, G. Metabolic flux distributions in *Corynebacterium*
glutamicum during growth and lysine overproduction. *Biotechnol. Bioeng.* **41**, 633-646

(1993).

23.   Pinu, F. R., Villas-Boas, S. G. & Aggio, R. Analysis of Intracellular Metabolites from Microorganisms: Quenching and Extraction Protocols. *Metabolites.* **7**, E53 (2017).

24.   Rabinowitz, J. D. & Kimball, E. Acidic acetonitrile for cellular metabolome extraction from Escherichia coli. *Anal. Chem.* **79**, 6167-6173 (2007).

25.   Annesley, T. M. Ion suppression in mass spectrometry. *Clin. Chem.* **49**, 1041-1044 (2003).

26.   Tanaka, K. et al. Quantification of NAD(P)H in cyanobacterial cells by a phenol extraction method. *Photosynth. Res.* **148**, 57-66 (2021).

27.   Shimakawa, G. et al. FLAVODIIRON2 and FLAVODIIRON4 proteins mediate an oxygen-dependent alternative electron flow in *Synechocystis* sp. PCC 6803 under CO_2-limited conditions. *Plant. Physiol.* **167**, 472-480 (2015).

28.   Wu, L. et al. Quantitative analysis of the microbial metabolome by isotope dilution mass spectrometry using uniformly 13C-labeled cell extracts as internal standards. *Anal. Biochem.* **336**, 164-171 (2005).

29.   Bennett, B. D. et al. Absolute metabolite concentrations and implied enzyme active site occupancy in *Escherichia coli*. *Nat. Chem. Biol.* **5**, 593-599 (2009).

30.   Dempo, Y., Ohta, E., Nakayama, Y., Bamba, T. & Fukusaki, E. Molar-based targeted metabolic profiling of cyanobacterial strains with potential for biological production. *Metabolites.* **4**, 499-516 (2014).

31.   Nishino, S., Okahashi, N., Matsuda, F. & Shimizu, H. Absolute quantitation of glycolytic intermediates reveals thermodynamic shifts in *Saccharomyces cerevisiae* strains lacking *PFK1* or *ZWF1* genes. *J. Biosci. Bioeng.* **120**, 280-286 (2015).

32.   Kauny, J. & Sétif, P. NADPH fluorescence in the cyanobacterium *Synechocystis* sp. PCC 6803: a versatile probe for in vivo measurements of rates, yields and pools. *Biochim. Biophys. Acta.* **1837**, 792-801 (2014).

33.   Yoo, S. H., Lee, B. H., Moon, Y., Spalding, M. H. & Jane, J. L. Glycogen synthase isoforms in *Synechocystis* sp. PCC6803: identification of different roles to produce glycogen by targeted mutagenesis. *PLoS One.* **9**, e91524 (2014).

34.   Hasunuma, T., Matsuda, M., Kato, Y., Vavricka, C. J. & Kondo, A. Temperature enhanced succinate production concurrent with increased central metabolism turnover in the cyanobacterium *Synechocystis* sp. PCC 6803. *Metab. Eng.* **48**, 109-120 (2018).

35.   Tamoi, M., Takeda, T. & Shigeoka, S. Functional analysis of Fructose-1,6-bisphosphatase isozymes (*fbp-I* and *fbp-II* gene products) in cyanobacteria. *Plant Cell Physiol.* **40**, 257-261 (1999).

36.   Miller, A. G. & Canvin, D. T. Glycolaldehyde Inhibits CO_2 Fixation in the Cyanobacterium *Synechococcus* UTEX 625 without Inhibiting the Accumulation of

Inorganic Carbon or the Associated Quenching of Chlorophyll *a* Fluorescence. *Plant.*

Physiol. **91**, 1044-1049 (1989).

37.   Tsukamoto, Y., Fukushima, Y., Hara, S. & Hisabori, T. Redox control of the activity of

phosphoglycerate kinase in *Synechocystis* sp. PCC6803. *Plant Cell. Physiol.* **54**, 484-491

(2013).

38.   McFarlane, C. R. et al. Structural basis of light-induced redox regulation in the

Calvin-Benson cycle in cyanobacteria. *Proc. Natl. Acad. Sci. USA.* **116**, 20984-20990

(2019).

39.   Feng, L. et al. Structural and biochemical characterization of

fructose-1,6/sedoheptulose-1,7-bisphosphatase from the cyanobacterium *Synechocystis*

strain 6803. *FEBS J.* **281**, 916-926 (2014).

40.   Mallén-Ponce, M. J., Huertas, M. J., Sánchez-Riego, A. M. & Florencio, F. J. Depletion of

m-type thioredoxin impairs photosynthesis, carbon fixation, and oxidative stress in

cyanobacteria. *Plant Physiol.* **187**, 1325-1340 (2021).

41.   Liang, F. & Lindblad, P. Effects of overexpressing photosynthetic carbon flux control

enzymes in the cyanobacterium *Synechocystis* PCC 6803. *Metab. Eng.* **38**, 56-64 (2016).

42.   Marcus, Y. & Gurevitz, M. Activation of cyanobacterial RuBP-carboxylase/oxygenase is

facilitated by inorganic phosphate via two independent mechanisms. *Eur. J. Biochem.* **267**,

5995-6003 (2000).

43.   Perchorowicz, J. T., Raynes, D. A. & Jensen, R. G. Light limitation of photosynthesis and

activation of ribulose bisphosphate carboxylase in wheat seedlings. *Proc. Natl. Acad. Sci.*

USA. **78**, 2985-2989 (1981).

44.   Seemann, J. R., Kirschbaum, M. U., Sharkey, T. D. & Pearcy, R. W. Regulation of

Ribulose-1,5-Bisphosphate Carboxylase Activity in *Alocasia macrorrhiza* in Response to

Step Changes in Irradiance. *Plant Physiol.* **88**, 148-152 (1988).

45.   Ito, S. & Osanai, T. Unconventional biochemical regulation of the oxidative pentose

phosphate pathway in the model cyanobacterium *Synechocystis* sp. PCC 6803. *Biochem.*

J. **477**, 1309-1321 (2020).

46.   Christodoulou, D. et al. Reserve Flux Capacity in the Pentose Phosphate Pathway Enables

Escherichia coli's Rapid Response to Oxidative Stress. *Cell Syst.* **6**, 569-578.e7 (2018).

47.   Maruyama, M. et al. Time-resolved analysis of short term metabolic adaptation at dark

transition in *Synechocystis* sp. PCC 6803. *J. Biosci. Bioeng.* **128**, 424-428 (2019).

48.   Gleizer, S. et al. Conversion of Escherichia coli to Generate All Biomass Carbon from

CO2. *Cell.* **179**, 1255-1263.e12 (2019).

49.   Gassler, T. et al. The industrial yeast Pichia pastoris is converted from a heterotroph into

an autotroph capable of growth on CO2. *Nat. Biotechnol.* **38**, 210-216 (2020).

50.   Williams, J. G. K. Construction of specific mutations in photosystem II photosynthetic
615   reaction center by genetic engineering methods in *Synechocystis* 6803. *Methods Enzymol.*
616   **167**, 766-778 (1988).
617   51.   Hasunuma, T. et al. Dynamic metabolic profiling of cyanobacterial glycogen biosynthesis
618   under conditions of nitrate depletion. *J. Exp. Bot.* **64**, 2943-2954 (2013).
619

620   **Acknowledgements**
621   This work was supported by the Mirai Program [grant number JPMJMI19E4] from the Japan
622   Science and Technology Agency (JST) of the Ministry of Education, Culture, Sports, Science, and
623   Technology (MEXT), Japan, and the Japan Society for the Promotion of Science (JSPS) Kakenhi
624   (Grants-in-Aid for Scientific Research) Grant Number 21J00113.
625

626   **Author contributions**
627   K.T. and T.H. designed the research. K.T. and M.M. performed research. K.T. and T.S. analyzed data.
628   K.T., T.S., A.K. and T.H. wrote the article.
629

630   **Competing interests**
631   The authors declare no competing interests.
632

 Figures

Fig. 1 Process for estimating temporal changes in metabolic flux. a, Relationship between concentration change and metabolic flux. $r_i(t)$ is the accumulation rate of metabolite M_i. x_n represents metabolic flux of the nth reaction. **b,** Experimental procedure for quantifying absolute metabolite concentration. **c,** Schematic example of isotope-labeled state during $^{13}CO_2$ incorporation. Empty and red-filled circles represent unlabeled and labeled carbon atoms, respectively. Abbreviations: CBB, Calvin-Benson-Bassham; CEMS, capillary electrophoresis-mass spectrometry; F6P, fructose 6-phosphate; IS, internal standard; RuBP, ribulose 1,5-bisphosphate; TCA, tricarboxylic acid; 2OG, 2-oxoglutarate; 3PG, 3-phosphoglycerate.

Fig. 2 Rapid temporal change of absolute metabolite concentration during photosynthesis start.
Intracellular metabolites were extracted at the indicated times after transition from darkness to
light-irradiated condition at the intensity of 200 μmol m^{-2} s^{-1} (red line) or 30 μmol m^{-2} s^{-1} (blue
line). Levels of glyceraldehyde 3-phosphate (GAP), erythrose 4-phosphate (E4P), and ribose
5-phosphate (R5P) were below the detection limit (0.001 nmol / mg DCW).
1,3-Bisphosphoglycerate (BPG) and sedoheptulose 1,7-bisphosphate (SBP) were not identified due
to lack of commercial reagent for use as a standard. Black arrows indicate metabolic pathways.
Abbreviations: DCW, dry cell weight; DHAP, dihydroxyacetonephosphate; FBP, fructose
1,6-bisphosphate; F6P, fructose 6-phosphate; G1P, glucose 1-phosphate; G6P, glucose 6-phosphate;

654   2OG, 2-oxoglutarate; PEP, phosphoenolpyruvate; 6PG, 6-phosphogluconate; 2PG,

655   2-phosphoglycerate; 3PG, 3-phosphoglycerate; RuBP, ribulose 1,5-bisphosphate; Ru5P, ribulose

656   5-phosphate; S7P, sedoheptulose 7-phosphate; Xu5P, Xylulose 5-phosphate. Values are mean ± SD

657   (bars) of three independent experiments. Significant differences between the two light intensities

658   were evaluated by a two-tailed non-paired Student's t test (* $p < 0.05$).

Fig. 3 Time course of ^{13}C fraction of metabolites following light irradiation at 200 µmol m^{-2} s^{-1} (red line) or 30 µmol m^{-2} s^{-1} (blue line). Values are mean ± SD (bars) of three biological replicates. Black arrows indicate metabolic pathways. Abbreviations are as in Figure 2 legend.

Fig. 4 Estimated change in metabolic flux after initiating light irradiation at 200 µmol m⁻² s⁻¹

(upper row) and 30 µmol m⁻² s⁻¹ (lower row). Metabolic fluxes are indicated as thickness of

arrows with the values in the scale bar in the lower right corner. Metabolite concentrations at the

beginning and end of each time interval are indicated by diameters of blue and orange circles,

respectively. Overlap between orange and blue circles (namely invariant concentration during a time

interval) is shown in gray. Abbreviations are as in Figure 2 legend.

674
Fig. 5 Oxygen evolution rate in the presence of glycolaldehyde (GA) and comparison with
metabolite variation rate. a, b, Oxygen evolution rate in the absence (black line) and presence (red
line) of GA after light irradiation at the intensity of 200 μmol m^{-2} s^{-1} (**a**) or 30 μmol m^{-2} s^{-1} (**b**). GA
(10 mM) was added before the initiation of light irradiation. **c, d,** Decreasing rate of 3PG (orange
line), PEP (yellow line), and 2PG (gray line), and the sum of these three metabolites (blue line) after
initiation of light irradiation at the intensity of 200 μmol m^{-2} s^{-1} (**c**) or 30 μmol m^{-2} s^{-1} (**d**). Since the
decreasing rates were derived from two consecutive concentration data points, values are plotted at a
time point representing the average (mean) of the two time points. For all panels, values are plotted
as mean ± SD (bars) of three biological replicates. Abbreviations are as in Figure 2 legend.
684
685

Extended Data Fig. 1 Time course of ^{13}C level in each metabolite (a, b) and total ^{13}C level (c, d).

Number of ^{13}C atoms was calculated as mean of ^{13}C enrichment multiplied by mean metabolite

concentration and the number of C atoms in the molecule for the two light intensities: (a, c) 200

μmol m^{-2} s^{-1}, (b, d) 30 μmol m^{-2} s^{-1}. (c, d) Number of ^{13}C atoms was summed for CBB cycle and

oxidative pentose phosphate pathway (OPPP) intermediates (orange line), TCA cycle, and glycolysis

intermediates (gray line), and all measured metabolites (blue line). Abbreviations are as in Figure 2

legend.

Extended Data Fig. 2 Effect of pre-illumination on the start of ^{13}C incorporation. ^{13}C fractions
of (a) 3-phosphoglycerate (3PG) and (b) Ribulose 1,5-bisphosphate (RuBP) after 12 s of illumination
(200 μmol m^{-2} s^{-1}) of the 12-h dark-adapted cells were lower than those at 10 s after addition of 1
mM NaH^{13}CO$_3$ to cells that had been pre-illuminated (200 μmol m^{-2} s^{-1}) for 5 min. Presumably, the
higher ^{13}C incorporation rate in pre-illuminated cells reflects the fact that CO$_2$ fixation already had
been activated (by pre-illumination) prior to the addition of NaH^{13}CO$_3$. Thus, lower ^{13}C
incorporation by dark-adapted cells can be attributed to the low activation state of CO$_2$ fixation
during photosynthetic induction. Values are mean ± SD (bars) of three biological replicates.
Significant differences were evaluated by a two-tailed non-paired Student's t test; *p* values were as
indicated.

Extended Data Fig. 3 Concentration transients of cofactors during photosynthetic induction.
Intracellular metabolites were extracted at the indicated times after transition from darkness to light-irradiated conditions at the intensity of 200 μmol m^{-2} s^{-1} (red line) or 30 μmol m^{-2} s^{-1} (blue line). Values are mean ± SD (bars) of three independent experiments. Significant differences between the two light intensities were evaluated by a two-tailed non-paired Student's t test (* $p <$ 0.05). Abbreviations: ADP, Adenosine diphosphate; AMP, adenosine monophosphate; ATP, adenosine triphosphate; GDP, Guanosine diphosphate; GMP, Guanosine monophosphate; GTP, Guanosine triphosphate; NAD(P), nicotinamide adenine dinucleotide (phosphate)

Extended Data Fig. 4 Concentration and ^{13}C fraction transients of amino acids during
photosynthetic induction of cells illuminated at the intensity of 200 μmol m^{-2} s^{-1}. Intracellular
metabolites were extracted at the indicated times after transition from darkness to light-irradiated
conditions at the intensity of 200 μmol m^{-2} s^{-1}. Red lines and yellow lines indicate metabolite
concentration and ^{13}C fraction, respectively. Simplified central metabolic pathways are indicated by
arrows. Values of concentrations and ^{13}C fractions are mean ± SD (bars) of three and two
independent experiments, respectively. Values of metabolite concentrations for individual
experiments are indicated by gray lines. Abbreviations are as in Figure 2 legend.

Extended Data Fig. 5 Concentration transients of intermediates in photorespiration pathway during photosynthetic induction. Intracellular metabolites were extracted at the indicated time after transition from darkness to light-irradiated conditions at the intensity of 200 μmol m^{-2} s^{-1} (red line) or 30 μmol m^{-2} s^{-1} (blue line). Values are mean ± SD (bars) of three independent experiments. DCW, dry cell weight

Extended Data Fig. 6 Short-term dynamic metabolic flux analysis (STD-MFA). (a) Metabolic

pathways incorporated into calculation for STD-MFA. The experimental data of absolute

concentration and ^{13}C incorporation change allowed us to simplify the metabolic model during

photosynthetic induction. x_i represents net flux of ith metabolic pathway. (b) Detailed form of the

mass balance equation used for STD-MFA. For more details, see Methods section. (c, d) Transients

of estimated metabolic fluxes during photosynthetic induction in illuminated conditions at (c) 200

µmol m^{-2} s^{-1} or (d) 30 µmol m^{-2} s^{-1}. Since the fluxes were estimated using metabolite accumulation

rates derived from two consecutive concentration data points, flux values are plotted at a time point

corresponding to the average (mean) of the two time points. Abbreviations are as in Figure 2 legend.

Extended Data Fig. 7 Measured and estimated metabolic accumulation rates. Measured accumulation rates at the light intensity of 200 μmol m^{-2} s^{-1} (red solid line) or 30 μmol m^{-2} s^{-1} (blue solid line) were derived from slope between two consecutive concentration data points shown in Fig. 2. Estimated accumulation rates at the light intensity of 200 μmol m^{-2} s^{-1} (orange dotted line) or 30 μmol m^{-2} s^{-1} (blue dotted line) are shown for validation of the short-term dynamic metabolic flux analysis (STD-MFA) result. Black arrows indicate metabolic pathways. Abbreviations are as in

753   Figure 2 legend. Values are plotted at the average (mean) time point of the two time points used for
754   the concentration data points. Values of measured accumulation rates are mean $\pm$ SD (bars) of
755   three independent experiments. Asterisk indicates that estimated value is out of range of the
756   measured mean $\pm$ SD.
757

Extended Data Fig. 8 Effect of CBB cycle inhibitor glycolaldehyde (GA) on metabolite
concentrations. Absolute concentrations of (a) 3-phosphoglycerate (3PG) and (b) Ribulose
1,5-bisphosphate (RuBP) before light illumination (dark condition; "0 s") and at 60 s after light
illumination at 200 μmol m^{-2} s^{-1} were analyzed in the absence (-) and presence (GA) of 10 mM GA.
Values are mean ± SD (bars) of three independent experiments. Significant differences were
evaluated by a two-tailed non-paired Student's t test, and *p* values were as indicated.
766

768   **Extended Data Fig. 9 Time course of reaction Gibbs free energy change ($\Delta_r G$).** $\Delta_r G$ values of

769   reactions (a) from 3-phosphoglycerate (3PG) to 2-phosphoglycerate (2PG) and (b) from 2PG to

770   phosphoenolpyruvate (PEP) were derived based on equation (2). Red and blue lines indicate $\Delta_r G$ at

771   the illuminated light intensity of 200 μmol m^{-2} s^{-1} and 30 μmol m^{-2} s^{-1}, respectively. Values are

772   mean $\pm$ SD (bars) derived from three independent experiments.

773