Moxifloxacin-mediated killing of *Mycobacterium tuberculosis* involves respiratory downshift, reductive stress, and ROS accumulation

Somnath Shee[1, 2]¶, Samsher Singh[1, 2]¶, Ashutosh Tripathi[1, 2], Chandrani Thakur[3], Anand Kumar T[4], Mayashree Das[1, 2], Vikas Yadav[1, 2], Sakshi Kohli[1, 2], Raju S. Rajmani[2], Nagasuma Chandra[3], Harinath Chakrapani[4], Karl Drlica[5]*, Amit Singh[1, 2]*

[1]Department of Microbiology and Cell Biology, [2]Centre for Infectious Disease Research, Indian Institute of Science, Bangalore 560012, Karnataka, India

[3]Department of Biochemistry, Indian Institute of Science, Bangalore 560012, Karnataka, India

[4]Department of Chemistry, Indian Institute of Science Education and Research (IISER) Pune, Pune 411 008, Maharashtra, India

[5]Public Health Research Institute and Department of Microbiology, Biochemistry & Molecular Genetics, New Jersey Medical School, Rutgers Biomedical and Health Sciences, Rutgers University, Newark, NJ 07103

Running title: Redox-mechanisms of moxifloxacin lethality in *M. tb*

¶ Equal Contribution

*For correspondence: *asingh@iisc.ac.in (Amit Singh), drlicaka@njms.rutgers.edu (Karl Drlica)

22 **Keywords:** antimycobacterial, ROS, oxidative stress, moxifloxacin, fluoroquinolone,

23 respiration, N-acetyl cysteine, redox biosensor, NADH, reductive stress, resistance

Abstract

Moxifloxacin is central to treatment of multidrug-resistant tuberculosis. Effects of moxifloxacin on *Mycobacterium tuberculosis* redox state were explored to identify strategies for increasing lethality and reducing the prevalence of extensively resistant tuberculosis. A non-invasive redox biosensor and an ROS-sensitive dye revealed that moxifloxacin induces oxidative stress correlated with *M. tuberculosis* death. Moxifloxacin lethality was mitigated by supplementing bacterial cultures with an ROS scavenger (thiourea), an iron chelator (bipyridyl), and, after drug removal, an antioxidant enzyme (catalase). Lethality was also reduced by hypoxia and nutrient starvation. Moxifloxacin increased the expression of genes involved in the oxidative stress response, iron-sulfur cluster biogenesis, and DNA repair. Surprisingly, and in contrast with *Escherichia coli* studies, moxifloxacin decreased expression of genes involved in respiration, suppressed oxygen consumption, increased the NADH/NAD$^+$ ratio, and increased the labile iron pool in *M. tuberculosis*. Lowering the NADH/NAD$^+$ ratio in *M. tuberculosis* revealed that NADH-reductive stress facilitates an iron-mediated ROS surge and moxifloxacin lethality. Treatment with N-acetyl cysteine (NAC) accelerated respiration and ROS production, increased moxifloxacin lethality, and lowered the mutant prevention concentration. Moxifloxacin induced redox stress in *M. tuberculosis* inside macrophages, and co-treatment with NAC potentiated the anti-mycobacterial efficacy of moxifloxacin during nutrient starvation, inside macrophages, and in mice where NAC restricted the emergence of resistance. Thus, oxidative stress, generated in a novel way, contributes to moxifloxacin-mediated killing of *M. tuberculosis.* The results open a way to make fluoroquinolones more effective anti-tuberculosis agents and provide a mechanistic basis for NAC-mediated enhancement of fluoroquinolone lethality *in vitro* and *in vivo.*

Author Summary

A new paradigm was revealed for stress-mediated bacterial death in which moxifloxacin treatment of *M. tuberculosis* decreases respiration rate (respiration <u>increases</u> in *E. coli*). Although moxifloxacin-induced, ROS-mediated bacterial death was observed, it derived from elevated levels of NADH and iron, a phenomenon not seen with antibiotic-treated *E. coli*. Nevertheless, stimulation of respiration and ROS by N-acetyl cysteine (NAC) enhanced moxifloxacin-mediated killing of *M. tuberculosis*, thereby reinforcing involvement of ROS in killing. NAC stimulation of moxifloxacin-mediated killing of *M. tuberculosis* and restriction of the emergence of resistance in a murine model of infection emphasize the importance of lethal action against pathogens. The work, plus published benefits of NAC to TB patients, encourage studies of NAC-based enhancement of fluoroquinolones.

Introduction

Antimicrobial resistance is a growing problem for the management of tuberculosis (TB). For example, between 2009 and 2016, the number of global cases of multidrug-resistant tuberculosis (MDR-TB), defined as resistance of *Mycobacterium tuberculosis* to at least rifampicin and isoniazid, increased annually by over 20% (Lange et al., 2018). Additional resistance to a fluoroquinolone and at least one of three injectable agents (kanamycin, amikacin, or capreomycin), termed extensively drug-resistant tuberculosis (XDR-TB), accounted for about 6% of MDR-TB cases in 2018 (WHO, 2019). Alarmingly, the annual XDR-TB cases reported worldwide increased almost 10-fold between 2011 and 2018 (WHO, 2013; WHO, 2019). Increased prevalence of XDR-TB is not surprising, since the fluoroquinolone-containing combination therapies used to halt progression to XDR-TB utilized early quinolones (ciprofloxacin, ofloxacin, and sparfloxacin) that are only modestly effective anti-TB agents (Shandil et al., 2007).

Efforts to find more active fluoroquinolones led to C-8 methoxy derivatives that exhibit improved ability to kill *M. tuberculosis in vitro* (Dong et al., 1998). Two of these compounds, moxifloxacin and gatifloxacin, have been examined as anti-TB agents (Malik et al., 2006; Rodríguez et al., 2001; Ruan et al., 2016). Preclinical studies with moxifloxacin in murine TB models demonstrate effective treatment with reduced relapse frequency as well as treatment-shortening properties (Nuermberger et al., 2004). Indeed, single-drug clinical studies indicate that the early bactericidal activity of moxifloxacin is similar to that of first-line anti-TB-drugs, such as isoniazid and rifampicin. Thus, moxifloxacin could be part of effective multidrug combination treatments to shorten therapy time (Dorman et al., 2021; Pletz et al., 2004). Shorter treatment could increase

treatment compliance, further limiting the progression of MDR- to XDR-TB. However,

multiple phase III clinical trials have failed to corroborate the preclinical efficacy data, as

moxifloxacin did not shorten treatment time (Gillespie et al., 2014; Jawahar et al., 2013;

Jindani et al., 2014). One approach for increasing moxifloxacin efficacy is to find ways to

increase its lethal action.

The fluoroquinolones have two mechanistically distinct anti-bacterial effects: 1) they

block growth by forming reversible drug-gyrase-DNA complexes that rapidly inhibit DNA

synthesis, and 2) they kill cells (Drlica and Zhao, 2020). Death appears to arise in two

ways, one through chromosome fragmentation and the other through accumulation of

reactive oxygen species (ROS) (Drlica and Zhao, 2020). The latter appear to dominate

when DNA repair is proficient (Hong et al., 2020; Hong et al., 2019). We are particularly

interested in fluoroquinolone lethality, because *M. tuberculosis* possesses a remarkable

ability to evade host immune pressures (Ehrt and Schnappinger, 2009). In the absence

of an effective immune response, bactericidal drugs are essential for clearing infection

quickly. If the redox-based mechanisms of fluoroquinolone lethality seen with *E. coli*

extend to *M. tuberculosis,* opportunities may exist for increasing ROS and lethality,

thereby reducing treatment duration and increasing cure rate for MDR-TB.

Several methods are available for assessing the contribution of ROS to bactericidal

activity. Among these are direct detection of ROS levels using ROS-sensitive dyes,

characterization of mutants known to alter ROS levels, and examining effects of agents

known to suppress ROS accumulation. The present work adds detection of antibiotic-

induced changes in redox physiology of *M. tuberculosis* using a non-invasive, genetic

biosensor (Mrx1-roGFP2) (Bhaskar et al., 2014). The sensor measures the redox

potential (E_{MSH}) of a major mycobacterial thiol buffer, mycothiol, thereby assessing

oxidative effects independent of radical-sensitive fluorescent dye methods whose

interpretation has been debated (Dwyer et al., 2014; Liu and Imlay, 2013). Mrx1-roGFP2

is well-suited for this work, as it has been used to measure E_{MSH} of *M. tuberculosis* during

in vitro growth, infection of macrophages, and exposure to non-quinolone antimicrobials

(Bhaskar et al., 2014; Mishra et al., 2019; Mishra et al., 2017; Nambi et al., 2015).

In the present work, we began by asking how well *M. tuberculosis* fits the paradigm

for ROS-mediated killing by antimicrobials as developed from studies using *E. coli*. We

used moxifloxacin as a lethal stressor, because the fluoroquinolones are among the better

understood agents and because moxifloxacin is potentially important for tuberculosis

control. *M. tuberculosis* deviated markedly from *E. coli* by suppressing respiration rather

than increasing it during fluoroquinolone-mediated stress. That raised questions about

the source of ROS accumulation, since with *E. coli* ROS levels correlate with increased,

not decreased, respiration. We found that an increase in NADH (reductive stress)

accounted for the increase in ROS and lethality. When we artificially raised respiration,

ROS levels increased, and we observed enhanced moxifloxacin lethality *in vitro,* inside

macrophages, and in a murine model of infection. Raising respiration by N-acetyl cysteine

(NAC) treatment also reduced the emergence of resistance. Thus, the work provides a

mechanistic basis for developing respiration enhancers to increase the lethal action of

moxifloxacin and perhaps other anti-TB agents.

Results

Moxifloxacin-mediated lethality associated with oxidative stress during aerobic growth. To obtain a reference point for *M. tuberculosis* susceptibility to moxifloxacin, we determined minimal inhibitory concentration (MIC) using the resazurin microtiter assay (REMA) (Padiadpu et al., 2016). MIC for moxifloxacin, which ranged from 0.125 µM to 0.5 µM, was similar for H37Rv and several drug-resistant isolates (Table S1; isolate similarity was also seen for levofloxacin and ciprofloxacin). Thus, strain H37Rv appeared to be representative and appropriate for subsequent experiments.

As an initial probe for moxifloxacin-induced oxidative stress, we examined a derivative of *M. tuberculosis* H37Rv that expresses the redox biosensor Mrx1-roGFP2 (strain *Mtb*-roGFP2). This biosensor reports the redox state of the mycothiol redox couple (reduced mycothiol [MSH]/oxidized mycothiol [MSSM]) in the bacterial cytosol **(Fig 1A)** (Bhaskar et al., 2014). Exposure of *Mtb*-roGFP2 to hydrogen peroxide (H_2O_2) led to rapid (5 min) oxidation of the biosensor in a concentration-dependent manner (Fig S1) in which a 2-fold increase in the biosensor ratio represented exposure to 500 µM of H_2O_2 (Fig S1).

Increasing moxifloxacin concentration, up to 2.5X MIC (1X MIC = 0.5 µM) during a 48-h incubation, increased the biosensor ratiometric signal by almost 2-fold (**Fig 1B,** S2A**,** S2C), indicating elevated levels of oxidized mycothiol (MSSM). Above 2.5X MIC, MSSM levels appeared to plateau. Measurement of ROS with CellROX Deep Red showed a similar increase to 2.5X MIC moxifloxacin, followed by a plateau of mean fluorescence intensity (**Fig 1C**). Moxifloxacin treatment damaged DNA and oxidized lipids (Fig S3), indicating that drug-induced oxidative stress kills *M. tuberculosis* by degrading essential biomolecules.

Maximum bacterial killing was observed at 10X MIC, a concentration at which survival

decreased by ≈ 2 $\log_{10}$-fold at treatment day 1 and by 6 $\log_{10}$-fold at treatment day 10 (**Fig**

1D, 1E). Survival dropped in a concentration-dependent manner, with a minimum being

reached at a higher concentration than observed for maximal oxidative stress (10X MIC,

Fig 1B, 1C, 1D). The discordance between the biosensor signal plateauing at 2.5X MIC

and killing continuing to 10X MIC (**Fig 1B**) may derive from the existence of two lethal

mechanisms. One mechanism, chromosome fragmentation (Dwyer et al., 2007; Malik et

al., 2006), likely dominates at high quinolone concentration, with an ROS-based

mechanism dominating at lower concentrations. Indeed, for norfloxacin treatment of *E.*

coli, oxidative effects on killing are seen only at low-to-moderate drug concentrations

(Malik et al., 2007). As expected, oxidative stress was not observed in *M. tuberculosis*

treated with sub-inhibitory concentrations of moxifloxacin (Fig S2A). The increase in

biosensor signal was observed as early as 12 h after initiating moxifloxacin treatment (half

the doubling time of untreated control cells); it then increased significantly at 48 h (Fig

S2A, S2B). Thus, ROS production precedes and contributes to the death of *M.*

tuberculosis. As expected, ROS levels in a moxifloxacin-resistant isolate of *M.*

tuberculosis remained low upon treatment with moxifloxacin (Fig S4A).

We also compared the ability of a weakly effective fluoroquinolone (ciprofloxacin, MBC

= 2 µM; Table S2), a moderately effective compound (levofloxacin, MBC = 1 µM), and

moxifloxacin (MBC = 0.5 µM) to oxidize the biosensor, all at 2.5-fold MBC. Moxifloxacin

induced biosensor oxidation after treatment for 12 h and 24 h (Fig S4B). In contrast,

ciprofloxacin did not induce biosensor oxidation, and levofloxacin triggered oxidative

stress only after a 24-h treatment (Fig S4B). Thus, ROS levels do not correlate with MBC,

a parameter obtained after long incubation time (see Discussion). The more active

quinolones are also more lethal at the same fold MIC with several bacterial species,

including mycobacteria (Dong et al., 1998; Zhao et al., 1998; Zhao et al., 1997), these

data corroborate the link between oxidative stress and moxifloxacin lethality.

High-dose chemotherapy is postulated to slow the emergence of drug resistance if the

drug concentration is above the mutant prevention concentration (MPC) (Drlica and Zhao,

2007). However bacterial strains can paradoxically show elevated survival levels at

extremely high drug concentrations. In the case of nalidixic acid at very high

concentration, *E. coli* survival can be 100% (Luan et al., 2018). Similarly, *M. tuberculosis*

survival increased dramatically at high moxifloxacin concentration (**Fig 1D)**.

Some aspects of high-concentration survival are discordant when *E. coli* and *M.*

tuberculosis are compared. For example, with *E. coli* this phenotype is associated with

reduced ROS levels (Luan et al., 2018); with *M. tuberculosis*, the biosensor and CellROX

signals remained high at high levels of moxifloxacin (**Fig 1B**, **1C**). This difference between

the organisms, which is unexplained, encouraged further comparisions.

Reduction of moxifloxacin-mediated killing of *M. tuberculosis* by ROS-

mitigating agents. Work with *E. coli* supports the idea that ROS contribute to quinolone-

mediated killing of bacteria (Drlica and Zhao, 2020). For example, superoxide ($O_2^{\cdot-}$)

damages Fe-S clusters and increases the free iron (Fe) pool (Keyer and Imlay, 1996),

which may then drive the generation of toxic hydroxyl radical ($HO^{\cdot}$) via the Fenton reaction

(Dwyer et al., 2007; Kohanski et al., 2007). To examine the effects of Fe, we assessed

ROS levels in *M. tuberculosis* grown in minimal medium under Fe-deficient and Fe-

excess conditions. Fe-overload raised ROS levels (Fig S5). Treatment with thiourea (TU),

a thiol-based scavenger of ROS, then reversed the Fe-induced ROS increase (Fig S5).

During moxifloxacin treatment of *M. tuberculosis*, a concentration-dependent increase in

free Fe occurs (Fig S6). Pretreatment with thiourea, at the non-toxic concentration of 10

mM (Nandakumar et al., 2014), increased survival of *M. tuberculosis* by almost 10-fold

during co-treatment with moxifloxacin (1X – 10X MIC) (**Fig 2A, 2B,** S7). These data

support the connection between ROS accumulation and moxifloxacin-mediated killing.

Another test involves chelating free ferrous Fe with bipyridyl, a high-affinity Fe^{2+}

chelator: with *E. coli*, bipyridyl lowers the lethal action of multiple lethal stressors (Hong

et al., 2017; Kohanski et al., 2007). When we treated *M. tuberculosis* cultures with a non-

inhibitory concentration (250 µM) of bipyridyl before moxifloxacin (**Fig 2A**), killing was

reduced by >100 fold (**Fig 2C**), and ROS accumulation was lowered (**Fig 2D**). Treatment

with bipyridyl or thiourea also protected from moxifloxacin-mediated DNA damage and

lipid peroxidation (Fig S3A, S3B). Collectively, the data indicate that Fe-mediated ROS

production is the underlying oxidative stress associated with moxifloxacin lethality.

Work with *E. coli* also indicates that ROS accumulates even after removal of a lethal

stressor (Hong et al., 2019). When we treated *M. tuberculosis* with moxifloxacin in liquid

medium and then plated cells on antibiotic-free 7H11 agar (see scheme, **Fig 2E**, we found

that addition of catalase to the agar reduced bacterial killing (**Fig 2F;** peroxide diffuses

freely across the cell membrane (Seaver and Imlay, 2001), allowing exogenous catalase

to reduce endogenous ROS levels). Replacement of catalase with 2.5% bovine serum

albumin, which is not expected to degrade peroxide, failed to protect *M. tuberculosis* from

moxifloxacin-mediated killing (Fig S8). Finding that an anti-ROS agent suppresses killing

after the removal of moxifloxacin indicates that the primary damage (rapid formation of

fluoroquinolone-gyrase-DNA complexes) can be insufficient to kill *M. tuberculosis*. Thus,

the post-stressor accumulation of ROS and death seen with *E. coli* occurs with *M.*

tuberculosis.

We note that shorter drug-treatment times in 7H9 broth (2 days) prior to catalase

treatment on drug-free agar demonstrated higher survival than longer times (6 to 10

days), but eventually there was little difference (**Fig 2F**). A similar phenomenon has been

observed with *E. coli* (Hong et al., 2019). These data likely reflect ROS acting rapidly to

cause accelerated lethality and then having little effect on long-term killing. Overall,

studies with ROS-mitigating agents indicate that ROS contribute causally to quinolone-

mediated lethality with both *E. coli* and *M. tuberculosis*.

Reduction of moxifloxacin lethality by nutrient starvation and hypoxia. Our data

support previous work (Gengenbacher et al., 2010) in which nutrient starvation blocked

the lethal action of moxifloxacin (Loebel cidal concentration (LCC_{90}) > 32 µM) (Fig S9A,

S9B, Table S2). This phenomenon has also been reported for older quinolones with *E.*

coli (Hong et al., 2020). With *E. coli*, these effects have been attributed to a metabolic

shift that suppresses respiration, which is thought to be a major source of ROS.

Previous work (Gengenbacher et al., 2010; Wayne and Hayes, 1996) also shows that

hypoxic conditions reduce the lethal action of older fluoroquinolones with *M. tuberculosis*.

We confirmed this observation by showing that moxifloxacin-mediated killing decreases

significantly under hypoxic conditions (Fig S9A, S9C, Table S2). With the Wayne Cidal

Concentration (WCC_{90}) assay (Gengenbacher et al., 2010), the moxifloxacin

concentration required to kill 90% of hypoxic bacteria after 5 days of drug treatment was

~10 µM (Table S2). This value corresponds to 20X MIC or 22-fold higher than the LD_{90}

(lethal dose) with *M. tuberculosis* (0.45 µM) cultured under aerobic growth conditions for

the same treatment time (Fig S10).

The hypoxia findings fit with moxifloxacin lethality being associated with an increase

in ROS, an event that should be suppessed by O_2 limitation (intracellular generation of

ROS may depend largely on the transfer of electrons directly to molecular O_2 (Imlay,

2013)). However, hypoxic suppression of moxifloxacin lethality could be due to decreased

respiration. To test this idea, we stimulated anaerobic respiration by providing nitrate as

an alternative electron acceptor (Wayne and Hayes, 1998), since that restores norfloxacin

lethality with anaerobic *E. coli* (Dwyer et al., 2014). Surprisingly, nitrate lowered residual

anaerobic lethality of moxifloxacin by several fold (Fig S9D). Nitrate was even more

protective with *M. tuberculosis* exposed to metronidazole (Fig S9D), an antimicrobial

known to be lethal in the absence of oxygen (Wayne and Sramek, 1994). The surprising

protective effect of nitrate, which is considered in more detail in the Discussion,

encouraged further comparisons with *E. coli***.**

Effect of moxifloxacin on the *M. tuberculosis* transcriptome. When we examined

the *M. tuberculosis* transcriptome using published data from a 16-h exposure to 2X, 4X,

and 8X MIC moxifloxacin (Ma et al., 2015), we found that 359 genes exhibited altered

expression (2-fold change across all three treatment conditions). Of these, 219 genes

were upregulated, and 140 were down-regulated (Dataset S1). We asked whether

expression after prolonged moxifloxacin exposure (16 h) is largely a secondary effect

from induction of other genes by comparing early changes (4-h moxifloxacin exposure) in

expression for a set of 28 genes deregulated at 16 h. We saw little difference (Fig S11A,

S11B). Thus, the patterns we observed at 16 h appear to largely reflect a primary

transcription response.

When we classified the differentially expressed genes (DEGs) according to annotated

functional categories (Kapopoulou et al., 2011), we found that "Information Pathways"

and "Insertion Sequences and Phages" were two-fold over-represented in the

moxifloxacin-treated *M. tuberculosis* transcriptome (Table S3). These data suggest that

the bacterium responds to the drug mainly by regulating DNA remodeling, transcription,

and translational machinery.

Increased expression was also seen with genes involved in redox homeostasis, such

as thioredoxin (*trxB1, trxB2,* and *trxC*), alkyl hydroperoxide reductase (*ahpC*), the

SigH/RshA system, the copper-sensing transcriptional regulator *csoR,* and the *whiB-*

family (*whiB4* and *whiB7*) (**Fig 3A**). Indeed, the transcriptome of moxifloxacin-treated *M.*

tuberculosis showed 67% overlap with the transcriptional signature of H_2O_2-treated cells

(**Fig 3B**) (Voskuil et al., 2011). Statistical analysis showed a signficant overlap between

the moxifloxacin transcriptome and the response to oxidative stress (H_2O_2; p = 5.39 e-14)

and nitrosative stress (NO; p = 2.13 e-8) (Fig S12, Table S4). An induction of oxidant-

responsive genes (*sox, sod, mar*) is also seen in norfloxacin-treated *E. coli* (Dwyer et al.,

2007). Other conditions, such as hypoxia and acidic pH, showed non-significant overlap

with the moxifloxacin transcriptome (Fig S12, Table S4). Collectively, the data are

consistent with the idea that fluoroquinolones stimulate ROS accumulation.

Several genes involved in repairing DNA (*recA/O/R/X, ruv ABC, uvrD, ung, dnaE2,*

xthA, radA, alkA, lexA, nei, ligB), in DNA metabolism (*nrdF2, nrdR, pyrC, pyrR*), and in

Fe-S cluster biogenesis (*sufR*) (Anand et al., 2021; Das et al., 2021) were also

upregulated (**Fig 3A**). *E. coli* treated with norfloxacin shows similar upregulation of DNA-

damage responses (*lexA, uvr, rec* systems and error-prone DNA polymerases IV and V),

nucleotide metabolism, and Fe-S cluster biogenesis (*IscRUSA*) (Dwyer et al., 2007). Such

results are expected from the known ability of ROS to damage DNA (Dwyer et al., 2015;

Foti et al., 2012).

With *M. tuberculosis*, moxifloxacin suppressed expression of an Fe-responsive

repressor (*hupB*) and an iron siderophore (*mbtF*), consistent with drug treatment

increasing the intrabacterial pool of labile Fe (Fig S6). Additionally, the microarray data

suggest that Fe–S clusters in *M. tuberculosis* are exposed to ROS during treatment with

moxifloxacin, resulting in Fe release from damaged clusters and increased expression of

Fe-S repair pathways (*sufR*). In contrast, *E. coli* upregulates Fe-uptake machinery

(Dwyer et al., 2007). Thus, *M. tuberculosis* and *E. coli* may generate ROS in different

ways.

In *M. tuberculosis*, moxifloxacin repressed the expression of energy-efficient

respiratory complexes, such as succinate dehydrogenase (*rv0248c, sdhC*), cytochrome

bc1 (*qcrB*), and type I NADH dehydrogenase (*nuo* operon) **(Fig 3A).** In contrast, the level

of the energetically inefficient, non-proton pumping *ndh2* was increased. These data

indicate that *M. tuberculosis* slows primary respiration and shifts to a lower energy state

in response to moxifloxacin. Indeed, several energy-requiring pathways (*e.g.,* cell wall

biosynthesis, cofactor biogenesis, cell division, transport, and ESX-secretion systems)

were downregulated (**Fig 3A** and Dataset S1). Moreover, genes coordinating alternate

carbon metabolism, such as the gluconeogenesis (*pckA*) and glyoxylate cycle (*icl1*), were

up-regulated (**Fig 3A**). Interestingly, Icl1 protects *M. tuberculosis* from anti-TB drugs

(isoniazid, rifampicin, and streptomycin), presumably by counteracting redox imbalance

induced by these antibiotics (Nandakumar et al., 2014). The key idea is that *M.*

tuberculosis enters into a quasi-quiescent metabolic state in response to stress from

moxifloxacin, opposite to fluoroquinolone effects with *E. coli* (Dwyer et al., 2014; Dwyer

et al., 2007).

Moxifloxacin slows respiration. As a direct test for decelerated respiration, we

measured the Extracellular Acidification Rate (ECAR) and Oxygen Consumption Rate

(OCR) for moxifloxacin-treated *M. tuberculosis* as readouts for proton-extrusion into the

extracellular medium (reflecting glycolysis and TCA cycle activity) and for oxidative

phosphorylation (OXPHOS), respectively, using a Seahorse XFp Analyzer (Lamprecht et

al., 2016; Mishra et al., 2019). We incubated *M. tuberculosis* in unbuffered 7H9 + glucose

medium in an XF microchamber, exposed the culture to 10X MIC moxifloxacin, and later

to the uncoupler CCCP (addition of CCCP stimulates respiration to the maximal level).

The difference between basal and CCCP-induced OCR estimates the spare (reserve)

respiratory capacity (SRC) available to counteract stressful conditions (Lamprecht et al.,

2016; Saini et al., 2016).

As expected for a growing, drug-free culture of *M. tuberculosis,* OCR showed a

gradual increase over the duration of the experiment (400 min); OCR increased further

upon uncoupling by CCCP (**Fig 4A**). In contrast, addition of 10X MIC of moxifloxacin

inhibited the time-dependent increase in basal OCR (**Fig 4A**), and the level of OCR upon

uncoupling by CCCP was markedly lower than in the absence of moxifloxacin (**Fig 4A**).

In parallel, we measured viability of *M. tuberculosis* and found that the bacterial culture

maintained 100% survival during the the entire incubation period and for an additional 4

h (Fig S13). Thus, the stalled oxygen-consumption effect of moxifloxacin cannot be attributed to dead cells.

The ECAR response increased for 60 min and then gradually dropped for untreated bacteria. This transient, early increase in ECAR was absent in cultures of moxifloxacin-treated *M. tuberculosis*, which exhibited a greater reduction of ECAR than untreated cultures at late times (**Fig 4B**). At the end of the experiment, moxifloxacin-treated cells were completely exhausted of glycolytic capacity (**Fig 4B).** The slowing in OCR and ECAR was also seen with treatment at lower concentrations of moxifloxacin (1X and 2.5X MIC) (Fig S14), while an anti-TB drug (ethambutol) that does not generate ROS, failed to significantly affect either OCR or ECAR (Fig S15).

In summary, moxifloxacin decelerates respiration and carbon catabolism in *M. tuberculosis*, which likely renders the bacterium metabolically quiescent and therefore less readily killed by moxifloxacin. In support of this idea, a recent study shows that pretreatment with the anti-TB drugs bedaquiline (an ATP synthase inhibitor) or Q203 (a cytochrome C oxidase inhibitor) reduces moxifloxacin-mediated killing of *M. tuberculosis* (Lee et al., 2019).

ROS derive from increased NADH. Since moxifloxacin slows respiratory metabolism (**Fig 4A**) while increasing ROS levels (**Fig 1B, 1C**), a source of ROS other than respiration must exist. Slowed respiration during anoxia or chemical hypoxia places the electron transport chain (ETC) in a reduced state, and NADH accumulates. This phenomenon is known as reductive stress (Mavi et al., 2019). Using a redox cycling assay**,** we detected an accumulation of NADH in 1X, 2.5X and 10X MIC moxifloxacin-treated *M. tuberculosis* at 48 h **(Fig 5A)**. NAD^+ was raised at 1X MIC moxifloxacin, but it showed no significant

difference from the untreated control at lethal, elevated moxifloxacin concentrations (**Fig**

5B); the total NAD/H pool decreased at high drug concentrations (2.5X and 10X MIC; Fig

S16). Thus, the NADH/NAD$^+$ ratio increased as moxifloacin concentration increased (**Fig**

5C), indicating reductive stress.

We also measured NADH-reductive stress by expressing a genetically encoded, non-

invasive biosensor of the NADH/NAD$^+$ ratio (Peredox) (Bhat et al., 2016) (Fig S17). A

ratiometric increase in Peredox fluorescence indicated an increase in the NADH/NAD$^+$

ratio upon moxifloxacin treatment (1X to 40X MIC) at 48 h. We observed a similar

increase in the Peredox ratio upon treatment of *M. tuberculosis* with bedaquiline, a drug

known to increase the NADH/NAD$^+$ ratio (Fig S17) (Bhat et al., 2016). As with NADH,

NADPH accumulated, and the NADPH/NADP$^+$ ratio increased in response to

moxifloxacin treatment (Fig S18). Thus, moxifloxacin preferentially triggers NAD(P)H

accumulation in *M. tuberculosis*.

In principle, stalled ETC and excessive reducing equivalents (NADH) can stimulate

ROS production in two ways. In one, NADH mobilizes bound iron, thereby increasing the

labile iron pool (Jaeschke et al., 1992). As previously mentioned, we found a drug-

concentration-dependent increase in the free iron pool in *M. tuberculosis* after

moxifloxacin treatment (Fig S6). Additionally, NADH autooxidation keeps iron in a

reduced state (Fe^{2+}); thus, NADH can drive the Fenton reaction towards hydroxyl radical

generation (Jaeschke et al., 1992). Second, reduced components of the electron

transport chain can directly transfer electrons to molecular oxygen to generate ROS

(Kareyeva et al., 2012; Vinogradov and Grivennikova, 2016). If these ideas are correct,

dissipation of the NADH overload, i.e., reductive stress, should lower the labile iron pool,

ROS surge, and moxifloxacin lethality.

We lowered the NADH/NAD$^+$ ratio with an *M. tuberculosis* strain that constitutively

expresses *Lactobacillus brevis* NADH oxidase (*Mtb-LbNox*) (**Fig 5C**, S17) (Titov et al.,

2016). In comparison to wild-type *M. tuberculosis*, *Mtb-LbNox* shows decreased levels of

NADH and increased NAD$^+$ without affecting the total pool of NAD$^+$ + NADH (**Fig 5A, 5B,**

5C, S19). As expected, moxifloxacin-treated *Mtb-LbNox* displayed a significantly reduced

free iron pool (Fig S6), ROS surge (**Fig 5D**), decreased DNA damage (Fig S6), and a 25-

to 30-fold higher surivival than wild-type cells (**Fig 5E**). Thus, increased NADH/NAD$^+$ ratio

occurring during metabolic quiescence caused by moxifloxacin accounts for the increased

labile iron pool, which in turn catalyzes Fenton-mediated production of ROS and cell

death.

The growth, OCR, and ECAR of *Mtb-LbNox* in 7H9 broth were the same as with wild-

type cells (Fig S20). Furthermore, MIC of moxifloxacin and other anti-TB drugs

(rifampicin, ethambutol, and bedaquiline) with *Mtb-LbNox* was similar to that of wild-type

M. tuberculosis (Table S1, S5), indicating that the interaction between drug and its

primary target is unaffected by overexpression of *LbNox.* In contrast, *Mtb-LbNox* exhibited

a 2-fold lower MIC for isoniazid, which is consistent with the interaction of isoniazid and

its primary target (enoyl-ACP reductase, InhA) depending on the NADH/NAD$^+$ ratio in *M*

.tuberculosis (Vilchèze et al., 2005). Thus, overexpression of *LbNox* appears to have no

effect on growth, metabolism, or respiration of *M. tuberculosis* that would complicate our

interpretation that NADH dissipation protects from moxifloxacin-mediated killing.

N-acetyl cysteine stimulates respiration, oxidative stress, and moxifloxacin lethality. Previous work showed that stimulating respiration using cysteine and N-acetyl cysteine (NAC) elevates the killing activity of combinations of anti-TB drugs (Vilchèze et al., 2017; Vilchèze and Jacobs, 2021). Since *M. tuberculosis* responded to moxifloxacin by dampening OXPHOS, we expected that countering the dampening with NAC would provide a way to increase ROS further and therefore enhance moxifloxacin lethality. We first measured time-dependent changes in OCR of *M. tuberculosis* upon exposure to a non-toxic dose of NAC (1 mM). Addition of NAC alone produced a sharp increase in OCR that reached its maximal level by 200 min (**Fig 4C**). At later times, OCR gradually declined; it did not increase upon addition of CCCP (**Fig 4C**), indicating that NAC-stimulated respiration exhausted the spare respiratory capacity of *M. tuberculosis*. Pre-treatment with NAC reversed moxifloxacin-mediated slowing of respiration (**Fig 4D**). As with NAC alone, the OCR of *M. tuberculosis,* treated with NAC plus moxifloxacin, increased for 200 min and then gradually dropped; CCCP remained ineffective at stimulating OCR (**Fig 4D**). These data indicate that NAC-stimulated oxygen consumption by *M. tuberculosis* overcomes the dampening effect of moxifloxacin on bacterial respiration.

Respiration, stimulated by NAC, augmented ROS accumulation upon moxifloxacin treatment and increased lethality (**Fig 6A).** NAC elevated the Mrx1-roGFP2-dependent redox signal for moxifloxacin-treated *M. tuberculosis*, confirming the increase in oxidative stress (**Fig 6B**, S2C, S2D). We emphasize that NAC had <u>no</u> effect on moxifloxacin MIC, as measured by REMA and 7H11 agar plate assay (Fig S21A, S21B); thus, the primary interaction between drug and DNA gyrase (cleaved complex formation) is unaffected by

NAC. Supplementation of moxifloxacin at 1X and 5X MIC with 1 mM of NAC reduced

survival by 21- and 11-fold, respectively (**Fig 6C**), levels that are consistent with recently

published results using drug combinations (Vilchèze and Jacobs, 2021).

As a complement to the suppression of killing by addition of catalase after drug

removal (**Fig 2F**), we treated *M. tuberculosis* cultures with 1X and 5X MIC of moxifloxacin

and then plated the cells on 7H11 agar with 1 mM NAC. Post-drug addition of NAC

blocked formation of colonies that would otherwise have been observed (Fig S22).

Extended incubation (>4 weeks) resulted in the appearance of small colonies on the NAC-

containing plates, which was likely due to instability/oxidation of NAC ($T_{1/2}$~ 14 h) under

aerobic conditions (Held and Biaglow, 1994; Jaworska et al., 1999; Sommer et al., 2020).

NAC also potentiated moxifloxacin lethality (5- to 7-fold) with nutrient-starved, dormant

M. tuberculosis (**Fig 6D**). These data agree with recent results in which activation of

oxidative metabolism and respiration, through addition of L-cysteine to nutrient-starved

cultures, reduced the fraction of persister subpopulations and enhanced the *in vitro* lethal

activity of isoniazid and rifampicin (Srinivas et al., 2020). As expected, a similar treatment

with NAC failed to enhance moxifloxacin lethality under hypoxic conditions (**Fig 6E**),

consistent with NAC functioning by inducing oxygen consumption and elevating ROS

levels.

To address the effect of NAC on the acquisition of moxifloxacin resistance, we

measured the mutant prevention concentration (MPC) i.e., the minimal concentration of

moxifloxacin at which no moxifloxacin-resistant clone emerges on moxifloxacin-

containing 7H11 agar inoculated with ~ 2.5 x 10^9 bacteria (Drlica and Zhao, 2007; Singh

et al., 2017). With a clinical MDR isolate of *M. tuberculosis* (NHN1664), moxifloxacin

exhibited an MPC of 4 µM that was reduced to 2 µM upon co-treatment with either 1 mM

or 2 mM NAC. For a lower moxifloxacin concentration (1 µM, 2X MIC), the fraction of cells

recovered decreased by 10-fold and 100-fold when co-plated with 1 mM and 2 mM NAC,

respectively (Table S6). Reduction of MPC is expected when lethal activity is increased

(Cui et al., 2006).

We also addressed the possibility that some effects of NAC derived from adducts

formed with moxifloxacin. A series of biochemical tests (thin layer chromatography, NMR,

fluorescence assays, and LC-MS) revealed no evidence for adduct formation (Fig S23).

Moxifloxacin-induced redox imbalance during infection of macrophages. The

Mrx1-roGFP2 biosensor previously showed that first-line anti-TB drugs cause an

oxidative shift in the E_{MSH} of *M. tuberculosis* residing inside macrophages (Bhaskar et al.,

2014). When we infected macrophages with *M. tuberculosis* H37Rv or an MDR clinical

isolate (strain NHN1664), each expressing the biosensor Mrx1-roGFP2, *M. tuberculosis*

displayed redox-heterogeneity (E_{MSH}-basal [-280 mV], E_{MSH}-oxidized [-240 mV], and

E_{MSH}-reduced [-320 mV]); the E_{MSH}-reduced subpopulation was predominant (Bhaskar et

al., 2014; Mishra et al., 2019; Mishra et al., 2021) (**Fig 7A**, S24). Treatment of *M.*

tuberculosis-infected THP-1 macrophages with various concentrations of moxifloxacin

significantly increased the E_{MSH}-oxidized subpopulation (fuchsia colored line, **Fig 7A,**

S24).

The shift toward the E_{MSH}-oxidized redox state was seen with moderate

concentrations of moxifloxacin (5X MIC) at an early time point (12 h post infection), before

M. tuberculosis survival started to drop (**Fig 7A, 7B**). Thus, moxifloxacin-mediated

oxidative stress precedes killing: oxidative stress is not simply a consequence of cell

death. Bacterial survival was reduced more by higher moxifloxacin concentrations (10X

and 20X MIC) or longer incubation times (24 h and 48 h) (**Fig 7B**).

We also assessed the effect of NAC on E_{MSH} and killing of *M. tuberculosis* inside

macrophages. NAC itself is not cytotoxic to macrophages up to at least 5-10 mM (Amaral

et al., 2016; Vilchèze et al., 2017), but at those elevated concentrations it exhibits anti-

mycobacterial properties (Amaral et al., 2016; Cao et al., 2018). Consequently, we

selected non-cytotoxic, non-anti-tuberculous concentrations of NAC (1 mM and 2 mM)

(Amaral et al., 2016) for our test with moxifloxacin. NAC supplementation induced an

oxidative shift in the E_{MSH} of *M. tuberculosis* that exceeded that seen with moxifloxacin

alone (**Fig 7C,** S24). NAC alone had no effect on survival of *M. tuberculosis* inside THP-

1 macrophages; however, the NAC plus moxifloxacin combination decreased bacterial

burden 5-10 times more than moxifloxacin alone in both concentration- and time-

dependent manners (**Fig 7D**). We note that for either isoniazid or an isoniazid-rifampicin

combination, NAC potentiates lethality only at late times (day 6 or 7) (Vilchèze et al., 2017;

Vilchèze and Jacobs, 2021); NAC acts more rapidly with moxifloxacin, as the killing effect

is evident at days 1 and 2 (**Fig 7D**). Thus, NAC may not function in the same way with all

anti-TB drugs inside macrophages.

Moxifloxacin also lowered the level of an E_{MSH}-reduced subpopulation inside

macrophages (blue line, **Fig 7A**, **7C**, S24); this subpopulation was further diminished by

NAC supplementation. E_{MSH}-reduced subpopulations are phenotypically tolerant to anti-

TB drugs (Bhaskar et al., 2014; Mishra et al., 2019) and to moxifloxacin (Fig S25). Overall,

results with infected macrophages demonstrate that accelerating respiration and

oxidative metabolism by NAC can enhance moxifloxacin lethality.

Potentiation of moxifloxacin lethality with NAC lowers *M. tuberculosis* burden in a murine model of infection and restricts the emergence of resistance. We next asked whether NAC-mediated enhancement of moxifloxacin lethality occurs *in vivo*. Recent work by Vilcheze *et. al.* (Vilchèze and Jacobs, 2021) directed our choice of murine models: they found little enhancement by NAC on the overall lethality of a combination of moxifloxacin and ethionamide when measured after long incubation using an acute model of murine tuberculosis. Since ROS accelerate killing *in vitro* without increasing the extent of killing (Liu et al., 2012), we shortened the drug treatment time. For this pilot test, we infected BALB/c mice with the clinical MDR strain, *M. tuberculosis* NHN1664. Three weeks after aerosol-mediated infection at a low dose (~100 bacilli), mice were treated once daily for ten days with a low dose of moxifloxacin (50 mg/kg body weight), NAC (500 mg/kg body weight), or moxifloxacin plus NAC at the same doses as for mono treatments (**Fig 8A**). After 10 days, moxifloxacin monotherapy reduced the bacillary load by 3-fold and 100-fold in lung and spleen, respectively; NAC alone exhibited no anti-bacterial activity (**Fig 8B, 8C**). The combination of moxifloxacin plus NAC reduced bacterial burden by another 4- and 12-fold beyond that observed for moxifloxacin alone for lung and spleen, respectively (**Fig 8B, 8C**). Thus, this preliminary finding indicates that NAC stimulates the lethal action of moxifloxacin *in vivo*.

As a proof of concept. we also measured the effect of NAC on the selection of moxifloxacin-resistant mutants in mice as described (Almeida et al., 2007; Ginsburg et al., 2005). We implanted ~ 4x 10^3 *M. tuberculosis* NHN1664 in the lungs of BALB/c mice, and at day 14 post-infection, when the mean lung $\log_{10}$ CFU reached ~ 10^7, mice received moxifloxacin, NAC, or moxifloxacin plus NAC. After 14 days of post-treatment, 66% of

moxifloxacin-treated mice harbored resistant bacteria as compared to 17% when NAC

was also present. The combination of moxifloxacin plus NAC brought the number of

mutants down to the level seen with untreated animals (Table S7). Enumeration of total

resistant colonies confirmed that while moxifloxacin treatment increases emergence of

resistance, NAC supplementaion reduces the recovery of moxifloxacin-resistant mutants

by 8-fold (Table S7) in our pilot experiment.

Discussion

The experiments described above show that *M. tuberculosis* responds to moxifloxacin through a lethal stress response characterized by suppressed respiration, elevated NADH levels, and ROS accumulation (shown schematically in **Fig 9**). Although ROS accumulation is also observed with *E. coli* during lethal stress, early steps in the *M. tuberculosis* stress response are distinct. *M. tuberculosis* enters a metabolically quiescent state characterized by increased NADH levels and NADH/NAD$^+$ ratio plus reduced respiratory and glycolytic rates. This response is opposite to that seen with *E. coli* (Dwyer et al., 2014; Kohanski et al., 2007; Lobritz et al., 2015). Decelerated respiration may be an adaptive strategy against ROS-inducing agents produced by the host. Production of ROS likely derives from increased NADH/NAD$^+$ ratio, which increases the labile, reduced form of Fe to fuel the Fenton reaction. Interfering with the NADH/NAD$^+$ increase dampened reductive stress and moxifloxacin lethality, suggesting that NADH disposal pathways could be targeted to enhance fluoroquinolone lethality with *M. tuberculosis*.

ROS contribution to lethality. Studies with *E. coli* indicate that ROS kill bacteria by damaging macromolecules *i.e.* DNA breakage, protein carbonylation, and lipid peroxidation (Drlica and Zhao, 2020). Our data support the general principle that ROS contribute to stress-mediated death of moxifloxacin-treated *M. tuberculosis*: 1) biosensors show an increase in ROS at lethal drug concentrations, 2) iron concentrations increase with moxifloxcin treatment, and 3) two ROS-mitigating agents, thiourea and bipyridyl, reduce oxidative stress and killing. However, the striking increase in survival for *E. coli* at very high quinolone concentrations is associated with a drop in ROS, while this is not the case with *M. tuberculosis*; this difference remains unexplained.

Another difference is the lowering of residual fluoroquinolone lethality by nitrate with

hypoxic *M. tuberculosis*: the opposite is observed with norfloxacin-treated *E. coli*.

Previous *M. tuberculosis* work has shown that nitrate protects hypoxic bacilli from killing

caused by exposure to reactive nitrogen intermediates (Tan et al., 2010), acid stress (Tan

et al., 2010), sudden anaerobiosis (Sohaskey, 2008), or inhibition of type II NADH-

dehydrogenase (*ndh-2*) by thioridazine (Sohaskey, 2008). Two phenomena may

contribute to nitrate-mediated protection. 1) *M. tuberculosis* reduces nitrate to nitrite

(Cunningham-Bussel et al., 2013), which oxidizes ferrous Fe to the ferric state, thereby

displacing Fe from critical Fe-sulfur clusters and inducing bacteriostasis (Cunningham-

Bussel et al., 2013). Nitrite also remodels the *M. tuberculosis* transcriptome similar to

several host-imposed stresses that suppress antibiotic-mediated lethality (Cunningham-

Bussel et al., 2013) and ensure bacterial survival. 2) Nitrate, as a terminal electron

acceptor, recycles or disposes of excess reducing equivalents that accumulate during

anaerobic conditions (Shapleigh, 2009). Discovering strategies to halt such adaptive

remodeling (*e.g.,* nitrate reduction) in *M. tuberculosis* could serve as adjunct therapy for

moxifloxacin with hypoxic bacilli.

Reductive stress as a source of ROS. When we dissipated NADH-reductive stress

by overexpression of *LbNox*, the ROS surge was mitigated, and moxifloxacin lethality was

lowered. Production of NADH can also be lowered by stimulating the glyoxylate shunt

via isocitrate lyase (*icl*) – deletion of *icl* increases the lethal action of isoniazid, rifampicin,

and streptomycin (Nandakumar et al., 2014), suggesting that NADH-mediated ROS

accumulation may also contribute to the lethal action of these agents. The observed

increase in NADPH/NADP$^+$ ratio in response to moxifloxacin exposure could assist *M.*

tuberculosis in mitigating oxidative stress by activating NADPH-dependent antioxidant

systems, thioredoxins, and mycothiol. As shown by microarray data, the increase in

NADPH/NADP$^+$ ratio could also be a consequence of down-regulation of NADPH-

requiring anabolic pathways, such as polyketide/lipid biogenesis, amino acid anabolism,

and *de novo* nucleotide synthesis.

In summary, *M. tuberculosis* appears to tolerate lethal stressors of the host immune

system by decelerating respiration coupled with dissipation of NADH-reductive stress.

The need to dissipate reductive stress may be of general importance, as this type of stress

is also generated and amplified by hypoxia, inhibition of aerobic respiration by NO,

and catabolism of fatty acids (Singh et al., 2009; Voskuil et al., 2003). *M. tuberculosis*

dissipates reductive stress through WhiB3-mediated anabolism of polyketide lipids (Singh

et al., 2009). Other bacterial species also dispose of excess reductant, which they do by

1) using nitrate as a terminal electron acceptor during hypoxia (Tan et al., 2010), 2) fueling

fermentation under nitrosative stress (Richardson et al., 2008), and 3) secreting

phenazines (redox-active polyketides) that generate NAD$^+$ from NADH (Price-Whelan et

al., 2007). Moxifloxacin is the first anti-mycobacterial agent shown to stimulate NADH-

reductive stress.

Lethality enhancement by N-acetyl cysteine*. Vilchèze and Jacobs (Vilchèze and

Jacobs, 2021) reported that NAC boosts the lethality of anti-TB drug combinations *in vitro*.

We emphasize that a general, enhancing effect on drug-target interactions by NAC is

unlikely, because such effects are usually detected by MIC measurement (Drlica and

Zhao, 2020): NAC has no effect on moxifloxacin MIC. This observation reinforces the

concept that drug-target interactions are mechanistically distinct from downstream ROS-

based killing (protection from killing, which is called tolerance when <u>no</u> change in MIC

occurs, is distinct from resistance, which is measured as an increase in MIC).

To simplify interpretation of killing data, we focused on a single drug, moxifloxacin,

rather than drug combinations. Previous demonstration that NAC stimulates respiration

(Vilchèze and Jacobs, 2021) made it a useful tool for examining killing mechanism. NAC

increased moxifloxacin-mediated ROS accumulation and lethality with cultured *M.*

tuberculosis. Increased killing was initially surprising, because NAC has anti-oxidant

properties expected to reduce moxifloxacin lethality (Ezeriṇa et al., 2018). It is likely that

the effect of NAC depends on the level of stress, since with other bacterial species, genes

(*mazF, cpx, lepA*) and treatments (chemical generation of superoxide, H_2O_2) are known

to be protective at low levels of stress but destructive at high ones (Dorsey-Oresto et al.,

2013; Li et al., 2014; Mosel et al., 2013; Rodríguez-Rojas et al., 2020). Indeed, treatment

with a sub-lethal concentration of H_2O_2 activates the OxyR-regulon in *E. coli* and "primes"

the bacterium to conteract subsequent oxidative-stress challenge. At higher

concentrations, H_2O_2 is cidal (Imlay and Linn, 1986). With the high level of stress

produced by moxifloxacin, the respiration-stimulating activity of NAC dominates and adds

to ROS derived from NADH-reductive stress.

As expected from *in vitro* studies, NAC increased moxifloxacin lethality with infected

macrophages and mice. The kinetic effects of ROS explain why our short-term treatment

model with infected mice showed stimulation by NAC, while the longer treatment used

previously with antimicrobial combinations did not (Vilchèze and Jacobs, 2021).

We addressed the potential of NAC to suppress the emergence of resistance by

measuring its effect on MPC. Although MPC is a bacteriostatic parameter (MIC of the

least susceptible mutant subpopulation), highly lethal agents lower it in an animal model

of infection, presumably by killing mutants (Cui et al., 2006). Thus, we were not surprised

that stimulation of moxifloxacin lethality by NAC lowered MPC. In a murine infection

model, NAC reduced the recovery of resistant mutants by 8-fold.

Although MPC for moxifloxacin is lower than concentrations achieved in serum by

approved doses (Dong et al., 2000), the clinical situation is likely complex: moxifloxacin

shows poor penetration into caseous regions of tubercular granulomas in a rabbit model

of experimental tuberculosis (Prideaux et al., 2015; Sarathy et al., 2019). Low, local

moxifloxacin concentrations may promote the emergence of fluoroquinolone resistance

(Davies Forsman et al., 2020). Thus, NAC supplementation could increase the probability

that moxifloxacin concentration is above MPC in vivo, but additional work is required to

test this idea.

NAC is a potential candidate as a lethality enhancer, because it is relatively safe for

humans (it has been used to alleviate drug-induced toxicity in TB-patients (Baniasadi et

al., 2010; Kranzer et al., 2015)). Furthermore, NAC by itself exhibits anti-mycobacterial

activity inside macrophages, mice, and guinea pigs (Amaral et al., 2016; Palanisamy et

al., 2011), and a small clinical trial in which NAC was administered to TB patients

improved host health, immunological response, and early sputum conversion

(Mahakalkar et al., 2017; Nagrale et al., 2013). The present work provides a mechanistic

foundation for refining NAC-based enhancement of anti-tuberculosis agents. Assessing

the general clinical significance of ROS enhancement is also complex, because ROS

accelerates killing without lowering bacterial survival at long incubations. This

phenomenon has been observed with *S. aureus* (Liu et al., 2012), *E. coli* (Hong et al.,

2019), and *M. tuberculosis* (**Fig. 2F**; S4B). With *M. tuberculosis*-infected mice, NAC

stimulates reduction of bacterial load at short but not long incubation times (Vilchèze and

Jacobs, 2021). Thus, an appropriate dosing interval must be determined, initially guided

by *in vitro* pharmacodynamic studies (Almeida et al., 2007; Firsov et al., 2003; Ginsburg

et al., 2005; Gumbo et al., 2004). Relevant tissue concentrations of NAC and moxifloxacin

for various doses over time must also be determined before clinical relevance can be

assessed. The present report, plus that of Vilchèze and Jacobs (Vilchèze and Jacobs,

2021), encourage such followup work.

Materials and Methods

Bacterial strains and culture conditions. *M. tuberculosis* H37Rv and drug-

resistant clinical isolates JAL 1934, JAL 2287, JAL 2261, and BND 320 were kind gifts

from Dr. Kanury V.S. Rao (International Centre for Genetic Engineering and

Biotechnology, Delhi, India). MDR strain NHN1664, originally isolated in China, was

obtained from BEI Resources, NIAID, NIH. Strain *Mtb*-roGFP2 was generated by

transforming *M. tuberculosis* strain H37Rv with an *E. coli*-mycobacterial shuttle vector,

pMV762 (Singh et al., 2006; Steyn et al., 2003). Plasmid pMV762 contains the Mrx1-

roGFP2 biosensor construct under control of the *M. tuberculosis hsp60* promoter and a

hygromycin-resistance gene as a selection marker. Hygromycin (MP Biomedical, cat. No.

0219417091, Santa Ana, CA) was used at a final concentration of 50 µg/mL. Plasmid

(pGMCgS-0×-Ptb38-LbNOX-FLAG-SD1) containing a *lbnox* construct was kind gift from

Dr. Dirk Schnappinger and Dr. Sabine Ehrt (Weill Cornell Medicine, New York, USA).

Plasmid was electroporated into wild-type *M. tuberculosis* H37Rv to create the *Mtb-*

LbNox strain. Streptomycin was used at a final concentration of 25 µg/mL. Plasmid

pMV762-Peredox-mcherry was a kind gift from Dr. Ashwani Kumar (Council of

Scientific and Industrial Research, Institute of Microbial Technology, Chandigarh , India).

All strains, including laboratory *M. tuberculosis* strain H37Rv and *M. tuberculosis*

expressing the Mrx1-roGFP2 redox biosensor, were grown in 7H9 broth supplemented

with 0.2% glycerol, 0.05% Tween-80, and ADS (0.5% albumin, 0.2% dextrose, and

0.085% NaCl) with shaking at 180 RPM in a rotary shaker incubator (Lab Therm LT-X;

Kuhner, Basel, Switzerland) or on 7H11 agar supplemented with ADS or OADC (ADS

plus 0.05% oleic acid and 0.004% catalase (Sigma Aldrich, cat. No. C9322, St Louis, MO)

or ADC (ADS plus 0.004% catalase) at 37°C. Bacterial strains expressing Mrx1-roGFP2

were grown in media containing hygromycin.

Determination of minimal inhibitory concentration (MIC) under aerobic growth

conditions. MIC was determined by a resazurin microtiter assay (REMA) using 96-well

flat-bottom plates (Padiadpu et al., 2016). *M. tuberculosis* strains were cultured in

7H9+ADS medium (Bhaskar et al., 2014) and grown to exponential phase (OD_{600} = 0.4

to 0.8). Approximately 1×10^5 bacteria per well were added in a total volume of 200 μL of

7H9+ADS medium. Wells lacking *M. tuberculosis* served as controls. Additional controls

consisted of wells containing cells without drug treatment (growth control). After 5 days

of incubation at 37°C in the presence of fluoroquinolone (or NAC), 30 μL of 0.02%

resazurin (Sigma-Aldrich, cat. No. R7017, St Louis, MO) was added, and plates were

incubated for an additional 24 h. Fluorescence intensity was measured using a

SpectraMax M3 plate reader (Molecular Devices, San Jose, CA) in the bottom-reading

mode with excitation at 530 nm and emission at 590 nm. Percent inhibition was calculated

from the relative fluorescence units compared with an untreated control culture; MIC was

taken as the lowest drug concentration that resulted in at least 90% reduction in

fluorescence compared to the untreated growth control.

Determination of bacterial survival. To determine the number of viable bacilli,

aliquots were removed from cultures, cells were concentrated by centrifugation (4200 g,

5 min) to remove drug or treatment compounds, and they were resuspended in an equal

volume of medium. Ten-fold dilutions were prepared, and 20 µL aliquots were spotted on

drug-free 7H11 agar plates containing 10% ADC or ADS. Plates were incubated for 3-4

weeks at 37°C for CFU enumeration by visual inspection.

Concentration-kill curves were obtained by treating exponentially growing cultures of

M. tuberculosis (OD_{600} of 0.3 or $\approx 5 \times 10^7$ cells/mL) with various concentrations of

moxifloxacin. Tubes were incubated with shaking at 180 RPM for 10 days at 37°C.

Aliquots were taken at various intervals, serially diluted, and plated on drug-free agar for

CFU enumeration.

LD_{90} (lethal dose) for moxifloxacin under *in vitro,* aerobic-culture conditions was

measured by treating 10-mL cultures in 50-mL centrifuge tubes under aerobic conditions

(shaking at 180 RPM, 37°C), followed by CFU measurement. A 90% reduction in CFU

on treatment day 5, compared with the input control, was defined as LD_{90}.

To determine the effect of an Fe-chelator on moxifloxacin-mediated lethality, 250 µM

(a non-inhibitory concentration) of bipyridyl (Sigma-Aldrich, cat.no. D216305, St Louis,

MO) was added to bacterial cultures 15 min prior to moxifloxacin addition; bipyridyl was

present throughout the experiment (i.e., 10 days).

For experiments with thiourea or N-acetyl cysteine (NAC), thiourea (10 mM) or NAC

(1 mM) was added 1 h prior to moxifloxacin and was maintained for 2 days or 10 days.

Aliquots were taken at indicated times, serially diluted, and plated for CFU enumeration

on drug-free agar.

Post-stressor bactericidal activity of moxifloxacin. *M. tuberculosis* cultures were

treated with moxifloxacin for 48 h, the drug was removed by washing, and cells were

plated on drug-free 7H11 agar with or without 1 mM NAC or with 2.5% bovine serum

albumen (BSA), followed by visual CFU determination.

Measurement of E_{MSH} using the Mrx1-roGFP2 redox biosensor. For intra-

mycobacterial E_{MSH} determination during *in-vitro* growth or during *ex-vivo* macrophage-

infection conditions, strain *Mtb*-roGFP2 was grown in 7H9 broth or in THP-1

macrophages, respectively. Cultures were then treated with moxifloxacin, and at various

times they were treated with 10 mM N-ethylmaleimide (NEM; Sigma-Aldrich, cat. No.-

E3876, St Louis, MO) for 5 min at room temperature followed by fixation with 4%

paraformaldehyde (PFA; Himedia, cat. No. GRM3660, Mumbai, India) for 1 h at room

temperature. Bacilli or infected macrophages were analyzed using a FACSVerse Flow

cytometer (BD Biosciences, San Jose, CA). Intramycobacterial E_{MSH} was determined

using the Nernst Equation as described previously (Bhaskar et al., 2014). E_{MSH} is defined

as the standard reduction potential of the MSH/MSSM redox couple (reduced mycothiol

to oxidized mycothiol*.*

The biosensor response was measured by analyzing the fluorescence ratio at a fixed

emission (510 nm) after excitation at 405 and 488 nm. Data were analyzed using the BD

FACSuite software. These ratiometric data were normalized to measurements with cells

treated with 10 mM cumene hydroperoxide (Sigma-Aldrich- cat. No. 247502, St Louis,

MO), which reports maximal oxidation of the biosensor, and 20 mM dithiothreitol (Sigma-

Aldrich, cat. No. D9779, St Louis, MO), which reports maximal reduction. 10,000 events

per sample were analyzed.

ROS measurement using CellROX Deep Red. Cultures of exponentially growing *M.*

tuberculosis at an initial OD_{600} of 0.3 were treated with various concentrations of

moxifloxacin for 48 h. Sterile 50 mL-capacity polypropylene centrifuge tubes (Abdos, cat

no. P10404, Roorkee, India), containing 10 mL of culture, were incubated in a shaker

incubator (180 RPM, 37°C). Then cells were harvested by centrifugation (4200 *g* for 5

min) and resuspended in 100 µL of growth medium. As per manufacturer's instructions,

CellROX Deep Red (Invitrogen, cat. No. C10422, Waltham, MA) was added to a final

concentration of 5 µM, and cells were agitated on a rocker (Biobee Tech, Bangalore,

India) for 30 min. After incubation, cells were washed to remove residual dye by

centrifugation (4200 *g* for 5 min). Cells were resuspended in 300 uL phosphate-buffered

saline, pH 7.4) and then fixed by addition of 4% paraformaldehyde (PFA) for 1 h at room

temperature. Fluorescence was measured at a fixed emission (670 nm) after excitation

with a red laser (640 nm) using a BD FACSVerse Flow cytometer (BD Biosciences, San

Jose, CA) with 10,000 events per sample. No autofluorescence was observed.

Determination of DNA damage by TUNEL assay. DNA damage was measured by

using In Situ Cell Death Detection Kit, TMR red (Roche Molecular Biochemicals,

Indianapolis, IN, Cat. No.- 12156792910), which is based on TUNEL (TdT-mediated

dUTP-X nick end labeling) assay (Fan et al., 2018; Vilchèze et al., 2013). An equal

number of cells (based on OD_{600}) were taken at various times of treatment with

moxifloxacin, washed once by centrifugation, and fixed in 2% paraformaldehyde (PFA).

PFA was removed by washing cells, followed by resuspension in 2% sodium dodecyl

sulfate (SDS) and a second wash by centrifugation. DNA double-strand breaks were

labeled in 100 µl of TUNEL reaction mix for 3-4 h. Cells incubated with label solution only

(no terminal transferase) were used as negative controls. Fluorescence was measured

at a fixed emission (585 nm) after excitation with green-yellow laser (561 nm) using a BD

FACSAria Flow cytometer (BD Biosciences, San Jose, CA). 10,000 events were acquired

per sample.

Lipid Peroxidation Assay. Cultures were grown to mid-log phase (OD_{600}= 0.6-0.8).

Lipid hydroperoxides were quantified from cell pellets after centrifugation (4000 *g,* 5 min),

using FOX2 reagent (Nambi et al., 2015). Briefly, cell pellets were resuspended in 1:2

chloroform/methanol and mixed by vortexing. Next, chloroform and water were added at

a 1:1 mixture The samples were then centrifuged to separate the water and organic

phases. The organic phase was collected and washed twice with water. 200 µl of the

organic phase was incubated with 1 ml of FOX2 reagent in the dark for 1 h at 22°C. Lipid

hydroperoxides were measured spectrophotometrically at 560 nm and normalized to

culture turbidity (OD_{600}).

Macrophage preparation and infection by *M. tuberculosis*. The human monocytic

cell line THP-1 was grown in RPMI1640 medium (Cell Clone, Manassas, VA)

supplemented with 10% heat-treated (55°C) fetal bovine serum (MP Biomedical, cat. no.

092916754, Santa Ana, CA). A total of 3×10^5 cells/well was seeded into a 24-well cell-

culture plate. THP-1 monocytes were stimulated to differentiate into macrophage-like

cells by treatment with 20 ng/mL phorbol 12-myristate 13-acetate (PMA; Sigma-Aldrich

Co., St Louis, MO) for 18-20 h at 37°C and incubated further for 48 h to allow

differentiation (Bhaskar et al., 2014). The resulting macrophage-like cells were infected

with *Mtb*-roGFP2 or MDR *M. tuberculosis* NHN1664 expressing Mrx1-roGFP2 at a

multiplicity of infection (MOI) of 10 and incubated for 4 h at 37 °C in 5% CO_2. After

Infection, extracellular bacteria were removed by washing three times (by centrifugation)

with phosphate-buffered saline (PBS; 137 mM NaCl, 2.7 mM KCl, 10 mM Na_2HPO_4, and

1.4 mM KH_2PO_4, pH 7.4). Moxifloxacin (Sigma-Aldrich, cat. No. PHR1542, St Louis, MO),

with or without NAC (Sigma-Aldrich, cat. No. A7250, St Louis, MO), was added to infected

cells that were incubated for various times. For CFU determination, infected cells were

lysed in 7H9 medium containing 0.06% sodium dodecyl sulfate (SDS); dilutions were

prepared using 7H9 medium, and aliquots were plated on 7H11+OADC agar plates

(Bhaskar et al., 2014). Plates were incubated at 37 °C for 3-4 weeks before colonies were

counted.

Nutrient starvation. Starvation was achieved as previously described

(Gengenbacher et al., 2010). Briefly*,* cultures of *M. tuberculosis* H37Rv were grown to

exponential phase in roller-culture bottles (Corning, cat no.- 430518, Corning, NY)

containing Middlebrook 7H9 medium, supplemented with ADS and 0.05 % Tween 80, at

37°C with rolling at 6 RPM in a roller incubator (120 Vac Roll-In Incubator; Wheaton,

Millville, NJ). Cultures grown to OD_{600} ≈ 0.2 were harvested by centrifugation (4000 *g* for

5 min) followed by two washes with phosphate-buffered saline (PBS; pH 7.4)

supplemented with 0.025 % Tween 80 (PBST). Bacterial cells were diluted to a final OD_{600}

of 0.1 in PBS. 50 mL of this suspension was transferred into a roller-culture bottle and

incubated for 14 days to achieve starvation conditions.

Hypoxia. For determination of moxifloxacin lethality under hypoxic conditions,

bacterial cultures (OD_{600} = 0.1) were placed in Vacutainer tubes (Becton Dickinson, cat.

no. 367812, Franklin Lakes, NJ) followed by incubation for 14-21 days at 37°C (Taneja

and Tyagi, 2007). A high cell density (OD_{600} = 0.1) was used for rapid achievement of

hypoxia, which was observed as decolorization of methylene blue (final concentration

was 1.5 μg/mL) in the culture medium. When hypoxia was established, moxifloxacin was

added to cultures anaerobically. Metronidazole at 10 mM and isoniazid at 10 μM were

used as positive and negative controls, respectively. Drugs were injected in volumes of

100 μL in phosphate-buffered saline following passage of argon through the drug solution

to remove residual oxygen. Hypoxic cultures were treated with drugs for five days, similar

to the incubation time for MIC determination with aerobically growing cells. After

treatment, Vacutainer tubes were unsealed, and end-point bacterial survival was

determined by plating on drug-free agar, incubating for 3-4 weeks at 37°C, and visually

enumerating colonies.

Effect of nitrate on survival of hypoxic bacteria. Survival during nitrate-dependent

respiration was achieved by supplementing *M. tuberculosis* cultures with 5 mM sodium

nitrate. Nitrate was added when cultures were placed in Vacutainer tubes before hypoxia.

All other conditions were as described above for hypoxia.

Microarray data analysis. Gene expression data from GSE71200 (Gene Expression

Omnibus, NCBI or GEO) was used for analyzing the response of *M. tuberculosis* to

moxifloxacin (Ma et al., 2015). The data were expressed as a two-channel microarray

with control and drug-exposed *M. tuberculosis* stained with Cy3 and Cy5 dyes,

respectively. We used GSM1829746, GSM1829747, and GSM1829748, which consist of

published data for *M. tuberculosis* exposed to 2X, 4X, or 8X MIC (1X MIC = 0.4 μM) of

moxifloxacin for 16 h. Normalized expression data for GSE71200 was downloaded from

GEO (Barrett et al., 2013; Edgar et al., 2002), and probe IDs were mapped to the respective gene-IDs. The expression levels for genes having multiple probes were averaged, and genes lacking data were removed from further analysis. Differentially expressed genes (DEGs) were defined as genes that were upregulated or downregulated by at least 2-fold in all 3 moxifloxacin treatment conditions.

For overlap analysis of differentially expressed genes in moxifloxacin-exposed bacteria compared with other stress-induced conditions (log_2 fold change >1 or <-1; $p <$ 0.01 are considered differentially expressed genes in stress conditions), GeneOverlap (v1.22.0) package from R (v3.6.3) was used (Core-Team, 2018; Li and Sinai, 2019). It uses Fisher's exact test to find statistical significance by calculating the p-value and the odds ratio for the overlap (p-value < 0.05 and an odds ratio of > 1 were taken as the significance thresholds).

OCR and ECAR measurements. To measure basal oxygen consumption rate (OCR) and extracellular acidification rate (ECAR), log-phase *M. tuberculosis* cultures (OD_{600}= 0.6-0.8) were briefly (one day) incubated in 7H9 medium containing the non-metabolizable detergent tyloxapol (MP Biomedical, cat. No.157162, Santa Ana, CA) and lacking ADS or a carbon source. These cultures were then passed 10 times through a 26-gauge syringe needle followed by centrifugation at 100 g for 1-2 mins to remove clumps of bacterial cells. The resulting single-cell suspensions of bacteria at 2×10^6 cells/well were placed in the bottom of wells of a Cell-Tak (Corning, cat. No. 354240, Corning, NY)-coated XF culture plate (Agilent/Seahorse Biosciences, Santa Clara, CA). Measurements were performed using a Seahorse XFp analyzer (Agilent/Seahorse Biosciences, Santa Clara, CA) with cells in unbuffered 7H9 growth medium (pH 7.35

lacking monopotassium phosphate and disodium phosphate) containing glucose (2

mg/mL) as a carbon source. OCR and ECAR measurements were recorded for ~21 min

(3 initial baseline readings) before addition of moxifloxacin (1X, 2.5X or 10X MIC; 1X MIC

= 0.5 µM), which was delivered automatically through the drug ports of the sensor

cartridge (Wave Software, Agilent Technologies, Santa Clara, CA). NAC or CCCP was

similarly added through drug ports at times indicated in figures. OCR and ECAR were

measured for an additional 6 h in the absence or presence of moxifloxacin and/or NAC.

Changes in OCR and ECAR readings triggered by moxifloxacin were calculated as a

percentage of the third baseline reading for OCR and ECAR taken prior to drug injection.

ROS measurement under Fe-depletion conditions. Iron starvation was as

described previously (Kurthkoti et al., 2017). Briefly, log-phase *M. tuberculosis* cultures

were grown in minimal medium (0.5% (wt/vol) asparagine, 0.5% (wt/vol) KH_2PO4, 2%

glycerol, 0.05% Tween 80, 10% ADS, 0.5 mg/L of sterile $ZnCl_2$, 0.1 mg/ L of $MnSO_4$, and

40 mg/L of $MgSO_4$) to early stationary phase (OD_{600} of ~1). To remove metal ions, the

minimal medium was treated with 5% Chelex-100 (Bio-Rad, cat. No. 142-2842, Hercules,

CA) for 24 h with gentle agitation. This Fe-depleted culture was diluted further to OD_{600}

of 0.1 in the same medium and allowed to grow to early stationary phase to deplete stored

Fe. The Fe-depleted cells were further treated with 50 µg/ml of freshly prepared

deferoxamine mesylate (Sigma Aldrich, cat. No. D9533, St Louis, MO)-containing minimal

medium for 6 days. The cells were washed and diluted in minimal medium containing 80

µM $FeCl_3$ or 80 µM $FeCl_3$ + 10 mM thiourea for an additional 4 days. ROS was quantified

by CellROX Deep Red dye using Flow cytometry as described in the section titled ROS

measurement using CellROX Deep Red.

Cellular Fe Estimation. Cell-free Fe levels were measured using the ferrozine-based

assay as described previously (Fish, 1988; Vilchèze et al., 2017). Briefly, *M. tuberculosis*

cultures were grown to an OD_{600} of 0.3-0.4 followed by treatment with 0X, 1X, 2.5X, or

10X MIC of moxifloxacin. After 48 h of treatment, cells were harvested and washed twice

with ice-cold PBS. The cell pellets were resuspended in 1 mL of 50 mM NaOH and lysed

using a bead beater. The cell lysate sample (300 µl) was mixed with 10 mM HCl (300 µL)

followed by addition of Fe-detection reagent (6.5 mM Ferrozine, 6.5 mM Neocuproine, 1

M ascorbic acid, and 2.5 M ammonium acetate in water) (90 µl). The reaction mix was

incubated at 37°C for 30 min, and then absorbance at 562 nm was measured. The cellular

Fe concentration was determined by plotting the absorbance values against a standard

curve of $FeCl_3$ concentration gradient and normalized to protein content. Protein

concentration was determined using the Pierce BCA Protein Assay Kit (Thermo Scientific,

cat. no. 23225, Rockford, IL).

Estimation of NAD^+, $NADP^+$, NADH, and NADPH. *M. tuberculosis* H37Rv was grown

to OD_{600} ~0.35 and treated with 1X, 2.5X or 10X MIC of moxifloxacin for 48 h. Pyridine

nucleotide levels were determined by a redox-cycling assay (Chawla et al., 2012; Singh

et al., 2009). Protein concentration of NAD^+ or NADH extracts was determined using the

Pierce BCA Protein Assay Kit (Thermo Scientific, cat. no. 23225, Rockford, IL) to

normalize NAD(P)H and $NAD(P)^+$ concentrations.

Analysis of mixtures containing moxifloxacin plus and minus NAC by thin layer

chromatography (TLC). Thin-layer chromatography (TLC) was performed using silica

gel 60 GF_{254} precoated aluminium backed plates (0.25 mm thickness; Merck, Darmstadt,

Germany), and visualization was accomplished by irradiation with UV light at 254 nm.

Stock solutions of moxifloxacin (10 mM) and NAC (0.2 M, 2 M and 20 M) were prepared

independently in DMSO and phosphate buffer (PB, pH 7.4, 10 mM), respectively. In a

typical incubation, moxifloxacin (100 µL at a final concentration of 2 mM) was

independently reacted with various concentrations of NAC (5 µL, final concentration 2

mM, 20 mM, or 200 mM) in 395 µL PB (pH 7.4, 10 mM). In a control experiment,

moxifloxacin (100 µL, final concentration 2 mM) was added to 400 µL PB (pH 7.4, 10

mM). The mixtures were incubated at 37 ºC on an Eppendorf ThermoMixer Comfort (800

rpm). The reactions were monitored by spotting aliquots (5 µL) from the incubation

mixtures onto the TLC plate at designated times. The solvent system used was MeOH

and $CHCl_3$ (1:9).

Analysis of mixtures containing moxifloxacin plus and minus NAC by NMR.

Deuterated phosphate buffer (PB, 10 mM) was prepared by dissolving monobasic

potassium phosphate (KH_2PO_4, 4 mg) and dibasic potassium phosphate (K_2HPO_4, 12

mg) in deuterated water (D_2O), and the pH was adjusted to 7.4 using 40% (w/w) sodium

deuteroxide solution in D_2O (Sigma-Aldrich, cat. No. 151882, St Louis, MO). Stock

solutions of moxifloxacin (10 mM) and NAC (0.2 M) were prepared in DMSO and

deuterated phosphate buffer (PB, pH 7.4, 10 mM) respectively. In a typical reaction,

moxifloxacin (200 µL, 2 mM final concentration) was incubated with NAC (10 µL, 2 mM

final concentration) in 790 µL deuterated PB (pH 7.4, 10 mM). In a control experiment,

moxifloxacin (200 µL, 2 mM final concentration) was added to 800 µL deuterated PB (pH

7.4, 10 mM). The incubation mixtures were incubated at 37 ºC in an Eppendorf

ThermoMixer Comfort (800 rpm). An aliquot (0.5 mL) of the incubation mixture was taken

after 1 h of incubation, and ^1H NMR and ^{19}F NMR spectra were recorded. ^1H NMR was

acquired with 64 scans on a Jeol 400 MHz spectrometer using deuterated water (D_2O;

Sigma-Aldrich) as an internal standard. Chemical shifts (δ) were reported in ppm

downfield from D_2O (δ = 4.79 ppm) for ^1H NMR. ^{19}F spectra was recorded on a JEOL (376

MHz) using an external reference (α, α, α-trifluorotoluene, δ_F = -63.72 ppm).

Fluorescence-based detection of intracellular levels of moxifloxacin in wild-

type *M. smegmatis*. Stock solutions of moxifloxacin (0.05 mM) and NAC (200 mM) were

prepared in DMSO and phosphate buffer (pH 7.4, 10 mM), respectively. Wild-type *M.*

smegmatis cultures were grown in Middlebrook 7H9 broth supplemented with glycerol

(0.2%) and Tween-80 (0.1%). Exponential-phase cultures of wild-type *M. smegmatis,*

grown to OD_{600} of 0.3 (985 µL), were transferred to 1.5 mL Eppendorf tubes and either

left untreated or treated with NAC (5 µL, final concentration 1 mM) for 1 h prior to addition

of 2x MIC of moxifloxacin (10 µL, final concentration 0.5 µM). Treated cells were

incubated with shaking at 180 rpm in a rotary shaker incubator at 37 °C for 48 h. The cell

suspension was then centrifuged at 9,391 *g* at 4 °C for 15 min, and the pellet was washed

twice with 1x phosphate buffer saline (PBS) and resuspended in 1 mL PBS in a

microcentrifuge tube. The cells were lysed by sonication using a 130-watt ultrasonic

processor (Vx 130W) by stepping a microtip with a 4 min pulse on-time (with 3s ON and

3s OFF pulse, 60% amplitude, 20 kHz frequency) under ice-cold conditions. An aliquot

(100 µL) of whole-cell lysate was withdrawn from the above samples and dispensed into

a 96-well microtiter plate. Fluorescence ascribed to moxifloxacin (λ_{ex} = 289 nm and λ_{em} =

488 nm), recovered in NAC-untreated or-pretreated *M. smegmatis* lysates, was recorded

using an Ensight Multimode Plate Reader (PerkinElmer, India). Readings were collected

from the top with 25 flashes per well and with a focus height adjusted to 9.5 mm.

Intracellular levels of moxifloxacin in _M. smegmatis_ using LC/MS. Stock

solutions of moxifloxacin (0.05 mM) and NAC (200 mM) were prepared as described

above. Whole-cell lysates of wild-type _M. smegmatis_ treated with moxifloxacin (0.5 µM,

2x MIC) plus or minus NAC (1 mM) were prepared as described in the fluorescence-

based method employed for the detection of moxifloxacin. An aliquot (100 µL) of whole-

cell lysate was withdrawn from the incubation mixtures, diluted with methyl alcohol (100

µL), and centrifuged at 9,390 x _g_ at 4 °C for 15 min. The supernatant fluids (50 µL) were

removed, diluted with methyl alcohol (50 µL), and analyzed by LC/MS. All measurements

were performed using the positive ion mode with high-resolution, multiple reaction

monitoring (MRM-HR) analysis with a Sciex X500R quadrupole time-of flight (QTOF)

mass spectrometer fitted with an Exion UHPLC system having a Kinetex 2.6 mm

hydrophilic interaction liquid chromatography (HILIC) column (100 Å particle size, 150

mm length, and 3 mm internal diameter; Phenomenex, Intek Chromasol Pvt. Ltd., India).

Nitrogen was used as the nebulizer gas, with nebulizer pressure set at 50 psi. MS was

calibrated in the positive mode, and samples were analyzed with the following

parameters: Mode: electrospray ionization (ESI), ion source gas 1 = 40 psi, ion source

gas 2 = 50 psi, curtain gas = 30, CAD gas = 7, spray voltage = 5500 V, and temperature

= 500 °C. The MRM-HR mass spectrometry parameters were moxifloxacin (Q1, M + H$^+$)

= 402.18, moxifloxacin-NAC adduct (Q2, M + H$^+$) = 565.21, declustering potential = 80

V, collision energy = 20 V, collision exit potential = 5 V, and accumulation time = 0.24 s.

The LC runs were for 30 min with a gradient of 100% solvent A (0.1% HCOOH in water)

for 5 min, linear gradient of solvent B (acetonitrile 0% to 100%) for 25 min followed by

100% solvent A for 5 min, all at a flow rate of 0.5 mL per min.

Determination of Mutant Prevention Concentration (MPC). MPC of moxifloxacin with *M. tuberculosis* NHN1664 was determined by methods described previously (Dong et al., 2000; Singh et al., 2017). Cultures of MDR strain *M. tuberculosis* NHN1664 were grown to OD_{600} 0.6-0.7. Approximately 2.5×10^9 bacilli were plated on 7H11 agar containing either moxifloxacin (2X, 4X, or 8X MIC) alone or in combination with NAC at either 1 mM or 2 mM. Resistant colonies were enumerated after incubation at 37°C for 8 weeks. The drug-resistant phenotype was confirmed by replating on drug-containing 7H11-agar plates. MPC was determined as the lowest concentration of drug that prevented bacterial colony formation when $>2 \times 10^9$ bacteria were plated on drug-containing 7H11 plates. Mutation frequency with moxifloxacin was calculated as the number of mutants appearing on drug-containing plates divided by the viable input bacteria.

Determination of drug-tolerant *M. tuberculosis* population *ex vivo*. Murine bone-marrow-derived macrophages (BMDMs) were infected with Mrx1-roGFP2-expressing *M. tuberculosis* H37Rv at a multiplicity of infection of 10. After washing off extracellular bacteria, infected macrophages were left untreated for 24 h for the emergence of the drug-tolerant population. At 24 h post-infection, macrophages harbouring E_{MSH}-reduced and E_{MSH}-basal *M. tuberculosis* were sorted using BD FACS Aria™ Fusion Flow Cytometer. The sorted, infected BMDM cells were seeded into 96-well plates followed by treatment with 3X MIC of moxifloxacin (1X MIC = 0.5 µM) for 48 h. After treatment, cells were lysed with 0.05% sodium dodecyl sulfate (SDS) in 7H9 medium, serially diluted, and plated on 7H11-OADC agar plates. Plates were incubated at 37°C for 3-4 weeks before colony forming units (CFUs) were enumerated.

***In vivo* drug efficacy.** All animal studies were executed as per guidelines prescribed

by the Committee for the Purpose of Control and Supervision of Experiments on Animals,

Government of India, with approval from the Institutional Animal Ethical Committee and

Biosafety Level-3 Committee. 6- to 8-week-old female pathogen-free BALB/c mice were

infected via a low-dose aerosol exposure to the *M. tuberculosis* MDR strain NHN1664

using a Madison chamber aerosol generation instrument. The short-course mouse model

of infection was performed as described previously (Lenaerts et al., 2003). At day one

post infection, three mice were sacrificed to verify implantation of ~100 CFU of bacteria

per mouse. Mice were randomly divided into 4 groups (n = 5 per group). Feed and water

were given *ad libitum*. Treatment with moxifloxacin and/or NAC started 21 days post-

infection and continued for 10 days; untreated control mice received water rather than the

two test compounds. Treated mice were administered moxifloxacin (50 mg/kg of body

weight), NAC (500 mg/ kg of body weight) and a combination of moxifloxacin and NAC at

50 mg/kg and 500 mg/kg body weight, respectively, by oral gavage, once daily. Five

infected mice were sacrificed at the start of treatment as pretreatment controls. After

treatment, mice were sacrificed, and the lungs and spleen were harvested for

measurement of bacterial burden. CFUs were determined by plating appropriate serial

dilutions on 7H11 (supplemented with OADC) agar plates and counting visible colonies

after 3–4 weeks of incubation at 37°C. Data were normalized to whole organ.

Detection of moxifloxacin-resistant mutants in murine model of infection. 6- to

8-week-old female, pathogen-free BALB/c mice were infected via a high-dose aerosol

exposure to the *M. tuberculosis* MDR strain NHN1664 using a Madison chamber aerosol

generation instrument. At day one post infection, three mice were sacrificed to determine

implantation of $>10^3$ CFU of bacteria per mouse. Mice were then randomly divided into 4 treatment groups, with 6 mice each in the vehicle control group, NAC-only group, groups receiving either moxifloxacin or a combination of NAC and moxifloxacin. Feed and water were given *ad libitum*. Treatment with moxifloxacin and/or NAC started 14 days post-infection and continued for 14 days; untreated control mice received water rather than the two test compounds. Treated mice were administered moxifloxacin (50 mg/kg of body weight), NAC (500 mg/ kg of body weight) and a combination of moxifloxacin and NAC at 50 mg/kg and 500 mg/kg body weight, respectively, by oral gavage, once daily. 5 infected mice were sacrificed at the start of treatment as pre-treatment controls. After treatment, mice were sacrificed, and the lungs and spleen were harvested together. Moxifloxacin-resistant mutants were selected by plating the undiluted homogenates (lung+spleen) on five 7H11 plates containing 0.5 µM of moxifloxacin (4X MIC). Plates were incubated for 3–4 weeks at 37˚C.

Statistical analysis. Statistical analysis was performed using GraphPad Prism version 8.4.3 software. Mean and standard deviation values were plotted as indicated in figure legends. A *p*-value of less than 0.05 was considered significant. Statistical significance was determined by unpaired two-tailed student's t-test; either one-way or two-way ANOVA was performed where comparison of multiple groups was made.

Acknowledgements

We thank the following for critical comments on the manuscript: Lanbo Shi, Bo Shopsin, and Xilin Zhao.

Declaration

1065

1066

1067

1068

1069

1070

1071

References

Almeida, D., E. Nuermberger, S. Tyagi, W.R. Bishai, and J. Grosset. 2007. *In vivo* validation of the mutant selection window hypothesis with moxifloxacin in a murine model of tuberculosis. *Antimicrob. Agents Chemother.* 51:4261-4266.

Amaral, E.P., E.L. Conceição, D.L. Costa, M.S. Rocha, J.M. Marinho, M. Cordeiro-Santos, M.R. D'Império-Lima, T. Barbosa, A. Sher, and B.B. Andrade. 2016. N-acetyl-cysteine exhibits potent anti-mycobacterial activity in addition to its known anti-oxidative functions. *BMC Microbiol.* 16:251.

Anand, K., A. Tripathi, K. Shukla, N. Malhotra, A.K. Jamithireddy, R.K. Jha, S.N. Chaudhury, R.S. Rajmani, A. Ramesh, V. Nagaraja, B. Gopal, G. Nagaraju, A.S. Narain Seshayee, and A. Singh. 2021. *Mycobacterium tuberculosis* SufR responds to nitric oxide via its 4Fe–4S cluster and regulates Fe–S cluster biogenesis for persistence in mice. *Redox Biol.* 46:102062.

Baniasadi, S., P. Eftekhari, P. Tabarsi, F. Fahimi, M.R. Raoufy, M.R. Masjedi, and A.A. Velayati. 2010. Protective effect of N-acetylcysteine on antituberculosis drug-induced hepatotoxicity. *Eur. J. Gastroenterol. Hepatol.* 22:1235-1238.

Barrett, T., S.E. Wilhite, P. Ledoux, C. Evangelista, I.F. Kim, M. Tomashevsky, K.A. Marshall, K.H. Phillippy, P.M. Sherman, M. Holko, A. Yefanov, H. Lee, N. Zhang, C.L. Robertson, N. Serova, S. Davis, and A. Soboleva. 2013. NCBI GEO: archive for functional genomics data sets--update. *Nucleic Acids Research* 41:D991-D995.

Bhaskar, A., M. Chawla, M. Mehta, P. Parikh, P. Chandra, D. Bhave, D. Kumar, K.S. Carroll, and A. Singh. 2014. Reengineering redox sensitive GFP to measure mycothiol redox potential of *Mycobacterium tuberculosis* during infection. *PLoS Pathog.* 10:e1003902.

Bhat, S.A., I.K. Iqbal, and A. Kumar. 2016. Imaging the NADH:NAD+ homeostasis for

understanding the metabolic response of mycobacterium to physiologically

relevant stresses. *Frontiers in Cellular and Infection Microbiology* 6:

Cao, R., G. Teskey, H. Islamoglu, R. Abrahem, S. Munjal, K. Gyurjian, L. Zhong, and V.

Venketaraman. 2018. Characterizing the effects of glutathione as an

immunoadjuvant in the treatment of tuberculosis. *Antimicrob. Agents Chemother.*

62:e01132-01118.

Chawla, M., P. Parikh, A. Saxena, M. Munshi, M. Mehta, D. Mai, A.K. Srivastava, K.V.

Narasimhulu, K.E. Redding, N. Vashi, D. Kumar, A.J.C. Steyn, and A. Singh. 2012.

Mycobacterium tuberculosis WhiB4 regulates oxidative stress response to

modulate survival and dissemination *in vivo*. *Mol. Microbiol.* 85:1148-1165.

Core-Team, R. 2018. A language and environment for statistical computing. R foundation

for statistical computing. Vienna, Austria. In.

Cui, J., Y. Liu, R. Wang, W. Tong, K. Drlica, and X. Zhao. 2006. The mutant selection

window in rabbits infected with *Staphylococcus aureus*. *J. Infect. Dis.* 194:1601-

1608.

Cunningham-Bussel, A., T. Zhang, and C.F. Nathan. 2013. Nitrite produced by

Mycobacterium tuberculosis in human macrophages in physiologic oxygen

impacts bacterial ATP consumption and gene expression. *Proc. Natl. Acad. Sci.*

U.S.A. 110:E4256-4265.

Das, M., A. Dewan, S. Shee, and A. Singh. 2021. The multifaceted bacterial cysteine

desulfurases: From metabolism to pathogenesis. *Antioxidants (Basel)* 10:

Davies Forsman, L., K. Niward, J. Kuhlin, X. Zheng, R. Zheng, R. Ke, C. Hong, J.

Werngren, J. Paues, U.S.H. Simonsson, E. Eliasson, S. Hoffner, B. Xu, J.-W.

Alffenaar, T. Schön, Y. Hu, and J. Bruchfeld. 2020. Suboptimal moxifloxacin and

levofloxacin drug exposure during treatment of patients with multidrug-resistant

tuberculosis: results from a prospective study in China. *Eur. Respir. J.* 57:2003463.

Dong, Y., C. Xu, X. Zhao, J. Domagala, and K. Drlica. 1998. Fluoroquinolone action

against mycobacteria: Effects of C-8 substituents on growth, survival, and

resistance. *Antimicrob. Agents Chemother.* 42:2978-2984.

Dong, Y., X. Zhao, B.N. Kreiswirth, and K. Drlica. 2000. Mutant prevention concentration

as a measure of antibiotic potency: studies with clinical isolates of *Mycobacterium*

tuberculosis. *Antimicrob. Agents Chemother.* 44:2581-2584.

Dorman, S.E., P. Nahid, E.V. Kurbatova, P.P.J. Phillips, K. Bryant, K.E. Dooley, M. Engle,

S.V. Goldberg, H.T.T. Phan, J. Hakim, J.L. Johnson, M. Lourens, N.A. Martinson,

G. Muzanyi, K. Narunsky, S. Nerette, N.V. Nguyen, T.H. Pham, S. Pierre, A.E.

Purfield, W. Samaneka, R.M. Savic, I. Sanne, N.A. Scott, J. Shenje, E. Sizemore,

A. Vernon, Z. Waja, M. Weiner, S. Swindells, and R.E. Chaisson. 2021. Four-

month rifapentine regimens with or without moxifloxacin for tuberculosis. *N. Engl.*

J. Med. 384:1705-1718.

Dorsey-Oresto, A., T. Lu, M. Mosel, X. Wang, T. Salz, K. Drlica, and X. Zhao. 2013. *YihE*

kinase is a central regulator of programmed cell death in bacteria. *Cell Reports*

3:528-537.

Drlica, K., and X. Zhao. 2007. Mutant selection window hypothesis updated. *Clin. Infect.*

Dis. 44:681-688.

Drlica, K., and X. Zhao. 2020. Bacterial death from treatment with fluoroquinolones and

other lethal stressors. *Expert Rev. Anti Infect. Ther.* 1-18.

Dwyer, D.J., P.A. Belenky, J.H. Yang, I.C. MacDonald, J.D. Martell, N. Takahashi, C.T.Y.

Chan, M.A. Lobritz, D. Braff, E.G. Schwarz, J.D. Ye, M. Pati, M. Vercruysse, P.S.

Ralifo, K.R. Allison, A.S. Khalil, A.Y. Ting, G.C. Walker, and J.J. Collins. 2014.

Antibiotics induce redox-related physiological alterations as part of their lethality.

Proc. Natl. Acad. Sci. U.S.A. 111:E2100-E2109.

Dwyer, D.J., J.J. Collins, and G.C. Walker. 2015. Unraveling the physiological

complexities of antibiotic lethality. *Annu. Rev. Pharmacol. Toxicol.* 55:313-332.

Dwyer, D.J., M.A. Kohanski, B. Hayete, and J.J. Collins. 2007. Gyrase inhibitors induce

an oxidative damage cellular death pathway in *Escherichia coli*. *Mol. Syst. Biol.*

3:91-91.

Edgar, R., M. Domrachev, and A.E. Lash. 2002. Gene Expression Omnibus: NCBI gene

expression and hybridization array data repository. *Nucleic Acids Research*

30:207-210.

Ehrt, S., and D. Schnappinger. 2009. Mycobacterial survival strategies in the phagosome:

defence against host stresses. *Cellular Microbiology* 11:1170-1178.

Ezeriņa, D., Y. Takano, K. Hanaoka, Y. Urano, and T.P. Dick. 2018. N-Acetyl Cysteine

functions as a fast-acting antioxidant by triggering intracellular H_2S and sulfane

sulfur production. *Cell Chem. Biol.* 25:447-459.e444.

Fan, X.Y., B.K. Tang, Y.Y. Xu, A.X. Han, K.X. Shi, Y.K. Wu, Y. Ye, M.L. Wei, C. Niu, K.W.

Wong, G.P. Zhao, and L.D. Lyu. 2018. Oxidation of dCTP contributes to antibiotic

lethality in stationary-phase mycobacteria. *Proc. Natl. Acad. Sci. U.S.A.* 201719627.

Firsov, A.A., S.N. Vostrov, I.Y. Lubenko, K. Drlica, Y.A. Portnoy, and S.H. Zinner. 2003. In vitro pharmacodynamic evaluation of the mutant selection window hypothesis using four fluoroquinolones against *Staphylococcus aureus*. *Antimicrob. Agents Chemother.* 47:1604-1613.

Fish, W.W. 1988. Rapid colorimetric micromethod for the quantitation of complexed iron in biological samples. *Methods Enzymol.* 158:357-364.

Foti, J.J., B. Devadoss, J.A. Winkler, J.J. Collins, and G.C. Walker. 2012. Oxidation of the guanine nucleotide pool underlies cell death by bactericidal antibiotics. *Science* 336:315-319.

Gengenbacher, M., S.P. Rao, K. Pethe, and T. Dick. 2010. Nutrient-starved, non-replicating *Mycobacterium tuberculosis* requires respiration, ATP synthase and isocitrate lyase for maintenance of ATP homeostasis and viability. *Microbiology (Reading)* 156:81-87.

Gillespie, S.H., A.M. Crook, T.D. McHugh, C.M. Mendel, S.K. Meredith, S.R. Murray, F. Pappas, P.P.J. Phillips, and A.J. Nunn. 2014. Four-month moxifloxacin-based regimens for drug-sensitive tuberculosis. *N. Engl. J. Med.* 371:1577-1587.

Ginsburg, A.S., R. Sun, H. Calamita, C.P. Scott, W.R. Bishai, and J.H. Grosset. 2005. Emergence of fluoroquinolone resistance in *Mycobacterium tuberculosis* during continuously dosed moxifloxacin monotherapy in a mouse model. *Antimicrob. Agents Chemother.* 49:3977-3979.

Gumbo, T., A. Louie, M.R. Deziel, L.M. Parsons, M. Salfinger, and G.L. Drusano. 2004.

Selection of a moxifloxacin dose that suppresses drug resistance in

Mycobacterium tuberculosis, by use of an in vitro pharmacodynamic infection

model and mathematical modeling. *J. Infect. Dis.* 190:1642-1651.

Held, K.D., and J.E. Biaglow. 1994. Mechanisms for the oxygen radical-mediated toxicity

of various thiol-containing compounds in cultured mammalian cells. *Radiat. Res.*

139:15-23.

Hong, Y., L. Li, G. Luan, K. Drlica, and X. Zhao. 2017. Contribution of reactive oxygen

species to thymineless death in *Escherichia coli*. *Nat. Microbiol.* 2:1667-1675.

Hong, Y., Q. Li, Q. Gao, J. Xie, H. Huang, K. Drlica, and X. Zhao. 2020. Reactive oxygen

species play a dominant role in all pathways of rapid quinolone-mediated killing. *J.*

Antimicrob. Chemother. 75:576-585.

Hong, Y., J. Zeng, X. Wang, K. Drlica, and X. Zhao. 2019. Post-stress bacterial cell death

mediated by reactive oxygen species. *Proc. Natl. Acad. Sci. U.S.A.* 116:10064-

10071.

Imlay, J.A. 2013. The molecular mechanisms and physiological consequences of

oxidative stress: lessons from a model bacterium. *Nat. Rev. Microbiol.* 11:443-454.

Imlay, J.A., and S. Linn. 1986. Bimodal pattern of killing of DNA-repair-defective or

anoxically grown Escherichia coli by hydrogen peroxide. *J Bacteriol* 166:519-527.

Jaeschke, H., C. Kleinwaechter, and A. Wendel. 1992. NADH-Dependent reductive stress

and ferritin-bound iron in allyl alcohol-induced lipid peroxidation *in vivo*: The

protective effect of vitamin E. *Chem. Biol. Interact.* 81:57-68.

Jawahar, M.S., V.V. Banurekha, C.N. Paramasivan, F. Rahman, R. Ramachandran, P.

Venkatesan, R. Balasubramanian, N. Selvakumar, C. Ponnuraja, A.S. Iliayas, N.P.

Gangadevi, B. Raman, D. Baskaran, S.R. Kumar, M.M. Kumar, V. Mohan, S.

Ganapathy, V. Kumar, G. Shanmugam, N. Charles, M.R. Sakthivel, K. Jagannath,

C. Chandrasekar, R.T. Parthasarathy, and P.R. Narayanan. 2013. Randomized

clinical trial of thrice-weekly 4-month moxifloxacin or gatifloxacin containing

regimens in the treatment of new sputum positive pulmonary tuberculosis patients.

PLoS One 8:e67030.

Jaworska, M., G. Szulińska, M. Wilk, and J. Tautt. 1999. Capillary electrophoretic

separation of N-acetylcysteine and its impurities as a method for quality control of

pharmaceuticals. *J. Chromatogr. A* 853:479-485.

Jindani, A., T.S. Harrison, A.J. Nunn, P.P.J. Phillips, G.J. Churchyard, S. Charalambous,

M. Hatherill, H. Geldenhuys, H.M. McIlleron, S.P. Zvada, S. Mungofa, N.A. Shah,

S. Zizhou, L. Magweta, J. Shepherd, S. Nyirenda, J.H. van Dijk, H.E. Clouting, D.

Coleman, A.L.E. Bateson, T.D. McHugh, P.D. Butcher, and D.A. Mitchison. 2014.

High-dose rifapentine with moxifloxacin for pulmonary tuberculosis. *N. Engl. J.*

Med. 371:1599-1608.

Kapopoulou, A., J.M. Lew, and S.T. Cole. 2011. The MycoBrowser portal: a

comprehensive and manually annotated resource for mycobacterial genomes.

Tuberculosis (Edinb) 91:8-13.

Kareyeva, A.V., V.G. Grivennikova, and A.D. Vinogradov. 2012. Mitochondrial hydrogen

peroxide production as determined by the pyridine nucleotide pool and its redox

state. *Biophys. Acta Bioenerg.* 1817:1879-1885.

Keyer, K., and J.A. Imlay. 1996. Superoxide accelerates DNA damage by elevating free-iron levels. *Proc. Natl. Acad. Sci. U.S.A.* 93:13635-13640.

Kohanski, M.A., D.J. Dwyer, B. Hayete, C.A. Lawrence, and J.J. Collins. 2007. A common mechanism of cellular death induced by bactericidal antibiotics. *Cell* 130:797-810.

Kranzer, K., W.F. Elamin, H. Cox, J.A. Seddon, N. Ford, and F. Drobniewski. 2015. A systematic review and meta-analysis of the efficacy and safety of N-acetylcysteine in preventing aminoglycoside-induced ototoxicity: implications for the treatment of multidrug-resistant TB. *Thorax* 70:1070-1077.

Kurthkoti, K., H. Amin, M.J. Marakalala, S. Ghanny, S. Subbian, A. Sakatos, J. Livny, S.M. Fortune, M. Berney, and G.M. Rodriguez. 2017. The capacity of *Mycobacterium tuberculosis* to survive iron starvation might enable it to persist in iron-deprived microenvironments of human granulomas. *mBio* 8:

Lamprecht, D.A., P.M. Finin, M.A. Rahman, B.M. Cumming, S.L. Russell, S.R. Jonnala, J.H. Adamson, and A.J.C. Steyn. 2016. Turning the respiratory flexibility of *Mycobacterium tuberculosis* against itself. *Nat. Comm.* 7:12393.

Lange, C., D. Chesov, J. Heyckendorf, C.C. Leung, Z. Udwadia, and K. Dheda. 2018. Drug-resistant tuberculosis: An update on disease burden, diagnosis and treatment. *Respirology* 23:656-673.

Lee, B.S., N.P. Kalia, X.E.F. Jin, E.J. Hasenoehrl, M. Berney, and K. Pethe. 2019. Inhibitors of energy metabolism interfere with antibiotic-induced death in mycobacteria. *J. Biol. Chem.* 294:1936-1943.

Lenaerts, A.J., V. Gruppo, J.V. Brooks, and I.M. Orme. 2003. Rapid *in vivo* screening of

experimental drugs for tuberculosis using gamma interferon gene-disrupted mice.

Antimicrob. Agents Chemother. 47:783-785.

Li, L., Y. Hong, G. Luan, M. Mosel, M. Malik, K. Drlica, and X. Zhao. 2014. Ribosomal

elongation factor 4 promotes cell death associated with lethal stress. *mBio*

5:e01708.

Li, S., and M. Sinai. 2019. GeneOverlap: Test and visualize gene overlaps. In.

Liu, Y., and J.A. Imlay. 2013. Cell death from antibiotics without the involvement of

reactive oxygen species. *Science* 339:1210-1213.

Liu, Y., X. Liu, Y. Qu, X. Wang, L. Li, and X. Zhao. 2012. Inhibitors of reactive oxygen

species accumulation delay and/or reduce the lethality of several

antistaphylococcal agents. *Antimicrob. Agents Chemother.* 56:6048-6050.

Lobritz, M.A., P. Belenky, C.B.M. Porter, A. Gutierrez, J.H. Yang, E.G. Schwarz, D.J.

Dwyer, A.S. Khalil, and J.J. Collins. 2015. Antibiotic efficacy is linked to bacterial

cellular respiration. *Proc. Natl. Acad. Sci. U.S.A.* 112:8173.

Luan, G., Y. Hong, K. Drlica, and X. Zhao. 2018. Suppression of reactive oxygen species

accumulation accounts for paradoxical bacterial survival at high quinolone

concentration. *Antimicrob. Agents Chemother.* 62:e01622-01617.

Ma, S., K.J. Minch, T.R. Rustad, S. Hobbs, S.L. Zhou, D.R. Sherman, and N.D. Price.

2015. Integrated Modeling of Gene Regulatory and Metabolic Networks in

Mycobacterium tuberculosis. *PLoS Comput. Biol.* 11:e1004543.

Mahakalkar, S.M., D. Nagrale, S. Gaur, C. Urade, B. Murhar, and A. Turankar. 2017. N-

acetylcysteine as an add-on to Directly Observed Therapy Short-I therapy in fresh

pulmonary tuberculosis patients: A randomized, placebo-controlled, double-blinded study. *Perspect Clin. Res.* 8:132-136.

Malik, M., S. Hussain, and K. Drlica. 2007. Effect of anaerobic growth on quinolone lethality with *Escherichia coli*. *Antimicrob. Agents Chemother.* 51:28-34.

Malik, M., X. Zhao, and K. Drlica. 2006. Lethal fragmentation of bacterial chromosomes mediated by DNA gyrase and quinolones. *Mol. Microbiol.* 61:810-825.

Mavi, P.S., S. Singh, and A. Kumar. 2019. Reductive stress: New insights in physiology and drug tolerance of mycobacterium. *Antioxid. Redox Signal.* 32:1348-1366.

Mishra, R., S. Kohli, N. Malhotra, P. Bandyopadhyay, M. Mehta, M. Munshi, V. Adiga, V.K. Ahuja, R.K. Shandil, R.S. Rajmani, A.S.N. Seshasayee, and A. Singh. 2019. Targeting redox heterogeneity to counteract drug tolerance in replicating *Mycobacterium tuberculosis*. *Sci. Transl. Med.* 11:eaaw6635.

Mishra, R., V. Yadav, M. Guha, and A. Singh. 2021. Heterogeneous host–pathogen encounters coordinate antibiotic resilience in *Mycobacterium tuberculosis*. *Trends in Microbiology* 29:606-620.

Mishra, S., P. Shukla, A. Bhaskar, K. Anand, P. Baloni, R.K. Jha, A. Mohan, R.S. Rajmani, V. Nagaraja, N. Chandra, and A. Singh. 2017. Efficacy of β-lactam/β-lactamase inhibitor combination is linked to WhiB4-mediated changes in redox physiology of *Mycobacterium tuberculosis*. *eLife* 6:

Mosel, M., L. Li, K. Drlica, and X. Zhao. 2013. Superoxide-mediated protection of *Escherichia coli* from antimicrobials. *Antimicrob. Agents Chemother.* 57:5755-5759.

Nagrale, D., S. Mahakalkar, and S. Gaur. 2013. Supplementation of N-acetylcysteine as

an adjuvant in treatment of newly diagnosed pulmonary tuberculosis patients: A

prospective, randomized double blind, placebo controlled study. *Eur. Respir. J.*

42:P2833.

Nambi, S., J.E. Long, B.B. Mishra, R. Baker, K.C. Murphy, A.J. Olive, H.P. Nguyen, S.A.

Shaffer, and C.M. Sassetti. 2015. The oxidative stress network of *Mycobacterium*

tuberculosis reveals coordination between radical detoxification systems. *Cell Host*

Microbe 17:829-837.

Nandakumar, M., C. Nathan, and K.Y. Rhee. 2014. Isocitrate lyase mediates broad

antibiotic tolerance in *Mycobacterium tuberculosis*. *Nature Communications*

5:4306.

Nuermberger, E.L., T. Yoshimatsu, S. Tyagi, R.J. O'Brien, A.N. Vernon, R.E. Chaisson,

W.R. Bishai, and J.H. Grosset. 2004. Moxifloxacin-containing regimen greatly

reduces time to culture conversion in murine tuberculosis. *Am. J. Respir. Crit. Care*

Med. 169:421-426.

Padiadpu, J., P. Baloni, K. Anand, M. Munshi, C. Thakur, A. Mohan, A. Singh, and N.

Chandra. 2016. Identifying and tackling emergent vulnerability in drug-resistant

mycobacteria. *ACS Infectious Diseases* 2:592-607.

Palanisamy, G.S., N.M. Kirk, D.F. Ackart, C.A. Shanley, I.M. Orme, and R.J. Basaraba.

2011. Evidence for oxidative stress and defective antioxidant response in guinea

pigs with tuberculosis. *PLoS One* 6:e26254.

Pletz, M.W.R., A. De Roux, A. Roth, K.-H. Neumann, H. Mauch, and H. Lode. 2004. Early

bactericidal activity of moxifloxacin in treatment of pulmonary tuberculosis: A

prospective, randomized study. *Antimicrob. Agents Chemother.* 48:780.

Price-Whelan, A., L.E.P. Dietrich, and D.K. Newman. 2007. Pyocyanin alters redox

homeostasis and carbon flux through central metabolic pathways in *Pseudomonas*

aeruginosa PA14. *J. Bacteriol.* 189:6372-6381.

Prideaux, B., L.E. Via, M.D. Zimmerman, S. Eum, J. Sarathy, P. O'Brien, C. Chen, F.

Kaya, D.M. Weiner, P.-Y. Chen, T. Song, M. Lee, T.S. Shim, J.S. Cho, W. Kim,

S.N. Cho, K.N. Olivier, C.E. Barry, and V. Dartois. 2015. The association between

sterilizing activity and drug distribution into tuberculosis lesions. *Nat. Med.*

21:1223-1227.

Richardson, A.R., S.J. Libby, and F.C. Fang. 2008. A nitric oxide-inducible lactate

dehydrogenase enables *Staphylococcus aureus* to resist innate immunity. *Science*

319:1672.

Rodríguez-Rojas, A., J.J. Kim, P.R. Johnston, O. Makarova, M. Eravci, C. Weise, R.

Hengge, and J. Rolff. 2020. Non-lethal exposure to H_2O_2 boosts bacterial survival

and evolvability against oxidative stress. *PLoS Genet.* 16:e1008649.

Rodríguez, J.C., M. Ruiz, A. Climent, and G. Royo. 2001. *In vitro* activity of four

fluoroquinolones against *Mycobacterium tuberculosis*. *Int. J. Antimicrob. Agents*

17:229-231.

Ruan, Q., Q. Liu, F. Sun, L. Shao, J. Jin, S. Yu, J. Ai, B. Zhang, and W. Zhang. 2016.

Moxifloxacin and gatifloxacin for initial therapy of tuberculosis: a meta-analysis of

randomized clinical trials. *Emerg. Microbes Infect.* 5:e12-e12.

Saini, V., Bridgette M. Cumming, L. Guidry, D.A. Lamprecht, John H. Adamson, Vineel P.

Reddy, Krishna C. Chinta, J.H. Mazorodze, Joel N. Glasgow, M. Richard-

Greenblatt, A. Gomez-Velasco, H. Bach, Y. Av-Gay, H. Eoh, K. Rhee, and

Adrie J.C. Steyn. 2016. Ergothioneine maintains redox and bioenergetic

homeostasis essential for drug susceptibility and virulence of *Mycobacterium*

tuberculosis. *Cell Rep.* 14:572-585.

Sarathy, J., L. Blanc, N. Alvarez-Cabrera, P. O'Brien, I. Dias-Freedman, M. Mina, M.

Zimmerman, F. Kaya, H.P. Ho Liang, B. Prideaux, J. Dietzold, P. Salgame, R.M.

Savic, J. Linderman, D. Kirschner, E. Pienaar, and V. Dartois. 2019.

Fluoroquinolone efficacy against tuberculosis is driven by penetration into lesions

and activity against resident bacterial populations. *Antimicrob Agents Chemother*

63:e02516-02518.

Seaver, L.C., and J.A. Imlay. 2001. Hydrogen peroxide fluxes and compartmentalization

inside growing *Escherichia coli*. *J. Bacteriol.* 183:7182-7189.

Shandil, R.K., R. Jayaram, P. Kaur, S. Gaonkar, B.L. Suresh, B.N. Mahesh, R. Jayashree,

V. Nandi, S. Bharath, and V. Balasubramanian. 2007. Moxifloxacin, ofloxacin,

sparfloxacin, and ciprofloxacin against *Mycobacterium tuberculosis*: evaluation of

in vitro and pharmacodynamic indices that best predict *in vivo* efficacy. *Antimicrob.*

Agents Chemother. 51:576.

Shapleigh, J.P. 2009. Dissimilatory and assimilatory nitrate reduction in the purple

photosynthetic bacteria. In The purple phototrophic bacteria. Advances in

photosynthesis and respiration. D.F. Hunter C.N., Thurnauer M.C., Beatty J.T.,

editor Springer, Dordrecht, 623-642.

Singh, A., D.K. Crossman, D. Mai, L. Guidry, M.I. Voskuil, M.B. Renfrow, and A.J.C.

Steyn. 2009. *Mycobacterium tuberculosis* WhiB3 maintains redox homeostasis by

regulating virulence lipid anabolism to modulate macrophage response. *PLoS*

Pathog. 5:e1000545.

Singh, A., D. Mai, A. Kumar, and A.J.C. Steyn. 2006. Dissecting virulence pathways of

Mycobacterium tuberculosis through protein–protein association. *Proc. Natl. Acad.*

Sci. U.S.A. 103:11346.

Singh, S., N.P. Kalia, P. Joshi, A. Kumar, P.R. Sharma, A. Kumar, S.B. Bharate, and I.A.

Khan. 2017. Boeravinone B, a novel dual inhibitor of NorA bacterial efflux pump of

Staphylococcus aureus and human p-glycoprotein, reduces the biofilm formation

and intracellular invasion of bacteria. *Front. Microbiol.* 8:1868.

Sohaskey, C.D. 2008. Nitrate enhances the survival of *Mycobacterium tuberculosis*

during inhibition of respiration. *J. Bacteriol.* 190:2981-2986.

Sommer, I., H. Schwebel, V. Adamo, P. Bonnabry, L. Bouchoud, and F. Sadeghipour.

2020. Stability of N-Acetylcysteine (NAC) in standardized pediatric parenteral

nutrition and evaluation of N,N-Diacetylcystine (DAC) formation. *Nutrients* 12:

Srinivas, V., M.L. Arrieta-Ortiz, A. Kaur, E.J.R. Peterson, and N.S. Baliga. 2020. PerSort

facilitates characterization and elimination of persister subpopulation in

mycobacteria. *m Systems* 5:e01127-01120.

Steyn, A.J.C., J. Joseph, and B.R. Bloom. 2003. Interaction of the sensor module of

Mycobacterium tuberculosis H37Rv KdpD with members of the Lpr family. *Mol.*

Microbiol. 47:1075-1089.

Tan, M.P., P. Sequeira, W.W. Lin, W.Y. Phong, P. Cliff, S.H. Ng, B.H. Lee, L. Camacho,

D. Schnappinger, S. Ehrt, T. Dick, K. Pethe, and S. Alonso. 2010. Nitrate

respiration protects hypoxic *Mycobacterium tuberculosis* against acid and reactive

nitrogen species stresses. *PLoS One* 5:e13356.

Taneja, N.K., and J.S. Tyagi. 2007. Resazurin reduction assays for screening of anti-

tubercular compounds against dormant and actively growing *Mycobacterium*

tuberculosis, *Mycobacterium bovis* BCG and *Mycobacterium smegmatis*. *J.*

Antimicrob. Chemother. 60:288-293.

Titov, D.V., V. Cracan, R.P. Goodman, J. Peng, Z. Grabarek, and V.K. Mootha. 2016.

Complementation of mitochondrial electron transport chain by manipulation of the

NAD+/NADH ratio. *Science* 352:231-235.

Vilchèze, C., T. Hartman, B. Weinrick, and W.R. Jacobs. 2013. Mycobacterium

tuberculosis is extraordinarily sensitive to killing by a vitamin C-induced Fenton

reaction. *Nature Communications* 4:1881.

Vilchèze, C., T. Hartman, B. Weinrick, P. Jain, T.R. Weisbrod, L.W. Leung, J.S.

Freundlich, and W.R. Jacobs. 2017. Enhanced respiration prevents drug tolerance

and drug resistance in *Mycobacterium tuberculosis*. *Proc. Natl. Acad. Sci. U.S.A.*

114:4495-4500.

Vilchèze, C., and W.R. Jacobs. 2021. The promises and limitations of N- acetylcysteine

as a potentiator of first-line and second-line tuberculosis drugs. *Antimicrob. Agents*

Chemother. AAC.01703-01720.

Vilchèze, C., T.R. Weisbrod, B. Chen, L. Kremer, M.H. Hazbón, F. Wang, D. Alland, J.C.

Sacchettini, and W.R. Jacobs, Jr. 2005. Altered NADH/NAD+ ratio mediates

coresistance to isoniazid and ethionamide in mycobacteria. *Antimicrob. Agents Chemother.* 49:708-720.

Vinogradov, A.D., and V.G. Grivennikova. 2016. Oxidation of NADH and ROS production by respiratory complex I. *Biophys. Acta Bioenerg.* 1857:863-871.

Voskuil, M., I. Bartek, K. Visconti, and G. Schoolnik. 2011. The response of *Mycobacterium tuberculosis* to reactive oxygen and nitrogen species. *Front. Microbiol.* 2:1-12.

Voskuil , M.I., D. Schnappinger , K.C. Visconti , M.I. Harrell , G.M. Dolganov , D.R. Sherman , and G. Schoolnik. 2003. Inhibition of respiration by nitric oxide induces a *Mycobacterium tuberculosis* dormancy program. *J. Exp. Med.* 198:705-713.

Wayne, L.G., and L.G. Hayes. 1996. An *in vitro* model for sequential study of shiftdown of *Mycobacterium tuberculosis* through two stages of nonreplicating persistence. *Infect. Immun.* 64:2062-2069.

Wayne, L.G., and L.G. Hayes. 1998. Nitrate reduction as a marker for hypoxic shiftdown of *Mycobacterium tuberculosis*. *Tuber Lung Dis.* 79:127-132.

Wayne, L.G., and H.A. Sramek. 1994. Metronidazole is bactericidal to dormant cells of *Mycobacterium tuberculosis*. *Antimicrob. Agents Chemother.* 38:2054-2058.

WHO. 2013. Global tuberculosis report 2013. World Health Organization, Geneva.

WHO. 2019. Global tuberculosis report 2019. World Health Organization, Geneva.

Zhao, X., J.Y. Wang, C. Xu, Y. Dong, J. Zhou, J. Domagala, and K. Drlica. 1998. Killing of *Staphylococcus aureus* by C-8-methoxy fluoroquinolones. *Antimicrob. Agents Chemother.* 42:956-958.

1430   Zhao, X., C. Xu, J. Domagala, and K. Drlica. 1997. DNA topoisomerase targets of the

1431   fluoroquinolones: a strategy for avoiding bacterial resistance. *Proc. Natl. Acad. Sci.*

1432   *U.S.A.* 94:13991-13996.

1433

1434

1435

1436

1437

1438

1439

1440

1441

1442

1443

1444

1445

1446

1447

1448

1449

1450

1451

1452

 Figure legends

 Fig 1. Effect of moxifloxacin on biosensor oxidation, ROS level, and bacterial

 survival. (**A**) Moxifloxacin increases oxidative stress that increases the ratio of oxidized

 (MSSM) to reduced mycothiol (MSH) via mycothiol-dependent peroxiredoxin (Prx) or

 peroxidase (Mpx). Oxidation of Mrx1-roGFP2 increases fluorescence intensity for

 excitation at ~400 nm and decreases it for excitation at ~490 nm. (**B**)   Strain *M.*

 tuberculosis-roGFP2 was exposed to the indicated concentrations of moxifloxacin (1X

 MIC = 0.5 µM) for 48 h, and the ratiometric response of the biosensor and survival

 following treatment were determined. (**C**) Cells were treated as in panel B, and ROS were

 quantified by flow-cytometry using CellROX Deep Red dye. 10 mM cumene

 hydroperoxide (CHP) served as a positive control. Data represent mean fluorescence

 intensity of the dye. (**D**) Exponentially growing *M. tuberculosis* H37Rv was treated with

 moxifloxacin at the indicated concentrations for the indicated times; survival was

 assessed by determining colony-forming units (CFU). (**E**) Time-kill curves for *M.*

 tuberculosis treated with 10X MIC of moxifloxacin. Error bars represent standard

 deviation from the mean. Data represent at least two independent experiments performed

 in at least duplicate. Statistical significance is calculated against the no-treatment control

 (****$p < 0.0001$, *** $p < 0.001$, and **$p < 0.01$, *$p < 0.05$).

 Fig 2. ROS mitigation reduces moxifloxacin-mediated killing of *M. tuberculosis*. (**A**)

 Plan for detecting thiourea (TU) and bipyridyl (BP) effects on moxifloxacin lethality. (**B**)

 Exponentially growing *M. tuberculosis* H37Rv cultures were either left untreated or treated

 with 10 mM TU for 1 h before addition of the indicated concentrations of moxifloxacin

 (MOXI; 1X MIC = 0.5 µM) for 10 days followed by determination of CFU. (**C**) Effect of

bipyridyl. *M. tuberculosis* as in *B* was untreated or treated with 250 µM BP for 15 min

prior to addition moxifloxacin as in *B*. (**D**)　The Mrx1-roGFP2 biosensor ratiometric

response was determined after 48 h treatment of *M. tuberculosis* cultures with the

indicated concentrations of MOXI alone or with 250 µM of BP. (**E**) *M. tuberculosis* cultures

were treated with moxifloxacin, the drug was removed by washing, and cells were plated

on drug-free 7H11 agar with or without catalase followed by CFU determination. (**F**)　*M.*

tuberculosis cultures were treated with 1X MIC of moxifloxacin for the indicated times,

washed, and plated with or without catalase (17.5 U/mL of agar). Percentage survival was

calculated relative to CFU of cultures at 0 h. Statistical significance was calculated

between drug-alone group with drug + catalase group. Statistical considerations were as

in Fig 1.

Fig 3. Whole-genome transcriptome profiling of *M. tuberculosis* treated with

moxifloxacin. (**A**) Heat-map showing expression changes due to 16-h treatment of *M.*

tuberculosis with moxifloxacin. Differentially expressed genes (DEGs) exhibited a 2-fold

change across all three treatment conditions (2X, 4X, and 8X MIC of moxifloxacin; 1X

MIC = 0.4 µM); color code for the fold change is at the bottom of the second column

(yellow: upregulated genes; turquois: down-regulated genes). Genes are grouped

according to function. For genes belonging to bioenergetics processes, color code for the

fold change is at the bottom of the fourth column. *$*sdhC$ and *cmtR* are deregulated in two

treatment conditions (2X and 4X) (**B**) Venn diagram showing transcriptome overlap

between moxifloxacin-mediated (green circle) and H_2O_2-mediated stress for *M.*

tuberculosis (Voskuil et al., 2011); DEGs obtained with treatment with 5 mM H_2O_2 for 40

min and 5 mM H_2O_2 for 4 h are shown in beige and blue circles, respectively.

Fig 4. Moxifloxacin-mediated respiration arrest reversed by NAC. (**A**) OCR (pmol/min) indicated oxygen consumption rate. Exponentially growing *M. tuberculosis* cultures were either left untreated (UT) or treated with 10X MIC of moxifloxacin (MOXI; 5 µM) for the indicated times; black dotted lines indicate when MOXI or NAC (1 mM) or CCCP (10 µM) were added. Determination was via Seahorse XFp Analyzer (**B**) ECAR (mpH/min) indicated H$^+$ production or extracellular acidification due to glycolytic and TCA flux. Determination was as in OCR with data representing percentage of third baseline value. NAC (1 mM) addition enhanced OCR of (**C**) untreated and (**D***)* MOXI-treated cells. Data shown are representative of two independent experiments performed in triplicate.

Fig 5. Dissipation of NADH reductive stress diminishes moxifloxacin-induced ROS increase and lethality with *M. tuberculosis.* Detection of **(A)** NADH or **(B)** NAD$^+$ levels. *M. tuberculosis* was treated with moxifloxacin (1X MIC = 0.5 µM), for 2 days, and NADH or NAD$^+$ levels were determined by an alcohol dehydrogenase-based redox cycling assay. **(C)** NADH/NAD$^+$ ratio. Untreated *M. tuberculosis* expressing *LbNox* served as a control. *p* was determined by unpaired two-tailed student's t-test analyzed relative to the untreated control. **(D)** ROS response to moxifloxacin. Wild-type *M. tuberculosis* H37Rv (WT *Mtb*) and cells expressing *Lbnox* (*LbNox*) were exposed to the indicated concentrations of moxifloxacin for 48 h, and ROS were quantified by flow-cytometry using CellROX Deep Red. **(E)** Cultures of exponentially growing wild-type *M. tuberculosis (*WT *Mtb)* and *LbNox* were treated with the indicated concentrations of moxifloxacin for 48 h, and survival was assessed by determining CFU. Statistical coniderations were as in Fig 1 (**** *p*< 0.0001, *** and ### *p* < 0.001, ** *p* < 0.01, # *p* < 0.05, ns indicates not significant).

Fig 6. N-acetyl cysteine increases oxidative stress and moxifloxacin-mediated killing of *M. tuberculosis*. NAC (1 mM) was administered 1 h before addition of moxifloxacin (MOXI; 1X MIC = 0.5 µM) at the indicated concentrations followed by 48 h incubation. **(A)** Experimental plan. **(B)** Oxidative stress, measured by the ratiometric response of the Mrx1-roGFP2 biosensor. **(C)** Bacterial survival, measured by plating on 7H11 agar. **(D)** Effect of NAC and moxifloxacin combination with dormant bacilli. *M. tuberculosis* cultures were starved for nutrients for 14 days and then treated with moxifloxacin for 5 days in the presence or absence of NAC (1 mM) before determination of survival. Rifampicin (25 µM) served as a positive control. **(E)** NAC (1 mM) was added when cultures were placed in Vacutainer tubes followed by the treatment conditions indicated in Fig S2A and S2D. Metronidazole (Mtz; 10 mM) served as a positive control. Statistical considerations were as in Fig 1.

Fig 7. Moxifloxacin-induced oxidative shift in E_{MSH} and killing of *M. tuberculosis* inside macrophages. THP-1 macrophages, infected with *Mtb*-roGFP2 (MOI = 1:10), were treated with moxifloxacin (MOXI; 1X MIC = 0.5 µM) immediately after infection and incubated for the indicated times. **(A)** ~10,000 infected macrophages were analyzed by flow cytometry to quantify changes in the E_{MSH} of *M. tuberculosis* subpopulations. **(B)** Bacterial survival kinetics after MOXI treatment of THP-1 macrophages infected with *Mtb*-roGFP2 (CFU determination). **(C)** *Mtb*-roGFP2-infected THP-1 macrophages were treated with MOXI at the indicated concentrations in presence or absence of NAC (1 mM) immediately after infection and incubated for the indicated times; analysis was as in *A*. **(D)** THP1 macrophages, infected by *Mtb*-roGFP2, were treated with NAC (1 mM or 2 mM), MOXI (10 µM), or the combination of NAC plus MOXI at those concentrations. After

the indicated incubation times, the bacterial load in the macrophages was determined by

plating on drug-free agar. *p* was determined by two-tailed student's t-test compared to

MOXI-alone treatment at each time-point. Statistical considerations were as in Fig 1.

Fig 8. NAC decreases MDR *M. tuberculosis* survival in mice when combined with

moxifloxacin. (A) Experimental protocol. **(B** and **C)** Bacterial CFUs were enumerated

from lungs and spleen at the indicated times. *p* was determined by unpaired two-tailed

student's t-test analyzed relative to vehicle control treatment (** $p \leq 0.01$, *** $p \leq 0.001$;

ns indicates not significant). Statistical significance between moxifloxacin (MOXI) alone

and MOXI + NAC treatment is also shown (# $p < 0.05$, ## $p \leq 0.01$). Error bars represent

standard deviation from the mean of bacterial burden in 5 mice per group.

Fig 9. Moxifloxacin-mediated killing of *M. tuberculosis* involves accumulation of

NADH-dependent ROS, which is further enhanced by NAC. (a) Moxifloxacin enters

M. tuberculosis and traps gyrase on DNA as reversible, bacteriostatic drug-enzyme-DNA

complexes in which the DNA is broken. The bacterium responds by down-regulating the

expression of genes involved in respiration. The transcriptional changes result in reduced

rate of respiration. NADH levels and the ratio of NADH to NAD^+ increase. NADH increases

the free Fe^{2+} pool by releasing Fe from ferritin-bound forms and keeps it in a reduced

state. ROS damage macromolecules in a self-amplifying process, as indicated by

exogenous catalase blocking killing when added after removal of moxifloxacin. (b)

Addition of N-acetyl cysteine to cells stimulates respiration and provides more ROS from

moxifloxacin-mediated lesions. NAC alone does not induce ROS or trigger death. The

additional ROS increases killing by moxifloxacin. Repair of moxifloxacin-mediated

lesions, NADH dissipation, Fe sequestration, and ROS detoxification mechanisms

contribute to survival.

1569

Figures

1571

1572

Fig 1. Effect of moxifloxacin on biosensor oxidation, ROS level, and bacterial

survival.

1575

1576

1577

1578

1579

1580

1581

1582

1583

Fig 2. ROS mitigation reduces moxifloxacin-mediated killing of *M. tuberculosis*.

1584

1585

1586

1587

1588

1589

1590

1591

1592

1593

1594

1596

Fig 3. Whole-genome transcriptomic profiling of *M. tuberculosis* treated with moxifloxacin.

1599

1600

1601

1602

1603

1604

1605

1606

1607

Fig 4. Moxifloxacin-mediated respiration arrest reversed by NAC.

1608

1609

1610

1611

Fig 5. Dissipation of NADH reductive stress diminishes moxifloxacin-induced

ROS increase and lethality with *M. tuberculosis.*

1612

1613

1614

1615

1616

Fig 6. N-acetyl cysteine increases oxidative stress and moxifloxacin-mediated

killing of *M. tuberculosis*.

1619

Fig 7. Moxifloxacin-induced oxidative shift in E_{MSH} and killing of *M. tuberculosis* inside macrophages.

1625

Fig 8. NAC decreases MDR *M. tuberculosis* survival in mice when combined with

moxifloxacin.

1628

1629

1630

1631

1632

1633

Fig 9. Moxifloxacin-mediated killing of *M. tuberculosis* enhanced by NAC.

1635