Title: Detection and molecular characterization of a phytoplasma associated with Cucumber (*Cucumis sativus*) and its first report in India

Authors: Mantesh Muttappagol[1], H. D. Vinay Kumar[1], Shridhar Hiremath[1], M. Nandan[1], C. R. Jahir Basha[2], K. S. Shankarappa[3], V. Venkataravanappa[4], C. N. Lakshminarayana Reddy[1]

[1]*Department of Plant Pathology, College of Agriculture, University of Agricultural Sciences, GKVK, Bangalore-560065, Karnataka, India*

[2]*Department of Plant Pathology, ARS, University of Agricultural Sciences (B), Rajavanthi, Pavagada, Tumkur (Dist.), Bangalore, Karnataka, India*

[3]*Department of Plant Pathology, College of Horticulture, University of Horticultural Sciences, Bagalkot, Bengaluru, Karnataka 560065, India*

[4]*Division of Plant Pathology, ICAR-Indian Institute of Horticultural Research, Hessaraghatta Lake PO, Bangalore-560089, Karnataka, India*

* Corresponding author's mailing address:
1. Dr. C. N. Lakshminarayana Reddy
        Department of Plant Pathology,
        College of Agriculture
        University of Agricultural Sciences, GKVK, Bangalore,
        Karnataka, India
        Email: cnl.reddy@uasbangalore.edu.in
                cnlreddy@gmail.com

2. Dr. C. R. Jahir Basha
        Department of Plant Pathology,
         ARS, University of Agricultural Sciences (B)
         Rajavanthi, Pavagada, Tumkur (Dist.), Bangalore,
         Karnataka, India
         Email: jahir_basha@rediffmail.com

Abstract

Cucumber (*Cucumis sativus*) plants exhibiting typical phyllody symptoms were collected from

farmer's field of Chintamani, Chickballapur districts of Karnataka (India). Disease incidence of

phyllody was 2–3%. The etiology of the cucumber phyllody phytoplasma (CuPP) was confirmed

by amplifying 16S rRNA gene from symptomatic plants using PCR followed by nested PCR

using universal primers pairs P1/P7 and R16F2n/R16R2. After general detection the non-

ribosomal *Sec*Y and *rp* (ribosomal protein) genes was amplified using specific primers. The PCR

amplified products 16SrRNA (~1.2 kb), SecY (~1.6 kb) rp gene (1.2 kb) was cloned and

sequenced. Sequence and phylogenetic analyses of 16S rRNA, secY and *rp* genes revealed that

the detected phytoplasma is a member of the 16SrI group ('*Candidatus* Phytoplasma asteris').

Further on the basis of computer-simulated RFLP (= *in silico* RFLP) analysis of amplified

F2n/R2 region of 16S rRNA gene indicates that, the detected phytoplasma was belongs to the

subgroup X (16SrII-X). This is the first report on phytoplasma associated with phyllody disease

of cucumber in India.

Key words: cucumber, phyllody, 16SrI phytoplasma, characterization, India

Introduction

Cucumber (*Cucumis sativus* L.) is an important vegetable crop belongs to family *Cucurbitaceae*

grown widely in warm regions of the world. It is the world's ancient vegetable crop originated in

India (Yamaguchi 1983) and popularly known as *'khira'* and gherkins. The crop is extensively

cultivated for its edible tender fruit, preferred as salad ingredient, pickles and also as cooked

vegetable (Navitha et al. 2019). In India cucumber is cultivated with in an area of 78,000 ha with

the production of 1,142,000 MT (Anonymous, 2018 NHM data base). The cucumber is mainly

gown during spring-summer and rainy season in commercial scale as well as kitchen gardens.

The cultivation of cucumber is threaten by number insects and diseases from germination till

harvest and causes heavy economic losses to the growers (Venkataravanappa et al. 2019).

Cucumber is reported to be attacked by plethora of viruses, bacteria and fungi in India,

but not yet by phytoplasmas. All over the world, phytoplasmas cause a multitude of diseases in

several hundred crop species including fruits, vegetables, cereals and trees (McCoy et al. 1989;

Lee et al. 2000). Many cucurbits belongs to the *Cucurbitaceous* family have been infected with

different phytoplasma has been well documented in different parts of the world that includes

witches' broom of *Cucurbita pepo* in India (Rao et al. 2017), cucumber phyllody in Iran

(Hosseini et al. 2015) and cucumber yellows (Valiunas et al. 2017). Yellows of *C. maxima*,

(McCoy et al., 1989), *C. moschata* (Montano et al. 2006), *Luffa cylindrica* L. (Lee et al. 1993;

Gundersen et al. 1994), *Momordica charantia* L. (Alves et al. 2017), *Sechium edule* (Jacquin)

Swartz (Villalobos et al. 2002), *Sicana odorifera* (Montano et al. 2007), *Trichosanthes*

cucumerina (Weng et al. 2021) and *C. moschata* L., (Xu et al. 2021).The phytoplasma is one of

the emerging pathogens in many vegetable crops and causing huge loss to various crops. Based

on the 16Sr RNA gene analysis, 16 different ribosomal groups and subgroups phytoplasmas

infecting different vegetable crops have been well documented across the world (Kumari et al.

2019). Considering nutritional benefits of cucumbers in human diet, the crop is protected against

biotic stress. With this backdrop, the present was attempted to identify and characterize strain of

the phytoplasma associated phyllody disease of cucumber based on PCR assays and 16S rRNA,

*Sec*Y and rp gene sequence comparison analysis.

Materials and Methods

Collection of disease samples

During April 2021, typical phytoplasma like symptoms were observed in cucumber in different

farmers fields in Chintamani (13.40°N 78.06°E.), Chickballapur (Dist.) Karnataka (India). The

infected cucumber plants displayed typical symptoms of phytoplasma infection such as witches'

broom, flower viresence, small leaves, and dense clusters of highly proliferating apical shoot.

Disease incidence was calculated as the percentage of symptomatic plants over total number of

plants observed in an area of 20 X 20 meter plot. Due to creeping growth habit of cucumber, care

was taken not to sample the same plant more than once, total five symptomatic and one

symptomatic leaf samples were collected from same location. A part of the collected leaf

samples were used for DNA isolation and the remaining sample was stored at

-80°C for further use.

DNA extraction and PCR amplification of 16SrRNA, *SecY* gene, and ribosomal protein (rp)

gene

Total genomic DNA of symptomatic and asymptomatic cucumber plants was extracted using

cetyl-trimethyl ammonium bromide (CTAB) method. (Dolye and Doyle 1990). The DNA

extracted from the phytoplasma infected bitter gourd (16SrI, KX179474) and bottle gourd

(16SrI, MT386510) was included as positive control in PCR assays (Venkataravanappa et al.

2017; Ashwathappa et al. 2020). Using universal 16Sr RNA gene specific primers pairs (P1/P7,

(R16mF2/R16mR2), the phytoplasma infection in cucumber was confirm in both direct and

nested PCR assays (Deng and Hiruki 1991; Schneider 1995); (Gundersen and Lee 1996).The

16S rRNA gene sequence is highly conserved and insufficient for finer level differentiation of

phytoplasma strains, therefore housekeeping genes such as translocase subunit *sec*Y and

ribosomal protein (*rp*) was amplified by PCR using primer pairs (SecYF1/SecYR1) (Lee et al.

2010) and (*rp*(II)F1A/*rp*(II)R1A) (Martini et al., 2013). PCR reactions were carried out in a

thermal cycler (Master cycler, Profelex, Hamsburg, Germany) and the cycling protocol used for

PCR assay was followed as earlier described (Venkataravanappa et al. 2017).Then five

microliters of PCR products were analysed in 0.8% (W/V) ethidium bromide stained agarose gel,

and visualized with a UV transilluminator. The amplified PCR products of 16SrRNA, *Sec*Y and

rp genes were purified from agarose gel using Qiaquick gel extraction kit (Qiagen, Hilder, USA)

and cloned into pTZ57R/T cloning vector according to manufacturer's directions (MBI

Fermentas, USA). The transformation was performed using *Escherichia coli* (DH5α) cells. The

recombinant plasmids were purified and confirmed with restriction digestion (*Pst*1 and *Sac*1

enzymes). The confirmed clones were sequenced in both directions at Eruofins Genomics Pvt.

Ltd., Bengaluru, India.

Sequence analysis

The full-length sequences of 16S rRNA, *secY* gene and *rp* genes obtained in the present study

were subjected to BLAST search for finding the similar sequences in the database. The

sequences showing maximum homology with 16SrRNA (Table 1), *secY* gene (Table 2) and *rp*

gene (Table 3) of cucumber phyllody phytoplasma were retrieved from NCBI database and

aligned using BioEdit (Hall 1999) and ClustalW (Thompson et al. 1997) program. MEGA X

software was used for constructed phylogeny using Neighbour-Joining method with 1000

bootstrapped replications (Kumar et al. 2016). Sequence of *Acholeplasma laidlawii* (Accession

number NR076550) for 16S rRNA*, Leptospira interrogans* (Accession number MH376290) for

secY gene sequences and *Acholeplasma laidlawii* (Accession number M81465) were used as out

groups in the phylogenetic tree construction.

Virtual RFLP analysis

Virtual RFLP (Restriction fragment length polymorphism) analysis was carried out for F2n/R2 region of the 16Sr RNA gene derived from cucumber phyllody phytoplasma using *iPhy*Classifier online tool (https://plantpathology.ba.ars.usda.gov/cgi-bin/resource/iphyclassifier.cgi) (Zhao et al. 2009) to identify the phytoplasma to group and subgroup level. The virtual RFLP patterns were compared, and the similarity coefficient was calculated (Wei et al. 2008). Restriction pattern obtained for the phytoplasmas associated with cucumber phyllody was compared with already published RFLP patterns of phytoplasma representatives of available ribosomal groups and 16SrI subgroups (Lee et al. 1998).

Results

The predominant symptoms observed on phytoplasma infected cucumber plants are little leaf, excessive proliferation along the stem, flower virescence and phyllody, internode shortening phyllody and witches' broom (Fig1a-d). These symptoms are similar to those described for cucumber phyllody disease reported to occur in other countries (Salehi et al. 2015). The cucumber phyllody disease incidence was ranged from 2-5%.

The resulting PCR amplicon of 1.2 kb corresponding to the phytoplasma 16Sr RNA gene, *secY* gene (1.6 kb) and *rp* gene (1.2 kb) were amplified in all four cucumber samples displaying phyllody disease symptoms and positive control samples. However, no such amplification was observed in asymptomatic plants with each of primer pair employed. Amplified PCR fragments of each gene was cloned separately and sequenced in both the direction. The consensus sequences of 16S rRNA (accession number MZ401356), *secY* gene (accession number MZ463022) and *rp* gene (accession number MZ463023) were deposited in GenBank.

BLASTn searches for homologous gene sequences of 16S rRNA gene sequence of cucumber phyllody isolate obtained in the study was more homologous to those of phytoplasmas classified in the, 16SrI group ('*Candidatus* Phytoplasma asteris') having the maximum nucleotide(nt) sequence identity of 99.13% with cucumber viresence from Taiwan (GU361755). The result was well supported in phylogenetic analysis of 16S rRNA gene with selected reference strains indicated that the cucumber phyllody phytoplasma is clustered together with several strains of phytoplasma belongs to the 16SrI subgroup (Fig. 2a).

The computer-simulated RFLP (*in silico* RFLP) analysis of F2n/R2 region derived from 16S rRNA gene sequence of cucumber phyllody phytoplasma was identical (similarity coefficient 0.98) to the reference pattern of 16Sr group I, subgroup X (GenBank accession number JF781308). Therefore, cucumber phyllody phytoplasma having similarity coefficient more than the threshold level (Wei et al. 2008) and it may be considered as a variant of 16SrI-X subgroup under "*Candidatus* Phytoplasma asteris" 16Sr I group.

The *secY* sequence of cucumber phyllody phytoplasma showed the maximum nt similarity of 98.75% to that of *Trema levigatum'* witches'-broom from China (GenBank accession number MW032211). Similarly, *rp* gene of cucumber phyllody phytoplasma shared maximum nt similarity of 99.34% to that rapeseed phyllody phytoplasma from Poland (accession number CP055264). This result was also well supported by the phylogenetic analysis, in which *secY* (MZ463022) and *rp* (MZ463023) gene sequences of the cucumber phyllody phytoplasma is closely cluster with several phytoplasmas belongs to *Candidatus* Phytoplasma asteris" 16Sr-I group (Fig. 2b and 2c).

Attempt to detect phytoplasma in the cucumber samples by LAMP assay using different LAMP primers specific to 16S RNA gene showed that CuP was successfully detected in the

infected cucumber samples (Fig. 7a). No amplification was observed from the DNA of negative

sample and water control. The LAMP amplicons (a ladderlike pattern) were observed on 1.5%

agarose gel from positive samples, but not from the negative sample and water control. The

positive reactions can be visualized directly by the naked eye with samples showing sky blue

color pattern, whereas a violet color change is observed in negative and water control (Fig. 7b).

Phytoplasma diseases cause significant economic losses in numerous *Cucurbitaceae*-

related vegetable crops, including *C. maxima* (Giant pumpkin), *C. mixta* (winter Squash), *C.*

pepo (summer squash) (Rao et al. 2017), *Lagenaria leucantha* (Bottle gourd) (Ashwathappa et

al. 2020; Tripathi et al. 2017), *C. moschata* (Butternut Pumpkin) (Montano et al. 2006; Xu et al.

2021), *Momordica charantia* (Bitter gourd) (Montano et al. 2007; Venkataravanappa et al.

2017). However, there is no information is available for phytoplasma infection in cucumber in

India. Literature survey showed that cucumber acts as natural host for different phytoplasma

groups, that includes 16SrII-M from Iran (Dehghan et al. 2015; Ghayeb Zamharir and Azimi

2018), 16SrVI-A (Salehi et al. 2015) from Iran. In the present study the cucumber phyllody

phytoplasma resulted from association of 16SrI group phytoplasma was confirmed based 16Sr

RNA gene characterization including *secY* and *rp* genes. This study represents the first recorded

of phytoplasma association with cucumber in Karnataka (India) is related to '*Ca*. P. asteris'-

related strain belonging to 16Sr I-X subgroup. This disease become major threat for cultivation

of cucumber as it causes 100% yield losses due to shoot and floral malformation and sterility.

Moreover, the crop is a regular and widely cultivated throughout India in large farming

community, therefore, further studies on epidemiology along with possible natural spread

sources and management is also a foremost need.

Contributions:

MM, HDV and MN performed the biological experiment, SH, VV and KSS involved in the bioinformatics analysis CRJB and CNL involved in experimental design, CNL conceptualized work and provided the overall direction, all authors are involved in literature mining and manuscript preparation. All authors read and approved the final manuscript

Ethics declarations

This article does not contain any studies requiring ethical approval.

Conflict of interests

The authors declare that they have no competing interests**.**

Human and animal rights

This article does not contain any studies with human or animal subjects performed by any of the authors.

Informed consent

Informed consent was obtained from all individual participants included in the study.

References

Alves M, Ribeiro G, Souza A, Carvalho C (2017) First report of a 16SrIII-B phytoplasma associated with *Momordica charantia* witches'-broom in Brazil. Plant Dis 101 (7):1314

Ashwathappa KV, Reddy L, Reddy CN, Venkataravanappa V, Sastry SV, Reddy MG, Rao GP (2020) A new strain of '*Candidatus* phytoplasma asteris' associated with bottlegourd phyllody disease in South India. Phytopathogenic Mollicutes 10 (1):43-49

Dehghan H, Salehi M, Khanchezar A, Afshar H (2015) Biological and molecular characterization of a phytoplasma associated with greenhouse cucumber phyllody in Fars province. Iranian J Plant Patholo50 (4):393-401

Deng S, Hiruki C (1991) Genetic relatedness between two nonculturable mycoplasmalike organisms revealed by nucleic acid hybridization and polymerase chain reaction. Phytopathology 81 (12):1475-1479

Dolye J, Doyle JL (1990) Isolation of plant DNA from fresh tissue. Focus (Madison) 12 (1):13-15

Ghayeb Zamharir M, Azimi H (2018) Detection and characterisation of a phytoplasma associated with cucumber (*Cucumis sativus*) regional yellows disease in Iran. Arch Phytopathol Plant Protect 51 (15-16):889-893

Gundersen D, Lee I-M (1996) Ultrasensitive detection of phytoplasmas by nested-PCR assays using two universal primer pairs. Phytopathol Mediterr:144-151

Gundersen DE, Lee I-M, Rehner SA, Davis RE, Kingsbury DT (1994) Phylogeny of mycoplasmalike organisms (phytoplasmas): a basis for their classification. J Bacteriol 176 (17):5244-5254

Hall TA BioEdit: a user-friendly biological sequence alignment editor and analysis program for Windows 95/98/NT. In: Nucleic acids symposium series, 1999. vol 41. [London]: Information Retrieval Ltd., c1979-c2000., pp 95-98

Kumar S, Stecher G, Tamura K (2018) MEGA7: molecular evolutionary genetics analysis version 7.0 for bigger datasets. Mol Biol Evol 33 (7):1870-1874

Kumari S, Nagendran K, Rai AB, Singh B, Rao GP, Bertaccini A (2019) Global status of phytoplasma diseases in vegetable crops. Front Microbiol 10:1349

Lee I-M, Bottner-Parker K, Zhao Y, Davis R, Harrison N (2010) Phylogenetic analysis and

delineation of phytoplasmas based on secY gene sequences. Int J Syst Evol Microbiol 60

(12):2887-2897

Lee I-M, Davis RE, Gundersen-Rindal DE (2000) Phytoplasma: phytopathogenic mollicutes.

Ann Rev Microbiol 54 (1):221-255

Lee I-M, Hammond R, Davis R, Gundersen D (1993) Universal amplification and analysis of

pathogen 16S rDNA for classification and identification of mycoplasmalike organisms.

Phytopathology 83 (8):834-842

McCoy R, Caudwell A, Chang C, Chen T, Chiykowski L, Cousin M, Dale J, de Leeuw G,

Golino D, Hackett K (1989) Plant Diseases Associated with Mycoplasma-Like

Organisms. Plant diseases associated with mycoplasma-like organisms In Whitcomb RF,

Tully JG, eds Mycoplasmas 5:545-640

Montano H, Brioso P, Pimentel J, Figueiredo D, Cunha Junior J (2006) *Cucurbita moschata,*

new phytoplasma host in Brazil. J Plant Pathol 88 (2)

Montano HG, Brioso PS, Junior JOC, Figueiredo DV, Pimentel JP (2007) First report of group

16SrIII phytoplasma in loofah (*Luffa cylindrica*). Bull Insectol 60 (2):277

Navitha P, Sujatha K, Beaulah A (2019) Effect Effect of fruit size on physiological seed quality

parameters of Cucumber (*Cucumis sativus*). J Appl Natur Sci 11 (2):394-397

Rao G, Gopala GS, Rao A (2017) First report of a'*Candidatus* Phytoplasma asteris'-related

strain (16SrI-B subgroup) associated with witches' broom disease in *Cucurbita pepo* in

India. New Dis Rep 35 (1):33-33

Salehi M, Siampour M, Esmailzadeh Hosseini S, Bertaccini A (2015) Characterization and
vector identification of phytoplasmas associated with cucumber and squash phyllody in
Iran. Bull Insectol 68 (2):311-319

Schneider B (1995) Phylogenetic classification of plant pathogenic mycoplasma-like organisms
or phytoplasma. Mol Diagno Proced Mycoplasmol 1:369-380

Thompson JD, Gibson TJ, Plewniak F, Jeanmougin F, Higgins DG (1997) The CLUSTAL_X
windows interface: flexible strategies for multiple sequence alignment aided by quality
analysis tools. Nucleic Acids Res 25 (24):4876-4882

Tripathi S, Thorat V, Verma R, Shouche Y, Yadav AJPD (2017) First report of '*Candidatus*
phytoplasma asteris'(subgroup 16SrI-X) associated with bottle gourd virescence and
phyllody disease in India. Plant Dis 101 (11):1949

Venkataravanappa V, Reddy C, Shankarappa K, Jayappa J, Pandey S, Krishna Reddy M (2019)
Characterization of *Tomato leaf curl New Delhi virus* and DNA-satellites association
with mosaic disease of cucumber. Int J Biotech & Bioeng 5:6-93

Venkataravanappa V, Reddy LRCN, Swarnalatha P, Shankarappa SK, Reddy MK (2017)
Detection and characterization of '*Candidatus* Phytoplasma asteris' associated with
littleleaf disease of bitter gourd from India by 16S rRNA phylogenetic and RFLP (in vitro
and virtual) analysis. Arch Biol Sci 69 (4):707-714

Villalobos W, Moreira L, Rivera C, Bottner K, Lee I-M (2002) First report of an aster yellows
subgroup 16SrI-B phytoplasma infecting chayote in Costa Rica. Plant Dis 86 (3):330-330

Wei W, Lee M, Davis RE, Suo X, Zhao YJ (2008) Automated RFLP pattern comparison and
similarity coefficient calculation for rapid delineation of new and distinct phytoplasma
16Sr subgroup lineages. Int J Syst Evol Microbiol 58 (10):2368-2377

Weng YY, Liou WC, Chien Y, Liao PQ, Wang CJ, Chiu YC, Chen YK, Yang JYJPD (2021)

First Report of 16SrII-V Peanut Witches' Broom Phytoplasma in Snake Gourd

(*Trichosanthes cucumerina* L.) in Taiwan. Plant Dis 101 (11):1949

Xu W, Chun-Guang W, Xiao-Yan L, Zheng-Nan L (2021) Molecular detection and identification

of a '*Candidatus* Phytoplasma solani'-related strain associated with pumpkin witches'

broom in Xinjiang, China. Phytopathol Mediterr 60 (1):63-68

Yamaguchi M (1983) Cucurbits. In: World Vegetables. Springer, pp 312-349

Zhao Y, Wei W, Lee M, Shao J, Suo X, Davis RE (2009) Construction of an interactive online

phytoplasma classification tool, iPhyClassifier, and its application in analysis of the

peach X-disease phytoplasma group (16SrIII). Int J Syst Evol Microbiol 59 (Pt 10):2582

Table:1 16Sr RNA gene sequences of phytoplasmas employed in analysis

Phytoplasma disease / country	16S rDNA group/subgroup	GenBank accession number	Identity with Cucumber phytoplasma
Cucumber phyllody phytoplasma/ Iran	16SrII-D	KR822804	67.30
Cucumis sativus' phyllody/Iran	16SrVI	KR633068	61.90
Cucumber phyllody phytoplasma/Iran	16SrII	JN574839	72.20
'*Candidatus* Phytoplasma trifolii'/Iran	16SrVI	KT807467	62.20
Cucumber viresence /Taiwan	16SrI	GU361755	99.10
Aster yellows/USA	16SrI-B	M30790	68.60
Potato purple top/Russia	16SrI-B	EU333396	68.60
Potato purple top/Russia	16SrI-P	EU333397	68.60
Apricot aster yellows/Spain	16SrI-F	AY265211	68.30
Aconitum proliferation/ Lithuania	16SrI-A	AF510323	68.40
Aster yellows-AV2192/Germany	16SrI-L	AY180957	68.60
Aster yellows-AVUT/Germany	16SrI-M	AY265209	68.70
Rhododendron x hybridum/ Czech	16SrI-C	KC009838	68.40

Republic			
Aster yellows-BB/USA	16SrI-A	AY180955	68.40
Blueberry stunt-BBS3/USA	16SrI-E	AY265213	68.40
Aster yellows Btsv2CarD3/USA	16SrI-B	AY180945	68.60
Cherry little leaf/ Lithuania	16SrI-Q	AY034089	68.50
Japanese maple witches' broom/	16SrI-B	JQ015183	68.70
Clover phyllody/Germany	16SrI-C	AY265217	68.40
Valeriana yellows/ Lithuania	16SrI-M	AY102274	68.10
Aster yellows / USA	16SrI-O	AF268405	68.70
Cucumis sativus' stunt virescence/ Lithuania	16SrI-A	KX298483	68.40
Cucumis. sativus' phytoplasma /Iran	16SrVI-A	MH004460	62.50
Cucumis sativus' phytoplasma /Iran	16SrVI-A	MH004459	39.40
Western X-disease/USA	16SrIII-A	L04682	61.90
Coconut lethal yellowing/USA	16SrIV-A	U18747	62.20
Elm yellows/USA	16SrV-A	AY197655	62.20
Jujube witches' broom/Korea	16SrV-B	AB052876	61.30
"Flavescence dorée/France	16SrV-C	AF176319	62.20
Clover proliferation/Canada	16SrVI-A	AY390261	62.20
Ash yellows/USA	16SrVII-A	AF092209	62.40
Apple proliferation/ Germany	16SrX-A	AJ542541	63.40
Spartium witches' broom/Italy	16SrX-B	X92869	63.50
Pear decline/ Germany	16SrX-C	AJ542543	63.60
European stone fruit yellows/ Germany	16SrX-F	AJ542544	63.50
Rice yellow dwarf/Germany	16SrXI-A	AB052873	60.80
Cucumis sativus phytoplasma /Iran	16SrVI-A	MH004458	39.40
Squash phyllody phytoplasma/Iran	16SrII-D	KR822805	67.40
Strawberry yellows / Lithuania	16SrXII-E	DQ086423	66.80
Mexican periwinkle virescence/Mexico	16SrXIII-A	AF248960	66.40
Bermudagrass white leaf/Italy	16SrXIV-A	AJ550984	62.00
Chestnut witches' broom/South Korea	16SrXIX-A	AB054986	61.40

Hibiscus witches' broom/Brazil	16SrXV-A	AF147708	61.50
Sugarcane yellow leaf/Cuba	16SrXVI-A	AY725228	64.80
American potato purple top wilt/USA	16SrXVIII-A	DQ174122	66.10
Buckthorn witches' broom/USA	16SrXX-A	X76431	63.10
Pine shoot proliferation/Europe	16SrXXI-A	AJ632155	62.00
Awka wilt/Nigeria	16SrXXII-A	Y14175	61.80
Buckland Valley grapevine yellows/ Australia	16SrXXIII-A	AY083605	66.70
Sorghum bunchy shoot/USA	16SrXXIV-A	AF509322	59.20
Cassia italica witches' broom/Oman	16SrXXIX-A	EF666051	61.40
Weeping tea witches' broom/Australia	16SrXXV-A	AF521672	61.20
Sugarcane yellows D3T1/ Mauritius	16SrXXVI-A	AJ539179	64.40
Salt cedar witches' broom/China	16SrXXX-A	FJ432664	63.80

292 **Table:2 *SecY* gene sequences of phytoplasmas employed in the analyses.**

Phytoplasma Species /Country	Sub-group	GenBank accession No.	Identity with Cucumber phytoplasma
Tomato big bud phytoplasma/USA	16SrI-A	AY803178	83.40
Chrysanthemum yellows phytoplasma/ Germany	16SrI-A	AY803170	83.20
Hydrangea phyllody phytoplasma/Belgium	16SrI-A	AY803181	83.20
Chrysanthemum yellows phytoplasma/Italy	16SrI-B	DQ787851	87.80
Trema levigatum' witches'-broom/China	16SI-B	MW032211	98.75
Primrose virescence phytoplasma/Germany	16SrI-B	AY803176	86.80
Clover phyllody phytoplasma/Germany	16SrI-C	AY803183	83.80
Paulownia witches-broom phytoplasma/Taiwan	16SrI-D	AY803184	86.50
Blueberry stunt phytoplasma/ Michigan, USA	16SrI-E	AY803169	84.20
Apricot chlorotic leaf roll phytoplasma/Spain	16SrI-F	AY803166	83.50
Strawberry multiplier phytoplasma/Florida, USA	16SrI-K	AY803180	83.50
Aster yellows phytoplasma/USA	16SrI-M	AY803168	86.60
Ipomoea witches-broom phytoplasma/Taiwan	16SrI-N	AY803182	98.58
Peanut witches-broom phytoplasma/Taiwan	16SrII-A	GU004331	26.80
Sesame phyllody phytoplasma/ Thailand	16SrII	GU004322	28.40
Sesame phyllody phytoplasma/ Thailand	16SrII	GU004362	37.50
Soybean phyllody phytoplasma/ Thailand	16SrII-C	GU004324	28.50
Picris echioides phytoplasma/Italy	16SrII-E	GU004348	37.80
Australian tomato big bud phytoplasma/ Australia	16SrII-D	GU004347	37.70

Sesame phyllody phytoplasma/ Myanmar	16SrII	AB703253	37.80
Milkweed yellows phytoplasma/ New York, USA	16SrIII-F	GU004340	30.00
Potato purple top phytoplasma/ Montana, USA	16SrIII-M	GU004333	29.90
Clover yellow edge phytoplasma/ Lithuania	16SrIII-B	GU004332	30.30
Spirea stunt phytoplasma/ New York, USA	16SrIII-E	GU004326	28.20
Poinsettia branch-inducing phytoplasma/USA	16SrIII-H	GU004328	30.00
Peach X-disease phytoplasma/Canada	16SrIII-A	GU004327	28.10
Walnut witches-broom phytoplasma/ Georgia, USA	16SrIII-G	GU004325	30.00
Elm yellows phytoplasma/ New York, USA	16SrV-A	AY197690	41.00
Cherry lethal yellows phytoplasma/China	16SrV-B	AY197693	41.30
Alder yellows phytoplasma/Germany	16SrV-C	AY197692	41.20
Rubus stunt phytoplasma/Italy	16SrV-E	AY197696	41.50
Flavescence doree phytoplasma/USA	16SrV-D	AY197685	41.40
Dry bean phyllody phytoplasma/ Washington, USA	16SrVI-A	GU004353	40.30
Ash yellows phytoplasma/ NY California, USA	16SrVI-A	GU004329	28.60
Brinjal little leaf phytoplasma/India	16SrVI-D	GU004356	30.30
Potato witches'-broom phytoplasma/Canada	16SrVI-A	GU004316	29.00
Clover phyllody phytoplasma/Canada	16SrVI-A	GU004315	29.10
Potato purple top phytoplasma / Alaska, USA	16SrVI-A	GU004343	29.20
Lucerne virescence phytoplasma/Frans	16SrVI-A	GU004318	30.00
Vinca virescence phytoplasma/ California, USA	16SrVI-A	GU004317	28.70
Potato purple top phytoplasma/ Alaska, USA	16SrVI-A	GU004344	37.30
Potato purple top phytoplasma/ Alaska, USA	16SrVI-A	GU004342	38.20
Potato purple top phytoplasma/ Alaska, USA	16SrVI-A	GU004351	38.70
Dry bean phyllody phytoplasma/USA	16SrVI-A	GU004352	40.00
Coconut lethal yellows phytoplasma/USA	16SrIV-A	GU004320	29.60
Candidatus Phytoplasma fraxini/USA	16SrVII-A	GU004329	28.60
Apple proliferation phytoplasma/Italy	16SrX-A	GU004335	29.60
Pear decline phytoplasma/Italy	16SrX-C	GU004363	36.60
Mexican periwinkle virescence /Meixco	16SrXIII-A	GU004336	29.10
American potato purple top wilt phytoplasma/USA	16SrXVIII-B	GU004338	29.50
Stolbur-It phytoplasma/Italy	16SrXII-A	GU004355	47.50
'Candidatus Phytoplasma americanum'/USA	16SrXVIII-B	MN227134	38.00

294 **Table.3 Rep gene sequences of phytoplasmas employed in analysis**

Phytoplasma Species/ Country	Sub-group	GenBank accession No.	Identity with Cucumber phytoplasma
Gynostemma pentaphyllum' witches'-broom/China	16SrI-B	MN543070	99.00
Aster yellows phytoplasma/Hungary	16SrI-B	MN526023	99.10

Rapeseed phyllody/Taiwan	16SrI	CP055264	99.30
Plum witches'-broom phytoplasma/Poland	16SrI-B	MH061366	96.10
Oilseed rape phytoplasma/China	16Sr1-A	KX551965	99.10
Melochia corchorifolia' phyllody/China	16Sr1-A	KX158198	99.10
Tilia platyphyllos' phytoplasma/ Lithuania	16SrI-B	JX184926	97.90
Salix tetradenia' witches'-broom/China	16SrI-B	KC117314	99.10
Periwinkle phyllody/Korea	16SrI-B	AB741642	98.20
Sesame phyllody/Korea	16SrI-L	AB741638	97.70
Larch phytoplasma strain / Lithuania	16SrI-B	JF767009	98.60
Periwinkle phyllody/China	16SrI-B	HQ285145	99.20
Iranian tomato phytoplasma /Iran	16SrI-B	HQ286478	99.10
Onion proliferation phytoplasma/ Lithuania	16SrI-A	GU228514	96.90
Aster yellows/USA	-	AY183710	96.90
Paulownia witches'-broom/Germany	-	AY264857	96.80
Aster yellows phytoplasma/Germany	-	AY183708	96.10
Green gram phyllody/ Myanmar	16SrII	AB703244	96.70
Black gram witches' broom/Myanmar	16SrII	AB703243	57.10
Parthenium weed witches' broom/ Yunnan	16SrII	KU958723	57.10
Sword bean witches' broom/ Yunnan	16SrII	KC953009	57.70
Peanut witches' broom/ Yunnan	16SrII	JX871469	57.60
Cucumber phyllody/Iran	16SrII	KY365523	57.50
Sesame phyllody/ Myanmar	16SrII	AB703247	56.80
Peach X-disease/USA	16SrIII-A	EF186813	55.60
Goldenrod yellows/USA	16SrIII-D	EF186810	57.80
Vaccinium witches' broom/ Germany	16SrIII-F	EF186809	57.70
Poinsettia branch inducing/ USA	16SrIII-H	EF186811	58.30
Gliricidia little leaf/ Honduras	16SrIX	EF186800	57.80
Knautia arvensis phyllody/ Italy	16SrIX	EF186801	60.40
Pigeon pea witches' broom/USA	16SrIX-A	EF183495	60.20
Almond witches' broom/ Lebanon	16SrIX-B	EF186803	59.70
Picris echioides phyllody/Italy	16SrIX-C	EF186802	59.90
Elm yellows/USA	16SrV-A	AY197675	59.40
Alder yellows/Italy	16SrV-C	AY197666	59.00
Spartium witches' broom/Italy	16SrV-C	AY197672	58.60
Hemp dogbane yellows/USA	16SrV-C	AY197674	58.20
"Flavescence dorée"/Italy	16SrV-D	AY197664	58.80
Catharanthus phyllody/Sudan	16SrVI	EF183494	58.60
Ash yellows/USA	16SrVII-A	EF183492	58.50
Apple proliferation/Italy	16SrX-A	EF193366	58.60
Pear decline/Germany	16SrX-C	EF193370	67.60
Mexican periwinkle virescence/Mexico	16SrXIII-A	EF193365	67.70
American potato purple top wilt/USA	16SrXVIII-A	EF193362	77.90
Tomato "stolbur"/Italy	16SrXII-A	EF193364	78.00
Palm lethal yellowing/Florida	16SrIV-A	DQ297677	80.10
Coconut lethal yellowing/ Honduras	16SrIV-A	DQ318238	47.40
Texas Phoenix palm phytoplasma/USA	16SrIV-A	EF050087	47.30
Lethal yellowing phytoplasma/Jamaica	16SrIV-A	EF186805	47.50
Lethal yellowing phytoplasma/USA	16SrIV-A	EF186804	60.60
Coconut lethal yellowing/USA	16SrIV-A	EF193382	60.50
Mexican periwinkle virescence/ Mexico	16SrXIII	EF193365	60.30
'*Candidatus* Phytoplasma meliae'/Argentina	16SrXIII	KU850947	77.90
'*Candidatus* Phytoplasma meliae'/South Korea	16SrXIII	KU850945	78.20
'*Candidatus* Phytoplasma fragariae'/USA	16SrXIII	KJ144893	78.20

Fig: 1 Symptoms on cucumber (*Cucumis sativus*) plant var. Rashmi under field condition
a; flower virescence with phyllody b; excessive shoot proliferation c; Little leaf d;
cucumber plant with floral virescence and small leaves, internode shortening e.

297
298
299

(a) 16Sr RNA

350 **b) sec Y gene**

(C) rp gene

Figure: 2 Phylogenetic tree constructed with MEGA X Software by utilizing 16S rRNA **(a)**
partial sec *Y* **(b)** and partial rp **(c)** gene sequences of cucumber phyllody phytoplasma (in
red) with phytoplasma strains listed in Tables 1, 2, and 3. The tree was rooted with
Acholeplasma laidlawii (GenBank accession number U14905), *Leptospira interrogans*
(GenBank accession number MH376290) and *Acholeplasma laidlawii* (GenBank

accession number M81465) respectively. A bootstrap analysis with 1,000 replicates was
performed and the bootstrap percent values above 50 are showed along the branches.

386