Emergence and clonal expansion of *Vibrio aestuarianus* lineages pathogenic for oysters in Europe.

Aurélie Mesnil[1,2], Maude Jacquot[1], Céline Garcia[1], Delphine Tourbiez[1], Lydie Canier[1], Audrey Bidois[2], Lionel Dégremont[1], Deborah Cheslett[3], Michelle Geary[3], Alessia Vetri[4], Ana Roque[5], Dolors Furones[5], Alison Garden[6], Petya Orozova[7], Isabelle Arzul[1], Mathieu Sicard[8], Guillaume Charrière[2], Delphine Destoumieux-Garzón[2], Marie-Agnès Travers[2*]

ASIM, Ifremer, Station de La Tremblade, avenue de Mus du Loup, 17390 La Tremblade, France

IHPE, Université de Montpellier, CNRS, IFREMER, Univ Perpignan Via Domitia, Montpellier, France

Marine Institute MI, Rinville, Oranmore, Co. Galway, H91 R673, Ireland

Istituto Zooprofilattico Sperimentale delle Venezie (IZSVe), Viale dell'Università 10, 35020 Legnaro (PD), Italy

IRTA La Ràpita. Aquaculture Program. 43540 – San Carles de la Ràpita, Spain.

Marine Laboratory, Marine Scotland Science, Aberdeen, UK.

National Diagnostic and Research Veterinary Medical Institute, Sofia, Bulgaria.

ISEM, Université de Montpellier, CNRS, IRD, Montpellier, France.

* Corresponding author: marie.agnes.travers@ifremer.fr

Competing interests: The authors declare no conflict of interest.

Abstract

Crassostrea gigas oysters represent a significant global food source, with 4.7 million tons

harvested per year. In 2001, the bacterium *V. aesturianus francensis* emerged as a pathogen

that causes adult oyster mortality in France and Ireland. Its impact on oyster aquaculture has

increased in Europe since its reemergence in 2012. To better understand the evolutionary

mechanisms leading to the emergence and persistence over time of this pathogen, we

conducted a survey of mollusk diseases through national reference laboratories (NRLs) across

Europe. We analyzed 54 new genomes of *V. aestuarianus* (*Va*) isolated from multiple

environmental compartments since 2001, in areas with and without bivalve mortalities. We

used a combination of comparative genomics and population genetics approaches to show

that *Va francensis* lineages have undergone clonal expansion in Europe, likely after a recent

selective bottleneck. Low mutation and recombination rates may have selected particular

virulent genotypes. Furthermore, we identified a specific *cus-cop*-containing island conferring

copper resistance to *Va francensis* whose acquisition may have favored the emergence of

pathogenic lineages adapted to oysters.

Introduction

Marine populations are under increasing threat of emerging infectious disease (EID) (1),

contributing to mass mortality events impacting corals, sea-stars, mollusks and fish (2-4).

Aquaculture is practiced in an open and highly connected environment and thus particularly

threatened by pathogen propagation (5). Moreover, aquacultural practices frequently result in

high population densities and many animal transfers, increasing the risk of pathogen

introduction and dispersal (6).

Oysters (*Crassostrea gigas*) are frequently impacted by bacterial pathogens of the genus

Vibrio (7-9). Vibrios are ubiquitously distributed in the oceans and show a wide range of niche

specialization, ranging from free-living forms to those attachment to biotic and abiotic surfaces,

from symbionts to pathogens, and from estuarine inhabitants to deep-sea piezophiles living under high pressure (10). Among them, *Vibrio aestuarianus* subsp. *francensis (Va francensis)* has caused recurrent mortality of adult oysters since 2001 (11). Having previously been restricted to France, this disease has now spread across European countries (France, Italy, Ireland, and Spain) (12, 13). This spread of *Va francensis* has resulted in mortality rates of approximately 25% and is a major economic concern as it impacts market-size oysters (14). Previous field studies found that pathogenic strains of *Va francensis* can colonize sentinel oysters, but are rare in samples from the local environment, *i.e.* sediment and water column. These data suggest that oysters are the preferred habitat of this pathogen (15). Moreover, it has been proposed that *Va francensis* can persist in oyster tissues during winter (15). These observations suggest *Va francensis* has adapted to oysters and raise questions about how this host specialization has occurred.

The emergence of pathogenic bacterial lineages can follow the acquisition of new traits which increase the ability to colonize a host (16). Trait acquisition can be achieved by mutation (17), but it occurs more frequently through horizontal transfer (through transformation, conjugation or transduction) of one or more genes favoring rapid adaptation to new niches and hosts (18). After emergence, the dynamics of pathogenic lineages are dictated by the relative importance of evolutionary forces: mutation, selection, genetic drift and recombination. Populations whose evolution is not or little influenced by recombination evolve clonally (19). One of the factors limiting the rates of horizontal transfer and recombination in a population is ecological isolation (20, 21) which limits opportunity to exchange genetic material with other populations.

Previous studies on *V. aestuarianus* diversity were based on a limited number of genomes, not reflecting overall diversity across diverse isolation contexts (22). This lack of genomic data hinders a complete understanding of the mechanisms governing the emergence and evolutionary dynamics of pathogenic lineages in *Va francensis.*

In the present study, we analyzed 54 genomes of isolates collected since 2001 across Europe.

We combined phylogenetic and population genetic approaches to decipher evolutionary

processes associated with the origin of the emergence of this oyster pathogen. Our study

shows that *Va francensis* strains exhibit numerous features commonly reported in specialized

pathogens. In particular, our data suggest the acquisition of genes involved in copper

resistance is one mechanism promoting *V. aestuarianus* adaptation to oysters.

Results

European spread of *V. aestuarianus* oyster pathogens

To assess the geographic distribution of pathogenic *V. aestuarianus* in Europe, a

questionnaire was sent in 2021 to all European laboratories in charge of the diagnosis and

surveillance of mollusk diseases (supplementary material). Eight out of the 12 respondent

laboratories reported that *V. aestuarianus* was detected during adult oyster mortality events.

The earliest detections were reported by France and Ireland in 2001, and the most recent

detection was reported by Denmark in 2021 (case used DNA detection only) (fig. 1, A). The

number of countries reporting detection has increased since 2011, with 8 European countries

currently reporting *V. aestuarianus*. Importantly the date of first detection may be under-

estimated, since surveillance and testing for the pathogen has varied between countries and

was only implemented in the majority of the laboratories following the development of a

diagnostic tool in 2009 (23).

*Figure 1 : Figure 1: European expansion of V. aestuarianus francensis. While the geographic distribution of V. aestuarianus strains involved in C. gigas mortalities expands, genetic diversity remains stable over time. A) The number of countries reporting the detection of V. aestuarianus was assessed through a questionnaire sent to European national reference laboratories (NRLs). *Note: Denmark case report corresponds to a suspected case without bacterial isolation. B) The two lineage expansions: 2 genotypes were determined based on a maximum likelihood phylogenetic tree from complete genome alignments (GTR substitution model).*

V. aestuarianus diversity is structured according to environmental origin and

isolation context.

In order to assess the overall population structure of *V. aestuarianus* in Europe, we studied

genetic diversity at the whole genome level using *V. aestuarianus* isolates from a broad variety

of origins. 54 strains were sequenced and genomes were *de novo* assembled. We obtained

15 4.5-Mb complete genomes (with 2 to 4 contigs) and 39 draft genomes (with up to 150

contigs) (table S1). Whole genomes were aligned, and relationships between isolates were

studied by building a maximum likelihood phylogeny (fig. 2). Overall, genomes were close to

each other in the tree and showed highly similar nucleotide compositions, showing average

nucleotide identities (ANIs) higher than 97.1% for all pairwise comparisons (table S2). The tree

showed a tightly clustered clade of oyster clinical isolates (*i. e.*, isolates from moribund oysters)

(n = 25) inside genetically diverse *V. aestuarianus* strains isolated outside of oyster mortalities

(n = 29), encompassing non-clinical isolates (i.e. strains isolated from sites without mortalities)

and strains isolated during cockles mortalities. All oyster clinical isolates are encompassed in

Va francensis subspecies (24) and all *Va francensis* isolates were found in oysters, and

showed higher virulence levels when assayed through injection into oyster muscle (p-value<

0.001). The reconstructed tree also confirmed that this subspecies is composed of two

lineages named A (n = 17) and B (n = 8) (22).

Figure 2 : V. aestuarianus diversity is structured according to host species and isolation context. Phylogenetic tree of V. aestuarianus species calculated with RAxML using a GTR substitution model and rooted with the midpoint rooting option from whole-genome alignments. Several layers of annotations are added: country (check mark), matrix of isolation (star), year of isolation (point), virulence

testing in the laboratory by injection into oysters (107 UFC/animal, duplicates of 10 animals) indicated with a bar plot on the right side representing mortality percentages, and context (strains isolated during oyster mortality events are indicated with a red arrow on the right). Va francensis lineages A and B are highlighted in blue and green, respectively. Their virulence is significantly higher compared to non clinical isolates (i. e. strains isolated on sites without mortalities) by Wilcoxon-Mann-Whitney test (p-value < 0.001). The absence of virulence data means the absence of a test for this strain.

Analysis of molecular variance (AMOVA) performed on the sequence dataset revealed that

the isolation context ("oyster mortality" or not) and environmental origin (five host species,

zooplankton and water) explained most significantly ($p < 0.005$) genetic variation (table 1).

AMOVA also indicated the absence of spatial and temporal delineation of the oyster pathogen

lineages. Altogether, our data reveal *V. aestuarianus* lineages linked European expansion of

this oyster pathogen.

Table 1 : Analysis of molecular variance supporting genetic partitioning according to environmental origin and isolation context

Factor	Variability (%)	p-value
Environmental origin	**17.8**	**0.002**
Isolation context	**16.0**	**0.001**
Country of isolation	7.1	0.074
Year of isolation	3.4	0.25

Temporal emergence of oyster pathogen lineages

Next, we used a Bayesian approach to date the emergence and reconstruct the temporal

evolution of oyster pathogens lineages. As a critical prerequisite, we checked whether the *Va*

francensis subspecies is a measurably evolving population. A temporal signal was observed

with a correlation between genetic distances from a reconstructed common ancestor and

strains sampling time, yielding a R² of 0.4236 (fig. S1A). The best-fitting evolutionary model

for the *Va francensis* population was obtained under a GMRF Bayesian skyline population

model with a relaxed clock, leading to a rate of 7.27×10^{-7} substitutions per position per year,

or 3.2 mutations per genome per year. The mean time of emergence of the most recent

common ancestor (TMRCA) of the lineage A was estimated to be 1956 (95% highest posterior

density (HPD): 1919–1972). At this time, a divergence occurred between strain 01-151, which

is non-virulent toward oysters, and other lineage A strains that are mostly pathogenic. The

TMRCA of lineage B maps to the late 1970s, albeit with uncertainty (95% HPD: 1898–1993).

According to the coalescence-based reconstruction of demographic variation, the *Va*

francensis population has slowly decline since its emergence, with accelerated decline since

2000 (fig S1B).

Genome stability in *V. aestuarianus francensis* lineages

Recombination can drive genetic diversity and evolutionary dynamics in bacterial populations.

We therefore quantified homologous recombination events using *V. aestuarianus* genome

alignments.

Recombination has little impact on oyster pathogen evolution. We found that

recombination (r) has a lower impact on the core genome substitution rate in *Va francensis*

lineages than the impact of mutation (m), with mean r/m = 0.36+/- 0.22 and 0.28+/- 0.09 in

lineages A and B respectively. The ratios of these lineages are significantly (student test p-

value < 0.005) smaller than in non-clinical *V. aestuarianus* isolates (i.e. strains isolated from

sites without mortalities) (r/m = 0.91 +/- 0.45). 7.8 and 3.3 recombination events were identified

on average for these lineages. Recombination in genomes can generate phylogenetic

inconsistencies between taxa that can emerge in phylogenetic networks as the presence of

alternative pathways (25). The network that we constructed from the whole-genome alignment

(fig. S2) confirmed that recombination does not affect *Va francensis* isolate relationships, with

few ambiguities among these taxa. One known consequence of the absence of recombination

impact is strong linkage between loci in non-recombinant genomes (26). In *Va francensis*

lineage A and, to a lesser extent, in lineage B, strong linkage between single nucleotide

polymorphisms (*SNPs*) was observed (fig. S3). In contrast, the recombination impact detected

in all other genomes was confirmed with phylogenetic networks, which showed ambiguities

between taxa (fig. S2) and low linkage between loci in these genomes (fig. S3). However,

recombination might have supported the emergence of the *Va francensis* group: 488

recombination events were detected at the node corresponding to the hypothetical common

ancestor of all pathogenic oyster isolates (fig. S4).

Core genome stability was observed in *Va francensis* lineages. Mutations also generated

little genetic diversity among *Va francensis*. Indeed, apart from the recombinant segments,

among the 19 775 variant genomic positions in the full dataset, 666 and 83 positions vary

among *Va francensis* A and B lineages. Finally, low impact of recombination and mutation on

Va francensis lineages results in core genome stability. We used Tajima's D to compare ob-

served genetic diversity in *V. aestuarianus* to that expected under neutral evolution and con-

stant population size. Deviation from neutrality was larger among the *Va francensis* lineages

A (Tajima's D = -2.3) and B (-2.5) than in the full dataset (-0.7), with less polymorphism than

expected under neutral evolution (fig. 3 A). To determine whether the low genetic diversity was

due to demographic effects (affecting whole genomes) or to selection effects (affecting partic-

ular genes), we adopted a gene-based approach to investigate selection pressures at the gene

level. While most of the 1,974 core genes were under neutral evolution outside of the *Va fran-*

censis subspecies, the vast majority were monomorphic within lineages A and B (fig. 3 B),

which is consistent with the purging of genetic diversity through bottlenecks during transmis-

sion dynamics.

Figure 3 : Support for a recent bottleneck that purged genetic diversity in V. aestuarianus francensis lineages. Genetic statistics in V. aestuarianus species (black), other than Va francensis (yellow), Va francensis, lineage A (blue) and lineage B (green). A) Whole genome scale. Even through the expected genetic diversity under neutral evolution and a stable population size in Va francensis is low (Watterson's theta), the observed diversity (Pi) is still lower, as shown by the Tajima's D value below - 2. Pi: observed genetic diversity; Theta_w: Watterson estimate of expected genetic diversity; Tajima_D: Tajima's D test statistic. The Y axis in the Pi and Watterson's theta plots is log10 scaled. B) Gene scale. Core genes show different selection patterns among groups of strains and among core genes within groups. While most core genes are under neutral evolution in isolates other than Va francensis, most genes are monomorphic in Va francensis lineage A and B. D>0 and D<0: significantly negative or positive Tajima's D value (p < 0.05) and D=0 (p > 0.05). A negative Tajima's D is indicative of purifying selection or recent population expansion, while a positive Tajima's D is indicative of diversifying selection or population contraction. A zero value is indicative of neutral evolution.

179   ***Va francensis* subspecies were found to have a closed pangenome**, with genes largely

180   shared among all isolates and almost no strain-specific genes, as determined by dataset

pangenome analysis (fig S5). In contrast, other *V. aestuarianus* isolates exhibited many strain-

specific genes. Altogether, our data support the clonal evolution of the two *Va francensis*

lineages.

Positive selection signatures of oyster pathogen lineages

We used *pcadapt,* which detects structure and clustering of individuals in a population, to

detect potential signatures of positive selection affecting oyster pathogen lineages. 2,925

SNPs were selected with an adjusted p-value <0.001. Based on principal component analysis,

principal component 1 (PC1) segregated the *Va francensis* lineages from the rest of *V.*

aestuarianus strain collection (fig. S6). 46 outlier SNPs were associated with PC1 including 17

non-synonymous and one stop-gained SNP potentially involved in diversifying selection.

These genes are listed in table S3 and notably include genes encoding flagellar proteins

(*pomA*) or related to chemiotaxis.

Oyster-pathogenic *V. aestuarianus* have genomic features of specialist

pathogens

According to low rates of homologous recombination and the closed pangenome, the two *Va*

francensis lineages can be considered as clonally evolving lineages. Furthermore, clonal

evolution, as well as the small number of known habitats (15), suggest that these lineages are

oyster specialists. Pathogen populations have often undergone a change of niche (passage

from a free-living lifestyle to an obligatory parasitic one), ultimately resulting in ecological

isolation and thus inhibition of DNA transfer with the parental population, or even with the

broader bacterial community (27, 28).

We further tested the hypothesis of specialization by analyzing genomic features known to be

enriched during transition to specialist lifestyle : insertion sequences (ISs) and pseudogene

(29). We found a higher number of ISs in the genomes of pathogenic oyster lineages A and B

than in strains isolated from contexts without bivalve mortality (p-value = 0.01 and 0.02,

respectively). Interestingly, in groups containing strains isolated during cockle mortality events,

ISs were more abundant than in pathogenic oyster genomes (p-value = 0.02 and 0.03,

respectively) (fig. 4). IS families were conserved between oyster pathogen genomes, with most

ISs belonging to the IS66, IS902 and ISAS1 families (table S4 and fig S7). IS conservation in

theses clades suggests that IS enrichment and expansion in genomes stopped after pathogen

emergence. The dynamics of the ISs appeared to be different in the cockle isolates. Indeed,

the number and families of ISs represented appeared to be poorly conserved. We suspect that

IS flux is active in cockle isolates, with ongoing horizontal exchange and intragenomic

expansion.

To check whether IS enrichment is involved in pseudogenization, we crossed IS and

pseudogene genomic coordinates and determined the number of ISs with pseudogenes

around them (from 2,000 bp before the IS start to 2,000 bp after the IS end) (table S5 and S6).

Regardless of the genome and chromosome considered, the vast majority (on average 86%)

of ISs had pseudogenes nearby. IS and pseudogene enrichment supports our hypothesis that

the *Va francensis* lineages are specialized toward oyster.

Figure 4 : Differential Insertion sequence (IS) enrichment is observed in Va francensis and cockle
associated isolates genomes. Distribution of the number of IS elements (complete and partial) in 16

Copper resistance acquisition: One potential step in the emergence of oyster-

pathogenic clades

Trait acquisition is often the first step preceding selective bottlenecks and clonal expansion of

host-adapted pathogens. To identify traits that may have preceded oyster pathogen

emergence, we studied the *V. aestuarianus* pangenome and detected 139 genes that were

shared exclusively by *Va francensis* genomes. Most of them (106) were annotated as

hypothetical proteins. The remaining 33 genes encode proteins with functions related to

copper export, cell-bacteria interaction (LPS/antigen O, cell wall or peptidoglycan), chemotaxis

and DNA repair. The sequences of these 139 genes were identical or well conserved between

strains (table S7).

Some oyster pathogen specific genes localize to a putative mobile genetic element (MGE), as

determined by crossing their coordinates with IS coordinates in an oyster pathogen genome

(02_041) (fig. 5 A). This putative MGE contains Cus operon genes (*cusABCF,S/R)* surrounded

by ISs from the IS66 family (length = 15 kb), as well as the copper export genes *copA* and

copG and the regulator *cueR*, located near tn7 transposon genes (tnA, B, C and E) (length =

10 kb), and is later referred to as a *cus-cop*-containing island.

The *Va francensis* ancestor could have acquired copper export genes by horizontal transfer.

Presence in an MGE could have facilitated integration of these genes into the bacterial

genome. To investigate the origin of *cus-cop*-containing islands, we used BLAST analyses of

each operon and of the entire region in 127 reference genomes of different *Vibrio* species

(table S8 and fig S8). The *cus-cop*-containing island presence is scattered along the *Vibrio*

phylogeny and is not specific to a monophyletic group. It appeared highly similar to some *V.*

fluvialis and *V. furnissii* genomic sequences with a nucleotide identity >95%. Moreover, this

region was close to plasmidic elements identified in *V. crassostreae* and *Vibrio* sp. dgh with a

nucleotide identity >89% (fig. 5B).

To test whether the genomic differences between the oyster pathogens and other strains could

lead to phenotypic differences, we tested copper resistance in 27 *V. aestuarianus* strains using

an agar diffusion assay. Supporting our hypothesis, all tested oyster pathogens (8 strains from

lineage A and 5 from lineage B) appeared significantly (p-value < 0.001) more resistant than

other strains to copper in radial diffusion assays (fig. 5C).

Figure 5 : A) Va francensis-specific locus containing copper resistance genes. Transposase and integrase are indicated in yellow, cus genes in orange, and cop genes in blue. B) Evolutionary history of the cus/cop locus. NJ, bootstrap test (500 replicates). The evolutionary distances were computed using the Kimura 2-parameter method and are shown as the number of base substitutions per site. This analysis involved 8 nucleotide sequences. All ambiguous positions were removed for each sequence pair (pairwise deletion option). There was a total of 29 555 positions in the final dataset. Evolutionary analyses were conducted in MEGA X [4]. C) Agar plate assay: bacterial growth inhibition diameter around a pellet impregnated with 50 mM CuSO4 solution. Va francensis linages A and B are significantly

more resistant to copper that other V. aestuarianus strains (Wilcoxon-Mann-Whitney test p value < 0.001).

Discussion

Here, we describe a genomic analysis of *V. aestuarianus* isolated from multiple European

countries, in different epidemic contexts and from multiple environments (five host species,

zooplankton and water) since 2001. We report the expansion and clonal evolution of two

oyster-pathogenic lineages from the *V. aestuarianus francensis* subspecies (11).

We estimate that 7.23×10^{-7} substitutions occur per site per year in *Va francensis*, similar to

the evolutionary rate inferred in *V. cholerae* during pandemic waves (30): this indicates that

pathogenic *Va francensis* lineages emerged during the 1990s, a decade when global oyster

production greatly increased according to Food and Agriculture Organisation. Indeed, oyster

production increased to over 140,000 tons per year in the 1990s (31). According to the

reconstructed demographic history, the effective population size of *Va francensis* is small and

showed a slowly decrease that accelerated in the 1990s. This sharper decreased may have

followed emergence of oyster-pathogenic lineages, indicating a possible bottleneck that

resulted in selection of oyster-adapted strains. This bottleneck may have purged

polymorphisms from this ancestral population, explaining the observed monomorphism in *Va*

francensis core genes. Indeed, some pathogenic monomorphic lineages may endure

population bottlenecks, supporting acquisition of new functions and colonization of new hosts

(32).

Several genomic modifications may help pathogenic oyster lineages adapt to their host,

including core genome substitutions and gene acquisition. We found non-synonymous

substitutions that were fixed in pathogenic oyster lineages. These occurred in genes involved

in host interaction or virulence, such as the *pomA* gene which encodes a flagellum motor

protein and the gene *luxP* linked to quorum-sensing. The functional consequences of these

substitutions remain to be explored.

Acquisition of copper resistance traits may also have contributed to the emergence of the

oyster-pathogenic lineages. Our data indicate that the *Va francensis* ancestor acquired a

genomic island carrying copper detoxification genes. Indeed, copper is a heavy metal found

at high levels in coastal waters because of agricultural and industrial activities (33-35). Copper

can accumulate in oyster tissues (up to 18,000 µg/g in China (36)), particularly at infection

sites (37) where it supports antimicrobial defense as copper can kill invading microorganisms

(37-39). Bacteria have, in turn, developed copper detoxification mechanisms which utilize the

cop and *cus* genes (40). *Cop* genes support initial Cu^+ translocation from the cytoplasm to the

periplasm*. Cus* genes encode a CusABC system that exports periplasmic copper into the

extracellular medium (41). Importantly, *copA* is essential for copper resistance, oyster

colonization and virulence in *V. tasmaniensis* LGP32, an oyster-pathogenic strain associated

with the Pacific Oyster Mortality Syndrome (42). Similarly, *copA* deletion in *V. crassostreae*

also attenuates virulence in oysters (7). Our data further support copper resistance as a shared

trait that allows pathogens to adapt to oysters as a host.

Our results suggest clonal evolution in *Va francensis* lineages. Recombination has a low

impact on genetic diversity. The r/m ratio, ~0.3 in both lineages, is similar to clonally evolving

pathogenic bacteria like the *V. cholerae* 7 pandemic lineage (r/m = 0.1 ; (43)), *M. tuberculosis*

complex (between 0,426 and 0,565, (44)) and lineage 1 of *Salmonella enterica* (r/m = 0.2 ;

(45)). We also found that *Va francensis* has a closed pangenome. This led us to formulate the

hypothesis of an ecological isolation of these lineages, a hallmark of specialized pathogens

that do not exchange genes with other bacterial populations. This hypothesis is also supported

by enrichment of insertion sequences and pseudogenization in both oyster-pathogenic

lineages. During specialization, pathogens transition from a free-living lifestyle to a permanent

association with a host. Through host association, pathogens can acquire compounds of

intermediate metabolism that allow genes associated with related biosynthetic pathways to

become facultative or redundant (25). Studies on pathogen specialization have largely focused

on highly specialized intracellular bacteria, which can undergo an extreme genome

rationalization process (26). This process implies pseudogenization of non-essential genes

that accumulate mutations or ISs and can ultimately result in gene loss (29, 46, 47). However,

genome size reduction was not observed, probably because *Va francensis* pathogenic

lineages are not obligate intracellular parasites. While oyster pathogens have a restricted

habitat, they can still replicate in the absence of their host, at least in rich medium. Moreover,

Va francensis can colonize diverse host tissues, including hemolymph, gills and digestive

connective tissues (15). As a consequence, *Va francensis* is adapted to diverse tissular micro-

environments, consistent with being less specialized than obligate intracellular bacteria.

Further studies of oyster pathogen microhabitats may help to elucidate their complete intra-

host life cycle, and the functions of specific genes linked to adaptation. Whether clades A and

B occupy similar niches in infected oysters and whether they have the potential to cooperate

remains unknown. Neither epidemiological parameter estimation (48) nor in-depth analysis of

their genomes identified lineage-specific functional traits.

Here we report the clonal expansion of *Va francensis* lineages across Europe. It has been

reported that another major pathogen, the herpes virus Ostreid OsHV-1, has also caused the

mortality of *C. gigas* (49) along the French coasts since the emergence of a new variant in

2008 (50). Double infections by OsHV-1 and *Va francensis* were reported in French oyster

farms in 2014 and 2015 (51, 52) and have also been observed in Ireland since 2008 (Cheslett,

pers comm). Although experimental infections combining the two pathogens at high doses in

spat and juvenile *C. gigas* have not demonstrated direct collaboration between them (53, 54),

at the population level the virus may still weaken animal defenses and thus amplify the impact

of *Va francensis*.

The global spread of pathogens is a recurrent problem in agriculture and aquaculture and is

often favored by anthropogenic introductions via infected hosts (55). The many transfers taking

place during the lifetime of an oyster probably explain most of the spread of pathogenic

lineages of *Va francensis* across Europe. Preventing the spread of pathogens during transfers

is particularly complicated in the case of *Va francensis* since the animals may be asymptomatic

(15). There is therefore an urgent need to develop sensitive detection assays to monitor the

spread of pathogens: the specific *Va francensis* genes revealed in this study, such as the

cusS/R genomic island, may provide new diagnostic tools, which may help to monitor and

prevent future mortality events.

Material & methods

Isolate selection

The study collection was designed to represent the temporal, geographic and clinical

heterogeneity of *V. aestuarianus* origins. However, ascertainment bias could occur due to

sampling strategy used. Indeed, strains were mainly isolated during bivalve mortalities. We

maximized recovery of the genetic diversity of oyster pathogenic *V. aestuarianus* by collecting

isolates from all European countries reporting mortalities related to *V. aestuarianus* or

presence of the bacteria. We also included strains isolated during field surveys of sites without

known mortality events. Strains were collected with the help of European NRLs responsible

for the diagnosis and surveillance of mollusk diseases: Ifremer (France), MI (Ireland), IZSVE

(Italy), IRTA (Spain), Marine Scotland Science (Scotland), and NRLFMC (Bulgaria). Sequence

comparisons of isolates from France, notably those isolated from moribund oysters, turned out

to be very similar. In order to equilibrate our sampling, we excluded 17 isolates identical with

included isolates. We selected approximately one strain per batch of moribund oysters (a batch

represents a sampling point) (table S9). The 54 isolates used in the study are summarized in

table S10.

Illumina, Ion torrent and PacBio whole-genome sequencing

Genome sequencing was performed between 2019 and 2021. Several sequencing

technologies were used depending on the objectives. We used a long-read sequencing

technology for some strains to improve assembly quality and study genomic features. DNA

was isolated from 35 strains using the NucleoSpin Tissue Kit (Macherey-Nagel) following the

manufacturer's guidelines. The concentration of each sample was adjusted to 0.2 ng/µl for

library preparation using the Illumina Nextera XT DNA Library Preparation Kit according to the

manufacturer's guidelines. For each library, paired-end reads of 150 bp were produced on the

Illumina NextSeq 550 platform (Bio Environnement, Perpignan, France). The quality of the

resultant fastq files was assessed using FastQC (56). Primers, Illumina adapter sequences,

low-quality bases at the 3′ end of reads with Phred-scaled quality scores less than 30 and

reads with a length <50 bp were removed using Trim Galore (57). Quality-controlled reads

were *de novo* assembled using the SPAdes genome assembler v3.13.1 (58) with the default

kmer size and a coverage cutoff value of 20 reads (59).

For 2 strains (04_091_1T1 and U8), enzymatic shearing was performed on 500ng of genomic

bacterial DNA using the NEBNext Fast DNA Library Prep kit (New Englands Biolabs) and

following the manufacturer's recommendations. Samples were barcoded with individual

IonCodes (Thermo). Purified libraries were then quantified by real-time PCR using the Ion

Universal Library Quantification Kit (Thermo) prior to be diluted and pooled at 25µM each. The

Ion Torrent S5 libraries were prepared using the "Ion 510™ & Ion 520™ & Ion 530™" for Ion

Chef Kit and sequenced on the Ion Torrent S5 using an Ion 530 semi-conductor sequencing

chip (Thermo).

DNA was isolated from other strains using a MagAttract HMW DNA Kit (Qiagen) following the

manufacturer's instructions. Library preparation and sequencing on a PacBio Sequel II

instrument were performed with the GENTYANE INRAe platform (Clermont-Ferrand, France).

De novo genome assemblies were obtained using FLYE v2.8.3 (60). One round of polishing

was performed using RACON v 1.4.3 (61).

The assembly was evaluated with QUAST, and genome completeness was checked using the

BUSCO v5.1.1 (62) and vibrionales_odb10 databases as lineage datasets, including 57

Vibrionales as detailed in table S11.

Assembly features and GenBank accession numbers are summarized in table S1.

Genome annotations

For each assembly, coding sequences (CDSs) were predicted using PROKKA v1.12 (63) with default databases and –compliant and –rfam flags and annotation of rRNAs, tRNAs and other ncRNAs enabled. Functional annotation of the detected CDSs was performed using EggNog-Mapper v2.1.0 (64, 65). ISs were detected in the genomes using ISEScan v1.7 (66). Pseudogene detection was performed with the GenBank files released from PROKKA with PSEUDOFINDER v1.0 (67), using the diamond flag and NCBI nonredundant database. These later analyses were performed only on resolved genomes to avoid obtaining false negatives in IS detection and false positives in pseudogene detection at the intercontig locations.

Phylogeny and recombination analysis

Pairwise ANIs (%) were calculated using FastANI (68). Whole-genome alignments of the 54 isolate contigs were made using Progressive Mauve v13/02/2015 (69). Whole-genome alignment was used to infer the maximum likelihood phylogeny using RAxML v8.2.12 with the TVM model of nucleotide substitution (transversion model, AG=CT and unequal base frequencies) and 100 bootstrap replicates, with strain 03_008T as the outgroup.

To identify polymorphic sites, the haploid variant calling pipeline snippy v4.6 (available at https://github.com/tseemann/snippy) was used with default parameters.

To assess the influence of various structuring factors on the observed genetic diversity, we performed AMOVA on the sequence dataset, with environmental origin, isolation context, sampling year, and country as structuring factors, using the R package *pegas*.

Recombination was evaluated across this genome alignment using ClonalFrameML v1.12 (70). We used an extension that allows different recombination parameters to be inferred on different branches of the clonal genealogy. Linear regression analysis of the root-to-tip distances against sampling time was performed using TempEst v1.5.3 (71). The substitution

rate was estimated under different demographic and clock models, using Beast v1.10.4, taking

advantage of the sampling timeframe between 2001 and 2021. We tested both a strict and a

relaxed molecular clock. Constant-sized, exponential-sized and GMRF Bayesian skyride plot

models (72, 73) were used. For each model, two independent chains were conducted for 200

million generations and convergence was assessed by checking effective sample size values

for key parameters using Tracer v1.7.1 (74). For robust model selection, both path sampling

and stepping-stone sampling approaches were applied to each BEAST analysis to estimate

the marginal likelihood (75, 76). The best model selected by Bayes factor comparison was a

relaxed molecular clock with a GMRF Bayesian skyride demographic model. For this model,

triplicate runs were combined using LogCombiner v1.10.4, with removal of 10% burn-in.

Selection analysis

Tajima's D was calculated considering core genome alignment using MEGA 11 Tajima's test

on neutrality. Core genome alignment were previously partitioned according to lineages or

subspecies, resulting in *Va francensis* lineage A alignment (n = 17 sequences), *Va francensis*

lineage B alignment (n = 8 sequences) and no *Va francensis* alignment (n = 29 sequences).

Since sample size is known to influence population genetic parameter estimations (77), we

calculated Tajima's D on subsamples of eight sequences. For *Va francensis* lineage A and no

Va francensis isolates, Tajima's D was calculated on ten different alignments constructed with

eight sequences randomly subsampled. Then we calculated mean of the Tajima's D obtained

from the ten alignments. For the *Va francensis* lineage B alignment, the eight sequences

represented the entire sampled set. Tajima's D was also calculated for each core genes,

considering the same partitions, employing the R package *pegas*.

In order to detect SNPs that may explain *Va francensis* emergence, we performed a genome

scan analysis using *pcadapt* package v.4.3.3 for R (78) that detect structure and clustering of

individuals in a population. Number of principal components was set to six. We used

LD.clumping option, with window size of 5 kb. LD.clumping uses p-value to sort the SNPs by

importance (e.g. keeping the most significant ones). It takes the first one (e.g. most significant

SNP) and removes SNPs that are too correlated with this one in a window around it. We also

removed SNP with minor allele frequency under 0.05. After computing the test statistic, we

used Bonferroni adjustment to correct for multiple comparisons, and threshold of 0.001 was

applied to identify outliers SNPs.

Bacterial virulence estimation by injection

Bacteria were grown with constant agitation at 22°C for 24 h in Zobell medium. Cultures were

washed by centrifugation (10 min, 1500 g) and resuspended in artificial, sterile seawater to

obtain OD_{600nm} = 0.1. Fifty microliters of bacterial suspensions (10^7 cfu, checked by plating on

Zobell agar) were injected intramuscularly into anesthetized specific pathogen-free (SPF)

oysters produced and maintained in the IFREMER facility [79]. After injection, the oysters were

transferred to aquaria (10 oysters per 2.5 L aquarium, duplicate aquaria per bacterial strain)

containing 2 L of aerated, UV-treated seawater at 22°C and kept under static conditions for 5

days.

Copper resistance phenotyping

Paper discs (6 mm) were loaded with $CuSO_4$ by soaking them overnight in 50 mM $CuSO_4$.

Cells from stationary phase cultures at DO = 1 were washed twice with SSW before being

spread onto agar plates containing 15 g.L^{-1} bactopeptone and 0.5 M NaCl. Using sterile

tweezers, copper-impregnated disks were placed on the bacterial lawn before incubation at

20°C. After 5 days of incubation, the diameter of the inhibition zone around the disk was

measured.

Statistic tests

Basic statistical analysis of all of the data were performed using R v4.1.2. Data were checked

for normality and homogeneity of variance before analysis. Wilcoxon-Mann-Whitney was

performed to evaluate if recombination ratios, IS number in genomes, virulence toward oysters

and halo size (representing strains sensitivity to copper) differed among *Va francensis*

lineages and between *Va francensis* lineages and other strains.

Acknowledgments

We first thank the NRLs for mollusk diseases that contributed to the questionnaire survey (Bojan Adžić, Raquel Aranguren, Giuseppe Arcangeli, Deborah Cheslett, Marc Engelsma, Mark Fordyce, Ana Grade, Michael Gubbins, Lone Madsen, Petya Orozova, and Ewa Paździor). We thank Nicolas Bierne for helpful discussion and Viviane Boulo, Jean-Michel Escoubas, Nicole Faury and Gaelle Courtay for their valuable technical assistance. We also thank Bruno Petton and staff of IFREMER Bouin, who contributed to animal production and delivery. We thank the Bio-Environment platform (University of Perpignan Via Domitia) and Jean-François Allienne for support in library preparation and sequencing. We are grateful to the Ifremer Bioinformatics Core Facility (SeBiMER; https://ifremer-bioinformatics.github.io/) for providing technical assistance and scientific support in bioinformatics analysis. We also acknowledge the Pôle de Calcul et de Données Marines (https://wwz.ifremer.fr/en/Research-Technology/Research-Infrastructures/Digital-infrastructures/Computation-Centre) for providing DATARMOR computing and storage resources. This work was funded by Europe DG-Sante (through European Union Reference Laboratory for mollusk diseases funding) and French Ministry DPMA (MIRAGE project) and DGAl through NRL funding. Moreover, A.M., PhD, was supported by IFREMER and Région Nouvelle Aquitaine. This study is part of the « Laboratoire d'Excellence (LabEx) » TULIP (ANR-10-LABX-41) framework.

Bibliographic references

1. Le Roux F, Wegner KM, Baker-Austin C, Vezzulli L, Osorio CR, Amaro C, Ritchie JM, Defoirdt T, Destoumieux-Garzón D, Blokesch M. 2015. The emergence of Vibrio pathogens in Europe: ecology, evolution, and pathogenesis (Paris, 11–12th March 2015). Frontiers in microbiology 6:830.

2. Frans I, Michiels CW, Bossier P, Willems KA, Lievens B, Rediers H. 2011. *Vibrio anguillarum* as a fish pathogen: virulence factors, diagnosis and prevention. Journal of Fish Diseases 34:643-661.

3. Hewson I, Button JB, Gudenkauf BM, Miner B, Newton AL, Gaydos JK, Wynne J, Groves CL, Hendler G, Murray M, Fradkin S, Breitbart M, Fahsbender E, Lafferty KD, Kilpatrick AM, Miner CM, Raimondi P, Lahner L, Friedman CS, Daniels S, Haulena M, Marliave J, Burge CA, Eisenlord ME, Harvell CD. 2014. Densovirus associated with sea-star wasting disease and mass mortality. Proceedings of the National Academy of Sciences of the United States of America 111:17278-17283.

4. Lafferty KD, Kuris AM. 1993. Mass mortality of abalone Haliotis cracherodii on the California Channel Islands: tests of epidemiological hypotheses. Marine Ecology-Progress Series 96:239-239.

5. Lafferty KD, Harvell CD, Conrad JM, Friedman CS, Kent ML, Kuris AM, Powell EN, Rondeau D, Saksida SM. 2015. Infectious Diseases Affect Marine Fisheries and Aquaculture Economics. Annual Review of Marine Science 7:471-496.

6. Murray AG, Peeler EJ. 2005. A framework for understanding the potential for emerging diseases in aquaculture. Preventive Veterinary Medicine 67:223-235.

7. Bruto M, James A, Petton B, Labreuche Y, Chenivesse S, Alunno-Bruscia M, Polz MF, Le Roux F. 2017. *Vibrio crassostreae,* a benign oyster colonizer turned into a pathogen after plasmid acquisition. The ISME Journal 11:1043-1052.

8. Lemire A, Goudenège D, Versigny T, Petton B, Calteau A, Labreuche Y, Le Roux F. 2015. Populations, not clones, are the unit of vibrio pathogenesis in naturally infected oysters. The ISME Journal 9:1523-1531.

9. Petton B, Bruto M, James A, Labreuche Y, Alunno-Bruscia M, Le Roux F. 2015. *Crassostrea gigas* mortality in France: the usual suspect, a herpes virus, may not be the killer in this polymicrobial opportunistic disease. Frontiers in microbiology 6:686.

10. Reen FJ, Almagro-Moreno S, Ussery D, Boyd EF. 2006. The genomic code: inferring Vibrionaceae niche specialization. Nat Rev Microbiol 4:697-704.

11. Garnier M, Labreuche Y, Nicolas JL. 2008. Molecular and phenotypic characterization of *Vibrio aestuarianus* subsp. *francensis* subsp. nov., a pathogen of the oyster *Crassostrea gigas*. Syst Appl Microbiol 31:358-65.

12. Lasa A, di Cesare A, Tassistro G, Borello A, Gualdi S, Furones D, Carrasco N, Cheslett D, Brechon A, Paillard C, Bidault A, Pernet F, Canesi L, Edomi P, Pallavicini A, Pruzzo C, Vezzulli L. 2019. Dynamics of the Pacific oyster pathobiota during mortality episodes in Europe assessed by 16S rRNA gene profiling and a new target enrichment next-generation sequencing strategy. Environ Microbiol 21:4548-4562.

13. Mandas D, Salati F, Polinas M, Sanna MA, Zobba R, Burrai GP, Alberti A, Antuofermo E. 2020. Histopathological and Molecular Study of Pacific Oyster Tissues Provides Insights into *V. aestuarianus* Infection Related to Oyster Mortality. Pathogens 9.

14. Dégremont L, Azéma P, Maurouard E, Travers M-A. 2020. Enhancing resistance to *Vibrio aestuarianus* in *Crassostrea gigas* by selection. Aquaculture 526:735429.

15. Parizadeh L, Tourbiez D, Garcia C, Haffner P, Dégremont L, Le Roux F, Travers M-A. 2018. Ecologically realistic model of infection for exploring the host damage caused by *Vibrio aestuarianus*. Environmental Microbiology 20:4343-4355.

16. Engering A, Hogerwerf L, Slingenbergh J. 2013. Pathogen-host-environment interplay and disease emergence. Emerging microbes & infections 2:e5-e5.

17. Sheppard SK, Guttman DS, Fitzgerald JR. 2018. Population genomics of bacterial
host adaptation. Nature Reviews Genetics 19:549-565.
18. Martínez JL. 2013. Bacterial pathogens: from natural ecosystems to human hosts.
Environmental Microbiology 15:325-333.
19. Tibayrenc M, Ayala FJ. 2012. Reproductive clonality of pathogens: A perspective on
pathogenic viruses, bacteria, fungi, and parasitic protozoa. Proceedings of the
National Academy of Sciences 109:E3305-E3313.
20. Cohan FM. 2002. Sexual isolation and speciation in bacteria. Genetics of Mate
Choice: From Sexual Selection to Sexual Isolation:359-370.
21. Polz MF, Alm EJ, Hanage WP. 2013. Horizontal gene transfer and the evolution of
bacterial and archaeal population structure. Trends Genet 29:170-5.
22. Goudenège D, Travers MA, Lemire A, Petton B, Haffner P, Labreuche Y, Tourbiez D,
Mangenot S, Calteau A, Mazel D, Nicolas JL, Jacq A, Le roux F. 2015. A single
regulatory gene is sufficient to alter *Vibrio aestuarianus* pathogenicity in oysters.
Environ Microbiol 17:4189-99.
23. Saulnier D, De Decker S, Haffner P. 2009. Real-time PCR assay for rapid detection
and quantification of Vibrio aestuarianus in oyster and seawater: A useful tool for
epidemiologic studies. Journal of Microbiological Methods 77:191-197.
24. Garnier M, Labreuche Y, Garcia C, Robert M, Nicolas JL. 2007. Evidence for the
involvement of pathogenic bacteria in summer mortalities of the Pacific oyster
Crassostrea gigas. Microb Ecol 53:187-96.
25. Moran NA. 2002. Microbial minimalism: genome reduction in bacterial pathogens.
Cell 108:583-6.
26. Floriano AM, Castelli M, Krenek S, Berendonk TU, Bazzocchi C, Petroni G, Sassera
D. 2018. The Genome Sequence of "Candidatus *Fokinia solitaria*": Insights on
Reductive Evolution in Rickettsiales. Genome Biol Evol 10:1120-1126.
27. Hauck S, Maiden MCJ. 2018. Clonally Evolving Pathogenic Bacteria, p 307-325. *In*
Rampelotto PH (ed), Molecular Mechanisms of Microbial Evolution doi:10.1007/978-
3-319-69078-0_12. Springer International Publishing, Cham.
28. Bergstrom CT, McElhany P, Real LA. 1999. Transmission bottlenecks as
determinants of virulence in rapidly evolving pathogens. Proceedings of the National
Academy of Sciences 96:5095-5100.
29. Bäumler A, Fang FC. 2013. Host Specificity of Bacterial Pathogens. Cold Spring
Harbor Perspectives in Medicine 3.
30. Mutreja A, Kim DW, Thomson NR, Connor TR, Lee JH, Kariuki S, Croucher NJ, Choi
SY, Harris SR, Lebens M, Niyogi SK, Kim EJ, Ramamurthy T, Chun J, Wood JL,
Clemens JD, Czerkinsky C, Nair GB, Holmgren J, Parkhill J, Dougan G. 2011.
Evidence for several waves of global transmission in the seventh cholera pandemic.
Nature 477:462-5.
31. FranceAgriMer. 2021. https://www.franceagrimer.fr/. https://www.franceagrimer.fr/.
Accessed
32. Namouchi A, Didelot X, Schöck U, Gicquel B, Rocha EP. 2012. After the bottleneck:
Genome-wide diversification of the Mycobacterium tuberculosis complex by mutation,
recombination, and natural selection. Genome Res 22:721-34.
33. Claisse D, Alzieu C. 1993. Copper contamination as a result of antifouling paint
regulations? Marine Pollution Bulletin 26:395-397.
34. Hall LW, Anderson RD. 1999. A Deterministic Ecological Risk Assessment for
Copper in European Saltwater Environments. Marine Pollution Bulletin 38:207-218.
35. Rossi N, Jamet J-L. 2008. *In situ* heavy metals (copper, lead and cadmium) in
different plankton compartments and suspended particulate matter in two coupled
Mediterranean coastal ecosystems (Toulon Bay, France). Marine Pollution Bulletin
56:1862-1870.
36. Weng N, Wang W-X. 2014. Variations of trace metals in two estuarine environments
with contrasting pollution histories. Science of The Total Environment 485-486:604-
614.

37. Shi B, Wang T, Zeng Z, Zhou L, You W, Ke C. 2019. The role of copper and zinc
accumulation in defense against bacterial pathogen in the fujian oyster (*Crassostrea
angulata*). Fish Shellfish Immunol 92:72-82.
38. Destoumieux-Garzón D, Canesi L, Oyanedel D, Travers M-A, Charrière GM, Pruzzo
C, Vezzulli L. 2020. Vibrio–bivalve interactions in health and disease. Environmental
Microbiology 22:4323-4341.
39. Yeh S-T, Liu C-H, Chen J-C. 2004. Effect of copper sulfate on the immune response
and susceptibility to *Vibrio alginolyticus* in the white shrimp *Litopenaeus vannamei*.
Fish & Shellfish Immunology 17:437-446.
40. Bondarczuk K, Piotrowska-Seget Z. 2013. Molecular basis of active copper
resistance mechanisms in Gram-negative bacteria. Cell Biol Toxicol 29:397-405.
41. Argüello JM, Raimunda D, Padilla-Benavides T. 2013. Mechanisms of copper
homeostasis in bacteria. Front Cell Infect Microbiol 3:73.
42. Vanhove AS, Duperthuy M, Charriere GM, Le Roux F, Goudenege D, Gourbal B,
Kieffer-Jaquinod S, Coute Y, Wai SN, Destoumieux-Garzon D. 2015. Outer
membrane vesicles are vehicles for the delivery of Vibrio tasmaniensis virulence
factors to oyster immune cells. Environ Microbiol 17:1152-65.
43. Didelot X, Pang B, Zhou Z, McCann A, Ni P, Li D, Achtman M, Kan B. 2015. The
Role of China in the Global Spread of the Current Cholera Pandemic. PLOS Genetics
11:e1005072.
44. Namouchi A, Didelot X, Schöck U, Gicquel B, Rocha EPC. 2012. After the bottleneck:
Genome-wide diversification of the Mycobacterium tuberculosis complex by mutation,
recombination, and natural selection. Genome Research 22:721-734.
45. Didelot X, Bowden R, Street T, Golubchik T, Spencer C, McVean G, Sangal V, Anjum
MF, Achtman M, Falush D, Donnelly P. 2011. Recombination and Population
Structure in Salmonella enterica. PLOS Genetics 7:e1002191.
46. Siguier P, Gourbeyre E, Chandler M. 2014. Bacterial insertion sequences: their
genomic impact and diversity. FEMS Microbiology Reviews 38:865-891.
47. Touchon M, Rocha EPC. 2007. Causes of Insertion Sequences Abundance in
Prokaryotic Genomes. Molecular Biology and Evolution 24:969-981.
48. Travers MA, Tourbiez D, Parizadeh L, Haffner P, Kozic-Djellouli A, Aboubaker M,
Koken M, Degremont L, Lupo C. 2017. Several strains, one disease: experimental
investigation of *Vibrio aestuarianus* infection parameters in the Pacific oyster,
Crassostrea gigas. Vet Res 48:32.
49. Garcia C, Thébault A, Dégrémont L, Arzul I, Miossec L, Robert M, Chollet B, François
C, Joly J-P, Ferrand S, Kerdudou N, Renault T. 2011. Ostreid herpesvirus 1 detection
and relationship with *Crassostrea gigas* spat mortality in France between 1998 and
2006. Veterinary Research 42:73.
50. Barbosa Solomieu V, Renault T, Travers MA. 2015. Mass mortality in bivalves and
the intricate case of the Pacific oyster, *Crassostrea gigas*. J Invertebr Pathol 131:2-
10.
51. Azéma P, Maurouard E, Lamy J-B, Dégrémont L. 2017. The use of size and growing
height to improve *Crassostrea gigas* farming and breeding techniques against OsHV-
1. Aquaculture 471:121-129.
52. Azema P, Travers MA, De Lorgeril J, Tourbiez D, Degremont L. 2015. Can selection
for resistance to OsHV-1 infection modify susceptibility to *Vibrio aestuarianus*
infection in Crassostrea gigas? First insights from experimental challenges using
primary and successive exposures. Vet Res 46:139.
53. Azema P, Lamy JB, Boudry P, Renault T, Travers MA, Degremont L. 2017. Genetic
parameters of resistance to *Vibrio aestuarianus,* and OsHV-1 infections in the Pacific
oyster, *Crassostrea gigas*, at three different life stages. Genet Sel Evol 49:23.
54. Azéma P, Travers M-A, Benabdelmouna A, Dégrémont L. 2016. Single or dual
experimental infections with *Vibrio aestuarianus* and OsHV-1 in diploid and triploid
Crassostrea gigas at the spat, juvenile and adult stages. Journal of Invertebrate
Pathology 139:92-101.

55. Anderson PK, Cunningham AA, Patel NG, Morales FJ, Epstein PR, Daszak P. 2004.
Emerging infectious diseases of plants: pathogen pollution, climate change and
agrotechnology drivers. Trends in Ecology & Evolution 19:535-544.
56. Wingett SW, Andrews S. 2018. FastQ Screen: A tool for multi-genome mapping and
quality control. F1000Res 7:1338.
57. Bush SJ. 2020. Read trimming has minimal effect on bacterial SNP-calling accuracy.
Microbial Genomics 6.
58. Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM,
Nikolenko SI, Pham S, Prjibelski AD, Pyshkin AV, Sirotkin AV, Vyahhi N, Tesler G,
Alekseyev MA, Pevzner PA. 2012. SPAdes: a new genome assembly algorithm and
its applications to single-cell sequencing. Journal of computational biology : a journal
of computational molecular cell biology 19:455-477.
59. Gurevich A, Saveliev V, Vyahhi N, Tesler G. 2013. QUAST: quality assessment tool
for genome assemblies. Bioinformatics 29:1072-5.
60. Kolmogorov M, Yuan J, Lin Y, Pevzner PA. 2019. Assembly of long, error-prone
reads using repeat graphs. Nature Biotechnology 37:540-546.
61. Vaser R, Sovic I, Nagarajan N, Sikic M. 2017. Fast and accurate de novo genome
assembly from long uncorrected reads. Genome Research
doi:10.1101/gr.214270.116.
62. Simão FA, Waterhouse RM, Ioannidis P, Kriventseva EV, Zdobnov EM. 2015.
BUSCO: assessing genome assembly and annotation completeness with single-copy
orthologs. Bioinformatics 31:3210-3212.
63. Seemann T. 2014. Prokka: rapid prokaryotic genome annotation. Bioinformatics
30:2068-2069.
64. Cantalapiedra CP, Hernández-Plaza A, Letunic I, Bork P, Huerta-Cepas J. 2021.
eggNOG-mapper v2: Functional Annotation, Orthology Assignments, and Domain
Prediction at the Metagenomic Scale. Molecular Biology and Evolution 38:5825-5829.
65. Huerta-Cepas J, Szklarczyk D, Heller D, Hernández-Plaza A, Forslund SK, Cook H,
Mende DR, Letunic I, Rattei T, Jensen Lars J, von Mering C, Bork P. 2018. eggNOG
5.0: a hierarchical, functionally and phylogenetically annotated orthology resource
based on 5090 organisms and 2502 viruses. Nucleic Acids Research 47:D309-D314.
66. Xie Z, Tang H. 2017. ISEScan: automated identification of insertion sequence
elements in prokaryotic genomes. Bioinformatics 33:3340-3347.
67. Syberg-Olsen MJ, Garber AI, Keeling PJ, McCutcheon JP, Husnik F. 2021.
Pseudofinder: detection of pseudogenes in prokaryotic genomes. bioRxiv
doi:10.1101/2021.10.07.463580:2021.10.07.463580.
68. Goris J, Konstantinidis KT, Klappenbach JA, Coenye T, Vandamme P, Tiedje JM.
2007. DNA-DNA hybridization values and their relationship to whole-genome
sequence similarities. Int J Syst Evol Microbiol 57:81-91.
69. Darling AE, Mau B, Perna NT. 2010. progressiveMauve: Multiple Genome Alignment
with Gene Gain, Loss and Rearrangement. PLOS ONE 5:e11147.
70. Didelot X, Wilson DJ. 2015. ClonalFrameML: Efficient Inference of Recombination in
Whole Bacterial Genomes. PLOS Computational Biology 11:e1004041.
71. Rambaut A, Lam TT, Max Carvalho L, Pybus OG. 2016. Exploring the temporal
structure of heterochronous sequences using TempEst (formerly Path-O-Gen). Virus
Evolution 2.
72. Minin VN, Bloomquist EW, Suchard MA. 2008. Smooth Skyride through a Rough
Skyline: Bayesian Coalescent-Based Inference of Population Dynamics. Molecular
Biology and Evolution 25:1459-1471.
73. Drummond AJ, Nicholls GK, Rodrigo AG, Solomon W. 2002. Estimating Mutation
Parameters, Population History and Genealogy Simultaneously From Temporally
Spaced Sequence Data. Genetics 161:1307-1320.
74. Rambaut A, Drummond AJ, Xie D, Baele G, Suchard MA. 2018. Posterior
Summarization in Bayesian Phylogenetics Using Tracer 1.7. Systematic Biology
67:901-904.

75. Baele G, Lemey P, Bedford T, Rambaut A, Suchard MA, Alekseyenko AV. 2012.
Improving the Accuracy of Demographic and Molecular Clock Model Comparison
While Accommodating Phylogenetic Uncertainty. Molecular Biology and Evolution
29:2157-2167.
76. Baele G, Li WLS, Drummond AJ, Suchard MA, Lemey P. 2012. Accurate Model
Selection of Relaxed Molecular Clocks in Bayesian Phylogenetics. Molecular Biology
and Evolution 30:239-243.
77. Subramanian S. 2016. The effects of sample size on population genomic analyses –
implications for the tests of neutrality. BMC Genomics 17:123.
78. Paradis E. 2010. pegas: an R package for population genetics with an integrated–
modular approach. Bioinformatics 26:419-420.
79. Petton B, Pernet F, Robert R, Boudry P. 2013. Temperature influence on pathogen
transmission and subsequent mortalities in juvenile Pacific oysters *Crassostrea*
gigas. Aquaculture Environment Interactions 3:257-273.
713