Emergence and clonal expansion in Europe of *Vibrio aestuarianus* lineages pathogenic for oysters.

Aurélie Mesnil[1,2], Maude Jacquot[1], Céline Garcia[1], Delphine Tourbiez[1], Lydie Canier[1], Audrey Bidois[2], Lionel Dégremont[1], Deborah Cheslett[3], Michelle Geary[3], Alessia Vetri[4], Ana Roque[5], Dolors Furones[5], Alison Garden[6], Petya Orozova[7], Isabelle Arzul[1], Mathieu Sicard[8], Guillaume Charrière[2], Delphine Destoumieux-Garzón[2], Marie-Agnès Travers[2*]

ASIM, Ifremer, Station de La Tremblade, avenue de Mus du Loup, 17390 La Tremblade, France

IHPE, Université de Montpellier, CNRS, IFREMER, Univ Perpignan Via Domitia, Montpellier, France

Marine Institute MI, Rinville, Oranmore, Co. Galway, H91 R673, Ireland

Istituto Zooprofilattico Sperimentale delle Venezie (IZSVe), Viale dell'Università 10, 35020 Legnaro (PD), Italy

IRTA La Ràpita. Aquaculture Program. 43540 – San Carles de la Ràpita, Spain.

Marine Laboratory, Marine Scotland Science, Aberdeen, UK.

National Diagnostic and Research Veterinary Medical Institute, Sofia, Bulgaria.

ISEM, Université de Montpellier, CNRS, IRD, Montpellier, France.

* Corresponding author: marie.agnes.travers@ifremer.fr

Competing interests: The authors declare no conflict of interest.

Abstract

Crassostrea gigas oysters, with 4.7 million tons harvested per year, represent a considerable food supply worldwide. Since 2001 *Vibrio aestuarianus* has emerged as a pathogenic bacterium causing adult oyster mortality events in France and Ireland, and its impact on oyster aquaculture has increased in Europe since its reemergence in 2012. To better understand the evolutionary mechanisms leading to the emergence and persistence over time of this pathogen, we conducted a survey with national reference laboratories (NRLs) at the European scale for mollusk diseases. We analyzed genomes of *V. aestuarianus* isolated from multiple countries, since 2001, with or without bivalve mortalities, and from multiple environmental compartments. Using a combination of comparative genomics and population genetics approaches, we show that two oyster pathogen lineages have almost clonally expanded in Europe, after a probable recent selective bottleneck. Low mutation and recombination rates, as well as purifying selective pressure, have preserved virulent selected genotypes. Furthermore, we identified a specific *cus-cop*-containing island conferring copper resistance to *V. aestuarianus* that could have favored the emergence of *V. aestuarianus* pathogenic lineages adapted to oysters.

Introduction

In an increasingly connected world, marine populations appear to be under increasing threat of emerging infectious disease (EID)(1), as illustrated by the mass mortality events currently affecting corals, sea-stars, mollusks and fish (2-4). Aquaculture is particularly threatened by EID, as it is practiced in an open and highly connected environment favoring the circulation of pathogens (5). Moreover, aquacultural practices frequently result in high population densities as well as many animal transfers, which increase the risk of regular pathogen lineage introductions and dispersal (6).

The emergence of new diseases is linked to changes in the interactions between pathogens,

hosts and the environment, which are usually caused by one or more of the following

situations: (a) an encounter between a pathogen and a new host, *i.e.,* cross-species

transmission, (b) access by a pathogen to a new area, *i.e.,* introduction, and (c) the

acquisition by a pathogen of a new trait that increases its capacity to colonize a host, *i.e.,* its

adaptation to a host (7). In bacteria, trait acquisition is often achieved by horizontal transfers

(through transformation, conjugation or transduction) of several genes favoring rapid

adaptation to new niches and hosts (8). For example, in the genus *Vibrio*, one ecological

population from the clade Splendidus that causes oyster spat mortalities have acquired a

mobilisable virulence plasmid (9). Mutations and genomic rearrangements can also influence

host adaptation by offering new evolutionary trajectory opportunities or by causing

progressive loss of gene function (10). Trait acquisition can be followed by "clonal

expansion", *i.e.,* evolution without recombination and the spatial and temporal expansion of

the lineage considered (11). The most documented example is related to *V. cholerae* clonal

emergence after acquisition of major virulence factors, cholera toxin (CT) and toxin

coregulated pilus (TCP) (12). One consequence of strict clonality is the strong effect of

population bottlenecks and selective sweeps (a new favorable substitution or set of

substitutions sweeps to fixation in a population and purges variation for the entire

chromosome in the clonal population) in reducing genetic diversity in the resulting population

(13).

Oysters (*Crassostrea gigas*) are farmed worldwide and are frequently impacted by bacterial

pathogens of the genus *Vibrio* (14-16). Among them, *Vibrio aestuarianus* subsp. *francensis*

has caused recurrent mortality of adult oysters since 2001 (17). Mortality rates of

approximately 25% (18) have had an important economic impact, as they affect market-size

oysters. Having previously been restricted to France, this disease has now spread across

European countries (France, Italy, Ireland, and Spain) (19, 20). Strains of another *V.*

aestuarianus subspecies (subsp. *cardii*) were isolated from diseased cockles. In addition

strains of *V. aestuarianus* subsp. *aestuarianus* were isolated from water, sediment,

zooplankton and bivalve tissues (including from oysters) outside disease outbreaks (21, 22).

First phenotypic and genetic studies allowed to describe *V. aestuarianus* subspecies without

clarifying population structure (21).

In previous field studies, pathogenic strains of *V. aestuarianus francensis* were found to be

able to colonize sentinel oysters, but they remained rare in other environmental

compartments sampled in the oyster surroundings (sediment and water column), suggesting

that oysters are its preferred habitat (23). Moreover, it has been proposed that *V.*

aestuarianus francensis can persist in oyster tissues during winter (23). These observations

suggest strong adaptation of *V. aestuarianus francensis* to its host and raise questions about

host specialization for this pathogen. However, the evolutionary processes that have led to

the emergence and persistence of pathogenic strains in *V. aestuarianus* remain unknown,

and only a limited number of genomes are available, which does not reflect overall diversity

across diverse isolation contexts (17, 22). This lack of genomic data hinders a complete

understanding of the genetic diversity and population structure of *V. aestuarianus,* and

prevents us from uncovering the evolutionary processed underpinning the emergence of

pathogenic *V. aestuarianus francensis.*

In the present study, we analyzed 54 genomes of isolates collected since 2001 across

Europe. We combined phylogenetic and population genetic approaches to decipher

evolutionary processes associated with the origin of the emergence of this oyster pathogen.

Our study shows that bacterial strains belonging to *V. aestuarianus francensis* exhibit

numerous features commonly reported in specialized pathogens. In particular, our data

suggest the acquisition of genes involved in copper resistance as one mechanism of *V.*

aestuarianus adaptation to oysters.

Results

European spread of *V. aestuarianus* oyster pathogens

To assess the geographic distribution of pathogenic *V. aestuarianus* in Europe, a
questionnaire survey was sent in 2021 to all European laboratories in charge of the
diagnosis and surveillance of mollusk diseases (supplementary material). Eight out of the 12
respondent laboratories reported the detection of *V. aestuarianus* during oyster mortality
events. The earliest detections were reported by France and Ireland in 2001, and most
recent detection by country is the suspicion of the pathogen reported by Denmark in 2021
(fig. 1, A). An increase in the number of countries in which the bacteria was detected was
observed after 2011, leading to a total of 8 European countries currently reporting *V.
aestuarianus*. However, it is important to notice that the reported date of first detection could
be under-estimated, as implementation of surveillance and testing for the pathogen has
varied between countries and was only implemented in the majority of the laboratories
following the development of a diagnostic tool (24).

**V. aestuarianus *diversity is structured according to environmental origin and isolation
context.*** To reconstruct the overall population structure of *V. aestuarianus* in Europe, we
assessed genetic diversity at the whole genome level using *V. aestuarianus* isolates that
were as diverse as possible. Strain selection was performed to equilibrate strain origins
(country, year, matrix and context): we used 54 isolates that originated from 7 countries, with
7 contrasting environmental origins (5 bivalve species, zooplankton and free-living bacteria
in the water column) and from different contexts (investigation of oyster mortalities or not).
The considered samples were isolated between 2001 and 2021 (plus the type strain isolated
in 1983 in the United States). Whole genomes were sequenced and *de novo* assembled.
This allowed us to obtain 15 4.5-Mb complete genomes (with 2 to 4 contigs) and 39 draft

genomes (with up to 150 contigs) (supplementary table 1). Whole genomes were aligned,

and relationships between isolates were studied by building a maximum likelihood

phylogeny, using strain 03_008T as an outgroup (fig. 2). Globally, genomes were close to

each other in the tree and showed highly similar nucleotide compositions, as indicated by

average nucleotide identities (ANIs) higher than 97.1% for all pairwise comparisons

(supplementary table 2).

To confirm population structure, we focused on allele frequency differences within *V.*

aestuarianus genomes. We applied the linkage model of Structure to our data and identified

six ancestral populations in our samples. Strains isolated during oyster mortality events

(hereafter referred to as oyster clinical isolates, OCIs) were grouped into a unique population

predicted to have descended from a unique ancestor (fig S1). Almost all other strains

descended from multiple ancestors, including the OCI ancestor.

The OCI group was composed of two sublineages supported by a bootstrap value of 100%

and named clades A (OCI_A, n = 17) and B (OCI_B, n = 8) as previously described (22) (fig.

2). To test whether OCI/No_OCI delineation correlated with *Vibrio* virulence potential, we

phenotyped most isolates by using oyster experimental infections. Virulence potential

perfectly matched with phylogenic delineation: the OCI clades A and B contained all virulent

isolates, which induced more than 50% mortality in oysters (fig. 2).

To determine the relationships between environmental factors and observed genetic

diversity, we performed principal component analysis of genetic polymorphism using 183 109

SNPs. Principal component 1 (PC1) segregated the isolates according to isolation context

and environmental origin (fig 3) but not according to isolation year or isolation country.

Furthermore, analysis of molecular variance (AMOVA) performed on the sequence dataset

revealed that isolation context and environmental origin explained the most significant ($p <$

0.005) amounts of genetic variation (table 1). This indicated the absence of spatial and

temporal delineation of the OCI group.

Altogether, our data show the European expansion of *V. aestuarianus* lineages pathogenic

to oysters that has spread without much genetic variation since the first associated oyster

mortality event in 2001.

Clonal expansion of *V. aestuarianus francensis* lineages

Recombination is an important source of genetic diversity in bacterial populations that may

drive the emergence of pathogens. We therefore quantified and localized recombination

events along *V. aestuarianus* genome alignments.

The impact of recombination on OCI diversity was weak. Recombination events were

almost exclusively detected outside OCIs, with, on average, 342.5 recombination events per

genome. In the NO_OCI alignment, 164 704 polymorphic sites are engendered by

recombination. In contrast, in OCI genomes, only 7.8 and 3.3 recombination events were

detected on average for clades A and B, respectively, generating only 5 207 and 1 232

polymorphic sites (0.1% of the genome nucleotides). Recombination in genomes is known to

generate phylogenetic inconsistencies between taxa that can be visible in a phylogenetic

network as the presence of alternative pathways (25). The network that we constructed from

the whole-genome alignment (fig S2) confirmed that recombination does not affect OCI

relationships, with few alternative pathways between these taxa. One known consequence of

the absence of recombination impacts is strong linkage between loci in non-recombinant

genomes (26). In OCI_A and, to a lesser extent, in OCI_B genomes, strong linkage between

single nucleotide polymorphisms (SNPs) was observed (fig S3). In contrast, the

recombination impact detected in all other genomes was confirmed with phylogenetic

networks, which showed inconsistencies between NO_OCI taxa (fig S2) and low linkage

between loci in these genomes (fig S3). However, recombination might have played a role in

the emergence of this pathogenic group, as 488 recombination events were detected at the

node corresponding to the hypothetical common ancestor of all OCI isolates (fig. 4).

Mutations also generated little genetic diversity among OCIs. Indeed, apart from the

recombinant segments, 666 and 83 polymorphic sites were observed in OCI_A and OCI_B,

respectively, compared with 19 775 in NO_OCI isolates.

Altogether, our data support the clonal expansion of two *V. aestuarianus francensis* lineages

that are pathogenic to oysters.

Bottlenecks and purifying selection may limit pathogenic *V. aestuarianus*

genetic diversity

Selection is a major evolutionary force that controls the genetic diversity generated by

mutation and recombination. Here we used Tajima's D to compare the genetic diversity

observed in *V. aestuarianus* to the genetic diversity expected under neutral evolution and

constant population size.

The deviation from neutrality was by far the largest among the OCI_A and OCI_B and could

be due to a combination of recent population expansion after selective bottlenecks and

purifying selection, as shown by the very negative values in these two clades (fig. 5, A). To

determine whether the low genetic diversity of OCIs was due to demographic effects

(affecting whole genomes) or to selection effects (affecting some of the genes), we adopted

a gene-based approach to investigate selection pressures at the gene level. While most of

the 1 974 core genes were under neutral evolution outside of the OCI clades, the vast

majority were monomorphic within OCI_A and OCI_B (fig. 5 B, supplementary table 3),

which is consistent with the low genetic diversity observed along the alignment of complete

genomes (fig. 5 A). Purifying selection could also act, amplifying low genetic diversity in OCI

clades. Indeed, contrasting selection patterns were observed among genes in OCIs, with a

small portion of genes showing polymorphisms, some of which were under neutral evolution

and others of which were under purifying selection (fig. 5 B, supplementary table 4).

Overall, our data indicate that pathogenic clades of *V. aestuarianus* have expanded following

a selective bottleneck, as revealed by a high number of monomorphic genes. However,

purifying selection pressure also tends to maintain the genotype of the ancestor of these

groups, as indicated by variation in gene diversity patterns.

Oyster-pathogenic *V. aestuarianus* have a conserved pangenome

Considering (i) the low genetic diversity of OCIs, (ii) their restricted (currently known)

habitats, and (iii) the scarcity of recombination events after the expansion of the two OCI

clades, we hypothesized that oyster-pathogen *V. aestuarianus* exhibit a closed pangenome

preserved from horizontal gene transfer (HGT).

OCIs were found to have a closed pangenome, mainly with genes being shared between

them and almost no genes specific to one strain, as determined by dataset pangenome

analysis (fig S4). In contrast, NO_OCI strains exhibited many strain-specific genes.

The absence of detection of homologous recombination and the closed pangenome indicate

that horizontal gene exchange in the OCI clades is currently absent or transparent (if

identical sequences are exchanged), supporting sexual isolation. In contrast, the numerous

homologous recombination events detected in NO_OCI genomes, as well as the large

quantity of specific genes that these genomes contain, showed that NO_OCI strains do not

come from strictly clonal reproduction but that HGT are still frequent.

According to these data, the two OCI clades can be considered pathogenic clonal complexes

that undergo clonal reproduction with limited mutation, recombination and HGT. Such sexual

isolation is a hallmark of specialist pathogens, which live in a very limited number of habitats

and have few opportunities to exchange genetic material with other populations. This

strongly suggests that oyster-pathogenic *V. aestuarianus* have evolved as oyster specialists.

Oyster-pathogenic *V. aestuarianus* have genomic features of specialist pathogens

We further tested the hypothesis of OCI specialization by analyzing the genomic features of specialists: insertion sequences (ISs) and pseudogene enrichment (27). We found higher numbers of ISs in the OCI_A and OCI_B genomes than in NO_OCI group 1 genomes corresponding to strains isolated in contexts without bivalve mortality (Kruskal test, $p = 0.01$ and 0.02, respectively). Interestingly, in NO_OCI group 2 containing strains isolated during cockle mortality events, ISs were more abundant than in the OCI_A and OCI_B genomes ($p = 0.02$ and 0.03, respectively) (fig. 6). IS families were conserved between OCI genomes, with most ISs belonging to the IS66, IS902 and ISAS1 families (supplementary table 5 and fig S5). IS conservation in OCI clades suggests that IS enrichment and expansion in OCI genomes stopped after pathogen emergence. The dynamics of the ISs appeared to be different in the cockle isolates. Indeed, the number and families of ISs represented appeared to be poorly conserved. We suspect that IS flux is active in cockle isolates, with both horizontal exchanges and intragenomic expansion.

To check whether IS enrichment is involved in pseudogenization, we crossed IS and pseudogene genomic coordinates and determined the number of ISs with pseudogenes around them (from 2 000 bp before the IS start to 2 000 bp after the IS end) (supplementary tables 6 and 7). Regardless of the genome and chromosome considered, the vast majority of ISs had pseudogenes nearby.

IS and pseudogene enrichment supports our hypothesis of specialization of the OCI clades.

Copper resistance acquisition: One potential step in the emergence of oyster-pathogenic clades

Trait acquisition is often the first step preceding selective bottlenecks and clonal expansion of host-adapted pathogens. To identify traits that may have allowed oyster pathogen emergence, we studied the *V. aestuarianus* pangenome and detected 139 genes that were

shared exclusively by OCI_A and OCI_B genomes. Most of them (106) were annotated as

hypothetical proteins. The remaining 33 genes encode proteins with functions related to

copper export, cell-bacteria interaction (LPS/antigen O, cell wall or peptidoglycan),

chemotaxis and DNA repair. The sequences of these 139 genes were identical or well

conserved between strains (supplementary table 8).

Some OCI-specific genes were located in a putative mobile genetic element (MGE), as

determined by crossing their coordinates with IS coordinates in an OCI genome (02_041)

(fig. 7 A). This putative MGE contains the genes of the Cus operon (*cusABCF,S/R)*

surrounded by ISs from the IS66 family (length = 15 kb), as well as the copper export genes

copA and *copG* and the regulator *cueR*, located near tn7 transposon genes (tnA, B, C and

E) (length = 10 kb), and is later referred to as a *cus-cop*-containing island.

The OCI ancestor could have acquired copper export genes by horizontal transfer, and

integration into its genome could be facilitated by its presence in the MGE. To investigate the

origin of *cus-cop*-containing islands, we used blast analyses of each operon and of the entire

region in 127 reference genomes of different *Vibrio* species (supplementary table 9 and fig

S6). The *cus-cop*-containing island presence is scattered along the Vibrio phylogeny and is

not specific of a monophyletic group. It appeared highly similar to some *V. fluvialis* and *V.*

furnissii genomic sequences with a nucleotide identity >95%. Moreover, this region was

close to plasmidic elements identified in *V. crassostreae* and *Vibrio* sp. dgh with a nucleotide

identity >89% (fig. 7B).

To test whether the genomic differences between the OCI and NO_OCI groups could lead to

phenotypic differences, we determined the copper resistance capacities of 27 *V.*

aestuarianus strains using an agar diffusion assay. Supporting our hypothesis, all OCIs (8

strains from OCI_ A and 5 from OCI_ B) appeared more resistant than NO_OCI strains to

copper in radial diffusion assays (fig. 7C).

Discussion

Here, we describe a genomic analysis of a *V. aestuarianus* collection isolated from multiple countries, in different epidemic contexts and from multiple environments since 2001. We report the European expansion and clonal evolution of two oyster-pathogenic lineages, belonging to the previously described *V. aestuarianus francensis* subspecies (17).

V. aestuarianus francensis pathogenic lineages OCI_A and OCI_B appear to have emerged as a consequence of the acquisition a specialized lifestyle trait. Indeed, we observed a markedly low level of genetic diversity among the 25 genomes of the OCI which was attributed to a low level of recombination. OCI were also characterized by a closed pangenome. This led us to formulate the hypothesis of a sexual isolation of OCI, a hallmark of specialized pathogens, which do not exchange genes with other bacterial populations. During specialization, pathogens make the transition from facultative parasitic life to a permanent association with a host. As they live continuously in a host, they can acquire compounds of intermediate metabolism from this host thus often discarding genes associated to biosynthetic pathways that have become facultative or redundant (25). Studies on pathogen specialization have mainly focused on highly specialized intracellular bacteria, often obligate parasite, which can undergo an extreme genome rationalization process (26). This process implies pseudogenization of genes that have become inessentials accumulating mutations or ISs, and can ultimately result in genome size reduction through gene loss (27-29). As a hallmark of host specialization, pseudogenization was observed in both OCI lineages. However, genome size reduction was not observed, probably because *V. aestuarianus francensis* pathogenic lineages are not obligate parasites. Indeed, whereas OCI have a restricted habitat (they are almost exclusively found in oysters), they have retained the ability to replicate in the absence of their host, at least in rich medium. Moreover, *V. aestuarianus francensis* have the capacity to colonize a diversity of tissues inside their host, including hemolymph, gills and digestive connective tissues (23). Hence *V.*

aestuarianus francensis is adapted to diverse tissular micro-environments in line with a lower

degree of specialization than highly specialized obligate intracellular bacteria. While being

strongly associated with oysters, OCI are probably not intracellular pathogens. Indeed,

Labreuche *et al.* showed that an OCI strain was able to inhibit phagocytosis by oyster

hemocytes (30). More in depth studies on OCI microhabitats within the oyster could help to

elucidate their complete intra-host life cycle, and the functional roles of the gene repertoire.

Whether OCI_A and OCI_B occupy similar niches in infected oysters and whether they have

the potential to cooperate is also unknown. Neither epidemiological parameter estimation

(31) nor in-depth analysis of their genomes has permitted the identification of functional traits

specific to each lineages yet.

We showed here that the two oyster pathogen lineages have almost clonally expanded in

Europe, after a probable recent selective bottleneck. A key event accompanying bottleneck

was the acquisition of copper resistance traits, which may have contributed to the

emergence of the two oyster-pathogenic lineages. Similar to what is observed in OCI

lineages, some pathogenic monomorphic lineages are thought to have endured population

bottlenecks associated with the acquisition of new functions (32). Such new functions could

allow a bacterial lineage to colonize a new host (32). Here we highlighted that OCI strains

have acquired a particular genomic island containing genes involved in copper detoxification.

Acquisition of this genomic island is likely to have allowed a better adaptation to life in

oysters. Indeed, copper is a heavy metal found at high levels in coastal waters because of

agricultural and industrial activities (33-35), and copper can accumulate at very high

concentrations in oyster tissues (up to 18,000 µg/g in China (36)). In oyster tissues, copper

can accumulate at sites of infection in oysters (37), where it plays a major role in the

antimicrobial defense of the infected host as copper can kill invading microorganisms (37-

39). Bacteria have developed copper detoxification mechanisms which utilize the *cop* and

cus genes (40). *Cop* genes encode for primary mediators allowing initial Cu^+ translocation

from the cytoplasm to the periplasm. *Cus* genes encode and regulate for a CusABC system

that transports periplasmic copper across the cell membrane toward the extracellular

medium (41). Importantly, *copA* is essential for copper resistance, oyster colonization and

virulence in the *V. tasmaniensis* LGP32, an oyster-pathogenic strain associated to the

Pacific Oyster Mortality Syndrome (42). Similarly, in another oyster pathogen, *V.*

crassostreae, the deletion of the *copA* also results in virulence attenuation (14). Together

with those previous observations, our data support copper resistance as a common trait

shared by different oyster bacterial pathogens whose acquisition may be required for

adaptation to the oyster as a niche.

Here we report the clonal expansion of *V. aestuarianus francensis* OCI_A and OCI_B across

Europe. One scenario which has been put forward to explain the increase of *V. aestuarianus*

associated mortalities over the last years implies the role of another major pathogen

affecting *C. gigas*, the Ostreid herpes virus OsHV-1. This virus has been regularly detected

during spat mortality events (43), and has been associated with higher mortality rate in all

growing areas of *C. gigas* along the French coasts (44), since the emergence of a new

variant in 2008. Recently, dual infections by OsHV-1 and *V. aestuarianus* were reported in

French oyster farms in 2014 and 2015 (45, 46) and have also been observed in Ireland since

2008 (Cheslett, pers comm). Even if experimental infections combining both pathogens at

high doses in spat and juvenile *C. gigas* did not demonstrate a direct collaboration between

them (47, 48), it can be hypothesized at a population level that the virus weakens animal

defenses thereby contributing to amplify *V. aestuarianus* local concentrations and initiate

epizootic events. Overall, the higher occurrence of *V. aestuarianus* since 2012 in Europe

could represent collateral damage resulting from the strong impact of the virus on French

and European oyster populations reported since 2008.

When the OCI pathogenic lineages emerged and how they have spread across Europe

remains to be elucidated. Based on genome sequencing data, phylodynamics makes it

possible to date outbreaks and investigate hypotheses of pathogen introduction and spread

(49). However, genomic data are still missing in order to date with confidence the emergence

of the 2 oyster-pathogenic lineages. Still, the clonality of the lineages that have emerged

concomitantly in Ireland and France suggests a possible transmission route from France to

Ireland, which is consistent with the fact that the Irish oyster industry relies massively on

French spat production. Global dissemination of pathogens is a recurring problem in

agriculture as in aquaculture and is often favored by anthropogenic introductions through

infected hosts (50). Preventing pathogen spread during transfers is particularly complicated

in the case of *V. aestuarianus* since animals can be asymptomatic (23). This is of particular

concern for the oyster industry which traditionally involves multiple transfers during the

lifetime of an oyster. Transfers of infected animals likely explain most of the spread of

pathogenic *V. aestuarianus* lineages across Europe. The lack of traceability in oyster farming

(52) is a major factor limiting the identification of epizootic sources, routes of dispersal and

application of control measures in many European countries (53). Therefore, beyond

regulating tranfers, there is an important need to develop very sensitive detection assays to

monitor animal sanitary status. We believe that some OCI specific genes revealed in this

study, like the *cusS/R* genomic island specific to pathogenic *V. aestuarianus* could be good

candidates to develop new early diagnostic tools.

Material & methods

Isolate selection

The *V. aestuarianus* isolates were collected during oyster mortality events in Europe

between 2001 and 2020 and independently of oyster mortality events between 1983 (type

strain isolated in the USA) and 2021. Among the isolates obtained independently of oyster

mortality episodes, a portion were isolated during cockle mortality episodes, which occurred

in France in 2012, 2015, 2018 and 2019.

Isolates were collected with the help of European NRLs responsible for the diagnosis and

surveillance of mollusk diseases: Ifremer (France), MI (Ireland), IZSVE (Italy), IRTA (Spain),

Marine Scotland Science (Scotland), and NRLFMC (Bulgaria). The 54 isolates used in the

study are summarized in the Supplementary table 10.

Illumina and PacBio whole-genome sequencing

DNA was isolated from 35 strains using the NucleoSpin Tissue Kit (Macherey-Nagel)

following the manufacturer's guidelines, and the concentration of each sample was adjusted

to 0.2 ng/µl for library preparation using the Illumina Nextera XT DNA Library Preparation Kit

according to the manufacturer's guidelines. For each library, paired-end reads of 150 bp

were produced on the Illumina NextSeq 550 platform (Bio Environnement, Perpignan,

France). The quality of the resultant fastq files was assessed using FastQC (54). Primers,

Illumina adapter sequences, low-quality bases at the 3′ end of reads with Phred-scaled

quality scores less than 30 and reads with a length <50 bp were removed using Trim Galore

(55). Quality-controlled reads were *de novo* assembled using the SPAdes genome

assembler v3.13.1 (56) with the default kmer size and a coverage cutoff value of 20 reads

(57).

DNA was isolated from other strains using a MagAttract HMW DNA Kit (Qiagen) following

the manufacturer's instructions. Library preparation and sequencing on a PacBio Sequel II

instrument were performed with the GENTYANE INRAe platform (Clermont-Ferrand,

France). *De novo* genome assemblies were obtained using FLYE v2.8.3 (58). One round of

polishing was performed using RACON v 1.4.3 (59).

The assembly was evaluated with QUAST, and genome completeness was checked using

the BUSCO v5.1.1 (60) and vibrionales_odb10 databases as lineage datasets, including 57

Vibrionales as detailed in Supplementary table 11.

Assembly features and GenBank accession numbers are summarized in Supplementary

table 1.

Genome annotations

For each assembly, coding sequences (CDSs) were predicted using PROKKA v1.12 (61) with default databases and –compliant and –rfam flags and annotation of rRNAs, tRNAs and other ncRNAs enabled. Functional annotation of the detected CDSs was performed using EggNog-Mapper v2.1.0 (62, 63). ISs were detected in the genomes using ISEScan v1.7 (64). Pseudogene detection was performed with the GenBank files released from PROKKA with PSEUDOFINDER v1.0 (65) , using the diamond flag and NCBI nonredundant database. These later analyses were performed only on resolved genomes to avoid obtaining false negatives in IS detection and false positives in pseudogene detection at the intercontig locations.

Phylogeny and recombination analysis

Pairwise ANIs (%) were calculated using FastANI (66). Whole-genome alignments of the 54 isolate contigs were made using Progressive Mauve v13/02/2015 (67). Whole-genome alignment was used to infer the maximum likelihood phylogeny using RAxML v8.2.12 with the GTRGAMMA model of nucleotide substitution and 100 bootstrap replicates, with strain 03_008T as the outgroup. Polymorphic sites along the alignment were detected using SNP sites (68).

To assess the influence of various structuring factors on the observed genetic diversity, we performed AMOVA and principal component analysis on the sequence dataset, with environmental origin, isolation context, sampling year, and country as structuring factors, using the R packages *pegas, factomineR* and *factoExtra*.

Recombination was evaluated across this genome alignment using ClonalFrameML v1.12 (69). We used an extension of the standard ClonalFrame model that allows different recombination parameters to be inferred on different branches of the clonal genealogy. 03_008T was used as a reference.

Selection analysis

Data were subjected to a neutrality test using Tajima's D (70) with the command tajima.test

in the R package *pegas* v1.1 (71). Tajima's D statistic measures the difference between two

statistics describing the diversity within the alignment (using π based on the average number

of pairwise differences among sequences and θ based on the number of segregating sites),

standardized by the variance of this difference. Under the neutral evolutionary model, which

assumes that polymorphisms segregate at mutation–drift equilibrium, Tajima's D is expected

to be zero. Positive Tajima's D values indicate that there is an excess of polymorphisms at

intermediate frequencies and is typically associated with diversifying selection. Conversely,

negative Tajima's D values can be the result of purifying selection or a selective sweep

(where a beneficial allele has recently reached fixation due to strong positive selection), both

of which result in an excess of weakly divergent alleles. However, both positive and negative

Tajima's D values can also result from demographic and epidemiological processes.

Bacterial virulence estimation by injection

Bacteria were grown with constant agitation at 22°C for 24 h in Zobell medium. Cultures

were washed by centrifugation (10 min, 1500 g) and resuspended in artificial, sterile

seawater to obtain OD_{600nm} = 0.1. Fifty microliters of bacterial suspensions (10^7 cfu, checked

by plating on Zobell agar) were injected intramuscularly into anesthetized specific pathogen-

free (SPF) oysters produced and maintained in the IFREMER facility (72). After injection, the

oysters were transferred to aquaria (10 oysters per 2.5 L aquarium, duplicate aquaria per

bacterial strain) containing 2 L of aerated, UV-treated seawater at 22°C and kept under static

conditions for 5 days.

Copper resistance phenotyping

Paper discs (6 mm) were loaded with $CuSO_4$ by soaking them overnight into a 50 mM $CuSO_4$

solution. Cells from stationary phase cultures at DO = 1 were washed twice with SSW before

462   being spread onto agar plates containing 15 g.L^{-1} bactopeptone and 0.5 M NaCl. Using

463   sterile tweezers, copper-impregnated disks were placed on the bacterial lawn before

464   incubation at 20°C. After 5 days of incubation, the diameter of the inhibition zone around the

465   disk was measured.

Bibliographic references

1.   Le Roux F, Wegner KM, Baker-Austin C, Vezzulli L, Osorio CR, Amaro C, et al. The emergence of Vibrio pathogens in Europe: ecology, evolution, and pathogenesis (Paris, 11–12th March 2015). Frontiers in microbiology. 2015;6:830.

2.   Frans I, Michiels CW, Bossier P, Willems KA, Lievens B, Rediers H. *Vibrio anguillarum* as a fish pathogen: virulence factors, diagnosis and prevention. Journal of Fish Diseases. 2011;34(9):643-61.

3.   Hewson I, Button JB, Gudenkauf BM, Miner B, Newton AL, Gaydos JK, et al. Densovirus associated with sea-star wasting disease and mass mortality. Proc Natl Acad Sci U S A. 2014;111(48):17278-83.

4.   Lafferty KD, Kuris AM. Mass mortality of abalone Haliotis cracherodii on the California Channel Islands: tests of epidemiological hypotheses. Marine Ecology-Progress Series. 1993;96:239-.

5.   Lafferty KD, Harvell CD, Conrad JM, Friedman CS, Kent ML, Kuris AM, et al. Infectious Diseases Affect Marine Fisheries and Aquaculture Economics. Annual Review of Marine Science. 2015;7(1):471-96.

6.   Murray AG, Peeler EJ. A framework for understanding the potential for emerging diseases in aquaculture. Preventive Veterinary Medicine. 2005;67(2):223-35.

7.   Engering A, Hogerwerf L, Slingenbergh J. Pathogen-host-environment interplay and disease emergence. Emerg Microbes Infect. 2013;2(2):e5-e.

8.   Martínez JL. Bacterial pathogens: from natural ecosystems to human hosts. Environmental Microbiology. 2013;15(2):325-33.

9.   Bruto M, Labreuche Y, James A, Piel D, Chenivesse S, Petton B, et al. Ancestral gene acquisition as the key to virulence potential in environmental Vibrio populations. The ISME Journal. 2018;12(12):2954-66.

10.   Sheppard SK, Guttman DS, Fitzgerald JR. Population genomics of bacterial host adaptation. Nature Reviews Genetics. 2018;19(9):549-65.

11.   Tibayrenc M, Ayala FJ. Reproductive clonality of pathogens: A perspective on pathogenic viruses, bacteria, fungi, and parasitic protozoa. Proceedings of the National Academy of Sciences. 2012;109(48):E3305-E13.

12.   Kumar A, Das B, Kumar N. Vibrio Pathogenicity Island-1: The Master Determinant of Cholera Pathogenesis. Frontiers in cellular and infection microbiology. 2020;10:561296.

13.   Smith NH, Gordon SV, de la Rua-Domenech R, Clifton-Hadley RS, Hewinson RG. Bottlenecks and broomsticks: the molecular evolution of *Mycobacterium bovis*. Nature Reviews Microbiology. 2006;4(9):670-81.

14.   Bruto M, James A, Petton B, Labreuche Y, Chenivesse S, Alunno-Bruscia M, et al. *Vibrio crassostreae,* a benign oyster colonizer turned into a pathogen after plasmid acquisition. The ISME Journal. 2017;11(4):1043-52.

15.   Lemire A, Goudenège D, Versigny T, Petton B, Calteau A, Labreuche Y, et al. Populations, not clones, are the unit of vibrio pathogenesis in naturally infected oysters. The ISME Journal. 2015;9(7):1523-31.

16.   Petton B, Bruto M, James A, Labreuche Y, Alunno-Bruscia M, Le Roux F. *Crassostrea gigas* mortality in France: the usual suspect, a herpes virus, may not be the killer in this polymicrobial opportunistic disease. Frontiers in microbiology. 2015;6:686.

17.   Garnier M, Labreuche Y, Nicolas JL. Molecular and phenotypic characterization of
Vibrio aestuarianus subsp. *francensis* subsp. nov., a pathogen of the oyster *Crassostrea*
gigas. Syst Appl Microbiol. 2008;31(5):358-65.
18.   Dégremont L, Azéma P, Maurouard E, Travers M-A. Enhancing resistance to *Vibrio*
aestuarianus in *Crassostrea gigas* by selection. Aquaculture. 2020;526:735429.
19.   Lasa A, di Cesare A, Tassistro G, Borello A, Gualdi S, Furones D, et al. Dynamics of
the Pacific oyster pathobiota during mortality episodes in Europe assessed by 16S rRNA
gene profiling and a new target enrichment next-generation sequencing strategy. Environ
Microbiol. 2019;21(12):4548-62.
20.   Mandas D, Salati F, Polinas M, Sanna MA, Zobba R, Burrai GP, et al.
Histopathological and Molecular Study of Pacific Oyster Tissues Provides Insights into *V.*
aestuarianus Infection Related to Oyster Mortality. Pathogens (Basel, Switzerland).
2020;9(6).
21.   Garcia C, Mesnil A, Tourbiez D, Moussa M, Dubreuil C, Gonçalves de Sa A, et al.
Vibrio aestuarianus subsp. *cardii* subsp. nov., pathogenic to the edible cockles
Cerastoderma edule in France, and establishment of *Vibrio aestuarianus* subsp.
aestuarianus subsp. nov. and *Vibrio aestuarianus* subsp. *francensis* subsp. nov. Int J Syst
Evol Microbiol. 2021;71(2):004654.
22.   Goudenège D, Travers MA, Lemire A, Petton B, Haffner P, Labreuche Y, et al. A
single regulatory gene is sufficient to alter *Vibrio aestuarianus* pathogenicity in oysters.
Environ Microbiol. 2015;17(11):4189-99.
23.   Parizadeh L, Tourbiez D, Garcia C, Haffner P, Dégremont L, Le Roux F, et al.
Ecologically realistic model of infection for exploring the host damage caused by *Vibrio*
aestuarianus. Environmental Microbiology. 2018;20(12):4343-55.
24.   Saulnier D, De Decker S, Haffner P. Real-time PCR assay for rapid detection and
quantification of Vibrio aestuarianus in oyster and seawater: A useful tool for epidemiologic
studies. Journal of Microbiological Methods. 2009;77(2):191-7.
25.   Moran NA. Microbial minimalism: genome reduction in bacterial pathogens. Cell.
2002;108(5):583-6.
26.   Floriano AM, Castelli M, Krenek S, Berendonk TU, Bazzocchi C, Petroni G, et al. The
Genome Sequence of "Candidatus *Fokinia solitaria*": Insights on Reductive Evolution in
Rickettsiales. Genome biology and evolution. 2018;10(4):1120-6.
27.   Bäumler A, Fang FC. Host Specificity of Bacterial Pathogens. Cold Spring Harbor
Perspectives in Medicine. 2013;3(12).
28.   Siguier P, Gourbeyre E, Chandler M. Bacterial insertion sequences: their genomic
impact and diversity. FEMS Microbiology Reviews. 2014;38(5):865-91.
29.   Touchon M, Rocha EPC. Causes of Insertion Sequences Abundance in Prokaryotic
Genomes. Molecular Biology and Evolution. 2007;24(4):969-81.
30.   Labreuche Y, Le Roux F, Henry J, Zatylny C, Huvet A, Lambert C, et al. *Vibrio*
aestuarianus zinc metalloprotease causes lethality in the Pacific oyster *Crassostrea gigas*
and impairs the host cellular immune defenses. Fish & shellfish immunology.
2010;29(5):753-8.
31.   Travers MA, Tourbiez D, Parizadeh L, Haffner P, Kozic-Djellouli A, Aboubaker M, et
al. Several strains, one disease: experimental investigation of *Vibrio aestuarianus* infection
parameters in the Pacific oyster, *Crassostrea gigas*. Vet Res. 2017;48(1):32.
32.   Namouchi A, Didelot X, Schöck U, Gicquel B, Rocha EP. After the bottleneck:
Genome-wide diversification of the Mycobacterium tuberculosis complex by mutation,
recombination, and natural selection. Genome Res. 2012;22(4):721-34.

33.   Claisse D, Alzieu C. Copper contamination as a result of antifouling paint
regulations? Marine Pollution Bulletin. 1993;26(7):395-7.
34.   Hall LW, Anderson RD. A Deterministic Ecological Risk Assessment for Copper in
European Saltwater Environments. Marine Pollution Bulletin. 1999;38(3):207-18.
35.   Rossi N, Jamet J-L. *In situ* heavy metals (copper, lead and cadmium) in different
plankton compartments and suspended particulate matter in two coupled Mediterranean
coastal ecosystems (Toulon Bay, France). Marine Pollution Bulletin. 2008;56(11):1862-70.
36.   Weng N, Wang W-X. Variations of trace metals in two estuarine environments with
contrasting pollution histories. Science of The Total Environment. 2014;485-486:604-14.
37.   Shi B, Wang T, Zeng Z, Zhou L, You W, Ke C. The role of copper and zinc
accumulation in defense against bacterial pathogen in the fujian oyster (*Crassostrea
angulata*). Fish & shellfish immunology. 2019;92:72-82.
38.   Destoumieux-Garzón D, Canesi L, Oyanedel D, Travers M-A, Charrière GM, Pruzzo
C, et al. Vibrio–bivalve interactions in health and disease. Environmental Microbiology.
2020;22(10):4323-41.
39.   Yeh S-T, Liu C-H, Chen J-C. Effect of copper sulfate on the immune response and
susceptibility to *Vibrio alginolyticus* in the white shrimp *Litopenaeus vannamei*. Fish &
shellfish immunology. 2004;17(5):437-46.
40.   Bondarczuk K, Piotrowska-Seget Z. Molecular basis of active copper resistance
mechanisms in Gram-negative bacteria. Cell biology and toxicology. 2013;29(6):397-405.
41.   Argüello JM, Raimunda D, Padilla-Benavides T. Mechanisms of copper homeostasis
in bacteria. Frontiers in cellular and infection microbiology. 2013;3:73.
42.   Vanhove AS, Rubio TP, Nguyen AN, Lemire A, Roche D, Nicod J, et al. Copper
homeostasis at the host vibrio interface: lessons from intracellular vibrio transcriptomics.
Environ Microbiol. 2016;18(3):875-88.
43.   Garcia C, Thébault A, Dégremont L, Arzul I, Miossec L, Robert M, et al. Ostreid
herpesvirus 1 detection and relationship with *Crassostrea gigas* spat mortality in France
between 1998 and 2006. Veterinary Research. 2011;42(1):73.
44.   Barbosa Solomieu V, Renault T, Travers MA. Mass mortality in bivalves and the
intricate case of the Pacific oyster, *Crassostrea gigas*. J Invertebr Pathol. 2015;131:2-10.
45.   Azéma P, Maurouard E, Lamy J-B, Dégremont L. The use of size and growing height
to improve *Crassostrea gigas* farming and breeding techniques against OsHV-1.
Aquaculture. 2017;471:121-9.
46.   Azema P, Travers MA, De Lorgeril J, Tourbiez D, Degremont L. Can selection for
resistance to OsHV-1 infection modify susceptibility to *Vibrio aestuarianus* infection in
Crassostrea gigas? First insights from experimental challenges using primary and
successive exposures. Vet Res. 2015;46:139.
47.   Azema P, Lamy JB, Boudry P, Renault T, Travers MA, Degremont L. Genetic
parameters of resistance to *Vibrio aestuarianus,* and OsHV-1 infections in the Pacific oyster,
Crassostrea gigas, at three different life stages. Genet Sel Evol. 2017;49(1):23.
48.   Azéma P, Travers M-A, Benabdelmouna A, Dégremont L. Single or dual
experimental infections with *Vibrio aestuarianus* and OsHV-1 in diploid and triploid
Crassostrea gigas at the spat, juvenile and adult stages. Journal of Invertebrate Pathology.
2016;139:92-101.
49.   Destoumieux-Garzón D, Matthies-Wiesler F, Bierne N, Binot A, Boissier J, Devouge
A, et al. Getting out of crises: Environmental, social-ecological and evolutionary research is
needed to avoid future risks of pandemics. Environment international. 2022;158:106915.

50.   Anderson PK, Cunningham AA, Patel NG, Morales FJ, Epstein PR, Daszak P.
Emerging infectious diseases of plants: pathogen pollution, climate change and
agrotechnology drivers. Trends in Ecology & Evolution. 2004;19(10):535-44.
51.   Buestel D, Ropert M, Prou J, Goulletquer P. History, status, and future of oyster
culture in France. Journal of Shellfish Research. 2009;28(4):813-20.
52.   Håstein T, Hill BJ, Berthe F, Lightner DV. Traceability of aquatic animals. Revue
scientifique et technique (International Office of Epizootics). 2001;20(2):564-83.
53.   Pernet F, Lupo C, Bacher C, Whittington RJ. Infectious diseases in oyster
aquaculture require a new integrated approach. Philosophical Transactions of the Royal
Society B: Biological Sciences. 2016;371(1689):20150213.
54.   Wingett SW, Andrews S. FastQ Screen: A tool for multi-genome mapping and quality
control. F1000Research. 2018;7:1338.
55.   Bush SJ. Read trimming has minimal effect on bacterial SNP-calling accuracy.
Microbial Genomics. 2020;6(12).
56.   Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, et al. SPAdes:
a new genome assembly algorithm and its applications to single-cell sequencing. J Comput
Biol. 2012;19(5):455-77.
57.   Gurevich A, Saveliev V, Vyahhi N, Tesler G. QUAST: quality assessment tool for
genome assemblies. Bioinformatics (Oxford, England). 2013;29(8):1072-5.
58.   Kolmogorov M, Yuan J, Lin Y, Pevzner PA. Assembly of long, error-prone reads
using repeat graphs. Nature Biotechnology. 2019;37(5):540-6.
59.   Vaser R, Sovic I, Nagarajan N, Sikic M. Fast and accurate de novo genome
assembly from long uncorrected reads. Genome Research. 2017.
60.   Simão FA, Waterhouse RM, Ioannidis P, Kriventseva EV, Zdobnov EM. BUSCO:
assessing genome assembly and annotation completeness with single-copy orthologs.
Bioinformatics (Oxford, England). 2015;31(19):3210-2.
61.   Seemann T. Prokka: rapid prokaryotic genome annotation. Bioinformatics (Oxford,
England). 2014;30(14):2068-9.
62.   Cantalapiedra CP, Hernández-Plaza A, Letunic I, Bork P, Huerta-Cepas J. eggNOG-
mapper v2: Functional Annotation, Orthology Assignments, and Domain Prediction at the
Metagenomic Scale. Molecular Biology and Evolution. 2021;38(12):5825-9.
63.   Huerta-Cepas J, Szklarczyk D, Heller D, Hernández-Plaza A, Forslund SK, Cook H,
et al. eggNOG 5.0: a hierarchical, functionally and phylogenetically annotated orthology
resource based on 5090 organisms and 2502 viruses. Nucleic Acids Research.
2018;47(D1):D309-D14.
64.   Xie Z, Tang H. ISEScan: automated identification of insertion sequence elements in
prokaryotic genomes. Bioinformatics (Oxford, England). 2017;33(21):3340-7.
65.   Syberg-Olsen MJ, Garber AI, Keeling PJ, McCutcheon JP, Husnik F. Pseudofinder:
detection of pseudogenes in prokaryotic genomes. bioRxiv. 2021:2021.10.07.463580.
66.   Goris J, Konstantinidis KT, Klappenbach JA, Coenye T, Vandamme P, Tiedje JM.
DNA-DNA hybridization values and their relationship to whole-genome sequence similarities.
Int J Syst Evol Microbiol. 2007;57(Pt 1):81-91.
67.   Darling AE, Mau B, Perna NT. progressiveMauve: Multiple Genome Alignment with
Gene Gain, Loss and Rearrangement. PLOS ONE. 2010;5(6):e11147.
68.   Page AJ, Taylor B, Delaney AJ, Soares J, Seemann T, Keane JA, et al. SNP-sites:
rapid efficient extraction of SNPs from multi-FASTA alignments. Microbial Genomics.
2016;2(4).

69.   Didelot X, Wilson DJ. ClonalFrameML: Efficient Inference of Recombination in Whole
Bacterial Genomes. PLOS Computational Biology. 2015;11(2):e1004041.
70.   Tajima F. Statistical method for testing the neutral mutation hypothesis by DNA
polymorphism. Genetics. 1989;123(3):585-95.
71.   Paradis E. pegas: an R package for population genetics with an integrated–modular
approach. Bioinformatics (Oxford, England). 2010;26(3):419-20.
72.   Petton B, Pernet F, Robert R, Boudry P. Temperature influence on pathogen
transmission and subsequent mortalities in juvenile Pacific oysters *Crassostrea gigas*.
Aquaculture Environment Interactions. 2013;3(3):257-73.
663

Table 1: **Analysis of molecular variance** supported genetic partitioning according to

environmental origin and isolation context

Factor	Variability (%)	p-value
Environmental origin	**17.8**	**0.002**
Isolation context	**16.0**	**0.001**
Country of isolation	7.1	0.074
Year of isolation	3.4	0.25

Figure 1: **European expansion of *V. aestuarianus francensis***. While the geographic

distribution of *V. aestuarianus* strains involved in *C. gigas* mortalities expands, genetic

diversity remains stable over time. A: The number of countries reporting the detection of *V.*

aestuarianus was assessed through a questionnaire sent to European national reference

laboratories (NRLs). *Note that the Denmark case report corresponds to a suspected case

without bacterial isolation. B: The two lineage expansions: 2 genotypes were determined

based on a maximum likelihood phylogenetic tree from complete genome alignments (GTR

substitution model).

Figure 2: *V. aestuarianus* **diversity is structured according to host species and**

isolation context. Phylogenetic tree of *V. aestuarianus* species calculated with RAxML

using a GTR substitution model and rooted with the 03_008T strain (outgroup) from whole-

genome alignments. Several layers of annotations are added: country (check mark), matrix

of isolation (star), year of isolation (point), virulence testing in the laboratory by injection into

oysters (10^7 UFC/animal) indicated with a bar plot on the right side representing mortality

percentages, and context (strains isolated during oyster mortality events are indicated with a

red arrow on the right). Oyster clinical isolate (OCI) clades A and B are highlighted in blue

and green, respectively. The absence of virulence data means the absence of a test for this

strain.

Figure 3: **Principal component analysis (PCA) of genetic polymorphism segregates _V._**

aestuarianus isolates according to environmental origin and isolation context,

according to genome-wide analysis of 54 genomes using 183 109 SNPs. PC1 and PC2

represent the first two eigenvectors from the PCA. Isolates are colored according to the

isolation context (A) and according to the environmental origin (B). PC1 and PC2 explain

16.7% and 9.2% of the variability between samples, respectively.

Figure 4: **A limited number of recombination events occurred in oyster clinical isolate**

(OCI) genomes after their expansion. Detected **recombination** events are shown by dark

blue horizontal bars. For a given branch, light blue sites indicate no substitution. All other

colors (from white to red) indicate a substitution. White represents a nonhomoplasic

substitution. An increasing level of redness means an increasing degree of homoplasy. OCI

clades A and B are represented in blue and green, respectively, and the NO_OCI group is

shown in yellow. For improved readability, some nodes of the tree as well as the row of the

recombination events matrix related to these nodes are indicated with points and dashed

lines. The strain 03_008T sequence was used as a reference.

Figure 5: **Strong impact of purifying selection on oyster pathogens.** Genetic statistics in

V. aestuarianus species (black), NO_OCI (yellow), oyster clinical isolates (OCIs), clade A

(blue) and clade B (green). **A: Whole genome scale**. Even through the expected genetic

diversity under neutral evolution and a stable population size in OCIs is low (Watterson's

theta), the observed diversity (Pi) is still lower, as shown by the Tajima's D value below -2.

For *V. aestuarianus*, OCI_A and NO_OCI, 10 random subsamples of 8 sequences in each

alignment were examined to avoid bias due to differences in the number of sequences. For

the OCI_B alignment, the 8 sequences represent the entire sampled set. Pi: observed

genetic diversity; Theta_w: Watterson estimate of expected genetic diversity; Tajima_D:

Tajima's D test statistic. The Y axis in the Pi and Watterson's theta plots is log10 scaled.

B: Gene scale. Core genes show different selection patterns among groups of strains and

among core genes within groups. While most core genes are under neutral evolution in

isolates other than OCIs, most genes are monomorphic in OCI clades A and B. D>0 and

D<0: significantly negative or positive Tajima's D value (p < 0.05) and D=0 (p > 0.05). A

negative Tajima's D is indicative of purifying selection or recent population expansion, while

a positive Tajima's D is indicative of diversifying selection or population contraction. A zero

value is indicative of neutral evolution.

Figure 6: **Differential Insertion sequence (IS) enrichment is observed in oyster clinical**

isolate (OCI) genomes and cockle associated NO-OCI. Distribution of the number of IS

elements (complete and partial) in 16 complete genomes (chromosomes A and B). Colors

represent IS families predicted by ISEScan. The genomes are those of 4 OCI clade A

strains, 3 OCI clade B strains, 5 NO_OCI strains isolated independent of bivalve mortality

(NO_OCI group 1), and 4 NO_OCI strains isolated during cockle mortality events (NO_OCI

group 2). The conservation of IS number and families between genomes of OCIs from clade

A or clade B indicates an ancestral origin of ISs. The genomes that contain the most ISs are

those of strains isolated during cockle mortality events (NO_OCI group 2). The diversity of IS

quantities and families represented suggests that IS enrichment is in progress. The number

of ISs in groups was significantly different between OCI and NO_OCI from group 1 and

group 2 (**: Kruskal test p value < 0.05).

Figure 7: A. Oyster clinical isolate (OCI)-specific locus containing copper resistance genes.

Transposase and integrase are indicated in yellow, cus genes in orange, and cop genes in

blue.

B. Evolutionary history of the cus/cop locus. NJ, bootstrap test (500 replicates). The

evolutionary distances were computed using the Kimura 2-parameter method [3] and are

shown as the number of base substitutions per site. This analysis involved 8 nucleotide

sequences. All ambiguous positions were removed for each sequence pair (pairwise deletion

option). There were a total of 29 555 positions in the final dataset. Evolutionary analyses

were conducted in MEGA X [4]. C. Agar plate assay: bacterial growth inhibition diameter

around a pellet impregnated with 50 mM $CuSO_4$ solution.