**Protein evidence of unannotated ORFs in *Drosophila* reveals unappreciated diversity in the**

**evolution of young proteins**

Eric B Zheng, Li Zhao[*]

**Affiliations:** Laboratory of Evolutionary Genetics and Genomics, The Rockefeller University, New York,

NY 10065, USA

[*]Correspondence to: lzhao@rockefeller.edu

**Abstract**

*De novo* gene origination, where a previously non-genic genomic sequence becomes genic through

evolution, has been increasingly recognized as an important source of evolutionary novelty across diverse

taxa. Many *de novo* genes have been proposed to be protein-coding, and in several cases have been

experimentally shown to yield protein products. However, the systematic study of *de novo* proteins has

been hampered by doubts regarding the translation of their transcripts without the experimental

observation of protein products. Using a systematic, ORF-focused mass-spectrometry-first computational

approach, we identify almost 1000 unannotated open reading frames with evidence of translation

(utORFs) in the model organism *Drosophila melanogaster*, 371 of which have canonical start codons. To

quantify the comparative genomic similarity of these utORFs across *Drosophila* and to infer

phylostratigraphic age, we further develop a synteny-based protein similarity approach. Combining these

results with reference datasets on tissue- and life-stage-specific transcription and conservation, we

identify different properties amongst these utORFs. Contrary to expectations, the fastest-evolving utORFs

are not the youngest evolutionarily. We observed more utORFs in the brain than in the testis. Most of the

identified utORFs may be of *de novo* origin, even accounting for the possibility of false-negative

similarity detection. Finally, sequence divergence after an inferred *de novo* origin event remains

substantial, raising the possibility that *de novo* proteins turn over frequently. Our results suggest that there

is substantial unappreciated diversity in *de novo* protein evolution: many more may exist than have been

previously appreciated; there may be divergent evolutionary trajectories; and *de novo* proteins may be

gained and lost frequently. All in all, there may not exist a single characteristic model of *de novo* protein

evolution, but rather complex origins and evolutionary trajectories for *de novo* proteins.

**Impact statement**

The analysis of mass-spectrometry data for all possible open reading frames reveals protein evidence for

evolutionarily young, unannotated proteins with distinct characters.

## Introduction

While the study of the evolution of entire species or the fixation of individual mutations has matured over decades, our understanding of how genes evolve is relatively less mature. The dominant mechanism is duplication and divergence (Ohno, 1970; Long et al., 2003), but the early years of the genomic era revealed a seeming paradox of species-specific "orphan genes". These genes lacked homologs even in closely related species, which made it necessary to invoke non-standard duplication and divergence mechanisms (Domazet-Loso and Tautz, 2003). Advances in both the sequencing of genomes and in comparative genomics soon led to an explosion of work on the origins of orphan genes, generally identified through lineage-specificity.

Lineage-specific orphan genes can originate from a variety of mechanisms, ranging from extreme, rapid divergence after duplication (Chen et al., 2010; Domazet-Loso and Tautz, 2003) to horizontal gene transfer (Verster et al., 2019) to *de novo* gene birth. In the latter mechanism, previously non-genic genomic sequence acquires genic function, thus yielding a *de novo* gene that is characterized by its lineage-specificity (Begun et al., 2006; Levine et al., 2006; McLysaght and Hurst, 2016; Schlötterer, 2015; Van Oss and Carvunis, 2019). *De novo* genes are identified from diverse taxa from fruit flies (Begun et al., 2006; Levine et al., 2006), to yeast (Cai et al., 2008) and even humans (Knowles and McLysaght, 2009; Xie et al., 2012). While it would seem unlikely for such *de novo* genes to gain viable function from presumably unoptimized sequence, empirical and population genetics studies have demonstrated that they have important functions (Cai et al., 2008; Lange et al., 2021; Levine et al., 2006; Reinhardt et al., 2013; Xie et al., 2019; Zhao et al., 2014). However, exactly what proportion of lineage-specific genes are contributed through *de novo* gene birth remains actively studied (Tautz and Domazet-Lošo, 2011; McLysaght and Guerzoni, 2015; Vakirlis et al., 2020).

Despite significant interest, the study of evolutionarily young genes is surprisingly conceptually difficult. While an empirical investigation of biological effect is ideal, designing such experiments for a specific evolutionarily young gene can be challenging due to the likely lack of homology to previously studied genes. Even identifying a given candidate genetic locus as an evolutionarily young gene can be fraught with challenges: evolutionary youth is functionally equivalent to concluding lineage-specificity, but methods like phylostratigraphy via BLAST (Domazet-Loso et al., 2007), while useful and intuitive, can be vulnerable to false-negative detections, especially for short genes (Moyers and Zhang, 2015, 2016). Moreover, while researchers generally agree that a gene should be functional, exactly how this should be defined and assessed can vary substantially (Van Oss and Carvunis, 2019). As an example, whether evolutionarily young genes function as RNAs or as proteins is often an area of debate. One of the earliest recognized young genes is an RNA (Wang et al., 2002), other studies also provided such support

(Cridland et al., 2022; Ruiz-Orera et al., 2015); in contrast, in several cases even *de novo* genes have been

empirically shown to yield protein products (Bungard et al., 2017; Cai et al., 2008; Knowles and

McLysaght, 2009; Li et al., 2010; Zhang et al., 2019). While defining a gene as one that can code for a

translated protein is convenient and intuitive, evidence for the translation into polypeptides of lineage-

specific genes and in particular, *de novo* genes as a class, is often lacking (Van Oss and Carvunis, 2019).

As a result, much effort remains needed for systematic approaches to studying evolutionarily young genes

and proteins.

Proteomics approaches like shotgun mass spectrometry (MS) can be a powerful tool for the rapid

identification of translational evidence for many potential proteins at once, and would be a good fit for

such a systematic study. However, standard shotgun MS analysis approaches require *a priori* assumptions

about the sequences of the proteins to be detected (Sinitcyn, Rudolph, and Cox 2018). In the case of

discovering possible lineage-specific proteins, this *a priori* requirement can be confounding, as lineage-

specific proteins may be systematically under-annotated due to their lack of homology.

Here, we employ a systematic, ORF-focused MS-first computational approach, to identify unannotated

open reading frames with evidence of translation (utORFs) in *Drosophila melanogaster*. We further

develop a synteny-based protein similarity approach to quantify the comparative genomic similarity of

these utORFs across *Drosophila*, to estimate their phylostratigraphic age, and finally to infer possible *de*

*novo* origin for the vast majority of these utORFs. Combined with reference datasets on tissue- and life

stage-specific transcription and conservation, this data suggests differences amongst these utORFs with

respect to their evolutionary and functional properties. Finally, we independently support a subset of these

utORFs via additional MS empirical data and ribosome profiling data.

**Results**

**A comprehensive database of all possible open reading frames in the *Drosophila melanogaster***
**genome**

As a first step towards the systematic discovery of unannotated proteins in *Drosophila melanogaster*, we

generated a comprehensive database of all possible open reading frames (ORFs) in the *D. melanogaster*

genome. This comprehensive database is the result of translating the entire annotated genome in all six

frames; we also included the annotated transcriptome to enable discovery of non-canonical translation

phenomena such as alternate frame ORFs (de Klerk and 't Hoen 2015). In this database, each potential

ORF is defined as a contiguous span of amino acid-coding codons; so, these potential ORFs do not

necessarily begin with or contain a canonical start codon. This is necessary in order to account for

unannotated splicing events, as a six-frame translation of the genome cannot correctly associate the exons

of a gene. Since no coding exon within the coding sequence of a gene can contain a stop codon, the set of

all codon spans between stop codons is a superset of the set of all exons in the genome. In addition to

annotated ORFs (FlyBase r6.15 release), this comprehensive potential ORF database contains 4,582,998

unique potential ORFs that are not otherwise annotated. Of these additional potential ORFs, the median

length is 32 amino acids and the 95th percentile is 96 amino acids. Considering solely the 2,597,794

potential ORFs that contain a canonical start codon, the median length following the start codon remains

32 amino acids, and the 95th percentile is 112 amino acids. In total, this suggests that the *D. melanogaster*

genome has substantial latent potential for generating new protein-coding genes. Of course, the

overwhelming majority of these potential ORFs are neither transcribed nor translated and are thus not

functional (Durand et al. 2019), but considering these will allow us to scan the potential proteomic

diversity in *D. melanogaster*.

**Two-round MS search improves discovery of unannotated ORFs without compromising FDR**

A comprehensive proteome of *D. melanogaster* was recently published (Casas-Vila et al. 2017), including

data from 15 stages of the whole lifecycle of the fruit fly (ranging from 0-2h embryos to adults, including

larval and pupal stages), as well as sharper time-resolution data covering 14 stages of embryogenesis

(hourly timepoints from 0-6 hours; bi-hourly timepoints from 6 to 20 hours). We analyzed the *Casas-*

*Villa et al.* data by searching for the complete potential ORF database described above to discover

evidence of translation of unannotated ORFs in *D. melanogaster*. Because the number of potential ORFs

is several orders of magnitude greater than the number of annotated proteins, the search space is

enormously expanded, and thus a traditional one-round search is likely to miss a very large number of

identifications at a given false-discovery rate (FDR). Accordingly, to improve total sensitivity while

maintaining an acceptable FDR, we used two rounds of analysis (**Methods**). We identify dramatically

more novel unannotated protein groups with statistical significance in this two-round approach than in a

traditional single-round search, while the number of false positive identifications is comparable

(**Supplemental File 1A**). The substantial enrichment (around ten-fold) of unannotated ORFs relative to

either the observed number of false positives or the desired FDR strongly supports the validity of the two-

round approach.

In total, after additional filtering (see **Methods**), our two-round search identified 3123 unannotated

unique peptides that underlie 993 unannotated ORFs without homology to annotated proteins (**Figure 1A;**

**Figure 1B; Figure 1–Source Data 1**; **Figure 1–Source Data 2**), which we hereafter refer to as

unannotated translated ORFs (utORFs). utORFs range in length from 14 amino acids (aa) to over 100 aa,

with the longest being 1748 aa (**Figure 1C**). The median length of a utORF is 37 aa, and the mean is 48.6

aa. Among these utORFs, 331 proteins have canonical start codons with 1027 peptides that originate from

sequence after the canonical start codon, implying that on average each protein has 3 supporting peptide-

spectrum matches. The mean and median lengths of a utORF starting with a canonical start codon are

53.5 aa and 37 aa, respectively, which is very similar to utORFs overall.  Unless specifically mentioned,

we analyze henceforth the set of 993 utORFs and not the 331 with canonical start codons.

Compared to the annotated *D. melanogaster* proteome, utORF sequences have similar amino acid

composition (**Figure 1D**; Spearman rank-order correlation of amino acid frequencies, utORFs to

annotated proteome: 0.767, $p$ = 7.98 E -5). The amino acid composition for those utORFs with start

codons is also highly similar. Some hydrophobic and positively-charged amino acids, such as leucine,

arginine, and tryptophan are overrepresented relative to annotated proteins, while negatively-charged

amino acids, such as aspartate and glutamate are underrepresented. Overall, the distribution of positively-

charged to negatively-charged amino acids is significantly different among utORFs versus annotated *D.*

*melanogaster* protein sequences (Chi-squared test with Yates's correction, Chi-sq = 1463.55, df=1, $p <$

machine epsilon). In addition, the distribution of hydrophobic amino acids versus polar amino acids is

also highly significantly different (same test, Chi-sq = 44.96, df=1, $p <$ 1E -10). These differences may be

explained by the amino acid composition of potential ORFs, as the implied amino acid composition of all

potential ORFs is more similar to utORFs than annotated proteins (Spearman rank-order correlation of

amino acid frequencies, potential ORFs to utORFs: 0.970, $p$ = 1.71 E -12; potential ORFs to annotated

proteome: 0.785, $p$ = 4.15 E -5).

All 993 utORFs have unique genomic locations (**Figure 1–Source Data 3**). Unannotated translated ORFs

reside in a range of genomic locations, including intergenic, intronic, or UTRs (**Figure 1E**).  About one-

third of utORFs are located intergenically; of these, about one-third again are longer than 50 aa and

contain a canonical start codon. We propose that these may be single-exon unannotated proteins. A

considerable number of utORFs overlap with annotated genes, with an even distribution between sense

and anti-sense directions relative to the annotation. Of the former, several are located either upstream or

downstream of the canonical CDS; these may be instances of uORFs or translation read-through, which

have been previously described as having important biological functions (Andrews and Rothnagel 2014).

Notably, with the exception of a modest reduction in the relative proportion of sense utORFs, the

distribution of locations is not different for those utORFs with a canonical start codon. The genomic

sequences that make up utORFs on average have greater GC content compared that of the overall *D.*

*melanogaster* genome (X and autosomes; mean utORF GC content 43.4% vs. X and autosome GC

content 42.15%; 1-sample two-tailed *t*-test, $p$<1E-4), but they have wide variety, with some with as much

as 62% GC content and others as few as 21% (**Figure 1G**). Intriguingly, the GC content of annotated

genes (annotated gene GC content 42.9%) is between that of the genome and utORFs, and utORFs do not

have significantly different GC content compared to annotated genes (1-sample two-tailed *t*-test, $p >$
0.05). This suggests either that a selective pressure may act towards increasing the GC content in utORFs,
or that utORFs may be more likely to originate from regions with higher GC content.

Approximately half of utORFs (566 of 993; 57.0%) have a majority coiled-coil secondary structure, with
the remainder a mixture of beta sheets and alpha helices. Notably, this remains true even when
normalizing for length, as the weighted proportion of sequence in a predicted coiled-coil structure is on
average about half (**Figure 1F**). Predictions of structural disorder of utORFs suggest that while they are
rather disordered, most retain a substantial proportion that is ordered. The median proportion of
disordered utORFs is 24.5%.

**Novel peptides identified are not artifacts caused by polymorphism in conserved genes**

It is possible that polymorphisms within a population can yield unique peptides due to amino acid
substitutions, thus resulting in inaccurate attribution of such peptides to utORFs (Faridi, Purcell, and Croft
2018). However, such polymorphisms can contribute at most a small fraction of the peptides that support
the identified utORFs. To quantify this, using a trie, we calculated the minimum distance between the
peptides that uniquely support utORFs and the expected set of peptides from the reference proteome (see
**Methods**). Of the 3123 tryptic peptides that are uniquely identified from utORFs, only 1 could possibly
result from a single amino acid substitution of a tryptic peptide from an annotated protein; none could
result from a single amino acid insertion or deletion. A simultaneous double amino acid substitution could
explain an additional 5 peptides; 24 more peptides could result from a double insertion or deletion. Thus,
single or double amino acid substitutions can explain at most 30 of 3123 unique tryptic peptides (< 1%).
As a result, it is exceedingly likely that the peptides detected here are the product of utORFs rather than
polymorphisms at annotated genes.

**Inferring ages of unannotated translated ORFs**

In order to compare the conservation of utORFs across species, it is necessary to find the orthologous
sequence with near base-pair-level resolution. Due to the algorithmic compromises necessary for whole-
genome searches, BLAST homology searches do not guarantee that they will return optimum results,
especially for short sequences that may be highly diverged. Moreover, any homology search that solely
relies on searching for a given query sequence necessarily neglects broader genomic context like synteny.
Conversely, common genomic utilities like UCSC's *liftOver* are explicitly not recommended for fine-
detailed genomic coordinate conversions. To address these issues, we used a pre-existing multiple-
sequence-alignment (MSA) to guide a maximally-optimal local genome alignment (**Figure 2 AB**). This

allows the combination of both whole genomic contexts as well as the algorithmic guarantees of the

Smith-Waterman alignment.

To estimate a null model of protein similarity scores, we calculated control distributions with varying

assumptions. Assuming amino acids drawn from the distribution of amino acids in the annotated *D.*

*melanogaster* proteome, 10000 simulated pairs of randomly-generated protein sequences have a mean

expected similarity score ranging from 1.7 to 2.0 (**Figure 2C**). We chose a conservative threshold of 2.5

points as a threshold for determining significant sequence similarity, as it is approximately at least 2

standard deviations above the mean expectation.

We used the above-described protein similarity score to identify significant orthologs across related

*Drosophila* and insect species using the UCSC 27-way MSA (**Figure 3A**). We inferred gene ages for

each utORF, where the inferred origin of a utORF corresponded to the last common ancestor between

*Drosophila melanogaster* and the most distantly related species where a significant ortholog was

identified. We find that utORFs have a wide range of inferred gene ages, ranging from utORFs specific

just to *D. melanogaster* to utORFs that may be conserved broadly through ancient insect taxa (**Figure**

**3B**). The plurality of utORFs likely predate the last common ancestor of *Drosophila* with the most-distant

significant ortholog in *D. grimshawi* or any of its sister species; the next largest group predate the last

common ancestor of the *Drosophila* species group, with the most-distant significant ortholog in *D.*

*ficusphila* or its sister species. Notably, gene age inferences are not dramatically affected by choice of

significance threshold (**Figure 3–figure supplement 1**).

**Latent class analysis reveals subpopulations of unannotated translated ORFs**

Latent class analysis (LCA) is a statistical method that supposes that the population under study is

comprised of a mixture of distinct subpopulations – the "latent" classes (Collins and Lanza 2009).

Importantly, this is the only major assumption necessary, so the analysis is robust to the shape of

distributions or correlations between variables. Moreover, LCA can be used both deductively and

inductively; its primary output is a table of probabilities for categorical variables conditional on class

membership. Thus, these conditional probabilities can be used deductively to describe differences in the

inferred classes, and they can be used inductively to predict class membership given an individual set of

observations.

Due to the unbiased strategy of identification, it is likely that the unannotated translated ORFs identified

here consist of a mixture of loci resulting from different, distinct evolutionary processes, ranging from

intergenic *de novo* genes to alternative open reading frames of conserved annotated genes (Samandi et al.

2017). In addition, certain key variables, like inferred gene age and genomic location, are inherently

categorical. Rather than simple inter-variable correlations, LCA is a valuable lens for understanding the
unannotated loci identified here.

Accordingly, latent class analysis suggests that the population of high-confidence unannotated translated
ORFs can be subdivided into five classes (Methods; **Table 1; Figure 4-Source Data 1**) and
interpretations assigned based off differences in the various factor probabilities conditional on class
membership (Collins and Lanza 2009).

Class 1 loci are notable for minimal transcription (**Figure 4C**) as well as a marked bias of being
intergenic or antisense (**Figure 4A**). They also are almost exclusively of intermediate length between 20
and 50 residues (**Figure 4B**). This supports an interpretation as putatively nonfunctional loci. Class 2 loci
are remarkable for monophyly (**Figure 4C**), likely due to very recent inferred emergence (**Figure 4D**),
suggesting that these may be *melanogaster*-specific loci. Dramatic differences in overall conservation
(**Figure 4E**) as well as subtler ones in inferred emergence times (**Figure 4D**) distinguish classes 3, 4, and
5. Class 5 is also remarkable for strong bias towards locations overlapping existing, annotated genes, as
well as high conservation, supporting an interpretation as ORFs in alternative reading frames. Finally,
class 3 can be distinguished from class 4 through a more recent inferred age of emergence approximately
at the last common ancestor of the *melanogaster* subgroup (**Figure 4D**) and through remarkably low
conservation that is even more frequent than that of *melanogaster*-specific ORFs. This supports a
distinction between class 3 as fast-evolving ORFs and class 4 as general unannotated ORFs.

With inferred classes in hand, it is possible to use the classes for forward inferences by assigning each
utORF to a class and comparing the classes. Importantly, while binning continuous variables is necessary
for conducting latent class analysis, comparisons between classes can be done on the original continuous
variables, removing that potential source of bias, and additional variables not originally used for assigning
classes can be used as well. The posterior probabilities of class membership for each utORF suggest
generally clean partitioning between classes, as each utORF generally has a high posterior probability of
belonging to only a single class (**Figure 4–figure supplement 1**). The exceptions are utORFs with
membership in either class 4 (unannotated ORFs) or class 5 (alternative-frame ORFs). However, allowing
an additional sixth class leads to overfitting (see **Methods**). Accordingly, we use the maximum posterior
probability per utORF to assign class membership for further analysis.

For certain parameters, differences by inferred class membership recapitulate expected trends. For
example, phastCons conservation scores vary by class, with the lowest conservation in fast-evolving and
*melanogaster*-specific utORFs (**Figure 5A**). Alternative-frame utORFs are very well conserved due to the
additional constraints imposed by their overlapping annotated genes. Despite not being included as a

variable in LCA, phyloP conservation scores also vary significantly by class (**Figure 5B**). Negative

phyloP scores are consistent with faster evolution; 61.4% (27 of 44) are from fast-evolving utORFs and

24.1% (15 of 44) are from *melanogaster*-specific utORFs. Importantly. while phastCons and phyloP

scores are significantly correlated, only phyloP scores can differentiate between relative low conservation

and accelerated evolution. This recapitulation of faster evolution supports the overall latent class analysis

approach. Finally, consistent with the assigned interpretation, *melanogaster*-specific and fast-evolving

utORFs are the youngest according to phylostratigraphy (**Figure 5C**). Thus, the fastest-evolving utORFs,

which are the fast-evolving utORFs with negative phyloP scores, are surprisingly not the youngest

utORFs, as the distinct *melanogaster*-specific utORFs are younger.

For other parameters, differences between inferred classes reveals surprising trends. Fast-evolving

utORFs and putatively nonfunctional utORFs are significantly shorter than all others (one-sided Brunner-

Munzel nonparametric test of lengths vs. all other utORFs, $p << 0.001$ for both). Interestingly,

*melanogaster*-specific utORFs are not significantly shorter (same test, $p=0.11$), despite prior reports that

younger genes tend to be shorter. With regards to maximum observed transcription, alternative-frame

utORFs tend to be significantly better-transcribed than all other utORFs (one-sided Brunner-Munzel

nonparametric test of log-transformed TPMs vs. all other utORFs, $p << 0.001$). This is likely due to active

transcription of the overlapping pre-existing loci. In contrast, fast-evolving utORFs are significantly less

well-transcribed (same test, $p < 0.001$), while differences in *melanogaster*-specific and utORFs are

statistically insignificant. Consistent with these trends, alternative-frame utORFs are less tissue- and

developmentally-specific (one-sided Brunner-Munzel test of tau across FlyAtlas2 tissues, $p < 0.001$; same

test of tau across modENCODE developmental stages, $p < 0.001$) while fast-evolving utORFs are more

specifically-expressed (tau across FlyAtlas2 tissues, $p = 0.0039$; tau across modENCODE developmental

stages, $p < 0.001$). Unusually, a plurality of fast-evolving utORFs (33 of 96, 34.4%) have their highest

expression in adult fly tissues in the brain, with the next most common location the testis (10 of 96,

10.4%). Moreover, while transcription of annotated genes is high in both the brain and testis, fast-

evolving utORFs are dramatically less-transcribed in the testis than in the brain (75th percentile $\log_{10}$ of

TPM in brain, fast-evolving utORFs: 0.172; vs. annotated genes: 1.320; in testis, fast-evolving utORFs: -

0.662; vs. annotated genes 1.177; delta $\log_{10}$ of TPM in testis versus in brain, fast-evolving utORFs

vs. annotated genes: one-sided Mann-Whitney U test $p = 7.5E-6$; **Figure 5–figure supplement 1**).

Indeed, with the exception of alternate-frame utORFs, most utORFs are not transcribed in the testis but

are transcribed in the brain, while most annotated genes are highly transcribed in both (**Figure 5–figure**

**supplement 1**). This is somewhat intriguing as previous work suggests that evolutionarily young genes –

particularly *de novo* genes – emerge in the testis.

When considering only utORFs with canonical start codons, the results are generally qualitatively similar
(**Figure 4–figure supplement 2, Figure 5–figure supplement 2**). Five latent classes remain the best fit,
as allowing a sixth class has unfavorable AIC and BIC effects. Class assignments can proceed with
similar logic; however, utORFs with start codons are longer overall (**Figure 4–figure supplement 2B**),
so some class interpretations change somewhat. Class 3 appears biased towards intergenic locations and
longer length, so we assign it an interpretation of "intergenic ORFs" (**Supplemental File 1B**).
Partitioning between classes remains good (**Figure 4–figure supplement 3**). Turning once again towards
forward inferences, we recapitulate the patterns seen for the conservation of *melanogaster*-specific and
alternative-frame ORFs. While differences in length are apparent between classes (**Figure 5–figure
supplement 2D**), they are not statistically significant; the *melanogaster*-specific class still is not
significantly shorter (one-sided Brunner-Munzel nonparametric test of lengths vs. all other utORFs:
intergenic ORFs, longer, $p = 0.11$; alternate-frame, shorter, $p = 0.15$; *melanogaster*-specific, shorter, $p =$
$0.36$; putatively nonfunctional, shorter, $p = 0.20$). Finally, utORFs with canonical start sites remain less
transcribed in the testis than the brain (delta log base 10 of TPM in testis versus in brain, utORFs with
canonical start sites vs. annotated genes: one-sided Mann-Whitney U test $p = 7.3E-21$). Non-canonical
start initiation is increasingly recognized, especially for short ORFs; evidence for translation initiation at
sites other than AUG has come from both ribosome profiling and mass spectrometry approaches from a
variety of model species (Chen et al., 2020; Ingolia et al., 2011; Ma et al., 2014; Wu et al., 2020). Since
we recapitulated similar results for utORFs with canonical start sites as for all utORFs, we continued with
all utORFs.

**Many unannotated translated ORFs have evidence consistent with a *de novo* origin**

A genetic locus emerging through *de novo* gene birth from nongenic sequence would be expected to be
detected in a lineage-specific manner with detectable genomic homology despite evidence inconsistent
with functional conservation. That is, a utORF may be of *de novo* origin if there exists at least one species
where an orthologous ORF is not significant despite genomic homology. Ideally, the strongest inferences
of *de novo* origin would rely on multiple outgroup species. This protects against an erroneous *de novo*
inference due to a homology detection failure rather than true functional changes, which has been
extensively described as a potential source of erroneous identifications (Moyers and Zhang 2015, 2016,
2018). Many of our utORFs have such stringent evidence consistent with a *de novo* origin (**Figure 6A**),
with at least two supporting outgroups originating from different phyla. Between 40 and 75% of
identified loci across all inferred classes may be *de novo*, with the greatest proportion in the fast-evolving,
*melanogaster*-specific, and unannotated ORFs. These differences between classes are statistically
significant (Pearson's Chi-squared test, 4 degrees of freedom, $X^2 = 31.941$, $p = 1.97E-6$). For all utORFs

with potential *de novo* origin, 394 (67.8%) have 3 or more such supporting outgroup species. The number

of such supporting outgroup species appears to be more affected by the availability of such than by

genomic divergence, as the youngest classes (*melanogaster*-specific and fast-evolving) have more

supporting outgroup species than the comparatively older alternative-frame class (**Figure 6B**).

Finally, we considered the frequency by which utORFs of potential *de novo* origin is lost. Since we

consider the protein sequence to be equivalent to the function, we use similar logic as for finding support

for *de novo* origin, but looking only at species descended from the inferred potential de novo origin event.

We find that losses of utORFs of potential *de novo* origin are extremely common; for all utORFs with

potential *de novo* origin, 470 (80.9%) have at least 1 potential inferred loss (**Figure 6C**).

**Observed retention times of many peptides underlying unannotated translated ORFs are similar to**
**those of peptides underlying annotated proteins**

One possible weakness of the methodology used in this study is that peptide-spectrum matches (PSMs)

are prone to false identifications (Nesvizhskii and Aebersold 2005). One method that has been proposed

to evaluate PSMs is to use the retention time of identified peptides on the liquid chromatography system

upstream of the mass spectrometer; retention time should vary as a function of the peptide's sequence.

While calculating retention time of a given peptide *a priori* remains an unsolved problem, substantial

progress has been made on predicting retention times of sets of peptides *a posteriori*. To evaluate the

validity of PSMs of utORFs, we used AutoRT (Wen *et al*. 2020) to predict retention times of peptides

supporting annotated FlyBase proteins as well as utORFs. While some peptides supporting utORFs have

drastically larger differences from the predicted retention times (**Figure 7–figure supplement 1**), a

sizeable number have observed retention times that are similar to peptides from annotated FlyBase

proteins. Specifically, 385 of 3123 peptides (12.3%) supporting utORFs have at least one PSM with a

delta retention time that is smaller than the 95th percentile of the delta retention times of all PSMs that

support annotated FlyBase proteins; within this subset, the difference between predicted and observed

retention times is modest (**Figure 7A**). These peptides are sufficient to support 270 utORFs, where

melanogaster-specific ORFs are slightly more likely and nonfunctional ORFs somewhat less likely to be

supported (**Supplemental File 1C**); however, this observation is not statistically significant (Chi-squared

test with 4 degrees of freedom; Chi-sq = 4.45; $p > 0.05$).

**utORFs can be independently validated**

To further validate that the discovered utORFs are not artifactual, we collected our own data to develop

independent empirical support for these utORFs. We collected targeted MS data from mixed 0-24 hour *D.*

*melanogaster* embryos (RAL517 strain), targeting a manually-curated set of 15 target utORFs chosen

both for suitability for targeted MS (e.g. few missed cleavages; no post-translational modifications) as

well as expression in embryo from modENCODE data. Of these 15 targeted utORFs, we were able to

detect supporting evidence for 1 through the observation of 3 transitions (**Figure 7B**). We also collected

shotgun MS data from mixed 0-24 hour embryos; we were able to detect supporting evidence for 3

additional utORFs from the shotgun MS data at a FDR of 1% (**Figure 7-Source Data 1**); this shotgun

dataset was fairly shallow, as we identified 1963 annotated proteins compared to the approximately 8000

protein groups reported by Casas-Vila et al. (2017). Given the low sensitivity of MS, having evidence for

utORFs from independent datasets supports an unneglectable translation level for at least a subset of the

utORFs. Furthermore, we re-analyzed published ribosome profiling data (Patraquim et al. 2020; Zhang et

al. 2018) and identified supporting evidence from an additional 243 utORFs. Of these, alternate-frame

ORFs are unsurprisingly the most frequently supported, perhaps due to translation of the canonical ORF,

but the other classes are well-represented with the exception of the putatively nonfunctional utORFs

(**Figure 7–figure supplement 2**). The biases across classes are statistically significant (Pearson's Chi-

squared test, 4 degrees of freedom, $X^2 = 41.569$, $p \ll 0.001$), and they are consistent with our proposed

interpretation of the classes. Finally, we specifically looked for evidence of framing in ribosome profiling

data by implementing the binomial statistical test used for the first report of framing in *Drosophila*

(Patraquim et al., 2020). Of 81 utORFs with more than 2 unique footprints, 17 utORFs (21.0%) have

statistically-significant biases towards the observed frame in at least 1 stage of embryogenesis, supporting

the notion of active translation (alpha 0.05; 11 utORFs at alpha 0.001).

Together, this evidence supports our identification of the pool of utORFs overall. We accordingly

integrated all data that we analyzed with respect to additional lines of evidence supporting our utORFs

and factors of interest in order to identify utORFs of highest priority for downstream functional analyses;

we used a combination of high transcription from RNAseq atlases, supporting validation from either our

MS experiments or framed ribosome profiling reads, and conservation metrics. By calculating ranks

across each of these factors and averaging ranks per utORF, we generated a priority list for our utORFs

(**Figure 7 - Source Data 2**). Of the top 50 utORFs, 24 are alternate-frame ORFs, 14 unannotated ORFs, 7

melanogaster-specific ORFs, and 4 fast-evolving ORFs. This is similar to the overall proportion of class

assignments, except alternate-frame ORFs are somewhat over-represented due a higher incidence of

framed ribosome profiling reads. These utORFs may be the most promising candidates for future

experimental work.

**Discussion**

**Identification of hundreds of potential protein products of *de novo* origin**

In this work, we identify almost one thousand possible utORFs from a systematic, ORF-focused MS-first

computational approach. These utORFs are unlikely to be annotation failures due to the lack of sequence

homology to annotated proteins. Instead, like many similar "orphan genes", these may be of *de novo*

origin: the vast majority have a syntenic sequence conservation pattern that is consistent with *de novo*

origin (**Figure 6AB**). The total number of identifications is larger than previous studies of *de novo* genes

within a relatively short timescale in *Drosophila* (Begun et al. 2006; Chen, Zhang, and Long 2010; Zhao

et al. 2014), and is consistent with the identification of thousands of possible *de novo* polypeptides from

systematic studies in other biological systems (Ruiz-Orera et al. 2018; Stein et al. 2018; Durand et al.

2019). Moreover, while evidence of translation of proposed *de novo* genes is generally lacking

(McLysaght and Hurst 2016), in this case, the method of identification allows a stronger inference of

translation.

Calling *de novo* origin for a given gene is challenging, as the call is almost wholly dependent on negative

results in the outgroup lineages. For example, in small phylogenies with a single outgroup, it is possible

for a single false negative call to lead to an incorrect inference of *de novo* origin. Recent studies have

demonstrated that false negatives may be alarmingly likely, especially when using simple BLAST-based

phylostratigraphy (Moyers and Zhang 2015, 2016). Here, our method of calling *de novo* origin employs

proposed best practices of incorporating syntenic information (Arendsee et al. 2019; Vakirlis and

McLysaght 2019) and is robust to individual sequence homology calls, improving our confidence in our

inferences. Importantly, our inferences of gene age and potential *de novo* origin do not rely at all on the

absence of sequence from the MSA, which would be analogous to the lack of a significant BLAST hit.

Instead, our inferences of *de novo* origin rely on protein similarity scores below our threshold of

significance (**Figure 6B**), which ensures that these inferences are not simply due to extensive genomic

divergence beyond the limit of detection. If anything, our formulation of the protein similarity score may

be vulnerable to calling spurious similarity due to shared genomic ancestry between closely related

species. However, such errors would cause us to infer systematically older *de novo* origin dates for each

gene; given that we still conclude that many utORFs are of potential *de novo* origin, the impact of such

possible errors is low.

Other mechanisms beyond *de novo* origin could potentially produce "orphan" / lineage-specific genes.

For example, horizontal gene transfer from microorganisms like viruses has been reported in *Drosophila*

(Verster et al., 2019), and much attention has been focused on sequence divergence (e.g. after

pseudogenization) as an alternative explanation for lineage-specific genes (Moyers and Zhang, 2015

Weisman et al., 2020).  When, as is often the case for our utORFs, there are many supporting outgroups,

multiple pseudogenization events are necessary by parsimony to produce the observed pattern; while this

seems unlikely, we cannot exclude the possibility of repeated substantial divergence. However, work

explicitly focusing on the possibility that sequence divergence is responsible for lineage-specific genes

recently concluded that extensive divergence was not likely the dominant mechanism (Vakirlis et al.,

2020).

Our confidence in our identifications of utORFs is supported by our confirmatory results from mass

spectrometry and published ribosome profiling, including the observation of statistically significant

framing from ribosome-profiling data. While the absolute number of recapitulated utORF detections is

relatively low, mass spectrometry has notoriously low power, even in well-studied proteomes like humans

(Baker et al. 2017). Thus, it is not particularly surprising that we were unable to recapitulate most utORFs

in a relatively heterogenous sample, as we expect utORFs to be of low abundance and potentially only

expressed under specific time- and tissue- limited conditions. Unlike sequencing approaches, the

empirical readout of mass spectrometry is not directly mappable to information (i.e. the underlying

nucleotide sequence), so false identifications of peptide-spectrum matches are a notorious problem in the

field (Nesvizhskii and Aebersold, 2005). However, the peptides underlying our identification of utORFs

are not artifacts caused by polymorphism in conserved genes. Moreover, our results suggest that the

reverse-decoy approach for controlling the number of false-positive identifications remains valid for our

two-round method: the power for the identification of annotated Drosophila genes is nearly the same

between a standard MS approach and our two-round approach (**Supplemental File 1A**). Further

functional investigation of utORF products would be a potential extension of this work (**Figure 7 -**

**Source Data 2**).

**Class inference reveals heterogenous evolutionary dynamics**

In contrast to previous work, our approach of applying latent class analysis allows additional refinement

beyond that achievable by simple per-variable analyses. For example, previous studies have described an

association between gene age and length: namely, that younger genes tend to be shorter (Carvunis et al.

2012; Neme and Tautz 2013; Wilson et al. 2017). Here, we find that the youngest class of utORFs (the

*melanogaster* specific class) are not significantly shorter; instead, the fast-evolving and the nonfunctional

classes are the shortest overall (**Figure 5C**). Importantly, while length is an input into the latent class

analysis, it is not the most obvious differentiator between the classes, nor is it the primary basis upon

which we interpreted the classes. Since the fast-evolving class is still quite young, it is possible that

previous reports of an association between gene age and length are related to a conflation of fast evolution

with gene age. Importantly, our analysis is not able to evaluate whether there is a direction in the

historical evolutionary trajectory of the length of a given utORF. Thus, we cannot distinguish between the

possibilities that shorter utORFs are faster evolving or that faster evolving ORFs require a shorter length.

This question merits further examination in future.

Previous studies, including our own, have suggested that evolutionarily young genes tend to be expressed

in the testis (Levine et al. 2006; Begun et al. 2007; Soumillon et al. 2013; Zhao et al. 2014; Luis

Villanueva-Cañas et al. 2017; Witt et al. 2019), though other tissues, like the brain, can give rise to

evolutionarily young genes (Chen et al., 2012). In contrast to that, the utORFs identified here are

expressed more in the brain than in the testis (**Figure 5–figure supplement 1**) in each inferred class.

Moreover, while most annotated genes are highly expressed in the testis, most utORFs are not expressed

at all in the testis, yet they are expressed in the brain (**Figure 5–figure supplement 1**). Considering that

most previously-reported *de novo* genes also tend to be expressed in the testis (Levine et al. 2006; Begun

et al. 2007; Zhao et al. 2014; Luis Villanueva-Cañas et al. 2017; Witt et al. 2019), our results may suggest

that the proportion of protein-coding *de novo* genes might be higher in brain than in testis, or that our MS-

first approach allows the detection of otherwise-missed protein-coding *de novo* genes. Furthermore, these

utORFs may have biologically significant effects in the brain. For example, the maximum observed

expression of a plurality of the relatively short, fast-evolving ORFs is in the brain, suggesting that some

could act as neuropeptides.

Despite the success of our latent class analysis approach in elucidating distinct evolutionary trajectories,

half of our identified utORFs of interest remain consolidated in a single class. However, additional classes

cannot be safely supported due to overfitting. Additional approaches may be necessary in order to further

characterize these utORFs.

**Turnover in utORFs may be frequent**

While *de novo* gene birth has been increasingly well-studied over recent years, *de novo* gene death has

been examined less-closely. It is obvious that *de novo* gene death must be relatively frequent, as otherwise

the net size of proteomes would be unrealistically large relative to the annotation. Consistent with this

intuition, we observe that most *de novo* genes diverge by enough to be indistinguishable from random

sequences in at least part of the clade descended from the last common ancestor after the *de novo* gene

birth event (**Figure 6C**). Since we consider conservation of the protein-coding sequence to be

approximately equivalent to the function of the locus, we consider such sequence divergence as evidence

of gene death.

An implication of most models of *de novo* gene origins is that *de novo* gene birth is rare at the individual

gene level and thus occurs only once in evolutionary history. However, the very large number of gene

death events suggests a second possibility. It is possible that the rates of gene birth and death may be

more balanced; that is, instead of being born once and recurrently lost, a subset of *de novo* genes may be

born, die, and subsequently reborn (Schmitz et al., 2018; Stein et al., 2018; Durand et al., 2019). Thus,

some *de novo* gene evolution may not proceed unidirectionally, but rather like a random walk along a

continuum. This is consistent with the proto-gene model (Carvunis et al. 2012) or the pre-adaptation

model (Wilson et al. 2017). Under the proto-gene model, the frequent deaths of our utORFs would imply

that they are proto-genes and not genes. However, with the exception of the *melanogaster*-specific

utORFs, our utORFs are often conserved across species with significant divergence times, which implies

that some may have evolutionary constraints and may in fact be functional genes. It is difficult to make

strong conclusions here without getting trapped in the philosophical quagmire of the nature of "gene-

ness"; nevertheless, it is possible that no single "grand unified theory" of *de novo* gene origination can

describe all phenomena.

**Outlook**

Together, our results show that evolution of young proteins may progress along different, distinct

trajectories in *Drosophila*. Whether similarly distinct trajectories are apparent in other model species such

as yeast or mammals remains to be seen. Of note, *Drosophila* is a taxon of multicellular organisms with a

large effective population size, so selective processes are more efficient; mammals – especially primates

and *Homo* – are evolutionarily young and have a smaller effective population size, while yeasts are

unicellular. If these factors affect general evolutionary properties, such as the selective cost of translation

of lowly functional proteins and the probability of fixation by drift, it is possible that they may affect the

evolution of *de novo* proteins. In the case of *Homo*, all these factors may be more favorable to the fixation

of new *de novo* proteins, and the availability of broad and varied -omics data is unparalleled. It would

therefore be an obvious extension to employ a similar approach to investigate possible utORFs and *de*

*novo* proteins in humans.

**Methods**

**Comprehensive *Drosophila melanogaster* ORFeome**

To generate the comprehensive database of potential ORFs in *Drosophila melanogaster*, we translated the

repeat-masked genome (UCSC Aug. 2014 BDGP Release 6 + ISO1 MT/dm6) and transcriptome

(FlyBase, r6.15). We used the repeat-masked genome to control the number of transposable elements and

other degenerate candidates. We retained only ORFs of at least 8 residues; for ORFs not containing a

canonical start codon, only those of at least 20 residues were retained. We deduplicated potential ORF

sequences, leaving 4,583,941 unique potential ORFs.

**Discovery of utORFs in published shotgun MS data**

To identify evidence of utORFs, we reanalyzed a published proteome of *D. melanogaster* (PRIDE accession numbers PXD005691 and PXD005713; respectively, whole lifecycle and hourly/bihourly embryogenesis stages; (Casas-Vila et al. 2017)) using MaxQuant v. 1.6.1.0 (Tyanova, Temu, and Cox 2016). To speed computation, we divided analysis of the overall proteome into 4 subsets (**Figure 1 - Source Data 4**). As in (Casas-Vila et al. 2017), we used standard parameters (namely, methionine oxidation and N-terminal acetylation as variable modifications; carbamidomethylation as a fixed modification; tryptic digestion; minimum peptide length 7).

*Two-round MS search*

In the two-round MS search, we used two rounds of analysis per data subset to improve total sensitivity while maintaining an acceptable false discovery rate (**Figure 1A**). In the first "discovery" round, we set the peptide and protein group FDRs in MaxQuant to 0.2, with the protein search sequences set to be the annotated *D. melanogaster* proteome (FlyBase, r6.15), supplemented with the comprehensive potential ORF database. We then identified candidate ORFs from any of the four data subsets (**Figure 1-Source Data 4**) to develop a filtered subset of potential ORFs with possible mass spectrometry evidence. These relaxed parameters allow us to identify many more potential ORFs, but with a corresponding increase in false positive identifications. In the second "verification" round, we set the peptide and protein group FDRs to 0.01, with the protein search sequences set to be the annotated *D. melanogaster* proteome (FlyBase, r6.15), supplemented with only that filtered subset of potential ORFs identified in the first round (around 1000-fold fewer). We subsequently performed additional manual curation of the identified hits to arrive at our final set of utORFs; in brief, we dropped utORFs that were shorter than 14 residues, that had detectable sequence homology to annotated proteins via BLASTP, and the shorter of multiple possible utORFs that were identified from the same protein group by MaxQuant.

*Evaluation of peptide and protein polymorphisms*

Evaluating whether a given peptide could have arisen from a polymorphism is equivalent to calculating the number of substitutions from said peptide to the nearest peptide expected from the annotated proteome. Variations of this problem are well-studied in computer science, and trie data structures are commonly used to solve them efficiently. We used a Python implementation of tries (https://github.com/jfjlaros/dict-trie) to build a trie of all tryptic peptides that would be expected from the annotated *D. melanogaster* proteome. For each peptide uniquely identified from a utORF, we calculated Hamming (the number of substitutions without the possibility of indels) and Levenshtein (the combined number of substitutions and indels) distances.

*Genomic analysis of utORFs*

To map utORF sequences back to their genomic locations, we used BLAT version 36 (Hinrichs et al.

2006) to identify the locations of utORF sequences that were supported in the discovery round. We

reformatted the output to a GTF and made certain manual corrections; this file, containing the final

locations, is appended as **Figure 1–Source Data 3**. We excluded those loci whose genomic location

could not be determined from genomic analyses. For those loci where multiple genomic locations were

possible, we used the first location identified by BLAT.

We subsequently classified the genomic locations of these utORF sequences using custom scripts relying

on the Bioconductor libraries in R (Huber et al. 2015). Specifically, we classified a location as sense or

antisense if any part of the location overlapped with an annotated gene in the appropriate direction. We

used the FlyBase r6.15 annotations. We classified a location as intergenic if and only if it was neither

sense nor antisense as defined above.

We computed conservation scores by using the 27-way phastCons and phyloP bigWig tracks provided by

the UCSC Genome Browser (Hinrichs et al. 2006). For a given genomic location, we calculated the

conservation score by averaging the score through the relevant interval.

*Published sequencing data*

We used a number of published sequencing datasets to characterize utORFs. To evaluate developmental

stage- and tissue-specific transcription levels, we used modENCODE (modENCODE Consortium et al.

2010) and FlyAtlas2 (Leader et al. 2018) data.

**Gene age inference**

*Multiple sequence alignment and genomic similarity scoring*

We used BioPython v. 1.69 (Cock et al. 2009) to subset the UCSC 27-way insect multiple-sequence

alignment (MSA) to a smaller MSA that corresponded to the location in the *D. melanogaster* genome of

each unannotated translated ORF plus the flanking 9bp on each side. In a pairwise fashion for each

species against *D. melanogaster*, we extracted genomic sequences from the MSA, dropped gaps in the

sequences, and calculated identity scores.

*Modified Smith-Waterman protein similarity scoring*

We calculated protein similarity scores using the Smith-Waterman algorithm with the BLOSUM62

matrix (Henikoff and Henikoff 1992), no gap or extension penalties, and we normalized by the length of

the ORF in *D. melanogaster*. To ensure score stability with respect to minor indels, we calculated protein

similarity scores in all three reading frames, keeping only the frame that scored best. In doing so, we

considered the entirety of the syntenic block irrespective of the formal boundaries of ORFs in the related

species; this allows us to capture the locally-optimal (within the block) similarity and tolerate recent

acquisition of nonsense or frameshift mutations.

*Phylostratigraphy*

Using the reference phylogeny corresponding to the UCSC 27-way insect MSA, we inferred approximate

gene emergence dates from the pattern of modified Smith-Waterman protein similarity scores. We

defined significance as a score greater than 2.5 points. In turn, we estimated the inferred age of a given

ORF as the age of the last common ancestor between *D. melanogaster* and the most distant insect species

for which a significant modified Smith-Waterman protein similarity score was calculated.

**Latent Class Analysis**

We performed latent class analysis (Collins and Lanza 2009) using the *poLCA* R package (Linzer and

Lewis 2011). Specifically, we used 100 iterations of a latent class model with no covariates and with

factors as follows: inferred gene age; monophyly; genomic location (classified as solely intergenic,

antisense, sense, or any combination); phastCons score (binned as [0, 0.2), [0.2, 0.8), and [0.8, 1.0]);

length (binned as [0, 20), [20, 50), and [50, infinity); transcription (binned as TPM > 0.1 or $\leq$ 0.1 from

maximum per-sample TPM) tissue specificity (binned as *tau* > 0.8 or $\leq$ 0.8 from FlyAtlas tissues)

developmental specificity (binned as *tau* > 0.8 or $\leq$ 0.8 from modENCODE developmental stages). We

excluded loci with missing values from latent class analysis. We chose the number of classes (five) based

on a combination of minimizing the BIC and AIC (**Supplemental File 1D and E**) as well as model

interpretation, similar to recommended procedures (Collins and Lanza 2009). For the analysis of all

utORFs, models with 6 classes did not converge on a single maximum-likelihood solution, suggesting

potential under-identification, and were thus not considered.

We subsequently used the resulting five-class latent class model to predict the posterior class membership

probabilities for each classified candidate locus. We used the modal probability to assign class

membership. We also performed a parallel analysis of utORFs with canonical start codons, which led to

similar conclusions.

**Validation of utORFs**
*Retention time validation*

We used AutoRT (Wen et al. 2020) to predict retention times of identified peptides from both annotated

FlyBase proteins and utORFs. To train the model, we conducted transfer learning per each source run

using the published AutoRT model (originally trained on PXD006109) with 40 epochs on either an

NVIDIA K80 GPU or an NVIDIA V100 GPU; the training set consisted of only those annotated FlyBase

peptides with posterior error probabilities of 0.01 or smaller as estimated by MaxQuant. We then used the

per-run trained models with default parameters to predict the retention times of the peptides resulting

from either the annotated FlyBase proteins or utORFs.

*Empirical validation*

We collected protein samples from *D. melanogaster* 0–24-hour embryos (RAL517 strain) for validation

of utORFs via mass spectrometry. In brief, we homogenized dechorionated embryos with a mortar and

pestle in ice-cold lysis buffer (50mM Tris HCl, pH7.5; 137 mM NaCl; 0.25% NP-40; Roche cOmplete

protease inhibitors) before centrifugation at 18,000 rpm at 4°C and filtration through a 0.45 micron PVDF

membrane.

Purified protein was acetone-precipitated, reduced, and alkylated before digestion with LysC and trypsin,

acidified, and solid-phase extraction. The resulting peptides were analyzed by LC-MS/MS (120min

gradient, 25cm EasySprayer column in high res./high mass accuracy mode) on a ThermoFisher Lumos

instrument. In addition, the embryo sample was analyzed on the same instrument by PRM-MS, targeting

a set of 16 tryptic peptides. We manually chose these targeted peptides based on suitability of the peptide

for targeting (i.e., few/no missed cleavages or modifications), expression of the utORF in embryos, and

the strength of the prior detection from the Casas-Vila datasets.

We analyzed the shotgun data using ProteomeDiscoverer 1.4 and searched via Mascot against the Uniprot

D. melanogaster database concatenated with common contaminants and our set of utORFs with a 1%

peptide FDR. We analyzed the targeted data using SkylineMS (v. 21.2.0.369 (2efacf038)). We deposited

the raw data in PRIDE under accession PXD032197.

We additionally reanalyzed published ribosome profiling data downloaded from the NCBI Sequence

Read Archive (SRA) project PRJNA306373 (Zhang et al. 2018) and project SRP254283 (Patraquim et al.

2020). For the Zhang dataset, we downloaded raw reads from the above SRA accessions, aligned them to

the genome using hisat2 v.2.2.1 (Kim et al. 2019), and calculated TPM using stringtie v.1.3.4d (Pertea et

al. 2016). We handled the Patraquim dataset similarly, except that we additionally trimmed the raw reads

using trimmomatic 0.38 (Bolger, Lohse, and Usadel 2014) and filtered out reads aligning to annotated

RNAs using bowtie2 v. 2.3.5 (Langmead and Salzberg 2012) before alignment. For each utORF, we

calculated the maximum TPM observed across either dataset and called the utORF as supported if the

TPM was greater than 0.2.

To identify utORFs with statistically significant framing of ribosome-profiling reads, we used a binomial test as in Patraquim et al. (2020). We aligned the trimmed, filtered reads to the set of sequences corresponding to our utORFs, plus an additional 30bp up- and down-stream, using bowtie v. 1.3.1 (Langmead et al., 2009) with options --tryhard --suppress 1,6,7,8. We analyzed the resulting aligned footprints using the R package riboSeqR v 1.20.0. For each utORF with at least two unique footprints, we calculated the binomial probability of observing the number of footprints in the utORF's frame versus the other two frames at an uncorrected alpha of 0.05. We pooled footprints across replicates and across read lengths for a given utORF in each sample.

## Data Availability

Raw MS data are deposited in PRIDE under accession number PXD032197. Relevant scripts and intermediate files can be found in our Github repository https://github.com/LiZhaoLab/utORF_mass_spec.

## Acknowledgements

We thank members of the Zhao laboratory for helpful discussions during the work. We thank Henrik Molina and the Proteomics Resource Center at the Rockefeller University for the help with mass spectrometry data.

## Funding

This work was supported by National Institutes of Health (NIH) MIRA R35GM133780, the Robertson Foundation, a Rita Allen Foundation Scholar Program, and a Vallee Scholar Program (VS-2020-35), a Monique Weill-Caulier Career Scientist Award, and an Alfred P. Sloan Research Fellowship (FG-2018-10627) to L.Z. E.B.Z. was supported by a Medical Scientist Training Program grant from the National Institute of General Medical Sciences of the National Institutes of Health under award number T32GM007739 to the Weill Cornell/Rockefeller/Sloan Kettering Tri-Institutional MD-PhD Program. The content of this study is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health.

## Competing interest statement

The authors declare no competing interests.

 **Table and Figure Legends**

668 **Table 1: Latent class analysis of all unannotated translated ORFs.**

| Class | Interpretation | Estimated Percent | Number |
|---|---|---|---|
| 1 | putatively nonfunctional loci | 4.35% | 41 |
| 2 | melanogaster-specific ORFs | 5.71% | 54 |
| 3 | fast-evolving ORFs | 12.03% | 96 |
| 4 | unannotated ORFs | 57.61% | 591 |
| 5 | alternative frame ORFs | 20.30% | 161 |

**Figure 1. Identification of utORFs and their properties**. **A**. We detected utORFs by searching via a
two-round approach through a comprehensive database of potential ORFs that was generated from a six-
frame translation of the transcriptome and genome (see Methods). **B**: an example utORF. The full
sequence of utORF 633.1 is shown. The peptide supporting the inference of this utORF is bolded and
colored red. **C**: distribution of lengths of utORFs. **D**: distribution of amino acids in utORFs (blue) and in
the reference D. melanogaster proteome (orange). Standard single-letter abbreviations for amino acids are
used. **E**: genomic locations of utORFs. Since a utORF can overlap multiple items, e.g., an ORF that
overlaps with an annotated intron, exon, and another gene antisense to the first, categories are not
exclusive. **F**: left: secondary structure composition of utORFs. Approximately half of utORFs have a
coiled-coil secondary structure, with the remainder a mixture of beta sheets and alpha helices. right:
predicted structural disorder of utORFs. **G**: Percent GC content of utORFs and GC content of the X and
autosomes of the *D. melanogaster* genome (red line).

**Figure 2. Synteny-based orthology detection and protein sequence similarity quantification. A**:
When performing a simple homology search for a locus of interest (red arrow) across a given genome
(blue line), the search space is orders of magnitude larger, requiring heuristic shortcuts to reduce the
computational complexity. **B**: Using a multiple sequence alignment, we can simply find the block that
contains the locus of interest and use that to evaluate potential orthologs in other genomes. The search
space is approximately a similar size as that of the original locus, so an optimal search is computationally
feasible. Syntenic information (i.e., flanking genes, gray arrows) is encoded within the multiple sequence
alignment. **C**: We calculated the pairwise protein similarity score (see **Methods**) across 10000 pairs of
randomly-generated sequences of lengths 20-150 with amino acid frequencies matching that of the
annotated *D. melanogaster* proteome. The mean score (blue line) and 2 standard deviations (shading) are
shown. Our significance threshold of 2.5 lies beyond 2 standard deviations from expectation.

**Figure 3: Inferred gene ages of the utORFs. A**: The reference phylogenetic tree used for these analyses

(UCSC 27-way insect alignment). Abbreviations are as follows: *D. mel*: *Drosophila melanogaster*, *D.*

*sim*: *D. simulans*, *D. sec*: *D. sechellia*, *D. yak*: *D. yakuba D. ere*: *D. erecta*, *D. bia*: *D. biarmipes*, *D. suz*:

*D. suzukii, D. ana*: *D. ananassae*, *D. bip*: *D. bipectinata*, *D. eug*: *D. eugracilis*, *D. ele*: *D. elegans*, *D. kik*:

*D. kikkawai, D. tak*: *D. takahashii, D. rho*: *D. rhopaloa, D. fic*: *D. ficusphila, D. pse*: *D. pseudoobscura*,

*D. per*: *D. persimilis, D. mir*: *D. miranda, D. wil*: *D. willistoni, D. vir: D. virilis, D. moj: D. mojavensis*,

*D. alb: D. albomicans, D. gri*: *D. grimshawi, M. dom*: *Musca domestica, A. gam*: *Anopheles gambiae*,

*Apis mel*: *Apis mellifera, T. cas*: *Tribolium castaneum*. **B**: The most distantly related species in which a

significant ortholog of a utORF exists varies. The red line illustrates the cumulative distribution of loci.

For convenience, sister species are grouped together under one species (e.g., *D. virilis* with *D. grimshawi*;

*D. eugracilis, D. elegans, D. kikkawai*, and *D. takahashii* with *D. rhopaloa*; etc.). Abbreviations as in **A**.

**Figure 4. Latent class analysis of the utORFs reveals differences between classes**. **A**: Class 1 is

notably distinct for strong bias towards intergenic and antisense locations at the expense of sense

locations. Class 2 is notable for being relatively unbiased, and for being the only class with appreciable

members in a combination of locations. Class 5 is strongly skewed towards antisense and sense locations.

**B**: Class 1 is almost exclusively of intermediate length. Class 4 has the greatest length bias, followed by

class 2. tiny: fewer than 20 residues; medium: from 20 to 49 residues; long: 50 or more residues. **C**: Class

1 is notably distinct from the others for minimal transcription. Low tissue and developmental specificity

may be an effect of minimal transcription. Class 2 is remarkable for being entirely monophyletic. Class 5

has slightly lower tissue and developmental specificities than classes 2-4. sigTPM: maximum per-sample

TPM 0.1; tau: tissue specificity 0.8; dev_tau: developmental specificity 0.8. **D**: Class 2 is by far the

youngest. Class 3 tends to be of an intermediate age, with inferred emergence at around the LCA of the

*melanogaster* subgroup (droRho). In contrast, class 4 emergence is distributed throughout the LCA of the

*melanogaster* subgroup and the *Drosophila* genus. **E**: Class 2 is notable for overall low conservation.

Class 3 is remarkably even less conserved. Class 4 and 5 are distinguished through differences in

intermediate vs. significant conservation. unconserved: phastCons score < 0.2; amb: phastCons score ≥

0.2 and < 0.8; significant: phastCons score ≥ 0.8.

**Figure 5. Differences between inferred classes recapitulate expected trends in age and conservation**

**and reveal surprising trends in lengths and expression**. **A**: As expected, phastCons conservation scores

vary by class. Scores near 0 indicate low conservation while scores near 1 indicate high conservation.

Note that fast-evolving and mel-specific loci have dramatically lower conservation whereas alt-frame loci

are very well conserved. **B**: Despite not being included as a variable in LCA, phyloP conservation scores

also vary by class. Scores below 0 indicate potential fast evolution (acceleration) while scores above 0

indicate conservation. Note that the lowest scores predominantly occur in fast-evolving and mel-specific

loci, and that alt-frame loci remain the best-conserved. **C**: Distribution of phylostratigraphically-inferred

gene ages by inferred latent class. **D**: Class variations in length are generally minimal, with the exception

of utORFs being slightly longer than average, and fast-evolving and nonfunctional loci being shorter than

average. **E**: Maximum observed transcription across all FlyAtlas2 and modENCODE samples varies by

class. **F**: Tissue specificity (tau) calculated from FlyAtlas2 data shows that nonfunctional and fast-

evolving loci are the most specific, with alternate frame loci the least. **G**: Developmental specificity (tau)

calculated from modENCODE data similarly shows that nonfunctional and fast-evolving loci are the most

specific, with alternate frame loci the least.

**Figure 6. Many utORFs have evidence consistent with a de novo origin. A**: Proportion of utORFs by

inferred class with genomic conservation consistent with *de novo* origin. Box widths correlate with size of

class (**Table 1**). **B**: Number of supporting outgroups by inferred class for loci with potential *de novo*

origin. **C**: Number of losses inferred after potential de novo origin.

**Figure 7. Independent validation of utORF identification. A:** Cumulative distribution of differences

between observed and predicted retention times for peptide-spectrum matches of peptides supporting

annotated FlyBase proteins (orange) and peptide-spectrum matches of peptides supporting utORFs that

are no worse than the 95[th] percentile of annotated FlyBase PSMs (blue). **B:** Transitions detected via PRM

of the peptide GPGAAISQR from protein extracts of *D. melanogaster* embryos.

**Supplemental Figure Legends**

**Figure 3–figure supplement 1:** Robustness of gene age inferences with respect to significance threshold.

**A**: Figure 3A, but also showing the *melanogaster* species subgroup, species group, and *Drosophila* taxa.

**B**: Change in furthest significant ortholog (using a significance threshold of 2.3 instead of 2.5). Coloring

corresponds to panel A. Lowering the threshold causes more orthologs of utORFs to be called as

significant, increasing the inferred gene age. **C**: As in panel B, but with a significance threshold of 2.7

instead of 2.5. Increasing the threshold causes fewer orthologs of utORFs to be called as significant,

reducing the inferred gene age.

**Figure 4–figure supplement 1:** Posterior probabilities per unannotated translated ORF of class

membership inferred from latent class analysis for all utORFs.

**Figure 5–figure supplement 1:** Transcription of utORFs in selected tissues. Top panel: utORFs,

separated by inferred LCA class. Mean TPMs across the given tissue in FlyAtlas2 are log10-transformed

with a pseudocount of 1E-3. Horizontal line marks an expression threshold of TPM 0.1. Bottom panel: As

top, but for, in order, annotated FlyBase genes, separated between those with names, those with "CG"

identifiers, and those with "CR" identifiers, and all utORFs.

**Figure 4–figure supplement 2:** Latent class analysis of utORFs with canonical start sites reveals

differences between classes. **A-E:** same as Figure 4, but examining utORFs with canonical start sites.

**Figure 4–figure supplement 3:** Posterior probabilities per unannotated translated ORF of class

membership inferred from latent class analysis for utORFs with canonical start sites.

**Figure 5–figure supplement 2:** Differences between inferred classes for utORFs with canonical start

sites recapitulate expected trends. **A-G**: same as Figure 5, but examining utORFs with canonical start

sites.

**Figure 7–figure supplement 1**: Cumulative distribution of differences between observed and predicted

retention times for every peptide-spectrum match of peptides supporting all annotated FlyBase proteins

(blue) and all utORFs (orange).

**Figure 7–figure supplement 2**: Proportion of utORFs by inferred class with supporting evidence from

ribosome profiling. Box widths correlate with size of class (Table 1).

**Supplemental File 1A:** Two-round analysis improves identifications.

**Supplemental File 1B**: Latent class analysis of unannotated translated ORFs with canonical start sites.

**Supplemental File 1C**: AutoRT-supported identifications by inferred gene class.

**Supplemental File 1D**: Summary statistics used for selecting the number of latent classes when

considering all utORFs. Five latent classes were chosen as a balance of minimizing the AIC and BIC and

model interpretation. AIC: Akaike information criterion; BIC: Bayesian information criterion; $G^2$:

likelihood-ratio statistic; Chi-sq: Chi-squared. * : Models with 6 classes did not converge on a single

maximum-likelihood solution within 100 replicates.

**Supplemental File 1E**: Summary statistics used for selecting the number of latent classes when

considering only utORFs with canonical start codons. Five latent classes were chosen as a balance of

minimizing the AIC and BIC and model interpretation. AIC: Akaike information criterion; BIC: Bayesian

information criterion; $G^2$: likelihood-ratio statistic; Chi-sq: Chi-squared.

**Source Data Files**

Figure 1–Source Data 1 (csv): all utORFs sequences.

Figure 1–Source Data 2 (csv):  utORF supporting peptides.

Figure 1–Source Data 3 (gtf): utORF locations.

Figure 4 –Source Data 1 (csv): utORF inferred LCA classes.

Figure 7–Source Data 1 (xlsx): MASCOT search results from embryo MS data.

Figure 7–Source Data 2 (csv): rank of potentially biologically significant targets.

Figure 1–Source Data 4 (xlsx) - dataset subset mappings.

**References**

Andrews, S.J., and Rothnagel, J.A. (2014). Emerging evidence for functional peptides encoded by short
open reading frames. Nat Rev Genet *15*, 193–204. *https://doi.org/10.1038/nrg3520*.

Arendsee, Z., Li, J., Singh, U., Bhandary, P., Seetharam, A., and Wurtele, E.S. (2019). fagin: synteny-
based phylostratigraphy and finer classification of young genes. BMC Bioinformatics *20*, 440.
*https://doi.org/10.1186/s12859-019-3023-y*.

Baker, M.S., Ahn, S.B., Mohamedali, A., Islam, M.T., Cantor, D., Verhaert, P.D., Fanayan, S., Sharma,
S., Nice, E.C., Connor, M., et al. (2017). Accelerating the search for the missing proteins in the human
proteome. Nature Communications *8*, 1–13. *https://doi.org/10.1038/ncomms14271*.

Begun, D.J., Lindfors, H.A., Thompson, M.E., and Holloway, A.K. (2006). Recently evolved genes
identified from Drosophila yakuba and D. erecta accessory gland expressed sequence tags. Genetics *172*,
1675–1681. *https://doi.org/10.1534/genetics.105.050336*.

Begun, D.J., Lindfors, H.A., Kern, A.D., and Jones, C.D. (2007). Evidence for de novo evolution of
testis-expressed genes in the Drosophila yakuba/Drosophila erecta clade. Genetics *176*, 1131–1137.
*https://doi.org/10.1534/genetics.106.069245*.

Bolger, A.M., Lohse, M., and Usadel, B. (2014). Trimmomatic: a flexible trimmer for Illumina sequence
data. Bioinformatics *30*, 2114–2120. *https://doi.org/10.1093/bioinformatics/btu170*.

Bungard, D., Copple, J.S., Yan, J., Chhun, J.J., Kumirov, V.K., Foy, S.G., Masel, J., Wysocki, V.H., and
Cordes, M.H.J. (2017). Foldability of a Natural De Novo Evolved Protein. Structure *25*, 1687–1696.e4.
*https://doi.org/10.1016/j.str.2017.09.006*.

Cai, J., Zhao, R., Jiang, H., and Wang, W. (2008). De novo origination of a new protein-coding gene in
Saccharomyces cerevisiae. Genetics *179*, 487–496. *https://doi.org/10.1534/genetics.107.084491*.

Carvunis, A.-R., Rolland, T., Wapinski, I., Calderwood, M.A., Yildirim, M.A., Simonis, N., Charloteaux,
B., Hidalgo, C.A., Barbette, J., Santhanam, B., et al. (2012). Proto-genes and de novo gene birth. Nature
*487*, 370–374. *https://doi.org/10.1038/nature11184*.

Casas-Vila, N., Bluhm, A., Sayols, S., Dinges, N., Dejung, M., Altenhein, T., Kappei, D., Altenhein, B.,
Roignant, J.-Y., and Butter, F. (2017). The developmental proteome of Drosophila melanogaster. Genome
Research *27*, 1273–1285. *https://doi.org/10.1101/gr.213694.116*.

Chen, J., Brunner, A.-D., Cogan, J.Z., Nuñez, J.K., Fields, A.P., Adamson, B., Itzhak, D.N., Li, J.Y.,
Mann, M., Leonetti, M.D., et al. (2020). Pervasive functional translation of noncanonical human open
reading frames. Science *367*, 1140–1146. *https://doi.org/10.1126/science.aay0262*.

Chen, S., Zhang, Y.E., and Long, M. (2010). New genes in Drosophila quickly become essential. Science
*330*, 1682–1685. *https://doi.org/10.1126/science.1196380*.

Chen, S., Spletter, M., Ni, X., White, K.P., Luo, L., and Long, M. (2012). Frequent recent origination of
brain genes shaped the evolution of foraging behavior in Drosophila. Cell Rep *1*, 118–132.
*https://doi.org/10.1016/j.celrep.2011.12.010*.

Cock, P.J.A., Antao, T., Chang, J.T., Chapman, B.A., Cox, C.J., Dalke, A., Friedberg, I., Hamelryck, T.,
Kauff, F., Wilczynski, B., et al. (2009). Biopython: freely available Python tools for computational
molecular biology and bioinformatics. Bioinformatics *25*, 1422–1423.
*https://doi.org/10.1093/bioinformatics/btp163*.

Collins, L.M., and Lanza, S.T. (2009). *Latent Class and Latent Transition Analysis* (Hoboken, NJ, USA:
John Wiley & Sons, Inc.).

Cridland, J.M., Majane, A.C., Zhao, L., and Begun, D.J. (2022). Population biology of accessory gland-
expressed de novo genes in Drosophila melanogaster. Genetics *220*, iyab207.
*https://doi.org/10.1093/genetics/iyab207*.

de Klerk, E., and 't Hoen, P.A.C. (2015). Alternative mRNA transcription, processing, and translation:
Insights from RNA sequencing. Trends in Genetics *31*, 128–139.
*https://doi.org/10.1016/j.tig.2015.01.001*.

Domazet-Loso, T., and Tautz, D. (2003). An evolutionary analysis of orphan genes in Drosophila.
Genome Research *13*, 2213–2219. *https://doi.org/10.1101/gr.1311003*.

Domazet-Loso, T., Brajković, J., and Tautz, D. (2007). A phylostratigraphy approach to uncover the
genomic history of major adaptations in metazoan lineages. Trends Genet *23*, 533–539.
*https://doi.org/10.1016/j.tig.2007.08.014*.

Durand, É., Gagnon-Arsenault, I., Hallin, J., Hatin, I., Dubé, A.K., Nielly-Thibault, L., Namy, O., and
Landry, C.R. (2019). Turnover of ribosome-associated transcripts from de novo ORFs produces gene-like
characteristics available for de novo gene emergence in wild yeast populations. Genome Res *29*, 932–
943. *https://doi.org/10.1101/gr.239822.118*.

Faridi, P., Purcell, A.W., and Croft, N.P. (2018). In Immunopeptidomics We Need a Sniper Instead of a
Shotgun. Proteomics *18*, e1700464. *https://doi.org/10.1002/pmic.201700464*.

Henikoff, S., and Henikoff, J.G. (1992). Amino acid substitution matrices from protein blocks. Proc Natl
Acad Sci U S A *89*, 10915–10919. *https://doi.org/10.1073/pnas.89.22.10915*.

Hinrichs, A.S., Karolchik, D., Baertsch, R., Barber, G.P., Bejerano, G., Clawson, H., Diekhans, M.,
Furey, T.S., Harte, R.A., Hsu, F., et al. (2006). The UCSC Genome Browser Database: update 2006.
Nucleic Acids Research *34*, D590–8. *https://doi.org/10.1093/nar/gkj144*.

Huber, W., Carey, V.J., Gentleman, R., Anders, S., Carlson, M., Carvalho, B.S., Bravo, H.C., Davis, S.,
Gatto, L., Girke, T., et al. (2015). Orchestrating high-throughput genomic analysis with Bioconductor.
Nat Methods *12*, 115–121. *https://doi.org/10.1038/nmeth.3252*.

Ingolia, N.T., Lareau, L.F., and Weissman, J.S. (2011). Ribosome profiling of mouse embryonic stem
cells reveals the complexity and dynamics of mammalian proteomes. Cell *147*, 789–802.
*https://doi.org/10.1016/j.cell.2011.10.002*.

Kim, D., Paggi, J.M., Park, C., Bennett, C., and Salzberg, S.L. (2019). Graph-based genome alignment
and genotyping with HISAT2 and HISAT-genotype. Nat Biotechnol *37*, 907–915.
*https://doi.org/10.1038/s41587-019-0201-4*.

Knowles, D.G., and McLysaght, A. (2009). Recent de novo origin of human protein-coding genes.
Genome Research *19*, 1752–1759. *https://doi.org/10.1101/gr.095026.109*.

Lange, A., Patel, P.H., Heames, B., Damry, A.M., Saenger, T., Jackson, C.J., Findlay, G.D., and
Bornberg-Bauer, E. (2021). Structural and functional characterization of a putative de novo gene in
Drosophila. Nat Commun *12*, 1667. *https://doi.org/10.1038/s41467-021-21667-6*.

Langmead, B., and Salzberg, S.L. (2012). Fast gapped-read alignment with Bowtie 2. Nat Methods *9*,
357–359. *https://doi.org/10.1038/nmeth.1923*.

Langmead, B., Trapnell, C., Pop, M., and Salzberg, S.L. (2009). Ultrafast and memory-efficient
alignment of short DNA sequences to the human genome. Genome Biology *10*, R25.
*https://doi.org/10.1186/gb-2009-10-3-r25*.

Leader, D.P., Krause, S.A., Pandit, A., Davies, S.A., and Dow, J.A.T. (2018). FlyAtlas 2: a new version
of the Drosophila melanogaster expression atlas with RNA-Seq, miRNA-Seq and sex-specific data.
Nucleic Acids Research *46*, D809–D815. *https://doi.org/10.1093/nar/gkx976*.

Levine, M.T., Jones, C.D., Kern, A.D., Lindfors, H.A., and Begun, D.J. (2006). Novel genes derived from
noncoding DNA in Drosophila melanogaster are frequently X-linked and exhibit testis-biased expression.

Proceedings of the National Academy of Sciences of the United States of America *103*, 9935–9939. *https://doi.org/10.1073/pnas.0509809103*.

Li, D., Dong, Y., Jiang, Y., Jiang, H., Cai, J., and Wang, W. (2010). A de novo originated gene depresses budding yeast mating pathway and is repressed by the protein encoded by its antisense strand. Cell Res *20*, 408–420. *https://doi.org/10.1038/cr.2010.31*.

Linzer, D.A., and Lewis, J.B. (2011). poLCA : An R Package for Polytomous Variable Latent Class Analysis. Journal of Statistical Software *42*. *https://doi.org/10.18637/jss.v042.i10*.

Long, M., Betrán, E., Thornton, K., and Wang, W. (2003). The origin of new genes: glimpses from the young and old. Nature Reviews Genetics *4*, 865–875. *https://doi.org/10.1038/nrg1204*.

Luis Villanueva-Cañas, J., Ruiz-Orera, J., Agea, M.I., Gallo, M., Andreu, D., and Albà, M.M. (2017). New Genes and Functional Innovation in Mammals. Genome Biol Evol *9*, 1886–1900. *https://doi.org/10.1093/gbe/evx136*.

Ma, J., Ward, C.C., Jungreis, I., Slavoff, S.A., Schwaid, A.G., Neveu, J., Budnik, B.A., Kellis, M., and Saghatelian, A. (2014). Discovery of Human sORF-Encoded Polypeptides (SEPs) in Cell Lines and Tissue. Journal of Proteome Research *13*, 1757–1765. *https://doi.org/10.1021/pr401280w*.

McLysaght, A., and Guerzoni, D. (2015). New genes from non-coding sequence: the role of de novo protein-coding genes in eukaryotic evolutionary innovation. Philosophical Transactions of the Royal Society B: Biological Sciences *370*, 20140332. *https://doi.org/10.1098/rstb.2014.0332*.

McLysaght, A., and Hurst, L.D. (2016). Open questions in the study of de novo genes: what, how and why. Nature Reviews. Genetics *17*, 567–578. *https://doi.org/10.1038/nrg.2016.78*.

modENCODE Consortium, Roy, S., Ernst, J., Kharchenko, P.V., Kheradpour, P., Negre, N., Eaton, M.L., Landolin, J.M., Bristow, C.A., Ma, L., et al. (2010). Identification of functional elements and regulatory circuits by Drosophila modENCODE. Science (New York, N.Y.) *330*, 1787–1797. *https://doi.org/10.1126/science.1198374*.

Moyers, B.A., and Zhang, J. (2015). Phylostratigraphic bias creates spurious patterns of genome evolution. Mol Biol Evol *32*, 258–267. *https://doi.org/10.1093/molbev/msu286*.

Moyers, B.A., and Zhang, J. (2016). Evaluating Phylostratigraphic Evidence for Widespread De Novo Gene Birth in Genome Evolution. Mol Biol Evol *33*, 1245–1256. *https://doi.org/10.1093/molbev/msw008*.

Moyers, B.A., and Zhang, J. (2018). Toward Reducing Phylostratigraphic Errors and Biases. Genome Biol Evol *10*, 2037–2048. *https://doi.org/10.1093/gbe/evy161*.

Neme, R., and Tautz, D. (2013). Phylogenetic patterns of emergence of new genes support a model of frequent de novo evolution. BMC Genomics *14*, 117. *https://doi.org/10.1186/1471-2164-14-117*.

Nesvizhskii, A.I., and Aebersold, R. (2005). Interpretation of shotgun proteomic data: the protein inference problem. Mol Cell Proteomics *4*, 1419–1440. *https://doi.org/10.1074/mcp.R500012-MCP200*.

Ohno, S. (1970). *Evolution by Gene Duplication* (Berlin, Heidelberg: Springer Berlin Heidelberg).

Patraquim, P., Mumtaz, M.A.S., Pueyo, J.I., Aspden, J.L., and Couso, J.-P. (2020). Developmental
regulation of canonical and small ORF translation from mRNAs. Genome Biol *21*, 128.
*https://doi.org/10.1186/s13059-020-02011-5*.

Pertea, M., Kim, D., Pertea, G.M., Leek, J.T., and Salzberg, S.L. (2016). Transcript-level expression
analysis of RNA-seq experiments with HISAT, StringTie and Ballgown. Nature Protocols *11*, 1650–
1667. *https://doi.org/10.1038/nprot.2016.095*.

Reinhardt, J.A., Wanjiru, B.M., Brant, A.T., Saelao, P., Begun, D.J., and Jones, C.D. (2013). De Novo
ORFs in Drosophila Are Important to Organismal Fitness and Evolved Rapidly from Previously Non-
coding Sequences. PLoS Genetics *9*. *https://doi.org/10.1371/journal.pgen.1003860*.

Ruiz-Orera, J., Hernandez-Rodriguez, J., Chiva, C., Sabidó, E., Kondova, I., Bontrop, R., Marqués-Bonet,
T., and Albà, M.M. (2015). Origins of De Novo Genes in Human and Chimpanzee. PLoS Genet *11*,
e1005721. *https://doi.org/10.1371/journal.pgen.1005721*.

Ruiz-Orera, J., Verdaguer-Grau, P., Villanueva-Cañas, J.L., Messeguer, X., and Albà, M.M. (2018).
Translation of neutrally evolving peptides provides a basis for de novo gene evolution. Nature Ecology
and Evolution *2*, 890–896. *https://doi.org/10.1038/s41559-018-0506-6*.

Samandi, S., Roy, A.V., Delcourt, V., Lucier, J.-F., Gagnon, J., Beaudoin, M.C., Vanderperre, B., Breton,
M.-A., Motard, J., Jacques, J.-F., et al. (2017). Deep transcriptome annotation enables the discovery and
functional characterization of cryptic small proteins. Elife *6*, e27860. *https://doi.org/10.7554/eLife.27860*.

Schlötterer, C. (2015). Genes from scratch - the evolutionary fate of de novo genes. Trends in Genetics
*31*, 215–219. *https://doi.org/10.1016/j.tig.2015.02.007*.

Schmitz, J.F., Ullrich, K.K., and Bornberg-Bauer, E. (2018). Incipient de novo genes can evolve from
frozen accidents that escaped rapid transcript turnover. Nature Ecology and Evolution *2*, 1626–1632.
*https://doi.org/10.1038/s41559-018-0639-7*.

Sinitcyn, P., Rudolph, J.D., and Cox, J. (2018). Computational Methods for Understanding Mass
Spectrometry–Based Shotgun Proteomics Data. Annual Review of Biomedical Data Science *1*, 207–234.
*https://doi.org/10.1146/annurev-biodatasci-080917-013516*.

Soumillon, M., Necsulea, A., Weier, M., Brawand, D., Zhang, X., Gu, H., Barthès, P., Kokkinaki, M.,
Nef, S., Gnirke, A., et al. (2013). Cellular source and mechanisms of high transcriptome complexity in the
mammalian testis. Cell Rep *3*, 2179–2190. *https://doi.org/10.1016/j.celrep.2013.05.031*.

Stein, J.C., Yu, Y., Copetti, D., Zwickl, D.J., Zhang, L., Zhang, C., Chougule, K., Gao, D., Iwata, A.,
Goicoechea, J.L., et al. (2018). Genomes of 13 domesticated and wild rice relatives highlight genetic
conservation, turnover and innovation across the genus Oryza. Nat Genet *50*, 285–296.
*https://doi.org/10.1038/s41588-018-0040-0*.

Tautz, D., and Domazet-Lošo, T. (2011). The evolutionary origin of orphan genes. Nature Reviews.
Genetics *12*, 692–702. *https://doi.org/10.1038/nrg3053*.

Tyanova, S., Temu, T., and Cox, J. (2016). The MaxQuant computational platform for mass
spectrometry-based shotgun proteomics. Nature Protocols *11*, 2301–2319.
*https://doi.org/10.1038/nprot.2016.136*.

Vakirlis, N., and McLysaght, A. (2019). Computational Prediction of De Novo Emerged Protein-Coding
Genes. Methods Mol Biol *1851*, 63–81. *https://doi.org/10.1007/978-1-4939-8736-8_4*.

Vakirlis, N., Carvunis, A.-R., and McLysaght, A. (2020). Synteny-based analyses indicate that sequence
divergence is not the main source of orphan genes. Elife *9*, e53500. *https://doi.org/10.7554/eLife.53500*.

Van Oss, S.B., and Carvunis, A.-R. (2019). De novo gene birth. PLoS Genet *15*, e1008160.
*https://doi.org/10.1371/journal.pgen.1008160*.

Verster, K.I., Wisecaver, J.H., Karageorgi, M., Duncan, R.P., Gloss, A.D., Armstrong, E.E., Price, D.K.,
Menon, A.R., Ali, Z.M., and Whiteman, N.K. (2019). Horizontal Transfer of Bacterial Cytolethal
Distending Toxin B Genes to Insects. Mol Biol Evol *36*, 2105–2110.
*https://doi.org/10.1093/molbev/msz146*.

Wang, W., Brunet, F.G., Nevo, E., and Long, M. (2002). Origin of sphinx, a young chimeric RNA gene
in Drosophila melanogaster. Proc Natl Acad Sci U S A *99*, 4448–4453.
*https://doi.org/10.1073/pnas.072066399*.

Weisman, C.M., Murray, A.W., and Eddy, S.R. (2020). Many, but not all, lineage-specific genes can be
explained by homology detection failure. PLoS Biol *18*, e3000862.
*https://doi.org/10.1371/journal.pbio.3000862*.

Wen, B., Li, K., Zhang, Y., and Zhang, B. (2020). Cancer neoantigen prioritization through sensitive and
reliable proteogenomics analysis. Nat Commun *11*, 1759. *https://doi.org/10.1038/s41467-020-15456-w*.

Wilson, B.A., Foy, S.G., Neme, R., and Masel, J. (2017). Young genes are highly disordered as predicted
by the preadaptation hypothesis of de novo gene birth. Nature Ecology and Evolution *1*, 1–19.
*https://doi.org/10.1038/s41559-017-0146*.

Witt, E., Benjamin, S., Svetec, N., and Zhao, L. (2019). Testis single-cell RNA-seq reveals the dynamics
of de novo gene transcription and germline mutational bias in Drosophila. Elife *8*, e47138.
*https://doi.org/10.7554/eLife.47138*.

Wu, Q., Wright, M., Gogol, M.M., Bradford, W.D., Zhang, N., and Bazzini, A.A. (2020). Translation of
small downstream ORFs enhances translation of canonical main open reading frames. EMBO J *39*,
e104763. *https://doi.org/10.15252/embj.2020104763*.

Xie, C., Zhang, Y.E., Chen, J.-Y., Liu, C.-J., Zhou, W.-Z., Li, Y., Zhang, M., Zhang, R., Wei, L., and Li,
C.-Y. (2012). Hominoid-specific de novo protein-coding genes originating from long non-coding RNAs.
PLoS Genet *8*, e1002942. *https://doi.org/10.1371/journal.pgen.1002942*.

Xie, C., Bekpen, C., Künzel, S., Keshavarz, M., Krebs-Wheaton, R., Skrabar, N., Ullrich, K.K., and
Tautz, D. (2019). A de novo evolved gene in the house mouse regulates female pregnancy cycles. Elife *8*,
e44392. *https://doi.org/10.7554/eLife.44392*.

Zhang, H., Dou, S., He, F., Luo, J., Wei, L., and Lu, J. (2018). Genome-wide maps of ribosomal
occupancy provide insights into adaptive evolution and regulatory roles of uORFs during Drosophila
development. PLoS Biol *16*, e2003903. *https://doi.org/10.1371/journal.pbio.2003903*.

Zhang, L., Ren, Y., Yang, T., Li, G., Chen, J., Gschwend, A.R., Yu, Y., Hou, G., Zi, J., Zhou, R., et al.
(2019). Rapid evolution of protein diversity by de novo origination in Oryza. Nat Ecol Evol *3*, 679–690.
*https://doi.org/10.1038/s41559-019-0822-5*.

Zhao, L., Saelao, P., Jones, C.D., and Begun, D.J. (2014). Origin and spread of de novo genes in
Drosophila melanogaster populations. Science (New York, N.Y.) *343*, 769–772.
*https://doi.org/10.1126/science.1248286.*

Figure 1

Figure 2

## Figure 3

**A**

**B**

Figure 4

Figure 5

Figure 6

Figure 7

**A**

**B**

Figure 3 - Figure supplement 1

Figure 4 - Figure supplement 1

2962.1
1709.1
1004.1
130.1
2478.1
2430.1
1203.1
2645.1
4202.1
4880.1
2054.1
2176.1
630.1
2682.1
1318.1
2720.1
638.1
628.1
986.1
1524.1
3091.1
1888.1
1692.2
4838.1
2792.1
1901.1
782.1
2437.1
408.1
127.1
3472.1
335.1
2155.1
1884.1
829.1
1756.1
1733.1
4589.1
999.1
1301.1
318.1
2207.1
39.1

nonfunctional   mel-specific   fast-evolving   utORFs   alt-frame

Figure 4 - Figure supplement 2

Figure 5 - Figure supplement 2

# Figure 7 - Figure supplement 2