Original Articles

Unique genomic sequences in a novel *Mycobacterium avium* subsp. *hominissuis*

lineage enable fine scale transmission route tracing during pig movement

Tetsuya Komatsu[a,¶], Kenji Ohya[b,c,d¶*a], Atsushi Ota[e], Yukiko Nishiuchi[f], Hirokazu Yano[g],

Kayoko Matsuo[h], Justice Opare Odoi[c], Shota Suganuma[b*b], Kotaro Sawai[b,i], Akemi

Hasebe[j], Tetsuo Asai[c,d], Tokuma Yanai[b,c,k], Hideto Fukushi[b,c], Takayuki Wada[l], Shiomi

Yoshida[m], Toshihiro Ito[n], Kentaro Arikawa[o], Mikihiko Kawai[p], Manabu Ato[q], Anthony D.

Baughn[r], Tomotada Iwamoto[o], Fumito Maruyama[f,s,t#]

[a]Aichi Prefectural Tobu Livestock Hygiene Service Center, Toyohashi, Aichi, Japan.

[b]Faculty of Applied Biological Sciences, Gifu University, Gifu, Gifu, Japan.

[c]United Graduate School of Veterinary Sciences, Gifu University, Gifu, Gifu, Japan.

[d]Education and Research Center for Food Animal Health, Gifu University (GeFAH), Gifu,

Gifu, Japan.

[e]Data Science Center, Division of Biological Science, Nara Institute of Science and

Technology, Ikoma, Nara, Japan.

[f]Office of Academic Research and Industry-Government Collaboration, Hiroshima

University, Higashi-Hiroshima, Hiroshima, Japan.

[g]Graduate School of Life Sciences, Tohoku University, Sendai, Miyagi, Japan.

[h]Kumamoto Prefectural Aso Public Health Center, Aso, Kumamoto, Japan.

[i]Division of Transboundary Animal Disease Research, National Institute of Animal

Health, National Agriculture Research Organization, Tsukuba, Ibaraki, Japan.

[j]Toyama Prefectural Meat Inspection Center, Imizu, Toyama, Japan.

[k]Hiwa Natural History Museum, Shobara, Hiroshima, Japan.

[l]Graduate School of Human Life Science, Osaka City University, Osaka, Osaka, Japan.

[m]Clinical Research Center, National Hospital Organization Kinki-Chuo Chest Medical

Center, Sakai, Osaka, Japan.

[n]Laboratory of Proteome Research, Proteome Research Center, National Institutes of

Biomedical Innovation, Health and Nutrition, Ibaraki, Osaka, Japan.

[o]Department of Infectious Diseases, Kobe Institute of Health, Kobe, Hyogo, Japan

[p]Graduate School of Human and Environmental Studies, Kyoto University, Kyoto, Kyoto,

Japan

[q]Department of Mycobacteriology, Leprosy Research Center, National Institute of

Infectious Diseases, Higashimurayama, Tokyo, Japan

[r]Department of Microbiology and Immunology, University of Minnesota Medical School,

Minneapolis, Minnesota, USA

[s]Project Research Center for Holobiome and Built Environment (CHOBE), Hiroshima

University, Higashi-Hiroshima, Hiroshima, Japan

[t]Scientific and Technological Bioresource Nucleus, Universidad de La Frontera, Temuco,

Chile.

[*a]Current Address: National Institute of Health Sciences, Kawasaki, Kanagawa, Japan.

[*b]Current Address: Central Research Institute for Feed and Livestock of Zen-noh,

Tsukuba, Ibaraki, Japan.

#Corresponding author

fumito@hiroshima-u.ac.jp (FM)

¶These authors contributed equally to this work.

Running title: TRANSMISSION ROUTE TRACING VIA A NOVEL MAH LINEAGE

Abstract

Mycobacterium avium subsp. *hominissuis* (MAH) is one of the most prevalent

mycobacteria causing non-tuberculous mycobacterial disease in humans and animals.

Of note, MAH is a major cause of mycobacterial granulomatous mesenteric

lymphadenitis outbreaks in pig populations. To determine the precise source of infection

of MAH in a pig farm and to clarify the epidemiological relationship among pig, human

and environmental MAH lineages, we collected 50 MAH isolates from pigs reared in

Japan and determined draft genome sequences of 30 isolates. A variable number of

tandem repeat (VNTR) analysis revealed that most pig MAH isolates in Japan were

closely related to North American, European and Russian human isolates but not to

those from East Asian human. Historical recombination analysis revealed that most pig

isolates could be classified into SC2/4 and SC3, which contain MAH isolated from pig,

European human and environmental isolates. Half of the isolates in SC2/4 had many

recombination events with MAH lineages isolated from humans in East Asia. To our

surprise, four isolates belonged to a new lineage (SC5) in the global MAH population.

Members of SC5 had few footprints of inter-lineage recombination in the genome, and

carried 80 unique genes, most of which were located on lineage specific-genomic

islands. Using unique genetic features, we were able to trace the putative transmission

route via their host pigs. Together, we clarify the possibility of species-specificity of MAH

in addition to local adaptation. Our results highlight two transmission routes of MAH,

one exposure on pig farms from the environment and the other via pig movement.

Moreover, our study also warns that the evolution of MAH in pigs is influenced by MAH

in human living environments, even if the MAH are genetically distinct.

Highlights

● Variable number of tandem repeat (VNTR) analysis of *Mycobacterium avium*

subsp. *hominissuis* (MAH) isolated from pigs (*n*=50) were conducted.

● Draft genome sequences of MAH (*n*=30) and genome analysis were

conducted.

● Pig MAHs were genetically far from East Asian human isolates and close to

those of Western countries.

● Novel MAH lineage which were transmitted farms by pig movement was found.

● Human MAH isolates influenced the evolution of pig isolates.

Key words: MAH, VNTR, Draft genome sequences, pig, transmission route

1. Introduction

Mycobacterium avium complex (MAC) is the most common causative agent of

non-tuberculous mycobacterial disease in humans in Asia and Europe (1, 2). The

frequency of pulmonary MAC infection is increasing worldwide (3). *M. avium* subsp.

hominissuis (MAH) is the most prevalent subspecies of MAC clinical isolates in Asia and

Europe (1, 2).

MAH also causes lymphadenitis most often involving mesenteric lymph nodes in

pigs (4, 5). Pigs infected with MAH usually have no clinical signs and the lesions are

incidentally found during slaughter, bringing great financial losses in pig farm

management in the event of an outbreak (6, 7).

In previous studies, IS*1245* or IS*1311* restriction fragment length polymorphism

(RFLP) analysis revealed that environmental materials in pig farms such as sawdust (8)

and peat (9) are important sources of MAH infection in pigs (10). The pig flow and the

mother pigs are also important sources of MAH infection by RFLP analysis (11).

However, discrimination power of RFLP analysis is inferior to a variable number of

tandem repeat (VNTR) analysis (12).

Recently, several epidemiological studies using VNTR analysis of MAH have been

described from around the world (4, 12, 13, 14, 15, 16, 17, 18, 19). VNTR analysis

shows epidemiological relationships of MAH strains between humans and pigs (4, 16,

17, 18, 19). From these studies, it is suggested that MAH would have region-specific

genetic diversity and distribution. In East Asian countries, human MAH strains isolated

from patients with lung disease were not identical to MAH isolates of pigs (4, 16). On

the contrary, MAH strains of pigs were identical to those of humans from Europe (14,

18) and Japan (19). However, MAH strains of pigs were isolated from limited regions, in

addition to the limitation due to use of only 8 loci for VNTR, which was thought to be

less useful than 16 loci used in other analyses (12). To circumvent these shortcomings,

MAH isolates from additional regions are needed to provide a more robust view of MAH

distribution.

Phylogenetic and population structure analysis based on MAH genomes has

recently revealed that MAH can be divided into six major clades, including

MAHEastAsia1 (EA1), MAHEastAsia2 (EA2) and SC1 - 4, suggesting that MAH

adaptation is geographically restricted around the world (20, 21). For example, EA1 and

EA2 lineages were mainly distributed in humans in Japan and Korea although SC1-SC4

lineages were distributed in Europe and USA (20, 21). Although these studies utilized

many MAH genomes, only two pig isolates were included. Therefore, by using

comparative genome analysis of additional pig MAH isolates, it is possible to unravel

the relationship between human, pig and environment MAH isolates, and decipher

respective transmission routes.

In this study, we aimed to characterize the distribution of MAH lineages in farmed

pigs in Japan and their genetic relationship with human and environmental MAH

isolates for better understanding of its diversity and the way of infection. In addition, we

collected detailed epidemiological information for the determination of transmission

route of MAH for pigs.

2. Materials and methods

2.1. Bacterial isolation and identification. The approach used for isolation and

identification of MAH is described in our previous study (22). 50 isolates used for

VNTR analysis were isolated from lymph nodes, liver, feces, sawdust and farm soil

(Supplementary Table 1). All of them were isolated from pigs or farm environments

in Tokai area (Gifu, Aichi and Shiga Prefecture) and Hokuriku area (Toyama and

Ishikawa Prefecture), whose geographical relationship is depicted in

Supplementary Figure 1.

2.2. Mycobacterial Interspersed Repetitive Unit (MIRU) and *Mycobacterium avium*

Tandem Repeat (MATR) VNTR. Primer sets for 15 loci of MATR (MATR1-9, 11-16)

and 7 loci selected from MIRU loci (32, 292, X3, 25, 7, 10, 47) were used to generate

VNTR profiles, as previously reported (16). The data profiles of 50 MAH isolates in

this study and the number of VNTR profiles were available in Supplementary Table 2

and 3. From all data, we selected 13 MATR (MATR1-8, 11-13, 15-16) and 7 MIRU

profiles for minimum spanning tree (MST) analysis to compare with variable data of

past reports (4, 13, 14, 16, 19, 23, 24) using Bionumerics version 7.0 (Applied Maths,

Sint-Martens-Latem, Belgium). Clonal complex (CC) in MST of 13 loci was defined as

a different group when there were two or more differences in the VNTR profile of

adjacent nodes according to the past study with one modify, that a CC was composed

when it had over five genotypes (13). As the discrimination ability of MST analysis

based on MIRU-VNTR was too low, we were not able to divide the MAH isolates into

several CCs and therefore did not show a corresponding index.

2.3. Draft genome sequence. Draft genome sequences of 30 isolates were obtained

according to our past study (22).

2.4. Genome phylogenetic tree and SNP analysis of *cinA* and *sugA* gene.

Genome phylogenetic tree analysis based on 30 MAH isolates and 20 reference

strains (21) were performed via online tool CSI Phylogeny version 1.4 with default

parameters (25) and IQ-TREE (26) by maximum likelihood (ML) method. *M. avium*

subsp. *paratuberculosis* (MAP) was set as an outgroup. Obtained newick file was

visualized by Molecular Evolutionary Genetics Analysis (MEGA) 7.0. Alignment file

generated by CSI Phylogeny was used for phylogenetic analysis by the neighbor

joining (NJ) method with p-distance method via MEGA 7.0. Topological evaluation

of obtained newick file was conducted by IQ-TREE (26). All of the reference

genome sequences were retrieved from National Center for Biotechnology

Information (NCBI) genome database. The sequence of two epidemiological

marker genes, *cinA* (product name: cytochrome P-450) and *sugA* (product name:

trehalose ABC transporter), which are offered to be able to distinguish major six

MAH lineages (21), were retrieved from draft genome sequences. These

sequences and the reference strains were aligned and used for the phylogenetic

tree analysis by ML method via MEGA 7.0.

2.5. Bayesian analysis of population structure

BAPS and recombination analysis for determination of MAH lineage. MAH

lineage was predicted by using fastBAPS (27) and fastGEAR (28), which was able

to infer the MAH lineages, considering genetic linkage (21, 27). Used genome

sequences were the same as using for followed phylogenetic tree analysis except

for MAP. Filtered polymorphic sites in vcf file generated by CSI Phylogeny were

used for fastBAPS (27). In addition, these sites were combined with intervening

reference genome sequences TH135, and were organized in an alignment file (21)

and then used as an input for fastGEAR (28). Lineages were then generated by

hierarchical clustering using fastGEAR (28), which produced lineage classifications

that were highly consistent with previous reports (20, 21).

2.6. **Pan and core genome analysis.** Draft genome sequences, which were the same

as BAPS and recombination analysis, were re-annotated to obtain gff files by

Prokka v1.14.5 (29). Genes present or absent only in four isolates (GM10, GM12,

GM21, GM32), which we named SC5 as a new lineage below, were searched by

Roary v3.11.2 (30). Basic Local Alignment Search Tool (BLAST) analysis was

performed with the genes present only in SC5 and annotated the genes which had

high coverage and similarity (over 95%) with genes of MAH in NCBI gene

database. We also evaluated whether these genes were located on genomic

islands (GI) and/or phage regions as described below.

2.7. **Detection of mobile elements (Plasmid, phage and GI) in MAH genomes.**

Plasmids were identified from draft genome sequences by PlasmidFinder v 2.1

with the following settings (threshold for minimum % identity: 50%, minimum %

coverage: 20%) (31). Prophage sequences were identified by PHASTER (PHAge

Search Tool Enhanced Release) (32, 33). GIs were predicted by IslandViewer

version 4 (34).

2.8. **Epidemiological information of farms.** Epidemiological information of each farm

about bedding materials, introduction and shipping of pigs was obtained via

interviewing farm owners.

3. **Results**

3.1. **MIRU and MATR VNTR.**

3.1.1. VNTR based on 7 loci. MAH isolates were classified into three groups

(Supplementary Figure 2). Group 1 contained mainly the human isolates, although

almost all pig isolates in Japan were classified in Group 2. Several pig isolates were

classified in Group 1, however, no isolate was clustered with human isolates. The

pig isolates formed a cluster with environmental isolates in Group 2. The eight

isolates of including four SC5 isolates formed Group 3.

3.1.2. VNTR based on 13 loci. Isolates were divided into six CCs (Figure 1). Most

human isolates in Japan and Korea were classified into CC1. CC2 was composed

of MAH isolates of USA, European, Russian and some Korean human and

Japanese pig isolates. CC4 and CC5 were mainly composed of pig isolates from

Japan, although CC3 contained human and pig isolates from Japan. The isolates

same as group 3 in MST of 7 loci were classified into CC6 and were genetically far

from any adjacent node.

3.2. Phylogenetic tree analysis. As a result of topological evaluation of both

phylogenetic trees, the phylogenetic tree of CSI phylogeny (25) was considered to

be more appropriate than that of the IQ-TREE (26). 24 isolates were classified into

3 main lineages (SC2: 13, SC3: 1, SC4: 10), and 4 isolates were classified into a

new lineage (SC5) (Figure 2). SC5 was the earliest branch from the MAP (Figure 2,

Supplementary Figure 3 and 4). GM16 and OCU484 were unclassified and placed

between EA2 and SC3 (Figure 2, Supplementary Figure 3), however, they were

classified in EA2 by the IQ-TREE (Supplementary Figure 4).

3.3. Phylogenetic analysis for *cinA* and *sugA* genes. SC5 shared a unique SNP

pattern in both genes compared with the other lineage (Supplementary Figure 5 and

6). GM5 had a different SNP pattern from SC4 in *cinA* (Supplementary Figure 5).

GM16 had the same SNP patterns as EA1 in *cinA*. GM16 and OCU484 had unique

SNP patterns in *sugA*, respectively (Supplementary Figure 6). Interestingly, SNP in

the *sugA* gene of SC2 were divided into two patterns, SC2a and SC2b. SNP

patterns of SC2a were of a new type and had a characteristic SNP, C25T. The

patterns of SC2b were the same as *M. avium* subsp *avium* (MAA).

3.4. Bayesian analysis of population structure (BAPS) analysis and recombination

analysis for lineage classification. Pig isolates were divided into 4 lineages, SC5,

SC1, SC2 and large SC by fastBAPs (27) (Supplementary Figure 7). The

occurrence of inter-lineage recombination events assessed by fastGEAR (28) is

shown in Figure 3. The strains and isolates were divided into 5 lineages (SC5,

EastAsia, SC2/4, and SC3). Each of the pig isolates was classified into three

lineages (SC3: 1 isolate, SC2/4: 25, SC5: 4). SC5 had few recombination events

with any other lineages. Similarly, some isolates of lineage SC2/4, which were

classified into SC2a by SNP analysis, also had few occurrences of recombination

among lineages.

3.5. Pan and core genome analysis. Core and accessory gene content were 2,314

and 12,115, respectively. Six core genes were absent only in the SC5 and 97 genes

were characteristic in this lineage (Supplementary Table 4). From BLAST analysis,

80 genes of the SC5 had less than 95% coverage and similarity with the other MAH

isolates. Most of them were clustered and were located on GIs, which were not

previously reported (35).

3.6. Detection of mobile elements (Plasmid, phage and GI) in MAH genome. No

plasmids were detected in any isolates. Intact and questionable phage regions were

also not detected, although incomplete sequences of phage were detected in all

isolates. GIs were detected in all isolates, however, there was diversity in the

number of GIs (Table 1).

3.7. Farm relationship where the novel lineage MAH was isolated. The isolates of

SC5 were isolated from pigs reared in three farms located in Gifu, Shiga and Aichi

Prefecture (Supplementary Table 1). From interviewing farm owners, these farms

were related epidemiologically (Figure 4). Aichi B and Gifu A farms had introduced

boars and gilts from Aichi Y farm. Shiga A farm had introduced fattening pigs from

Aichi X farm, whose gilts were introduced from Aichi Y farm. There was nothing in

common about using animal bedding materials, such as sawdust, among farms.

Further, the other farms including in the Hokuriku area had not introduced pigs from

the Tokai area.

4. Discussion

In the present study, we revealed three critical points, the first was the genetic

relationship among MAH isolates of pigs, humans and the environment in Japan, the

second was the identification of a new MAH lineage and its genomic and

epidemiological features, and the third was putative transmission routes of pig MAH.

4.1. Genetic relationship of pig isolates in Japan. From VNTR analysis using both 7

and 13 loci, most pig isolates were grouped with those from pigs and the

environment in Japan and humans in Europe, Russia and USA, supporting previous

studies (4, 13, 14, 16). Nevertheless, pig MAH isolates from various regional origins

are closely genetically related, except for the new lineage. This observation

indicates that genetically close MAHs are widely distributed among Japanese pigs.

In addition, by phylogenetic tree analysis based on draft genome sequences, 28

isolates were classified into SC2, SC3, SC4, SC5, and two isolates were

unclassified. These results are consistent with previous reports (20, 21), but it is

worth noting that SC3 was isolated from pigs in Japan for the first time

(Supplementary Table 5). Recombination analysis clarified two previously

unclassified isolates as belonging to lineage SC2/4. These isolates had many

recombination events with 'EastAsia' and were classified into EA1 and SC4 by SNP

analysis of the *cinA* gene, respectively. Frequent recombination events and multiple

SNP in recombination-cold regions may contribute to the result of these isolates

being unclassified in genome-based phylogenetic tree analysis. These isolates can

be reclassified when several isolates with similar genetic features were found.

On the other hand, SNP analysis reveals there is diversity in pig isolates in

Japan at the level of a single gene. SC2/4 isolates were divided into 5 patterns by

phylogenetic analysis of *sugA*. They had been isolated from multiple farms across

the prefectures. Focusing on SC2a, the isolates had a unique SNP pattern in their

core gene and had few recombination events with the other lineages. This indicates

that SC2a has evolved through accumulation of genetic variations by microevolution

to facilitate to adapt to living pigs, rather than evolve through recombination events

in the environment.

In lineage SC4, most isolates had the same SNP pattern as the reference strain

and had relatively frequent recombination events among lineages, consistent with a

past study (21). SC4 was isolated from human and the environment in Europe, USA

and Russia (21) (Supplementary Table 5) and Japanese pig isolates are genetically

close to those of environmental isolates in these areas (13). According to a recent

annual report from the Animal Quarantine Service in Japan, most live pigs that are

imported to Japan originate from Canada (43.7%), USA (33.4%) and Denmark

(12.9%) in 2019 (https://www.maff.go.jp/aqs/tokei/toukeinen.html). These

observations suggest that SC4 may be transmitted from overseas countries by a pig

movement and has evolved through recombination events with the 'EastAsia'

lineage that is released from humans or human living environments in Japan.

4.2. Genomic and epidemiological features of the novel lineage. Four isolates of

SC5 had the same characteristic SNP patterns in the *cinA* and *sugA* genes, VNTR

profiles and were present in a very unique place in MST trees. These results

including genome based phylogenetic tree, BAPS and recombination analysis

indicated that this cluster was a novel lineage. SC5 isolates had 80 unique genes in

their genomes, most of which were clustered on GIs. In addition, these genes

contained mobile element associated genes, such as IS*481* family transposase

IS*3514* and putative prophage phiRv2 integrase. SC5 had few recombination

events with the other lineages and had characteristic SNPs in their core genes.

These results suggest that SC5 can be partially characterized by mobile elements

rather than recombination events (20, 21) and that they may adapt to the

environment in living pigs via accumulation of genetic variations by microevolution.

These unique genes may serve as the understanding of MAH survival strategies in

living pigs.

It was unknown why SC5 was isolated only from the Tokai area in Japan. In

general, the *Mycobacterium* genus is mainly present in the environment, such as in

water sources, soil and dust (36, 37), however, SC5 is genetically distinct from the

other MAH isolates including environmental, human and pig isolates from the world.

MAH is isolated from wild boars in Europe (38), however, MAH is not isolated from

wild animals in the same area of this study (39). Based on these observations, it is

unlikely that SC5 was introduced from other areas or the environment. The

epidemiological information suggests that SC5 is spreading via pig movement. This

hypothesis is also supported by the observation that there were few inter-lineage

recombination events in SC5, because it is thought that acquisition of genetic

variation through recombination (20) or plasmid transfer (40) occurs more frequently

in the environment than in living hosts. Our results also indicate that SC5 has

species-specificity for pigs, suggesting the possibility of adaptation of MAH for

specific environments including within living hosts and that there are likely additional

lineages that have yet to be identified.

4.3 Putative transmission route of pig MAH. We clarified two transmission route of

pig MAH by recombination and SNP analysis, from environment and via pig

movement (Figure 5). SC2a and SC5 would be transmitted farms via pig movement

because they had few recombination events and characteristic SNP, although SC4

would be invaded into farms from the environment. Our study highlights that the

elimination of subclinically infected pigs with MAH and disinfection of equipment

brought from outside the pigsty could be effective tools against MAH invasion.

5. Conclusion. We have clarified the relationship among MAH isolates of pigs and

humans in Japan and the world using VNTR and genome analysis. Our study

highlights that there are two MAH transmission routes in pigs, via pig movement and

from the environment. We have also identified that SC2a and SC5 can have evolved

in a pig-specific manner. Finally, our study emphasizes that MAH in humans and

human living environments has become a source of transmission to the environment

and is important for the evolution of pig MAH.

Data availability.

The draft genome sequence data used in this study have been submitted to the NCBI

genome database (https://www.ncbi.nlm.nih.gov/genome/) and all of accession number

was available in our past study (22).

Funding

This work was supported by a grant from the Japan Agency for Medical Research and

Development (AMED) (17fk0108116h040, 21fk0108129h0502), the Japan Racing

Association (JRA) Livestock Industry Promotion Project (H28-29_239, H29-30_7) of the

JRA, a grant for Meat and Meat Products (H28-130, H30-60) managed by the Ito

Foundation for Research in design study, collection, analysis; and was supported by

grants from the Japan Society for the Promotion of Science (JSPS) KAKENHI

(JP26304039, JP18K19674, 16H05501, 16H01782, 20H00562) and the Joint Research

Program of the Research Center for Zoonosis Control in Hokkaido University. JOO is a

recipient of a Japanese government (Ministry of Education, Culture, Sports, Science

and Technology) scholarship. All funders had no role in study design, data collection

and interpretation, or the decision to submit the work for publication.

CRediT authorship contribution statement

Tetsuya Komatsu: Methodology, Software, Validation, Formal analysis, Investigation,

Data Curation, Writing-Original Draft. Kenji Ohya: Conceptualization, Methodology,

Validation, Resource, Data Curation, Writing-Review & Editing, Supervision, Project

administration, Funding acquisition. Atsushi Ota: Software, Yukiko Nishiuchi:

Conceptualization, Methodology, Resource, Writing-Review & Editing. Hirokazu Yano:

Software, Writing-Review & Editing, Kayoko Matsuo: Resource, Data Curation,

Investigation. Justice Opare Odoi: Methodology, Software, Formal analysis,

Investigation, Writing-Review & Editing. Shota Suganuma: Methodology, Formal

analysis, Investigation. Kotaro Sawai: Methodology, Formal analysis, Investigation,

Writing-Review & Editing. Akemi Hasebe: Resource, Data Curation, Investigation.

Tetsuo Asai: Writing-Review & Editing, Supervision. Tokuma Yanai: Writing-Review &

Editing, Supervision. Hideto Fukushi: Writing-Review & Editing, Supervision. Takayuki

Wada: Conceptualization, Methodology, Supervision. Shiomi Yoshida:

Conceptualization, Methodology, Supervision. Toshihiro Ito: Software, Data Curation.

Kentaro Arikawa: Conceptualization, Methodology. Mikihiko Kawai: Software, Data

Curation, Writing-Review & Editing. Manabu Ato: Writing-Review & Editing, Supervision,

Project administration. Anthony D. Baughn: Writing-Review & Editing, Supervision,

Project administration. Tomotada Iwamoto: Conceptualization, Methodology, Writing-

Review & Editing, Supervision, Project administration, Fumito Maruyama:

Conceptualization, Methodology, Validation, Data Curation, Writing-Review & Editing,

Supervision, Project administration, Funding acquisition.

Declaration of Competing Interest

All authors reported no potential conflicts of interest.

Acknowledgments

We thank the member of Gifu Prefectural Chuo Livestock Hygiene Service Center for

sampling. We thank the Data Integration and Analysis Facility at the National Institute

for Basic Biology for providing some of the computational resources.

References

1. D.R. Prevots, T.K. Marras, Epidemiology of human pulmonary infection with

nontuberculous mycobacteria: a review. Clin. Chest. Med. 36 (2015) 13-34.

2. H. Namkoong, A. Kurashima, K. Morimoto, Y. Hoshino, N. Hasegawa, M. Ato, S.

Mitarai, Epidemiology of pulmonary nontuberculous mycobacterial disease, Japan.

Emerg. Infect. Dis. 22 (2016) 1116-1117.

3. C.L. Daley, *Mycobacterium avium* Complex Disease. Microbiol. Spectr. 5 (2017).

4. T. Adachi, K. Ichikawa, T. Inagaki, M. Moriyama, T. Nakagawa, K. Ogawa, Y.

Hasegawa, T. Yagi, Molecular typing and genetic characterization of *Mycobacterium*

avium subsp. *hominissuis* isolates from humans and swine in Japan. J. Med.

Microbiol. 65 (2016) 1289-1295.

5. A. Agdestein, T.B. Johansen, Ø. Kolbjørnsen, A. Jørgensen, B. Djønne, I. Olsen, A

comparative study of *Mycobacterium avium* subsp. *avium* and *Mycobacterium avium*

subsp. *hominissuis* in experimentally infected pigs. BMC. Vet. Res. 8 (2012) 11.

6. J. Álvarez , E. Castellanos, B. Romero, A. Aranaz, J. Bezos, S. Rodríguez, A.

Mateos, L. Domínguez, L. de Juan. Epidemiological investigation of a

Mycobacterium avium subsp. *hominissuis* outbreak in swine. Epidemiol. Infect. 139

(2011) 143-148. doi: 10.1017/S0950268810001779.

7. N. Hulinova Stromerova, M. Faldyna. *Mycobacterium avium* complex infection in pigs:

A review. Comp. Immunol. Microbiol. Infect. Dis. 57 (2018) 62-68. doi:

10.1016/j.cimid.2018.06.005.

8. L. Matlova, L. Dvorska, K. Palecek, L. Maurenc, M. Bartos, I. Pavlik, Impact of

sawdust and wood shavings in bedding on pig tuberculous lesions in lymph nodes,

and IS1245 RFLP analysis of *Mycobacterium avium* subsp. *hominissuis* of serotypes

6 and 8 isolated from pigs and environment. Vet. Microbiol. 102 (2004) 227-236.

9. A. Agdestein, T.B. Johansen, V. Polaček, B. Lium, G. Holstad, D. Vidanović, S.

Aleksić-Kovačević, A. Jørgensen, J. Žultauskas, S.F. Nilsen, B. Djønne B,

Investigation of an outbreak of mycobacteriosis in pigs. BMC. Vet. Res. 7 (2011) 63.

10. I.A. Gardner, D.W Hird. Environmental source of mycobacteriosis in a California

swine herd. Can. J. Vet. Res. 53 (1989) 33-37.

11. K. Hibiya, Y. Kazumi, Y. Nishiuchi, I. Sugawara, K. Miyagi, Y. Oda, E. Oda, J. Fujita.

Descriptive analysis of the prevalence and the molecular epidemiology of

Mycobacterium avium complex-infected pigs that were slaughtered on the main

island of Okinawa. Comp. Immunol. Microbiol. Infect. Dis. 33 (2010) 401-421.

12. T. Inagaki, K. Nishimori, T. Yagi, K. Ichikawa, M. Moriyama, T. Nakagawa, T.

Shibayama, K. Uchiya, T. Nikai, K. Ogawa. Comparison of a variable-number

tandem-repeat (VNTR) method for typing *Mycobacterium avium* with mycobacterial

interspersed repetitive-unit-VNTR and IS*1245* restriction fragment length

polymorphism typing. J. Clin. Microbiol. 47 (2009) 2156-2164.

13. K. Arikawa, T. Ichijo, S. Nakajima, Y. Nishiuchi, H. Yano, A. Tamaru, S. Yoshida, F.

Maruyama, A. Ota, M. Nasu, D.A. Starkova, I. Mokrousov, O.V. Narvskaya, T.

Iwamoto. Genetic relatedness of *Mycobacterium avium* subsp. *hominissuis* isolates

from bathrooms of healthy volunteers, rivers, and soils in Japan with human clinical

isolates from different geographical areas. Infect. Genet. Evol. 74 (2019) 103923.

14. K. Ichikawa, J. van Ingen, W.J. Koh, D. Wagner, M. Salfinger, T. Inagaki, K.I. Uchiya,

T. Nakagawa, K. Ogawa, K. Yamada, T. Yagi. Genetic diversity of clinical

Mycobacterium avium subsp. *hominissuis* and *Mycobacterium intracellulare* isolates

causing pulmonary diseases recovered from different geographical regions. Infect.

Genet. Evol. 36 (2015) 250-255.

15. B.R. Imperiale, R.D. Moyano, A.B. DI Giulio, M.A. Romero, M.F. Alvarado Pinedo,

M.P. Santangelo, G.E. ravería, N.S. Morcillo, M.I. Romano. Genetic diversity of

Mycobacterium avium complex strains isolated in Argentina by MIRU-VNTR.

Epidemiol. Infect. 145 (2017) 1382-1391.

16. T. Iwamoto, C. Nakajima, Y. Nishiuchi, T. Kato, S. Yoshida, N. Nakanishi, A. Tamaru,

Y. Tamura, Y. Suzuki, M. Nasu. Genetic diversity of *Mycobacterium avium* subsp.

hominissuis strains isolated from humans, pigs, and human living environment.

Infect. Genet. Evol. 12 (2012) 846-852.

17. A. Kalvisa, C. Tsirogiannis, I. Silamikelis, G. Skenders, L. Broka, A. Zirnitis, I.

Jansone, R. Ranka. MIRU-VNTR genotype diversity and indications of homoplasy in

M. avium strains isolated from humans and slaughter pigs in Latvia. Infect. Genet.

Evol. 43 (2016) 15-21.

18. C. Leão, A. Canto, D. Machado, I.S. Sanches, I. Couto, M. Viveiros, J. Inácio, A.

Botelho. Relatedness of *Mycobacterium avium* subspecies *hominissuis* clinical

isolates of human and porcine origins assessed by MLVA. Vet. Microbiol. 173 (2014)

92-100.

19. M. Subangkit, T. Yamamoto, M. Ishida, A. Nomura, N. Yasiki, P.E. Sudaryatma, Y.

Goto, T. Okabayashi. Genotyping of swine *Mycobacterium avium* subsp.

hominissuis isolates from Kyushu, Japan. J. Vet. Med. Sci. 81 (2019) 1074-1079.

20. H. Yano, T. Iwamoto, Y. Nishiuchi, C. Nakajima, D.A. Starkova, I. Mokrousov, O.

Narvskaya, S. Yoshida, K. Arikawa, N. Nakanishi, K. Osaki, I. Nakagawa, M. Ato, Y.

Suzuki, F. Maruyama. Population structure and local adaptation of MAC lung

disease agent *Mycobacterium avium* subsp. *hominissuis*. Genome. Biol. Evol. 9

(2017) 2403-2417.

21. H. Yano, H. Suzuki, F. Maruyama, T. Iwamoto. The recombination-cold region as an

epidemiological marker of recombinogenic opportunistic pathogen *Mycobacterium*

avium. BMC. Genomics. 20 (2019) 752.

22. T. Komatsu, K. Ohya, A. Ota, Y. Nishiuchi, Y. Hirokazu, K. Matsuo, J.O. Odoi, S.

Suganuma, K. Sawai, A. Hasebe, T. Asai, T. Yanai, H. Fukushi, T. Wada, S.

Yoshida, T. Ito, K. Arikawa, M. Kawai, M. Ato, A.D. Baughn, T. Iwamoto, F.

Maruyama. Genomic features of *Mycobacterium avium* subsp. *hominissuis* isolated

from pigs in Japan. Gigabyte (2021) 1.

23. T. Iwamoto, K. Arikawa, C. Nakajima, N. Nakanishi, N. Nishiuchi, Y. Yoshida, A.

Tamaru, Y. Tamura, Y. Hoshino, H. Yoo, Y.K. Park, H. Saito, Y. Suzuki. Intra-

subspecies sequence variability of the MACPPE12 gene in *Mycobacterium avium*

subsp. *hominissuis*. Infect. Genet. Evol. 21 (2014) 479-483.

24. D.A. Starkova, T.F. Otten, I.V. Mokrousov, A.A. Viazovaia, B.I. Vishnevskiĭ, O.V.

Narvskaia. Genotypic characteristics of *Mycobacterium avium* subsp. *hominissuis*

strains. Genetika. 49 (2013) 1048-1054.

25. R.S. Kaas, P. Leekitcharoenphon, F.M. Aarestrup, O. Lund. Solving the problem of

comparing whole bacterial genomes across different sequencing platforms. PLoS.

One. 11 (2014) e104984.

26. J. Trifinopoulos, L.T. Nguyen, A. von Haeseler, B.Q. Minh. W-IQ-TREE: a fast online

phylogenetic tool for maximum likelihood analysis. Nucleic. Acids. Res. 44 (2016)

W232-W235.

27. G. Tonkin-Hill, J.A. Lees, S.D. Bentley, S.D.W. Frost, J. Corander. Fast hierarchical

Bayesian analysis of population structure. Nucleic. Acids. Res. 47 (2019) 5539-5549.

28. R. Mostowy, N.J. Croucher, C.P. Andam, J. Corander, W.P. Hanage, P. Marttinen.

Efficient inference of recent and ancestral recombination within bacterial populations.

Mol. Biol. Evol. 34 (2017) 1167-1182.

29. T. Seemann. Prokka: rapid prokaryotic genome annotation. Bioinformatics. 30

(2014) 2068-2069.

30. A.J. Page, C.A. Cummins, M. Hunt, V.K. Wong, S. Reuter, M.T. Holden, M. Fookes,

D. Falush, J.A. Keane, J. Parkhill. Roary: rapid large-scale prokaryote pan genome

analysis. Bioinformatics. 31 (2015) 3691-3693.

31. A. Carattoli, E. Zankari, A. García-Fernández, M. Voldby Larsen, O. Lund, L. Villa, F.

Møller Aarestrup, H. Hasman. In silico detection and typing of plasmids using

PlasmidFinder and plasmid multilocus sequence typing. Antimicrob. Agents.

Chemother. 58 (2014) 3895-3903.

32. D. Arndt, J.R. Grant, A. Marcu, T. Sajed, A. Pon, Y. Liang, D.S. Wishart. PHASTER:

a better, faster version of the PHAST phage search tool. Nucleic. Acids. Res. 44

(2016) W16-W21.

33. Y. Zhou, Y. Liang, K.H. Lynch, J.J. Dennis, D.S. Wishart. PHAST: a fast phage

search tool. Nucleic. Acids. Res. 39 (2011) W347-352.

34. C. Bertelli, M.R. Laird, K.P. Williams, B.Y. Lau, G. Hoad, G.L. Winsor, F.S.L.

Brinkman. IslandViewer 4: expanded prediction of genomic islands for larger-scale

datasets. Nucleic. Acids. Res. 45 (2017) W30-W35.

35. A. Sanchini, T. Semmler, L. Mao, N. Kumar, F. Dematheis, K. Tandon, V.

Peddireddy, N. Ahmed, A. Lewin. A hypervariable genomic island identified in

clinical and environmental *Mycobacterium avium* subsp. *hominissuis* isolates from

Germany. Int. J. Med. Microbiol. 306 (2016) 495-503.

36. J. Makovcova, M. Slany, V. Babak, I. Slana, P. Kralik. The water environment as a

source of potentially pathogenic mycobacteria. J. Water. Health. 12 (2014) 254-263.

37. A. Lahiri, J. Kneisel, I. Kloster, E. Kamal, A. Lewin. Abundance of *Mycobacterium*

avium ssp. *hominissuis* in soil and dust in Germany - implications for the infection

route. Lett. Appl. Microbiol. 59 (2014) 65-70.

38. G. Ghielmetti, M. Hilbe, U. Friedel, C. Menegatti, L. Bacciarini, R. Stephan, G.

Bloemberg. Mycobacterial infections in wild boars (*Sus scrofa*) from Southern

Switzerland: diagnostic improvements, epidemiological situation and zoonotic

potential. Transbound. Emerg. Dis. 68 (2021) 573-586.

39. J.O. Odoi, K. Ohya, J. Moribe, Y. Takashima, K. Sawai, K. Taguchi, H. Fukushi, T.

Wada, S. Yoshida, T. Asai. Isolation and antimicrobial susceptibilies of

nontuberculous Mycobacteria from wildlife in Japan. J. Wildl. Dis. 56 (2020) 851-862.

40. S.A. Shoulah, A.M. Oschmann, A. Selim, T. Semmler, C. Schwarz, E. Kamal, F.

Hamouda, E. Galila, W. Bitter, A. Lewin. Environmental *Mycobacterium avium* subsp.

hominissuis have a higher probability to act as a recipient in conjugation than clinical

strains. Plasmid. 95 (2018) 28-35.

Supporting Information Legends

Supplementary Table 1. Information of MAH isolates in this study.

Supplementary Table 2. VNTR profiles of 50 MAH isolates in this study.

Supplementary Table 3. Information of MAH strains or isolates used for VNTR

analysis in this study.

Supplementary Table 4. Uniquely present or absent genes in new lineage (SC5).

Supplementary Table 5. Geographic and host features of MAH lineages.

Supplementary Figure 1. Location of each regions and prefectures in Japan. Pigs

from Tokai area (southern Gifu, Shiga and Aichi Prefecture) are slaughtered in Gifu

meat inspection center. Pigs from Hokuriku area (northern Gifu, Toyama and Ishikawa

Prefecture) are slaughtered in Toyama meat inspection center.

Supplementary Figure 2. Minimum spanning tree (MST) based on 7 loci MIRU-

VNTR genotyping of MAH isolates in Japan. Circles indicate different VNTR profiles.

The size of each circle depended on the number of isolates sharing the same profiles.

New lineages indicated by an arrow were clustered and consisted of isolates GM10,

GM11, GM12, GM21, GM23, GM24, GM25, GM32.

Supplementary Figure 3. Phylogenetic tree based on draft genome sequences

generated by CSIphylogeny. Maximum likelihood tree was generated by CSI

phylogeny version 1.4 and was visualized via MEGA 7.0. The area and Prefecture of

isolates in this study were indicated by the color dots. Lineage classification except for

new lineage was based on Yano et al., 2019. BMC Genomics.

Supplementary Figure 4. Phylogenetic tree based on draft genome sequences

generated by IQ-TREE. Maximum likelihood tree was generated by IQ-TREE and

visualized via MEGA 7.0. The Area and Prefecture of isolates in this study were

indicated by the color dots. Lineage classification except for new lineage was based on

Yano et al., 2019. BMC Genomics.

Supplementary Figure 5. Allelic variants in *cinA* gene of pig MAH. A) Maximum

likelihood tree generated by MEGA 7.0. Bootstrap values were calculated by 1,000

replications. The scale bar indicates genetic distance of each strain. Lineage

classification except for new lineage was based on Yano et al (25). B) SNP cites in pig

MAH. New lineage shared the same SNP patterns, which were highlighted by gray.

Some variations were detected in SC4. Reference SNP patterns were referred from

Yano et al (25).

Supplementary Figure 6. Allelic variants in *sugA* gene of pig MAH. A) Maximum

likelihood tree generated by MEGA 7.0. Bootstrap values were calculated by 1,000

replications. The scale bar indicates genetic distance of each strain. Lineage

classification except for new lineage was based on Yano et al (25). B) SNP cites in pig

MAH. New lineage shared the same SNP patterns, which were highlighted by gray. Two

type of SNP patterns were detected in SC2. Reference SNP patterns were referred from

Yano et al (25).

Supplementary Figure 7. Lineage classification by fastBAPS. Lineage classification

was predicted by fastBAPS (Tonkin-Hil et al., 2019 Nucleic Acids Res). Bootstrap

values were calculated by 1,000 replications. The scale bar indicates genetic distance

of each strain. Lineage classification in the right side except for new lineage was based

on Yano et al., 2019. BMC Genomics. Neighbor joining tree was generated by via

MEGA 7.0 using alignment file obtained from CSI phylogeny version 1.4.

585   **Figure Legends**

Figure 1. Minimum spanning tree (MST) based on 13 loci MATR-VNTR genotyping of MAH isolates from various geographic regions and sources. Circles indicate different VNTR profiles. The size of each circle depended on the number of isolates sharing the same profiles. Clonal complexes were highlighted in gray. Each clonal complex consisted of over five isolates.

Figure 2. Phylogenetic tree based on draft genome sequences. Neighbor joining tree was generated by MEGA 7.0 using alignment file obtained from CSI phylogeny version 1.4. Bootstrap values were calculated by 1,000 replications. The scale bar indicates genetic distance of each strain. Lineage classification except for new lineage was based on Yano et al (21).

Figure 3. Determination of MAH lineages. Left: Visualization of recent horizontal acquisition events generated by fastGEAR. Color of chromosomal regions expresses donor lineage of the regions. Right: Neighbor joining tree was generated by via MEGA 7.0 using alignment file obtained from CSI phylogeny version 1.4.

Figure 4. Relationship among farms where new lineage MAH was isolated. Aichi Y farm bred and supplied pigs to Aichi X, Aichi B and Gifu A farm, where these farms had the novel lineage (GM21, GM10 and GM32). Aichi X farm whose gilts were introduced from Aichi Y farm supplied pigs to Shiga A farm where GM12 was isolated. Finally, pigs of Gifu A, Aichi B and Shiga A farm were slaughtered in the same slaughtered plant (Gifu meat inspection center).

Figure 5. The differences in the way of MAH infection in pigs by lineages. SC2 and SC5 exist in live pigs and would evolve with the accumulation of genetic variations in

608   their genes by microevolution. They move from farms to farms via live pigs. SC3 and

609   SC4 evolve through numerous recombination events in the environment with the other

610   lineages released from various hosts. MAH that have completed recombination would

611   be supplied to pig farms from the environment.

Complex 6
New lineage

Complex 3

Complex 2

Complex 1

Complex 4

Complex 5

Area, origin and No. of isolates

- ■ Japan Gifu pigs (22)
- ■ Japan Toyama pigs (13)
- ■ Japan Ishikawa pigs (8)
- ■ Japan Kinki pigs (70)
- ■ Japan Okinawa pigs (19)
- ■ Japan Chubu human (191)
- ■ Japan Kinki human (93)

- ■ Japan Hokkaido human (31)
- ■ Japan bathroom human (60)
- □ Japan environment (29)
- ■ Korea human (175)
- ■ Russia human (90)
- ■ Europe human (37)
- ■ USA human (32)