CryoEM structures of the multimeric secreted NS1, a major factor for dengue hemorrhagic fever

Authors: Bo Shu, Justin S.G. Ooi, Aaron W.K Tan, Thiam-Seng Ng, Wanwisa Dejnirattisai, Juthathip Mongkolsapaya, Guntur Fibriansah, Jian Shi, Victor A. Kostyuchenko, Gavin Screaton, Shee-Mei Lok*

*Correspondence: Shee-Mei Lok (sheemei.lok@duke-nus.edu.sg)

Abstract

Dengue virus infection can cause dengue hemorrhagic fever (DHF). Dengue NS1 is

multifunctional: the intracellular dimeric NS1 (iNS1) forms part of the viral replication

complex, the extracellular multi-oligomeric secreted NS1 (sNS1) is a major factor

contributing to DHF. The structure of the iNS1 is well studied but not sNS1. Here we show

the tetrameric (stable and loose conformation) and hexameric structures of sNS1. Stability of

the stable and loose tetramers is determined by the conformation of their N-terminal domain

– elongated β-sheet or β-roll. Binding of an anti-NS1 Fab breaks the loose tetrameric and

hexameric sNS1 into dimers, whereas the stable tetramer remains largely unbound. Our

results show detailed quaternary organization of different oligomeric states of sNS1 and will

contribute towards the design of dengue therapeutics.

Main

Dengue is a positive sense single-stranded RNA enveloped virus. It belongs to the family *Flaviviridae*, which also includes significant human pathogens like West Nile (WNV), yellow fever (YF) and Zika virus (ZIKV). Dengue virus causes disease that ranges from mild dengue fever to the severe dengue hemorrhagic fever (DHF) and the dengue shock syndrome (DSS)[1]. There are four DENV serotypes. Dengvaxia, which is currently the only licensed dengue vaccine, has an overall efficacy of only 45%, and there is a possibility of priming the previously immune naïve children to develop severe disease in subsequent natural infection after vaccination [2]. It has proven to be difficult to develop an effective vaccine using whole dengue virus or its surface E protein, as the vaccine would need to stimulate equally strong immune responses simultaneously against all four dengue serotypes or else it can lead to a possibility of antibody-dependent enhancement (ADE) of infection[3]. This could lead to severe DHF/DSS.

Recently, dengue non-structural protein 1 (NS1) has been investigated as an alternative protein candidate for vaccine development. It has been shown to play an important role in causing the vascular leakage symptoms in DHF/DSS[4]. NS1 is secreted alongside dengue virus particles during an infection. It is a multifunctional protein which exists in different oligomeric forms intracellularly and extracellularly[5], and they play distinctly different roles. The intracellular NS1 (iNS1), exists as membrane associated dimeric form and is a part of the viral replication complex. The extracellular secreted NS1 (sNS1), exists in higher oligomeric forms. sNS1 likely causes vascular leakage symptoms in DHF/DSS via two pathways – indirectly, by stimulating the innate immune response through activation of complement and Toll-like receptors[6], and directly, by binding to endothelial cells causing endothelial hyperpermeability. sNS1 when bound directly to endothelial cells causes barrier dysfunction of the cell monolayer as demonstrated in an *in vitro* trans-

endothelial electrical resistance (TEER) assay[4,7]. *In vivo* studies show that the sNS1 is present

at very high concentrations in patient sera and correlates closely with the severity of disease [8].

Mice injected with sNS1 along with a sublethal dose of DENV were observed to succumb to

infection. Injection of sNS1 alone also causes vascular leakage in mice[4]. NS1-immune

polyclonal serum and anti-NS1 monoclonal antibodies (MAbs) can protect mice against

lethal DENV2 challenge [4]. sNS1 was also found to enhance viral infection in mosquitoes by

downregulating mosquito midgut immune genes[9]. sNS1 is therefore an important viral

protein both for viral infection (mammalian cells and mosquitoes) and pathogenesis, and it is

important to understand the structure of the sNS1 protein.

iNS1 (45 kDa) is first made in the endoplasmic reticulum (ER) as a monomer. It then

forms dimers and is glycosylated in the ER and the trans-Golgi network. It is eventually

transported to the cell membrane where some is secreted outside the cell, becoming sNS1.

There is a high-resolution crystal structure of full length dimeric dengue iNS1[10] produced by

baculovirus expression and subsequent extraction from cellular membrane compartments .

There are also crystal structures of the dimeric form of NS1 protein from other flaviviruses[11]

and they are largely similar to that of the DENV iNS1. Each protomer of the iNS1 contains

three domains (Extended data Fig. 1A): 1) the N-terminus contains the β-roll domain (amino

acids 1-30) that interacts with the same domain from the other protomer within the dimer to

form a roll-like structure (named β-roll). This domain is generally hydrophobic and is thought

to interact with cellular membranes. 2) Following the β-roll domain is the wing domain

(amino acids 31-181). The wing domain is the most flexible domain as it differs between

different flavivirus NS1 crystal structures [11]. 3) the β-ladder domain (amino acids 182-352),

consists of a continuous arrangement of 10 β-strands. The β-ladder domains from the two

iNS1 protomers line up with each other to form a long ladder-like pattern. One side of the β-

ladder faces the hydrophobic β-roll, while the other side is decorated with its highly charged

spaghetti loops. The interaction between protomers within the dimer is stabilized by a large

number of hydrogen bonds[12].

About a decade ago, two very low resolution (23 Å and 30 Å) EM structures of sNS1

were determined: a cryoEM structure reconstructed from images acquired by using an 120kV

LaB$_6$-equipped electron microscope (Philip CM12 TEM)[13] and the other, a negative stained

room temperature EM structure (Extended data Fig. 1B)[14]. Both maps had an overall hollow

barrel-like structure lacking fine structural details. Since then, the availability of 300kV

transmission electron microscopes equipped with direct electron detectors and field emission

gun electron sources, as well as improved cryoEM reconstruction methods, have made more

routine determination of cryoEM structures of small proteins to high resolution possible.

Here we report the cryoEM structures of sNS1 tetramers (dimer of dimers) and hexamers

(trimer of dimers) from a DENV2 clinical strain PVP94/07. We observe sNS1 exists

predominantly in the tetrameric form. There are two tetrameric states - stable and loose, and

some of these structures are determined to ~3.5Å resolution. The most obvious and dramatic

difference between the stable and loose tetramers is the organization of their N-terminal

domains – elongated β-sheet and β-roll, respectively. Only a minority of the sNS1 population

is hexameric and this cryoEM structure was determined to 8Å resolution. We also determined

a structure of sNS1 complexed with a Fab 5E3 to 3.5Å resolution. Most of the sNS1 (loose

tetramers and hexamers) dissociates into dimers when bound to Fab, while the stable tetramer

remains largely unbound. This is consistent with our results showing the ability of the Fab

5E3 to only partially inhibit sNS1-induced endothelial cell permeability. This study shows

sNS1 exists in a heterogenous mixture of oligomerization states, and we present the first high

resolution structures of sNS1.

sNS1 from DENV2 contains heterogenous oligomeric states with tetramers as the predominant population.

His-tagged NS1 (DENV2 clinical strain PVP94/07) was expressed in human embryonic kidney (HEK) cells. We harvested the sNS1 from tissue culture supernatant and after purification, we analyzed it using coomassie blue stained semi-denaturing SDS-PAGE gel (the sample was not boiled nor exposed to DTT). The gel shows clear bands corresponding to dimers and tetramers (Fig. 1A). As the sample was exposed to SDS, a detergent, the presence of tetramer band suggests that the tetramer is resistant to detergent. We then performed cross-linking of sNS1 with bis(sulfosuccinimidyl) suberate (BS^3) before analyzing it by denaturing SDS-PAGE. This showed the presence of aggregates, which could be due to BS^3 non-specifically crosslinking proteins, as well as bands for hexamer, tetramer, dimer and monomer (Fig. 1B). These results also showed that the tetrameric form is the predominant oligomerization state in the NS1 protein sample.

We collected 7714 cryoEM micrographs (Table S1) and observed particles with distinct shapes (Fig. 1C). We then performed several cycles of 2D classification and alignment of the boxed particles. The 2D class averages show particles with different features (Fig. 1D) - some appearing like stable and loose tetramers and others like hexamers. We did not observe any sNS1 dimers. This suggests that sNS1 consists of heterogenous populations of particles with likely tetrameric and hexameric states.

Fig. 1 Profile of the sNS1 from DENV2 (PVP94/07) clinical strain. **(A)** Semi-denaturing SDS-PAGE gel (non-boiling and no DTT) shows the clear dimer and tetramer bands, along with a very faint hexamer band. **(B)** Western blot denaturing SDS-PAGE gel (using Anti-polyHistidine−Peroxidase antibody) analysis of sNS1 cross-linked with BS^3 at different incubation times. Aggregates, hexamers, tetramers, dimers and monomers were seen. **(C)** A cryoEM micrograph of the sNS1 showing distinct-shaped particles, e.g., butterfly-shaped (red circles) and cube-shaped particles (purple circles). Scale bar is 10 nm. **(D)** The 2D class averages indicate that the particles are heterogenous, with some likely existing as tetramers (stable and loose forms) and others as hexamers. **(E)** The distribution of particles in different oligomerization states determined after 3D classification and refinement. Most of the particles are tetramers (loose and stable) and hexamers are a minority in the population.

CryoEM reconstruction of different oligomeric states of sNS1

As sNS1 consists of a heterogenous population of particles of different oligomerization states, we performed a 3D classification using the program Relion [15] (Extended data Fig. 2). The first round of unsupervised classification was done assuming no symmetry (C1) to avoid bias towards any specific oligomerization states. We then performed rough fitting with a dimeric NS1 crystal structure into these maps to determine their oligomerization states (Extended data Fig. 3). We observed two maps that correspond to tetramers: one with clear densities suggesting that the overall structure is quite stable (Extended data Figs. 2-3), while the other with looser interactions between the two dimers and poorer map resolution (Extended data Fig. 2). We refer the former as stable tetramer and the latter, as loose tetramer. We also observed a 3D class that is hexameric in structure (Extended data Figs. 2-3). Consistent with the 2D class averages, the 3D classification also did not yield any dimeric structures. We then used the maps from 3D classification in Relion as initial models for further classification and refinement in the program CryoSPARC [16]. For the stable tetramers, we performed multiple cycles of heterogenous and non-uniform refinement [17] by imposing D2 symmetry and obtained a final map of 3.5Å resolution (Fig. 2A and Extended data Fig. 4), as measured by gold standard FSC curve cutoff at 0.143. For the loose tetramers, we performed heterogenous and non-uniform refinement with no symmetry imposed, and we determined one subclass of the loose tetramer to 8 Å resolution (Fig. 3A(i) and Extended data Fig. 5A). We then performed local refinement on one of its dimers and obtained a 3.4 Å cryoEM density map (Fig. 3A(ii) and Extended data Fig. 5B). For the hexamers, since we only had a limited number of particles (~30,000 particles), we performed only non-uniform refinement with D3 symmetry imposed and obtained an 8Å map (Figs. 3B and Extended data Fig.6). The cryoEM reconstruction flow chart is summarized in Extended data Fig. 2 and validation statistics in Extended data Table 1.

Structure of sNS1 stable tetramers

The 3.5Å resolution sNS1 stable tetramer cryoEM map shows clear separation between β-strands (Extended data Figs. 4C-D). We also calculated the local resolution (Extended data Fig. 4A) of the map. While most of the density map is ~3.5 Å resolution, parts of the map (corresponding to the β-ladder) are at ~3.0Å resolution with some of the side chain densities resolved (Extended data Fig. 4D). The resolution of the wing domain and the β-roll was lower (4.0 to 4.5 Å). The density of both N- (residues 1-11) and C- terminal ends (residues 341-352) is missing from the map.

We interpreted the stable tetramer (dimer of dimers) cryo-EM density by fitting in the zika dimeric NS1 crystal structure (PDB:5GS6), because the structure is more complete than the dengue iNS1, and then mutated it into dengue NS1 sequence (Figs. 2A-B and Extended data Fig.4B). After fitting, we compared one of our sNS1 dimer in the stable tetramer to the dengue iNS1 dimer (PDB:4O6B) and it shows the wing domain of our sNS1 dimer is rotated by 6.8° with respect to the β-ladder. The β-roll structures of the two dengue iNS1 dimers cannot fit well into the cryoEM density of the stable tetramer and they clash with each other. Indeed, the space (25Å) between two dimers within the stable tetramers is not big enough to accommodate the two iNS1 β-rolls. We observed that the β-roll densities, instead of concentrating at the middle of the dimers as observed in iNS1 (Extended data Fig. 1), have an elongated shape that stretches across the two protomers within a dimer (Figs. 2A and B). This suggests that they have very different tertiary structures. Fitting of the density map shows that residues 15 to 24 form a long β-strand which pairs with the same β-strand of the other protomer within the dimer to form a long anti-parallel β-sheet (hereafter, referred to as elongated β-sheet) (Figs 2A-B). As the resolution of this region is ~4.5 Å, we analyzed their interaction by using a distance cut off of 8 Å between the Cα backbone. Using this criterion, we observed that this elongated β-sheet from one dimer likely interacts with the same β-sheet

from the opposite dimer via electrostatic interactions or hydrogen bonds at both ends of each

respective β-sheet (Fig. 2C(i)-(ii)). Charge analysis in UCSF Chimera [18] shows that these

two ends have high charge complementarity between dimers (Fig. 2C (iii)), suggesting

possible hydrogen bonds or electrostatic interactions. Hydrophobic analysis shows that these

β-sheets have an overall highly hydrophobic surface that also plays an important role in

stabilizing their interactions.

We next compared the β-roll domain of iNS1 with the elongated β-sheet of the sNS1

stable tetramer. The iNS1 β-roll domain contains three β-strands (residues 1-7, 12-16 and 18-

22), whereas we were unable to observe any density for residues 1-11 in the sNS1 stable

tetramer, probably due to disorder in this region, and residues 15-24 form a long, continuous

β-strand. This raises a question: is the sNS1 tetramer assembled from dimers of iNS1? If so,

is it possible for the iNS1 β-roll structure to rearrange into the elongated β-sheet seen in the

stable tetramer? If this were to happen, it would require the second β-strand of iNS1 to rotate

by ~120° with respect to the third β-strand to form the long, continuous β-strand we observed

(Fig. 2D). Whether the iNS1 dimer is a precursor of the sNS1 stable tetramer is unknown. It

is also possible that the sNS1 stable tetramer is assembled during translation of the NS1

protein by a pathway independent of iNS1 dimer formation, before being secreted.

Stable tetramer

A

B Elongated β sheet density

Elongated β sheet

90°

C (i) **(ii)** **(iii)** Charge distribution   Hydrophobicity

N   N

K11   29T 30E   24N   15C
28W   17S
22T   18G
21I   19I
20F   20F
K11   19I   21I
30E   29T   18G   22T
28W   17S   24N
15C

Top view   Side view   Side view

D

Dengue iNS1 dimer   ?   sNS1 stable tetramer

Fig. 2 Structure of the stable sNS1 tetramer. (**A)** Two NS1 dimeric structures fitted into the cryoEM map (reduced sharpening was applied to make the density of the elongated β-sheet clearer). One dimer is colored in pink and light green and the other in yellow and light blue. (**B)** Left: the overall structure of the sNS1 dimer is similar to that of iNS1 except for the β-roll region (residues 1-30) in iNS1, which is an elongated β-sheet (residues 15-24) in sNS1. View from the elongated β-sheet of the black boxed region in (A). The respective elongated β-strand of each protomer is highlighted in a darker shade of the same color (red and dark green). Right: surface density map with the regions corresponding to the elongated β-sheet colored in green, and the rest of the map in yellow. (**C)** Interaction of the elongated β-strands between the two dimers (red/green and yellow/blue). (i) Top view showing the elongated β-sheets of the red/green dimer interacting with that of the yellow/blue dimer (only its elongated β-sheet is shown). (ii) Side view showing possible hydrogen bonds or electrostatic interactions between the two elongated β-sheets from opposite dimers (dotted black lines).

These interactions are identified by the distance between Cα backbone (< 8Å) and also their charge characteristics. (iii) The interacting residues are complementary in both charges and hydrophobicity. Open book representation of (ii): (Left) charge, and (Right) hydrophobicity distributions on the surface of the two elongated β-sheets. For charge, positive charges are colored in blue while negative charges are red. For hydrophobicity, cyan color indicates hydrophobic residues. Possible interacting residues are labelled. **(D)** Comparison of iNS1 β-roll with sNS1 stable tetramer elongated β sheet and how it might change from the β-roll structure (dotted arrow on the iNS1 structure) into the elongated β sheet structure.

CryoEM structure of loose sNS1 tetramers

We also determined an 8Å resolution loose sNS1 tetramer structure (Figs. 3A(i) and Extended data Fig.5A) and by focused refinement on one of its constituent dimers, we obtained a 3.4Å resolution cryoEM map (Figs. 3A (ii) and Extended data Fig.5B (i-ii)). The overall structure of the dimer and its β-roll (Fig. 3A (ii)) is very similar to that of iNS1 (RMSD 0.8 Å), unlike the stable sNS1 tetramer. The overall 8Å resolution loose sNS1 tetramer structure shows that the two dimers are rotated by ~35° with respect to each other and extra density is visible between the β-rolls of these two dimers (Fig. 3A(i)). Our high resolution sNS1 dimer structure within the loose tetramer also shows some strong extra densities above the β-roll (Fig. 3A(iii)). Previous work from other research groups [13,14] show that there are lipids associated with sNS1 that may help stabilize the overall structure, and which may contribute to these uninterpreted densities.

The difference between the stable and the loose tetramers is whether the residues near the N-terminus form a β-roll or an elongated β-sheet. Hence, this could be the determinant that defines stability.

An existing crystal structure of a ZIKV sNS1 [11] (PDB code: 5GS6), obtained by purifying sNS1 produced using a baculovirus expression system, is a dimeric structure. However, upon examination of the crystal packing in this structure, sNS1 appears to have a tetrameric structure similar to our loose tetramer (RMSD 4.2 Å) (Fig. 3A(iv)). This may suggest that at least one of the ZIKV sNS1 oligomers might be the loose tetramer structure.

A **sNS1 loose tetramer**

(i) 8Å resolution loose tetramer

(ii) 3.4 Å resolution structure by focused refinement of one dimer within the loose tetramer

Lipids?

β roll density

Crystal packing ZIKV sNS1

Loose tetramer

(iii) Lipids?

Zoom-in side view of the β roll

(iv)

B **sNS1 hexamer**

(i) 8Å resolution sNS1 Hexamer

Lipids?

Top view   Side view

(ii) Dengue iNS1 hexamer due to crystal packing(PDB:4O6B)

Top view   Side view

C

Loose tetramer   +   Dimer   ?   sNS1 hexamer

Fig. 3 CryoEM structures of the sNS1 loose tetramer and hexamer. **(A)** (i) The fit of sNS1 into the 8 Å resolution loose tetramer density map. (ii) Focused refinement of one dimer of the loose tetramer yields a 3.4Å resolution map. An iNS1 dimer fits well into the density showing that the loose tetramer contains a β-roll structure similar to that of iNS1. Left: cryoEM map with the density corresponding to the β-roll colored in green and the rest in

yellow. Right: the fit of the red/green dimer into the density. (iii) Zoom-in side view of the β-roll density (from (ii)) showing uninterpreted density that might belong to lipid molecules. (iv) Superposition of our loose tetramer with the tetramer observed in the crystal packing of ZIKV sNS1 structure (PDB: 5GS6) shows that they have similar structures. **(B)** Comparison of our sNS1 hexamer with the dengue iNS1 hexamer from the crystal packing. (i) Top and side views of the 8Å resolution sNS1 hexamer structure. Left: fit of three dimers into the cryoEM map. We are unable to discern whether sNS1 adopts a β-roll or an elongated β-sheet structure, as the interior of the hexamer contains unfeatured density. It may also contain lipid cargo as suggested by Gutsche and colleagues [14] (ii) The previously published dengue iNS1 structure [10] shows hexamers due to crystal packing; however, there is little interaction between the iNS1 dimers (left; top view). The dimers in the hexamer (right; side view) are also oriented differently to our hexameric structure (in (i), side view). **(C)** Interaction of a loose tetramer with a dimer could lead to the formation of a hexamer.

CryoEM structure of the sNS1 hexamer

We obtained an 8Å resolution sNS1 hexamer map (Figs. 3B(i) and S6A-B). The low resolution structure suggests that the NS1 molecules in the hexameric state are interacting loosely with each other. Fitting of iNS1 (Extended data Fig. 6C) suggests that three dimers interact with each other via their wing domain and the distal end of the β-ladder. This interaction is similar to that in the loose tetramer. The density in the central core of the hexamer is weak (Extended data Fig. 6B) and we were therefore unable to determine if the dimers have a β-roll or elongated β-sheet conformation. It is also possible that the interior of this hexamer contains lipid cargo, as suggested by other research groups [13,14] and this may prevent determination of the structure of this region.

The crystal structure of dengue dimeric iNS1 [10] shows that the crystal packing contains a trimer of dimers (i.e., hexamer) organization; however, there are very few interactions between the three dimers (Fig. 3B(ii)). Their positions in the crystal lattice are mainly supported by their interactions with other iNS1 molecules inside the crystal unit cell. This suggests that the hexameric packing of iNS1 may not be physiological. Comparison of our cryoEM sNS1 hexamer to this crystal hexamer packing shows a very different organization of dimers –they are rotated by ~90° relative to each other (Fig. 3B(i) and (ii), right).

It is possible that the loose tetrameric structure is a transition intermediate of the hexameric structure (Fig. 3C). However, we observed loose tetramers forming ~74% of the particle population (Fig. 1D), while hexamers only form ~3%, suggesting that the loose tetramer may be a preferred or a lower energy oligomerization state.

It is possible that both the loose tetrameric and the hexameric structures are stabilized by lipids. To test this, we incubated the NS1 sample with detergent (0.05% DDM) and imaged it by cryoEM SPA. We observed that the percentage of loose tetramers and hexamers

276 decreased dramatically after detergent treatment (Extended data Table 2), compared to

277 untreated NS1 (Fig. 1E). This suggests that the lipids inside the loose tetramers and hexamers

278 could be important for their structural integrity. A caveat to this conclusion is the detergent

279 could also disrupt the hydrophobic interactions between the dimers.

280 CryoEM structure of sNS1 complexed with Fab 5E3

We also investigated the cryoEM structure of sNS1 complexed with the Fab fragment

of an anti-NS1 antibody 5E3. The 2D class averages (Fig. 4A) of the boxed particles from the

cryoEM micrographs suggest that, when bound by Fabs, most sNS1 particles become dimeric.

Unbound stable tetramers were also present (Fig. 4A) and formed ~18% of the total particles.

It suggests that the Fab 5E3 can readily dissociate the sNS1 loose tetrameric and hexameric

structures. The stable tetrameric structure, on the other hand, is more resistant to Fab binding.

The structure of the Fab 5E3:sNS1 dimer complex was determined to an overall

resolution of 3.5Å (Fig. 4B, and Extended data 7A), as measured by gold standard FSC curve

cutoff at 0.143 (Extended data Fig. 7B). The sNS1 dimer:Fab complex has an overall arch

shape, where the Fabs bind to both ends of the β-ladders of the sNS1 dimer (Fig. 4B and

Extended data Fig.7A) and are tilted towards the β-roll side of the sNS1.

Fab 5E3 interacts with the end of the β-ladder of NS1 (Fig. 4B, Extended data Figs. 7-

8) via mainly the Fab heavy chain (Extended data Table 3). The 5E3 epitope is similar to two

other previously published Fab:NS1 dimer complex [19,20] – Fabs 2B7 and 1G5.3 (Extended

data Fig. 8). Comparison of these structures shows that these three Fabs bind at different

angles to the NS1 β-ladder. It has been postulated previously[10] that the hydrophobic β-roll

may bind directly to cellular membranes (Extended data Fig. 1), which could be important for

its pathogenesis, and therefore the binding orientation of Fab 5E3 may hinder this interaction.

A similar neutralization mechanism was suggested for Fab 2B7 [20].

We observed that these NS1 dimers have β-roll structures (Figs. 4B and Extended data Fig.7C). This suggests that they are likely not derived from the stable tetramer which has an elongated β-sheet structure. Hence, they could be a broken down product of Fab binding to sNS1 loose tetramers (which contain β-roll structures) and perhaps also the hexamers. We superimposed the Fab:dimer complexes into the loose tetramer and hexamer, and they showed Fabs when bound, could clash with the neighbouring NS1 dimer (Extended data Fig. 9A-B). Analysis of the solvent accessibility of the epitopes on the loose tetramer (Extended data Fig. 9C) shows two of the epitopes (orange spheres) are partially exposed (37.5%), while the other two (cyan spheres), are fully exposed. All epitopes in the hexamer (Extended data Fig. 9D) are partially exposed (37.5%). Since Fab likely could bind to the loose tetramer and hexamer. This suggests that the loose tetramers and hexamers likely undergo motions that will at some time point, have their epitopes fully exposed for Fab binding, and once the Fab has bound, the loose tetramer and hexamers could be dismantled into dimers.

We reconstructed the unbound stable tetramer to 3.9Å resolution (Fig. 4C (i)), it is identical to the uncomplexed stable tetramer (Fig. 2). Analysis of the epitopes on all NS1 molecules in the unbound stable tetramer shows they are only partially exposed (37.5%) (Extended data Fig. 9E), thus discouraging Fab binding. Superposition of the Fab 5E3:dimer complex with one of the dimers in the stable tetramer structure (Fig. 4C (ii)) also shows that the Fab if bound, will clash with the wing domain of the neighbouring dimer. While the Fab is able to cause dissociation between dimers in the loose tetramer and hexamer, it is unable to do the same to the stable tetramer. This suggests that the two dimers in the stable tetramer have tighter interactions, thus disallowing antibody binding. The presence of unbound stable tetramer is consistent with the results of the TEER assay (Fig. 4D), showing Fab can only partially inhibit sNS1 induced endothelial cell hyperpermeability.

324   In conclusion, since the loose tetramer and hexamers are less stable, their quaternary

325   structure can be easily disrupted by antibody binding. This may form part of the antibody

326   neutralization mechanism. The stable tetramer is however more resistant to Fab 5E3 binding,

327   and this may help sNS1 to evade the host immune response.

Fig. 4 CryoEM structure of sNS1 complexed with Fab 5E3. **(A)** Some of the 2D class averages showing Fab bound dimers and also unbound stable tetramers. **(B)** The 3.5Å resolution structure of dimer NS1(orange) complexed with Fab 5E3 (grey) shows that they form an arch shape – the Fabs are bending towards the β-roll side of the sNS1 dimer. This NS1 dimer contains the β-roll structure which is also present in the loose tetramer and the

hexamer. The Fabs bind to the ends of the β-ladder. The 5E3 epitope is shown as blue
spheres. **(C) (i)** The 3.9Å resolution sNS1 unbound tetramer map. The two dimers of the
tetramer are colored in blue and yellow. **(ii)** Superposition of Fab 5E3:dimer onto one dimer
(yellow) of the stable tetramer shows that Fab (transparent grey surface) when bound will
clash (red arrow) with the neighboring dimer (blue). **(D)** TEER assay showing Fab 5E3 can
only partially prevent sNS1 from inducing human umbilical vein endothelial cell
hyperpermeability. Data are the mean ± SEM from three independent experiments.
Statistically significant differences between distinct groups (all time points combined for
each group) compared to the untreated groups were determined by a two-way ANOVA
analysis using Dunnett's test for multiple comparisons. (***$p < 0.001$ and **$p < 0.01$).

Discussion

Previous studies [13,14,21] and our cross-linking experiments of sNS1 (Fig. 1B) consistently showed the presence of dimers, tetramers, and hexamers in semi- and fully denaturing SDS PAGE gel. Syzdykova *et al*. also showed by native gel that yellow fever virus sNS1 consists of mainly tetramers with little hexamers [22]. Our cryoEM study shows the vast majority of sNS1 particles from a DENV2 clinical strain (PVP 94/07) exist as tetramers, while the hexamer only forms a minor population (Fig. 1E). Flamand and colleagues [21] performed small angle X-ray scattering (SAXS) experiments on sNS1 and observed particles with a homogenous gyration radius of 10nm (100 Å), which they concluded were likely hexamers. When we measured the dimensions of our cryoEM sNS1 tetrameric and hexameric structures (Extended data Fig. 3), their longest dimensions are the same (~100 Å), thus correlating well with size of particles detected by the SAXS[21]. This suggests that SAXS may not be able to effectively differentiate between the heterogenous populations of the particles within the sNS1 sample.

NS1 plays multiple roles in dengue virus infection of mammalian cells and also mosquitoes. It was shown that sNS1 is one of the major factors contributing to the development of DHF/DSS in patients. Our cryoEM studies show structural details of this important sNS1 protein – (1) the quaternary organization of the different native oligomerization states, (2) how an antibody could bind and dismantle the loose tetrameric and hexameric sNS1 structures and also (3) how sNS1 stable tetramers could resist this antibody-induced disassembly. These findings will contribute to the design of therapeutics and vaccines against severe dengue disease.

Methods

sNS1 cloning, expression and purification

DENV 2 PVP94/07 NS1-HisTag was cloned into the pcDNA 3.4-TOPO (Invitrogen) and the plasmid was transfected into Expi293 HEK cells (Thermo Fisher). After expression in HEK cells, we harvested the supernatant containing sNS1. It was then clarified by centrifuging at 9000 x g at 4°C for 30 min. The sNS1 protein was purified by using a HisTrap column (GE Healthcare). It was then further purified through a gel-filtration chromatography (Superdex 200 10/300 GL column (GE Healthcare)) in buffer (50 mM Tris 8.0 and 150 mM NaCl). Protein purity was then assessed by SDS-PAGE.

BS^3 cross-linking of protein

BS^3, a cross-linker with an 11.4 Å spacer arm, were added to NS1 in PBS to a final concentration of 5mM and the samples were incubated for 10, 20, 30 mins at room temperature. Reactions were stopped by adding Tris·HCl, pH 7.5 to a final concentration of 50mM and then the sample was added the SDS-PAGE loading buffer, boiled for 2 min and then analyzed on a 4–20 % SDS-PAGE gradient gels (Bio-Rad). The protein bands were then transferred to a nitrocellulose membrane, probed with anti-polyHistidine−peroxidase antibody and visualized by chemiluminescence.

CryoEM sample preparation

Quantifoil Au R1.2/1.3 grids were glow-discharged at 5mA for 1min using an Emitech K100X Glow Discharge Unit. About 2.1 μl of sNS1 (0.25 mg/mL) / sNS1 complexed with Fab 5E3 (molar ratio 1:1.2, final concentration of sNS1 is 0.25 mg/mL) were applied to the grids, blotted with a filter paper (Ted Pella Standard Vitrobot filter paper, Grade 595) for 2s (Blot force 2, Blot Time 2 seconds, drain time 0.5 seconds, Humidity 100%) to remove excess sample, and then flash frozen in liquid ethane by using the Vitrobot Mark IV plunger (FEI, Netherlands). To ensure temperature consistency, the temperature of the Vitrobot humidity chamber was adjusted to 4 °C for blotting.

CryoEM image acquisition and reconstruction procedure

CryoEM micrographs of sNS1/ sNS1 complex were collected using a Titan Krios transmission electron microscope operated at 300kV and equipped with a Gatan K3 direct electron detector and Gatan GIF post-column energy filter operated in zero-loss mode with a slit width of 20 eV. Images were recorded at a nominal magnification of 130,000.

SerialEM was used to carry out the automated data collection. Image shift was used to target 9 exposures per stage position with an image shift of 1.6 um. Images were recorded by movie mode, with similar total exposure of 4.9s, 58 frames per movie and total dose of 81.5 e-/Å2. The frames from each movie were aligned using MotionCor2 [23] to produce full dose images used for particle selection and orientation search. The images were taken at underfocus in the 0.8-2.0 μm range. In total, 7714 micrographs were collected for the sNS1. The astigmatic

CTF parameters were estimated with Patch CTF estimation in CryoSPARC [16] and were
accounted for during orientation search. We performed automated particle picking using
Warp [24] and CryoSPARC. A total of 2,342,754 particles were selected. These particles were
binned 4x before going through 20 iterations of 2D classification (CryoSPARC) to produce
2D class averages. The starting model was generated by the CryoSPARC Ab-Initio
Reconstruction. We then performed 25 iterations of 3D classification and reconstruction
using C1 symmetry using the program Relion [15] to remove broken and distorted particles. We
then switched to CryoSPARC to perform further 3D classification. For the class with stable
tetramers, we performed multiple cycles of heterogenous and non-uniform refinement [16] by
imposing D2 symmetry. For the loose tetramers, we performed heterogenous, non-uniform
refinement without any symmetry imposed. We then performed local refinement on one of its
dimers. For the hexamers, we performed only non-uniform refinement with D3 symmetry
imposed. The cryoEM reconstruction flow chart is summarized in Extended data Fig. 2.
Resolution was determined by using gold standard Fourier shell correlation and local map
resolution was estimated with ResMap [25]. The refinement statistics are shown in Extended
data Table 1.

Protein structure building

The sNS1 structures were first interpreted by approximate fitting using the ZIKA NS1 dimer
structure (PDB 5GS6) as rigid bodies into the density using "fit-in-map" function in Chimera
[18]. We then used COOT to mutate the residues to those of dengue virus NS1 sequence and
then fit individual residues into densities [26]. We then refined the structures using the
"phenix.real_space_refine" procedure in the Phenix software package, with default
parameters and rigid body refinement using secondary-structure and torsion angle restraints [27].
The final coordinates containing an asymmetric unit (i.e., a protomer in the stable tetramer
structure) were then used to build the biological unit (full tetramer) by using the command
"sym" in UCSF Chimera [18]. The final coordinates of the asymmetric units were checked
using MolProbity [28]. Maps and structures shown in the Figs were generated by using UCSF
Chimera and Coot.

TEER assay

A monolayer of Human Umbilical Vein endothelial cells (80%–90% confluency) cultured in
T75 flasks were detached using trypsin-EDTA (0.25%). The cells were then resuspended
using fresh culture media and then counted using an automated cell counter. 60,000 cells (300
μL per well) were seeded onto the apical side of Transwell inserts. Each Transwell was
transferred inside a 24-well format plate containing 1.5 mL of endothelial cell culture media.
Transwells containing endothelial cells were incubated at 37°C and 5% CO_2 for 3~4 days,
and 50% of the culture medium was changed in each well 48 hours post-seeding. Cells were
grown until the TEER values reached ~ 160 to 170 Ohms (Ω), indicating 100% cell
confluency. After this, 3 μg NS1 protein / NS1_Fab5E3 complex (molar ratio 1:1.2) was
added to the apical side of the Transwell insert. Electrical resistance values, measured in
Ohms (Ω), were recorded at every 2-hour time-points following the addition of the proteins
using an Epithelial Volt Ohm Meter with the "chopstick" electrodes. Endothelial permeability

was expressed as relative TEER which represents a ratio of resistance values (Ω) as follows:
(Ω experimental condition - Ω medium alone)/(Ω nontreated endothelial cells - Ω medium
alone).

Data availability

458 The cryoEM map and the atomic models of the all sNS1 and sNS1:Fab structures have been
459 deposited in the Electron Microscopy Data Bank and the Protein Data Bank - stable tetramer
460 (EMD: 32841, PDB: 7WUT), loose tetramer (EMD: 32842 and PDB: 7WUU), Dimer of
461 Loose tetramer (EMD: 32840 and PDB: 7WUS), hexamer (EMD: 32842 and PDB: 7WUV)
462 and sNS1:Fab5E3(EMD: 32839 and PDB: 7WUR).

References

1   Gubler, D. J. Dengue and dengue hemorrhagic fever. *Clin Microbiol Rev* **11**, 480-496,
doi:10.1128/CMR.11.3.480 (1998).
2   Thomas, S. J. & Yoon, I. K. A review of Dengvaxia(R): development to deployment. *Hum*
Vaccin Immunother **15**, 2295-2314, doi:10.1080/21645515.2019.1658503 (2019).
3   Halstead, S. B. Neutralization and antibody-dependent enhancement of dengue viruses. *Adv*
Virus Res **60**, 421-467, doi:10.1016/s0065-3527(03)60011-4 (2003).
4   Beatty, P. R. *et al.* Dengue virus NS1 triggers endothelial permeability and vascular leak that
is prevented by NS1 vaccination. *Sci Transl Med* **7**, 304ra141,
doi:10.1126/scitranslmed.aaa3787 (2015).
5   Carpio, K. L. & Barrett, A. D. T. Flavivirus NS1 and Its Potential in Vaccine Development.
Vaccines (Basel) **9**, doi:10.3390/vaccines9060622 (2021).
6   Modhiran, N. *et al.* Dengue virus NS1 protein activates immune cells via TLR4 but not TLR2
or TLR6. *Immunol Cell Biol* **95**, 491-495, doi:10.1038/icb.2017.5 (2017).
7   Puerta-Guardo, H., Glasner, D. R. & Harris, E. Dengue Virus NS1 Disrupts the Endothelial
Glycocalyx, Leading to Hyperpermeability. *PLoS Pathog* **12**, e1005738,
doi:10.1371/journal.ppat.1005738 (2016).
8   Libraty, D. H. *et al.* High circulating levels of the dengue virus nonstructural protein NS1 early
in dengue illness correlate with the development of dengue hemorrhagic fever. *J Infect Dis*
186, 1165-1168, doi:10.1086/343813 (2002).
9   Liu, J. *et al.* Flavivirus NS1 protein in infected host sera enhances viral acquisition by
mosquitoes. *Nat Microbiol* **1**, 16087, doi:10.1038/nmicrobiol.2016.87 (2016).
10   Akey, D. L. *et al.* Flavivirus NS1 structures reveal surfaces for associations with membranes
and the immune system. *Science* **343**, 881-885, doi:10.1126/science.1247749 (2014).
11   Xu, X. *et al.* Contribution of intertwined loop to membrane association revealed by Zika virus
full-length NS1 structure. *EMBO J* **35**, 2170-2178, doi:10.15252/embj.201695290 (2016).
12   Wang, D. *et al.* A Mutation Identified in Neonatal Microcephaly Destabilizes Zika Virus NS1
Assembly in Vitro. *Sci Rep* **7**, 42580, doi:10.1038/srep42580 (2017).

13 Muller, D. A. *et al.* Structure of the dengue virus glycoprotein non-structural protein 1 by
electron microscopy and single-particle analysis. *J Gen Virol* **93**, 771-779,
doi:10.1099/vir.0.039321-0 (2012).
14 Gutsche, I. *et al.* Secreted dengue virus nonstructural protein NS1 is an atypical barrel-
shaped high-density lipoprotein. *Proc Natl Acad Sci U S A* **108**, 8003-8008,
doi:10.1073/pnas.1017338108 (2011).
15 Scheres, S. H. RELION: implementation of a Bayesian approach to cryo-EM structure
determination. *J Struct Biol* **180**, 519-530, doi:10.1016/j.jsb.2012.09.006 (2012).
16 Punjani, A., Rubinstein, J. L., Fleet, D. J. & Brubaker, M. A. cryoSPARC: algorithms for rapid
unsupervised cryo-EM structure determination. *Nat Methods* **14**, 290-296,
doi:10.1038/nmeth.4169 (2017).
17 Punjani, A., Zhang, H. & Fleet, D. J. Non-uniform refinement: adaptive regularization
improves single-particle cryo-EM reconstruction. *Nat Methods* **17**, 1214-1221,
doi:10.1038/s41592-020-00990-8 (2020).
18 Pettersen, E. F. *et al.* UCSF Chimera--a visualization system for exploratory research and
analysis. *J Comput Chem* **25**, 1605-1612, doi:10.1002/jcc.20084 (2004).
19 Modhiran, N. *et al.* A broadly protective antibody that targets the flavivirus NS1 protein.
Science **371**, 190-194, doi:10.1126/science.abb9425 (2021).
20 Biering, S. B. *et al.* Structural basis for antibody inhibition of flavivirus NS1-triggered
endothelial dysfunction. *Science* **371**, 194-200, doi:10.1126/science.abc0476 (2021).
21 Flamand, M. *et al.* Dengue virus type 1 nonstructural glycoprotein NS1 is secreted from
mammalian cells as a soluble hexamer in a glycosylation-dependent fashion. *J Virol* **73**, 6104-
6110, doi:10.1128/JVI.73.7.6104-6110.1999 (1999).
22 Syzdykova, L. R. *et al.* Fluorescent tagging the NS1 protein in yellow fever virus: Replication-
capable viruses which produce the secretory GFP-NS1 fusion protein. *Virus Res* **294**, 198291,
doi:10.1016/j.virusres.2020.198291 (2021).
23 Zheng, S. Q. *et al.* MotionCor2: anisotropic correction of beam-induced motion for improved
cryo-electron microscopy. *Nat Methods* **14**, 331-332, doi:10.1038/nmeth.4193 (2017).
24 Tegunov, D. & Cramer, P. Real-time cryo-electron microscopy data preprocessing with Warp.
Nat Methods **16**, 1146-1152, doi:10.1038/s41592-019-0580-y (2019).
25 Kucukelbir, A., Sigworth, F. J. & Tagare, H. D. Quantifying the local resolution of cryo-EM
density maps. *Nat Methods* **11**, 63-65, doi:10.1038/nmeth.2727 (2014).
26 Emsley, P. & Cowtan, K. Coot: model-building tools for molecular graphics. *Acta Crystallogr D
Biol Crystallogr* **60**, 2126-2132, doi:10.1107/S0907444904019158 (2004).
27 Adams, P. D. *et al.* PHENIX: a comprehensive Python-based system for macromolecular
structure solution. *Acta Crystallogr D Biol Crystallogr* **66**, 213-221,
doi:10.1107/S0907444909052925 (2010).
28 Williams, C. J. *et al.* MolProbity: More and better reference data for improved all-atom
structure validation. *Protein Sci* **27**, 293-315, doi:10.1002/pro.3330 (2018).

Acknowledgments

We thank Valerie S-Y Chew, Xinni Lim and Qunfei Zhou for help in experiments. This work was supported by Duke-NUS Signature Research Programme funded by the Ministry of Health, Singapore and National Research Foundation Investigatorship award (NRF-NRFI2016-01) awarded to SML. AWKT is supported by a Khoo Postdoctoral Fellowship Award (Duke-NUS-KPFA/2021/0044) from Duke-NUS Medical School and the Estate of Tan Sri Khoo Teck Puat.

Author information

Affiliations

Program in Emerging Infectious Diseases, Duke-National University of Singapore Medical School, Singapore 169857, Singapore.

Bo Shu, Justin S.G. Ooi, Aaron W.K Tan, Thiam-Seng Ng, Guntur Fibriansah, Victor A. Kostyuchenko, Shee-Mei Lok

Centre for Bio-Imaging Sciences, Department of Biological Sciences, National University of Singapore, Singapore 117557, Singapore.

Bo Shu, Aaron W.K Tan, Thiam-Seng Ng, Guntur Fibriansah, Victor A. Kostyuchenko, Shee-Mei Lok

Center for Bioimaging Sciences, National University of Singapore, Singapore, 117557, Singapore.

Jian Shi, Shee-Mei Lok

Medical Sciences Division, University of Oxford, Oxford OX3 9D

Wanwisa Dejnirattisai, Juthathip Mongkolsapaya, Gavin Screaton

Contributions

S.M.L. supervised the project. S.M.L. and B.S. designed research studies; J.S.G.O. produced sNS1. W.D., J.M. and G.S. produced Fab. B.S. and J.S.G.O. conducted experiments. T.S.N., A.W.K.T. and J.S. collected cryo-EM data; S.M.L., B.S., G.F. and V.A.K. analyzed data; S.M.L. and B.S. wrote the manuscript.

Corresponding author

Correspondence to Shee-Mei Lok.

Ethics declarations

Competing interests

The authors declare no competing interests.

Additional information

Extended Data Fig. 1

Extended Data Fig. 1 Previously published structures of the dengue NS1. (A) Crystal structure of dengue dimeric iNS1 (PDB:5GS6). Left: top view, looking from the β-roll of the iNS1 dimer. One protomer contains three domains: starting from the N-terminus- β-roll (blue), wing domain (yellow) and the β-ladder (red). It interacts with the other protomer (colored in grey) through extensive hydrogen bonds. Right: side view of the iNS1 dimer. The β-roll is highly hydrophobic and is predicted to interact with the cellular membrane. (B) A previously published low resolution EM reconstruction of the sNS1 (EMDB:2073). This map is reconstructed from negatively stained sNS1 and C3 symmetry was imposed [13].

 Extended Data Fig. 2

Extended Data Fig. 2 The cryoEM reconstruction workflow for uncomplexed sNS1.
594

Extended Data Fig. 3

Extended Data Fig. 3 The initial sNS1 cryoEM maps after 3D classification using C1 symmetry. Dimensions of the stable, loose tetramers and hexamers are shown. Fitted models are shown on the bottom row.

Extended Data Fig. 4

Extended Data Fig. 4 The 3.5Å resolution sNS1 stable tetrameric structure. (A) Different
views of the cryoEM map displayed at high contour level. The densities are colored
according to their local resolution. (B) The fitted structure show in the same view as the
cryoEM map directly above (in (A)). There are two dimers facing each other. One dimer is
colored in pink and green, while the other is light blue and yellow. (C) Fourier shell
correlation plot showing resolution. (D) Zoom-in views of density map showing the β-strands
densities are well resolved (left). Some of the side chains densities (right) can also be
observed.

Extended Data Fig. 5

Extended Data Fig. 5 CryoEM maps of the sNS1 loose tetramer. (A) 8Å cryoEM map of the loose
tetramer. (i) Different views of the fitted structure of the loose tetramer into the density map
(transparent grey). (ii) FSC plot showing resolution. (B) The 3.4Å cryoEM map of one dimer of the
loose tetramer after focused refinement. (i) Different views of the cryoEM map. Density is colored
according to local resolution. (ii) Up: FSC plot showing resolution of the map: Down: zoom-in view
showing that some side chain densities are well resolved. (iii) The fitted structure of one of the dimers
(pink and light green) is very similar to that of dengue iNS1. The β-roll in each protomer is colored in
a darker shade of the respective protomer's color.

Extended Data Fig. 6

Extended Data Fig. 6 The 8Å cryoEM map of the sNS1 hexamer. (A) FSC plot showing
resolution of the map. (B) The hexamer density displayed at different contour levels. At lower
contour levels, densities at the interior of the hexamer can be seen. (C) Different views of the
fit of the three dimers into the hexamer density map.
631

Extended Data Fig. 7

Extended Data Fig. 7 3.5Å resolution cryoEM structure of Fab 5E3:sNS1 dimer. (A)
Side and top views of the cryoEM map of the Fab 5E3:sNS1 dimer. NS1 is colored in orange
while Fabs in grey. (B) Left: FSC plot showing the resolution of the map. Right: zoom-in
view showing well-resolved β-strands. (C) Zoom-in view of the fitted β-roll into its
corresponding density.

Extended Data Fig. 8

Extended Data Fig. 8 Comparison of the binding (top panels) and epitope (bottom panels) of Fab 5E3 to other previously published Fab:NS1 complexes [19,20]**.**

Extended Data Fig. 9

Extended Data Fig. 9 Superposition of Fab 5E3:dimer into the (A) loose tetramer and (B)
hexamer and the analysis of percentage exposure of epitopes (C-E) in the stable, loose
tetramers and hexamers. (A-B) Superposition of Fab 5E3:dimer onto one dimer (yellow) of
the loose tetramer and hexamer. Fab is shown as transparent grey surface and clashes with
neighboribg NS1 dimer is indicated with red arrows. (C-E) The percentage exposure of
epitopes in the oligomers. The two epitopes within one dimer are colored in orange and cyan.
(C) In the loose tetramer, 37.5% of the orange epitope is exposed while the cyan epitope is
fully exposed (100%). (D) In the hexamer, 37.5% of both the orange and cyan epitopes are
exposed. (E) In the stable tetramer, the two epitopes are both 37.5% exposed.

	DENV2_NS1_Tetramer	DENV2_NS1_dimer	DENV_NS1_Fab5E3
Data collection and processing	(EMDB-32841) (PDB 7WUT)	(EMDB-32840) (PDB 7WUS)	(EMDB-32839) (PDB 7WUR)
Microscope	FEI Titan Krios	FEI Titan Krios	FEI Titan Krios
Camera	Gatan K3	Gatan K3	Gatan K3
Magnification	130,000 ×	130,000 ×	130,000 ×
Voltage (kV)	300	300	300
Electron exposure (e$^-$/Å^2)	81.5	81.5	80.1
Defocus range (μm)	-0.8 to -2.0	-0.8 to -2.0	-0.8 to -2.0
Exposure rate (e$^-$/A^2/sec)	8.74	8.86	8.35
Number of frames per movie	58	58	58
Pixel size (Å)	0.66	0.66	0.66
Energy filter slit width (eV)	20	20	20
Automation software	SerialEM	SerialEM	SerialEM
Symmetry imposed	D2	C1	C1
Micrographs used	7714	7714	7145
Total number of extracted particles	2,342,754	2,342,754	1,264968
Total number of refined particles	278,366	454,477	273,074
Number of particles in final map	80,674	181,587	95,895
Map resolution (Å)	3.5	3.3	3.5
FSC threshold	0.143	0.143	0.143
Refinement			
Initial model used (PDB code)	5GS6	5GS6	5GS6
Refinement package, (resolution cutoff)	Phenix, real-space refinement (3.5)	Phenix, real-space refinement (3.3)	Phenix, real-space refinement (3.5)

Resolution of unmasked and masked reconstructions at 0.5 and 0.143 FSC	Unmasked 3.3/3.9 Masked 3.3/3.8	Unmasked 3.5/4.0 Masked 3.5/3.9	Unmasked 3.5/4.1 Masked 3.4/3.9
Map sharpening B factor (Å^2)	-230	-210	-105.9
Model composition			
Non-hydrogen atoms	11,176	5,456	8,928
Protein residues	1400	682	1134
Ligands	0	0	0
Global CC (CCvol)	0.88	0.85	0.79
Local CC (CCmask)	0.77	0.84	0.78
B factors (Å^2)			
Protein	88.90	93.61	134.83
Ligand	-	-	-
R.m.s. deviations			
Bond lengths (Å)	0.009	0.005	0.009
Bond angles (°)	1.116	1.042	1.000
Validation			
MolProbity score	2.70	2.35	2.49
Clashscore	31.67	18.75	24.01
Poor rotamers (%)	0.00	0.00	0.00
C-beta deviations	0.00	0.00	0.00
Ramachandran plot			
Favored (%)	81.54	88.94	87.25
Allowed (%)	18.46	11.06	12.75
Disallowed (%)	0	0	0
CaBLAM outliers (%)	8.89	6.08	5.05

EMRinger score	2.70	2.26	2.65

Extended Data Table 1 Cryo-EM data collection, refinement and validation statistics

	Stable Tetramer	Loose Tetramer	Hexamer
Particles	126130	90928	0
Percentage	58%	42%	0 %

665 **Extended Data Table 2 The percentage of particles in different oligomerization states**
666 **determined after 3D classification of NS1 treated with detergent (0.05% DDM).**

NS1	Fab Heavy Chain	Fab Light Chain
138		27(CDR1)
269		94(CDR3)
281	52/53/54/55/56(CDR2)	
299	102/103(CDR3)	
301	33(CDR1)	
302	103(CDR3)	
303	103(CDR3)	
304	32/106(CDR1/CDR3)	
305	31/33/103(CDR1/CDR3)	
306	31(CDR1)	
307	31/53(CDR1/CDR2)	
324	56(CDR2)	
325	57(CDR2)	
326	57/102/100(CDR2/CDR3)	
327	100/102(CDR3)	94(CDR3)
328	100(CDR3)	

668   **Extended Data Table S3 List of hydrogen bonds/electrostatic interactions between Fab**
669   **5E3 (heavy and light chains) with the NS1 protein. The interacting CDRs of the Fabs**
670   **are indicated.**