Extracting multiple surfaces from 3D microscopy images in complex biological tissues with the Zellige software tool.

Céline Trébeau[1], Jacques Boutet de Monvel[1], Gizem Altay[1], Jean-Yves Tinevez[2]* and Raphaël Etournay[1]*.

[1]Institut Pasteur, Université Paris Cité, Inserm, Institut de l'Audition, F-75012 Paris, France
[2]Institut Pasteur, Université Paris Cité, Image Analysis Hub, F-75015 Paris, France
* To whom correspondence should be addressed, tinevez@pasteur.fr, raphael.etournay@pasteur.fr

Abstract.

Efficient tools allowing the extraction of 2D surfaces from 3D-microscopy data are essential for studies aiming to decipher the complex cellular choreography through which epithelium morphogenesis takes place during development. Most existing methods allow for the extraction of a single and smooth manifold of sufficiently high signal intensity and contrast, and usually fail when the surface of interest has a rough topography or when its localization is hampered by other surrounding structures of higher contrast. Multiple surface segmentation entails laborious manual annotations of the various surfaces separately. As automating this task is critical in studies involving tissue-tissue or tissue-matrix interaction, we developed the Zellige software, which allows the extraction of a non-prescribed number of surfaces of varying inclination, contrast, and texture from a 3D image. The tool requires the adjustment of a small set of control parameters, for which we provide an intuitive interface implemented as a Fiji plugin. As a proof of principle of the versatility of Zellige, we demonstrate its performance and robustness on synthetic images and on four different types of biological samples, covering a wide range of biological contexts.

Introduction.

The interplay between gene regulatory networks and physical forces in driving collective cell behaviors is key to tissue morphogenesis during development and to tissue homeostasis throughout life. Recent quantitative studies of epithelial morphogenesis have begun to unravel the basic cellular and physical principles of tissue development, by providing the tools to integrate multiple scales of tissue dynamics [1–4]. These tools are instrumental to quantify how cell shape changes, cell divisions, cell rearrangements and cell extrusions contribute to tissue remodeling, and to establish data-driven computational models of tissue morphogenesis.

Quantitative analysis of an epithelium starts with the extraction of its apical surface from 3D-microscopy images (z-stacks of xy-optical sections) encompassing the volume immediately surrounding the epithelium. However, this is a difficult task because this surface is usually not flat (it is best modelled as a curved surface, or 2D submanifold embedded in 3D space), and is often surrounded by other biological structures such as cell layers, acellular membranes, extracellular matrix, and vesicles that hamper its visualization and reconstruction. To make the surface extraction tractable, these studies rely on specific preparations of the specimen, allowing to expose the entire epithelial surface labeled with junctional fluorescent markers to reveal the network formed by epithelial cell-cell contacts. Once the epithelial surface has been extracted, automated cell segmentation and cell contour tracking tools can be used to follow the dynamics of every cell within the epithelium.

Another challenging experimental limitation of these studies is that some of the structures surrounding the epithelium exert external physical constraints that are known to critically affect epithelial morphogenesis by directing cellular dynamics and signaling pathways [1, 5–7]. To understand the physical forces controlling tissue morphogenesis, it is thus essential to also characterize how the dynamics of these extra-epithelial surfaces relate to that of the epithelium (see [8] for review). This calls for the development of dedicated tools allowing the automated extraction of information from several surfaces of interest in a given sample, since the sheer volume of the data precludes any attempt at a manual analysis.

Several surface extraction tools have been developed, some of which are available as open access software, such as PreMosa [9], FastSME [10], and LocalZProjector [11]. These tools focus on the extraction of a single, near-horizontal epithelial layer, which is assumed to (i) be sufficiently smooth, (ii) show enough contrast against surrounding background signals, and (iii) cover the entire image field-of-view. Specifically, it is assumed that the fluorescent marker used to label the epithelial cell network should provide the highest contrast in the image and allow to select it out from autofluorescent extracellular structures such as the cuticle in flies, or other acellular membranes in mammalian epithelia. The surface is then localized using heuristic algorithms based on the detection of the pixels of maximum contrast and/or brightness. However, applying these tools on more complex biological images with several epithelia of weaker contrast often leads to incorrect localization of the surface of interest, and its blending with the nearby unwanted biological structures.

MinCostZ on the other hand, is the only available open-source tool that allows the extraction of up to two surfaces from a 3D stack, and imposes explicit continuity constraints on the reconstructed surfaces. MinCostZ surface extraction relies on a previously developed formulation of the problem as a graph-cut optimization [12]. It is implemented as an ImageJ plugin [13], taking as control parameters, the number of surfaces to extract, the maximum slope and the range of distances allowed between the surfaces, as well as some user-defined cost function that should reflect the characteristics of the surfaces in term of signal intensity, contrast and texture. Despite its interest, this approach remains computationally costly and difficult to apply in practice due to the non-trivial choice of the cost function and the need to know beforehand the relative positions of the surfaces to be extracted.

Alternatively, one can segment the surfaces of interest by using supervised machine learning tools
such as the software solutions Weka [14] or Ilastik [15], as proposed in the ImSAnE surface reconstruction
framework [16]. A deep learning approach, using a network of the U-net type to segment the pixels
belonging to a single surface of interest, has also recently been reported [17]. While promising as they can
provide state of the art segmentations of epithelial surfaces in difficult imaging conditions, machine learning
approaches require the prior manual annotation of a sufficiently large set of surfaces to generate suitable
training sets. This process can be very time consuming, often necessitating several rounds of trials and
errors to obtain satisfactory results, without guarantees to be generic, *i.e.*, to generalize to a wide range of
datasets. So far, no solution to the multiple surface extraction problem has been proposed, which is
satisfactory both in terms of genericity and ease of use.

However, such a tool is highly desirable for modern biology studies. Indeed, tissue organization in
the context of developmental biology emerges from the interaction of several neighboring structures [8]
through the interplay of molecular signals [18, 19], as well as electrical [20, 21], hydraulic [22, 23] and
mechanical contact interactions [24, 25]. The importance of such interaction is exemplified in embryonic
explanted tissue cultures that develop abnormally when separated from their neighboring structures [26].
Similarly, in the context of tissue engineering, stem-cell derived aggregates harbor various types of tissues
surrounding the genuine organoid, and these tissues presumably influence organoid shape, fate and
differentiation (see [27] for review). The ability to simultaneously study the dynamics of neighboring
structures together with the structure of interest is therefore essential for an integrated understanding of
tissue development, and for any attempt to harness tissue self-organization *in vitro*.

Here, we introduce Zellige, a tool based on a novel constructive approach that allows the automatic
extraction of a non-prescribed number of surfaces from a 3D image. To do this, the user is only required to
adjust a small set of intuitive control parameters, a task largely facilitated by a user-friendly interface
implemented as a plugin for the open-source Fiji platform [28]. We tested the performance and robustness
of Zellige for multiple surface extraction by applying it to synthetic images and 3D microscopy images from
four different types of biological samples, containing multiple surfaces of interest of widely varying texture
and contrast. These experiments demonstrate the ability of the approach to extract several (up to 4)
surfaces of potentially very low contrast, selectively from other highly contrasted and complex structures,
with a single set of reconstruction parameters. A sensitivity analysis also reveals a high robustness of Zellige
against small variations of these parameters. This will make it a tool of choice in terms of versatility and
ease of use for the investigation of biological surfaces.

Results & discussion.

Proof of concept of multiple surface extraction on a synthetic image.

The implementation of Zellige is summarized in **Figure 1** and in the *Methods* section (**Figure S1**). **Figure 2** shows the results produced by Zellige on a phantom 3D image [29] containing three distinct synthetic surfaces generated as described in Supplementary note 2. The three surfaces are extracted with little errors. We assessed the quality of the reconstruction by comparing each of the height-maps produced by Zellige to the corresponding ground truth (GT) height-map, which is exactly known in this case (**Figure 2A-C**, and Supplemental note 3). For the three surfaces, the reconstruction has subpixel accuracy over >99% of the GT pixels (**Figure 2D-E**), with a root mean square error (RMSE) of ≤ 0.6 in pixel units, showing that the surface localization is highly accurate. In addition, the coverage, which measures the proportion of the reconstructed surface relative to the GT, is near 100% for the three surfaces. To achieve these results, the control parameters of the two steps of the surface extraction were adjusted manually to some adequate reference values (see Supplementary Table S1) using the Zellige Fiji interface. Only the parameters controlling the pixel classification step (amplitude and Otsu threshold parameters T_A and T_{otsu}, minimal island size S_{min}, and smoothing parameters σ_{xy} and σ_z) did actually require a modest adjustment. The parameters of the surface assembly step (parameters T_{OSE1}, R_1, C_1 and T_{OSE2}, R_2, C_2 of the 1st and 2nd construction rounds, respectively) were set to their default reference values (see **Figure S2** and Supplemental note 4) and did not need to be adjusted.

Thus, using a single set of control parameters, Zellige can extract multiple surfaces of various shapes and textures with virtually no error, without requiring the user to provide information about their number or relative position, nor about their shape or texture characteristics.

Performance of Zellige on biological samples.

Example 1: Extracting multiples surfaces from an image of a pupal fly specimen.

Over the past few decades, the *Drosophila* model has been invaluable to decipher the molecular and cellular mechanisms underlying organ embryogenesis [30, 31]. Epithelium morphogenesis studies not only revealed the importance of mechanical stresses (including stress boundary conditions) and planar polarity signaling on cell dynamics to generate tissues of reproducible sizes and shapes, it also highlighted the importance of extracellular matrix attachments in constraining the tissue stresses that guide patterning [1, 32]. At the pupal stage, the fly undergoes dramatic remodeling of its larval organs into adult organs. Large scale tissue flows initiate at a timing that coincides with molting, when the epithelium contracts away from the overlaying cuticular sac, a protective acellular membrane that imposes mechanical boundary conditions to the tissue.

Figure 3 shows the results of applying Zellige on a 3D image of a *Drosophila* pupa acquired with a spinning disk confocal microscope [33]. The sample expresses Ecadherin-GFP, a fluorescent marker of cell-cell junctions, and encompasses a portion of the pupa's abdomen and a small portion of its wing. Four surfaces of interest can be identified, with varying signal intensities, noise levels and features (**Figures 3A-B**). The abdomen is formed of an epithelium (surface S2) overlaid by a cuticle (surface S1). Lying just beneath these two surfaces, one can observe globular structures showing in some places a higher intensity than the signal coming from the surfaces. The wing also consists of an epithelium of low intensity signal (surface S4), and an overlying cuticle (surface S3). These two surfaces are relatively flat, except for surface S3 which is very steep near one of its edges.

Figure 3C shows a 3D graphical representation of the height-maps reconstructed by Zellige (green)
and those reconstructed by an expert biologist (blue), taken as ground truth (GT). While these height-maps
clearly show greater roughness than those of the synthetic surfaces presented earlier, they could again be
obtained with a single set of control parameters that were adjusted manually with the Zellige interface (see
Supplementary Table S1). We observe an excellent match between the four reconstructed and
corresponding GT height-maps, despite the rather complex topography of surfaces S1 and S2 (with slopes
reaching up to 45°), and the near-vertical inclination of surface S3 at its boundary. Yet, small deviations may
be seen in the regions of highest slope of the surfaces. Some of these deviations are likely attributable to
uncertainties in the definition of the GT height-maps, whose accuracy depends on the expert.

Figure 3D shows the differences between the reconstructed and GT height-maps, plotted as color-
coded error maps. These differences are <2 (in pixel units) for most pixels, while some regions of higher
error values can be seen locally in surfaces S1 and S2, and at the boundary of surface S3. Note that for
surfaces S2 and S4, which contain junctional epithelial meshes composed of larger and smaller cells,
respectively, the GT height-map encompass not only the mesh but also the interior of the cells, where no
junctional signal is detected. The distance calculated inside the cells is thus more subjected to intensity
fluctuations, especially for surface S2. Nonetheless, the RMSEs of surfaces S1, S2 and S4 are less than 1,
showing that on average the reconstructed height-maps match the corresponding ground truth with
subpixel accuracy. The higher RMSE (1.25) of surface S3 is largely due to the region of steep region at the
edge of this surface (yellow region on the error map for this surface, **Figure 3D**). The coverage of the
reconstructed height-maps is excellent (≥ 96%) for surfaces S1 and S2, and slightly lower, but still very good
(≥ 93%) for the smaller surfaces S3 and S4. **Figure 3E** shows the 2D projections of the 3D image obtained
for each of the reconstructed surfaces and for the corresponding ground-truths. The inaccuracies visible on
the error maps (see **Figure 3D**) do not significantly impact these projections, which appear very similar to
the projections obtained with the corresponding ground-truths. Thus, while the biological sample contains
significant noise and shows a much more variable contrast (especially with the presence of high intensity
globular structures near surfaces S1 and S2), Zellige makes it possible to segment these surfaces selectively,
with a quality of segmentation comparable to that obtained by manual expert segmentation.

This possibility brings several perspectives that are not offered by single surface extraction
algorithms. First it opens the possibility to systematically study the tissue axial movements (along z) relative
to the cuticle during molting, allowing for example to gain insights into the early tissue contraction of the
wing hinge that acts as a mechanical inducer over the wing blade [8]. Second, Zellige makes it possible to
automatically extract structures such as the abdomen epithelium, which is usually segmented manually
[34], due to the difficulty to separate the large larval cells from the cuticle mesh and from other globular
structures (such as fat bodies or macrophages) present underneath the epithelium. All these structures
become intertwined when using other extracting tools. In this context, Zellige opens new opportunities to
study collective cell behavior during epithelial morphogenesis *in vivo*, and to integrate in the analysis the
surrounding surface-like structures involved in the mechanics of the system.

Example 2: Extracting a thin cochlear epithelium surface from a multilayer dataset.

As the first model in which planar cell polarity signalling was shown to be conserved in vertebrates [35], the
mammalian auditory organ, the cochlea, is arguably our most valuable model to study epithelial patterning
and morphogenesis beyond the fly and zebrafish [36, 37]. Cochlear morphogenesis involves complex and
tightly controlled patterning processes during which the cochlear sensory epithelium extends and develops
its characteristic coiled snail shape, while adopting a striking cellular mosaic organization, with graded
changes of morphogenetic parameters along the cochlea [38, 39]. These morphogenetic processes are well
recapitulated in organotypic cultures, on the condition that the mesenchyme that underlies the epithelium

be preserved. The cultures are then amenable to live imaging [37, 40], pharmacological [41] and genetic manipulations.

Figure 4A shows a confocal swept field microscope acquisition of an embryonic mouse cochlea [42]. The sample contains only one surface of interest, the cochlear epithelium, but this surface lays on top of a thick tangled mesh of non-epithelial cells originating from the mesenchyme. The whole biological tissue is stained for filamentous actin (F-actin) using phalloidin. The epithelium surface presents a non-uniform signal included in a small z-range ($6 \leq z \leq 10$), and a mesh of very heterogeneous size. Between sections $z = 10$ and $z = 14$ one can observe the basolateral region of the epithelial cells, also stained for F-actin. he particularity of this sample is that the mesenchyme presents an intense and contrasted signal over a wide range of z-values ($14 \leq z \leq 43$). This makes it challenging to extract the surface of the epithelium, which is characterized by low intensity and low contrast.

Figure 4B shows a 3D representation of the height-map reconstructed by Zellige and the corresponding GT height-map (again reconstructed manually by an expert). On the corresponding error map (**Figure 4C**), most (~83%) pixels of the reconstructed height-map show subpixel accuracy (with distances < 1 to the corresponding pixels of the GT height-map). The errors are greater in regions where the cell size is larger, as well as in the area where the signal intensity is very low. However, they remain smaller than 2 for > 95% of the pixels. This result is consistent with the low value of the RMSE (1.1). The surface is also reconstructed with an excellent coverage (> 99%, **Supplementary Table S1**).

As the sample contains a single epithelial surface of interest, we compared the performance of Zellige with three other software that can extract only a single surface (**Figure 4D**). The projections of PreMosa, FastSME and LocalZProjector completely miss the epithelium. Only regions of high contrast corresponding to the mesenchyme are projected. In contrast, Zellige generates a projection very close to the ground truth. This demonstrates the efficiency of Zellige to selectively extract a low contrast surface, despite the presence of several structures of higher contrast. Indeed, Zellige detects every structure as a possible surface seed without any assumption on its contrast, and only extends this seed into a surface if enough spatial continuity is found in the surrounding signal. This feature allows to separate individual surfaces from other structures spatially, which should greatly facilitate the analysis of live imaging experiments.

Example 3: Extracting a single bronchial epithelial surface rendered abnormally rough by SARS-CoV-2.

Recently, we used Zellige to extract the surface of a primary culture of bronchial epithelial cells following infection by the SARS-CoV-2 virus [43]. The infection causes the surface of the epithelium to become abnormally rough due to cell damages as seen from discontinuities within the cell layer. The sample we chose from this study is a 3D confocal image of the epithelium responding to SARS-CoV-2 infection [44] (**Figure 5A**). The surface of interest in this image corresponds to the layer of epithelial cells stained for the tight junction protein Zona Occludens-1 (ZO-1). The surface roughness causes the network of junctions to extend over the height of the z-stack, with a signal of varying intensity (**Figure 5A**). In addition, the junctional network remains non-planar even at the level of a single cell, hence violating the smoothness condition commonly assumed to hold in the context of epithelial surface extraction. We also observe the presence of nearby punctiform structures of high contrast that are mainly located outside of the epithelium surface. This sample therefore provides an example of a surface with a complex landscape, interspersed with a constellation of signals which may interfere with the surface segmentation.

The 3D representation of the reconstructed and corresponding GT height-maps (**Figure 5B**) makes it possible to appreciate the roughness of the surface of interest. The two height-maps overlap quite satisfactorily. As shown on the error map (**Figure 5C**), a large majority (71.1%) of pixels of the reconstructed

height-map show errors smaller than 1 pixel (~ 96% of them showing errors smaller than 2 pixels). The error is however larger in regions where the cell size is larger, as well as in areas where the ZO-1 signal intensity is very low, preventing a complete reconstruction of the junctions. Nevertheless, the overall RMSE remains small, with a value of 0.81 (**Figure 5**). The coverage of the reconstructed surface is also excellent (98% of the GT height-map), despite the above-mentioned discontinuities. **Figure 5D** shows the comparison of the projection generated by Zellige to those produced by PreMosa, FastSME, and LocalZProjector. The projection generated by PreMosa misses many junctions of the epithelial network, but it removes quite well the punctiform signal originating from other optical sections. FastSME performs better than PreMosa in reconstructing the junctions, but they produce a projection where the punctiform signal remains strong. In contrast, Zellige and LocalZProjector manage to both reconstruct the surface well and to filter out the punctiform signal quite effectively. This result demonstrates the efficiency of Zellige to extract a surface with complex topography by excluding intense and contrasted spurious signals away from the epithelium surface.

Example 4: Extracting the apical and basal layers of a dome-shaped epithelium (developing inner ear organoid).

Organoids are stem cell-derived and self-organizing 3D tissue structures that can mimic certain organ structures. They have emerged as promising *in vitro* models for developmental biology research, as well as biomedical translational research applications. Here we take the example of mouse stem cell derived inner ear organoids that form vesicular structures composed of an epithelium harboring sensory cells. These organoids are part of a cellular aggregate that also contains other tissues such as the mesenchyme [45] adjacent to the organoids. The epithelial cells of the forming inner ear organoids acquire a basal-apical polarity, with their apical side facing the lumen of the organoid, and their basal side facing outwards. The apical junctional network of the epithelium is difficult to visualize in microscopy images as it is seen from below, through the basal layer. Another difficulty is the spherical geometry of the vesicle system, which makes the epithelial surface of interest difficult to extract in regions of high inclination relative to the focal plane.

Figure 6 shows the result of applying Zellige on a 3D confocal microscopy image of half of a developing inner ear organoid at 14 days of culture, a stage at which markers characteristic of the mouse otic vesicle can be detected [46]. The sample was fixed and stained for F-actin to visualize all cellular structures including the epithelium. Two surfaces of interest can be identified, namely the basal side of the epithelium and the apical junctional network (**Figure 6A-B**). Both surfaces are mesh-like structures of high inclination, high signal intensity and high contrast. The vesicle lumen also contains cell debris of high intensity and contrast that are not part of any surface of interest.

Figure 6C shows a 3D graphical representation of the height-maps reconstructed by Zellige and those reconstructed by an expert biologist, taken as ground truth (GT). Due to their dome-shaped topography, the manual segmentation of these surfaces was rather laborious, and is more likely prone to errors in the regions of high inclination. Despite this, we observe an excellent match between the two reconstructed and corresponding GT height-maps (**Figure 6D).** The distance between the two height maps is <2 for the large majority (> 92%) of pixels, while larger error values occur locally in regions of near-vertical inclination of the surfaces. The RMSEs of both apical and basal surfaces are close to 1, showing that on average the reconstructed height-maps match the corresponding GT with about pixel accuracy. The coverage of the reconstructed height-map is nearly 100% for the basal surface, and ~88% for the apical surface. **Figure 6E** shows the 2D projections of the 3D image obtained for each of the reconstructed surfaces, as compared to the projected GT height maps. Thus, for the extraction for these highly inclined

surfaces, Zellige produces height maps of a quality comparable to those obtained from manual expert
segmentation.

In this example, Zellige could be combined with a 2D cell tracking framework such as TissueMiner
[3] to perform cell dynamics analysis. Note that geometric distortions introduced in the projected surfaces
by the epithelium inclination could be corrected for, using complementary tools such as DProj [11]. This
approach could provide a means to quantitatively address how an inner ear organoid epithelium patterns
at the cellular and organoid scales, while quantifying the epithelial thickness changes due to cellular
intercalation or cell shape changes in the depth of the epithelium. This would also permit to better
characterize the variability of inner ear organoids within in a given aggregate, and it could allow one to
explore how the organoid interacts with surrounding tissues and how these interactions influence the
differentiation of their constituent sensory cells.

A sensitivity analysis reveals the robustness of Zellige in extracting surfaces from biological images.

To evaluate further the quality and robustness of the segmentation obtained by Zellige, we carried out a
sensitivity analysis of the reconstruction on each of the samples tested. This analysis consisted in varying
one control parameter at a time (**Figures S2-S6**, **Figure 7**, Supplemental note 4), while keeping the other
parameters fixed at a nominal value (Supplementary Table S1). The RMSE and coverage of each of the
reconstructed surfaces were evaluated and plotted as a function of the value of the parameter that was
varied.

Figure S3 shows the results of the sensitivity analysis carried out on the image of **Figure 3**
(Example 1, pupal fly specimen), when varying the parameters controlling step 1, *i.e.* the selection of
putative surface pixels (parameters T_A, T_{otsu}, S_{min}, σ_{xy} and σ_z). As can be seen, the variations of the two
classification threshold parameters T_A, T_{otsu} and of the minimum island size S_{min} in their respective intervals
does not substantially modify the RMSE and the coverage of the reconstructed surfaces, whose values
remain roughly constant for $T_A \leq 8$, $T_{otsu} \leq 12$ and over the entire S_{min} interval (**Figure S3A**). Within these
intervals, the RMSEs of all the reconstructed surfaces remain ≤1.5, while the coverage values are > 95% for
surfaces S1 and S2, and > 85% for surfaces S3 and S4. The surfaces S1 and S3 show lower signal intensity
and lower contrast than surfaces S2 and S3, making them more difficult to extract. Surface S4 has the lowest
contrast, and fails to be reconstructed if the classification threshold values are too stringent (namely for
$T_{otsu} > 12$, or $T_A > 8$). Nevertheless, the intervals of stability of T_A and T_{otsu} (that is, the intervals of values
over which a high-quality extraction of all surfaces is obtained) remain relatively wide (cf. **Figure 7**). The
smoothing parameters σ_{xy} and σ_z also have some effect on the reconstruction of the surfaces. When σ_{xy} is
less than 3, the RMSE is higher for the mesh-like epithelial surfaces S2 and S4 (formed by the junctional
network of the epithelium). A minimal smoothing along the axial direction is also important to ensure that
the reconstructed surfaces are not too fragmented, preventing their complete reconstruction. Yet, σ_z
should not be chosen too large either, to avoid merging nearby surfaces along the z axis. In this case, the
closely positioned surfaces S1 and S2 are well-separated if setting σ_z close to 1, but they become merged
when $\sigma_z > 2$. In general, the surface construction parameters have little effect on RMSE and coverage
(**Figure S3B**). The sensitivity analysis on this challenging specimen shows a good robustness of Zellige to
extract the four surfaces with a single set of parameters, each of which can be chosen in a reasonably wide
interval considering the other fixed.

The results of the sensitivity analyses performed with the other biological image stacks
(examples 2, 3 and 4 described above) are shown in **Figures S4**, **S5** and **S6** (Supplemental note 4). These
results are summarized in **Figure 7**, which show the stability intervals over which the extracted height-maps

satisfy the criteria RMSE ≤ 1.5 and coverage ≥ 85%, for which the reconstruction can be considered of high
quality. Overall, the stability intervals for the two classification threshold parameters (T_A and T_{otsu}) are
narrower for specimens containing a surface of low signal intensity and low contrast, while still covering
about 1/4 of their respective width. The graphical user interface of Zellige allows the user to adjust T_A and
T_{otsu} interactively, making it intuitive to search for reasonable values. We found that $2 \leq \sigma_{xy} \leq 3$ and $\sigma_z = 1$
generally give high quality results for all tested specimens (**Figure 7A**). We therefore expect only little
adjustment to be required by the user on the smoothing parameters from their default permissive values
(set to $\sigma_{xy} = 1$ and $\sigma_z = 1$). Values of σ_z that are too large may lead to the merging of a surfaces with a nearby
structure of high contrast (surface or else), as it happens for the epithelium surface in the cochlea specimen,
which merges with the underlying mesenchyme signal when σ_z is greater than 2 **(Figure 4 and Figure S4).**
The effect is even more pronounced for surfaces of high inclination (**Figures 2, 6** and **Figures S2, S6**), or
presenting a particularly rough texture (**Figures 5** and **Figures S5**).

Regarding the control parameters of step 2 (the assembly step), for the four examples presented
except for example 1, the quality of the reconstructed surfaces is stable and high over the main part of their
respective intervals of variation, deteriorating occasionally only when extreme values ($\simeq 1$) are used for
these parameters (see **Figures S2-S6, Figure 7B** and Supplemental note 4). Example 1 poses particularly
stringent constraints on the control parameters of the reconstruction due to the requirement of
reconstructing the 4 surfaces of different contrast and texture present in this sample using the same set of
parameters.

Finally, the computation times for running Zellige on a given dataset ranged between a few seconds
and a minute on a standard PC computer (see **Figure S7** and Supplementary note 5), except in a few
exceptional cases corresponding to extreme values of the control parameters. As a safeguard, a stopping
criterion could be implemented so as to exit the run (declaring the current parameter values invalid) if the
surface assembly computation exceeds a user-prescribed duration.

Overall, the sensitivity analysis indicates that the surface extraction performed by Zellige is robust
to variations of the control parameters of step 1 (surface pixel selection step). In general, the reconstruction
is more sensitive to the amplitude threshold parameter (T_A), and the Otsu threshold parameter (T_{otsu})
should be kept sufficiently low for samples containing surfaces of intensity close to the background.
However, in some cases such as in our example 3, the opposite is true. Thus, the two threshold parameters
play somewhat complementary roles, and the possibility to adjust them independently is useful in practice
to be able to cover as many cases as possible. A smoothing along *xy* appears necessary to correctly
reconstruct the surfaces supported by a junctional mesh. Not surprisingly, best results are obtained when
the radius of the gaussian filter used for this (parameter σ_{xy}) is adapted to the mesh (or cell) size. Likewise,
a smoothing along z is beneficial, but the extent of this smoothing (parameter σ_z) should not be too large
to avoid causing the fusion of nearby surfaces. With a few exceptions, the values of the RMSE and coverage
show little sensitivity to the values of the parameters controlling step 2 (surface assembly step), at least
once putative surface pixels have been properly selected. In the presence of several surfaces of potentially
very different sizes, the parameter controlling the fraction of OSE sizes allowed for OSE seeds (parameter
T_{OSE1}) should be relatively large (≥ 0.5 or greater, *i.e.* allowing more than 50% of the largest OSE sizes for
seeds) to allow the extraction of a surface of small size (for example, to extract the surface n° 4 of
example 1, which covers less than 20% of the xy-field of view, T_{OSE1} must be larger than 0.6). Extreme values
(close to 1) for the connectivity rates (C1 and C2) are too stringent and lead to a drop in the coverage of the
reconstructed surfaces. To sum up, we see that the most critical parameters for a satisfactory extraction of
the different surfaces are those controlling step 1. In most cases the parameters controlling step 2 do not
need to be adjusted and can be fixed to their default reference values.

Conclusion.

We have developed Zellige, a new tool to extract multiple surfaces from 3D fluorescence microscopy images. Zellige automatically finds surfaces by first identifying putative pixels that are likely to belong to a biological surface, and second by assembling a surface through connection of adjacent pixels satisfying natural proximity constraints. By using Zellige on synthetic epithelium images we have shown that it accurately reconstructs a surface with excellent performances in terms of both the distance to the ground-truth height-map and the surface coverage (**Figure 2**). Zellige can deal with complex images containing multiple surfaces, with computation times not exceeding a few tens of seconds on a standard computer. Importantly, the user is not required to specify the number of surfaces to be extracted. In the *Drosophila* specimen (**Figure 3**), the software readily extracts the 4 surfaces of interest that could be identified. Since Zellige detects putative surface pixels in the first step by combining local and global thresholds, it can deal with images where the multiple surfaces display different features, such as in the mouse cochlear embryo (**Figure 3**). With this difficult dataset, we could also confirm Zellige's robustness against very low signal-to-noise levels. The constructive approach of surface region growing used by Zellige in its second step enables it to circumvent the surface smoothness requirement, that is classically assumed by other surface extraction tools. For instance, it could reconstruct the highly irregular surface of a bronchial tissue infected by SARS-CoV-2 (**Figure 5**).

The robustness and flexibility of Zellige come at a price, namely, the requirement to specify 12 parameters when running the surface extraction. However, the sensitivity analysis we performed shows that adjusting only 4 of these parameters is enough in practice to handle a wide range of image types. These parameters correspond to intuitive notions (*e.g.,* thresholding and smoothing), which makes Zellige particularly easy to use. The Fiji interface that we implemented to perform this adjustment should make Zellige even more user-friendly and effective for biological applications.

To our knowledge, Zellige is the only open-source tool that can extract an unspecified number of epithelial surfaces from a 3D volume, possibly larger than two. This unique feature is especially useful in complex images that could be processed only by specialized tools before. For instance, Zellige can extract surfaces with projections on the *xy* plane that completely overlap, such as the basal and apical epithelia in the organoid image of **Figure 6**. Previously, such an image could be processed only by tools that relied on segmenting a mesh around the object surface, such as MorphoGraphX or ImSAnE [16, 47].

The flexibility and robustness of Zellige should allow to considerably relax the constraints that were previously imposed on the sample preparation and the image acquisition steps by the subsequent analysis. Indeed, Zellige can accommodate any number of surfaces in the acquired volume, overlapping or not, and of different contrast features. Zellige also showed excellent robustness against image noise. This should make it particularly useful in imaging contexts that are not easily amenable to automated analysis, such as intravital imaging. Finally, it is worth noting that Zellige is a generalist and modular method. With some adaptation of the surface pixel selection step, it could be used with imaging modalities beyond the scope of this article, for instance, in extracting the irregular and noisy surfaces of biological objects imaged with 3D electron microscopy images.

Methods.

Implementation.

Zellige was devised with the goal of achieving accurate segmentation of multiple biological surfaces from 3D confocal images. Unlike other existing surface extraction tools, it makes no assumption on the number of surfaces to be extracted and does not require the surfaces of interest to be the structures of highest contrast in the image. Zellige is written in Java, relying on the ImgLib2 library [48] and is distributed as a Fiji plugin, with a graphical user interface (GUI) designed to allow users to quickly find a good set of extraction parameters for a given image.

Zellige extracts each surface present in the image in the form of a height-map (or z-map), that is, a mapping

$$z: (x,y) \to z(x,y),$$

which associates to each point (x,y) over which the surface projects, the z-coordinate of the unique pixel (x,y,z) belonging to the surface. Each extracted height-map is then used to produce a 2D projection of the 3D stack restricted to a small sub-volume (of user-selected width) centered around the corresponding surface. To achieve a robust extraction, Zellige proceeds in two algorithmic steps, which are only outlined below (see Supplementary note 1 for implementation details).

In the first step, or *surface pixel selection* step, a segmentation is applied to the 3D image to select pixels that likely belong to a surface of interest (**Figure 1 and Figure S1A**). These putative surface pixels are detected as local maxima of image intensity along the z-axis, after using two independent binary classifiers, one based on pixel contrast and the other one on pixel intensity. Five adjustable parameters control the selection step: two threshold parameters (T_A and T_{otsu}) control the strength of the binary classifiers applied on contrast and intensity, respectively, and three parameters (S_{min}, σ_{xy} and σ_z) control clean-up operations applied at the end of the classification (removal of small isolated spots, and local averaging along the *xy* plane and the z axis, respectively).

In the second step, or *surface assembly step*, an iterative algorithm is used to extract the height-maps of each of the surfaces present in the image (**Figure 1 and Figure S1A**). The assembly starts by grouping neighboring putative surface pixels together within each orthogonal (*xz* or *yz*) section of the 3D image, in order to form a set of building blocks referred to as *orthogonal surface elements* (OSEs). These building blocks are then used to assemble the surfaces, in a process analogous to jigsaw puzzles, where OSEs adjacent to the surface boundary are added if they match this boundary, and rejected otherwise, until no matching OSE can be found (**Figure S1**). In order to increase the robustness of the assembly step, Zellige applies it in two rounds, proceeding along different axes during the first and second rounds. Each round is controlled by 3 adjustable parameters: a threshold parameter ($0 \le T_{OSE} \le 1$) sets a minimum size for the building blocks that can be used as seeds to initiate the assembly of a surface; and two other parameters ($0 \le R \le 50$ and $0 \le C \le 1$) set the matching constraints used to accept or reject the addition of OSEs to a surface. The assembly step is thus controlled overall by 6 parameters, *i.e.* two groups of 3 parameters (T_{OSE1}, R_1, C_1) and (T_{OSE2}, R_2, C_2) controlling the first and second assembly rounds, respectively.

Finally, the height-maps of each of the reconstructed surfaces are used to obtain a corresponding 2D projection (**Figure 1**). In practice a maximum projection restricted to a subvolume of width δz (δz being a user-defined parameter) centered around the surface of interest is performed.

Availability of data and code.

The project homepage below contains the source code, installation instructions, and documentations.

Zellige

- Project name: Zellige.
- Project homepage: https://gitlab.pasteur.fr/ida-public/zellige-core
- URL for the Fiji plugin (Update > Manage update sites): https://sites.imagej.net/Zellige/
- Gitlab Branch: master
- Operating systems: Platform independent.
- Programming language: Java.
- Compiled in Java8
- Other requirements : Runs from Fiji [28].
- License: BSD 2
- Any restrictions to use by non-academics: None.

Scripts to create Phantoms

- Project name: Phantoms.
- Project homepage: https://doi.org/10.5281/zenodo.6414596
- Operating systems: Platform independent.
- Programming language: MATLAB.
- License: BSD 2
- Any restrictions to use by non-academics: None.

Data sets.

Image 3D stacks, ground-truth height map and height maps produced by Zellige are available on Zenodo under the CC-BY license: https://zenodo.org/communities/zellige/ [29, 33, 42, 44, 46].

Human bronchial epithelium imaging.

The data used in Figure 5 were taken from the recent study [43] to which we refer for the preparation and imaging of human bronchial epithelium cultures. In brief, MucilAirTM were purchased from Epithelix (Saint-Julien-en-Genevois, France) and cultured for at least 4 weeks to reconstruct a differentiated human bronchial epithelium *in vitro* and stained as previously described. Images of the cultures were acquired using an inverted Zeiss LSM 710 confocal microscope controlled by the ZEN pro 2.3 software. Z-stack images of whole-mount samples were acquired with a Zeiss Plan Apochromat 63x oil immersion lens (NA=1.4). The image used here was published in Robinot *et al.* [43] under the CC-BY-4 license.

Drosophila imaging.

Flies were raised at 25°C under standard conditions. Pupae were collected for imaging as described previously [49]. Ecad::GFP flies [50] were used for live imaging as previously described [1]. In brief, images were acquired with a spinning disk microscope from Gataca Systems driven by the MetaMorph software. The system is equipped with an inverted Nikon TI2E stand, a motorized XYZ stage, and a Nikon Plan Apo 60x oil immersion (NA=1.4) lens and with a Prime95B camera.

Cochlea imaging.

The inner ears from wild-type (C57BL/6) mice were rapidly dissected from temporal bones at embryonic stages E14.5 in HEPES-buffered (10 mM, pH 7.4) Hanks' balanced salts solution and fixed in 4% paraformaldehyde, 1 hour at room temperature. Specimen were permeabilized and stained for phalloidin-Atto 565 (Sigma) as previously described [39]. Fluorescence images were obtained with a swept-field confocal microscope (Opterra II) from Brucker. This system is equipped with a Nikon Plan Fluor 60x oil immersion lens (NA=1.4).

Biologists in this study hold a designer certificate of animal experimentation (level 1), allowing them to perform experimental work on animals in strict accordance with the European directive 2010/60/EU, and French regulations. The Ethics Committee of the Institut Pasteur (Comité d'Ethique en Experimentation Animale - CETEA) has approved this study with the project identifier dha170006. This approval is based on careful compliance to the 3Rs principle in the care and use of animals (Annex IV - 2010/60/EU).

Inner ear organoid imaging.

ESCs derived from blastocyst-stage embryos of R1 mice (mESCs) (ATCC, SCRC-1036) were maintained in feeder-free culture on 0.1% w/v gelatin (Sigma) coated substrates using LIF-2i medium as established previously [51]. The organoids were generated following the previously published protocol [45, 51]. Aggregates were harvested at day 14 and fixed in 4% v/v PFA (Electron Microscopy Sciences) overnight at 4°C. After blocking (PBS; 10% v/v normal goat serum; 0.1% v/v Triton X-100), the aggregates were stained for phalloidin Atto 565 (1:1000) (Sigma) overnight at RT on a shaker, and washed three times with PBS containing 0.1% v/v Triton X-100 for 1 h each at RT. Prior to imaging, the aggregates were incubated in a modified version of ScaleS solution containing 4 M Urea (Sigma), 40% w/v D-Sorbitol (Sigma), and 0.1% v/v Triton X-100, for 3-5 days to clarify the tissue. Finally, the aggregates were whole-mounted using the ScaleS solution and imaged using a confocal laser scanning microscope (A1R HD25, Nikon) equipped with a Nikon 25x silicon oil immersion lens (NA=1.05).

Acknowledgements.

We thank Maia Brünstein of the Hearing Institute Bioimaging Core Facility - C2RT/C2RA and Florian Rückerl of the Photonic BioImaging - C2RT/C2RA, for sharing their expertise on light microscopy. We thank Romain Levayer for sharing flies and lab space with us. We thank Gilles Trébeau (https://fr.linkedin.com/in/gilles-trebeau-427601212) for helpful discussions regarding the Java frontend deployment of Zellige. We thank Lisa Chakrabarti, Rémy Robinot and Vincent Michel for sharing data. This work was supported by *Fondation pour l'Audition* (FPA-IDA-STARTING-GRANT); *Institut Pasteur* (PTR#272); DIM ELICIT's grant from *Région Ile-de-France* (QP-IDF-DIM-ELICIT-2019) ; The French *Agence Nationale de la Recherche* (ANR-21-CE13-0038-013-01 and LabEx LIFESENSES ANR-10-LABX-65).

References.

1. Etournay R, Popović M, Merkel M, Nandi A, Blasse C, Aigouy B, et al. Interplay of cell dynamics and epithelial tension during morphogenesis of the Drosophila pupal wing. eLife. 2015;4:e07090. https://doi.org/10.7554/eLife.07090.

2. Guirao B, Rigaud SU, Bosveld F, Bailles A, López-Gay J, Ishihara S, et al. Unified quantitative characterization of epithelial tissue development. eLife. 2015;4:e08519. https://doi.org/10.7554/eLife.08519.

3. Etournay R, Merkel M, Popović M, Brandl H, Dye NA, Aigouy B, et al. TissueMiner: A multiscale

analysis toolkit to quantify how cellular processes create tissue dynamics. eLife. 2016;5. https://doi.org/10.7554/eLife.14334.

4. Saadaoui M, Rocancourt D, Roussel J, Corson F, Gros J. A tensile ring drives tissue flows to shape the gastrulating amniote embryo. Science. 2020;367:453–8. https://doi.org/10.1126/science.aaw1965.

5. Butler LC, Blanchard GB, Kabla AJ, Lawrence NJ, Welchman DP, Mahadevan L, et al. Cell shape changes indicate a role for extrinsic tensile forces in Drosophila germ-band extension. Nat Cell Biol. 2009;11:859–64. https://doi.org/10.1038/ncb1894.

6. Collinet C, Rauzi M, Lenne P-F, Lecuit T. Local and tissue-scale forces drive oriented junction growth during tissue extension. Nat Cell Biol. 2015;17:1247–58. https://doi.org/10.1038/ncb3226.

7. Kong D, Wolf F, Großhans J. Forces directing germ-band extension in Drosophila embryos. Mech Dev. 2017;144:11–22. https://doi.org/10.1016/j.mod.2016.12.001.

8. Villedieu A, Bosveld F, Bellaïche Y. Mechanical induction and competence in epithelial morphogenesis. Curr Opin Genet Dev. 2020;63:36–44. https://doi.org/10.1016/j.gde.2020.03.003.

9. Blasse C, Saalfeld S, Etournay R, Sagner A, Eaton S, Myers EW. PreMosa: Extracting 2D surfaces from 3D microscopy mosaics. Bioinforma Oxf Engl. 2017. https://doi.org/10.1093/bioinformatics/btx195. https://doi.org/10.1093/bioinformatics/btx195.

10. Basu S, Rexhepaj E, Spassky N, Genovesio A, Paulsen RR, Shihavuddin A. FastSME: Faster and Smoother Manifold Extraction from 3D Stack. In: 2018 IEEE/CVF Conference on Computer Vision and Pattern Recognition Workshops (CVPRW). 2018. p. 2362–23628. https://doi.org/10.1109/CVPRW.2018.00305.

11. Herbert S, Valon L, Mancini L, Dray N, Caldarelli P, Gros J, et al. LocalZProjector and DeProj: a toolbox for local 2D projection and accurate morphometrics of large 3D microscopy images. BMC Biol. 2021;19:136. https://doi.org/10.1186/s12915-021-01037-w.

12. Li K, Wu X, Chen DZ, Sonka M. Optimal Surface Segmentation in Volumetric Images-A Graph-Theoretic Approach. IEEE Trans Pattern Anal Mach Intell. 2006;28:119–34. https://doi.org/10.1109/TPAMI.2006.19.

13. Lombardot B. Min Cost Z Surface Projection Fiji. Java. 2017. https://imagej.net/Minimum_Cost_Z_surface_Projection.

14. Hall M, Frank E, Holmes G, Pfahringer B, Reutemann P, Witten IH. The WEKA data mining software: an update. ACM SIGKDD Explor Newsl. 2009;11:10–8. https://doi.org/10.1145/1656274.1656278.

15. Sommer C, Straehle C, Kothe U, Hamprecht FA. Ilastik: Interactive learning and segmentation toolkit. In: 2011 IEEE International Symposium on Biomedical Imaging: From Nano to Macro. 2011. p. 230–3. https://doi.org/10.1109/ISBI.2011.5872394.

16. Heemskerk I, Streichan SJ. Tissue cartography: compressing bio-image data by dimensional reduction. Nat Methods. 2015;12:1139–42. https://doi.org/10.1038/nmeth.3648.

17. Haertter D, Wang X, Fogerson S, Ramkumar N, Crawford J, Poss K, et al. DeepProjection: Rapid and structure-specific projections of tissue sheets embedded in 3D microscopy stacks using deep learnin. 2021. https://doi.org/10.1101/2021.11.17.468809.

18. Green JBA, Sharpe J. Positional information and reaction-diffusion: two big ideas in developmental biology combine. Development. 2015;142:1203–11. https://doi.org/10.1242/dev.114991.

19. Miller CJ, Davidson LA. The interplay between cell signalling and mechanics in developmental processes. Nat Rev Genet. 2013;14:733–44. https://doi.org/10.1038/nrg3513.

20. Levin M, Martyniuk CJ. The bioelectric code: An ancient computational medium for dynamic control of

growth and form. Biosystems. 2018;164:76–93. https://doi.org/10.1016/j.biosystems.2017.08.009.

21. Duclut C, Prost J, Jülicher F. Hydraulic and electric control of cell spheroids. Proc Natl Acad Sci. 2021;118. https://doi.org/10.1073/pnas.2021972118.

22. Ruiz-Herrero T, Alessandri K, Gurchenkov BV, Nassoy P, Mahadevan L. Organ size control via hydraulically gated oscillations. Development. 2017;144:4422–7. https://doi.org/10.1242/dev.153056.

23. Dumortier JG, Le Verge-Serandour M, Tortorelli AF, Mielke A, de Plater L, Turlier H, et al. Hydraulic fracturing and active coarsening position the lumen of the mouse blastocyst. Science. 2019;365:465–8. https://doi.org/10.1126/science.aaw7709.

24. Mammoto T, Mammoto A, Ingber DE. Mechanobiology and developmental control. Annu Rev Cell Dev Biol. 2013;29:27–61. https://doi.org/10.1146/annurev-cellbio-101512-122340.

25. Ladoux B, Mège R-M. Mechanobiology of collective cell behaviours. Nat Rev Mol Cell Biol. 2017;18:743–57. https://doi.org/10.1038/nrm.2017.98.

26. Närhi K. Embryonic Explant Culture: Studying Effects of Regulatory Molecules on Gene Expression in Craniofacial Tissues. In: Seymour GJ, Cullinan MP, Heng NCK, editors. Oral Biology: Molecular Techniques and Applications. New York, NY: Springer; 2017. p. 367–80. https://doi.org/10.1007/978-1-4939-6685-1_21.

27. Zeevaert K, Elsafi Mabrouk MH, Wagner W, Goetzke R. Cell Mechanics in Embryoid Bodies. Cells. 2020;9:2270. https://doi.org/10.3390/cells9102270.

28. Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, et al. Fiji: an open-source platform for biological-image analysis. Nat Methods. 2012;9:676–82. https://doi.org/10.1038/nmeth.2019.

29. Boutet de Monvel J, Trébeau C, Altay G, Tinevez J-Y, Etournay R. Zellige example dataset: synthetic image dataset. 2022. https://doi.org/10.5281/zenodo.6376584.

30. Jülicher F, Eaton S. Emergence of tissue shape changes from collective cell behaviours. Semin Cell Dev Biol. https://doi.org/10.1016/j.semcdb.2017.04.004. https://doi.org/10.1016/j.semcdb.2017.04.004.

31. Morata G, Lawrence P. An exciting period of Drosophila developmental biology: Of imaginal discs, clones, compartments, parasegments and homeotic genes". Dev Biol. 2022;484:12–21. https://doi.org/10.1016/j.ydbio.2022.01.008.

32. Ray RP, Matamoro-Vidal A, Ribeiro PS, Tapon N, Houle D, Salazar-Ciudad I, et al. Patterned Anchorage to the Apical Extracellular Matrix Defines Tissue Shape in the Developing Appendages of Drosophila. Dev Cell. 2015;34:310–22. https://doi.org/10.1016/j.devcel.2015.06.019.

33. Trébeau C, Boutet de Monvel J, Altay G, Tinevez J-Y, Etournay R. Zellige example dataset: Drosophila pupal wing and abdomen. 2022. https://doi.org/10.5281/zenodo.6376566.

34. Prat-Rojo C, Pouille P-A, Buceta J, Martin-Blanco E. Mechanical coordination is sufficient to promote tissue replacement during metamorphosis in Drosophila. EMBO J. 2020;39:e103594. https://doi.org/10.15252/embj.2019103594.

35. Zallen JA. Planar Polarity and Tissue Morphogenesis. Cell. 2007;129:1051–63. https://doi.org/10.1016/j.cell.2007.05.050.

36. Montcouquiol M, Rachel RA, Lanford PJ, Copeland NG, Jenkins NA, Kelley MW. Identification of Vangl2 and Scrb1 as planar polarity genes in mammals. Nature. 2003;423:173–7. https://doi.org/10.1038/nature01618.

37. Cohen R, Amir-Zilberstein L, Hersch M, Woland S, Loza O, Taiber S, et al. Mechanical forces drive ordered patterning of hair cells in the mammalian inner ear. Nat Commun. 2020;11:5137. https://doi.org/10.1038/s41467-020-18894-8.

38. McKenzie E, Krupin A, Kelley MW. Cellular growth and rearrangement during the development of the
mammalian organ of Corti. Dev Dyn. 2004;229:802–12. https://doi.org/10.1002/dvdy.10500.

39. Etournay R, Lepelletier L, Boutet de Monvel J, Michel V, Cayet N, Leibovici M, et al. Cochlear outer
hair cells undergo an apical circumference remodeling constrained by the hair bundle shape. Dev Camb Engl.
2010;137:1373–83. https://doi.org/10.1242/dev.045138.

40. Driver EC, Northrop A, Kelley MW. Cell migration, intercalation and growth regulate mammalian
cochlear extension. Dev Camb Engl. 2017;144:3766–76. https://doi.org/10.1242/dev.151761.

41. Qian D, Jones C, Rzadzinska A, Mark S, Zhang X, Steel KP, et al. Wnt5a functions in planar cell polarity
regulation in mice. Dev Biol. 2007;306:121–33. https://doi.org/10.1016/j.ydbio.2007.03.011.

42. Trébeau C, Boutet de Monvel J, Altay G, Tinevez J-Y, Etournay R. Zellige example dataset: mouse
embryonic cochlea stained for F-actin. 2022. https://doi.org/10.5281/zenodo.6376542.

43. Robinot R, Hubert M, de Melo GD, Lazarini F, Bruel T, Smith N, et al. SARS-CoV-2 infection induces
the dedifferentiation of multiciliated cells and impairs mucociliary clearance. Nat Commun. 2021;12:4354.
https://doi.org/10.1038/s41467-021-24521-x.

44. Robinot R, Chakrabarti L, Michel V, Trébeau C, Boutet de Monvel J, Altay G, et al. Zellige example
dataset: primary culture of human bronchial cells infected by SARS-CoV-2. 2022.
https://doi.org/10.5281/zenodo.6376594.

45. Koehler KR, Mikosz AM, Molosh AI, Patel D, Hashino E. Generation of inner ear sensory epithelia from
pluripotent stem cells in 3D culture. Nature. 2013;500:217–21. https://doi.org/10.1038/nature12298.

46. Altay G, Trébeau C, Boutet de Monvel J, Tinevez J-Y, Etournay R. Zellige example dataset: inner ear
organoid epithelium. 2022. https://doi.org/10.5281/zenodo.6376582.

47. Barbier de Reuille P, Routier-Kierzkowska A-L, Kierzkowski D, Bassel GW, Schüpbach T, Tauriello G,
et al. MorphoGraphX: A platform for quantifying morphogenesis in 4D. eLife. 2015;4:e05864.
https://doi.org/10.7554/eLife.05864.

48. Pietzsch T, Preibisch S, Tomančák P, Saalfeld S. ImgLib2—generic image processing in Java.
Bioinformatics. 2012;28:3009–11. https://doi.org/10.1093/bioinformatics/bts543.

49. Classen A-K, Aigouy B, Giangrande A, Eaton S. Imaging Drosophila Pupal Wing Morphogenesis. In:
Dahmann C, editor. Drosophila: Methods and Protocols. Totowa, NJ: Humana Press; 2008. p. 265–75.
https://doi.org/10.1007/978-1-59745-583-1_16.

50. Huang J, Zhou W, Dong W, Watson AM, Hong Y. Directed, efficient, and versatile modifications of the
Drosophila genome by genomic engineering. Proc Natl Acad Sci. 2009;106:8284–9.
https://doi.org/10.1073/pnas.0900641106.

51. Nie J, Koehler KR, Hashino E. Directed Differentiation of Mouse Embryonic Stem Cells Into Inner Ear
Sensory Epithelia in 3D Culture. In: Organ Regeneration. Humana Press, New York, NY; 2017. p. 67–83.
https://doi.org/10.1007/978-1-4939-6949-4_6.

Figure 1. Flowchart of Zellige's algorithmic steps. Surface pixel selection (step 1), surface assembly in the form of a height map (step 2), and subsequent projection localized to the height map, are schematically depicted in the case of a 3D image containing 4 surfaces of interest.

Figure 2. Multiple surface extraction on a synthetic 3D image. (A-B) The image contains 3 phantom surfaces (S1, S2, S3) of different shapes (sinusoidal, flat, and paraboloidal, respectively), and different textures (surface S1 has constant intensity, while surfaces S2 and S3 are supported by Voronoi meshes of different cell-sizes). **(C)** 3D representations of the height maps extracted by Zellige (in green) and of the ground truth (GT, in blue) height maps of surfaces S1, S2, and S3. **(D)** Error maps displaying the distance along the z-axis between the reconstructed and GT height maps for surfaces S1, S2, and S3. **(E)** Projections of the 3D image localized to the different surfaces S1-S3 (maximum intensity projections over a subvolume of a width δz=1 pixel above or below the corresponding height-maps). Upper and lower panels show the projections based on the GT and the reconstructed height-maps, respectively.

Figure 3. Fly specimen. (A,B) Volume rendering (A) and orthogonal sections (B) of a 3D image of fly embryo taken around 24h after puparium formation, covering a portion of the abdomen (showing histoblast cells and larval cells), and a portion of the developing wing. Scale bar 50 μm. Four surfaces of interest may be identified in the dataset (of dimensions 1200 × 1200 × 51 pixels): surfaces S1 and S2 are relatively close to one another and located within overlapping z-ranges ($8 \leq z \leq 50$ and $20 \leq z \leq 50$, respectively). Surfaces S3 and S4 (located in the z-ranges $42 \leq z \leq 50$ and $9 \leq z \leq 50$, respectively) are relatively far from each other and can nearly be separated by a plane. **(C)** 3D representations of the height maps extracted by Zellige (in green) and of the ground truth height maps (GT, in blue) of surfaces S1-S4. The reconstructed height-maps of all surfaces S1-S4 cover >93% of the area of the corresponding GT (cf. Figure S2 and Table S1). To reduce the staircase artifacts (more or less visible depending on the surface) due to the digitization of the GT and reconstructed height-maps, all height-maps were smoothed with a 2D gaussian filter with a standard radius of 5 pixels (cf. Supplemental note 1). **(D)** Error maps (color-coded distance along the z-axis between the reconstructed and the GT height-maps) plotted for each of the reconstructed surfaces. The large majority of pixels on the reconstructed height-maps (98%, 96%, 91%, and 99% for surfaces S1 to S4, respectively) display errors of <2 pixels. The height-maps of surfaces S1, S2, S4 show subpixel accuracy on average (RMSE < 1), while that of surface S3 is slightly less accurate (RMSE = 1.25). **(E)** Projections of the 3D image localized to the different surfaces S1-S4 (in this and all subsequent figures, these are maximum intensity projections over a subvolume of width $\delta z = \pm 1$ pixel above or below the corresponding height-maps). Upper and lower panels show the projections based on the GT and the reconstructed height-maps, respectively. Scale bar 50 μm.

Figure 4. Cochlea specimen. (A) Volume rendering of a 3D confocal swept-field image of the mouse cochlear embryo on embryonic day E14.5. The dataset (of dimensions 1024 × 1024 × 45 pixels) shows a portion of the sensory epithelium (at the topmost sections of the stack) and the underlying non-cellular layer of mesenchyme on which the organ develops. Both structures are stained with phalloidin to reveal F-actin. Scale bar 40 µm. The surface of interest is the epithelium surface, harboring the sensory and supporting cells under differentiation. The mesenchyme layer is not (strictly speaking) assimilable to a surface, but it produces a strong background signal nearby the surface of interest, hampering its extraction. **(B)** 3D representations of the height map extracted by Zellige (in green) and the GT height map (in blue), of the epithelium surface. **(C)** Color-coded error map of the reconstructed height-map, which shows subpixel accuracy (errors <1) over a large majority (83%) of pixels, as well as on average (RMSE ~ 1.1). **(D)** Projections localized to the GT height-map of the epithelium surface (left most panel), and to the height-maps extracted with the four different algorithms: FastSME, LocalZProjector, PreMosa, and Zellige. Only Zellige correctly extracts the surface of the epithelium in this example. Scale bar 40 µm.

Figure 5. Human bronchial epithelial cells infected by SARS-CoV-2. (A) Volume rendering and individual sections of a confocal 3D image of a primary culture of bronchial epithelial cells 4 days after it was infected by the SARS-CoV-2 virus. The dataset (of dimensions 1024 × 1024 × 15 pixels) covers a portion of the epithelium immunostained for the tight junction protein ZO-1. Notice the roughness of the epithelium surface and the presence of anomalous bulges (arrows) resulting from the SARS-CoV-2 infection. Scale bar 10 μm. **(B)** 3D representations of the height map extracted by Zellige (in green) and the GT height map (in blue), of the epithelium surface. **(C)** Color-coded error map of the reconstructed height-map. Despite its roughness, the surface of interest is reconstructed with subpixel accuracy over the majority (71%) of pixels, as well as on average (RMSE ~ 0.8). **(D)** Projections localized to the GT height-map of the epithelium surface (leftmost panel), and the height-maps extracted with the four different algorithms: FastSME, LocalZProjector, PreMosa, and Zellige. Scale bar 30 μm.

Figure 6. Organoid specimen. (A-B) Volume rendering (A) and orthogonal sections (B) of a confocal 3D image of a (half of) inner ear organoid, which has been fixed and stained with phalloidin to reveal F-actin. The dataset (of dimensions 520 × 465 × 35 pixels) includes two dome-shaped epithelial surfaces of interest, forming the apical (inward) and basal (outward) sides of the organoid. **(C)** 3D representations of the height map extracted by Zellige (in green) and the GT height map (in blue), of the epithelium surface. **(D)** Color-coded error maps of the reconstructed height-maps for the apical (left) and basal (right) epithelial surfaces of the organoid. The surfaces of interest are reconstructed with an error of < 2 pixels over a large majority (96% and 93% for the apical and basal surfaces, respectively) of pixels, as well as on average (RMSE ~ 0.8 and 1.1 for the apical and the basal surfaces, respectively). **(E)** Projections localized to the GT height-maps of the epithelium surface (panels on the left), and the height-maps extracted by Zellige (panels on the right). Scale bar 100 μm

Figure 7. Summary of the sensitivity analysis. The intervals indicated in grey for each parameter and each of the images tested correspond to the parameter values for which the reconstruction satisfies high quality criteria defined by RMSE ≤ 1.5 and coverage ≥ 85%. Black marks indicate the reference value obtained by manual adjustment for each image (cf. Supplemental note 2). **(A)** Parameters of the surface selection step. **(B)** Parameters of the surface assembly step.

FIGURE 1
Flowchart of Zellige's algorithmic steps. Surface pixel selection (step 1), surface assembly in the form of a height map (step 2), and subsequent projection localized to the height map, are schematically depicted in the case of a 3D image containing 4 surfaces of interest.

FIGURE 2
Multiple surface extraction on a synthetic 3D image. (A) The image contains 3 phantom surfaces (S1, S2, S3) of different shapes (sinusoidal, flat, and paraboloidal, respectively), and different textures (surface S1 has constant intensity, while surfaces S2 and S3 are supported by Voronoi meshes of different cell-sizes). (B) 3D representations of the height maps extracted by Zellige (in green) and of the ground truth (GT, in blue) height maps of surfaces S1, S2, and S3. (C) Error maps displaying the distance along the z-axis between the reconstructed and GT height maps for surfaces S1, S2, and S3. (E) Projections of the 3D image localized to the different surfaces S1-S3 (maximum intensity projections over a subvolume of a width $\delta z=1$ pixel above or below the corresponding height-maps). Upper and lower panels show the projections based on the GT and the reconstructed height-maps, respectively.

FIGURE 3
(A,B) Volume rendering (A) and orthogonal sections (B) of a 3D image of fly embryo taken around 24h after puparium formation, covering a portion of the abdomen (showing histoblast cells and larval cells), and a portion of the developing wing. Scale bar 50 μm. Four surfaces of interest may be identified in the dataset (of dimensions 1200 × 1200 × 51 pixels): surfaces S1 and S2 are relatively close to one another and located within overlapping z-ranges ($8 \leq z \leq 50$ and $20 \leq z \leq 50$, respectively). Surfaces S3 and S4 (located in the z-ranges $42 \leq z \leq 50$ and $9 \leq z \leq 50$, respectively) are relatively far from each other and can nearly be separated by a plane.
(C) 3D representations of the height maps extracted by Zellige (in green) and of the ground truth height maps (GT, in blue) of surfaces S1-S4. The reconstructed height-maps of all surfaces S1-S4 cover >93% of the area of the corresponding GT (cf. Figure S2 and **Table S1**). To reduce the staircase artifacts (more or less visible depending on the surface) due to the digitization of the GT and reconstructed height-maps, all height-maps were smoothed with a 2D gaussian filter with a standard radius of 5 pixels (cf. Supplemental note 1).
(D) Error maps (color-coded distance along the z-axis between the reconstructed and the GT height-maps) plotted for each of the reconstructed surfaces. The large majority of pixels on the reconstructed height-maps (98%, 96%, 91%, and 99% for surfaces S1 to S4, respectively) display errors of <2 pixels. The height-maps of surfaces S1, S2, S4 show subpixel accuracy on average (RMSE < 1), while that of surface S3 is slightly less accurate (RMSE = 1.25).
(E) Projections of the 3D image localized to the different surfaces S1-S4 (in this and all subsequent figures, these are maximum intensity projections over a subvolume of width $\delta z= \pm 1$ pixel above or below the corresponding height-maps). Upper and lower panels show the projections based on the GT and the reconstructed height-maps, respectively.

FIGURE 4
(A) Volume rendering of a 3D confocal swept-field image of the mouse cochlear embryo on embryonic day E14.5. The dataset (of dimensions 1024 × 1024 × 45 pixels) shows a portion of

the sensory epithelium (at the topmost sections of the stack) and the underlying non-cellular layer of mesenchyme on which the organ develops. Both structures are stained with phalloidin to reveal F-actin. Scale bar 40 µm. The surface of interest is the epithelium surface, harboring the sensory and supporting cells under differentiation. The mesenchyme layer is not (strictly speaking) assimilable to a surface, but it produces a strong background signal nearby the surface of interest, hampering its extraction.

(B) 3D representations of the height map extracted by Zellige (in green) and the GT height map (in blue), of the epithelium surface.

(C) Color-coded error map of the reconstructed height-map, which shows subpixel accuracy (errors <1) over a large majority (83%) of pixels, as well as on average (RMSE ~ 1.1).

(D) Projections localized to the GT height-map of the epithelium surface (left most panel), and to the height-maps extracted with the four different algorithms: FastSME, LocalZProjector, PreMosa, and Zellige. Only Zellige correctly extracts the surface of the epithelium in this example.

FIGURE 5

(A) Volume rendering and individual sections of a confocal 3D image of a primary culture of bronchial epithelial cells 4 days after it was infected by the SARS-CoV-2 virus. The dataset (of dimensions 1024 × 1024 × 15 pixels) covers a portion of the epithelium immunostained for the tight junction protein ZO-1. Notice the roughness of the epithelium surface and the presence of anomalous bulges (arrows) resulting from the SARS-CoV-2 infection. Scale bar 10 µm.

(B) 3D representations of the height map extracted by Zellige (in green) and the GT height map (in blue), of the epithelium surface.

(C) Color-coded error map of the reconstructed height-map. Despite its roughness, the surface of interest is reconstructed with subpixel accuracy over the majority (71%) of pixels, as well as on average (RMSE ~ 0.81).

(D) Projections localized to the GT height-map of the epithelium surface (leftmost panel), and the height-maps extracted with the four different algorithms: FastSME, LocalZProjector, PreMosa, and Zellige. Scale bar 30 µm

FIGURE 6

(A-B) Volume rendering (A) and orthogonal sections (B) of a confocal 3D image of a (half of) inner ear organoid, which has been fixed and stained with phalloidin to reveal F-actin. The dataset (of dimensions 520 × 465 × 35 pixels) includes two dome-shaped epithelial surfaces of interest, forming the apical (inward) and basal (outward) sides of the organoid.

(C) 3D representations of the height map extracted by Zellige (in green) and the GT height map (in blue), of the epithelium surface.

(D) Color-coded error maps of the reconstructed height-maps for the apical (left) and basal (right) epithelial surfaces of the organoid. The surfaces of interest are reconstructed with an error of < 2 pixels over a large majority (96% and 93% for the apical and basal surfaces, respectively) of pixels, as well as on average (RMSE ~ 0.8 and 1.1 for the apical and the basal surfaces, respectively).

(E) Projections localized to the GT height-maps of the epithelium surface (panels on the left), and the height-maps extracted by Zellige (panels on the right). Scale bar 100 μm.

FIGURE 7
Summary of the sensitivity analysis. The intervals indicated in grey for each parameter and each of the images tested correspond to the parameter values for which the reconstruction satisfies high quality criteria defined by RMSE ≤ 1.5 and coverage ≥ 85%. Black marks indicate the reference value obtained by manual adjustment for each image (cf. Supplemental note 2). (A) Parameters of the surface selection step. (B) Parameters of the surface assembly step.