<u>**Title**</u>

Identification of Antimalarial Compounds that Inhibit Apicomplexan AP2 Proteins in the

Human Malaria Parasite *Plasmodium falciparum*

Word Count: 9239

<u>**Authors**</u>

Timothy Russell[1,2,3,4], Erandi K. De Silva[6], Valerie Crowley[1], Kathryn Shaw-Saliba[7], Namita
Dube[1], Gabrielle Josling[1,3,4], Charisse Flerida A. Pasaje[8], Irene Kouskoumvekaki[9], Gianni
Panagiotou[10,11], Jacquin C. Niles[8], Marcelo Jacobs-Lorena[7], C. Denise Okafor[1,5], Francisco-
Javier Gamo[12], Manuel Llinás[1,2,3,4,5]

1.   Department of Biochemistry and Molecular Biology, Pennsylvania State University.

2.   Huck Institutes Center for Eukaryotic Gene Regulation (CEGR), Pennsylvania State
University.

3.   Huck Institutes Center for Malaria Research (CMaR), Pennsylvania State University.

4.   Huck Institutes Center for Infectious Disease Dynamics, Pennsylvania State University.

5.   Department of Chemistry, Pennsylvania State University.

6.   Lewis-Singler Institute for Integrative Genomics, Princeton University.

7.   Department of Molecular Biology and Immunology, Malaria Research Institute, Johns
Hopkins Bloomberg School of Public Health.

8.   Department of Biological Engineering, Massachusetts Institute of Technology.

9.   Department of Systems Biology, Technical University of Denmark, Kgs. Lyngby,
Denmark.

10. Systems Biology and Bioinformatics, Leibniz Institute for Natural Products Research and
Infection Biology, Hans Knöll Institute.

11. Department of Medicine, the University of Hong Kong, Hong Kong SAR, China

12. Global Health Medicines, GlaxoSmithKline, Tres Cantos, Spain.

<u>**Abstract**</u>

Plasmodium parasites are reliant on the Apicomplexan AP2 (ApiAP2) transcription

factor family to regulate gene expression programs. AP2 DNA binding domains have no

homologs in the human or mosquito host genomes, making them potential antimalarial

drug targets. Using an *in-silico* screen to dock thousands of small molecules into the

crystal structure of the AP2-EXP (Pf3D7_1466400) AP2 domain (PDB:3IGM), we

identified compounds that interact with this domain. Four compounds were found to

compete for DNA binding with AP2-EXP and at least one additional ApiAP2 protein. Our

top ApiAP2 competitor compound perturbs the transcriptome of *P. falciparum*

trophozoites and results in a decrease in abundance of log_2 fold change > 2 for 50%

(46/93) of AP2-EXP target genes.   Additionally, two ApiAP2 competitor compounds

have multi-stage anti-*Plasmodium* activity against blood and mosquito stage parasites.

In summary, we describe a novel set of antimalarial compounds that are targeted

against the ApiAP2 family of proteins. These compounds may be used for future

chemical genetic interrogation of ApiAP2 proteins or serve as starting points for a new

class of antimalarial therapeutics.

<u>Author Summary</u>

Plasmodium parasites are the causative agent of malaria, which resulted in over

600,000 deaths in 2021. Due to resistance arising for every antimalarial therapeutic

deployed to date, new drug targets and druggable pathways must be explored. To

address this concern, we used a molecular docking screen to predict competitors of

DNA binding by the parasite specific family of Apicomplexan AP2 (ApiAP2) transcription

factor proteins for testing *in vitro* and *in vivo*. We find that ApiAP2 competing

compounds have antimalarial activity consistent with the disruption of gene regulation.

This work will further our understanding of both the biological role and targetability of

parasite transcriptional regulation.

<u>Introduction</u>

Malaria is a disease caused by intracellular parasites from the genus *Plasmodium* that

represents a significant health and economic burden worldwide[1]. The most virulent of

the human infectious malaria parasites is *Plasmodium falciparum*, which caused more

than 200 million cases of malaria and resulted in over 600 thousand deaths in 2021[1].

Resistance has been reported for every antimalarial therapeutic deployed to date,

necessitating the need for malaria drugs that target new parasite processes[2] . To

successfully proliferate, *P. falciparum* parasites must develop through a complex

lifecycle that includes intracellular and extracellular stages in both the human and

Anopheles mosquito hosts[3]. The clinical symptoms of malaria are caused by the

intraerythrocytic development cycle (IDC), a 48-hour cyclic asexual proliferation that

results in the destruction of red blood cells (RBCs). During the asexual IDC, malaria

parasites progress through three morphological stages referred to as ring, trophozoite,

and schizont. Parasites transmit from human to mosquito following differentiation and

maturation into sexual stage gametocytes. Once ingested by the mosquito, gametes

sexually reproduce and ultimately develop into sporozoites, which can be transmitted

back to the human host to initiate a liver stage infection which precedes the asexual

blood stages[3].

Up to 80% of *P. falciparum* protein coding transcripts are developmentally regulated

during the IDC(4–8) as part of a 'just in time' cascade(5, 8, 9). This is predicted to be

principally driven by the 27 Apicomplexan Apetala AP2 (ApiAP2) proteins(10–12), which

are the major family of sequence specific transcription factors encoded in the

Plasmodium genome. ApiAP2 proteins contain one to three AP2 DNA binding domains

that are analogous to the plant APETALA2/Ethylene Responsive Factor (AP2/ERF)

domain and therefore have no homologs in the human or mosquito genome(13, 14).

Over half of the ApiAP2 proteins are predicted to be essential for the IDC in *P.*

falciparum(15–17). In-depth studies of ApiAP2 proteins to date have demonstrated roles

in transcriptional activation or repression affecting diverse processes including

invasion(18, 19), heterochromatin maintenance(17, 20–23), sexual and mosquito stage

differentiation(16, 24–34), and heat stress tolerance(35). Despite these properties which

make ApiAP2 proteins potential drug targets, no efforts to date have focused on

targeting the sequence specific DNA binding ApiAP2 transcription factors.

In this study we conducted an *in silico* chemical screen against the crystal structure of

the AP2-EXP AP2 domain(36) to predict competitors of DNA binding activity. AP2-EXP

is predicted to be essential for the *P. falciparum* asexual blood stage,(16, 17, 21) and its

orthologue in the rodent malaria parasite *P. berghei* (PbAP2-Sp) is a master regulator of

sporogony(32). Several compounds identified in the screen can compete for DNA

binding activity with the AP2-EXP AP2 domain *in vitro*. We also find that modifications to

the substitution state of an AP2 domain competitor compound can modulate its ability to

compete DNA binding. Our lead compound kills *P. falciparum* parasites in concordance

with the stage specific expression of AP2-EXP. We measured AP2-EXP genomic

occupancy by Chromatin immunoprecipitation followed by high throughput sequencing

(ChIP-seq) and found high correlation between dysregulated mRNA transcripts in the

presence of our lead compound and AP2-EXP gene targets, corroborating the

hypothesis that AP2-EXP is inhibited *in vivo*. Finally, two AP2 domain competitor

compounds are active against *P. berghei* development in the mosquito stage. Overall,

our results demonstrate the potential to chemically compete the DNA binding activity of

ApiAP2 proteins.

Results

***In silico* Prediction of AP2-EXP Competitors**

To identify competitors of DNA binding by an ApiAP2 protein, we used AP2-EXP

(PF3D7_1466400), which is the only ApiAP2 protein whose AP2 domain is structurally

characterized (PDB accession: 3IGM)(36). *In silico* molecular docking was run on

AUTODOCK(37) using over ten thousand small molecules from both the Tres Cantos

Antimalarial Set (TCAMS)(38) and the Drug Bank(39) against AP2-EXP. Docking hits

were prioritized based on the free energy of interaction and proximity to amino acids

necessary for DNA contact **(Fig S1A)**. In some cases, the compound that generated a

high docking score was not available for purchase, so alternative choices with >0.9

Tanimoto similarity score were identified, computationally docked, and used for further

evaluation **(All compounds listed in Figure S1B).** These docking simulations resulted

in a final list of nine top scoring predicted competitors of DNA binding by AP2-EXP.

Each compound was assigned an alphabetical identifier A-I **(Fig 1A, Fig S2A, B)** for

use in this study. We determined that all nine compounds kill asexual *P. falciparum*

parasites at micromolar concentrations (range 11-170µM) in a growth inhibition assay

(Fig 1A, Fig S3).

<u>Four Compounds Have Activity Against AP2-EXP *in vitro*</u>

The nine predicted competitors of DNA binding by AP2-EXP were tested in a

competitive electrophoretic mobility shift assay (EMSA). The minimum mass of AP2-

EXP required to visualize DNA binding **(Fig S4A)** was mixed with each compound and

incubated prior to adding the DNA oligonucleotide **(Table S1)**. Four of the nine

compounds (A, B, C and I) **(Fig 1B)** were able to effectively compete for DNA binding

by AP2-EXP *in vitro*. Three of the four AP2 domain competitors (Compounds A, B and

C) have a benzoxazole core moiety, while Compound I is made up of planar rings **(Fig

1C)**.

To assess whether DNA binding competition was specific to the AP2-EXP AP2 domain,

we repeated competitive EMSAs with three different purified *P. falciparum* AP2

domains: AP2-I Domain 3 (AP2-I D3)(12, 18) **(Fig S4B)**, AP2-HS Domain 1 (AP2-HS

D1)(12, 35) **(Fig S4C)**, and PfSIP2 Domain 1 (PfSIP2 D1)(20) **(Fig S4D)**, again using

the minimum amount of protein needed to visualize interaction with the DNA

oligonucleotide **(Table S1)**. We found that Compounds A, B, C, and I can compete DNA

binding by AP2-I **(Fig S5A)**, Compounds B and I compete DNA binding by AP2-HS D1

(Fig S5B) and Compounds B, C and I compete DNA binding by PfSIP2 D1 **(Fig S5C)**.

As a control for compound specificity, we performed competitive EMSAs using two non-

Plasmodium AP2 domain proteins: the *Arabidopsis thaliana* Ethylene Response Factor

1(40) (AtERF1) **(Fig S4E)** and the human High Mobility Group Box (HMGB) domain

protein SOX2 **(Fig S4F)**(41). Only Compound I was able to compete AtERF1 **(Fig S6A)**,

while SOX2 was competed by Compound A only **(Fig S6B)**. Therefore, Compounds B

and C demonstrate specificity for Apicomplexan AP2 domains *in vitro*. (EMSA Results

Summarized in **Table S2**).

To eliminate the possibility that ApiAP2 competing compounds interfere non-specifically

with DNA binding proteins by intercalating with DNA, we conducted an ethidium

bromide exclusion assay(42). Compared to the positive control DRAQ5(42), only

Compound I was able to effectively displace ethidium bromide from DNA **(Fig S7)**.

Based on these results, we focused on Compounds B and C for further study.

<u>Structural Analogs of Compound B Differ in DNA Binding Competition Activity</u>

We tested the four closest available analogs to Compound B from the TCAMS based on

similarity score(38) (Designated as Compounds B-1, B-2, B-3 and B-4) for competition

against AP2-EXP. Compounds B-1 and B-4 compete DNA binding by AP2-EXP **(Fig**

2A) and AP2-I D3 **(Fig S8A)**, while Compounds B-2 and B-3 do not compete DNA

binding by either AP2-EXP or AP2-I D3 **(Fig 2A, S8A)**. Compound B-1 is the least

effective competitor relative to Compounds B and B-4 **(Fig 2B)**. Non-AP2 domain

competitor Compounds B-2 and B-3 each have halogen atoms substituted on the

benzene ring. Conversely, Compounds B, B-1, and B-4, have methyl or ethyl groups

substituted onto the benzene ring and can compete DNA binding **(Fig 2C)**. Compound

B-1 has both a methyl group and a chlorine atom substitution. If the halogen atom

substitution decreases DNA binding competition, this mix of substitutions is consistent

with its lower efficacy of DNA binding competition relative to Compounds B and B-4.

Compounds B, B-1 and B-4 have no difference in DNA intercalation ability measured by

ethidium bromide exclusion **(Fig S8B)**. A molecular dynamics simulation of AP2-EXP

docked with Compound B and each analogue predicts that Compounds B, B-1, B-2, and

B-4 dock stably with AP2-EXP, while non-AP2 competitor Compound B-3 moves away

from the protein **(Supplemental Movies 1-5)**. In summary, substituting bulky,

electronegative groups onto Compound B can abolish competition for DNA binding with

AP2-EXP.

<u>Compound C Activity Coincides With AP2-EXP Expression</u>

To determine the stage in the IDC at which AP2-EXP is expressed, we used Selection

Linked Integration (SLI)(43) to create a parasite line with endogenously GFP tagged

AP2-EXP (AP2-EXP::GFP) **(Fig S9A)**. We observed maximum abundance of AP2-

EXP::GFP in 30 hpi trophozoites by both western blot and live fluorescence microscopy

(Fig 3A, Fig S10, S11). Maximum protein abundance of AP2-EXP in mid-trophozoites

was independently confirmed in a parasite line expressing AP2-EXP tagged with 2XHA

and the TetR:DOZI mRNA repression aptamer (AP2-EXP::HA) **(Fig S9B, Fig S12)**.

AP2-EXP was not detected in maturing Stage III gametocytes **(Fig 3A)**.

To assess whether Compounds B and C are likely to share a common ApiAP2 target,

we determined their parasite killing phenotypes against *P. falciparum* during the IDC.

Compound H was used as a control since it does not compete AP2-EXP **(Fig 1B)**. Each

compound was spiked (40µM) into synchronous asexual blood stage *P. falciparum*

parasites at 24 hpi and parasite morphology was determined at 24 and 48 hours post

spike-in **(Fig S13A)**. At 24 hours post spike-in, parasites in the presence of either

Compounds B or C had failed to progress past the mid-trophozoite stage and did not

morphologically progress any further by 48 hours post spike-in **(Fig 3B)**. In contrast,

parasites spiked with Compound H progressed through one IDC before failing to

reinvade in the next cycle **(Fig 3B)**. As expected, each compound exposure prevented

further parasite growth within 48 hours of spike-in **(Fig S13B)**.

Due to its commercial availability, we continued with Compound C as the lead

compound for phenotypic and molecular characterization. To determine the precise

timing of Compound C antimalarial activity, synchronous parasites were monitored

throughout the 48-hour IDC in the presence of either 40µM Compound C or DMSO

control after spiking in Compound C at 6 hpi. Compound C spiked parasites failed to

progress beyond the mid-trophozoite stage at 24-30 hours post invasion **(Fig S14)**. To

further narrow down the timing of the anti-*Plasmodium* action of Compound C, 40µM

Compound C was washed out of a parasite culture at 8-hour intervals beginning at 14

hpi. Growth was rescued, albeit with diminishing effectiveness, until 30 hpi **(Fig 3C)**.

Spiking 40µM Compound C into culture at 8-hour intervals beginning at 14 hpi

effectively killed the parasite culture at each time point except for 46 hpi. **(Fig 3D)**.

Taken together, these results suggest that Compound C maximally inhibits the IDC

progression of *Plasmodium* parasites between 30-46 hpi. These results show that the

maximum abundance of AP2-EXP at 30 hpi coincides with the timing of action for

Compound C.

Compound C Disrupts the _P. falciparum_ Transcriptome with Stage Specificity

Since AP2-EXP is predicted to act as a transcription factor[17, 21] we hypothesized that

changes in parasite transcript abundance should occur when AP2-EXP DNA binding is

competed by Compound C. We cultured parasites in 12µM ($0.66xIC_{50}$) Compound C or

DMSO vehicle and measured RNA abundance at seven time points during the asexual

blood stage. As a quality control for normal IDC progression, we found high correlation

for a set of "control" genes[44, 45] with an established 48-hour periodicity in the IDC

between our DMSO and Compound C samples **(Fig S15A-D)**. Since 12µM Compound

C kills fewer than 50% of parasites within one IDC **(Fig S16A, B)**, this demonstrates

that there is not an overall perturbation to the IDC transcriptome upon Compound C

incubation. The Spearman Correlation between total transcriptome samples deviated

most significantly at 30 hpi, (Corr. = -0.16) **(Fig 4A)** as expected from the compound

activity measured **(Fig 3C, D, Gene Expression values in Table S3)**.

Next, we quantified differences in transcript abundance that are unique to the

progression from 24 to 30 hpi for Compound C vs. DMSO vehicle control populations

using Linear Models for Microarray Data (LIMMA[46]) **(Differentially Expressed genes**

in Table S3). Overall, 463 RNA transcripts are decreased in abundance by log_2 fold

change >2 between 24-30 hpi in the presence of Compound C. Gene ontology (GO)

analysis[47] **(Full GO Terms in Table S3)** revealed an enrichment for genes that

encode proteins important for the parasite to invade or modify red blood cells among

decreased transcripts (e.g., RON3, GAP45, RhopH2, RhopH3, MSP1, MSP6). We

independently determined differential transcript abundance over the entire time course

using the RNA time course specific software Rnits(48) and found high overlap with our

LIMMA analysis **(Overlaps in Table S3)**. Therefore, the maximal perturbation of the

transcriptome **(Fig 4A)**, action of Compound C **(Fig 3C, D)**, and maximal abundance of

AP2-EXP **(Fig 3A)** all occur at roughly 30 hpi in the IDC.

AP2-EXP Gene Targets Correlate with Differentially Abundant Transcripts

We used ChIP-seq to determine the genome wide binding occupancy of AP2-EXP. A

total of three samples from highly synchronous trophozoites were collected at 30 hpi

using both the AP2-EXP::GFP (2 replicates) and AP2-EXP::HA (1 replicate) parasite

lines **(Peaks in Table S4)**. As a quality control, we determined that ChIP recovers intact

AP2-EXP **(Fig S17A)**, and a co-immunoprecipitation blot confirmed that AP2-EXP

interacts with Histone H3 in the nucleus **(Fig S17B)**. In aggregate, AP2-EXP binds 240

genomic loci, corresponding to 101 total target genes **(Genes in Table S4)**. Enrichment

of AP2-EXP was well conserved between the GFP and HA tagged parasite lines, as

indicated by nearly identical coverage in a metagene plot of AP2-EXP target genes **(Fig

4B)**. A 'no epitope control' ChIP done on a wild type Pf3D7 parasite line with the anti-

GFP antibody detected only 2 peaks, neither of which overlapped with those of AP2-

EXP::GFP or AP2-EXP::HA **(Figure 4B, Table S4)**. AP2-EXP peaks were highly

enriched for the known AP2-EXP and PbAP2-Sp DNA sequence motif CATGCA(17, 49)

(Fig 4B, Fig S18A-C).

Out of the 463 transcripts decreased in abundance by $\log_2$ fold change > 2 in the

presence of Compound C, 46 are AP2-EXP targets **(Table S3)**. This represents 50%

(46/93) of AP2-EXP target genes detected in the RNA time course **(Fig 4C, Figure**

S19). In general, the transcripts with the greatest decrease in abundance are AP2-EXP

targets. We found that 7/14 total transcripts decreased in abundance by $\log_2$ fold

change > 5, and 16/33 decreased in abundance by $\log_2$ fold change > 4, are AP2-EXP

targets **(Fig 4C, Table S3)**. AP2-EXP targets that are dysregulated at 24-30 hpi have

functions related to red blood cell invasion and host remodeling **(Fig 4D, Table S3)**. To

assess whether AP2-EXP DNA binding is impacted by Compound C, parasite cultures

were spiked with either 40μM Compound C or DMSO vehicle control at 30 hpi and AP2-

EXP occupancy was measured by ChIP-quantitative PCR (ChIP-qPCR). AP2-EXP DNA

binding decreases at five specific peaks of occupancy (GAP45, SIP2, RON3, RALP,

and AMAI) in the presence of Compound C **(Fig S20)**.

We then compared AP2-EXP genomic occupancy with several published datasets in

order to further evaluate its function as a sequence specific transcription factor. AP2-

EXP occupancy is correlated with a nucleosome depleted region **(Fig S21A, B)**(50) and

activating chromatin marks(51, 52) **(Fig S22A, B)**, and is proximal to target gene

transcription start sites (TSS)(53) **(Fig S23)**. The majority of target genes increase in

abundance starting at 32 hpi **(Fig S24)**(54). AP2-EXP target gene function is enriched

for invasion and red blood cell modification, **(Table S3)** as was found for the genes that

decrease in abundance in the presence of Compound C **(Fig 4D, Table S3)**. In

aggregate, these findings suggest that Compound C inhibits the function of AP2-EXP as

a sequence specific transcription factor.

ApiAP2 Competitor Compounds B and C are Active Against Mosquito Stage

***Plasmodium* Parasites**

Elucidation of the role of essential *P. falciparum* ApiAP2 proteins in the IDC is

challenging due to the inability to knock out asexual blood stage essential genes. We

did not recover transgenic parasites after attempting to completely disrupt the coding

sequence of AP2-EXP using the targeted gene disruption (pSLI-TGD) system(43) in

three independent attempts, suggesting that AP2-EXP is essential for the IDC (not

shown). We then attempted to disrupt AP2-EXP protein abundance with the conditional

knockdown approaches Knock Sideways(43), TetR:DOZI mRNA repression(55), and

glms ribozyme mediated cleavage(56). Each of these genetic systems failed to mediate

protein knockdown or mislocalization **(Fig S9A, B, Fig S25, Fig S26)**. To account for

this limitation, we tested the ApiAP2 competitor Compounds B and C against rodent

infectious *Plasmodium berghei* parasites due to the well-characterized genetic

phenotype of the AP2-EXP orthologue PbAP2-Sp (89% amino acid identity to the AP2-

EXP AP2 domain)(32, 49) as a master regulator of sporogenesis.

We injected the measured IC_{50} **(Figure 1A)** of Compounds B, C, or F into the midgut of

Anopheles mosquitoes infected with *P. berghei* parasites to determine their effect on

Plasmodium mosquito stage development **(Fig S27)**. Compound F **(Fig 1B)** was used

as a non AP2 domain competing control. Mosquitoes injected with Compounds F or C

developed comparable oocyst numbers to the DMSO control **(Fig 6A)** while Compound

B prevented the development of midgut oocysts entirely **(Fig 6A)**. Compound C treated

mosquitoes developed 10-fold fewer midgut sporozoites than the control **(Fig 6B)**.

Therefore, consistent with their hypothesized *in vitro* activity against ApiAP2 proteins,

Compounds B and C are strong inhibitors of *P. berghei* sporozoite development *in vivo*

(parasite counts in **Table S5)**.

<u>Discussion</u>

ApiAP2 transcription factors are unique to Apicomplexan parasites due to their plant-like

domain architecture(13, 14) and many are essential to asexual blood stage

development, making them valuable as potential drug targets. It is therefore desirable to

discover chemical scaffolds that can target essential ApiAP2 proteins. Furthermore,

chemical inhibition of ApiAP2 proteins may be used in future studies to uncover details

about their true biological functions during parasite development. Using a combination

of *in silico*, biochemical, and genetic approaches, we identified a series of compounds

(Compounds A, B, C, and I) that compete DNA binding by ApiAP2 proteins. Compounds

A, B, and C all have the same core benzoxazole moiety but vary in their measured IC_{50},

indicating that changes which do not affect ApiAP2 DNA binding competition can alter

the potency against parasites. Furthermore, Compounds B and C can inhibit *P. berghei*

parasite development without killing mosquitoes, suggesting that ApiAP2 competitor

compounds may also be tolerated by the host and may serve as transmission blocking

agents(57).

Compounds B and C have no activity against the plant encoded AtERF1 or human

encoded SOX2, supporting their selectivity for Apicomplexan AP2 domains. Although

the mode of DNA binding is shared between AP2-EXP and AtERF1(40), the specific

amino acids that contact DNA are not strictly conserved(36). Lack of activity against

AtERF1 may be the result of filtering molecular docking hits based on proximity to DNA

base contacting amino acids. The DNA binding competition capacity of Compound B

analogs varies *in vitro* based on substitutions to the benzene ring. In a molecular

dynamics simulation, Compound B remains stably associated with AP2-EXP, while the

non ApiAP2 competitor Compound B-3 does not. This suggests that a steric clash

between AP2-EXP and the bromine atom on Compound B-3 is responsible for its lack of

DNA binding competition. By extension, this may explain the lower efficacy of DNA

binding competition observed for Compounds B-1, B-2, and B-3 compared to B and B-4.

Based on the description of its genome-wide DNA binding sites and transcript

dysregulation in the presence of Compound C, new inferences can be made about the

biological function of AP2-EXP. This demonstrates the utility of combining chemical and

traditional genetics, because previous attempts to genetically characterize AP2-EXP

have not provided a complete picture of its function in the asexual blood stage. The

ap2-exp AP2 domain coding region has previously been shown to be essential for the

asexual blood stage by both saturating mutagenesis and targeted deletion attempts(15–

17, 21). Conversely, the coding region beyond the AP2 domain was truncated in two

studies(15, 21). Transcriptomic profiling of a truncated AP2-EXP parasite line revealed

that many transcripts encoding exported proteins were dysregulated(21). Unexpectedly,

AP2-EXP target genes identified in this study overlap poorly with the differentially

regulated transcripts reported as a result of truncation of AP2-EXP **(Table S4)**[21].

There is greater overlap between our study and AP2-EXP ChIP-seq recently described

by Shang *et al*[17], with 50/101 target genes conserved **(Table S4)**. This may explain

the apparent essentiality of the AP2 domain, while the full-length AP2-EXP has roles

that impact a different subset of non-essential genes. A comparison of transcripts

differentially abundant in the presence of Compound C to the DNA binding occupancy

of AP2-I[18] revealed that 54/85 AP2-I target genes are decreased in abundance by

$\log_2$ fold change >2 at 24-30 hpi. Interestingly, 22/55 AP2-I target genes that decrease

in abundance when parasites are exposed to Compound C are also AP2-EXP targets,

implying the potential for co-regulation of certain gene subsets. This overlap is

consistent with our EMSA results and corroborates the hypothesis that Compound C

competes both AP2-EXP and AP2-I *in vivo*. Only 3/22 PfSIP2 target genes are

decreased in abundance in the presence of Compound C **(ApiAP2 Target Gene**

Results Summarized in Table S3). This may reflect the cryptic relationship between

PfSIP2 DNA binding and transcriptional control, as was noted when PfSIP2 was

originally characterized as playing a role in genome integrity and heterochromatin

formation[20]. Several ApiAP2 proteins have been implicated in oocyst and sporozoite

development in a *P. berghei* ApiAP2 knockout screen[24]. Since Compounds B and C

both inhibit mosquito stage development of *P. berghei* with different phenotypes, it is

possible that they target different sets of ApiAP2 proteins *in vivo*. Since PbAP2-Sp is

required for sporogony, Compounds B and C should minimally inhibit sporozoite

development, which is consistent with our results. Overall, these data support a

potential for multi-AP2 domain competition by Compounds B and C *in vivo*. Due to their

inclusion in the TCAMS, Compounds B, B-1, B-2, B-3, and B-4 have all been tested for

activity against the human HEPG2 cell line(38). ApiAP2 competitor Compounds B and

B-4 inhibit HEPG2 cell growth by just 4 and 8%, respectively(38). Therefore, either drug

may potentially be prioritized for further development based on selectivity for

Plasmodium parasites.

AP2-EXP remains the only ApiAP2 protein for which the structure has been solved.

Surprisingly, over a decade later, the field still lacks a clear understanding of the

biological role of AP2-EXP or insights into the druggability of the AP2 domain. Our study

has enabled us to make inferences about the role of AP2-EXP and set a proof of

principle for targeting the highly unique Apicomplexan AP2 DNA binding proteins as a

new antimalarial strategy.

<u>Materials and Methods</u>
Parasite lines

Parasites were grown at 37°C, 5% O2, 7% CO2 using RPMI1640 media supplemented

with hypoxanthine and .5% Albumax II (Thermo). The parasite lines used in this study

were AP2-EXP::GFP, AP2-EXP::HA, AP2-EXP::*glms*, and wild Type Pf3D7 (Malaria

Research and Reference Reagents Repository). The human biological samples were

sourced ethically and their research use was in accord with the terms of the informed

consents under an IRB/EC approved protocol.

To create AP2-EXP::GFP the C-Terminal coding region of AP2-EXP was cloned **(Table**

S1) into the plasmid pSLI::2xFKBP(43) for endogenous tagging by single homologous

recombination **(Fig S9A)**. Transgenic parasites with the correct C-Terminal tag were

further transfected with pLyn-FRB-mCherry(43) for inducible mislocalization using the

same method. Parasites were maintained in media with 2.5nM WR99210. All ChIP and

western blot experiments using AP2-EXP::GFP except **Figure S26C** were collected

using AP2-EXP::GFP without the pLyn mislocalization plasmid.

To create AP2-EXP::HA we used the pSN054 vector system(58). The right homology

region (RHR) was amplified by PCR **(Table S1)**, and the recodonized left homology

region (LHR) was synthesized using the BioXP™ 3200 System. Single guide RNA

fragments synthesized using the BioXP™ 3200 System **(Table S1)** were cloned into the

linearized pSN054 donor vector(58) The parental parasite line used for transfection

expresses Cas9 and T7 RNA polymerase(59) **(Fig S9B)**. Cell cultures were maintained

in 500nM anhydrotetracycline (aTc, Sigma-Aldrich 37919) and 2.5 mg/mL of Blasticidin

S.

To create AP2-EXP::*glms*::HA, the C-terminal coding region of AP2-EXP **(Table S1)**

was cloned into the plasmid pSLI::3xHA::*glms*(60) **(Figure S25)**. Parasite cultures were

maintained in media with 2.5nM WR99210.

All *P. falciparum* transfection were performed as described(43, 61). All parasite strains

were cloned by limiting dilution and genotyped using PCR. AP2-EXP::GFP and AP2-

EXP::HA were also genotyped using whole genome sequencing **(NCBI SRA:**

PRJNA818769).

Genomic DNA Isolation

Parasite cultures were lysed with .1% Saponin in 1xPBS and collected by centrifugation

at 1500 RPM, then resuspended in 1xPBS. Genomic DNA was then isolated using the

Qiagen DNeasy nucleic acid isolation kit according to the manufacturer's instructions.

AP2-EXP Knockdown Assays

For AP2-EXP::GFP + pLyn, synchronous parasites were split into two populations.

250nM Rapalog was added to one parasite group for 48 hours as described(43) while

the second group was used as a control. Addition of Rapalog did not affect parasite

growth (not shown). After 48 hours parasite protein was collected in nuclear and

cytosolic fractions and AP2-EXP::GFP, Histone H3, and Pf Aldolase were detected by

western blot.

For AP2-EXP::HA parasites were grown routinely in the presence of 500nM aTc. aTc

was washed out of one group, while the second group was maintained with aTc as a

control for 120 hours before protein harvest. Parasite growth was not affected by

removal of aTc (not shown). AP2-EXP::HA and Histone H3 were detected by western

blot.

For AP2-EXP::*glms*::HA, synchronous parasites were split into two populations. 5mM

glucosamine was added to one parasite group for 72 hours as described(56) while the

second group was used as a control. Addition of glucosamine did not affect parasite

growth (not shown). After 72 hours parasite protein was harvested for detection of AP2-

EXP::*glms*::HA and Histone H3 by Western blot.

***In silico* docking screen**

Three dimensional structures of every available molecule from the Tres Cantos

Antimalarial Set(38) and the DrugBank(39) were created using BALLOON and the

Merck Molecular Force Field. The crystal structure of AP2-EXP (PDB:3IGM) was

modeled using AutoDockTools(62) and a molecular docking screen was run using

AutoDock(37). Details about the preparation of ligands, the protein macromolecule, and

the docking screen criteria are provided in the SI text.

IC_{50} Determination

The IC_{50} of each putative ApiAP2 competing compound was determined using a 48-

hour growth inhibition assay as described(63). Parasites were seeded at .5%

parasitemia and 4% hematocrit on a 96 well plate. Drug or DMSO vehicle control was

added in triplicate to each well and parasites were incubated in standard culture

conditions for 48 hours. Growth values were normalized to the vehicle control. IC_{50}

values were calculated and growth inhibition curves were plotted using GraphPad

Prism.

Recombinant Protein Expression

Recombinant AP2 domains AP2-EXP, AP2-I D3, AP2-HS D1, and PfSIP2 D1 were

overexpressed and purified from BL21 PlysS *E. coli* as described previously(12).

Recombinant AtERF1 was purified using the same method after cloning the AtERF1

AP2 domain(40) into the pGex4t-1 overexpression vector **(Table S1)** Recombinant

protein was quantified using Braford reagent (Pierce). Input, flowthrough, and eluate

fractions were analyzed by SDS-PAGE to ensure recovery of the full- length

recombinant protein. Recombinant full-length SOX2 protein was purchased from Abcam

(ab169843).

Electrophoretic Mobility Shift Assays

478 EMSAs were performed using the Thermo Light Shift EMSA kit with recommended

479 buffer components. Gel shifts were visualized using the Light Shift detection reagents

480 and imaged using a BioRad Chemiluminescence imager. Additional details are provided

481 in the SI text.

Ethidium Bromide Exclusion Assay

10µM of double stranded DNA (sequence TGCATGCA, purchased from IDT) in .01mM

EDTA, 9.4mM NaCl, 2mM HEPES buffer pH 7.9 was incubated for 10 minutes in the

presence of 100nM ethidium bromide. Fluorescence was measured at

excitation/emission 546nm/595nm for each well. Following the baseline reading, each

putative ApiAP2 competing compound was added. DRAQ5 nucleic acid dye was used

as a positive control for knockdown of ethidium bromide fluorescence by DNA

intercalation. Ethidium bromide exclusion assays were performed in technical triplicate

using a 96 well plate.

Molecular Dynamics Simulations

Five Compounds (B, B-1, B-2, B-3, and B-4) were simulated using AMBER(64) for

100ns of interaction with the AP2-EXP AP2 domain. The initial position of each

compound used was identical to the predicted docking conformation of Compound B

(Figure S2A). Additional details are provided in the SI text.

AP2 Domain Competitor Phenotyping Assays

For Compound B, C, and H spike in phenotyping **(Fig 3B)**, highly synchronous Pf3D7

wild type parasites were spiked with 40µM of each drug at 24 hpi. An equivalent volume

of DMSO was used as a control. Parasites were morphologically assessed by Giemsa

staining at 24 and 48 hours post spike in (48 and 72 hpi).

For continuous Compound C exposure phenotyping **(Fig S14)**, highly synchronous

Pf3D7 wild type parasites were spiked with 40µM Compound C or equivalent DMSO

vehicle control at 6 hpi. Parasite progression was then monitored through the IDC by

Giemsa staining at 6-hour intervals.

For Compound C spike in and washout assays **(Fig 3C, D)**, 40µM of Compound C was

spiked into highly synchronous Pf3D7 wild type parasites at 8-hour intervals starting at

14 hpi. For Compound C washout assays, 40µM of Compound C was washed out of

highly synchronous culture at 8-hour intervals starting at 14 hpi. For each assay an

equivalent volume of DMSO vehicle was used as a control for normal growth.

40µM of compound was used for all phenotyping assays because it was the maximum

concentration achievable in culture after resuspending each compound in 100% DMSO.

Higher concentrations of compound caused parasite mortality due to the high volume of

DMSO required (>1% volume/volume). All assays were performed in triplicate and

parasitemia was counted by Giemsa-stained slides. Error bars represent the standard

deviation of the mean.

Preparation of RNA for DNA Microarray

DNA microarrays were prepared using the protocol described in(44). After collecting a

control timepoint at 6 hpi, parasites were cultured with either 12µM Compound C

(0.66xIC_{50}) or DMSO vehicle control. Parasite RNA was then harvested at 12, 18, 24,

30, 42, and 48 hpi. cDNA was synthesized using SuperScript II, hybridized onto Agilent

DNA microarrays(44), and scanned using an Axon 4200A Scanner. Agilent Feature

Extraction Software version 11.0.1.1 with the protocol GE2-

v5_95_Feb07_nospikein was used to extract signal intensities. Raw signal intensities for

all microarrays are available in **Table S6**.

Analysis of DNA Microarrays

Microarray data were extracted and normalized using the R packages LIMMA(46) and

Rnits(48). LIMMA was used to analyze differential transcript abundance specific to the

24-30 hpi time points, while Rnits was used to assess differential transcript abundance

across the entire IDC. Additional details are provided in the SI text.

ChIP Sample Preparation

ChIP-seq was performed as described in(19). Highly synchronous parasites were grown

up to 5-10% parasitemia and AP2-EXP ChIP was performed at 30 hpi for all replicates.

Immunoprecipitation was performed overnight with .5mg/mL 3F10 anti-HA (Sigma) or

.1mg/mL Ab290 (Abcam) anti-GFP.

DNA Library Preparation

ChIP-seq DNA libraries were prepared as described in(19) using the NEBNext II DNA

library kit (New England Biolabs) according to the manufacturer's instructions. Quality

was assessed using an Agilent 2100 Bioanalyzer or TapeStation. Libraries were

sequenced using a HiSeq 2500 (Replicate GFP1) or NextSeq 550 (Replicates GFP2

and HA3) Illumina sequencer. AMPure XP beads (Beckmann Coulter) were used to size

select and purify DNA between NEBNext II library preparation steps. Whole genome

sequencing DNA libraries were prepared using the Illumina Tru-Seq PCR free DNA

library kit and sequenced using a HiSeq 2500.

qPCR

ChIP-qPCR samples were collected from 30-35 hpi trophozoites following two hours of

Compound C (40µM) or DMSO vehicle control spike in. Primer pairs to be used for

ChIP-qPCR were first evaluated to check for 80-110% efficiency using sonicated

genomic DNA. RT-qPCR was carried out using Sybr Green Polymerase master mix

(Thermo) with the specified primer concentration **(Table S1)**. The Ct was calculated

using SDSv1.4 (Applied Biosystems) software, averaged over technical triplicate. The

percent of input per immunoprecipitated DNA fraction was calculated using the delta Ct

method. Each assay was performed in biological triplicate, with the exception of the

ama1 primer pair, where n =2. Error bars represent standard deviation of the mean.

Data was obtained using an Applied Biosystems 7300 Real-Time PCR Machine.

ChIP-seq data analysis

ChIP-seq reads were mapped to generate bam and bigwig coverage files as

described(19). Peaks of occupancy were called using MACS2(65) and peak intervals

were analyzed for overlap and gene proximity using BedTools(66). DeepTools(67) and

cegr-tools: https://github.com/seqcode/cegr-tools were used to compare AP2-EXP

genomic occupancy to previously published datasets. BedTools(66) was used to

compare AP2-EXP peaks determined in this study to AP2-EXP occupancy determined

in Shang *et al*(17). Additional details are available in the SI text.

Western blot

Full parasite protein western blot samples were collected by lysing RBCs with .1%

saponin and boiling protein in Loading Buffer (50mM Tris-Cl pH 8.0, 20% SDS, 1%

Bromophenol Blue). Fractionated parasite protein was prepared as described in(68).

Blots were performed as described(19). Primary antibodies used were: 1/1000 rat ant-

HA (Roche 3F10), 1/1000 mouse anti-GFP (Roche), 1/3000 rabbit anti-aldolase

conjugated to HRP (Abcam ab38905), or 1/3000 mouse anti-H3 (Abcam ab10799).

Secondary antibody concentrations used were 1/3000 goat anti-rat HRP conjugate

(Millipore), 1/3000 goat anti-mouse HRP conjugate, or 1/10,000 (Pierce) goat anti-rabbit

HRP conjugate (Millipore). ECL reagent (Pierce) was used to detect HRP signal. Blots

were exposed to autoradiography film (VWR) and visualized using an autoradiography

developer.

Protein Pulldown

Parasite nuclear protein was isolated as described(68). Following the final centrifugation

step, the supernatant was collected and diluted by 1:3 in Dilution Buffer (30% glycerol,

20mM HEPES pH 7.8). GFP tagged AP2-EXP was pulled down using Chromotek anti-

GFP or mock immunoprecipitated using Chromotek negative control magnetic beads.

Beads were washed twice in Wash Buffer prior to use (20mM HEPES pH 7.4, 250mM

NaCl, 1mM EDTA, 1mM TCEP, .05% NP-40). Protein and beads were incubated

together for 1 hour at 4°C. The beads were then washed twice with Wash Buffer and

bound protein was collected in Loading Buffer by boiling at 95°C for 10 minutes.

Fluorescent Microscopy

Samples were prepared by incubating packed infected red blood cells with DRAQ5 dye

(Thermo) for 15 minutes. Parasites were then washed in 1xPBS to remove excess dye

and immediately placed on a glass slide for imaging. Fluorescent microscopy images

were acquired using an Olympus Bx61 fluorescent microscope. All images were

processed using SlideBook 5.0.

Mosquito Stage *P. berghei* AP2 Competitor Growth Assay

A female Swiss Webster mouse was inoculated with Plasmodium berghei ANKA 2.34 from frozen stock. Once the parasitemia reached 15%, the blood was harvested by heart puncture, washed twice with 1xPBS and resuspended to 10 mL in 1xPBS. 5 female Swiss Webster mice were infected with 500µL of the resuspended blood. 3-4 days after blood passage, exflagellation of the *P. berghei* gametocytes was assayed. Briefly, a drop of tail vein blood was incubated in RPMI 1640 (Invitrogen) containing 1µM xanthurenic acid (Sigma) for ~12-15 minutes. The mixture was added to a slide and observed under a light microscope at 40x. Ten or more fields were observed. Mice with 0.3-0.7 exflagellations per field were anesthetized and fed on 3-day post emergence *Anopheles stephensi* mosquitoes. Mosquitoes were maintained at 19°C and fed 10% sucrose. Bloodfed mosquitoes were separated 30 hours post-bloodfeeding. 10 days post-bloodfeeding, 5-10 mosquitoes were dissected and oocysts were counted by mercurochrome staining and light microscopy to ensure *P. berghei* oocyst development. The remaining mosquitoes were injected by standard mouth pipette technique with different small molecule inhibitors or a mixture of PBS-DMSO for control. The surviving injected mosquitoes were dissected 14 days post-bloodfeeding. Oocyst numbers were counted as before. Pooled groups of midguts were ground using a pestle, centrifuged at 7,000 rpm for 5 minutes and resuspended in 20µL 1xPBS to release developing mid-gut sporozoites. Sporozoites were counted on a haemocytometer. All *in vivo* studies were conducted in accordance with the GSK Policy on the Care, Welfare and Treatment of

Laboratory Animals and were reviewed by the ethical review process at the institution

where the work was performed.

<u>Data Availability</u>

All sequencing data for AP2-EXP ChIP-seq and whole genome sequencing for

transgenic parasite lines AP2-EXP::GFP and AP2-EXP::HA have been deposited in the

SRA under **PRJNA818769**.

Raw microarray data for the Compound C and DMSO control RNA time course is

available in **Table S6**.

<u>Author Contributions</u>

T.J.R., E.K.D.S., K.S.S., N.D., C.F.A.P., G.P., M.J.L., C.D.O., J.C.N. and M.L. designed

experiments. T.J.R., E.K.D.S., V.C., K.S.S., N.D., G.J., C.F.A.P., and I.K. performed the

experiments. T.J.R. and N.D. analyzed the data. T.J.R. generated figures. T.J.R. and

M.L. wrote the manuscript in collaboration with the other authors

<u>Acknowledgements</u>

This work was funded through NIH/NIAID R01AI076276 (M.L.), R01AI125565 (M.L.),

and with support from the Center for Quantitative Biology (P50 GM071508) (M.L.).

T.J.R. was supported by NIH T32 Predoctoral Training Grant (5T32GM125592-01)

awarded to the Center for Eukaryotic Gene Regulation (CEGR) at The Pennsylvania

State University. G.A.J. is a recipient of the Sir Keith Murdoch Fellowship from the

American Australian Association and a Postdoctoral Research Grant from the American

Heart Association (16POST26420067).

Conflict of Interest Statement

F.J.G. is a GlaxoSmithKline employee and own shares of the company

REFERENCES

1.   WHO, *Word Malaria Report 2021* (2021).

2.   M. M. Ippolito, K. A. Moser, J.-B. B. Kabuya, C. Cunningham, J. J. Juliano, Antimalarial Drug
Resistance and Implications for the WHO Global Technical Strategy. *Curr. Epidemiol. Reports* **8**,
46–62 (2021).

3.   A. F. Cowman, J. Healer, D. Marapana, K. Marsh, Malaria: Biology and Disease. *Cell* **167**, 610–624
(2016).

4.   G. Zanghì, *et al.*, A Specific PfEMP1 Is Expressed in P. falciparum Sporozoites and Plays a Role in
Hepatocyte Infection. *Cell Rep.* **22**, 2951–2963 (2018).

5.   Z. Bozdech, *et al.*, The transcriptome of the intraerythrocytic developmental cycle of Plasmodium
falciparum. *PLoS Biol.* **1**, 85–100 (2003).

6.   X. M. Lu, *et al.*, Nascent RNA sequencing reveals mechanisms of gene regulation in the human
malaria parasite Plasmodium falciparum. *Nucleic Acids Res.* **45**, 7825–7840 (2017).

7.   H. J. Painter, M. Carrasquilla, M. Llinás, Capturing in vivo RNA transcriptional dynamics from the
malaria parasite Plasmodium falciparum. *Genome Res.* **27**, 1074–1086 (2017).

8.   K. G. Le Roch, *et al.*, Discovery of gene function by expression profiling of the malaria parasite life
cycle. *Science* **301**, 1503–1508 (2003).

9.   B. J. Foth, *et al.*, Quantitative time-course profiling of parasite and host cell proteins in the
human malaria parasite Plasmodium falciparum. *Mol. Cell. Proteomics* **10**, 1–16 (2011).

10.   T. Hollin, K. G. Le Roch, From Genes to Transcripts, a Tightly Regulated Journey in Plasmodium.
Front. Cell. Infect. Microbiol. **10**, 1–13 (2020).

11.   H. J. Painter, T. L. Campbell, M. Llinás, The Apicomplexan AP2 family: Integral factors regulating
Plasmodium development. **176**, 1–7 (2012).

12.   T. L. Campbell, E. K. de Silva, K. L. Olszewski, O. Elemento, M. Llinás, Identification and Genome-
Wide Prediction of DNA Binding Specificities for the ApiAP2 family of regulators from the malaria
parasite. *PLoS Pathog.* **6**, e1001165 (2010).

13.   S. Balaji, M. Madan Babu, L. M. Iyer, L. Aravind, Discovery of the principal specific transcription
factors of Apicomplexa and their implication for the evolution of the AP2-integrase DNA binding
domains. *Nucleic Acids Res.* **33**, 3994–4006 (2005).

14.   M. Jeninga, J. Quinn, M. Petter, ApiAP2 Transcription Factors in Apicomplexan Parasites.
Pathogens **8** (2019).

15.   M. Zhang, *et al.*, Uncovering the essential genes of the human malaria parasite Plasmodium
falciparum by saturation mutagenesis. *Science* **360**, eaap7847 (2018).

16.   X. Shang, *et al.*, A cascade of transcriptional repression determines sexual commitment and

692   development in Plasmodium falciparum. *Nucleic Acids Res.* **49**, 9264–9279 (2021).

693 17. X. Shang, *et al.*, Genome-wide landscape of ApiAP2 transcription factors reveals a
694   heterochromatin-associated regulatory network during Plasmodium falciparum blood-stage
695   development. *Nucleic Acids Res.* **50**, 3413–3431 (2022).

696 18. J. M. Santos, *et al.*, Red Blood Cell Invasion by the Malaria Parasite Is Coordinated by the PfAP2-I
697   Transcription Factor. *Cell Host Microbe* **21**, 731–741 (2017).

698 19. G. A. Josling, *et al.*, Dissecting the role of PfAP2-G in malaria gametocytogenesis. *Nat. Commun.*
699   **11**, 1503 (2020).

700 20. C. Flueck, *et al.*, A major role for the Plasmodium falciparum ApiAP2 protein PfSIP2 in
701   chromosome end biology. *PLoS Pathog.* **6**, e1000784 (2010).

702 21. R. M. Martins, *et al.*, An ApiAP2 member regulates expression of clonally variant genes of the
703   human malaria parasite Plasmodium falciparum. *Sci. Rep.* **7**, 14042 (2017).

704 22. E. Carrington, *et al.*, The ApiAP2 factor PfAP2-HC is an integral component of heterochromatin in
705   the malaria parasite Plasmodium falciparum. *iScience* **24**, 102444 (2021).

706 23. M. Sierra-Miranda, *et al.*, PfAP2Tel, harbouring a non-canonical DNA-binding AP2 domain, binds
707   to Plasmodium falciparum telomeres. *Cell. Microbiol.* **19**, e12742 (2017).

708 24. K. Modrzynska, *et al.*, A Knockout Screen of ApiAP2 Genes Reveals Networks of Interacting
709   Transcriptional Regulators Controlling the Plasmodium Life Cycle. *Cell Host Microbe* **21**, 11–22
710   (2017).

711 25. Z. Li, *et al.*, Plasmodium transcription repressor AP2-O3 regulates sex-specific identity of gene
712   expression in female gametocytes. *EMBO Rep.*, e51660 (2021).

713 26. Y. Xu, *et al.*, PfAP2-G2 Is Associated to Production and Maturation of Gametocytes in Plasmodium
714   falciparum via Regulating the Expression of PfMDV-1. *Front. Microbiol.* **11**, 631444 (2021).

715 27. S. Singh, *et al.*, The PfAP2-G2 transcription factor is a critical regulator of gametocyte maturation.
716   *Mol. Microbiol.* **115**, 1005–1024 (2021).

717 28. C. Zhang, *et al.*, Systematic CRISPR-Cas9-Mediated ApiAP2 Genes Reveal Functional Insights into
718   Parasite Development. *MBio* **8**, 1–17 (2017).

719 29. A. Sinha, *et al.*, Europe PMC Funders Group Europe PMC Funders Author Manuscripts A cascade
720   of DNA binding proteins for sexual commitment and development in Plasmodium. **507**, 253–257
721   (2014).

722 30. B. F. C. Kafsack, *et al.*, A transcriptional switch underlies commitment to sexual development in
723   malaria parasites. *Nature* **507**, 248–52 (2014).

724 31. S. Iwanaga, I. Kaneko, T. Kato, M. Yuda, Identification of an AP2-family Protein That Is Critical for
725   Malaria Liver Stage Development. *PLoS One* **7**, e47557 (2012).

726 32. M. Yuda, S. Iwanaga, S. Shigenobu, T. Kato, I. Kaneko, Transcription factor AP2-Sp and its target
727   genes in malarial sporozoites. *Mol. Microbiol.* **75**, 854–863 (2010).

728 33. M. Yuda, *et al.*, Identification of a transcription factor in the mosquito-invasive stage of malaria
729   parasites. *Mol. Microbiol.* **71**, 1402–1414 (2009).

730 34. M. Yuda, I. Kaneko, S. Iwanaga, Y. Murata, T. Kato, Female-specific gene regulation in malaria
731   parasites by an AP2-family transcription factor. *Mol. Microbiol.* **113**, 40–51 (2020).

732 35. E. Tintó-font, L. Michel-todó, T. J. Russell, N. Casas-, A heat-shock response regulated by the
733   PfAP2-HS transcription factor protects human malaria parasites from febrile temperatures. **6**,
734   1163–1174 (2021).

735 36. S. E. Lindner, E. K. De Silva, J. L. Keck, M. Llinás, Structural Determinants of DNA Binding by a P.
736   falciparum ApiAP2 Transcriptional Regulator. *J. Mol. Biol.* **395**, 558–567 (2010).

737 37. G. M. Morris, *et al.*, Automated docking using a Lamarckian genetic algorithm and an empirical
738   binding free energy function. *J. Comput. Chem.* **19**, 1639–1662 (1998).

739 38. F. J. Gamo, *et al.*, Thousands of chemical starting points for antimalarial lead identification.

Nature **465**, 305–310 (2010).

39.   D. S. Wishart, *et al.*, DrugBank 5.0: A major update to the DrugBank database for 2018. *Nucleic*
Acids Res. **46**, D1074–D1082 (2018).

40.   M. D. Allen, K. Yamasaki, M. Ohme-Takagi, M. Tateno, M. Suzuki, A novel mode of DNA
recognition by a β-sheet revealed by the solution structure of the GCC-box binding domain in
complex with DNA. *EMBO J.* **17**, 5484–5496 (1998).

41.   A. J. Geall, I. S. Blagbrough, A comparison of the properties of Sox-3 with Sry and two related
genes, Sox-1 and Sox-2. *J. Pharm. Biomed. Anal.* **22**, 849–859 (2000).

42.   Y. Wang, H. Schellenberg, V. Walhorn, K. Toensing, D. Anselmetti, Binding Mechanism of
Fluorescent Dyes to DNA Characterized by Magnetic Tweezers. *Mater. Today Proc.* **4**, S218–S225
(2017).

43.   J. Birnbaum, *et al.*, A genetic system to study Plasmodium falciparum protein function. *Nat.*
Methods **14**, 450–456 (2017).

44.   B. F. C. Kafsack, H. J. Painter, M. Llinás, New Agilent platform DNA microarrays for transcriptome
analysis of Plasmodium falciparum and Plasmodium berghei for the malaria research community.
Malar. J. **11**, 187 (2012).

45.   M. Llinás, Z. Bozdech, E. D. Wong, A. T. Adai, J. L. DeRisi, Comparative whole genome
transcriptome analysis of three Plasmodium falciparum strains. *Nucleic Acids Res.* **34**, 1166–1173
(2006).

46.   G. K. Smyth, Linear models and empirical Bayes methods for assessing differential expression in
microarray experiments. *Stat. Appl. Ge- netics Mol. Biol.* **3**, Article 3 (2011).

47.   A. Bahl, *et al.*, PlasmoDB: The Plasmodium genome resource. A database integrating
experimental and computational data. *Nucleic Acids Res.* **31**, 212–215 (2003).

48.   D. P. Sangurdekar, The Rnits package for normalization and inference of differential expression in
time series microarray data . *Rvignette*, 1–13 (2014).

49.   M. Yuda, I. Kaneko, Y. Murata, S. Iwanaga, T. Nishi, Targetome analysis of malaria sporozoite
transcription factor AP2-Sp reveals its role as a master regulator. *bioRxiv* (2022)
https:/doi.org/https://doi.org/10.1101/2022.02.23.481739.

50.   P. R. Kensche, *et al.*, The nucleosome landscape of Plasmodium falciparum reveals chromatin
architecture and dynamics of regulatory sequences. *Nucleic Acids Res.* **44**, 2110–2124 (2015).

51.   R. Bártfai, *et al.*, H2A.Z demarcates intergenic regions of the Plasmodium falciparum epigenome
that are dynamically marked by H3K9ac and H3K4me3. *PLoS Pathog.* **6**, e1001223 (2010).

52.   G. A. Josling, *et al.*, A Plasmodium Falciparum Bromodomain Protein Regulates Invasion Gene
Expression. *Cell Host Microbe* **17**, 741–751 (2015).

53.   L. Chappell, *et al.*, Refining the transcriptome of the human malaria parasite Plasmodium
falciparum using amplification-free RNA-seq. *BMC Genomics* **21**, 395 (2020).

54.   C. G. Toenhake, *et al.*, Chromatin Accessibility-Based Characterization of the Gene Regulatory
Network Underlying Plasmodium falciparum Blood-Stage Development. *Cell Host Microbe* **23**,
557–569 (2018).

55.   S. M. Ganesan, A. Falla, S. J. Goldfless, A. S. Nasamu, J. C. Niles, Synthetic RNA-protein modules
integrated with native translation mechanisms to control gene expression in malaria parasites.
Nat. Commun. **7**, 10727 (2016).

56.   P. Prommana, *et al.*, Inducible Knockdown of Plasmodium Gene Expression Using the glmS
Ribozyme. *PLoS One* **8**, e73783 (2013).

57.   D. G. Paton, *et al.*, Exposing Anopheles mosquitoes to antimalarials blocks Plasmodium parasite
transmission. *Nature* **567**, 239–243 (2019).

58.   A. S. Nasamu, *et al.*, An integrated platform for genome engineering and gene expression
perturbation in Plasmodium falciparum. *Sci. Rep.* **11**, 342 (2021).

59. J. M. Murithi, *et al.*, The Plasmodium falciparum ABC transporter ABCI3 confers parasite strain-
dependent pleiotropic antimalarial drug resistance. *Cell Chem. Biol.*, 1–16 (2021).
60. C. S. Simon, *et al.*, An extended DNA-free intranuclear compartment organizes centrosome
microtubules in malaria parasites. *Life Sci. Alliance* **4**, e202101199 (2021).
61. Y. Wu, L. A. Kirkman, T. E. Wellems, Transformation of Plasmodium falciparum malaria parasites
by homologous integration of plasmids that confer resistance to pyrimethamine. *Proc. Natl.*
Acad. Sci. U. S. A. **93**, 1130–4 (1996).
62. R. Huey, G. M. Morris, Using AutoDock 4 with AutoDockTools : A Tutorial. 1–56.
63. L. Abrahams, M. Burke, w o T w o. 20–36 (2010).
64. A. W. Götz, *et al.*, Routine microsecond molecular dynamics simulations with AMBER on GPUs. 1.
generalized born. *J. Chem. Theory Comput.* **8**, 1542–1555 (2012).
65. J. M. Gaspar, Improved peak-calling with MACS2. *bioRxiv*, 1–16 (2018).
66. A. R. Quinlan, I. M. Hall, BEDTools: A flexible suite of utilities for comparing genomic features.
Bioinformatics **26**, 841–842 (2010).
67. F. Ramírez, F. Dündar, S. Diehl, B. A. Grüning, T. Manke, DeepTools: A flexible platform for
exploring deep-sequencing data. *Nucleic Acids Res.* **42**, 187–191 (2014).
68. S. C. Oehring, *et al.*, Organellar proteomics reveals hundreds of novel nuclear proteins in the
malaria parasite Plasmodium falciparum. *Genome Biol* **13**, R108 (2012).
806
807

<u>Figure Legends</u>

Figure 1. Nine putative competitors of DNA binding by AP2-EXP were identified by an *in-

silico screen and tested *in vitro* and *in vivo***

A) Each compound that was prioritized based on the *in-silico* screen was assigned an identifier

(A-I) and tested for anti-*Plasmodium* activity in a 48-hour growth inhibition assay against

asexual blood stage *P. falciparum*.

B) Each putative AP2-EXP competitor was added to an EMSA containing the AP2-EXP AP2

domain. DNA binding competition will result in a loss of the shifted DNA probe. DMSO vehicle

was used as a control for normal DNA binding by AP2-EXP. 150 fmoles of DNA probe, 125ng of

AP2-EXP, and 300µM of each compound were used for each lane.

C) Chemical structures of AP2-EXP competing Compounds A, B, C, and I. Three of the four

compounds that compete DNA binding *in vitro* (Compounds A, B, and C) have a benzoxazole

core moiety, denoted by a red box.

Figure 2. Analogues of Compound B compete AP2-EXP DNA binding with differing efficacy

A) Compounds B, B-1, B-2, B-3, and B-4 were added to an EMSA with AP2-EXP. DNA binding

competition will result in the loss of the shifted probe. 40 fmoles of DNA probe, 125ng of AP2-

EXP, and 300µM of each compound were used for each lane.

B) Compounds B, B-1 and B-4 were titrated at 25, 50, and 150µM into an EMSA with AP2-EXP

to determine whether there are differences in DNA binding competition. DMSO vehicle was

used as a control for normal DNA binding. 40 fmoles of DNA probe and 125ng of AP2-EXP were

used for each lane. The concentration increase for each compound is indicated by the triangle

from left to right.

C) Chemical structures of Compounds B, B-1, B-2, B-3, and B-4. Compounds that compete

DNA binding by AP2-EXP are highlighted in black. Each compound has a different substitution

pattern on the right-side benzene ring, denoted by a red box. The IC_{50} against asexual *P.*

falciparum of compounds B-1, B-2, B-3, and B-4 measured by Gamo *et al*(38) is indicated at the

bottom of each identifier. The IC_{50} of Compound B was determined in this study.

Figure 3. AP2 Domain Competing Compounds B and C affect *P. falciparum* at in the mid trophozoite stage, coinciding with the maximum expression of AP2-EXP

A) AP2-EXP protein expression in the asexual blood stages ring (R), trophozoite (T), and

schizont (S), and Stage III gametocytes (Gam), was probed by western blot against AP2-EXP

endogenously tagged with GFP (AP2-EXP::GFP). The expected molecular weight of AP2-

EXP::GFP (147kDa) is indicated by an arrow. Wild type Pf3D7 protein was used as a negative

control. The full length western blot is provided in **Figure S10**.

B) Representative Giemsa-stained microscopy images of *P. falciparum* parasites spiked with

40µM Compounds B, C, or H. DMSO vehicle was used as a control for normal progression

through the asexual blood stage at 24 and 48 hours post spike-in.

C) Highly synchronous asexual blood stage *P. falciparum* parasites were cultured with 40µM

Compound C, which was washed out of the media at eight-hour intervals. IDC progression was

measured by counting ring stage parasites at 54 hpi. Each assay was done in triplicate. Error

bars represent standard deviation.

D) Highly synchronous asexual blood stage *P. falciparum* parasites were spiked with 40µM

Compound C at eight-hour intervals in the IDC. IDC progression was measured by counting ring

stage parasites at 54 hpi Each assay was done in triplicate. Error bars represent standard

deviation.

Figure 4. Compound C disrupts the *P. falciparum* transcriptome specifically at 30hpi, with

bias towards AP2-EXP targets predicted by ChIP-seq

A) *P. falciparum* parasites were spiked with 12µM ($0.66 \times IC_{50}$) Compound C or DMSO vehicle

control at 10 hpi. Total RNA was harvested at 6, 12, 18, 24, 30, 42, and 48 hpi for quantification

by DNA microarray. The full transcriptome Spearman Correlation between Compound C and

DMSO control spiked parasites was plotted as a correlogram. A * indicates *p-value* < 0.05.

B) Three replicates of AP2-EXP Chip-seq were collected at 30 hpi using two genetically tagged

parasite lines (AP2-EXP::GFP and AP2-EXP::HA). A no epitope control was collected using wild

type Pf3D7 parasites and anti-GFP antibodies. Log_2 enrichment of the immunoprecipitate (ChIP)

over Input DNA for one replicate of AP2-EXP::GFP, AP2-EXP::HA, and No Epitope Control is

plotted relative to the coding sequence start site of AP2-EXP target genes identified in at least

2/3 replicates. CATGA is the most overrepresented DNA motif within the peaks of AP2-EXP

occupancy conserved in at least 2/3 ChIP-seq experiments.

C) A volcano plot of the changes in abundance for AP2-EXP target genes at 24-30 hpi in the

presence of Compound C. 46/93 detected AP2-EXP target transcripts decrease in abundance

by log_2 fold change > 2.

D) GO-term analysis of transcripts that are both AP2-EXP targets and decreased in abundance

with respect to Compound C (Bonferroni *p-value* cutoff < 0.05).

Figure 5. Compounds B and C are active against mosquito stage *Plasmodium berghei*

parasites

A) Midgut oocyst counts per mosquito for *P. berghei* infected mosquitoes on day 14 post

infection following compound injection on day 10 post infection. Compound identity is indicated

in the legend. Rep. 1 and Rep. 2 correspond to biological replicates one and two, respectively.

B) Midgut sporozoite counts per five infected mosquitoes on day 14 post infection following

compound injection on day 10 post infection. Compound identity is indicated in the legend. Rep.

1 and Rep. 2 correspond to biological replicates one and two, respectively.

Figure 1

A

Letter Code	PubChem ID	IC$_{50}$ (µM)
A	CID1365835	11
B	CID1365471	80
C	CID5750730	18
D	CID4541005	22
E	CID2343	170
F	CID4437	40
G	CID11390	34
H	CID6914273	40
I	CID25863	125

B

AP2-EXP

Compound ID: A B C D E F G H I DMSO

Shifted Probe ▶

Free Probe ▶

C

CID1365853
Compound A

CID1365471
Compound B

CID124220
Compound C

CID25863
Compound I

Figure 2

CID1365471
IC$_{50}$: 80μM

CID1104215
IC$_{50}$: 1.13μM

CID1931935
IC$_{50}$: 0.16μM

CID1356045
IC$_{50}$: 0.96μM

CID15991863
IC$_{50}$: 0.87μM

Figure 3

Figure 4

Figure 5

SI Text

Extended Methods

In Silico **Docking Screen**

The Tres Cantos Antimalarial Set of 13533 small molecules was retrieved from the

supplementary material of Gamo *et al*(1) and downloaded in Spatial Data File (SDF) format from

BatchEntrez. 4603 small molecule structures were downloaded from DrugBank(2) version 2.5 in

SDF format. Using the SDF library, a 3-dimensional MOL2 library was created with the use of

BALLOON and the Merck Molecular Force Field. BALLOON was used to calculate up to 20

conformations using a genetic algorithm and to select the conformation with the lowest energy

of conformation. The 3-dimensional MOL2 library was then prepared for AutoDock using the

python script "ligand_prepare4.py" from AutoDockTools(3). The prepared ligands were then

saved in PDBQT file format. The python script "ligand_prepare4.py" automatically defines the

torsions of the molecule.

The homodimerized AP2-EXP AP2 domain was downloaded from the solved crystal structure

(PDB ID 3IGM). AutoDockTools was used to remove water molecules, add hydrogens, and

calculate charges. Residues Arg-88 and Asn-118 were found to be flexible using FlexPred and

set as flexible residues for docking. The flexible residues were saved as flexible residues in a

flexible residue file and the remaining molecule was saved as rigid residues in a rigid residue file

as specified in the AutoDock manual(3). The rigid residues file was used to pre-calculate an

energy grid of the macromolecule using autogrid4(3). A set of 14 grid map types S, Cl, F,A, Br,

N, P, OA, SA, C, I, HD, Ca, NA were calculated for the macromolecule using AutoGrid(3).

From the prepared ligands a docking set was prepared for AutoDock with the macromolecule

and pre-calculated energy grids. For each docking set a docking parameter file was prepared,

with the Lamarckian Genetic Algorithm (LGA) specified as the search algorithm for AutoDock.

To allow for high throughput parallel processing PERL was used on top of AutoDock to manage

the dockings. The algorithm was initially set to run with a maximum of 25000 energy evaluations

and 20 repeats. Since the calculations are computationally intensive, this setting was used as

initial screening of the ligands, to identify a subset of ligands for a more thorough evaluation(3).

The 1000 best hits from the GSK compound evaluation and the DrugBank compound evaluation

were selected for a more thorough evaluation. The docking parameters were reconfigured to

250000 energy evaluations and 100 repeats and docking repeated. The docking results were

examined with PERL, to provide an automated approach to the interpretation of the docking

results. The docking results were evaluated using two different approaches. The top candidates

for competition of DNA were selected using a PERL script to filter based on having a geometric

center within 10 Å of the location of sense-strand DNA binding. Hits were then filtered based on

having a predicted free energy of interaction < -5 kJ/Mol.

Electrophoretic Mobility Shift Assay

EMSAs were run in DNA binding buffer (10mM Tris pH 7.5, 50mM KCl, 1mM DTT, 6mM MgCl2,

60ng/µL Poly DiDC, 65ng BSA). Recombinant proteins were titrated to empirically determine the

minimum mass required for DNA binding and this mass **(Fig S4)** was used for each gel-shift

unless otherwise specified. PAGE purified DNA probes with a 5' biotin ligated on the forward

DNA strand, along with an unlabeled complementary stand, were purchased from IDT **(Table**

S1). DNA probes were double stranded by heating to 95°C, followed by stepwise cooling in

Annealing Buffer (10mMTtris-Cl pH7.5, 1mM EDTA, 10mM NaCl). Each recombinant protein

was incubated in DNA binding mixture plus competitor compound for 15 minutes prior to

addition of the cognate double stranded DNA oligonucleotide. Protein, competitor, and DNA

oligonucleotide were incubated together for an additional 5 minutes. The mixtures were

separated on a .5x TBE polyacrylamide gel, transferred to a nylon membrane (Amersham) at 50

Volts for 30 minutes, and probed using the Light Shift nucleic acid detection module

(Thermo) according to the manufacturer's protocol. All gels were imaged using a Bio Rad

chemiluminescence imager.

Molecular Dynamics Simulations

Five compounds were prepared for molecular dynamics (MD) simulation: B, B-1, B-2, B-3, and

B-4. In preparation for MD, parameters for the compounds were obtained using

Antechamber(4–6) with the Generalized Amber Force Field. Starting conformations for each

compound bound to AP2-EXP were based on predictions from docking. All complexes were

prepared using the tleap module of AmberTools(7) with the protein.ff14SB forcefield(8). Each

complex was solvated in an octahedral box of TIP3P water with a 10 -Å buffer around the

protein complex. Na^+ and Cl^- ions were added to neutralize the protein and achieve

physiological conditions. All MD minimizations and simulations were performed using Amber

with GPU acceleration(9, 10). First, complexes minimized with 5000 steps each of steepest

decent and conjugate gradient minimization with 500 kcal/mol·$Å^2$ restraints on all complex

atoms. Restraints were reduced to 100 kcal/mol·$Å^2$ and the minimization protocol was repeated.

Restraints were then retained only on the compound for a final minimization step. Following

minimization, all complexes were heated from 0 to 300 K using a 100-ps run with constant

volume periodic boundaries and 10 kcal/mol·$Å^2$ restraints on all protein and compound atoms.

To equilibrate complexes, 10 ns of MD was performed first with 10 kcal/mol·$Å^2$, then with 1

kcal/mol·$Å^2$ restraints on protein and compound atoms using the NPT ensemble. With

kcal/mol·$Å^2$ restraints retained on complexes, 500 ns production simulations were performed. A

2-fs timestep was used and all bonds between heavy atoms and hydrogens were fixed with the

SHAKE algorithm(11). A cut-off distance of 10 Å was used to evaluate long-range electrostatics

with Particle Mesh Ewald (PME) and for van der Waals forces. The 'strip' and 'trajout'

commands of the CPPTRAJ module(12) were used to remove solvent atoms and extract 50,000

evenly spaced frames from each simulation for analysis.

Analysis of DNA Microarrays

LIMMA(13) was used to normalize and extract signal intensities. Arrays were normalized using

robust splines normalization and within arrays parameters selected. Average expression values

were calculated per gene by averaging the $\log_2$ Cy5 (cDNA)/Cy3 (Reference Pool) signal

intensity across all probes. Correlation plots for the Compound C vs. DMSO control total

transcriptome and control genes were made using the $\log_2$ signal intensities for all detected

genes using the R package CorrPlot downloaded from: https://github.com/taiyun/corrplot. The

LIMMA eBayes function was used on average gene abundance data to determine changes in

transcript abundance between 24 and 30hpi that occur differentially in DMSO control vs.

Compound C dosed parasites. Volcano plots were made with LIMMA eBayes data using the

Enhanced Volcano package in R downloaded from:

https://github.com/kevinblighe/EnhancedVolcano. The Rnits(14) R package was used to model

differences in gene expression across the entire time course between the DMSO control and

Compound C spike in parasites. Rnits was run using the parameters: center genes, normalize

by intensity, and background filter probes. The Rnits model for differential expression was fit at

the gene level. All data from LIMMA, eBayes, and Rnits is provided in **Table S3**. Heat maps for

control genes and AP2-EXP target genes were made using Java Treeview(15) after means

centering and log transformation of transcript abundance values with Cluster3.0(16).

ChIP-seq Data Analysis

First, reads in each DNA library were trimmed to remove low quality base calls and Illumina

adaptor sequences using Trimmomatic(17) with a quality score cutoff of 20. FastQC(18) was

used to assess the quality of each DNA library following this step. Reads were then mapped to

the *Plasmodium falciparum* genome version 36 downloaded from PlasmoDB using BWA-

Mem(19). Multiply mapped reads were filtered out using Samtools(20). Filtered and mapped

Input and immunoprecipitate bam files were used for peak calling by MACS2(21) with the

following parameters: effective genome size 20000000, No Model, q 0.01. Peaks called in each

ChIP-seq replicate were overlapped using Bedtools(22) Intersect, and peaks that occur in a

minimum of 2/3 replicates were used for downstream analysis. For data visualization using IGV,

.bam files were converted to $\log_2$ Immunoprecipitate fraction/Input fraction bigwigs using

DeepTools(23) BamCompare. Bedtools(22) ClosestBed was used to correlate regions called as

peaks of occupancy by MACS2 to the closest gene. Results were filtered based on 1.5kb

proximity to the MACS2 peak and strandedness of the putative target gene. For comparison of

ChIP-seq library coverages to Transcription Start Sites and Start Codons, DeepTools

PlotHeatmap was used. Matrix files underlying PlotHeatmap were created using DeepTools

ComputeMatrix. A four-color plot of conserved AP2-EXP DNA binding sites was made using the

program cegr-tools four color plot downloaded from https://github.com/seqcode/cegr-

tools/tree/master/src/org/seqcode/cegrtools. DNA motifs enriched in AP2-EXP ChIP-seq peaks

were determined using the DREME Suite(24). DNA motifs were sorted from highest to lowest

level of conservation to the aggregate AP2-EXP DNA motif using FIMO(25).

For comparison of AP2-EXP peaks with existing post-translational histone marks, chromatin

post translational modification datasets(26, 27) were downloaded from

https://github.com/Daread/plasmodiumExprPrediction(28). Chromatin reader and nucleosome

occupancy datasets(29–31) were downloaded from NCBI. ChIP-seq data for BDP1 and HP1

was processed as described above. Nucleosome occupancy data was downloaded from NCBI

and processed as described above, with the exception that bam files were turned into bigwig

using BamtoBigWig with the --mnase parameter selected. DeepTools ComputeMatrix was used

to format the data. DeepTools PlotHeatmap was used to plot coverage of each dataset within

10kb of AP2-EXP peaks. For negative control coverage plots, Bedtools(22) ShuffleBed was

used to create random genomic intervals on the same chromosome and the same length as

AP2-EXP peaks of occupancy.

For comparison of AP2-EXP peaks identified by Shang *et al*(32) to this study, AP2-EXP peaks

of occupancy in the trophozoite stage were downloaded from

https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE184658 . Peak summits from each

replicate were expanded to 200 nucleotide windows using Bedtools SlopBed(22). Conserved

peaks in 2/2 replicates were identified using Bedtools intersect(22), then AP2-EXP peaks from

this study and Shang *et al* were overlapped using Bedtools Intersect(22).

SI Figures

A

AP2-EXP Crystal Structure   >10,000 Small Molecules

Computational Docking

B

Letter Code	Compound Used Originally for *in silico* Docking	Compound Name Used for Purchase	PubChem ID of Compound used in this study
A	GSK-like (TCMDC-124220)	CB5768506*	CID1365835
B	GSK (TCMDC-123924)	CB5842949	CID1365471
C	GSK-like (TCMDC-124220)	ChemDiv-8002-1285*	CID5750730
D	GSK-like (TCMDC-124220)	ChemDiv-8004-0752*	CID4541005
E	Drug Bank (DB00562)	Benzathiazide	CID2343
F	Drug Bank (DB04409)	Napthalene Trisulfonate	CID4437
G	Drug Bank (DB04640)	2,6 Napthalene Sulfonate	CID11390
H	Drug Bank (DB01219)	Dantrolene	CID6914273
I	Drug Bank (DB02633)	Procion Blue	CID25863

Figure S1. Putative AP2-EXP competitors were identified using computational docking

A) The crystal structure of AP2-EXP (PDB:3IGM)(33) was used as a template to

computationally dock thousands of small molecules *in silico* using AutoDock. Results were

filtered for compounds that dock within 10 Angstroms of DNA binding residues with a free

energy less than -5kJ/mol. Compounds matching these criteria were sourced and used for

further testing.

B) Seven compounds were identified as putative ApiAP2 competitors in an *in-silico* screen

(Column 2). Five of these were available for direct purchase (Column 3). For the two remaining

compounds, four alternate choices with a Tanimoto similarity score of .9 or greater were

purchased (denoted by an asterisk in Column 3). The PubChem ID used to purchase each

compound in this study is listed in Column 4.

Figure S2. Docking conformations for Compounds A-I

A) The spatial conformation of each of the nine compounds (A-I) that dock within 10 Angstroms

of the DNA binding pocket of AP2-EXP with a free energy less than -5kJ/mol is depicted above.

B) Chemical structures of each compound (A-I) corresponding to the molecular docking results

in panel A.

Figure S3. IC50 assays for Compounds A-I

165   A-I) 48-hour Sybr Green growth inhibition assays were conducted for each of the nine putative

166   ApiAP2 competitor compounds in order to determine IC50 values against asexual *P. falciparum*.

167   All growth assays were performed in triplicate. All compounds kill asexual stage *P. falciparum*

168   parasites in the micromolar (11-170µM) concentration range.

Figure S4. Titration of recombinant DNA binding domains to optimize competition

electrophoretic mobility shift (EMSA) assays

A-F) DNA binding domains AP2-EXP, AP2-I D3, AP2-HS D1, PfSIP2 D1, AtERF1, and full

length SOX2 were titrated against DNA oligos containing their respective binding motifs

(highlighted in red) in an EMSA. Unless otherwise specified, the minimum mass of each

recombinant DNA binding domain required to visualize DNA binding (denoted by an arrow) was

used in competitive EMSAs with putative ApiAP2 competitor compounds.

Figure S5. Putative ApiAP2 competitor compounds were tested against three asexual

blood stage essential *P. falciparum* AP2 domains in addition to AP2-EXP in a competitive

EMSA

183 AP2-I D3 (A) DNA binding activity is competed by Compounds A, B, C and I. AP2-HS D1 (B) is

184 competed by Compounds B and I, and PfSIP2 D1 (C) is competed by Compounds B, C, and I.

185 Compounds A, B, C and I all compete at least one AP2 domain in addition to AP2-

186 EXP. Cognate DNA motifs for each protein are highlighted in red. 300µM of each compound

187 was used per lane.

189

Figure S6. Putative ApiAP2 competitor compounds were tested against off target DNA

binding domains in a competitive EMSA

A) The plant encoded *Arabidopsis thaliana* AP2 domain from Ethylene Response Factor 1

(AtERF1) is competed by Compound I. The *Plasmodium* AP2 domain competitors Compound A,

B, and C do not compete AtERF1. The cognate AtERF1 DNA motif is highlighted in red. 300µM

of each compound was used per lane.

B) The human encoded High Mobility Group Box Domain transcription factor SOX2 is competed

by Compound A. Due to the lack of homology between SOX2 and AP2 domain proteins, this

result indicates that Compound A's DNA binding competition activity is not unique to the AP2

domain. The cognate SOX2 DNA motif is highlighted in red. 300µM of each compound was

used per lane.

Letter Code	AP2-EXP	AP2-I D3	PfSIP2 D1	AP2-HS D1	AtERF1	SOX2
A	+	+		-	-	+
B	+	+	+	+	-	-
C	+	+	+	-	-	-
D	-	-		-	-	-
E	-	-		-	-	-
F	-	-		-	-	-
G	-	-		-	-	-
H	-	-		-	-	-
I	+	+	+	+	+	-

Table S2. Summary of the Competitive EMSA results for each DNA binding domain

tested

Compounds A, B, C, and I all can compete DNA binding by *Plasmodium* AP2 domains.

Compounds A and I each compete one off-target protein domain.

Figure S7. Compounds A-I were tested for DNA intercalation in an ethidium bromide

exclusion assay

Each putative ApiAP2 competitor compound was added into a mixture containing double

stranded DNA and ethidium bromide. The positive control DNA major groove intercalator

DRAQ5 knocks down ethidium bromide fluorescence nearly completely relative to the DNA and

ethidium bromide control. The legend indicates the cumulative result for each compound in

competitive EMSAs. Each assay was performed in triplicate. Error bars represent standard

deviation of the mean.

Figure S8. Compound B analogues were tested against AP2-I D3 in a competitive EMSA

and checked for DNA intercalation ability

A) Compounds B, B-1, B-2, B-3, and B-4 were added to an EMSA with AP2-I D3 to check

whether their DNA binding competition is consistent with AP2-EXP. The cognate AP2-I D3 DNA

motif is highlighted in red. 300µM of each compound was used per lane.

B) Compounds B, B-1, B-2, B-3, and B-4 were tested for DNA major groove intercalation in an

ethidium bromide exclusion assay. DRAQ5 was used as a positive control for intercalation.

Each assay was performed in triplicate. Error bars represent standard deviation of the mean.

Movies S1-S5 Molecular dynamics simulations for Compound B and Compound B

analogues B-1, B-2, B-3, and B-4 interaction with AP2-EXP

Compound B (Movie S1), Compound B-1 (Movie S2), Compound B-2 (Movie S3), Compound B-

3 (Movie S4), and Compound B-4 (Movie S5) interactions with AP2-EXP were each simulated

using molecular dynamics. Each molecule was initially started in the location predicted for

Compound B by molecular docking.

Figure S9. Creation of endogenously tagged parasite lines AP2-EXP::GFP and AP2-

EXP::HA

A) The Selection Linked Integration system was used to add 2xFKBP inducible mislocalization

protein and GFP to AP2-EXP by single homologous recombination. Successful integration was

confirmed by genotyping PCR. Genomic DNA from the wild type Pf3D7 parental line was used

as a control. In order to test the efficacy of the knock sideways system, the pLyn mislocalizer

plasmid was added to AP2-EXP::GFP and confirmed by PCR. DNA kb are indicated by the

marks to the left of the gel.

B) The PSN054 TetR:DOZI plasmid was used to add the TetR:DOZI mRNA repression module

and endogenous 2xHA tag to AP2-EXP by double homologous recombination. Correct

integration was confirmed by genotyping PCR. The parental NF54 parasite line was used as the

unedited control. Clonal populations one and two are indicated as C1 and C2, respectively.

Clone one had correct integration and complete absence of the wild type *ap2-exp* DNA locus

and was used for further experiments. DNA kb are indicated by the marks to the right of the gel.

Figure S10. AP2-EXP protein expression in the AP2-EXP::GFP endogenously tagged

parasite line (related to figure 3A)

AP2-EXP expression was tracked throughout the IDC by harvesting protein from highly

synchronous asexual blood stage parasites followed by a western blot against the GFP tag.

Histone H3 was used as a loading control.

Figure S11. Fluorescent microscopy tracking of AP2-EXP::GFP expression

AP2-EXP::GFP expression was monitored in a highly synchronous parasite population by

fluorescent microscopy across the IDC. DRAQ5 was used as a nuclear stain for parasites.

Figure S12. AP2-EXP protein expression in the AP2-EXP::HA endogenously tagged

parasite line

AP2-EXP expression was tracked throughout the IDC by harvesting protein from highly

synchronous asexual blood stage parasites followed by a western blot against the 2xHA tag.

Histone H3 was used as a loading control.

Figure S13. Parasite counts for 48-hour Compounds B, C, and H phenotyping time

course, related to figure 3B

A) Schematic for AP2 domain competitor compound phenotyping time course.

B) 40µM of ApiAP2 competitor Compounds B or C, or non ApiAP2 competitor Compound H,

was spiked into highly synchronous wild type Pf3D7 parasites at 24 hours post invasion. DMSO

vehicle was used as a reference for normal growth. All growth assays were performed in

triplicate. Error bars represent standard deviation of the mean.

Figure S14. Compound C causes parasites to stop progressing in the early trophozoite

stage at 30 hpi

Highly synchronous wild type Pf3D7 parasites were spiked with DMSO vehicle control (A) or

40µM Compound C (B) and monitored at 6-hour intervals throughout the IDC. Each growth

assay was performed in triplicate. Error bars represent standard deviation of the mean. Giemsa

stained parasite images are representative of each population at the specified time point. R, ET,

T, LT, and S indicate ring, early trophozoite, late trophozoite, or schizont morphologies,

respectively.

Figure S15. Quality control of DNA microarray data for DMSO vehicle control and

Compound C parasites, related to figure 4

A) DNA microarray data from(34) for a set of highly periodic control genes expressed in the

IDC.

B-C) The same set of highly periodic control genes as in panel A were plotted for the DMSO

control and 12μM Compound C spiked parasites in order to compare parasite staging between

the two experiments.

D) Correlogram depicting the Spearman Correlation value between control gene expression for

DMSO (Panel B) and Compound C (Panel C) samples. A * indicates p value < .05.

Figure S16. A 48-hour time course to determine the phenotype for 12µM Compound C, related to Figure 4A

A) Highly synchronous asexual blood stage Pf3D7 parasites were spiked with DMSO vehicle control, 12µM Compound C, or 40µM Compound C. 40µM Compound C was used as a positive control for complete parasite death at 48 hpi. Each growth assay was performed in triplicate. Error bars represent standard deviation of the mean.

B) Representative images of each parasite population (DMSO, 12µM Compound C, 40µM Compound C) at 24 and 48 hpi.

Figure S17. ChIP-seq protein quality control, related to figure 4B

A) Crosslinked nuclear material was blotted after sonication to ensure recovery of the full-length

AP2-EXP protein during chromatin immunoprecipitation. Full length AP2-EXP is recovered,

indicating that the protocol is suitable to analyze AP2-EXP DNA binding *in vivo*. Crosslinked

nuclear material from the wildtype Pf3D7 parental parasite line was used as a negative control.

B) Anti-GFP beads were used to pull down GFP tagged AP2-EXP from AP2-EXP::GFP.

Flowthrough (FT), Wash (W1 and W2) and Eluate fractions were saved and analyzed by

western blot. The presence of AP2-EXP and Histone H3 in the Eluate lane indicates that AP2-

EXP interacts with chromatin in the nucleus. The non-immune negative control beads do not

enrich AP2-EXP or Histone H3 in the eluate.

Figure S18. ChIP-seq extended data

A) Log2 immunoprecipitate/input ChIP-seq data from each replicate (2x AP2-EXP::GFP and

1xAP2-EXP::HA) of AP2-EXP ChIP-seq visualized by IGV at a representative DNA locus. The

location of a conserved MACS2 called peak of occupancy and the TGCATGCA DNA motif is

indicated by the bottom tracks. The (-) control lane is the coverage resulting from a no-epitope

control ChIP-seq.

B) The top three ranked DNA motifs present within peaks of occupancy for each ChIP-seq

replicate as determined by DREME(24). The core DNA motif CATGCA is overrepresented

within each individual replicate.

C) The top overrepresented DNA motif within AP2-EXP peaks of occupancy conserved in 2/3

replicates of ChIP-seq as determined by DREME(24) plotted at the primary DNA sequence

level. DNA sequences were sorted from highest to lowest degree of motif conservation using

FIMO.(25)

Figure S19. Comparison of AP2-EXP target genes with Compound C induced changes in

transcript abundance

The total overlap between AP2-EXP gene targets detected in the Compound C RNA time

course and global decrease in transcript abundance at 24-30 hpi by $\log_2$ fold change >2.

Figure S20. ChIP-Quantitative PCR to assess Compound C impact on AP2-EXP genomic occupancy

AP2-EXP::GFP parasites were spiked with 40µM Compound C or DMSO vehicle control at 30 hpi for two hours. ChIP samples were collected for each population using either anti-GFP or negative control IgG antibodies. The percent of input was determined by RT-qPCR. Each experiment was performed in triplicate. The presence or absence of an AP2-EXP peak of occupancy at each DNA locus based on ChIP-seq is indicated by a (+) or (-), respectively. Each assay was done in triplicate with the exception of AMAI, where n =2.

Figure S21. Nucleosome occupancy is depleted at AP2-EXP DNA binding sites

Mnase-seq data(30) was plotted against DNA binding sites conserved in 2/3 replicates of AP2-

EXP ChIP-seq (A) or random genomic intervals of equal length from the same chromosome as

the original peak (B).

Figure S22. Histone post translational modifications and chromatin reader occupancy at

AP2-EXP peaks

The occupancy of histone variant H2A.z, histone modifications H3K9ac, H3K4me3(27),

H3K9me3, H3K36me2/3(26), and chromatin readers BDP1(29) and HP1(31) were plotted

against AP2-EXP peaks of occupancy conserved in 2/3 ChIP-seq replicates (A) or random

genomic intervals of the same length (B), taken from the same chromosome on which the AP2-

EXP peak originally occurred.

Figure S23. AP2-EXP DNA occupancy with respect to the Transcription Start Site (TSS) of

target genes

Log_2 immunoprecipitate (ChIP)/Input ChIP-seq coverage for each replicate of AP2-EXP ChIP-

seq and the no epitope control was plotted against the TSS(35) of each target gene conserved

in 2/3 ChIP-seq replicates.

Figure S24. Normal Transcript Abundance of AP2-EXP Target Genes

AP2-EXP target genes were determined by ChIP-seq and their transcript abundance data

during the 48-hour IDC was plotted using data from Chappell *et al*(35).

Figure S25. Creation of a *glms* ribozyme based knockdown line for AP2-EXP

Wildtype Pf3D7 *Plasmodium falciparum* parasites were transfected with pSLI:AP2-

EXP*::glms*::HA to endogenously tag the AP2-EXP DNA locus with an inducible *glms* ribozyme

and HA epitope tag. Successful integration to create AP2-EXP::*glms*::HA by single crossover

homologous recombination was confirmed by genotyping PCR. C1 and C2 represent clonal

populations selected for integration. WR represents a parasite population selected for the

plasmid but not for integration. WT represents Pf3D7 wild type control parasite gDNA.

Figure S26. Western blot phenotyping of attempts to genetically knockdown AP2-EXP

A) To assess genetic knockdown of AP2-EXP by *glms* ribozyme tag in the parasite line AP2-

EXP::*glms*::HA, highly synchronous parasites were spiked with 5mM glucosamine or vehicle

control for 72 hours and AP2-EXP quantity was determined by anti-HA western blot. Histone H3

was used as a loading control. Glucosamine treatment did not impact the amount of AP2-EXP

protein present.

B) Genetic knockdown of AP2-EXP by the TetR:DOZI mRNA repression module was assessed

in the parasite line AP2-EXP::HA by washing anhydrotetracycline (aTc) out of the media for 72

hours. AP2-EXP quantity was determined by anti-HA western blot, and Histone H3 was used as

a loading control. Removal of aTc from the media did not impact the amount of AP2-EXP

protein present.

C) Genetic knockdown of AP2-EXP via protein mislocalization was assessed for the parasite

line AP2-EXP::GFP. 250nM rapalog was added to the media for 48 hours and AP2-EXP protein

localization was assessed by ant-GFP western blot. Histone H3 and Aldolase were used as

nuclear and cytosolic markers, respectively. The addition of rapalog did not cause any

detectable mislocalization of AP2-EXP from the nucleus to the cytosol. N indicates the nuclear

protein fraction, and C indicates the cytosolic fraction.

Figure S27. Mosquito stage *P. berghei* inhibition assay schematic

A. stephensi mosquitoes were infected with *Plasmodium berghei* parasites. On day 10 post

infection, mosquito midguts were injected with the measured IC50 of Compounds B, C, F, or

DMSO vehicle control. On day 14 post infection mosquitoes were dissected to count oocysts

and midgut sporozoites. For Compound C and DMSO vehicle control, each experiment was

performed in duplicate. Compounds B and F phenotyping were performed as a single

experiment.

References

1.   F. J. Gamo, *et al.*, Thousands of chemical starting points for antimalarial lead identification. *Nature* **465**, 305–
310 (2010).

2.   D. S. Wishart, *et al.*, DrugBank 5.0: A major update to the DrugBank database for 2018. *Nucleic Acids Res.*
46, D1074–D1082 (2018).

3.   R. Huey, G. M. Morris, Using AutoDock 4 with AutoDockTools : A Tutorial. 1–56.

4.   J. Wang, R. M. Wolf, J. W. Caldwell, P. A. Kollman, D. A. Case, Development and testing of a general Amber
force field. *J. Comput. Chem.* **25**, 1157–1174 (2004).

5.   J. Wang, W. Wang, P. A. Kollman, D. A. Case, Automatic atom type and bond type perception in molecular

mechanical calculations. *J. Mol. Graph. Model.* **25**, 247–260 (2006).

6. P. A. K. and D. A. C. Wang, J., W. Wang, Antechamber: an accessory software package for molecular mechanical calculations. *J. Am. Chem. Soc* **222**, U403 (2001).

7. D. Y. and P. K. Case, D., K. Belfon, I. Ben-Shalom, S. Brozell, D. Cerutti, T. Cheatham III, V. Cruzeiro, T. Darden, R. Duke, G. Giambasu, M. Gilson, H. Gohlke, A. Goetz, R. Harris, S. Izadi, S. Izmailov, K. Kasavajhala, A. Kovalenko, R. Krasny, T. Kurtzman, T. Lee, S. L, AMBER 2020 (2020).

8. A. Pérez, *et al.*, Refinement of the AMBER force field for nucleic acids: Improving the description of α/γ conformers. *Biophys. J.* **92**, 3817–3829 (2007).

9. A. W. Götz, *et al.*, Routine microsecond molecular dynamics simulations with AMBER on GPUs. 1. generalized born. *J. Chem. Theory Comput.* **8**, 1542–1555 (2012).

10. W. R. Salomon-Ferrer R, Götz AW, Poole D, Le Grand S, Routine Microsecond Molecular Dynamics Simulations with AMBER on GPUs. 2. Explicit Solvent Particle Mesh EwaldNo Title. *J Chem Theory Comput.* **10**, 3878–3888 (2013).

11. J. P. Ryckaert, G. Ciccotti, H. J. C. Berendsen, Numerical integration of the cartesian equations of motion of a system with constraints: molecular dynamics of n-alkanes. *J. Comput. Phys.* **23**, 327–341 (1977).

12. D. R. Roe, T. E. Cheatham, PTRAJ and CPPTRAJ: Software for processing and analysis of molecular dynamics trajectory data. *J. Chem. Theory Comput.* **9**, 3084–3095 (2013).

13. G. K. Smyth, Linear models and empirical Bayes methods for assessing differential expression in microarray experiments. *Stat. Appl. Ge- netics Mol. Biol.* **3**, Article 3 (2011).

14. D. P. Sangurdekar, The Rnits package for normalization and inference of differential expression in time series microarray data . *Rvignette*, 1–13 (2014).

15. A. J. Saldanha, Java Treeview - Extensible visualization of microarray data. *Bioinformatics* **20**, 3246–3248 (2004).

16. M. J. L. de Hoon, S. Imoto, J. Nolan, S. Miyano, Open source clustering software. *Bioinformatics* **20**, 1453–1454 (2004).

17. A. M. Bolger, M. Lohse, B. Usadel, Trimmomatic: A flexible trimmer for Illumina sequence data. *Bioinformatics* **30**, 2114–2120 (2014).

18. S. Andrews, FastQC: A Quality Control Tool for High Throughput Sequence Data. http://www.bioinformatics.babraham.ac.uk/projects/fastqc (2010).

19. H. Li, Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM. **00**, 1–3 (2013).

20. H. Li, *et al.*, The Sequence Alignment/Map format and SAMtools. *Bioinformatics* **25**, 2078–2079 (2009).

466   21.   J. M. Gaspar, Improved peak-calling with MACS2. *bioRxiv*, 1–16 (2018).

467   22.   A. R. Quinlan, I. M. Hall, BEDTools: A flexible suite of utilities for comparing genomic features. *Bioinformatics*

468   **26**, 841–842 (2010).

469   23.   F. Ramírez, F. Dündar, S. Diehl, B. A. Grüning, T. Manke, DeepTools: A flexible platform for exploring deep-

470   sequencing data. *Nucleic Acids Res.* **42**, 187–191 (2014).

471   24.   T. L. Bailey, DREME: Motif discovery in transcription factor ChIP-seq data. *Bioinformatics* **27**, 1653–1659

472   (2011).

473   25.   C. E. Grant, T. L. Bailey, W. S. Noble, FIMO: Scanning for occurrences of a given motif. *Bioinformatics* **27**,

474   1017–1018 (2011).

475   26.   L. Jiang, *et al.*, PfSETvs methylation of histone H3K36 represses virulence genes in Plasmodium falciparum.

476   *Nature* **499**, 223–227 (2013).

477   27.   R. Bártfai, *et al.*, H2A.Z demarcates intergenic regions of the Plasmodium falciparum epigenome that are

478   dynamically marked by H3K9ac and H3K4me3. *PLoS Pathog.* **6**, e1001223 (2010).

479   28.   D. F. Read, K. Cook, Y. Y. Lu, K. G. Le Roch, W. S. Noble, Predicting gene expression in the human malaria

480   parasite Plasmodium falciparum using histone modification, nucleosome positioning, and 3D localization

481   features. *PLoS Comput. Biol.* **15**, e1007329 (2019).

482   29.   G. A. Josling, *et al.*, A Plasmodium Falciparum Bromodomain Protein Regulates Invasion Gene Expression.

483   *Cell Host Microbe* **17**, 741–751 (2015).

484   30.   E. M. Bunnik, *et al.*, DNA-encoded nucleosome occupancy is associated with transcription levels in the

485   human malaria parasite Plasmodium falciparum. *BMC Genomics* **15**, 347 (2014).

486   31.   S. A. Fraschka, *et al.*, Comparative Heterochromatin Profiling Reveals Conserved and Unique Epigenome

487   Signatures Linked to Adaptation and Development of Malaria Parasites. *Cell Host Microbe* **23**, 407–420

488   (2018).

489   32.   X. Shang, *et al.*, Genome-wide landscape of ApiAP2 transcription factors reveals a heterochromatin-

490   associated regulatory network during Plasmodium falciparum blood-stage development. *Nucleic Acids Res.*

491   **50**, 3413–3431 (2022).

492   33.   S. E. Lindner, E. K. De Silva, J. L. Keck, M. Llinás, Structural Determinants of DNA Binding by a P.

493   falciparum ApiAP2 Transcriptional Regulator. *J. Mol. Biol.* **395**, 558–567 (2010).

494   34.   M. Llinás, Z. Bozdech, E. D. Wong, A. T. Adai, J. L. DeRisi, Comparative whole genome transcriptome

495   analysis of three Plasmodium falciparum strains. *Nucleic Acids Res.* **34**, 1166–1173 (2006).

496   35.   L. Chappell, *et al.*, Refining the transcriptome of the human malaria parasite Plasmodium falciparum using

497   amplification-free RNA-seq. *BMC Genomics* **21**, 395 (2020).