Title

Identification of Antimalarial Compounds that Inhibit Apicomplexan AP2 Proteins

Word Count: 11,627

Authors

Timothy Russell[1,2,3,4], Erandi K. De Silva[6], Valerie Crowley[1], Kathryn Shaw-Saliba[7], Namita
Dube[1], Gabrielle Josling[1,3,4], Charisse Flerida A. Pasaje[8], Irene Kouskoumvekaki[9], Gianni
Panagiotou[10,11], Jacquin C. Niles[8], Marcelo Jacobs-Lorena[7], C. Denise Okafor[1,5], Francisco-
Javier Gamo[12], Manuel Llinás[1,2,3,4,5]

1.   Department of Biochemistry and Molecular Biology, Pennsylvania State University.

2.   Huck Institutes Center for Eukaryotic Gene Regulation (CEGR), Pennsylvania State
University.

3.   Huck Institutes Center for Malaria Research (CMaR), Pennsylvania State University.

4.   Huck Institutes Center for Infectious Disease Dynamics, Pennsylvania State University.

5.   Department of Chemistry, Pennsylvania State University.

6.   Lewis-Singler Institute for Integrative Genomics, Princeton University.

7.   Department of Molecular Biology and Immunology, Malaria Research Institute, Johns
Hopkins Bloomberg School of Public Health.

8.   Department of Biological Engineering, Massachusetts Institute of Technology.

9.   Department of Systems Biology, Technical University of Denmark, Kgs. Lyngby,
Denmark.

10.   Systems Biology and Bioinformatics, Leibniz Institute for Natural Products Research and
Infection Biology, Hans Knöll Institute.

11.   Department of Medicine, the University of Hong Kong, Hong Kong SAR, China

12.   Global Health Medicines, GlaxoSmithKline, Tres Cantos, Spain.

Abstract

Plasmodium parasites are reliant on the Apicomplexan AP2 (ApiAP2) transcription

factor family to regulate gene expression programs. The AP2 DNA binding domains

have no homology to the human or mosquito hosts, making them potential drug targets.

Using an *in-silico* screen to dock thousands of small molecules into the crystal structure

of the AP2-EXP (Pf3D7_1466400) AP2 domain (PDB:3IGM), we identified compounds

that interact with this domain. Four compounds were found to compete for DNA binding

with AP2-EXP and at least one additional ApiAP2 protein. Our top ApiAP2 competitor

compound perturbs the transcriptome of *P. falciparum* trophozoites and results in a

decrease in abundance of $\log_2$ fold change > 2 for 50% (46/93) of AP2-EXP target

genes identified. Additionally, two ApiAP2 competitor compounds have anti-

Plasmodium activity against *P. berghei* mosquito stage parasites. In summary, we

describe a novel set of antimalarial compounds that are targeted against the ApiAP2

family of proteins. These compounds may be used for future chemical genetic

interrogation of ApiAP2 proteins or serve as starting points for a new class of

antimalarial therapeutics.

Introduction

Malaria is a disease that is caused by intracellular parasites of the genus *Plasmodium*

and represents a significant health and economic burden worldwide[1]. The most virulent

of the human infectious malaria parasites is *Plasmodium falciparum*, which caused

more than 200 million cases and resulted in over 600 thousand deaths in 2021[1].

Unfortunately, resistance has been reported for every antimalarial therapeutic deployed

to date, necessitating the need for malaria drugs that target new parasite processes[2].

To successfully proliferate, *P. falciparum* parasites must develop through a complex

lifecycle that includes intracellular and extracellular stages in the human and *Anopheles*

mosquito hosts[3]. To initiate an infection, *Plasmodium* sporozoites are deposited into the

bloodstream of the human host when an infected mosquito takes a blood meal.

Sporozoites invade hepatocytes and replicate into daughter merozoites, which egress to

invade red blood cells (RBCs). Inside RBCs the parasite replicates asexually over the

course of ~48 hours, developing through three major morphological stages referred to

as: ring, trophozoite, and schizont. Schizonts rupture at 48 hours post invasion (hpi),

releasing 16-32 daughter merozoites into the blood stream. Merozoites reinvade new

RBCs and begin replicating again. This iterative invasion and replication, called the

intraerythrocytic development cycle (IDC), is responsible for the periodic fevers

associated with malaria due to destruction of RBCs[3]. In each round of replication up to

10% of parasites will commit to a sexual development program which results in non-

replicating gametocytes[4]. Gametocytes are the only blood stage that are competent to

re-establish infection in a mosquito upon ingestion in the blood meal. Male and female

gametocytes develop into gametes that fuse to become a zygote, which ultimately

develops back into sporozoites, thereby completing the *Plasmodium* lifecycle.

Transcriptomics-based evidence has shown that up to 80% of *P. falciparum* protein

coding transcripts are developmentally regulated during the IDC, with peaks of

expression occurring at specific IDC stages[5–9]. This pattern has been described as a

"just-in-time" cascade due to the observation that for many transcripts, the timing of

maximal abundance precedes the developmental stage at which the protein product is

required for progression through the IDC[6,9,10]. Transcriptional regulation in *Plasmodium*

parasites is driven through a combination of modulation of the chromatin landscape[11–17],

the general transcriptional machinery[18,19], and sequence-specific transcription

factors[20,21]. Although many of the general transcription factors (GTFs) required for RNA

Polymerase II activity are conserved between *Plasmodium* and model eukaryotes[19,22,23],

Plasmodium parasites encode seemingly few sequence-specific DNA binding

transcription factors.

The major family of sequence specific transcription factors is the 27-member family of

Apicomplexan Apetala AP2 (ApiAP2) proteins[21,24,25]. ApiAP2 proteins all contain one to

three ~60 amino acid AP2 DNA binding domains that are analogous to the plant

APETALA2/Ethylene Responsive Factor (AP2/ERF) domain[22]. Their plant-like origin

distinguishes them from all other known transcription factors in humans or

mosquitoes[20]. Over half of the ApiAP2 proteins are predicted to be essential for the IDC

in *P. falciparum* based on a random mutagenesis screen[26], and in targeted knockout

screens only 11/23 ApiAP2 coding sequence disruptions attempted were successful[27–

29]. In the rodent malaria model species *P. berghei*[30] and *P. yoelii*[31], 11 ApiAP2 proteins

have developmental lethal phenotypes in the sexual blood or mosquito stages. ApiAP2

transcription factors can cause activation or repression of transcription of specific

genes, and they are involved in diverse lifecycle processes including invasion, mutually

exclusive antigenic gene expression, sexual differentiation, and heat stress

tolerance[27,32–40]. Chromatin interacting proteins such as histone deacetylases and

methyltransferases have received recent attention as potential drug targets against

malaria parasites[41,42]. No efforts to date have focused on targeting the sequence

specific DNA binding ApiAP2 transcription factors, which are potential drug targets due

to their essentiality for the asexual blood stage and lack of conservation with the host.

In this study we conducted an *in silico* chemical screen against the published crystal

structure of the AP2-EXP AP2 domain[43] to predict competitors of DNA binding activity.

AP2-EXP is predicted to be essential for the *P. falciparum* asexual blood stage[27,29,33]

and its orthologue *P. berghei* AP2-Sp is a master regulator of sporogony[44]. Several

compounds identified in the screen can compete for DNA binding activity with the AP2-

EXP AP2 domain *in vitro*. Each of these compounds competes DNA binding by at least

one additional ApiAP2 protein, suggesting the potential for multi-ApiAP2 inhibition. We

also find that modifications to the substitution state of an AP2 domain competitor

compound can modulate its ability to compete DNA binding. Our lead compound kills *P.*

falciparum parasites in concordance with the stage specific expression of AP2-EXP. We

measured AP2-EXP genomic occupancy by ChIP-seq and found high correlation

between dysregulated mRNA transcripts in the presence of our lead compound and

AP2-EXP bound genes, corroborating the hypothesis that AP2-EXP is inhibited *in vivo*.

Finally, we find that two AP2 domain competitor compounds are also active against

Plasmodium parasite development in the mosquito stage. Overall, our results

demonstrate the potential to chemically compete the DNA binding activity of ApiAP2

proteins for the purposes of providing proof-of-concept for a new antimalarial strategy,

and interrogation of the function of *P. falciparum* blood stage essential ApiAP2 proteins.

Results

***In silico* Computational Modeling of AP2-EXP Competitors**

To identify competitors of DNA binding by an ApiAP2 protein, we used AP2-EXP

(PF3D7_1466400), which is the only ApiAP2 protein whose AP2 domain is structurally

characterized (PDB accession: 3IGM). *In silico* molecular docking was run on

AUTODOCK[45] using over ten thousand small molecules from both the Tres Cantos

Antimalarial Set (TCAMS)[46] and the Drug Bank[47] against the crystal structure of the

AP2-EXP AP2 protein domain (**Fig 1A, Fig S1**). Docking hits were prioritized based on

the free energy of interaction and proximity to amino acids necessary for base-specific

DNA contacts. In some cases, the compound that generated a high docking score was

not available for purchase, so alternative choices with >0.9 Tanimoto similarity score

were identified, computationally docked, and used for further evaluation **(All**

compounds listed in Table S1). These docking simulations resulted in a final list of

nine top scoring predicted competitors of DNA binding by AP2-EXP. Each compound

was assigned an alphabetical identifier A-I **(Fig 1B)** for use in this study. The list

contains both known antimalarials from the Tres Cantos Antimalarial Set and molecules

from the Drug Bank that had not been previously tested for antimalarial activity. To

assess antimalarial activity, we assayed *P. falciparum* parasite viability when exposed

to each of the putative AP2-EXP competitors and determined that all nine compounds

kill asexual *P. falciparum* parasites at micromolar concentrations (range 11-170µM) in a

48-hour asexual stage growth inhibition assay **(Fig 1B, Fig S2)**.

Four Predicted AP2-EXP Competitors Compete DNA Binding *in vitro*

The nine predicted competitors of DNA binding by the AP2-EXP AP2 domain were

tested in a competitive electrophoretic mobility shift assay (EMSA). The minimum mass

of AP2-EXP required to visualize DNA binding **(Fig S3A)** was mixed with each

compound and incubated prior to adding the DNA oligonucleotide **(DNA Oligos in**

Table S2). Competition of DNA binding activity should result in a loss of signal for the

higher molecular weight DNA probe which is bound by AP2-EXP. Four of the nine

compounds (A, B, C and I) **(Fig 2A)** were able to effectively compete for DNA binding

by AP2-EXP *in vitro*. Three of the four AP2 domain competitors (Compounds A, B and

C) have a benzoxazole core moiety, while Compound I is made up of planar rings **(Fig**

2B).

To assess whether DNA binding competition was specific to the AP2-EXP AP2 domain,

we repeated competitive EMSAs with three different purified *P. falciparum* AP2

domains: AP2-I Domain 3 (AP2-I D3)[25,32] **(Fig S3B)**, AP2-HS Domain 1 (AP2-HS

D1)[25,36] **(Fig S3C)**, and PfSIP2 Domain 1 (PfSIP2 D1)[35] **(Fig S3D)**, again using the

minimum amount of protein needed to visualize binding of each ApiAP2 protein's

cognate DNA oligonucleotide **(Table S2)**. AP2-I D3[32] and PfSIP2 D1[35] are essential for

the IDC, while *ap2-hs* deletion is tolerated when parasites are grown below

physiological temperatures[36]. We found that Compounds A, B, C, and I can compete

DNA binding by AP2-I **(Fig S4A)**, Compounds B and I compete DNA binding by AP2-

HS D1 **(Fig S4B)** and Compounds B, C and I compete DNA binding by PfSIP2 D1 **(Fig**

S4C).

As a control for compound specificity, we performed competitive EMSAs using the non-

Plasmodium AP2 domain protein *Arabidopsis thaliana* Ethylene Response Factor 1[48]

(AtERF1) **(Fig S3E)** and the High Mobility Group Box (HMGB) domain protein SOX2

(Fig S3F)[49]**.** SOX2 is a human encoded transcription factor and has no homology to the

AP2 domain. Only Compound I can compete AtERF1 **(Fig S5A)**. SOX2 is competed by

Compound A only **(Fig S5B)**. Overall, Compounds B and C have specificity for

Apicomplexan AP2 domains *in vitro*. (EMSA Results Summarized in **Table S3**.)

To test whether the ApiAP2 competing compounds interfere non-specifically with DNA

binding proteins by intercalating with DNA, we conducted an ethidium bromide

exclusion assay[50]. In this assay, if the molecule of interest can compete away ethidium

bromide from intercalating DNA, it will reduce ethidium bromide fluorescence.

Compared to the positive control DRAQ5[50], only Compound I was able to effectively

displace ethidium bromide **(Fig S6)**. Based on these results, we focused on Compounds

B and C due to their specificity for *P. falciparum* AP2 domain proteins and their lack of

DNA intercalation.

<u>Structural Analogs of Compound B Differ in DNA Binding Competition Activity</u>

We leveraged the availability of compounds with high structural similarity to Compound

B due to its origin in the Tres Cantos Antimalarial Set[46] to determine whether specific

features of the chemical structure impact DNA binding competition. We tested the 4

closest available analogs to Compound B from the Tres Cantos Antimalarial Set based

on similarity score[46] (Designated as Compounds B-1, B-2, B-3 and B-4) **(Fig 3A)** for

competition against AP2-EXP. Compounds B-1 and B-4 compete DNA binding by AP2-

EXP **(Fig 3B)** and AP2-I D3 **(Fig S7A)**, while Compounds B-2 and B-3 do not compete

DNA binding by either AP2-EXP or AP2-I D3 **(Fig 3B, S7A)**. Compound B-1 is the least

effective competitor relative to Compounds B and B-4 **(Fig 3C)**. Non-AP2 domain

competitor Compounds B-2 and B-3 each have halogen atoms substituted on the

benzene ring in the para and ortho, or para positions of the benzene ring, respectively.

Conversely, Compounds B, B-1, and B-4, have methyl or ethyl groups substituted onto

the benzene ring and can compete DNA binding **(Fig 3D)**. Compound B-1 has both a

methyl group and a chlorine atom substitution. If the halogen atom substitution

decreases DNA binding competition, this mix of substitutions is consistent with its lower

efficacy of DNA binding competition relative to Compounds B and B-4**.** Compounds B,

B-1 and B-4 have no difference in DNA intercalation ability as measured by ethidium

bromide exclusion **(Fig S7B)**. We followed up on these results by running molecular

dynamics simulations on AP2-EXP docked with Compounds B,B-1, B-2, B-3, and B-4.

While Compound B remains stably docked with AP2-EXP over the course of the

simulation, Compound B-3 moves away from the protein and does not re-dock

(Supplemental Movies 1-5). Compounds B-1, B-2, and B-4 remain stably associated

with AP2-EXP in the simulation **(Supplemental Movies 1-5)**. In summary, substituting

bulky, electronegative groups onto Compound B can abolish its ability to compete for

DNA binding with AP2-EXP, potentially due to a steric clash.

Compound C Activity Coincides With the Expression of AP2-EXP

To assess whether Compounds B and C are likely to share a common ApiAP2 target,

we determined their parasite killing phenotypes against *P. falciparum* during the IDC.

Compound H was used as a control since it does not compete AP2-EXP AP2 domain

DNA binding **(Fig 2A)**. Each compound was spiked (40μM) into synchronous asexual

blood stage *P. falciparum* parasites at 24 hours post invasion and parasite progression

was determined at 24 and 48 hours post spike-in **(Fig 4A)**. At 24 hours post spike-in,

parasites in the presence of either Compounds B or C had failed to progress past the

mid-trophozoite stage and did not morphologically progress any further by 48 hours post

spike-in **(Fig 4B)**. In contrast, parasites spiked with Compound H progressed through

one IDC before failing to reinvade in the next cycle **(Fig 4B)**. Compounds B and C

therefore may have a common protein target as they exhibit the same morphological

impact on the asexual blood stage and similar timing of action. As expected, each

compound exposure prevented further parasite growth within 48 hours of spike-in **(Fig**

S8).

Due to its commercial availability, we then continued with Compound C as the lead

compound for phenotypic and molecular characterization. To determine the precise

timing of Compound C antimalarial activity, synchronous parasites were monitored

throughout the 48-hour IDC in the presence of either 40µM Compound C or DMSO

control after spiking in Compound C at 6 hpi. While the control parasites progressed

normally, the Compound C spiked parasites failed to progress beyond the mid-

trophozoite stage at 24-30 hours post invasion **(Fig S9)**. To further narrow down the

timing of the anti-*Plasmodium* action of Compound C, 40µM of Compound C was spiked

into a parasite culture at 8-hour intervals beginning at 14 hpi. Parasite growth was

quantified in the next cycle by counting ring stage parasites at 54 hpi. This growth assay

revealed that Compound C has no antimalarial effect after 46 hpi **(Fig 4C).** In support of

this result, washing 40µM Compound C out of culture at 8-hour intervals beginning at 14

hpi only rescued parasite growth until 30 hpi, albeit with diminishing effectiveness as the

parasites approached 30 hpi **(Fig 4D).** Taken together, these results suggest that

Compound C maximally inhibits of IDC progression of *Plasmodium* parasites between

30-46 hpi.

To validate the stage in the IDC at which AP2-EXP is expressed, we used Selection

Linked Integration (SLI)[51] to introduce an endogenous C-terminal tag containing a

2XFK506-Binding Protein (FKBP) dimerization domain and GFP **(Fig S10A)** to create

AP2-EXP::GFP. We observed maximal abundance of AP2-EXP::GFP in 30 hpi

trophozoites by both Western blot and live fluorescence microscopy **(Fig 4E, F).**

Maximal protein abundance of AP2-EXP in mid-trophozoites was independently

confirmed in a different parasite line expressing AP2-EXP tagged with 2XHA and the

TetR:DOZI mRNA repression aptamer (AP2-EXP::HA) **(Fig S10B, Fig S11)**. AP2-EXP

was not detected in maturing Stage III gametocytes **(Fig 4E)**. These results

demonstrate that the maximal protein abundance of AP2-EXP at 30hpi coincides with

the timing of action for Compound C.

Compound C Disrupts the Parasite Transcriptome with Stage Specificity

Since AP2-EXP is predicted to act as a transcription factor[29,33] we hypothesized that

changes in parasite transcript abundance should occur when AP2-EXP DNA binding is

competed by Compound C. To test this hypothesis, we cultured parasites in 12µM

($0.66 \times IC_{50}$) Compound C or DMSO vehicle and measured comparative RNA abundance

by DNA microarray at seven time points during the asexual blood stage. As a quality

control for normal IDC progression, we found a high correlation between a set of

"control" genes[52,53] with an established 48-hour periodicity with maxima at different

times throughout the 3D7 *P. falciparum* IDC[6] in our DMSO control and Compound C

mRNA abundance profiles **(Fig S12A-D)**. Phenotypically, 12μM of Compound C kills

less than 50% of parasites in one asexual cycle **(Fig S13A, B)**. To identify perturbations

to parasite RNA abundance, we compared all detected transcripts at each time point

assayed between the DMSO control and Compound C dosed parasites **(Expression**

values in Table S4). The Spearman Correlation between samples deviated most

significantly at 30hpi, (Corr. = -0.16) **(Fig 5A)** as expected from the compound activity

measured **(Fig 4B-4D)**.

To follow up on this observation, we quantified differences in transcript level that are

unique to the progression from 24 to 30 hpi timepoints for Compound C or DMSO

vehicle control populations using Linear Models for Microarray Data (LIMMA[54])

(Differentially Expressed genes in Table S4). Overall, 463 RNA transcripts are

decreased in abundance by $\log_2$ fold change >2 between 24-30 hpi in the presence of

Compound C. Gene ontology term (GO-term) analysis[55] of the differentially regulated

transcripts **(Full GO Terms in Table S4)** revealed an enrichment for genes that encode

proteins important for the parasite to invade or modify red blood cells (e.g., RON3,

GAP45, RhopH2, RhopH3, MSP1, MSP6). We independently determined differential

expression over the entire time course using the RNA time course specific software

RNITS[56] and found high overlap with our LIMMA analyses **(Overlaps in Table S4)**.

Therefore, the maximal perturbation of the transcriptome **(Fig 5A)**, action of Compound

C **(Fig 4B)**, and maximal abundance of AP2-EXP **(Fig 4E, F)** all occur at roughly 30 hpi

in the IDC.

AP2-EXP Genomic Occupancy Correlates with Compound C Induced Transcript

Dysregulation

We used chromatin immunoprecipitation followed by high throughput sequencing (ChIP-seq) to determine the genome wide binding occupancy of AP2-EXP. A total of three ChIP-seq samples from highly synchronous trophozoites were collected at 30 hpi using both the AP2-EXP::GFP (2 replicates) and AP2-EXP::HA (1 replicate) parasite lines. Immunoprecipitation of full length AP2-EXP was confirmed by Western blot **(Fig S14A)**, and a co-immunoprecipitation blot confirmed that AP2-EXP interacts with Histone H3 **(Fig S14B)**. Peaks of occupancy were determined for each replicate using the MACS2 peak calling program[57] **(List of Peaks in Table S5)**. Peaks detected in 2/3 ChIP-seq replicates were considered for downstream analysis. In aggregate, AP2-EXP binds 240 genomic loci, corresponding to 101 total target genes **(Genes in Table S5)**. Enrichment of AP2-EXP was well conserved between the GFP and HA tagged parasite lines, as indicated by nearly identical coverage in a metagene plot of conserved AP2-EXP target genes **(Fig 5B)**. A 'no epitope control' ChIP done on a wild type 3D7 parasite line with the anti-GFP antibody detected only 2 peaks, neither of which overlapped with those of AP2-EXP::GFP or AP2-EXP::HA. Both the conserved AP2-EXP peaks and individual replicates were highly enriched for the known AP2-EXP and PbAP2-Sp DNA sequence motif CATGCA[29,58] **(Fig 5C, Fig S15A-C).**

We then determined whether there is overlap between the transcripts dysregulated in the presence of Compound C and AP2-EXP target genes identified by ChIP-seq. Out of the 463 transcripts decreased in abundance by $\log_2$ fold change > 2 in the presence of

Compound C, 46 are AP2-EXP targets **(Fig 5D)**. This represents 50% (46/93) of AP2-

EXP target genes detected in the RNA time course **(Fig 5D)**. In general, the transcripts

with the greatest decrease in abundance are AP2-EXP targets. We found that 7/14 total

transcripts decreased in abundance by $\log_2$ fold change > 5, and 16/33 decreased in

abundance by $\log_2$ fold change > 4, are AP2-EXP targets **(Fig 5E, Table S4)**. Using

GO-term analysis[55], we found that AP2-EXP targets that are dysregulated at 24-30 hpi

have functions related to red blood cell invasion and host remodeling **(Fig 5F, Table**

S4). To assess whether AP2-EXP DNA binding at ChIP-seq defined target promoter

regions is impacted by Compound C, parasite cultures were spiked with either 40µM

Compound C or DMSO vehicle control at 30 hpi and AP2-EXP occupancy was

measured by ChIP-quantitative PCR (ChIP-qPCR). AP2-EXP DNA binding decreases at

five specific peaks of occupancy (GAP45, SIP2, RON3, RALP, and AMAI) in the

presence of Compound C **(Fig S16)**, supporting a link between AP2-EXP inhibition and

Compound C.

We also compared AP2-EXP genomic occupancy (ChIP-seq) with several previously

published chromatin datasets in order to further probe its function as a sequence

specific transcription factor. AP2-EXP binds chromatin primarily in nucleosome depleted

regions[59] compared to randomized control loci **(Fig S17A, B)**. AP2-EXP occupies sites

near the Transcription Start Site (TSS) of target genes[60] **(Fig S18)**. A comparison of

RNA abundance data[12] to AP2-EXP target genes revealed that many target gene

mRNA transcripts are increased in abundance at 24-30hpi, while a subset changes to

low abundance **(Fig S19)**. Overall, AP2-EXP genomic occupancy is highly predictive of

enrichment for activating histone marks H3K9ac[17] and H3K4me3[17] as well as the

activating chromatin reader Bromodomain Protein 1 (BDP1)[15], again in comparison to

randomized control loci **(Fig S20A, B).** Although the known repressing histone markers

H3K9me3 and H3K36me2/3[13] do not correlate well with AP2-EXP occupancy,

enrichment of the Heterochromatin Protein 1 (HP1)[61] repressing chromatin reader does

overlap modestly with AP2-EXP **(Fig S20A, B)**. A GO-term analysis[55] of all AP2-EXP

target genes revealed enrichment for invasion and red blood cell modification, **(Table**

S4) as was found for the genes that decrease in abundance in the presence of

Compound C **(Fig 5F, Table S4)**. In aggregate, these findings suggest that Compound

C is able to disrupt the function of AP2-EXP as a sequence specific transcription factor.

ApiAP2 Competitor Compounds B and C are Active Against Mosquito Stage

***Plasmodium* Parasites**

Elucidation of the role of essential *P. falciparum* ApiAP2 proteins in the IDC is

challenging due to the inability to knockout asexual blood stage essential genes from

the parasite's haploid genome. Previous attempts to knockout AP2-EXP[27–29,33] failed to

produce parasite lines in which the AP2 domain coding sequence was disrupted. We

similarly did not recover transgenic parasites after attempting to completely disrupt the

coding sequence for AP2-EXP using the targeted gene disruption (pSLI-TGD) system[51]

in three independent attempts, suggesting that AP2-EXP is essential for the IDC (not

shown). We also attempted to disrupt AP2-EXP protein expression with the conditional

knockdown approaches Knock Sideways[51], TetR:DOZI mRNA repression[62], and *glms*

ribozyme mediated cleavage[63]. Each of these genetic systems failed to mediate protein

knockdown or mislocalization despite successful integration into the parasite genome

(Fig S10A, B, Fig S21, Fig S22). To account for this limitation, we tested the ApiAP2

competitor Compounds B and C against rodent infectious *Plasmodium berghei*

parasites due to the well-characterized genetic phenotype of the AP2-EXP orthologue

PbAP2-Sp[44,58]. PbAP2-Sp is a master regulator of sporogony and therefore has a

phenotype that can be correlated to the antimalarial activity of our lead compounds. We

hypothesized that if Compounds B and C can compete DNA binding by the AP2 domain

of PbAP2-Sp (89% identity (49/55 amino acids) to the AP2-EXP AP2 domain[55]), they

should affect the development of sporozoites in the mosquito stage of

Plasmodium berghei.

We infected *Anopheles stephensi* mosquitoes with *P. berghei* parasites and allowed the

parasites to develop for 10 days to the oocyst stage. At this time, mosquito midguts

were injected with the IC_{50} measured for asexual blood stage *P. falciparum* of

Compounds B, C or F **(Fig 1B)**. Compound F **(Fig 2A, B)** was used as a non AP2

domain competing control. On day 14 post infection, surviving mosquitoes were

dissected and both oocysts and midgut sporozoites were counted **(Fig 6A)**. Mosquitoes

injected with Compounds F and C developed comparable oocyst numbers to the DMSO

control **(Fig 6B)** while Compound B prevented the development of midgut oocysts

entirely **(Fig 6B)**. Compound C treated mosquitoes developed 10-fold fewer midgut

sporozoites than the control **(Fig 6C)**. Therefore, consistent with their hypothesized *in*

vitro activity against ApiAP2 proteins, Compounds B and C are strong inhibitors of

P. berghei mosquito stage development into sporozoites *in vivo.*

Discussion

ApiAP2 transcription factors are unique to Apicomplexan parasites due to their plant-like domain architecture[20,22] and many are essential to asexual blood stage development, making them valuable as potential drug targets. It is therefore desirable to discover chemical scaffolds that can target ApiAP2 proteins and to uncover details about their biological function. Additionally, phenotypic characterization has been especially challenging for essential asexual blood stage ApiAP2 proteins as no complete inducible knockdown has been described to date. Here, we used *in silico*, biochemical, and genetic approaches to demonstrate that ApiAP2 proteins can serve as drug targets against *Plasmodium* parasites. We have identified a starting point for chemical-based competition of ApiAP2 proteins and demonstrated a specific effect on parasite development and transcription by Compound C, a benzoxazole moiety compound which competes DNA binding by AP2-EXP, AP2-I D3, and PfSIP2 D1 *in vitro*.

Our results demonstrate the utility of *in-silico* screening to rapidly search a large and diverse set of compounds to target a specific protein domain. This approach allowed us to identify nine potential ApiAP2 competitors from a library of over 10,000 compounds. Compounds B and C both lack off-target activity against AtERF1 and SOX2 DNA binding, indicating selective activity against the AP2 domains of ApiAP2 proteins. Although the mode of DNA binding is shared between AP2-EXP and AtERF1[48], the specific amino acids that mediate DNA binding are not strictly conserved[43]. Since the predicted mode of docking for Compounds A and B would interfere with the AP2-EXP dimerization pocket in addition to DNA binding residues **(Fig S1)**, disruption of

protein:protein interaction may be their mode of action. As AtERF1 is not known to

homodimerize, disruption of protein interaction would not impact its DNA binding.

Compound I has the greatest magnitude of DNA major groove intercalation ability and

does not compete DNA binding by SOX2, consistent with the observation that HMGB

family proteins bind DNA *via* the minor groove[64].

We find that the DNA binding competition capacity of Compound B analogs varies *in*

vitro based on substitutions to the benzene ring, thereby illuminating a possible

determinant of ApiAP2 DNA binding competition. In a molecular dynamics simulation,

Compound B remains stably associated with AP2-EXP, while the non ApiAP2

competitor Compound B-3 does not. This suggests that a steric clash between AP2-

EXP and the bromine atom on Compound B-3 is responsible for its lack of DNA binding

competition. By extension, this may explain the lower efficacy of DNA binding

competition observed for Compounds B-1, B-2, and B-3 compared to B and B-4.

ApiAP2 competitor Compounds B and C both cause asexual *P. falciparum* parasites to

stop morphologically progressing in the trophozoite stage. Importantly, we observed that

more than 50% of parasites successfully reinvade in the presence of $0.66xIC_{50}$ (12μM)

Compound C, which we used for our RNA time course. Therefore, the changes in

transcript abundance measured after exposing parasites to Compound C are likely

specific to the time points maximally affected, rather than a cumulative effect of

completely halting parasite progression in the IDC. Compound C induced growth

disruption results in a decrease in the abundance of many transcripts that are AP2-EXP

targets. Affected transcripts are enriched for pathways important for parasites to invade

new RBCs and modify the RBC surface.

Our ChIP-seq datasets demonstrate that AP2-EXP preferentially binds the DNA

sequence CATGCA *in vivo*. This is the same core DNA motif that was previously shown

to be bound by AP2-EXP and PbAP2-Sp *in vitro*[44,65]. Many transcripts that correspond

to AP2-EXP target genes are activated, and a small subset are downregulated, at 25-35

hpi in the IDC when AP2-EXP is maximally expressed. Transcripts encoding proteins

important for parasite RBC modification and invasion (EBA-181, GAP45, MSP6, EBA-

140), and genome integrity (H2A, H2B, H4, DNA Mismatch Repair Protein) are

represented among those upregulated and targeted by AP2-EXP. Transcripts

downregulated following AP2-EXP occupancy primarily code for proteins of unknown

function, or proteins that are expressed outside of the asexual blood stage (*e.g.*,

PHISTb, P113). AP2-EXP occupancy is well correlated with the activating chromatin

reader PfBDP1[15]. This coincides with results reported for PfAP2-I, which binds many of

the same genomic loci as AP2-EXP and BDP1[11,29,32]. Together, these results point to

transcription activation *via* chromatin modulation as a key role of AP2-EXP in the

asexual blood stage. Future studies are needed to understand the precise mechanisms

by which AP2-EXP and other ApiAP2 proteins regulate transcription.

New inferences can be made about the function of AP2-EXP based on the description

of its genome-wide DNA binding sites and transcript dysregulation in the presence of

Compound C. This demonstrates the utility of combining chemical and traditional

genetics because previous attempts to genetically characterize AP2-EXP have not

provided a complete picture of its function in the asexual blood stage. The *ap2-exp* AP2

domain coding region has been shown to be essential for the asexual blood stage by

both saturating mutagenesis and targeted deletion attempts[26,27,29,33]. Conversely, the

coding region beyond the AP2 domain was truncated in two studies[26,33]. Transcriptomic

profiling of a truncated AP2-EXP parasite line revealed that many transcripts encoding

exported proteins were dysregulated[33]. Unexpectedly, AP2-EXP target genes identified

in our study overlap poorly with the differentially regulated transcripts reported as a

result of truncation of the AP2-EXP protein **(Table S5)**[33]. There is greater overlap

between our study and the AP2-EXP ChIP-seq in trophozoites recently described by

Shang *et al*[29], with 50/101 target genes overlapping **(Table S5)**. It is therefore possible

that the truncated AP2-EXP retains some critical function that relies on DNA binding.

This would explain the apparent essentiality of the AP2 domain, while the full-length

protein has other roles beyond DNA binding that impact a different subset of non-

asexual blood stage essential genes.

Since Compounds B and C each compete AP2 domain proteins in addition to AP2-EXP

in vitro, they could potentially be used to assay the function of blood stage essential

ApiAP2 proteins in addition to AP2-EXP. A comparison of transcripts downregulated in

the presence of Compound C to the DNA binding occupancy of AP2-I[32], for which

knockdown has not been reported, revealed that 54/85 AP2-I target genes are

decreased in abundance by $\log_2$ fold change >2 at 24-30 hpi. Interestingly, 22/55 AP2-I

bound genes that decrease in abundance when parasites are exposed to Compound C

are also AP2-EXP targets, implying the potential for co-regulation of certain gene

subsets. This overlap is consistent with our EMSA results and corroborates the

hypothesis that Compound C competes both AP2-EXP and AP2-I *in vivo*. Previously,

PfSIP2 knockdown has been shown to impact a small subset of genes[35]. Only 3/22 of

these are decreased in abundance in the presence of Compound C **(ApiAP2 Target**

Gene Results Summarized in Table S4). This may reflect the cryptic relationship

between PfSIP2 DNA binding and transcriptional control, as was noted when PfSIP2

was originally characterized as playing a role in genome integrity and heterochromatin

formation[35].

Several ApiAP2 proteins have been implicated in oocyst and sporozoite development in

a *P. berghei* ApiAP2 knockout screen[30]. Since Compounds B and C both inhibit

mosquito stage development of *P. berghei* with different phenotypes, it is possible that

they target different ApiAP2 proteins *in vivo*. The AP2-EXP orthologue PbAP2-Sp acts

as the master regulator of infectious sporozoite formation by binding upstream of many

genes essential for sporogony[44,58]. Therefore, Compounds B and C should minimally

inhibit sporozoite development, which is consistent with our results. Overall, these data

support a potential for multi-AP2 domain competition by Compounds B and C *in vivo*.

The IC_{50} values against *P. falciparum* for the ApiAP2 domain competitors identified in

this study are all in the micromolar range. Therefore, in order for these compounds to be

effective antimalarial drugs, future work will be needed to increase parasite killing

potency. Encouragingly, Compounds A, B, and C all have the same core benzoxazole

moiety and compete DNA binding by ApiAP2 proteins, but vary widely in their measured

$IC_{50}s$. This indicates that changes which do not affect ApiAP2 DNA binding competition

can change the potency against parasites. Encouragingly, our present results indicate

that Compounds B and C can inhibit *P. berghei* parasite development without killing

mosquitoes, suggesting that ApiAP2 competitor compounds may be tolerated by the

host and could serve as transmission blocking agents[66]. Due to their inclusion in the

Tres Cantos Antimalarial Set, Compounds B, B-1, B-2, B-3, and B-4 have all been

tested for activity against the human HEPG2 cell line[46]. ApiAP2 competitor Compounds

B and B-4 inhibit HEPG2 cell growth by just 4 and 8%, respectively[46]. Therefore, either

drug may be prioritized for further development based on their selectivity for

Plasmodium parasites.

AP2-EXP was originally characterized at the genetic level over a decade ago, and it is

the only ApiAP2 protein for which the structure has been solved. Surprisingly, the field

still lacks a clear understanding of the biological role of AP2-EXP in the asexual blood

stage. Our study has enabled us to make inferences about the role of AP2-EXP and set

a proof of principle for targeting the highly unique Apicomplexan AP2 DNA binding

proteins as a new antimalarial strategy.

<u>Materials and Methods</u>

Parasite lines

Parasites were grown at 37°C, 5% O2, 7% CO2 using RPMI1640 media supplemented

with hypoxanthine and .5% Albumax II (Thermo). The parasite lines used in this study

were AP2-EXP::GFP, AP2-EXP::HA, AP2-EXP::*glms*, and wild Type Pf3D7 (Malaria

Research and Reference Reagents Repository).

To create AP2-EXP::GFP the C-Terminal region of AP2-EXP was cloned **(Table S2)**

and inserted into the plasmid pSLI::2xFKBP[51] for endogenous tagging by single

homologous recombination. Red blood cells were pre-loaded with 100μg of plasmid as

previously described[67] and Pf3D7 wild type parasites were added at 1% parasitemia.

After 24 hours, cultures were maintained in media with 2.5nM WR99210 until parasites

were detectable by Giemsa stained slides, at which time they were switched to G418

selection at 400μg/mL as described[51]. Parasites were cloned by limiting dilution and

screened for correct genomic integration **(Fig S10A)**. Transgenic parasites with the

correct C-Terminal tag were further transfected with pLyn-FRB-mCherry[51] for inducible

mislocalization by preloading red blood cells with 100μg plasmid and selection using

1μg/mL blasticidin to create AP2-EXP::GFP + pLyn. All components were confirmed by

genotyping PCR. From this point onward, parasites were maintained in media with

2.5nM WR99210. All ChIP and Western blot experiments using AP2-EXP::GFP except

Figure S22 were collected using AP2-EXP::GFP without the pLyn mislocalization

plasmid.

To create AP2-EXP::HA we used the pSN054 vector system[68]. The right homology

region (RHR) was amplified by PCR **(Table S2)**, and the recodonized left homology

region (LHR) was synthesized using the BioXP™ 3200 System. Single guide RNA

fragments synthesized using the BioXP™ 3200 System (**Table S2)** were cloned into the

linearized pSN054 donor vector[68] that incorporates a C-terminal V5 and 2x-

hemagglutinin (HA) tags, 10x RNA aptamer array, the TeTR-DOZI expression cassette

containing the blasticidin S-deaminase gene for selection in the parasites, and the

reporter gene Renilla luciferase (RLuc). The 3'-end was re-codonized to avoid further

modifications of the edited locus in the parasite via removal of the PAM site. Donor

vector generation was carried out via Gibson assembly, and the final constructs were

sequence verified and further confirmed by restriction digests. Transfections into Cas9-

and T7 RNA polymerase-expressing NF54 parasites[69] were carried out by preloading

erythrocytes with linearized vectors as described[70]. Cell cultures were maintained in 500

nM anhydrotetracycline (aTc, Sigma-Aldrich 37919), drug selection with 2.5 mg/mL of

Blasticidin S was initiated four days after transfection, and recovered parasites were

monitored via Giemsa smears and RLuc measurements. Parasites were then cloned by

limiting dilution and PCR genotyped **(Fig S10B)**.

To create AP2-EXP::*glms*::HA, the C-terminal coding region of AP2-EXP was amplified

(Table S2) and cloned into the plasmid pSLI*::*3xHA::*glms*[71] to create pSLI:AP2-

EXP*::glms*::HA*.* Red blood cells were pre-loaded with 100µg of plasmid as described[67]

and 3D7 wild type parasites were added in at 1% parasitemia. Cultures were

maintained on 2.5nM WR99210 until parasites were detectable by Giemsa stained

smear, then switched to 400µg/mL G418 as described[51]. Surviving parasites were

cloned by limiting dilution, then genotyped **(Figure S21)**. From this point on AP2-

EXP::*glms*::HA parasites were maintained in media with 2.5nM WR99210. The human

biological samples were sourced ethically and their research use was in accord with the

terms of the informed consents under an IRB/EC approved protocol.

Genomic DNA Isolation

Parasite cultures were lysed with .1% Saponin in 1xPBS and collected by centrifugation at 1500 RPM, then resuspended in 1xPBS. Genomic DNA was then isolated using the Qiagen DNeasy nucleic acid isolation kit according to the manufacturer's instructions.

AP2-EXP Knockdown Assays

For AP2-EXP::GFP + pLyn, synchronous parasites were split into two populations. 250nM Rapalog was added to one parasite group for 48 hours as described[51] while the second group was used as a control. Addition of Rapalog did not affect parasite growth (not shown). After 48 hours parasite protein was collected in nuclear and cytosolic fractions and AP2-EXP::GFP, Histone H3, and Pf Aldolase were detected by Western blot.

For AP2-EXP::HA parasites were grown routinely in the presence of 500nM Anhydrotetracycline (Atc). Synchronous parasites were split into two groups. Atc was washed out of one group, while the second group was maintained with Atc as a control. Parasites were maintained for 120 hours before protein harvest. Parasite growth was not affected by removal of Atc (not shown). AP2-EXP::HA and Histone H3 were detected by western blot.

For AP2-EXP::*glms*::HA synchronous parasites were split into two populations. 5mM glucosamine was added to one parasite group for 72 hours as described[63] while the second group was used as a control. Addition of glucosamine did not affect parasite

growth (not shown). After 72 hours parasite protein was harvested for detection of AP2-

EXP::*glms*::HA and Histone H3 by western blot.

In silico docking screen

The Tres Cantos Antimalarial Set of 13533 small molecules was retrieved from the

supplementary material of Gamo *et al*[46] and downloaded in Spatial Data File (SDF)

format from BatchEntrez. 4603 small molecule structures were downloaded from

DrugBank[47] version 2.5 in SDF format. Using the SDF library, a 3-dimensional MOL2

library was created with the use of BALLOON and the Merck Molecular Force Field.

BALLOON was used to calculate up to 20 conformations using a genetic algorithm and

to select the conformation with the lowest energy of conformation. The 3-dimensional

MOL2 library was then prepared for AutoDock using the python script

"ligand_prepare4.py" from AutoDockTools[72]. The prepared ligands were then saved in

PDBQT file format. The python script "ligand_prepare4.py" automatically defines the

torsions of the molecule.

The homodimerized AP2-EXP AP2 domain was downloaded from the solved crystal

structure (PDB ID 3IGM). AutoDockTools was used to remove water molecules, add

hydrogens, and calculate charges. Residues Arg-88 and Asn-118 were found to be

flexible using FlexPred and set as flexible residues for docking. The flexible residues

were saved as flexible residues in a flexible residue file and the remaining molecule was

saved in as rigid residues in a rigid residue file as specified in the AutoDock manual[72].

The rigid residues file was used to pre-calculate an energy grid of the macromolecule

using autogrid4[72]. A set of 14 grid map types S, Cl, F,A, Br, N, P, OA, SA, C, I, HD, Ca,

NA were calculated for the macromolecule using AutoGrid[72].

From the prepared ligands a docking set was prepared for AutoDock with the

macromolecule and pre-calculated energy grids. For each docking set a docking

parameter file was prepared, with the Lamarckian Genetic Algorithm (LGA) specified as

the search algorithm for AutoDock. To allow for high throughput parallel processing

PERL was used on top of AutoDock to manage the dockings. The algorithm was initially

set to run with a maximum of 25000 energy evaluations and 20 repeats. Since the

calculations are computationally intensive, this setting was used as initial screening of

the ligands, to identify a subset of ligands for a more thorough evaluation[72]. The 1000

best hits from the GSK compound evaluation and the DrugBank compound evaluation

were selected for a more thorough evaluation. The docking parameters were

reconfigured to 250000 energy evaluations and 100 repeats and docking repeated. The

docking results were examined with PERL, to provide an automated approach to the

interpretation of the docking results. The docking results were evaluated using two

different approaches. The top candidates for competition of DNA were selected using a

PERL script to filter based on having a geometric center within 10 Å of the location of

sense-strand DNA binding. Hits were then filtered based on having a predicted free

energy of interaction < -5 kJ/Mol.

IC$_{50}$ Determination

The IC_{50} of each putative ApiAP2 competing compound was determined using a 48-

hour growth inhibition assay adapted from[73]. Parasites were seeded at .5% parasitemia

and 4% hematocrit on a 96 well plate. Drug or DMSO vehicle control was added in

triplicate to each well and parasites were incubated in standard culture conditions for 48

hours. Parasite cultures were then frozen overnight and resuspended in 1:1 volume of

Sybr Green Lysis Buffer (20mM Tris-HCl, pH 7.5; 5mM EDTA; 0.08% Triton X-100;

0.008% saponin, 1xPBS). After 1 hour of incubation at room temperature, plates were

read at excitation/emission 485/535nm. Growth values were normalized to the vehicle

control. IC_{50} values were calculated and growth curves were plotted using GraphPad

Prism.

Recombinant Protein Expression

Recombinant AP2 domains AP2-EXP, AP2-I D3, AP2-HS D1, and PfSIP2 D1, were

overexpressed and purified from BL21 PlysS *E. coli* as described previously[25], using the

same plasmid constructs. Recombinant AtERF1 was purified using the same method

after cloning the AtERF1 AP2 domain[48] into the pGex4t-1 overexpression vector **(Table**

S2) Briefly, *E. coli* cultures grown to optical density .6 were induced to overexpress

recombinant protein by adding 40mM IPTG overnight at room temperature. Bacteria

pellets were collected by centrifugation and frozen overnight at –80°C. The pellets were

then lysed by sonication in sonication buffer (25mM Tris-Cl pH8.0, 150mM NaCl). GST

tagged fusion proteins were purified overnight by incubation with GST affinity beads

(Clontech) according to the manufacturer's protocol. Recombinant protein was

recovered by elution with 10mM reduced glutathione and quantified by Braford reagent

(Pierce). Input, flowthrough, and eluate fractions were analyzed by SDS-PAGE to

ensure recovery of the full- length recombinant protein. Recombinant full-length SOX2

protein was purchased from Abcam (ab169843).

Gel-Shifts

Gel-shifts were done using the Thermo Light Shift EMSA kit with recommended buffer

components (10mM Tris pH 7.5, 50mM KCl, 1mM DTT, 6mM MgCl2, 60ng/µL Poly

DiDC, 65ng BSA). Recombinant proteins were titrated to empirically determine the

minimum mass required for DNA binding and this mass **(Fig S3)** was used for each gel-

shift unless otherwise specified. PAGE purified DNA probes with a 5' biotin ligated on

the forward DNA strand, along with an unlabeled complementary stand, were

purchased from IDT **(Table S2)**. DNA probes were double stranded by heating to 95°C,

followed by stepwise cooling in annealing buffer (10mMTtris-Cl pH7.5, 1mM EDTA,

10mM NaCl). Each recombinant protein was incubated in DNA binding mixture plus

competitor compound for 15 minutes prior to addition of the cognate double stranded

DNA oligonucleotide. Protein, competitor, and DNA oligonucleotide were incubated

together for an additional 5 minutes. The mixtures were separated on a .5x TBE

polyacrylamide gel, transferred to a nylon membrane (Amersham) at 50 Volts for 30

minutes, and probed using the Light Shift nucleic acid detection module

(Thermo) according to the manufacturer's protocol. All gels were imaged using a Bio

Rad chemiluminescence imager.

Ethidium Bromide Exclusion Assay

10µM of double stranded DNA (sequence TGCATGCA, purchased from IDT) in .01mM

EDTA, 9.4mM NaCl, 2mM HEPES buffer pH 7.9 was incubated for 10 minutes in the

presence of 100nM ethidium bromide. DNA with no ethidium bromide was used in

parallel as a background control. Fluorescence was measured at

excitation/emission 546nm/595nm for each well. Following the baseline reading, each

putative AP2 competing compound was added. DRAQ5 DNA dye was used as a

positive control for knockdown of ethidium bromide fluorescence by DNA intercalation.

Ethidium bromide exclusion assays were performed in technical triplicate using a 96

well plate.

Molecular Dynamics Simulations

Five compounds were prepared for molecular dynamics (MD) simulation: B, B-1, B-2, B-

3, and B-4. In preparation for MD, parameters for the compounds were obtained using

Antechamber[74–76] with the Generalized Amber Force Field. Starting conformations for

each compound bound to AP2-EXP were based on predictions from docking. All

complexes were prepared using the tleap module of AmberTools[77] with the

protein.ff14SB forcefield[78]. Each complex was solvated in an octahedral box of TIP3P

water with a 10 -Å buffer around the protein complex. Na^+ and Cl^- ions were added to

neutralize the protein and achieve physiological conditions. All MD minimizations and

simulations were performed using Amber with GPU acceleration[79,80]. First, complexes

minimized with 5000 steps each of steepest decent and conjugate gradient minimization

with 500 kcal/mol·Å^2 restraints on all complex atoms. Restraints were reduced to 100

kcal/mol·Å^2 and the minimization protocol was repeated. Restraints were then retained

only on the compound for a final minimization step. Following minimization, all

complexes were heated from 0 to 300 K using a 100-ps run with constant volume

periodic boundaries and 10 kcal/mol·$Å^2$ restraints on all protein and compound atoms.

To equilibrate complexes, 10 ns of MD was performed first with 10 kcal/mol·$Å^2$, then

with 1 kcal/mol·$Å^2$ restraints on protein and compound atoms using the NPT ensemble.

With kcal/mol·$Å^2$ restraints retained on complexes, 500 ns production simulations were

performed. A 2-fs timestep was used and all bonds between heavy atoms and

hydrogens were fixed with the SHAKE algorithm[81]. A cut-off distance of 10 Å was used

to evaluate long-range electrostatics with Particle Mesh Ewald (PME) and for van der

Waals forces. The 'strip' and 'trajout' commands of the CPPTRAJ module[82] were used

to remove solvent atoms and extract 50,000 evenly spaced frames from each simulation

for analysis.

AP2 Domain Competitor Phenotyping Assays

For Compound B, C, and H spike in phenotyping **(Fig 4B)**, highly synchronous 3D7 wild

type *P. falciparum* parasites were spiked with 40µM of each drug at 24 hpi. An

equivalent volume of DMSO was used as a control. Parasites were morphologically

assessed by Giemsa staining at 24 and 48 hours post spike in (48 and 72 hpi).

For continuous Compound C exposure phenotyping **(Fig S9)**, highly synchronous 3D7

wild type *P. falciparum* parasites were spiked with 40µM Compound C or equivalent

DMSO vehicle control at 6 hpi. Parasite progression was then monitored through the

IDC by Giemsa staining at 6-hour intervals.

For Compound C spike in and washout assays **(Fig 4C, D)**, 40µM of Compound C was

spiked into highly synchronous 3D7 wild type *P. falciparum* parasites at 8-hour intervals

starting at 14 hpi. For Compound C washout assays, 40µM of Compound C was

washed out of highly synchronous cultures at 8-hour intervals starting at 14 hpi. For

each assay an equivalent volume of DMSO vehicle control was used as a control for

normal growth.

40µM of compound was used for all phenotyping assays because it was the maximum

concentration achievable in culture after resuspending each compound in 100% DMSO.

Higher concentrations of compound caused parasite mortality due to the high volume of

DMSO required (>1% volume/volume). All assays were performed in triplicate and

parasitemia was counted by Giemsa-stained slides. Error bars represent the standard

deviation of the mean.

Preparation of RNA for DNA Microarray

DNA microarrays were prepared using the protocol described in[52]. Briefly, RNA was

collected from highly synchronous 3D7 wild type *P. falciparum* parasites using TRIzol

Reagent in accordance with manufacturer's protocols. After collecting a control timepoint

at 6 hpi, parasites were cultured with either 12µM Compound C ($0.66 \times IC_{50}$) or DMSO

vehicle control. Parasite RNA was then harvested at 12, 18, 24, 30, 42, and 48

hpi. Purified RNA was reverse transcribed using SuperScript II according to the

manufacturer's protocol, with dUTP-aminoallyl and a mixture of nonamer and oligo-dT

primers added. cDNA was concentrated using DNA Clean and Concentrator-5

kits. cDNA was labeled using Amersham CyDye Reactive Dyes and re-purified. cDNA for

each sample was labeled using Cy5, while Cy3 was used for the reference pool made

up of mixed stage asexual cDNA. cDNA was hybridized onto Agilent DNA microarrays[52],

washed, and scanned using an Axon 4200A Scanner. Agilent Feature Extraction

Software version 11.0.1.1 with the protocol GE2-v5_95_Feb07_nospikein was used to

extract signal intensities. Raw signal intensities for all microarrays and the .gal file

3D7v7.1+apimito used for feature extraction are available in **Table S7**.

Analysis of DNA Microarrays

Microarray data were extracted and normalized using the R package LIMMA[54]. Arrays

were normalized using robust splines normalization and within arrays parameters

selected. Average expression values were calculated per gene by averaging the $\log_2$

Cy5 (cDNA)/Cy3 (Reference Pool) signal intensity across all probes. Correlation plots

for the Compound C vs. DMSO control total transcriptome and control genes were

made using the $\log_2$ signal intensities for all detected genes using the R package

CorrPlot downloaded from: https://github.com/taiyun/corrplot. The LIMMA eBayes

function was used on average gene abundance data to determine changes in transcript

abundance between 24 and 30hpi that occur differentially in DMSO control vs.

Compound C dosed parasites. Volcano plots were made with LIMMA eBayes data

using the Enhanced Volcano package in R downloaded from:

https://github.com/kevinblighe/EnhancedVolcano. The RNITS[56] R package was used to

model differences in gene expression across the entire time course between the DMSO

control and Compound C spike in parasites. RNITS was run using the parameters:

center genes, normalize by intensity, and background filter probes. The RNITS model

for differential expression was fit at the gene level. All data from LIMMA, eBayes, and

RNITS is provided in **Table S4**. Heat maps for control genes and AP2-EXP target genes

were made using Java Treeview[83] after means centering and log transformation of

transcript abundance values with Cluster3.0[84].

ChIP Sample Preparation

ChIP-seq was performed as described in[38]. Highly synchronous parasites were grown

up to 5-10% parasitemia and AP2-EXP ChIP was performed at 30 hpi for all replicates.

Parasites were first chemically crosslinked by addition of 1% formaldehyde for 10

minutes at 37°C, followed by 5 minutes of quenching in 125mM Glycine on ice. The

supernatant was removed by centrifugation and parasites were resuspended in 1xPBS

with .1% saponin for 5 minutes at 4°C to lyse red blood cells. Following lysis, parasites

were washed in cold 1xPBS and lysed in Lysis Buffer (10 mM HEPES pH 7.9, 10 mM

KCl, 0.1 mM EDTA pH 8.0, 0.1 mM EGTA pH 8.0, 1 mM DTT, 1x Pierce Protease

Inhibitor) on ice for 30 minutes. NP-40 was added to a final concentration of .25% and

parasites were homogenized in a douncer homogenizer using 100 strokes per 2X10^9

parasites. Nuclei were separated from the supernatant by centrifugation at 14000 RPM

for 10 minutes.

Nuclei were then sonicated in Shearing Buffer (0.1% SDS, 10 mM Tris pH 8.1, 1 mM

EDTA, 1X Pierce Protease Inhibitor) using an M220 Covaris Sonicator with the following

settings: duty cycle 5%, peak incident power 75 W, cycles per burst 200, temperature

7°C, time 300 seconds. Sonicated material was diluted 1:5 in Dilution Buffer, (0.01%

SDS, 1.1% Triton X-100, 1.2 mM EDTA, 16.7 mM Tris-HCl pH 8.1, 150 mM NaCl) pre-

cleared using Magna ChIP Protein A+G Magnetic Beads (Sigma) for 2 hours, then the

Input fraction was separated and set aside. Immunoprecipitation was performed

overnight with .5mg/mL 3F10 anti-HA (Sigma) antibody or .1mg/mL Ab290 (Abcam)

anti-GFP and immune complexes were collected using Magna ChIP Protein A+G

Magnetic Beads. Immunoprecipitated material was washed five times total, once with

Low Salt at 4°C (0.1% SDS, 1% Triton X-100, 2 mM EDTA, 20 mM Tris-HCl pH 8.1, 150

mM NaCl), High Salt at 4°C (0.1% SDS, 1% Triton X-100, 2 mM EDTA, 20 mM Tris-HCl

pH 8.1, 500 mM NaCl), Lithium Chloride Buffer at 4°C (0.25 M LiCl, 1% NP-40, 1%

Deoxycolate, 1mM EDTA, 10 mM Tris-HCl pH 8.1), and twice at room temperature with

1xTE buffer. Immune complexes were then eluted from the beads using Elution Buffer

(1% SDS, .1M NaHCO3). Crosslinking was reversed by overnight incubation with .2M

NaCl at 65°C, then RNA and protein were removed by sequential treatments with

RNAseA and Proteinase K. DNA was isolated using a Minelute PCR Purification Column

(Qiagen) and DNA concentration was measured using a Qbit dsDNA HS assay kit (Life

Technologies).

DNA Library Preparation

ChIP-seq DNA libraries were prepared as described in[38] using the NEBNext II DNA

library kit (New England Biolabs) according to the manufacturer's instructions. Briefly,

DNA ends were blunted, and a single A-tail overhang was added. Next, barcoded DNA

adaptors for Illumina sequencing (Bioo Scientific) were ligated to the sample and DNA

was amplified by 11 cycles of PCR using Kapa HiFi DNA Polymerase. DNA libraries

were size selected for 250nt fragments and library quality was assessed using an

Agilent 2100 Bioanalyzer or TapeStation. Libraries were quantified by Qbit dsDNA HS

assay kit and sequenced using a HiSeq 2500 (Replicate GFP1) or NextSeq 550

(Replicates GFP2 and HA3) Illumina sequencer. AMPure XP beads (Beckmann

Coulter) were used to size select and purify DNA between NEBNext II library

preparation steps.

Whole genome sequencing DNA libraries were prepared using the Illumina Tru-Seq

PCR free DNA library kit and sequenced using a HiSeq 2500.

qPCR

ChIP-qPCR samples were collected from 30-35 hpi trophozoites following two hours of

Compound C (40µM) or DMSO vehicle control spike in. Primer pairs to be used for

ChIP-qPCR were first evaluated to check for 80-110% efficiency using sonicated

genomic DNA. RT-qPCR was carried out using Sybr Green Polymerase master mix

(Thermo) with the specified primer concentration **(Table S2)**. The Ct was calculated

using SDSv1.4 (Applied Biosystems) software, averaged over technical triplicate. The

percent of Input per immunoprecipitated DNA fraction was calculated using the delta Ct

method. Each assay was performed in biological triplicate, with the exception of the

ama1 primer pair, where n =2. Error bars represent standard deviation. Data was

obtained using an Applied Biosystems 7300 Real-Time PCR Machine.

ChIP-seq data analysis

First, reads in each DNA library were trimmed to remove low quality base calls and

Illumina adaptor sequences using Trimmomatic[85] with a quality score cutoff of 20.

FastQC[86] was used to assess the quality of each DNA library following this step. Reads

were then mapped to the *Plasmodium falciparum* genome version 36 downloaded from

PlasmoDB using BWA-Mem[87]. Multiply mapped reads were filtered out using

Samtools[88]. Filtered and mapped Input and immunoprecipitate bam files were used for

peak calling by MACS2[57] with the following parameters: effective genome size

20000000, No Model, q 0.01. Peaks called in each ChIP-seq replicate were overlapped

using Bedtools[89] Intersect, and peaks that occur in a minimum of 2/3 replicates were

used for downstream analysis. For data visualization using IGV, .bam files were

converted to $\log_2$ Immunoprecipitate fraction/Input fraction bigwigs using DeepTools[90]

BamCompare. Bedtools[89] ClosestBed was used to correlate regions called as peaks of

occupancy by MACS2 to the closest gene. Results were filtered based on 1.5kB

proximity to the MACS2 peak and strandedness of the putative target gene. For

comparison of ChIP-seq library coverages to Transcription Start Sites and Start

Codons, DeepTools PlotHeatmap was used. Matrix files underlying PlotHeatmap were

created using DeepTools ComputeMatrix. A four color plot of conserved AP2-EXP DNA

binding sites was made using the program cegr-tools four color plot downloaded from

https://github.com/seqcode/cegr-tools/tree/master/src/org/seqcode/cegrtools. DNA

motifs enriched in AP2-EXP ChIP-seq peaks were determined using the DREME

Suite[91]. DNA motifs were sorted from highest to lowest level of conservation to the

aggregate AP2-EXP DNA motif using FIMO[92].

For comparison of AP2-EXP peaks with existing post-translational histone marks,

chromatin post translational modification datasets[13,17] were downloaded from

https://github.com/Daread/plasmodiumExprPrediction[14]. Chromatin reader and

nucleosome occupancy datasets[61,93,94] were downloaded from NCBI. ChIP-seq data for

BDP1 and HP1 was processed as described above. Nucleosome occupancy data was

downloaded from NCBI and processed as described above, with the exception that bam

files were turned into bigwig using BamtoBigWig with the --mnase parameter selected.

DeepTools ComputeMatrix was used to format the data. DeepTools PlotHeatmap was

used to plot coverage of each dataset within 10kB of AP2-EXP peaks. For negative

control coverage plots, Bedtools[89] ShuffleBed was used to create random genomic

intervals on the same chromosome and the same length as AP2-EXP peaks of

occupancy.

For comparison of AP2-EXP peaks identified by Shang *et al*[29] to this study, AP2-EXP

peaks of occupancy in the trophozoite stage were downloaded from

https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE184658 . Peak summits from

each replicate were expanded to 200 nucleotide windows using Bedtools SlopBed[89].

Conserved peaks in 2/2 replicates were identified using Bedtools intersect[89], then AP2-

EXP peaks from this study and Shang *et al* were overlapped using Bedtools Intersect[89].

Western blot

For full parasite lysate Western blots, parasite protein was collected by lysing RBCs

with .1% saponin on ice. After lysis, parasite pellets were washed by centrifugation with

cold 1xPBS. Parasite pellets were resuspended in 50µL Loading Buffer (50mM Tris-Cl

pH 8.0, 20% SDS, 1% Bromophenol Blue) and boiled at 95°C for 10 minutes. For

fractionated parasite lysate, red blood cells were saponin lysed and parasite material

was washed as described. Fractionated parasite protein was prepared as described

in[95]. Briefly, parasites were resuspended in Lysis Buffer (20mM HEPES pH 7.8, 10mM

KCl, 1mM EDTA, 1mM DTT, 1x Pierce Protease Inhibitor) and incubated for 15 minutes

at 4°C. NP-40 was added to a final concentration 0.65% and nuclear material was

pelleted by centrifugation at 2500xg at 4°C for 10 minutes. The cytosolic protein was

collected and boiled in 1:1 volume of Loading Buffer at 95°C for 10 minutes. Pelleted

nuclei were then resuspended in 1mL Extraction Buffer (20mM HEPES pH 7.8, 800mM

KCl, 1mM EDTA, 1mM DTT, 1x protease inhibitor) per 1x10^9 cells for 1 hour at 4°C.

Soluble nuclear protein was separated by centrifugation at 2500xg for 10 minutes at

4°C. Nuclear protein was then boiled in 1:1 volume of Loading Buffer at 95°C for 10

minutes.

Protein was run on a 6% SDS PAGE at 100V for 3 hours, then transferred overnight at

4°C and 10 Volts to a nitrocellulose membrane. Membranes were blocked using 5%

milk in 1xTBS-.1% tween for 1 hour. Primary antibodies used were: 1/1000 rat ant-HA

(Roche 3F10), 1/1000 mouse anti-GFP (Roche), 1/3000 rabbit anti-aldolase conjugated

to HRP (Abcam ab38905), or 1/3000 mouse anti-H3 (Abcam ab10799). Membranes

were washed three times with 1xTBS-.1% tween between primary and secondary

antibody incubation. Secondary antibody concentrations used were 1/3000 goat anti-rat

HRP conjugate (Millipore), 1/3000 goat anti-mouse HRP conjugate, or 1/10,000

(Pierce), goat anti-rabbit HRP conjugate (Millipore). ECL reagent (Pierce) was used to

detect HRP signal. Blots were exposed to autoradiography film (VWR) and imaged

using an autoradiography developer.

Protein Pulldown

Parasite nuclear protein was isolated as described above for fractionated Western blots.

Following the final centrifugation step, the supernatant was collected and diluted by 1:3

in Dilution Buffer (30% glycerol, 20mM HEPES pH 7.8). GFP tagged AP2-EXP was

pulled down using Chromotek anti-GFP or mock immunoprecipitated using Chromotek

negative control magnetic beads. Beads were washed twice in Wash Buffer prior to use

(20mM HEPES pH 7.4, 250mM NaCl, 1mM EDTA, 1mM TCEP, .05% NP-40). Protein

and beads were incubated together for 1 hour at 4°C. The beads were then washed

twice with Wash Buffer and bound protein was collected in Sample Buffer by boiling at

95°C for 10 minutes.

Fluorescent Microscopy

Samples were prepared by incubating packed infected red blood cells with DRAQ5 dye

(Thermo) for 15 minutes. Parasites were then washed in 1xPBS to remove excess dye

and immediately placed on a glass slide for imaging. Fluorescent microscopy images

were acquired using an Olympus Bx61 fluorescent microscope. All images were

processed using SlideBook 5.0.

Mosquito Stage *P. berghei* AP2 Competitor Growth Assay

A female Swiss Webster mouse was inoculated with Plasmodium berghei ANKA 2.34

from frozen stock. Once the parasitemia reached 15%, the blood was harvested by

heart puncture, washed twice with 1xPBS and resuspended to 10 mL in 1xPBS. 5

female Swiss Webster mice were infected with 500µL of the resuspended blood. 3-4

days after blood passage, exflagellation of the *P. berghei* gametocytes was assayed.

Briefly, a drop of tail vein blood was incubated in RPMI 1640 (Invitrogen) containing

1µM xanthurenic acid (Sigma) for ~12-15 minutes. The mixture was added to a slide

and observed under a light microscope at 40x. Ten or more fields were observed. Mice

with 0.3-0.7 exflagellations per field were anesthetized and fed on 3-day post

emergence *Anopheles stephensi* mosquitoes. Mosquitoes were maintained at 19°C and

fed 10% sucrose. Bloodfed mosquitoes were separated 30 hours post-bloodfeeding. 10

days post-bloodfeeding, 5-10 mosquitoes were dissected and oocysts were counted by

mercurochrome staining and light microscopy to ensure *P. berghei* oocyst development.

The remaining mosquitoes were injected by standard mouth pipette technique with

different small molecule inhibitors or a mixture of PBS-DMSO for control. The surviving

injected mosquitoes were dissected 14 days post-bloodfeeding. Oocyst numbers were

counted as before. Pooled groups of midguts were ground using a pestle, centrifuged at

7,000 rpm for 5 minutes and resuspended in 20µL 1xPBS to release developing mid-gut

sporozoites. Sporozoites were counted on a haemocytometer. All in vivo studies were

conducted in accordance with the GSK Policy on the Care, Welfare and Treatment of

954 Laboratory Animals and were reviewed by the ethical review process at the institution

955 where the work was performed.

Data Availability

958 All sequencing data for AP2-EXP ChIP-seq and whole genome sequencing for

959 transgenic parasite lines AP2-EXP::GFP and AP2-EXP::HA has been deposited in the

960 SRA under **PRJNA818769**.

962 Raw microarray data for the Compound C and DMSO control RNA time course is

963 available in **Table S7**.

Author Contributions

966 T.J.R., E.K.D.S., K.S.S., N.D., C.F.A.P., G.P., M.J.L., C.D.O., J.C.N. and M.L. designed

967 experiments. T.J.R., E.K.D.S., V.C., K.S.S., N.D., G.J., C.F.A.P., and I.K. performed the

968 experiments. T.J.R. and N.D. analyzed the data. T.J.R. generated figures. T.J.R. and

969 M.L. wrote the manuscript in collaboration with the other authors

Acknowledgements

972 This work was funded through NIH/NIAID R01AI076276 (M.L.), R01AI125565 (M.L.),

973 and with support from the Center for Quantitative Biology (P50 GM071508) (M.L.).

974 T.J.R. was supported by NIH T32 Predoctoral Training Grant (5T32GM125592-01)

975 awarded to the Center for Eukaryotic Gene Regulation (CEGR) at The Pennsylvania

976 State University. G.A.J. is a recipient of the Sir Keith Murdoch Fellowship from the

American Australian Association and a Postdoctoral Research Grant from the American

Heart Association (16POST26420067).

Conflict of Interest Statement

FJG is a GlaxoSmithKline employee and own shares of the company

REFERENCES

1. WHO. *Word Malaria Report 2021*. *Word Malaria report Geneva: World Health Organization. (2021). Licence: CC* (2021).
2. Ippolito, M. M., Moser, K. A., Kabuya, J.-B. B., Cunningham, C. & Juliano, J. J. Antimalarial Drug Resistance and Implications for the WHO Global Technical Strategy. *Curr. Epidemiol. Reports* **8**, 46–62 (2021).
3. Cowman, A. F., Healer, J., Marapana, D. & Marsh, K. Malaria: Biology and Disease. *Cell* **167**, 610–624 (2016).
4. Josling, G. a & Llinás, M. Sexual development in Plasmodium parasites: knowing when it's time to commit. *Nat. Rev. Microbiol.* **13**, 573–87 (2015).
5. Zanghì, G. *et al.* A Specific PfEMP1 Is Expressed in P. falciparum Sporozoites and Plays a Role in Hepatocyte Infection. *Cell Rep.* **22**, (2018).
6. Bozdech, Z. *et al.* The transcriptome of the intraerythrocytic developmental cycle of Plasmodium falciparum. *PLoS Biol.* **1**, 85–100 (2003).
7. Lu, X. M. *et al.* Nascent RNA sequencing reveals mechanisms of gene regulation in the human malaria parasite Plasmodium falciparum. *Nucleic Acids Res.* **45**, 7825–7840 (2017).
8. Painter, H. J., Carrasquilla, M. & Llinás, M. Capturing *in vivo* RNA transcriptional dynamics from the malaria parasite *Plasmodium falciparum*. *Genome Res.* gr.217356.116 (2017) doi:10.1101/gr.217356.116.
9. Le Roch, K. G. *et al.* Discovery of gene function by expression profiling of the malaria parasite life cycle. *Science (80-.).* **301**, 1503–1508 (2003).
10. Foth, B. J. *et al.* Quantitative time-course profiling of parasite and host cell proteins in the human malaria parasite Plasmodium falciparum. *Mol. Cell. Proteomics* **10**, M110.006411 (2011).
11. Hoeijmakers, W. A. M. *et al.* Epigenetic reader complexes of the human malaria parasite, Plasmodium falciparum. *Nucleic Acids Res.* **47**, 11574–11588 (2019).
12. Toenhake, C. G. *et al.* Chromatin Accessibility-Based Characterization of the Gene Regulatory Network Underlying Plasmodium falciparum Blood-Stage Development. *Cell Host Microbe* **23**, 557-569.e9 (2018).
13. Jiang, L. *et al.* PfSETvs methylation of histone H3K36 represses virulence genes in Plasmodium falciparum. *Nature* **499**, 223–227 (2013).
14. Read, D. F., Cook, K., Lu, Y. Y., Le Roch, K. G. & Noble, W. S. Predicting gene expression in the human malaria parasite Plasmodium falciparum using histone modification, nucleosome positioning, and 3D localization features. *PLoS Comput. Biol.* **15**, 1–23 (2019).
15. Josling, G. A. *et al.* A Plasmodium Falciparum Bromodomain Protein Regulates Invasion Gene

Expression. *Cell Host Microbe* **17**, 741–751 (2015).

16. Pérez-Toledo, K. *et al.* Plasmodium falciparum heterochromatin protein 1 binds to tri-methylated
histone 3 lysine 9 and is linked to mutually exclusive expression of var genes. *Nucleic Acids Res.*
37, 2596–2606 (2009).

17. Bártfai, R. *et al.* H2A.Z demarcates intergenic regions of the Plasmodium falciparum epigenome
that are dynamically marked by H3K9ac and H3K4me3. *PLoS Pathog.* **6**, (2010).

18. Ruvalcaba-Salazar, O. K. *et al.* Recombinant and native Plasmodium falciparum TATA-binding-
protein binds to a specific TATA box element in promoter regions. *Mol. Biochem. Parasitol.* **140**,
183–196 (2005).

19. Coulson, R., Hall, N. & Ouzounis, C. Comparative genomics of transcriptional control in the
human malaria parasite Plasmodium falciparum. *Genome Res.* 1–7 (2004)
doi:10.1101/gr.2218604.

20. Jeninga, M., Quinn, J. & Petter, M. ApiAP2 Transcription Factors in Apicomplexan Parasites.
Pathogens **8**, 47 (2019).

21. Painter, H. J., Campbell, T. L. & Llinás, M. The Apicomplexan AP2 family: Integral factors
regulating Plasmodium development. *Mol. Biochem. Parasitol.* **176**, 1–7 (2011).

22. Balaji, S., Madan Babu, M., Iyer, L. M. & Aravind, L. Discovery of the principal specific
transcription factors of Apicomplexa and their implication for the evolution of the AP2-integrase
DNA binding domains. *Nucleic Acids Res.* **33**, 3994–4006 (2005).

23. Aravind, L., Iyer, L. M., Wellems, T. E. & Miller, L. H. Plasmodium Biology: Genomic Gleanings. *Cell*
115, 771–785 (2003).

24. Hollin, T. & Le Roch, K. G. From Genes to Transcripts, a Tightly Regulated Journey in Plasmodium.
Front. Cell. Infect. Microbiol. **10**, 1–13 (2020).

25. Campbell, T. L., de Silva, E. K., Olszewski, K. L., Elemento, O. & Llinás, M. Identification and
Genome-Wide Prediction of DNA Binding Specificities for the ApiAP2 family of regulators from
the malaria parasite. *PLoS Pathog.* **6**, (2010).

26. Zhang, M. *et al.* Uncovering the essential genes of the human malaria parasite Plasmodium
falciparum by saturation mutagenesis. *Science (80-.).* **360**, (2018).

27. Shang, X. *et al.* A cascade of transcriptional repression determines sexual commitment and
development in Plasmodium falciparum. *Nucleic Acids Res.* **49**, 9264–9279 (2021).

28. Zhang, M. *et al.* Uncovering the essential genes of the human malaria parasite Plasmodium
falciparum by saturation mutagenesis. *Science (80-.).* **360**, (2018).

29. Shang, X. *et al.* Genome-wide landscape of ApiAP2 transcription factors reveals a
heterochromatin-associated regulatory network during Plasmodium falciparum blood-stage
development. 1–19 (2022).

30. Modrzynska, K. *et al.* A Knockout Screen of ApiAP2 Genes Reveals Networks of Interacting
Transcriptional Regulators Controlling the Plasmodium Life Cycle. *Cell Host Microbe* **21**, 11–22
(2017).

31. Zhang, C. *et al.* Systematic CRISPR-Cas9-Mediated ApiAP2 Genes Reveal Functional Insights into
Parasite Development. *MBio* **8**, 1–17 (2017).

32. Santos, J. M. *et al.* Red Blood Cell Invasion by the Malaria Parasite Is Coordinated by the PfAP2-I
Transcription Factor. *Cell Host Microbe* **21**, 731-741.e10 (2017).

33. Martins, R. M. *et al.* An ApiAP2 member regulates expression of clonally variant genes of the
human malaria parasite Plasmodium falciparum. *Sci. Rep.* **7**, 1–10 (2017).

34. Sierra-Miranda, M. *et al.* PfAP2Tel, harbouring a non-canonical DNA-binding AP2 domain, binds
to Plasmodium falciparum telomeres. *Cell. Microbiol.* (2017) doi:10.1111/cmi.12742.

35. Flueck, C. *et al.* A major role for the Plasmodium falciparum ApiAP2 protein PfSIP2 in
chromosome end biology. *PLoS Pathog.* **6**, (2010).

36.   Tintó-font, E., Michel-todó, L., Russell, T. J. & Casas-, N. The PfAP2-HS transcription factor
1068   protects malaria parasites from febrile temperatures. (2021).
1069   37.   Kafsack, B. F. C. *et al.* A transcriptional switch underlies commitment to sexual development in
1070   malaria parasites. *Nature* **507**, 248–52 (2014).
1071   38.   Josling, G. A. *et al.* Dissecting the role of PfAP2-G in malaria gametocytogenesis. *Nat. Commun.*
1072   **11**, 1–13 (2020).
1073   39.   Xu, Y. *et al.* PfAP2-G2 Is Associated to Production and Maturation of Gametocytes in Plasmodium
1074   falciparum via Regulating the Expression of PfMDV-1. *Front. Microbiol.* **11**, 1–11 (2021).
1075   40.   Singh, S. *et al.* The PfAP2-G2 transcription factor is a critical regulator of gametocyte maturation.
1076   *Mol. Microbiol.* **115**, 1005–1024 (2021).
1077   41.   Coetzee, N. *et al.* Epigenetic inhibitors target multiple stages of Plasmodium falciparum parasites.
1078   *Sci. Rep.* **10**, 1–11 (2020).
1079   42.   Vanheer, L. N., Zhang, H., Lin, G. & Kafsack, B. F. C. Activity of Epigenetic Inhibitors against
1080   Plasmodium. 1–6 (2020).
1081   43.   Lindner, S. E., De Silva, E. K., Keck, J. L. & Llinás, M. Structural Determinants of DNA Binding by a
1082   P. falciparum ApiAP2 Transcriptional Regulator. *J. Mol. Biol.* **395**, 558–567 (2010).
1083   44.   Yuda, M., Iwanaga, S., Shigenobu, S., Kato, T. & Kaneko, I. Transcription factor AP2-Sp and its
1084   target genes in malarial sporozoites. *Mol. Microbiol.* **75**, 854–863 (2010).
1085   45.   Morris, G. M. *et al.* Automated docking using a Lamarckian genetic algorithm and an empirical
1086   binding free energy function. *J. Comput. Chem.* **19**, 1639–1662 (1998).
1087   46.   Gamo, F. J. *et al.* Thousands of chemical starting points for antimalarial lead identification.
1088   *Nature* **465**, 305–310 (2010).
1089   47.   Wishart, D. S. *et al.* DrugBank 5.0: A major update to the DrugBank database for 2018. *Nucleic*
1090   *Acids Res.* **46**, D1074–D1082 (2018).
1091   48.   Allen, M. D., Yamasaki, K., Ohme-Takagi, M., Tateno, M. & Suzuki, M. A novel mode of DNA
1092   recognition by a β-sheet revealed by the solution structure of the GCC-box binding domain in
1093   complex with DNA. *EMBO J.* **17**, 5484–5496 (1998).
1094   49.   Tintó-font, E. *et al.* A comparison of the properties of Sox-3 with Sry and two related genes, Sox-1
1095   and Sox-2. *Plant Cell* **9**, 1–6 (2000).
1096   50.   Wang, Y., Schellenberg, H., Walhorn, V., Toensing, K. & Anselmetti, D. Binding Mechanism of
1097   Fluorescent Dyes to DNA Characterized by Magnetic Tweezers. *Mater. Today Proc.* **4**, S218–S225
1098   (2017).
1099   51.   Birnbaum, J. *et al.* A genetic system to study Plasmodium falciparum protein function. *Nat.*
1100   *Methods* **14**, 450–456 (2017).
1101   52.   Kafsack, B. F. C., Painter, H. J. & Llinás, M. New Agilent platform DNA microarrays for
1102   transcriptome analysis of Plasmodium falciparum and Plasmodium berghei for the malaria
1103   research community. *Malar. J.* **11**, 1–9 (2012).
1104   53.   Llinás, M., Bozdech, Z., Wong, E. D., Adai, A. T. & DeRisi, J. L. Comparative whole genome
1105   transcriptome analysis of three Plasmodium falciparum strains. *Nucleic Acids Res.* **34**, 1166–1173
1106   (2006).
1107   54.   Smyth, G. K., Ritchie, M. & Thorne, N. Linear Models for Microarray Data User ' s Guide.
1108   *Bioinformatics* **20**, 3705–3706 (2011).
1109   55.   Bahl, A. *et al.* PlasmoDB: The Plasmodium genome resource. A database integrating experimental
1110   and computational data. *Nucleic Acids Res.* **31**, 212–215 (2003).
1111   56.   Sangurdekar, D. P. The Rnits package for normalization and inference of differential expression in
1112   time series microarray data . *Rvignette* 1–13 (2014).
1113   57.   Gaspar, J. M. Improved peak-calling with MACS2. *bioRxiv* 1–16 (2018) doi:10.1101/496521.
1114   58.   Yuda, M., Kaneko, I., Murata, Y., Iwanaga, S. & Nishi, T. Targetome analysis of malaria sporozoite

1115   transcription factor AP2-Sp reveals its role as a master regulator. (2022).

1116 59. Kensche, P. R. *et al.* The nucleosome landscape of Plasmodium falciparum reveals chromatin
1117   architecture and dynamics of regulatory sequences. *Nucleic Acids Res.* **44**, 2110–2124 (2015).

1118 60. Chappell, L. *et al.* Refining the transcriptome of the human malaria parasite Plasmodium
1119   falciparum using amplification-free RNA-seq. *BMC Genomics* **21**, 1–19 (2020).

1120 61. Fraschka, S. A. *et al.* Comparative Heterochromatin Profiling Reveals Conserved and Unique
1121   Epigenome Signatures Linked to Adaptation and Development of Malaria Parasites. *Cell Host*
1122   *Microbe* **23**, 407-420.e8 (2018).

1123 62. Ganesan, S. M., Falla, A., Goldfless, S. J., Nasamu, A. S. & Niles, J. C. Synthetic RNA-protein
1124   modules integrated with native translation mechanisms to control gene expression in malaria
1125   parasites. *Nat. Commun.* **7**, 1–10 (2016).

1126 63. Prommana, P. *et al.* Inducible Knockdown of Plasmodium Gene Expression Using the glmS
1127   Ribozyme. *PLoS One* **8**, 1–10 (2013).

1128 64. Štros, M. HMGB proteins: Interactions with DNA and chromatin. *Biochim. Biophys. Acta - Gene*
1129   *Regul. Mech.* **1799**, 101–113 (2010).

1130 65. De Silva, E. K. *et al.* Specific DNA-binding by apicomplexan AP2 transcription factors. *Proc. Natl.*
1131   *Acad. Sci.* **105**, 8393–8 (2008).

1132 66. Paton, D. G. *et al.* Exposing Anopheles mosquitoes to antimalarials blocks Plasmodium parasite
1133   transmission. *Nature* **567**, 239–243 (2019).

1134 67. Wu, Y., Kirkman, L. A. & Wellems, T. E. Transformation of Plasmodium falciparum malaria
1135   parasites by homologous integration of plasmids that confer resistance to pyrimethamine. *Proc.*
1136   *Natl. Acad. Sci. U. S. A.* **93**, 1130–4 (1996).

1137 68. Nasamu, A. S. *et al.* An integrated platform for genome engineering and gene expression
1138   perturbation in Plasmodium falciparum. *Sci. Rep.* **11**, 1–15 (2021).

1139 69. Murithi, J. M. *et al.* The Plasmodium falciparum ABC transporter ABCI3 confers parasite strain-
1140   dependent pleiotropic antimalarial drug resistance. *Cell Chem. Biol.* 1–16 (2021)
1141   doi:10.1016/j.chembiol.2021.06.006.

1142 70. Deitsch, K. W., Driskill, C. L. & Wellems, T. E. Transformation of malaria parasites by the
1143   spontaneous uptake and expression of DNA from human erythrocytes. *Nucleic Acids Res.* **29**,
1144   850–853 (2001).

1145 71. Simon, C. S. *et al.* An extended DNA-free intranuclear compartment organizes centrosome
1146   microtubules in malaria parasites. *Life Sci. Alliance* **4**, 1–15 (2021).

1147 72. Huey, R. & Morris, G. M. Using AutoDock 4 with AutoDockTools : A Tutorial. 1–56.

1148 73. Abrahams, L. & Burke, M. w o T w o. 20–36 (2010).

1149 74. Wang, J., Wolf, R. M., Caldwell, J. W., Kollman, P. A. & Case, D. A. Development and testing of a
1150   general Amber force field. *J. Comput. Chem.* **25**, 1157–1174 (2004).

1151 75. Wang, J., Wang, W., Kollman, P. A. & Case, D. A. Automatic atom type and bond type perception
1152   in molecular mechanical calculations. *J. Mol. Graph. Model.* **25**, 247–260 (2006).

1153 76. Wang, J., W. Wang, P. A. K. and D. A. C. Antechamber: an accessory software package for
1154   molecular mechanical calculations. *J. Am. Chem. Soc* **222**, U403 (2001).

1155 77. Case, D., K. Belfon, I. Ben-Shalom, S. Brozell, D. Cerutti, T. Cheatham III, V. Cruzeiro, T. Darden, R.
1156   Duke, G. Giambasu, M. Gilson, H. Gohlke, A. Goetz, R. Harris, S. Izadi, S. Izmailov, K. Kasavajhala,
1157   A. Kovalenko, R. Krasny, T. Kurtzman, T. Lee, S. L, D. Y. and P. K. AMBER 2020. (2020).

1158 78. Pérez, A. *et al.* Refinement of the AMBER force field for nucleic acids: Improving the description
1159   of α/γ conformers. *Biophys. J.* **92**, 3817–3829 (2007).

1160 79. Götz, A. W. *et al.* Routine microsecond molecular dynamics simulations with AMBER on GPUs. 1.
1161   generalized born. *J. Chem. Theory Comput.* **8**, 1542–1555 (2012).

1162 80. Salomon-Ferrer R, Götz AW, Poole D, Le Grand S, W. R. Routine Microsecond Molecular

1163   Dynamics Simulations with AMBER on GPUs. 2. Explicit Solvent Particle Mesh EwaldNo Title. *J*
1164   *Chem Theory Comput.* **10**, 3878–3888 (2013).

1165   81.   Ryckaert, J. P., Ciccotti, G. & Berendsen, H. J. C. Numerical integration of the cartesian equations
1166   of motion of a system with constraints: molecular dynamics of n-alkanes. *J. Comput. Phys.* **23**,
1167   327–341 (1977).

1168   82.   Roe, D. R. & Cheatham, T. E. PTRAJ and CPPTRAJ: Software for processing and analysis of
1169   molecular dynamics trajectory data. *J. Chem. Theory Comput.* **9**, 3084–3095 (2013).

1170   83.   Saldanha, A. J. Java Treeview - Extensible visualization of microarray data. *Bioinformatics* **20**,
1171   3246–3248 (2004).

1172   84.   de Hoon, M. J. L., Imoto, S., Nolan, J. & Miyano, S. Open source clustering software.
1173   *Bioinformatics* **20**, 1453–1454 (2004).

1174   85.   Bolger, A. M., Lohse, M. & Usadel, B. Trimmomatic: A flexible trimmer for Illumina sequence
1175   data. *Bioinformatics* **30**, 2114–2120 (2014).

1176   86.   Andrews, S. FastQC: A Quality Control Tool for High Throughput Sequence Data.
1177   http://www.bioinformatics.babraham.ac.uk/projects/fastqc (2010).

1178   87.   Li, H. Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM. **00**, 1–3
1179   (2013).

1180   88.   Li, H. *et al.* The Sequence Alignment/Map format and SAMtools. *Bioinformatics* **25**, 2078–2079
1181   (2009).

1182   89.   Quinlan, A. R. & Hall, I. M. BEDTools: A flexible suite of utilities for comparing genomic features.
1183   *Bioinformatics* **26**, 841–842 (2010).

1184   90.   Ramírez, F., Dündar, F., Diehl, S., Grüning, B. A. & Manke, T. DeepTools: A flexible platform for
1185   exploring deep-sequencing data. *Nucleic Acids Res.* **42**, 187–191 (2014).

1186   91.   Bailey, T. L. DREME: Motif discovery in transcription factor ChIP-seq data. *Bioinformatics* **27**,
1187   1653–1659 (2011).

1188   92.   Grant, C. E., Bailey, T. L. & Noble, W. S. FIMO: Scanning for occurrences of a given motif.
1189   *Bioinformatics* **27**, 1017–1018 (2011).

1190   93.   Josling, G. A., Selvarajah, S. A., Petter, M. & Duffy, M. F. The role of bromodomain proteins in
1191   regulating gene expression. *Genes (Basel).* **3**, 320–343 (2012).

1192   94.   Bunnik, E. M. *et al.* DNA-encoded nucleosome occupancy is associated with transcription levels in
1193   the human malaria parasite Plasmodium falciparum. *BMC Genomics* **15**, 1–15 (2014).

1194   95.   Oehring, S. C. *et al.* Organellar proteomics reveals hundreds of novel nuclear proteins in the
1195   malaria parasite Plasmodium falciparum. *Genome Biol* **13**, R108 (2012).

1196
1197

1198   **<u>Figure Legends</u>**

1199

1200   **Figure 1 Nine putative competitors of DNA binding by AP2-EXP were identified by an *in-***

1201   ***silico* screen and prioritized for further testing**

1202

1203   A) The crystal structure of AP2-EXP (PDB:3IGM) was used as a template to computationally

1204   dock thousands of small molecules *in silico* using AutoDock. Results were filtered for

1205   compounds that dock within 10 Angstroms of DNA binding residues with a free energy less than

1206   -5kJ/mol. Compounds matching these criteria were sourced and used for further testing.

1207

1208   B) Each compound that was prioritized based on the *in-silico* screen was assigned an identifier

1209   (A-I) and tested for anti-*Plasmodium* activity in a 48-hour growth inhibition assay against

1210   asexual blood stage *P. falciparum*.

1211

1212   **Figure 2 Four of nine predicted AP2-EXP competitors compete AP2-EXP DNA binding in**

1213   **an Electrophoretic Mobility Shift Assay (EMSA)**

1214

1215   A) Each putative AP2-EXP competitor was added to an EMSA containing the AP2-EXP AP2

1216   domain with its cognate DNA oligo. DNA binding competition will result in a loss of the shifted

1217   probe. DMSO vehicle was used as a control for normal DNA binding by AP2-EXP. 150 fmoles of

1218   DNA probe, 125ng of AP2-EXP, and 300µM of each compound were used for each lane.

1219

1220   B) Chemical structures of Compounds A-I. Compounds highlighted in black (A, B, C and I)

1221   compete DNA binding by AP2-EXP. Three of the four compounds that compete DNA binding *in*

1222   *vitro* (Compounds A, B, C and D) have a benzoxazole core moiety, denoted by a red box.

1223

1224   **Figure 3 Analogues of Compound B compete AP2-EXP DNA binding with differing**

1225   **efficacy**

1226

1227   A) The four closest structural analogues to Compound B were sourced through the Tres Cantos

1228   Antimalarial Set. These compounds were given the identifiers B-1, B-2, B-3, and B-4. Each

1229   compound kills *P. falciparum* parasites at low micromolar (0.16-1.13µM) concentrations as

1230   measured by Gamo *et al*[46].

1231

1232 B) Compounds B, B-1, B-2, B-3, and B-4 were added to an EMSA with AP2-EXP. DNA binding

1233 competition will result in the loss of the shifted probe. 40 fmoles of DNA probe, 125ng of AP2-

1234 EXP, and 300µM of each compound were used for each lane.

1235

1236 C) Compounds B, B-1 and B-4 were titrated at 25, 50, and 150µM into an EMSA with AP2-EXP

1237 to determine whether there are differences in DNA binding competition. DMSO vehicle was

1238 used as a control for normal DNA binding. 40 fmoles of DNA probe and 125ng of AP2-EXP were

1239 used for each lane. The concentration increase for each compound is indicated by the triangle

1240 from left to right.

1241

1242 D) Chemical structures of Compound B and B-1, B-2, B-3, and B-4. Compounds that compete

1243 DNA binding by AP2-EXP are highlighted in black. Each compound has a different substitution

1244 pattern on the right-side benzene ring, denoted by a red box.

1245

1246 **Figure 4 AP2 Domain Competing Compounds B and C affect *P. falciparum* at in the mid**

1247 **trophozoite stage, coinciding with the maximal expression of AP2-EXP**

1248

1249 A) Schematic for ApiAP2 domain competitor compound phenotyping time course. Highly

1250 synchronous asexual *P. falciparum* parasites were spiked with 40µM of AP2 domain competing

1251 Compounds B and C, non-ApiAP2 domain competitor Compound H, or DMSO vehicle control at

1252 24 hpi and parasite progression was monitored for 48 hours to determine the parasite killing

1253 phenotype.

1254

B) Representative Giemsa-stained microscopy images of *P. falciparum* parasites spiked with Compounds B, C, or H as described in panel A. DMSO vehicle was used as a control for normal progression through the asexual blood stage.

C) Highly synchronous asexual blood stage *P. falciparum* parasites were spiked with 40μM Compound C at eight-hour intervals in the IDC. Each assay was done in triplicate. Error bars represent standard deviation.

D) Highly synchronous asexual stage *P. falciparum* parasites were cultured with 40μM Compound C, which was washed out of the media at eight-hour intervals. Each assay was done in triplicate. Error bars represent standard deviation.

E) AP2-EXP protein expression in the asexual blood stages ring, trophozoite, and schizont, as well as Stage III gametocytes, was probed by Western blot against AP2-EXP endogenously tagged with GFP (AP2-EXP::GFP). The expected molecular weight of AP2-EXP::GFP (147kDa) is indicated by an arrow.

F) Live fluorescent microscopy of AP2-EXP::GFP was used to further probe AP2-EXP protein expression and localization. DRAQ5 was used as a nuclear stain for parasites.

Figure 5 Compound C disrupts the *P. falciparum* transcriptome specifically at 30hpi, with bias towards AP2-EXP targets predicted by ChIP-seq

A) *P. falciparum* parasites were spiked with 12μM ($0.66 \times IC_{50}$) Compound C or DMSO vehicle control at 10 hpi. Total RNA was harvested at 6, 12, 18, 24, 30, 42, and 48 hpi for quantification

1280   by DNA microarray. The full transcriptome Spearman Correlation between Compound C and

1281   DMSO control spiked parasites was plotted as a correlogram. A * indicates *p-value* < 0.05.

1282

B) Three replicates of AP2-EXP Chip-seq were collected at 30 hpi using two genetically tagged

parasite lines (AP2-EXP::GFP and AP2-EXP::HA). A no epitope control was collected using wild

type 3D7 parasites and anti-GFP antibodies. Log_2 enrichment of the immunoprecipitate (ChIP)

over Input DNA for one replicate of AP2-EXP::GFP, AP2-EXP::HA, and No Epitope Control is

plotted relative to the coding sequence start site of AP2-EXP target genes identified in at least

2/3 replicates by the MACS2 peak calling software.

1289

1290   C) CATGA is the most overrepresented DNA motif within the 240 peaks of AP2-EXP occupancy

1291   conserved in at least 2/3 ChIP-seq experiments. No significant DNA motif was recovered from

1292   the No Epitope Control.

1293

1294   D) Compound C induced changes in transcript abundance at 24-30hpi compared to transcripts

1295   bound by AP2-EXP. Of the 463 genes decreased in abundance by log_2 fold change > 2, 46 are

1296   AP2-EXP targets.

1297

1298   E) Volcano plot displaying the changes in abundance for AP2-EXP target genes at 24-30 hpi.

1299   46/93 detected AP2-EXP target transcripts decrease in abundance by log_2 fold change > 2 in

1300   the presence of Compound C.

1301

1302   F) GO-term analysis of transcripts that are both AP2-EXP targets and decreased in abundance

1303   with respect to Compound C (Bonferroni *p-value* cutoff < 0.05).

1304

1305 **Figure 6 Compounds B and C are active against mosquito stage *Plasmodium berghei***

1306 **parasites**

1307

A) *A. stephensi* mosquitoes were infected with *Plasmodium berghei* parasites. On day 10 post

infection, mosquito midguts were injected with the measured IC_{50} of Compounds B, C, F, or

DMSO vehicle control. On day 14 post infection mosquitoes were dissected to count oocysts

and midgut sporozoites. For Compound C and DMSO vehicle control, each experiment was

performed in duplicate. Compounds B and F phenotyping were performed as a single

experiment.

1314

1315 B) Midgut oocyst counts per mosquito for *P. berghei* infected mosquitoes on day 14 post

1316 infection. Compound identity is indicated in the legend. R1 and R2 correspond to biological

1317 replicates one and two, respectively.

1318

1319 C) Midgut sporozoite counts per five infected mosquitoes on day 14 post infection. Compound

1320 identity is indicated in the legend. R1 and R2 correspond to biological replicates one and two,

1321 respectively.

1322

A

AP2-EXP Crystal Structure   >10,000 Small Molecules

Computational Docking

B

Letter Code	PubChem ID	IC$_{50}$ (µM)
A	CID1365835	11
B	CID1365471	80
C	CID5750730	18
D	CID4541005	22
E	CID2343	170
F	CID4437	40
G	CID11390	34
H	CID6914273	40
I	CID25863	125

Figure 1 A-B

Figure 2 A-B

68 Figure 3 A-D

92   Figure 4 A-F

Figure 5A-F

135 Figure 6 A-C