Fragmented habitat compensates for the adverse effects of genetic bottleneck

Ari Löytynoja[1,*], Pasi Rastas[1], Mia Valtonen[1,4], Juhana Kammonen[1], Liisa Holm[1,2], Morten Tange Olsen[5], Lars Paulin[1], Jukka Jernvall[1,3], and Petri Auvinen[1]

[1]Institute of Biotechnology, HiLIFE, University of Helsinki, Finland

[2]Organismal and Evolutionary Biology Research Program, Faculty of Biosciences, University of Helsinki, Finland

[3]Department of Geosciences and Geography, Faculty of Science, University of Helsinki, Finland

[4]Current address: Natural Resources Institute Finland (LUKE), Helsinki, Finland

[5]Section for Molecular Ecology and Evolution, Globe Institute, University of Copenhagen, Denmark

[*]Corresponding author: ari.loytynoja@helsinki.fi

Abstract

In the face of human-caused biodiversity crisis, understanding the theoretical basis of conservation efforts of endangered species and populations has become increasingly important. According to population genetics theory, population subdivision helps organisms retain genetic diversity, crucial for adaptation in a changing environment. Habitat shape is thought to be important for generating and maintaining population subdivision, but empirical cases are needed to test this assumption. We studied Saimaa ringed seals, landlocked in a labyrinthine lake and recovering from a drastic bottleneck, by whole-genome sequencing 105 individuals and additional individuals from three other ringed seal subspecies. We analyzed the distribution of variation and genetic relatedness among the individuals in relation to the habitat shape. Despite a severe history of a genetic bottleneck with prevalent homozygosity in Saimaa ringed seals, we found evidence for population structure mirroring the subregions of the lake. Highlighting the significance of habitat connectivity in conservation biology and the power of genomic tools in understanding its impact, genome-wide analyses showed that the subpopulations had retained unique variation and largely complementary patterns of homozygosity. Integration of genetic analyses in conservation decisions gives hope to Saimaa ringed seals and other endangered species in fragmented habitats.

Introduction

Human activities have had profound impacts on the diversity of terrestrial and marine organisms in the form of habitat fragmentation, overexploitation and disturbance. Conservation efforts aim at retaining and increasing the census size (N) of endangered species and populations, but the significance of N_e, the effective population size, is also widely acknowledged (1–3), e.g., when designing protected areas or deciding to translocate animals. The concept of N_e was developed by Wright (4) and Fisher (5) and refers to the size of an idealized population that behaves genetically similarly to the real-life population under study. In conservation biology, the significance of N_e is that it determines the populations' ability to respond to evolutionary forces, e.g., to adapt to a changing environment or to survive under negative selection (6). While N_e can be affected by sex ratio, age structure and differences in reproductive success, one of its key determinants is the population structure. Wright (4; 7) showed that the metapopulation effective size, N_{eT}, is a function of N_T, the total population size, and F_{ST}, a differentiation between subpopulations, $N_{eT} \approx \frac{N_T}{1-F_{ST}}$. Although this suggests that the effective population size increases in structured populations, Wright made strong underlying assumptions about the nature of subpopulations, and in realistic scenarios, population subdivision is thought to reduce rather than increase the global effective size (6). Many results of Wright and Fisher's early work have later been derived from coalescence theory (8). There, N_e is defined as the coalescence time of sequences and is always increased by population subdivision.

Lake Saimaa, Finland, provides a unique opportunity to empirically study the effects of population substructure on the preservation of genetic diversity and to learn about the significance of the environment shape for the conservation efforts of endangered species in general. The Saimaa ringed seals (*Pusa hispida saimensis*) are a land-locked population of ringed seals, and along with the Arctic (*P. h. hispida*), Okhotsk (*P. h. ochotensis*), Baltic (*P. h. botnica*), and Ladoga (*P. h. ladogensis*) ringed seals, one of the five recognized subspecies of the ringed seals. According to estimates (9), the census sizes of Okhotsk and Arctic ringed seals are counted in hundreds of thousands and millions, while the census size of the Baltic population is around 10,000 and that of the Ladoga popu-

lation 3,000-5,000 individuals. While the population size of Saimaa ringed seals has recovered from 100-160 individuals in the early 1980s to more than 400 individuals in 2020 (10), the consequences of the extreme bottleneck to the population's adaptive potential and long-term prospects are unclear. Lake Saimaa is highly labyrinthine with a shore length of 13,700 km for an area of 4,400 km^2, and is further fragmented by ca. 14,000 islands. This is in stark contrast to neighboring Lake Ladoga, housing the only other freshwater subspecies of ringed seals, with a shore length 1,570 km for an area of 17,700 km^2. Ringed seals from different parts of Lake Saimaa have been shown to be genetically differentiated (11; 12), but the magnitude of this population stratification and its impact on the amount of genetic diversity and the subspecies' adaptive potential has not been assessed in detail.

Results

Since 1980s, tissue samples of deceased Saimaa ringed seals have been collected for research needs. 19 of these were whole-genome sequenced at medium (mean 11.2X) and 86 at low sequencing coverage (mean 5.2X; Table S1). To place them into a global context, 9-19 samples were sequenced from the Ladoga, Baltic, and Arctic subspecies (mean 8.3X; Fig. 1A; Table 1; Table S1). We used the 2.350 Gbp contig assembly for the Saimaa ringed seal and in most analyses focused on 55.9% of the genome defined as callable (13) and unique (14); a subset of analyses were performed on the 100 longest contigs, the shortest of these being 5.48 Mbp and them in total covering 804 Mbp (34%) of the genome. Using bwa mem (15) and samtools (16) for read mapping, ANGSD (17) and beagle (18) for genotyping and phasing, we separately processed the Saimaa samples ('Saimaa data'; n=105) and the medium coverage samples of the four subspecies ('subspecies data'; n=59). Among the 1.313 Mbp sites, we found in total 5,894,190 single nucleotide polymorphisms (SNPs) in the Saimaa data and 9,830,328 SNPs in the subspecies data.

Saimaa subspecies is genetically diverged from the others

In the principal component analysis (PCA) of the subspecies data, the first PC explains 20.8% of total genetic variation and separates the Saimaa from all others; the second PC explains 4.3% and separates the three other subspecies (Fig. 1A). Despite smaller sample sizes (n=9-19), the Arctic, Baltic and Ladoga subspecies contain significantly greater numbers of variants, from 7.98×10^6 to 8.73×10^6 SNPs each, than the Saimaa samples (n=19), in total 4.92×10^6 SNPs (Table 1). 32.9% of the variants are shared between all subspecies, while an additional 37.0% is shared between the three non-Saimaa

Table 1: Sample size, sequencing coverage, and number of variants for the four ringed seal subspecies.

	Arctic	Baltic	Ladoga	Saimaa
samples	19	12	9	19
coverage	5.1X	11.3X	10.8X	11.2X
total SNPs	8,609,960	8,731,833	7,979,288	4,922,410
private SNPs	79,879	24,304	20,265	257,481
private fixed	17	9	71	6,495

Fig. 1: Genetic divergence and demographic histories of the four ringed seal subspecies. (A) Dot plot shows the first two axes of PCA of genetic variation (dots), and background map the origin of the samples (stars). (B) The lines indicate inferred ancestral population sizes (N_e) across time. For each subspecies, the analysis was performed for ten randomly chosen sample pairs.

subspecies (Fig. S1). Consistent with this, the Saimaa sample contains more than 257,000 private SNPs and 6,485 fixed differences, while the other three subspecies have 20,200-69,900 private SNPs and 8-71 fixed differences (Table 1, Fig. S1).

PCA and SNP statistics indicate that the Saimaa population has diverged from other ringed seals and lost genetic diversity in comparison to other subspecies. To understand this in more detail, we inferred the demographic histories

Table 2: Heterozygosity of individuals (H), nucleotide diversity within populations (π), genetic divergence between populations (d_{xy}; above diagonal) and population differentiation (F_{ST}; below diagonal) for the four ringed seal subspecies.

		Arctic	Baltic	Ladoga	Saimaa
	H	0.00190	0.00211	0.00203	0.00118
	π	0.00189	0.00210	0.00204	0.00125
d_{xy}	Arctic	–	0.00206	0.00207	0.00209
or	Baltic	0.0435	–	0.00214	0.00217
F_{ST}	Ladoga	0.0750	0.0494	–	0.00218
	Saimaa	0.3343	0.3206	0.3459	–

¹²³ with MSMC2 (19) using the mutation rate estimated for polar
¹²⁴ bears, 1.826×10^{-8} substitutions per site per generation (20),
¹²⁵ generation time of 10 years (21) and ten four-chromosome
¹²⁶ sets from each subspecies. The N_e trajectories for Arctic,
¹²⁷ Baltic and Ladoga subspecies are very similar until 25,000
¹²⁸ years ago (ya; Fig. 1B); then, the Arctic population grows
¹²⁹ strongly while the Baltic population increases slightly and
¹³⁰ the Ladoga population decreases in size. The trajectories of
¹³¹ Saimaa individuals separate from others around 1,000,000 ya
¹³² and show a clearly higher N_e until 100,000 ya. After that,
¹³³ the Saimaa N_e dives and eventually drops below 1,000, more
¹³⁴ than an order of magnitude lower than for any other sub-
¹³⁵ species (Fig. 1B).

¹³⁶   Three points are noticeable. First, the diverging estimates
¹³⁷ of recent N_e for Arctic and Baltic (Fig. 1B) disagree with their
¹³⁸ very similar SNP statistics (Table 1) and suggest that, due
¹³⁹ to the low sequencing coverage, our SNP counts miss recent
¹⁴⁰ variants, affecting most the Arctic population. The MSMC
¹⁴¹ method uses the lengths and ages of coalescent blocks (19)
¹⁴² and is not significantly affected. Second, a growth in MSMC
¹⁴³ estimates can be explained either by an increase in the actual
¹⁴⁴ census size or by population structure and admixture between
¹⁴⁵ distinct lineages (4; 7; 6; 8), and the bump in the Saimaa N_e
¹⁴⁶ before 100,000 ya (Fig. 1B) possibly reflects the population
¹⁴⁷ deriving some of its ancestry from elsewhere. Similar signals
¹⁴⁸ are seen in other aquatic organisms in the Baltic Sea region
¹⁴⁹ (22–24), and an origin in an eastern freshwater refugium has
¹⁵⁰ been proposed also for Saimaa ringed seals based on their
¹⁵¹ unique mitochondrial haplotypes (25). Finally, conversion of
¹⁵² coalescent estimates into years and numbers of individuals
¹⁵³ requires a good understanding of the genomic mutation rate,
¹⁵⁴ and lacking that, our estimates should be considered approx-
¹⁵⁵ imate. This does not invalidate the clear qualitative differ-
¹⁵⁶ ences between Saimaa and three other subspecies, however.

¹⁵⁷ **Saimaa seals have lost much of their genetic variation**

¹⁵⁸ Genetic variation can be quantified as heterozygosity of indi-
¹⁵⁹ viduals (H), nucleotide diversity within populations (π) and

Fig. 2: Nucleotide diversity and runs of homozygosity across the genomes. (A, B) Number of 250 kbp genomic blocks (y axis) with nucleotide diversity in given bracket (x axis) for (A) individuals (heterozygosity; H) and (B) populations (π). Different subspecies are indicated with colors. (C) Proportion of the genome (y axis) within ROHs of at least 100 kbp in length for individuals of different subspecies. Vertical lines indicate the sample means.

genetic divergence between populations (d_{xy}). Genome-wide ¹⁶⁰
mean estimates confirm the lower variation in the Saimaa ¹⁶¹
subspecies, with H and π of 0.0012 and 0.0013 differences per ¹⁶²
site, respectively, being slightly more than half of that of the ¹⁶³
Baltic and Ladoga subspecies (Table 2). On the other hand, ¹⁶⁴
the highest d_{xy} estimates are for pairs involving Saimaa, con- ¹⁶⁵
sistent with the population containing private variation. Due ¹⁶⁶
to the lower sequencing coverage (Table 1), the diversity es- ¹⁶⁷
timates for the Arctic subspecies are likely downward biased ¹⁶⁸
(Fig. S2). The distributions of local estimates H and π, com- ¹⁶⁹
puted in 250 kbp genomic windows, peak around 0.002 dif- ¹⁷⁰
ferences per site for the three non-Saimaa subspecies and are ¹⁷¹

nearly indistinguishable from each other (Fig. 2A,B). The picture is very different for Saimaa and, while the distribution of π peaks around 0.0012 (Fig. 2B), H is concentrated close to zero with another gently sloping bump above 0.002 (Fig. 2A). This indicates that the genomes of individual Saimaa seals contain long segments with virtually no variation, but copies of these low-variation regions tend to differ among the seals.

The lack of genetic variation implicates inbreeding and formation of runs of homozygosity (ROH) in the genome. Consistent with their largest N_e, Arctic individuals have the smallest amount, on average less than 1.44%, of their genome within ROHs longer than 100 kbp (Fig. 2C). The Ladoga seals have a higher proportion of the genome covered by ROHs than the Baltic ones (7.53% vs. 4.07%), though neither show signs of contemporary inbreeding (26). In contrast to this, in Saimaa individuals between 33.2 and 52.9% of the genome are within ROHs greater than 100 kbp (Fig. 2C).

In addition to the lower absolute sequencing coverage for the Arctic samples, variant imputation (applied to compensate for the low coverage) and phasing may perform differently on populations with vastly different effective sizes. This has no impact on the big picture, though, and Saimaa clearly has a deep independent history and its size has remained small for a long time. However, the lake is extremely fragmented and previous studies have shown that seals from different parts of the lake are genetically diverged (11; 12). For the conservation efforts, it is crucial to understand the impact of the small population potentially breaking into even smaller subpopulations.

The small Saimaa population is highly structured

The presence of substructure within Lake Saimaa was confirmed with a PCA of the 98 Saimaa individuals with known geographic information: the first two PCs explain 8.6 and 4.7% of genetic variation and separate, respectively, the individuals from Northern Saimaa (NS) and Southern Saimaa (SS) from all others, leaving Central Saimaa (CS) to connect the different clusters (Fig. 3A). With one exception, the three groups occupy distinct lake regions connected by narrow straits (Fig. 3B). Additional PCs separate individuals from specific regions such as Kolovesi, the most fragmented and labyrinthine part of the lake (Fig. S3A-E).

To test whether the distribution of variation is explained by simple isolation by distance, we computed water distances between the map locations and the genetic similarities between the samples, using coancestry as defined by fineSTRUCTURE (27). The logarithmic transformation of coancestry (27) correlates strongly with geographic distances (r=-0.441, $p < 2.2 \times 10^{-16}$). However, analysis of partitioned

Table 3: Heterozygosity of individuals (H), nucleotide diversity within populations (π), genetic divergence between populations (d_{xy}; above diagonal) and population differentiation (F_{ST}; below diagonal) across the whole genome for the three Saimaa ringed seal subpopulations. Only the 22 samples born in the 2010s were included.

		Northern	Central	Southern
	H	0.00122	0.00119	0.00120
	π	0.00124	0.00116	0.00116
d_{xy}	Northern	–	0.00125	0.00125
or	Central	0.1140	–	0.00120
F_{ST}	Southern	0.0893	0.0862	–

data reveals that this correlation is driven by both genetic similarities and physical distances being very different within and between different lake regions (Fig. 3C): within regions, the correlation is strong (cor=−0.200, $p < 2.2 \times 10^{-16}$), but for sample pairs coming from different lake regions, it is almost non-existent (cor =−0.070, $p < 4.7 \times 10^{-8}$). Consistent with this, the coancestry matrix contains three large blocks of high relatedness, the corresponding individuals coming from the three distinct parts of the lake, but also blocks of a few individuals with very high relatedness (Fig. 3D). In addition to blocks, the matrix contains stripes indicating genetic identity to many other seals. One of these individuals, '2569', is the clearest deviation among the three groups and is firmly amongst the Northern samples in the PCA (Fig. 3A) despite being originally sampled deep within the Central region of the lake (Fig. 3B). '2569' shows coancestry to different lake regions (Fig. S4A) and this mixed ancestry produces one of the highest levels of genomic heterozygosity (Table S2).

Saimaa subpopulations contain private variation

To quantify the current variation within subpopulations, we extracted the 22 Saimaa samples born in the 2010s (Fig. S5). Sample '2569' was excluded as an apparent migrant and two others due to atypically high homozygosity, possibly contributing to their deaths as young pups. Consistent with the coancestry estimates, Northern samples contain more genetic variation than Central and Southern samples, and each subpopulation contains some private variation, as indicated by d_{xy} being higher than π (Table 3). The F_{ST} estimates, measuring population differentiation, between the subpopulations vary from 0.086-0.114 (Table 3), and are thus of the same magnitude as between the human superpopulations (28). F_{ST} analysis is known to depend heavily on SNP ascertainment and estimates from different data sets are not fully comparable (28). Nevertheless, in our analyses the Saimaa subpopulations (Table 3) show greater population differentiation than the three other ringed seal subspecies (Table 2). The F_{ST}

Fig. 3: Genetic variation within Saimaa. (A) First two PCs explain 8.6 and 4.7% of the variation among the 98 samples. (B) The map of Saimaa shows the fragmentation and the labyrinthine shape of the lake. The sampling locations of the studied seals are indicated with dots matching the colors in the PCA plot. Orange star indicates the living area of a seal translocated in 1992 and black asterisks mark individuals '2569' and '2575'. (C) Log10-transformed coancestry estimates and water distance between sample pairs correlate but this is largely explained by pairs coming either from the same or from different lake regions, indicated by red and blue points, respectively. Histograms on top and right show the distribution within and between lake regions. The black, red and blue regression lines are fitted to full data, to red dots and to blue dots, respectively. (D) The coancestry matrix contains three blocks of higher similarity (scale on the right with higher values indicating greater genetic similarity) that, with the exception of '2569', consist of samples from northern (top, indicated by green line), central (middle, yellow) and southern (bottom, magenta) part of the lake. The dendrogram indicates the inferred grouping of the samples, and the axis label colors (left, bottom) match those in the PCA plot.

estimates are not markedly different with the full 98 sample set, varying from 0.082-0.106.

In analyses of 250kbp windows, the distributions resemble those of the four subspecies analysis, and H concentrates close to zero (Fig. 4A) and π peaks around 0.0012 (Fig. 4B). The peak around zero for π, meaning complete loss of variation within a window, is the greatest in Southern individuals, re-flecting the low historical and current population size in that part of the lake. Crucially, π for the whole Saimaa shows significantly fewer genomic blocks void of any variation, a drop of 44.2% for windows with π below 0.0001 in comparison to the subregion mean, and in average 10.4% increase for windows with π between 0.0009 and 0.0015 (Fig. 4B,C). This shows that the Saimaa subpopulations have lost variation in

A

other   2575

H

B

NS   CS   SS   Regions mean   All Saimaa

π

C

$\Delta\pi$

Fig. 4: Nucleotide diversity within 250 kbp genomic blocks in Saimaa. (A) Number of genomic blocks (y axis) with heterozygosity in given bracket (x axis) for '2575' (blue) and all other Saimaa individuals (light gray) and the variation of the latter (red boxplots). (B) Number of genomic blocks with π in given bracket for all Saimaa (dark gray) and the average (orange) of different Saimaa regions (NS, CS, SS). (C) Change in number of windows falling in each diversity bracket in the total population and the three subpopulations, i.e., the difference between the gray and orange lines.

different parts of the genome and contain private variation not present in other subpopulations.

We evaluated the significance of population substructuring using synthetic data reflecting the properties of the Saimaa data (Fig. S6). Overall, whereas increasing number of subpopulations with limited migration leads to the loss of genetic variation in individual subpopulations, the total amount of variation retained in the whole population actually increases (Fig. S7). Interestingly, three distinct subpopulations, in-

dicating a relatively low migration rate, observed in Lake Saimaa seem to be close to the sweet spot where the subpopulations do not accumulate too many ROHs while the total population retains more variation than one panmictic population (Fig. S7). With only two subpopulations and high migration, the benefits of substructuring over a panmictic population are negligible; with greater numbers of subpopulations or very low migration rates, the high homozygosity starts to threaten the survival of the subpopulations (Fig. S7).

Coancestry reveals a past translocation experiment?

One case of migration within Saimaa is empirically known as an adult female was translocated from the north-western part to the southern part of the lake in 1992 (29). No tissue sample is available, but the female is known to have reproduced in the area since 1994 (29) and was still alive in 2020 (10). Interestingly, the PCA shows the seals around her living area to be genetically intermediate while the most distinctively "southern" seals are from the narrow entry through which the possible natural migration has to enter that part of the lake (Fig. 3A,B). There are 73 sample pairs with coancestry above 2000, all within the distinct parts of the lake (Fig. 3D, Fig. S4B). On the other hand, there are 29 cross-region sample pairs with coancestry above 1500 (Fig. S4C) and these include '2575' from the immediate vicinity of the translocated seal's living area (Fig. 3B), showing the highest genetic similarity from the southern part to the northern part of the lake (Fig. S4D). A closer look at the distribution of H reveals that '2575' has a notably lower proportion of genomic segments with no variation (Fig. 4A), consistent with the individual having mixed ancestry and containing genetic variation from different parts of the lake.

Discussion

Lake Saimaa is unique in its fragmentation and labyrinthine outline. For a comparable area, the lake shoreline is roughly 58 times longer than if the lake would be a perfect circle. The labyrinthine shape of Lake Saimaa, together with its isolation, provides a natural setting to examine how population substructure responds to habitat fragmentation. Especially considering large mammals that typically require broad home ranges, Saimaa ringed seals offer a test case to measure the genetic effects of population subdivision.

Our analyses show that Saimaa ringed seals are unique among the studied ringed seal subspecies and contain a high number of private polymorphic sites and fixed nucleotide changes, consistent with the proposed long isolation and population bottleneck in Saimaa (25). Our demographic analyses reveal the severity of this bottleneck, give an approximate

timing for the start of isolation and suggest a complex ancestry for the Saimaa population. Although the impact of the population bottleneck is evident in the total lack of variation in parts of the Saimaa genomes, a larger sample reveals a division into three strongly differentiated subpopulations. Importantly, individuals and subpopulations have lost variation across different genomic regions and, as a total, the Saimaa population retains some variation throughout most of the genome.

Detailed analyses show that subpopulation divisions can be linked to narrow straits that decrease migration. In the analysis of genetic variation, the first principal component separates the northern subpopulation from the rest, the split coinciding with the narrow straits surrounded by the city of Savonlinna. The second principal component separates the southern subpopulation, leaving the central part to connect the two extremities; other components separate geographically isolated individuals (cf. 12). While the subpopulation structure is suggestive of a star-like formation where peripheric parts of the lake are connected through a central population, the subpopulation in central Saimaa does not show the greater genetic variation as suggested by the simulation, possibly reflecting a lower historical population size.

The loss of genetic variation in Lake Saimaa seems drastic, but the situation could be even worse. Our simulations examining different degrees of migration between subpopulations suggest that the Saimaa population is close to the sweet spot where the population as a whole is retaining or even gaining variation due to substructuring. Although the unique shape of Lake Saimaa and its impact on population structure and migration patterns are natural, one can draw parallels to extensive work on ecosystems and the importance of the number and connectivity of habitat patches (e.g. 30). We emphasise that our study should not be interpreted as an endorsement of artificial division of large habitats but rather as a further example of the importance of migration and the value of even small subpopulations. Although Saimaa ringed seals are a very special case of habitat fragmentation with some migration, our findings should have both theoretical and practical significance in conservation genetics. Specifically, compared to previous reports on the substructuring of the Saimaa population (12) and population analyses in general (31), our analyses also demonstrate that simple metrics such as F_{ST} alone provide limited information. Deciphering the genetic details underlying differentiation among subpopulations and individuals can reveal crucial knowledge for a species' conservation.

Saimaa-ringed seals appear to be a success story of conservation biology and the population size of the subspecies once listed as "Critically Endangered" (32) has grown from 100-160 individuals in the early 1980s to over 400 in 2020 (10). Despite recent growth, the population remains highly vulnerable and especially warmer winters and shorter ice and snow cover pose a serious risk to the breeding and early life of the pups (10). Nevertheless, our analyses indicate that the labyrinthine shape of Lake Saimaa has benefitted the retention of genetic diversity of the population as a whole, a result that can be useful in population management strategies that include translocations.

Methods

Draft reference genome. The Saimaa reference individual was sequenced with Pacific Biosciences RSII SMRT sequencing to 50X coverage, and assembled into 2,183 contigs using FALCON (v.0.3.0; 33). The contig assembly was further error corrected by mapping 100X coverage of Illumina short reads into the assembly and running the Pilon (v.1.16) improvement pipeline (34). A detailed description of the Saimaa ringed seal reference genome will be given by Kammonen et al. (in preparation).

Callable regions. The callable regions were defined with SNPable (13) and RepeatMasker (14). The SNPable mask file was generated using bwa (v.0.7.16; 15), k-mers of 35 bp and threshold of 50%. The fasta-formatted mask was converted to a bed-formatted positive mask using makeMappabilityMask.py from the MSMC tools (19). The positive mask covered 86.7% of the genome. RepeatMasker (v.open-4.0.5) was run using the rmblastn (v.2.6.0 +) search engine and RepBase Update 20150807, and query species "Canis lupus familiaris". RepeatMasker masked 42.50% of the genome as repeats. After combining the two masks with bedtools, 55.9% of the genome was defined as callable and unique.

Population sequencing. DNA sequencing of Baltic, Ladoga and Saimaa samples was performed on NextSeq 500 and NovaSeq 6000 Illumina platforms at the DNA Sequencing and Genomics Laboratory, Institute of Biotechnology, University of Helsinki. Arctic samples were sequenced on Illumina 4000 platform at the GeoGenetics Sequencing Core, Globe Institute, University of Copenhagen.

Variant calling. Bam files were constructed with bwa mem followed by GATK IndelRealigner (v.3.8-0; 35). Variant calling was done with ANGSD (v.0.930; GATK genotyping mode: -GL 2; 17) and PCANGSD (v.0.981; number of PCA axes: -e 2 for the Saimaa dataset, -e 4 for the four subspecies dataset; 17) following the program documentation. The output of PCANGSD was converted to a vcf file using a custom script. Beagle (v.4.1; 36) was run on the vcf file in its low coverage mode (gl=input.vcf.gz) and then in its genotype mode

(gt=beagle_output.vcf.gz); for the Saimaa data, Beagle was run on all samples and for each subpopulation separately. The subsets of sites were extracted with bcftools (v.1.9-87; 16).

Runs of homozygosity. ROHs were called with bcftools roh (v.1.9-87; 37) using the complete data and assuming a PL of 30 for the two least likely genotypes. The coverage and overlap of ROH segments of different length was computed with R package GenomicRanges (38). For the sum of ROHs, only contigs with length over 100 kbp were included, covering 98.9% of the total length of the assembly.

Nucleotide diversity and F_{ST}. Genome-wide heterozygosity per sample and genome-wide nucleotide diversity for the full population was computed using bcftools and custom R code. Estimates were corrected for callable sites within windows of 250,000 bp, retaining windows with at least 40% of sites inferred as callable. Genome-wide mean heterozygosity and mean nucleotide diversity were computed across all callable sites. F_{ST} was computed with vcftools (v.0.1.13; 39) across all callable sites.

Coancestry, IBS distances and principal component analysis. Coancestry (chromosome sharing) was computed with fineSTRUCTURE (v.4.1.1; 27) using the full data for the 100 longest contigs. The principal component analysis (PCA) was performed with smartpca (v.16000) from the EIGENSOFT package (40) using the 100 longest contigs. The geographic data for the background map was from the rnaturalearth R package (https://github.com/ropensci/rnaturalearth).

Water distances. The lake Saimaa map was downloaded from the Finnish Environment Institute open data service (http://paikkatieto.ymparisto.fi/lapio/latauspalvelu.html) and the locations of the seal carcasses were received from the Natural Resources Institute Finland. The Saimaa map was converted to 576x717 raster (0.25km spacing), using R packages raster (https://rspatial.org/raster) and fasterize (https://github.com/ecohealthalliance/fasterize), and the shortest paths between the carcass locations were computed with R package gdistance (https://github.com/DistanceDevelopment/Distance), allowing eight movement directions.

Data simulation. Data simulations were performed with msprime (v.0.7.4; 41). Simulated populations started with 10,000 individuals and were reduced to 1,000 individuals 1600 generations ago; after 1200 generations the populations were split into N (N=1-4) subpopulations and, during the last 400 generations, reduced to 300 individuals; in the end, 120 individuals were sampled. (In this setup, one "subpopulation" represents the panmictic scenario.) We assumed mutation and recombination rates of 5×10^{-8} and 1×10^{-8} events per site per generation, migration rate (i.e. the fraction of population migrating on each generation) of 0.001, 0.005, 0.01 and 0.05, and simulated 25 replicates, each consisting of 330 pieces of 3.75 Mbp long sequence (in empirical data 50% sequence is in contigs longer than 3.715 Mbp). This gave a total length of 1237.5 Mbp (empirical 1231.5 Mbp), and in total 8,250 pieces of 150 kbp genomic blocks (in empirical data, 8,478 blocks and in average 145,260 sites). From each simulation outcome, we computed π within 150 kbp blocks for the subpopulations and for the total population, and compared the distributions of counts of genomic windows falling within each bracket of nucleotide diversity, reflecting the analyses in Fig. 4.

Data and code availability

DNA sequencing data and draft contig assembly have been submitted to ENA under accessions PRJEB51012 and PR-JEB51013, respectively. The code for msprime simulation and computation of nucleotide diversity estimates is available at https://github.com/ariloytynoja/saimaa-population-structure.

References

1. Frankham, R. Effective population size/adult population size ratios in wildlife: a review. *Genet. Res.* **66**, 95–107 (1995).

2. Palstra, F. P. & Ruzzante, D. E. Genetic estimates of contemporary effective population size: what can they tell us about the importance of genetic stochasticity for wild population persistence? *Mol. Ecol.* **17**, 3428–3447 (2008).

3. Hare, M. P. *et al.* Understanding and estimating effective population size for practical application in marine species management. *Conserv. Biol.* **25**, 438–449 (2011).

4. Wright, S. Evolution in mendelian populations. *Genetics* **16**, 97–159 (1931).

5. Fisher, R. A. The distribution of gene ratios for rare mutations. *Proceedings of the Royal Society of Edinburgh* **50**, 204–219 (1931).

6. Waples, R. S. Spatial-temporal stratifications in natural populations and how they affect understanding and estimation of effective population size. *Mol. Ecol. Resour.* **10**, 785–796 (2010).

7. Wright, S. Size of population and breeding structure in relation to evolution. *Science* **87**, 430–431 (1938).

8. Wakeley, J. *Coalescent Theory: An Introduction* (W. H. Freeman, 2008), 1 edn.

9. NAMMCO. Ringed seals. https://nammco.no/topics/ringed-seal.

10. Kunnasranta, M. *et al.* Sealed in a lake — biology and conservation of the endangered Saimaa ringed seal: A review. *Biol. Conserv.* **253**, 108908 (2021).

11. Valtonen, M., Palo, J. U., Ruokonen, M., Kunnasranta, M. & Nyman, T. Spatial and temporal variation in genetic diversity of an endangered freshwater seal. *Conserv. Genet.* **13**, 1231–1245 (2012).

12. Valtonen, M. *et al.* Causes and consequences of fine-scale population structure in a critically endangered freshwater seal. *BMC Ecol.* **14**, 22 (2014).

13. Li, H. SNPable. http://lh3lh3.users.sourceforge.net/snpable.shtml. Accessed: 2021-10-20.

14. Smit, A., Hubley, R. & Green, P. RepeatMasker open-3.0 (1996).

15. Li, H. Aligning sequence reads, clone sequences and assembly contigs with bwa-mem (2013). http://arxiv.org/abs/1303.3997.

16. Danecek, P. *et al.* Twelve years of SAMtools and BCFtools. *Gigascience* **10**, giab008 (2021).

17. Korneliussen, T., Albrechtsen, A. & Nielsen, R. ANGSD: Analysis of next generation sequencing data. *BMC Bioinformatics* **15**, 356 (2014).

18. Browning, B. L., Tian, X., Zhou, Y. & Browning, S. R. Fast two-stage phasing of large-scale sequence data. *Am. J. Hum. Genet.* **108**, 1880–1890 (2021).

19. Schiffels, S. & Wang, K. MSMC and MSMC2: The multiple sequentially Markovian coalescent. *Methods Mol. Biol.* **2090**, 147–166 (2020).

20. Liu, S. *et al.* Population genomics reveal recent speciation and rapid evolutionary adaptation in polar bears. *Cell* **157**, 785–794 (2014).

21. Palo, J. U., Mäkinen, H. S., Helle, E., Stenman, O. & Väinölä, R. Microsatellite variation in ringed seals (*Phoca hispida*): genetic structure and history of the Baltic Sea population. *Heredity* **86**, 609–617 (2001).

22. Koskinen, M. T. *et al.* Genetic lineages and postglacial colonization of grayling (thymallus thymallus, salmonidae) in europe, as revealed by mitochondrial DNA analyses. *Mol. Ecol.* **9**, 1609–1624 (2000).

23. Nilsson, J. *et al.* Matrilinear phylogeography of atlantic salmon (salmo salar l.) in europe and postglacial colonization of the baltic sea area. *Mol. Ecol.* **10**, 89–102 (2001).

24. Säisä, M. *et al.* Population genetic structure and postglacial colonization of Atlantic salmon (*Salmo salar*) in the Baltic Sea area based on microsatellite DNA variation. *Can. J. Fish. Aquat. Sci.* **62**, 1887–1904 (2005).

25. Heino, M. T. *et al.* Museum specimens of a landlocked pinniped reveal recent loss of genetic diversity and unexpected population connections (2022).

26. Ceballos, F. C., Joshi, P. K., Clark, D. W., Ramsay, M. & Wilson, J. F. Runs of homozygosity: windows into population history and trait architecture. *Nat. Rev. Genet.* **19**, 220–234 (2018).

27. Lawson, D. J., Hellenthal, G., Myers, S. & Falush, D. Inference of population structure using dense haplotype data. *PLoS Genet.* **8**, e1002453 (2012).

28. Bhatia, G., Patterson, N., Sankararaman, S. & Price, A. L. Estimating and interpreting FST: the impact of rare variants. *Genome Res.* **23**, 1514–1521 (2013).

29. Koskela, J. T., Kunnasranta, M., Hämäläinen, E. & Hyvärinen, H. Movements and use of haul-out sites of radio-tagged Saimaa ringed seal (*Phoca hispida saimensis* Nordq.) during the open-water season. *Ann. Zool. Fennici* **39**, 59–67 (2002).

30. Wintle, B. A. *et al.* Global synthesis of conservation studies reveals the importance of small habitat patches for biodiversity. *Proc. Natl. Acad. Sci. U. S. A.* **116**, 909–914 (2019).

31. Nyman, T. *et al.* Demographic histories and genetic diversities of Fennoscandian marine and landlocked ringed seal subspecies. *Ecol. Evol.* **4**, 3420–3434 (2014).

32. Kovacs, K. M. *et al.* Global threats to pinnipeds. *Mar. Mamm. Sci.* **28**, 414–436 (2012).

33. Chin, C.-S. *et al.* Phased diploid genome assembly with single-molecule real-time sequencing. *Nat. Methods* **13**, 1050–1054 (2016).

34. Walker, B. J. *et al.* Pilon: an integrated tool for comprehensive microbial variant detection and genome assembly improvement. *PLoS One* **9**, e112963 (2014).

35. McKenna, A. *et al.* The genome analysis toolkit: a MapReduce framework for analyzing next-generation DNA sequencing data. *Genome Res.* **20**, 1297–1303 (2010).

36. Browning, B. L. & Browning, S. R. Genotype imputation with millions of reference samples. *Am. J. Hum. Genet.* **98**, 116–126 (2016).

37. Narasimhan, V. *et al.* BCFtools/RoH: a hidden Markov model approach for detecting autozygosity from next-generation sequencing data. *Bioinformatics* **32**, 1749–1751 (2016).

38. Lawrence, M. *et al.* Software for computing and annotating genomic ranges. *PLoS Comput. Biol.* **9**, e1003118 (2013).

39. Danecek, P. *et al.* The variant call format and VCFtools. *Bioinformatics* **27**, 2156–2158 (2011).

40. Patterson, N., Price, A. L. & Reich, D. Population structure and eigenanalysis. *PLoS Genet.* **2**, e190 (2006).

41. Kelleher, J., Etheridge, A. M. & McVean, G. Efficient coalescent simulation and genealogical analysis for large sample sizes. *PLoS Comput. Biol.* **12**, e1004842 (2016).

Acknowledgements

We acknowledge University of Eastern Finland and Metsä-hallitus for the Saimaa samples, M. Verevkin for the Ladoga samples, M. Kunnasranta, P. Timonen, S. Oksanen, and J. Vierimaa for the Baltic samples, and A. Rosing-Asvid, S. Ferguson and R. Dietz for the Arctic samples; the DNA sequencing and genomics laboratory for DNA isolation and sequencing; CSC – IT Center for Science Ltd. for computational resources; and Mervi Kunnasranta, Claudius Kratochwil and Atte Moilanen for discussions and comments. JJ and PA are funded by the Jane and Aatos Erkko Foundation (4-2013, 5-2017) and PA by the LIFE programme (LIFE19NAT/FI/000832).

Authors contributions

AL, PA and JJ conceived the paper. AL designed and performed the analyses, and wrote the first draft. PR imputed and genotyped the data. MV advised in analyses of Saimaa population structure. JK assembled the reference genome. LH contributed to genome annotation. MOT provided the Arctic data. LP and PA coordinated DNA sequencing. AL and JJ finalized the text. JJ and PA head the Saimaa ringed seal genome project. All authors commented on the manuscript.

Competing interests

The authors declare no competing interests.

Fig. S1: SNP sharing among the four subspecies. (A) Sharing of the 10,123,096 polymorphic sites. (B) Sharing of the 9,059 fixed sites. Together the two sets make the 10,132,155 variable sites after MAF 2.5% filtering.

Fig. S2: Low sequencing coverage reduces the power to identify rare variants, affecting especially the heterozygosity (H) estimates. Dots show H and mean sequencing coverage for individuals of different populations, and the dashed lines the linear models fitted to each subset. For the Arctic samples, the correlation is significantly different from zero (p = 0.0001, Pearson's product-moment correlation); for the Baltic, Ladoga and Saimaa samples, the correlation is weak or absent (p = 0.0271, p = 0.1570 and p = 0.8543, respectively).

Fig. S3: Distribution of PCs and coancestry on the map. (A-E) The distribution of the first five PCs in different subpopulations (left) and on the map (right). (F) The mean coancestry with the five NS samples identified as a diverged clade in the fineSTRUCTURE analysis. The dots on the map are colored by the PC/coancestry value, from yellow (low) to red (high).

Fig. S4: Samples with very high coancestry estimates. (A) Sample pairs of '2569' with coancestry above 1500. (B) 25, 24 and 24 sample pairs, connected with lines, in the northern, central and southern part of Saimaa show coancestry above 2000. Line colors match the scale in Fig. 3D. (C) 29 sample pairs show coancestry above 1500 despite coming from different parts of the lake. Lines are colored by the subpopulations. (D) Sample pairs of '2575' with coancestry above 1500.

Fig. S5: Saimaa seals born in the 2010s chosen to represent the current population.

Fig. S6: Simulation of substructured populations. (A) The Saimaa ringed seal population shows strong differentiation with limited historical migration between the subpopulations. Due to the fragmented shape of the lake, the central subpopulation (c) connects the two peripheric (p) parts of the lake. (B) We simulated synthetic data for a population divided into N=1..4 subpopulations that were either connected with migration ($m > 0$) or evolving in isolation ($m = 0$). The population was founded by an initial population of 1,000 individuals 1600 generations ago and this was then reduced to 300 individuals during the last 400 generations. (C) Schematic view of the star-shaped migration patterns. Peripheric (p) and central (c) subpopulations behave differently when the number of subpopulation is three or higher.

Fig. S7: Preservation of variation in structured populations. (A) Distribution of nucleotide diversity (π) within 250kbp genomic blocks. Boxplots in the upper row show the number of windows (y axis) in each category of π (x axis) for the total population (red) and the average of two subpopulations. The panels (from left to right) show the results for migration rates 0.05, 0.01, 0.005 and 0.001, each consisting of 25 replicates. Boxplots in the lower row show the difference in number of windows between the two categories. Negative numbers indicate that the total population has fewer counts in that category than the subpopulations; positive numbers indicate increase in the total population. (B) Boxplots in the upper row show the number of windows in each category of π for the substructured population in total (red) and an undivided population (black). As an undivided population cannot contain migration, the same black distribution is repeated in different panels. Boxplots in the lower row show the difference in number of windows between the two categories.

Fig. S7: Continued from the previous page. (C) Distribution of nucleotide diversity (π) within 250kbp genomic blocks when the population is divided into three subpopulations. (D) Comparison to an undivided population.

Fig. S7: Continued from the previous page. (E) Distribution of nucleotide diversity (π) within 250kbp genomic blocks when the population is divided into four subpopulations. (F) Comparison to an undivided population.

Table S1: Sequencing coverage of the four-subspecies data (left) and the Saimaa data (right). The coverage was computed for the first contig (20.5 Mbp) from bam files, including all sites.

Sample	Population	bam depth
PH001	Arctic/Ittoqqortoormiit	3.442
PH002	Arctic/Ittoqqortoormiit	2.643
PH003	Arctic/Ittoqqortoormiit	4.254
PH006	Arctic/Ittoqqortoormiit	3.759
PH044	Arctic/Ittoqqortoormiit	5.888
PH067	Arctic/Qaanaaq	6.703
PH083	Arctic/Ittoqqortoormiit	5.110
PH086	Arctic/Ittoqqortoormiit	5.514
PH087	Arctic/Ittoqqortoormiit	5.230
PH090	Arctic/Ittoqqortoormiit	5.183
PH109	Arctic/Qeqertarsuaq	4.884
PH112	Arctic/Qeqertarsuaq	6.248
PH116	Arctic/Qeqertarsuaq	5.202
PH130	Arctic/Kangia	5.975
PH157	Arctic/Holman	5.316
PH158	Arctic/Holman	4.912
PH161	Arctic/Holman	5.088
PH176	Arctic/Ulukhaktok	6.220
PH185	Arctic/Ulukhaktok	5.523
BG-01	Baltic	9.497
BG-02	Baltic	8.727
BG-03	Baltic	25.012
BG-09	Baltic	11.576
BG-10	Baltic	8.583
BG-12	Baltic	10.462
H2210	Baltic	19.518
N160	Baltic	2.993
N210	Baltic	10.508
N3-09	Baltic	9.255
N7-07	Baltic	8.377
NN3-07	Baltic	11.388
L1501	Ladoga	10.545
L1683	Ladoga	12.949
L1684	Ladoga	4.410
LG-01	Ladoga	26.864
LG-02	Ladoga	8.870
LG-03	Ladoga	8.174
LG-04	Ladoga	8.630
LG-05	Ladoga	8.490
LG-06	Ladoga	8.645
258	Saimaa	13.787
292	Saimaa	13.671
819	Saimaa	11.491
1090	Saimaa	8.663
1271	Saimaa	7.451
1482	Saimaa	13.241
1983	Saimaa	22.659
2259	Saimaa	6.670
2343	Saimaa	8.569
2357	Saimaa	8.169
2399	Saimaa	11.892
2436	Saimaa	8.646
2449	Saimaa	12.885
2474	Saimaa	11.430
2496	Saimaa	8.309
2528	Saimaa	13.333
2564	Saimaa	8.831
2585	Saimaa	10.677
2614	Saimaa	12.772

Sample	bam depth	Sample	bam depth
297	3.643	2405	1.798
298	6.986	2411	4.682
1235	4.652	2421	2.225
1262	6.206	2422	5.195
1270	6.638	2426	4.626
1488	9.300	2427	10.610
1552	8.174	2432	7.058
1956	3.269	2433	5.591
1957	7.442	2440	6.053
1977	1.260	2445	4.460
2119	2.696	2452	2.797
2221	1.209	2454	6.967
2232	8.215	2456	2.619
2233	7.270	2472	5.434
2277	6.831	2477	3.938
2327	1.113	2485	1.176
2332	2.466	2497	3.276
2333	0.434	2499	5.857
2337	5.460	2500	4.045
2340	7.151	2501	6.541
2341	4.643	2504	3.594
2342	7.152	2506	8.741
2348	5.958	2509	5.502
2350	6.187	2511	5.242
2352	0.836	2513	6.235
2353	2.023	2514	5.089
2354	1.210	2527	2.055
2356	5.142	2549	4.784
2363	4.621	2550	7.326
2375	6.257	2551	6.412
2376	3.183	2555	1.772
2377	0.558	2557	5.416
2379	8.209	2563	5.927
2380	5.550	2565	8.138
2382	7.823	2566	8.144
2383	5.900	2567	6.741
2392	8.311	2568	6.452
2393	7.478	2569	9.112
2395	1.123	2570	8.488
2396	5.503	2572	5.062
2397	8.362	2573	4.459
2402	3.998	2574	8.231
2404	4.262	2575	0.342

Table S2: For each sample, sampling region and year, age at death, and heterozygosity across the longest 100 contigs.

Sample	Region	Year	Age	H	Sample	Region	Year	Age	H
819	NS	1985	0	0.001207	2411	SS	2006	0	0.001323
1090	CS	1986	0	0.001541	2421	CS	2006	16	0.001376
1235	SS	1987	0	0.001337	2422	NS	2007	0	0.001338
1262	NS	1988	0	0.001373	2426	NS	2007	0	0.001319
1482	NS	1994	0	0.001322	2427	CS	2007	0	0.001365
1488	SS	1994	21	0.001412	2432	NS	2007	0	0.001298
1552	CS	1995	0	0.001207	2433	NS	2007	0	0.001456
1956	NS	1997	0	0.001383	2436	SS	2007	0	0.001139
1957	SS	1997	0	0.001084	2440	NS	2007	0	0.001350
1977	CS	1997	0	0.001341	2445	CS	2007	0	0.001364
1983	NS	1997	0	0.001047	2449	SS	2007	6	0.001084
2119	CS	1998	6	0.001070	2452	NS	2008	0	0.001134
2221	NS	2000	28	0.001379	2454	NS	2008	0	0.001297
2232	NS	2000	5	0.001375	2456	CS	2008	0	0.001330
2233	CS	2000	0	0.001578	2472	CS	2008	0	0.001192
2259	SS	2001	0	0.001429	2474	NS	2009	2	0.001480
2277	NS	2001	NA	0.001508	2477	CS	2009	0	0.001256
2327	SS	2002	0	0.001241	2485	NS	2009	5	0.001438
2332	SS	2003	1	0.001563	2496	CS	2009	0	0.001142
2333	NS	2003	NA	0.001378	2497	CS	2010	3	0.001305
2337	NS	2003	12	0.001373	2499	NS	2010	0	0.001119
2340	NS	2003	0	0.001234	2500	CS	2010	0	0.001309
2341	NS	2003	0	0.001175	2501	NS	2010	6	0.001417
2342	NS	2003	0	0.001227	2504	NS	2010	33	0.001305
2343	CS	2003	0	0.001352	2506	NS	2010	0	0.001424
2348	NS	2003	1	0.001378	2509	CS	2010	0	0.001311
2350	NS	2003	0	0.001393	2511	NS	2010	0	0.001244
2352	NS	2003	0	0.001435	2513	CS	2010	0	0.001191
2353	SS	2004	1	0.001507	2514	CS	2011	20	0.001244
2354	SS	2004	1	0.001529	2527	CS	2011	0	0.001201
2356	NS	2004	0	0.001360	2528	CS	2011	2	0.001331
2357	SS	2004	1	0.001083	2549	SS	2012	0	0.001081
2363	NS	2004	0	0.001234	2550	CS	2012	0	0.001394
2375	CS	2005	0	0.001302	2551	NS	2012	0	0.001073
2376	CS	2005	0	0.001351	2555	SS	2012	0	0.001276
2377	NS	2005	0	0.001433	2557	NS	2012	0	0.001406
2379	NS	2005	0	0.001526	2563	CS	2013	NA	0.001046
2380	NS	2005	0	0.000997	2564	CS	2013	NA	0.001403
2382	NS	2005	0	0.001415	2565	CS	2013	NA	0.001232
2383	CS	2005	NA	0.001410	2566	SS	2013	0	0.001305
2392	NS	2005	0	0.001301	2567	CS	2013	0	0.001345
2393	NS	2005	0	0.001483	2568	NS	2013	0	0.001479
2395	CS	2006	1	0.001477	**2569**	**CS**	**2013**	**0**	**0.001667**
2396	NS	2006	5	0.001114	2570	CS	2013	0	0.001357
2397	CS	2006	0	0.001210	2572	CS	2013	0	0.001226
2399	NS	2006	0	0.001128	2573	SS	2013	0	0.001193
2402	CS	2006	0	0.001023	2574	CS	2013	0	0.001375
2404	CS	2006	0	0.001289	2575	SS	2013	0	0.001536
2405	CS	2006	0	0.001276	2585	NS	2013	0	0.001432