1 **Microclimate is a strong predictor of the native and invasive plant-associated soil**
2 **microbiota on San Cristóbal Island, Galápagos archipelago**
3

4 Alexi A. Schoenborn*,[1], Sarah M. Yannarell*,[1], Caroline T. MacVicar[2], Noelia N. Barriga-
5 Medina[5,6], Meng Markillie[7], Hugh Mitchell[7], Kevin S. Bonham[2], Antonio Leon-Reyes[5,6], Diego
6 Riveros-Iregui[3,4,5], Vanja Klepac-Ceraj[2,#], Elizabeth A. Shank[8,#]
7
8 * Co-first authors
9 # Co-corresponding Authors
10
[1] Department of Microbiology and Immunology, University of North Carolina at Chapel Hill,
Chapel Hill, NC, 27599
[2] Department of Biological Sciences, Wellesley College, Wellesley, MA, 02481
[3] Department of Geography, University of North Carolina at Chapel Hill, Chapel Hill, NC, 27599
[4] Center for Galápagos Studies, University of North Carolina at Chapel Hill, Chapel Hill, NC,
27599
[5] Galápagos Science Center, San Cristóbal Island, Galápagos Archipelago, Ecuador 200101
[6] Laboratorio de Biotecnología Agrícola y de Alimentos-Agronomía, Universidad San Francisco
de Quito USFQ, Quito, Ecuador 170901
[7] Environmental Molecular Sciences Laboratory, Pacific Northwest National Laboratory,
Richland WA, 99352 USA
[8] Department of Systems Biology, University of Massachusetts Chan Medical School,
Worcester, MA 01605
24
25 Running Title: *Galápagos Island Soil Microbiome*
26
ORCIDs:
Alexi A. Schoenborn: 0000-0003-2998-638X
Sarah M. Yannarell: 0000-0003-4504-9193
Caroline T. MacVicar: 0000-0001-8415-8291
Noelia N. Barriga Medina: 0000-0003-4995-3737
Meng Markillie: 0000-0001-8275-4882
Hugh Mitchell: 0000-0003-0143-8461
Kevin S. Bonham: 0000-0003-3200-7533
Antonio Leon-Reyes: 0000-0001-9142-9694
Diego Riveros-Iregui: 0000-0003-0919-2988
Vanja Klepac-Ceraj: 0000-0001-5387-5706
Elizabeth Shank: 0000-0002-4804-1966
39
40 Address Correspondence to: Elizabeth A. Shank, Elizabeth.Shank@umassmed.edu, 774-455-
41 6624, 368 Plantation St., ASC-5.1059, Worcester, MA 01605 or Vanja Klepac-Ceraj,
42 vklepacc@wellesley.edu, 781-253-3541, 160 Central Street, Wellesley, MA 02481

43
44

45 **Keywords:** microclimate, microbiome, invasive plant rhizosphere, soil microbial community,
46 biodiversity, 16S rRNA gene amplicons, fungal diversity, bacterial diversity

47
48
49
50

51 **Author Contributions:**
52 C = Conceptualization; DC = Data curation; FOA = Formal Analysis; FUA = Funding acquisition;
53 I = Investigation; M = Methodology; PA = Project administration; R = Resources; SO = Software;
54 SU = Supervision; VAL = Validation; VIS = Visualization; WO = Writing – original draft; WR =
55 Writing – review & editing
56
Alexi A. Schoenborn: C, DC, FOA, FUA, I, M, VIS, WO, WR
Sarah M. Yannarell: C, DC, FUA, I, M, WR
Caroline T. MacVicar: FOA, DC, VIS, WO, WR
Noelia N. Barriga Medina: I, WR
Hugh Mitchell: DC, R
Meng Markillie: DC, R
Kevin S. Bonham: I
Antonio Leon-Reyes: I, R, WR
Diego Riveros-Iregui: I, R, WR
Vanja Klepac-Ceraj: DC, FOA, M, SU, VIS, WR
Elizabeth Shank: C, DC, FUA, PA, SU, VIS, WR
68
69

ABSTRACT

Understanding the major drivers that influence soil bacterial and fungal communities is essential
to mitigate the impacts of human activity on vulnerable ecosystems, like those found on the
Galápagos Islands. Located ~1000 km off the coast of Ecuador, the volcanically formed islands
are situated within distinct oceanic currents, which provide seasonal weather patterns and
unique microclimates within small spatial scales across the islands. Although much is known
about the impacts of human activity, such as climate change and invasive plant species, on
above ground biodiversity of the Galápagos Islands, little is known about the resident soil
microbial communities and the drivers that shape these communities. Here, our goal was to
investigate the bacterial and fungal communities found in soil located in three distinct
microclimates: Mirador (arid), Cerro Alto (transition zone), and El Junco (humid), and associated
with native and invasive plant types. At each site, we collected soil at three depths (rhizosphere,
5 cm, and 15 cm) associated with the invasive plant, *Psidium guajava* (guava), and native plant
types. We determined that the sampling location (microclimate) was the strongest driver of both
bacterial and fungal communities (74 and 38%, respectively), with additional minor but
significant impacts from plant type and soil depth. This study highlights the continued need to
explore microbial communities across diverse environments and demonstrates the weight of
different abiotic and biotic factors impacting soil microbial communities across San Cristóbal
Island in the Galápagos archipelago.

IMPORTANCE/SIGNIFICANCE

Human activity such as climate change, pollution, introduction of invasive species, and
deforestation, poses a huge threat to biodiverse environments. Soil microbiota are an essential
component to maintaining healthy ecosystems. However, a greater understanding of factors that
alter these microbial communities is needed in order to find ways to mitigate and reverse the
impacts imposed by human activity. The Galápagos Islands are a unique real-world laboratory,
in that the islands' biogeography and physical locations in the Pacific Ocean provide distinct
microclimates within small geographic distances. Harnessing these distinct environments
allowed us to investigate the influence of microclimates, soil depth, and vegetation cover on
bacterial and fungal community composition.

INTRODUCTION

Biodiversity is essential for sustaining proper ecosystem function[1]. However, due to global
climate change, human impact (deforestation, introduction of invasive species, mining,
population growth), and other factors, biodiversity across a vast array of environments is
decreasing at an alarming rate[2]. Invasive species are one of the largest human-induced sources
of global environmental change[3,4]. Much of the habitable world suffers from the introduction of
invasive species; however, there are key locations that are particularly vulnerable[5,6]. The Pacific
Islands, including the Galápagos Islands, are under intense threat from invasive plant species[7].
Roughly 1,400 invasive species have become naturalized in the Galápagos Islands, with the
majority of them being plants or plant-associated organisms[8].

The Galápagos Islands are a chain of islands of volcanic origin located ~1000 km west of the
coast of Ecuador. One of the oldest islands in the archipelago is San Cristóbal. Due to their
location along the equatorial Pacific, the islands experience distinct weather changes
throughout the year, with major influences coming from the El Niño-Southern Oscillation[9] and
the Intertropical Convection Zone (ITCZ)[10]. The ITCZ largely dictates the seasonal weather
patterns observed on the islands: January - June is the hot/wet season with large rain storms
occurring across the entire islands while July - December is the cooler/dry season, with coastal

zones experiencing rainfall and cloudy conditions at elevations around 700 meters above sea
level (masl)[11,12]. This seasonal shift results in unique microclimates that are strongly dependent
on elevation, ranging from arid near the coast to "transition zone" elevations to humid at the
volcano summits[12]. These unique microclimates create distinct ecosystems over small
geographic scales, which has led to speciation and the endemism of much of the Islands' native
flora and fauna. The high levels of endemism make the Galápagos Islands particularly
susceptible to invasive species, where perturbations can quickly alter resident biodiversity and
harm fragile ecosystems[13].
127
Environmental microbes in soil and surrounding plants have the potential to greatly alter
ecosystem functions[14,15]; however, microbes are often overlooked when discussing how to
preserve biodiversity. Microbes play a pivotal role in global carbon and nitrogen cycling[16], as
well as in maintaining plant and soil health[17,18] through the prevention of plant pathogens and
pests[19] as well as the secretion of metabolites that enhance plant growth[20,21]. Soil perturbations,
such as those induced in agriculture and crop production, have been shown to reduce soil
microbial diversity[22-24]. Although there have been extensive investigations into the factors that
drive microbial communities, such as pH[25-27], salinity[28-30], soil depth[31], soil organic carbon
content[32], temperature[33], soil moisture[34], redox status[35] and plant communities[36-38], these factors
are likely environment specific; thus, continued investigation into the different drivers that apply
to unique environments and ecosystems are still needed.
139
Unlike endemic flora, which are specialized for and adapted to specific microclimates, invasive
flora are often able to tolerate a range of environmental conditions, providing them with a growth
advantage that allows them to spread quickly. Beyond outcompeting endemic flora, invasive
plants can negatively impact the soil nutrient levels as well as inhibit carbon and nitrogen cycling
[39]. However, while much is known about the impacts invasive plant species have on above-
ground flora and fauna, we are still limited in our understanding of the role invasive plant
species have on below-ground communities such as resident soil microbiota; many of plant-
microbe interactions are expected to be plant-species specific[40-44]. The invasive plant *Psidium
guajava* (common name: guava), is one of the most detrimental invasive plants on the
Galápagos Islands. Unlike the native plants on San Cristóbal, which typically thrive in only a
single microclimate, *P. guajava* can be found in most areas of San Cristóbal as well as other
inhabited islands, except in the most arid locations on the coasts[45].
152
The Galápagos Islands are uniquely suited to investigate the interplay of multiple factors on the
composition of soil microbial communities. The physical features of the island landscape allow
the impact of microclimate and plant species on soil microbial communities to be examined
across distinct environments within a small geographic location, removing confounding variables
typically present when larger physical distances must be examined to access distinct
microclimates. In this study, we investigated soil microbial community structure on San Cristóbal
Island in the Galápagos archipelago across arid, transition zone, and humid microclimates as
monitored at the end of the wet season. We characterized and compared both the bacterial and
fungal communities associated with the rhizosphere (the region of soil surrounding plant roots)
of invasive and native plant types from three distinct microclimates, designing a study to
examine these environmental variables as well as sampling depth. Our goal was to better
understand the biotic and abiotic factors impacting the composition of soil microbial
communities and their potential function. Such an understanding may allow us to harness the
biological potential of these microbes to mitigate invasive plant species or alter the impacts of
human activity on microbial communities. We determined that the strongest driver influencing
both bacterial and fungal communities in the unique landscape of San Cristóbal was
microclimate, with soil depth and plant species both modestly but significantly influencing these

communities.
171
172   **RESULTS**
173
Sampling sites on San Cristóbal Island
To determine the effects of distinct microclimates and invasive plant species on resident soil
microbiota in unique geographic locations, this study was conducted on the Island of San
Cristóbal in the Galápagos Archipelago. We selected three sampling sites that encompassed
three distinct microclimates within four miles of each other. The sites were: Mirador (leeward
side, 320 masl, arid), Cerro Alto (leeward side, 520 masl elevation, transition zone), and El
Junco (windward side, 690 masl elevation, very humid) (Figure 1A and Supplemental Figure
1A). These sites were chosen not only because they embody distinct environmental

Figure 1

Figure 1. San Cristóbal Island in the Galápagos Archipelago exhibits geographically close sampling sites with distinct microclimates. (A) The relative locations of our three field sites on San Cristóbal in the Galápagos. (B) Temperature, % relative humidity, and precipitation data from weather stations installed in 2015. Data are the average of all collected over the given season. Due to technical issues arising from their remote locations, some weather station data is absent. (C) Tables detailing the microclimates (top) and soil properties (bottom) at Mirador, Cerro Alto, and El Junco. (D) Schematic of soil sampling scheme: rhizosphere, 5 cm, or 15 – 20 cm below the plant.

microclimates that are geographically
close to one another, but also
because they house weather stations
that have been continuously collecting
environmental variables since 2015,
including temperature (°C), % relative
humidity, and average rainfall (Figure
1B). Historically, Mirador is the
warmest site in both rainy (January -
June) and dry (July - December)
seasons compared to the other two
sites (Figure 1B). Cerro Alto
experiences moderate temperatures
and moderate rainfall throughout the
year, while El Junco has the lowest
average temperatures, highest %
humidity, and highest average rainfall
(Figure 1B).
200
In addition to the three field sites
having distinct microclimates, the
vegetation, soil properties, and plants
also varied with each site (Fig. 1).
Prior to 2015, the land surrounding
Mirador was used for agricultural
purposes; it is now used as a
conservation farm to grow native
plants. The plot of land sampled at
Mirador had several citrus trees and
shrub grasses present with larger
bushes, guava, and scalasia spaced
out across the plot (Supplemental
Figure 1A). The Cerro Alto site was
densely packed with a range of plants

Supplemental Figure 1

**Supplementary Figure 1: Images of field sites and plant
species located at each site. (**A) Field photos taken in June-July
2019 of sampling sites and plants sampled at each location; Mirador
(guava and scalesia), Cerro Alto (guava and verbena), and El Junco
(guava and miconia). (B) Image of root-associated soil. C) Images
of the different soils from each sampling site.

216 including tall grasses, guava, and herbs; this land has not previously been used for agriculture,
217 but it is near land with cow pastures. The land surrounding El Junco is protected by the
218 Galápagos National Park; we sampled a plot containing guava, miconia, ferns, and small
219 grasses (Supplemental Figure 1A).
220
Soil sampling scheme
We performed soil collections at the end of the wet season, in June of 2019. At the Mirador and
Cerro Alto sites, we obtained soil samples from six native and six non-native plants, while at El
Junco we sampled from twelve of each plant type. The non-native plant, *Psidium guajava*
(guava), was present at all sites. However, there was not a single native plant species
sufficiently abundant to consistently sample at all three sites. Therefore, for each sampling site
we selected a distinct native plant species that was similar in size to guava, abundant at that
site, and within the same 50 m x 50 m sampling plot as the guava (Figure 1C): *Scalesia gordilloi*
at Mirador, *Verbena* spp. at Cerro Alto, and *Miconia robinsoniana* at El Junco (Figure 1D and
Supplemental Figure 1A). From each plant, we collected soil at three depths: (a) rhizosphere,
the soil particles adhered to the plant root and shaken off (Supplemental Figure 1B), (b) 5 cm
below surface from where the plant emerged from the soil, and (c) 15 to 20 cm below the plant

(hereafter simply as 15 cm) (Figure
1D). The soil collected at each site
differed visually in color, water
content, and texture (Supplemental
Figure 1C); we did not observe visual
differences between different soil
depths within a single sample site.
We therefore collected 144 soil
samples (Mirador and Cerro Alto
each having six native and six non-
native plants sampled at three depths
and El Junco having 12 native and 12
non-native plants sampled at three
depths).
247

Soil carbon and nitrogen are
influenced by plant type and soil
depth
To better quantify the physiochemical
differences between the soils at each
site, we conducted soil analyses on
pooled samples. From each site we
pooled soils from the same plant type

Sample	Cl⁻ (mg/kg)	SO₄²⁻(mg/kg)	%C	%N	%TOC
Mirador Rhizo G	6.99	31.0	6.24	0.58	6.12
Mirador 5 cm G	14.15	18.6	6.35	0.58	6.38
Mirador 15 cm G	11.53	<14.5	3.47	0.34	3.45
Mirador Rhizo S	6.44	27.0	4.71	0.42	4.72
Mirador 5 cm S	9.45	19.3	4.53	0.43	4.51
Mirador 15 cm S	16.20	17.4	3.69	0.35	3.72
Cerro Alto Rhizo G	9.91	17.4	7.71	0.67	6.38
Cerro Alto 5 cm G	54.59	21.9	6.63	0.61	4.91
Cerro Alto 15 cm G	7.93	<13.2	4.93	0.47	3.25
Cerro Alto Rhizo V	23.32	18.2	5.38	0.51	4.17
Cerro Alto 5 cm V	13.92	13.4	4.38	0.43	3.68
Cerro Alto 15 cm V	11.14	<19.1	3.21	0.34	2.60
El Junco Rhizo G	26.08	33.5	7.25	0.51	6.05
El Junco 5 cm G	14.03	27.1	6.39	0.47	5.06
El Junco 15 cm G	10.30	22.8	4.94	0.32	3.52
El Junco Rhizo M	20.84	26.7	6.37	0.46	5.09
El Junco 5 cm M	12.28	31.5	5.19	0.37	4.45
El Junco 15 cm M	11.01	22.8	4.94	0.30	3.25

Supplemental Table 1. Soil nutrients above detectable limits.
Samples from each site, plant, and depth were pooled, resulting in
18 samples total**.** Soil analyses were performed to measure Cl⁻, SO₄⁻
², PO₄⁻³, NO₃N, F⁻, %TOC, %C, and %N. (PO₄⁻³, NO₃N, and F⁻ were
measured but below detectable limits). Rhizo, 5 cm, and 15 cm refer
to sample depth. Plant type is indicated by: G = guava, S = Scalesia,
V = Verbena, and M = Miconia.

and sample depth (i.e., all guava from 5 cm; all miconia from rhizosphere, etc.) which led to 18
samples for soil analysis. When the data were analyzed by averaging all of the results from
samples from the same site, we did not observe significant differences for %TOC (total organic
carbon), %C, %N, or Cl⁻ mg/kg, but we did observe significant differences in SO₄²⁻ mg/kg and
pH (Figure 2A, Supplemental Table 1). El Junco had significantly elevated levels of sulfate and
lower soil pH than the other sites (Figure 2A). When we then compared samples collected from

guava versus the native
plants across all sites,
we observed
significantly (*P* <0.05)
elevated %TOC, %C,
and %N in samples
from guava compared
to the native plant
samples (Figure 2B).
Additionally, we
observed significantly
lower %TOC, %N, and
%C for the 15 cm
samples compared to
either the rhizosphere
or 5 cm depth samples
(Figure 2C). These data
indicate that the
invasive plant guava
may either directly or
indirectly impact soil
carbon levels compared

Figure 2

Figure 2. Soil properties vary by site, plant type, and soil depth. Samples from
within each site, plant, and depth were pooled, resulting in 18 samples. A)
Comparison of soil by site, averaged across other variables. B) Comparison of soil by
plant type (guava vs. native; all native samples were averaged). C) Comparison of
soil by depth (all samples were averaged by depth regardless of plant type or site).
Each dot represents a independent sample. * denotes *P* <0.05; * above a group
indicates that it is significantly different from all others on the graph. M = Mirador, CA
= Cerro Alto, EJ = El Junco. R = rhizosphere, 5 = 5 cm depth, and 15 = 15 cm.

284 to the native flora. Additionally, soil
285 closest to the plant, regardless of plant
286 type, had higher levels of %TOC, N, and
287 C compared to the 15 cm samples.
288
The dominant bacterial and fungal taxa
show distinct distribution patterns
across sites
From the soil samples collected, we
identified the soil microbial communities
present using 16S rDNA and ITS (internal
transcribed spacer) amplicon sequencing.
Rarefaction analysis indicated sufficient
read depth was achieved during
sequencing (Supplementary Figure 2).
We assigned bacterial taxonomy to
amplicon sequence variants (ASVs) using
a Naïve-Bayes classifier compared
against a SILVA reference database that
was trained on the V3-V4 region of the
16S rRNA gene[46]. When we assessed the
relative abundances of bacteria and fungi
at the phylum, order, and genus levels,
we saw distinct distribution patterns of
community composition at each site

Supplemental Figure 2

Supplementary Figure 2. Rarefaction analysis of amplicon sequences. Rarefaction curves of A) Bacterial (16S) and B) Fungal (ITS) denoting analysis at each site and each plant were sufficiently sampled and reached asymptote.

309 (Supplemental Figure 3). Furthermore, the overall patterns of both bacterial and fungal relative
310 abundances were similar across all samples from a particular site, regardless of plant type or
311 soil depth from which they were collected, indicating that each site has distinct microbial
312 communities associated with it.
313
Across all three sites, the three dominant bacterial phyla were Proteobacteria (26%),
Acidobacteriota (also known as Acidobacteria; 25%), and Actinobacteriota (also known as
Actinobacteria; 20%) (Figure 2A). The most abundant phyla showed distinct distribution patterns
across sites. The relative abundances of the Actinobacteriota, Methylomirabilota, Myxococcota,
and Firmicutes were all highest at Mirador, intermediate at Cerro Alto, and then lowest at El
Junco (Figure 3A). The Acidobacteriota and Proteobacteria exhibited the opposite pattern, with
lowest relative abundances in the Mirador samples and highest in those from El Junco (Figure
3A). In contrast, the relative abundances of the Verrucomicrobiota and Bacteroidota were
significantly higher at Cerro Alto than at Mirador or El Junco, while the relative abundances of
the Plantcomycetota and Chloroflexi were lowest at Cerro Alto compared to both Mirador and El
Junco (Figure 3A).
325
We similarly assessed the relative abundance of the dominant fungal groups across the three
sampling sites. The Ascomycota (56%), Basidomycota (20%), and Chytridiomycota (11%) were
the dominant fungal phyla at all three sites, with the relative abundance of the Ascomycota
being significantly higher at Mirador (78%) compared to Cerro Alto (54%) and El Junco (46%)
(Figure 3B). The Basidiomycota and Mortierellomycota exhibited the opposite pattern, with their
relative abundances being highest at El Junco (Figure 3B). Finally, the relative abundance of
the Chytridiomycota was highest in Cerro Alto compared to the other two sites (Figure 3B).

Supplemental Figure 3

Supplementary Figure 3: Relative abundances of bacterial and fungal communities at each sampled site. Samples are grouped by site, plant, and depth. Stacked bar plots show the relative frequency of 16S rRNA (left) and ITS (right) gene sequences assigned to each (A, D) phylum, (B, E) order, and (C, E) genus across samples. Each color represents a specific phylogenetic designation. Only taxa with relative abundances above 1% are shown; all others are grouped into the Other category. Blank bars represent missing data.

Figure 3

Figure 3: **Dominant taxa have distinct distribution patterns**. Relative abundances of (A) the eight most abundant bacterial phyla, (B) the seven most abundant fungal phyla, and C) two archaeal phyla from all samples from Mirador, Cerro Alto, and El Junco. Kruskal-Wallis ANOVA was used for initial statistical comparisons across all sites; once significance was identified, Mann-Whitney t-tests were used to compare between two sites. Asterisk (*) denotes significance *P* <0.05, if the asterisk does not have lines denoting the groups being compared, the group with the * is significantly different from all other groups in the graph.

Figure 4

Figure 4: El Junco soil samples have a lower alpha diversity compared to samples from Mirador and Cerro Alto. Bacterial (A) and fungal (B) communities across sampling sites, and between the native and invasive plant samples within each individual site for bacterial (C) and fungal (D) communities. Alpha diversity was determined by measuring the Shannon Entropy for bacterial and fungal communities in all of our samples (Mirador n=36, Cerro Alto n=34, and El Junco n=72). Higher values indicate greater alpha diversity. Sites are coded by colors. Asterisk designates significance $P < 0.01$; if there is no bar between comparisons, this means that the group with the asterisk is significantly different than all others in that graph.

335   Although we were not targeting archaea, we did detect two phyla using 16S rRNA sequences,
336   the Thermoplasmatota and Crenarchaeota at El Junco, with undetectable levels at either
337   Mirador or Cerro Alto (Figure 3C).
338
The microbial communities at El Junco are the least diverse of the three sites
We next investigated the impact sampling site and plant type had on alpha diversity, or mean
species richness. The Shannon entropy was similar at Mirador and Cerro Alto for both bacterial
and fungal communities ($P = 0.23$ and $P = 0.72$, respectively) but it was significantly lower for
the El Junco samples ($P < 0.001$ for bacterial and $P < 0.01$ for fungal communities) (Figure 4A),
indicating that El Junco has significantly lower microbial alpha diversity than either Mirador or
Cerro Alto.
346
We then compared the alpha diversity of samples grouped by plant type (invasive or native) to
investigate if the invasive guava had lower or higher diversity compared to the native plants at
each site. At Mirador and Cerro Alto, there were no significant differences in bacterial alpha
diversity between samples from each plant type (Figure 4C). However, at El Junco, guava-
associated soil had significantly higher Shannon entropy (or bacterial diversity) than the native
miconia-associated soil ($P <0.001$) (Figure 4A, right). In contrast, the Shannon entropy for the
fungal communities at Cerro Alto and El Junco were similar, while at Mirador the native-
associated soil had a significantly higher fungal diversity compared to guava-associated
samples ($P = 0.025$) (Figure 4D). This indicates that, although microbial alpha diversity is not
uniformly influenced by plant type, the bacterial communities at El Junco and the fungal
communities at Mirador have distinct Shannon entropy values depending on the microbial
association with the particular plant species.
359
Geographic location predicts soil microbial community composition
To determine the major drivers of the microbial communities, we explored the variation in
community composition across 142 samples (two Cerro Alto samples were excluded due to
quality concerns). We visualized the relationships between the composition of all samples using
non-metric multidimensional scaling (NMDS) plots based on the Bray-Curtis dissimilarity of the
samples (Figures 5A and 5B). Samples in both the bacterial and fungal NMDS plots (Figure 5)
clustered by geographic location, with 'site' explaining 74% of the observed bacterial variance
and 38% of fungal variance (PERMANOVA, $P < 0.001$ for both). The microbial communities of

El Junco soils were
most distinct from those
at Mirador and Cerro
Alto, which had some
minimal cluster overlap
(Figure 5A and 5B). We
observed significantly
different bacterial and
fungal beta diversity at
each site
(PERMANOVA, *P* =
0.001; Figure 5C) and
an increasing taxonomic
divergence between
Mirador and the other
two sites,Cerro Alto, and
El Junco
(PERMANOVA, *P* =
0.001) (Figure 5C), with
the most dissimilar
community from Mirador
occurring at El Junco.
Altogether, these data
indicate that the
composition of both the
bacterial and fungal
communities strongly
correlates with site.
396
Soil depth and plant
type also influence the
soil microbiota
We then assessed the
difference in beta
diversity between
samples associated with

Figure 5

Figure 5. All three sites have significantly different beta diversity. We compared
all (A) bacterial and (B) fungal communities. Ellipses indicate 95% confidence
intervals. All sites are significantly different from each other (*P* < 0.05) by
PERMANOVA. Beta diversity of (C) bacterial and fungal communities across
microclimates, and (D) between the native plant and guava within each individual
microclimate for bacterial and fungal communities. Beta diversity was determined by
measuring the Bray-Curtis dissimilarity for bacterial and fungal communities in all of
our samples (Mirador n = 36, Cerro Alto n = 34, and El Junco n = 72) and comparing
the obtained values to Mirador. Higher values indicate greater beta diversity. The
Bray-Curtis value of 0 indicates that the two communities are identical and share all
taxa, and the value of 1 indicates that the two communities do not share any taxa.

either the native or invasive (guava) plants and sampling depth at each site. We did not observe
a significant impact of plant type on either the bacterial or fungal beta diversity (Figure 5D).
However, the variance of the communities was significantly impacted by plant type, ranging
from 8 - 16% (based on PERMANOVA analysis), with El Junco having the greatest plant-type-
induced variance for both bacterial and fungal communities (Supplemental Figure 4).
Additionally, comparing soil depth within each site indicated that soil depth did not influence
Cerro Alto's bacterial or fungal communities, but did impact Mirador and El Junco's bacterial
and fungal communities, explaining 7.8 - 31% of their variance (Supplemental Figure 5). The
rhizosphere and 5 cm samples were more similar to each other than either was to the 15 cm
samples (Supplemental Figure 5). Overall, these data indicate that while the bacterial and
fungal communities are primarily influenced by geographic location, soil depth and plant species
also affect their composition, with the extent of that effect dependent on sampling site.
416
417 ***Dominant phyla are distinctly distributed between native and non-native sites***
418 Having observed the strongest plant-induced variance at El Junco, we next wanted to compare

Supplemental Figure 4

Supplementary Figure 4. Plant species influence soil microbiota. NMDS plots of Bray-Curtis distances to compare all samples by plant type at all sites. Bacterial communities are shown on the left and fungal communities on the right. Ellipses indicate 95% confidence intervals for samples from each plant type. Community variance attributable to soil depth and its associated *P* and NMDS-stress values are shown in the bottom corner of each graph.

Supplemental Figure 5

Supplementary Figure 5: Soil depth influences soil microbiota. (Top) NMDS plots of Bray-Curtis distances comparing all samples by sample depths. Bacterial communities are shown on the left and fungal communities on the right. (Bottom) NMDS plots of Bray-Curtis distances comparing sample depths within each indicated site. Solid symbols = native plant; open symbols = guava. Ellipses indicate 95% confidence intervals and are grouped by depth. Community variance attributable to soil depth and its associated *P* and NMDS-stress values are shown in the bottom corner of each graph.

419   the most abundant microbial phyla between
420   guava and miconia samples at El Junco. We examined the ten most abundant bacterial and six
421   most abundant fungal phyla in the rhizosphere and 5 cm soil samples (since these depths are
422   most impacted by plant type, Supplemental Figure 5). The majority of the bacterial (8 out of 10)
423   and fungal phyla (5 out of 6) relative abundances were significantly (*P* < 0.05) different between

Supplemental Figure 6

Supplementary Figure 6: Relative abundances of bacterial and fungal phyla from guava and miconia at El Junco. (A) The ten most abundant bacterial and (B) six most abundant fungal phyla of samples from the rhizosphere and 5 cm soil associated with either guava or miconia at El Junco. (C) Relative abundance of Glomeromycota from the rhizosphere and 5 cm soil samples from either guava or native plant at Mirador, Cerro Alto, and El Junco. (D) Principal component analysis of ASVs in the Glomeromycota phylum. The relative abundance was compared using a Mann-Whitney test. Asterisk (*) denotes significance P <0.05.

guava and miconia as determined by
Mann-Whitney t-tests (Supplemental
Figure 6A and 6B).

The phylum Glomeromycota was
notable based on previous work on
another Galápagos Island showing
that invasive plants have a higher
abundance of these fungi compared to
native flora[47][48]. To determine if this
trend was consistent across all sites,
we compared the relative abundance
of Glomeromycota from the
rhizosphere and 5 cm-associated
samples of guava and native plants at
Mirador and Cerro Alto. However,
there were no significant differences
between the relative abundances of
Glomeromycota based on plant type at
either of these two sites
(Supplemental Figure 6C). The
relative abundance of Glomeromycota
was similar in guava-associated
samples across all sites
(Supplemental Figure 6C), but a
principal component analysis of the
Glomeromycota ASVs showed that
different genera of these fungi are
present at each site (Supplemental
Figure 6D)

Predicted functional and ecological
roles of the bacterial and fungal
communities across the different
sites
A major caveat to studying microbial
communities is that differences in
community composition may not lead
to environmentally relevant functional
differences. To gain insight into the
functional and ecological potential of
these soil communities, we used the
predictors FARPROTAX for bacteria[49],
and FUNguild for fungi[50]. These
predictors found that the dominant
bacterial function were nitrogen
fixation, nitrate reduction,
chemoheterotrophy, and
photoheterotrophy (Figure 6A and
Supplemental Figure 7A) while the
dominant fungal ecological guilds were

Figure 6

Figure 6: **Functional and ecological potential of microbial communities across sampling sites.** (A) Relative abundances of predicted functional potential of the bacterial communities and (B) fungal guilds communities at Mirador (M), Cerro Alto (CA), and El Junco (EJ). * = *P* < 0.05 for sites significantly different from the other two sites unless connected with lines. Significance was first determined by comparing all sites using Kruskal-Wallis ANOVA, and then pairwise comparisons using Mann-Whitney t-test.

Supplemental Figure 7

Supplementary Figure 7: (A) Relative abundances of predicted functional potential of the bacterial communities at Mirador (M), Cerro Alto (CA), and El Junco (EJ). (B) Relative abundances and variation of fungal guilds across different sampling depths, plants, and sites based upon fungal *genus*-level classification. Each color represents a fungal guild. Samples are grouped by plant within each panel. (C) Relative abundances of fungal guilds at M, CA, and EJ. Asterisk (* = *P* < 0.05) denotes the site that is significantly different from the other two sites, unless two sites are connected with lines. Significance was determined by first comparing all sites using Kruskal-Wallis ANOVA, and then performing pairwise comparisons using Mann-Whitney t-test.

saprotrophs, epiphytes, endophytes, and mycorrhizae (Figure 6B and Supplemental Figure 7B).
These assigned categories were predominantly driven by sam pling site. Relative to the other
two sites, El Junco was predicted to have the highest abundance of nitrogen fixers, with lower
levels of nitrate reducers or denitrifiers, whereas Mirador was predicted to be dominated by
nitrate reducers; Cerro Alto was predicted to have the highest abundance of denitrifiers, with
moderate levels of nitrogen fixers or nitrate reducers (Figure 6A). In terms of carbohydrate
degradation, each site was predicted to have distinct abundances of xylanolysis-active bacteria,

Figure 7. Distinct associations of six nitrogen fixing genera: *Bradyrhizobium*, *Allorhizobium-Neorhizobium-Pararhizobium-Rhizobium, Rhodomicrobium, Roseiarcus (Bekjerinckiaceae* family*), MND1 (Nitrosomonadaceae* family*)*, and *Frankia*. For each genus, the graph on the left indicates the relative abundances from all samples from Mirador, Cerro Alto, and El Junco; the graph on the right shows relative abundances of the genus for each sampling depth and plant type. Kruskal-Wallis ANOVA was used for initial statistical comparisons across sites. Mann-Whitney t-tests were sued to compare between two sites. Asterix (*) denotes significance P<0.05, the line denotes two groups being compared.

while El Junco was distinct in its abundance of bacteria capable of cellulolysis, chitinolysis, and
fermentation (Figure 6A).

For the fungal communities, the FUNguild predictions identified a number of prevalent
saprotrophic operational taxonomic units (OTUs) that were distinctly abundant at multiple sites
(Figure 6B). Multiple of these saprophyte OTUs (OTUs 446, 128, 113, 448) as well as other
fungal guilds (mycorrhizal, epiphytic, and endophytic) were significantly enriched at El Junco
relative to the other two sites (Figure 6B). In contrast, when we assessed the fungi predicted to
be present at Mirador and Cerro Alto, they were instead dominated by fungi that fell into animal
and plant pathogen guilds (Supplemental Figure 7B). Overall, these data suggest that not only
are the microbial communities compositionally distinct at these three sample sites, but also that

their predicted functional and
ecological capabilities are
distinct as well.
496
El Junco guava-associated
soil has an increased
prevalence of free-living
nitrogen fixers compared to
miconia-associated soil
Based on El Junco having the
highest predicted abundance
of microbes involved in
nitrogen fixation (Figure 6),
we elected to further probe
the microbial composition of
the samples at this time.
Specifically, since the guava-
associated soil was found to
have a high percentage of
total nitrogen (Figure 2), we
delved into the relative
abundances of genera
suggested to be free-living
nitrogen fixers, including:
Rhizobiums,
Rhodomicrobiums,
Roseiarcus, MND1, and
Frankia. All of the genera
investigated, with the
exception of
Rhodomicrobium, had
significantly elevated relative
abundances in guava-
associated soil compared to
miconia-associated soil at El
Junco ($P < 0.05$) (Figure 7).
529

Supplementary Figure 8. Taxonomic and functional profiling comparing native and invasive species rhizobiomes. Three samples each from guava and miconia were analyzed using shotgun metagenomic sequencing. A) Principal coordinate analysis (PCoA) using classical multidimensional scaling of Bray-Curtis dissimilarity in taxonomic profiles. Fourteen species in total were identified, 7 of which were found in all samples (mean prevalence = 64%). Three species (*Dyella* sp. DHG54, *Nitrolancea hollandica*, and *Limnochorda pilosa*) were found only on miconia, and 3 (*Actinospica robiniae*, *Bradyrhizobium erythrophlei*, Candidatus *Solibacter usitatus*) were found only on guava. B) PCoA of functional profiles, after removing reads that could not be assigned to any known gene function (mean 84.7%). Of 837,903 identified genes, 206,699 (24.7%) were found in all samples (mean prevalence 53.2%), 3,846 (0.46%) were found only in miconia, and 13,020 (1.55%) were found only in guava. C) Genes (matching reads per kilobase per million reads, RPKM) identified as *nif* (nitrogen fixation), *ure* (urease - nitrogen recycling), or other nitrogen reduction genes**.**

While computational predictors such as those based on 16S rRNA gene and ITS used above
are suitable proxies for obtaining insights into the function of microbial communities,
metagenomes, or all the genes present in a sample, are capable of providing deeper insights
into the functions of the microorganisms associated with each sample. To further explore the
functional component of the microbial communities associated with El Junco's native and
invasive plants, we therefore sequenced metagenomes from three randomly selected miconia
and three randomly-selected guava-associated rhizosphere samples. Taxonomically, there were
no differences detected between these six randomly selected samples, regardless of whether
taxa identifications were derived from the 16S or ITS amplicon data or from the metagenomic
data. We also did not see any statistical differences in canonical nitrogen-fixation genes.
However, when looking at the predicted functions from the metagenomic samples, the samples
associated with guava were distinct from those associated with miconia (Supplementary Figure
8).
543

DISCUSSION

Elevation gradients are often exploited to investigate the environmental impacts on biodiversity. Previous work characterizing microbial community diversity has demonstrated conflicting impacts of elevation; some studies found higher microbial diversity at higher elevations[51,52], while others found microbial diversity did not change with elevation[53,54]. These discrepancies in how elevation impacts soil microbiota may reflect the different environmental factors found at the specific sites and elevations being investigated[55]. Some of the most analogous studies to ours have been conducted in Hawaii. Both the Galápagos and Hawaiian Islands are volcanic in origin, forming as a result of hotspots beneath their respective tectonic plates. Although both of the above-sea level islands of these archipelagos are similar in age[56,57], the soils at our sites are ~800,000 - 1,000,000 years in age[58], while one Hawaiian study examining a temperature gradient was examining very young soils (~20,000 years old)[59], and another examining an extensive elevation-precipitation gradient examined soils ~150,000 years in age[52]. In addition, the Galápagos tends to be cooler than expected for its equatorial location and the climate in the Galápagos is drier than in Hawaii, with fewer dust depositions from Asia[57]. Notably, our elevation resolution is sparse (since we only sampled from three elevations) compared to the larger number of samples taken in these Hawaiian studies[52,59]. Nevertheless, a comparison of our results is informative. One Hawaiian study examined sites where the mean annual temperature (as well as the carbon flux, but not the total carbon storage) varied, but where vegetation, soil type, moisture, and pH were constant[59]. No differences in bacteria richness was observed at these sites, indicating that other factors (i.e. pH, plants, precipitation) appear more important than temperature in driving bacterial community composition[59]. In the study of elevation-precipitation gradients on Hawaii, they observed that fungal richness increased with increasing precipitation, while bacterial richness appeared unimodal[52]. This is in contrast to our results, where both bacterial and fungal diversity metrics were lowest at the highest elevation (El Junco) compared to lower elevations. The relationship between pH and microbial diversity was also different between our work and that in Hawaii: in our study, higher pH correlated with higher bacterial and fungal diversity, while in Hawaii, fungal diversity was decreased at higher pH[52]. Overall, as a review by Looby and Martin 2020 points out, differences in biodiversity across topographic gradients are likely due to variations in a range of abiotic factors such as precipitation, landscape, and temperature[55].

Abiotic factors previously shown to influence microbial community diversity and composition include pH[27,60,61], temperature, soil moisture[27], and organic content[27,6263]. Here we determined that, overall, the site from which a sample is obtained (Mirador, Cerro Alto, or El Junco) accounts for ~70% of the bacterial and ~40 % of the fungal variance observed. El Junco having the lower overall diversity is consistent with prior studies indicating that acidic soils lead to lower bacterial richness[27]. In addition, El Junco is significantly more humid than the other two sites, and samples from this site exhibited the most distinct communities compared to Cerro Alto (transition zone) and Mirador (arid). Previous work exploring the influence of precipitation on microbial communities has demonstrated inconsistent patterns. In some cases, more humid environments have increased biodiversity compared to more arid sites[64,65], while other studies have seen no changes in biodiversity metrics across precipitation gradients[66,67]. Cerro Alto, the site with the highest bacterial and fungal diversity, receives a moderate and fluctuating amount of precipitation. Previous work exploring wet-dry cycles and transition zones has indicated that differing microbial communities are present depending on whether the location is in the wet or dry part of the cycle[68-70]; this could explain the enhanced biodiversity observed at Cerro Alto. One limitation of environmental DNA sequencing is that sampling captures all DNA present in the sample regardless of whether the cells are dead, dormant, or alive. Thus, future work specifically investigating which microbial members are metabolically active during different

594   seasons and environmental conditions (for instance with BONCAT or stable isotope probing[71,72]
595   along with manipulative experiments (such as altering water availability in soil cores) could
596   provide additional insights into the influence of moisture and precipitation on microbial
597   communities at these sites.
598
In addition to site-specific differences, we identified plant type as a minor but significant
additional driver influencing microbial community variance between samples. Bacterial
community composition varied between plant types in samples from Mirador (arid) and El Junco
(humid), while the fungal community was influenced by plant type at all three sites. Plant-
microbe interactions are a significant contributor to ecosystem functions and processes[73], and
the microbes involved in these interactions have the potential to enhance plant health[74-76].
However, how plant-microbe interactions shape the microbial community composition remains
unclear, with some studies demonstrating plant type influences community[36-38,77], while others
do not observe a plant influence[78-81], with one study showing that fungal endophytes varied over
an elevation gradient, even from within a single plant host[82]. Plant roots excrete chemicals,
nutrients, and carbon sources into the soil, which can alter microbial communities within the
rhizosphere[83]. In our study, except for the bacterial communities at Cerro Alto, where the plant
type does not significantly contribute to the observed community variance, we saw that
approximately 9% of the bacterial and 12% of the fungal variance is explained by 'plant type',
and that these values increase when selectively analyzing the samples closest to the plant root
(rhizosphere and 5 cm).
615
Consistent with other soil microbiota studies[84,85], the dominant bacterial phyla we observed were
Acidobacteria, Actinobacteria, Bacteroidetes, Proteobacteria and Verrucomicrobia, and the
abundance of these taxa tracked with different environmental variables present at each site.
Actinobacteria, which form spores and thus are abundantly found in deserts[84,86] were most
abundant at Mirador, our most arid site. At El Junco, our most humid site, Proteobacteria and
Acidobacteria were abundant, consistent with data showing they are less abundant in arid
environments[84]. The acidic soil of El Junco (pH ~4.4) is consistent with the high prevalence of
the acid-loving Acidobacteria found there[25,84,87,88]. In terms of fungi, the Ascomycota and
Basidiomycota, frequently dominant phyla in soils[89-92], were also abundant in our samples.
Ascomycota are more abundant in more arid climates[89,93], and were found at high levels at
Mirador, while Basidiomycota are more abundant in high precipitation areas[52] and were most
abundant at El Junco. Thus, our data supports the idea that environmental factors such as soil
moisture and pH are important determinants of microbial community composition.
629
Glomeromycota, which include the arbuscular mycorrhizal fungi (AMF)[94,95], represented ~3% of
the overall fungal abundance observed per site. AMF form an intricate symbiotic relationship
with plants, enhancing the nutrient and water uptake ability of plants and their ability to tolerate
stress, while the plant provides carbon to the AMF[96-101]. Approximately 90% of terrestrial plants
worldwide form symbiotic relationships with AMF[102,103]. However, the native plants on the
Galápagos Islands have significantly lower levels of AMF compared to similar plants on
mainland Ecuador[48,104] and invasive plants on Santa Cruz Island (another island in the
Galápagos archipelago), had higher levels of AMF compared to native plants there; this was
particularly true for guava[47,48]. Consistent with this, at the El Junco site, we found that soil
associated with the invasive guava had significantly elevated levels of Glomeromycota
compared to the native miconia; similar levels of Glomeromycota were observed between plants
at Mirador and Cerro Alto. Although the addition of AMF is typically considered beneficial to the
environment by restoring soil health and enhancing plant growth[105-107], it is possible that in this
environment, AMF could instead enhance the survival of invasive guava and thus be detrimental
to native plants.

645
In addition to examining microbial diversity and abundance, it is possible to use amplicon data
to gain insights into the potential functional roles of these communities using predictors such as
FARPROTAX and FUNguild. While caveats of these approaches exist, in the absence of
extensive (and expensive) SIP-metagenomic or meta-transcriptomic studies, these are
reasonable analyses to obtain insights into these communities. Functional approaches also
enable an examination of the potential functional redundancy existing within the population, a
phenomenon where distinct species perform the same ecosystem function[108,109]; such functional
redundancy is expected to increase with an increase in population biodiversity[110][111]. Our
observation that plant and animal pathogens were predicted to be more abundant at Mirador
and Cerro Alto relative to El Junco is consistent with Mirador previously being used for
agricultural purposes and Cerro Alto being in close proximity to a current agricultural site,
whereas El Junco is protected land distant from human activity. We also identified multiple
congruent predictions at our most humid site, El Junco. Denitrification, nitrate reduction, and
fermentation are all metabolic processes that occur in anaerobic conditions. All of these
processes were predicted to be low at El Junco, indicating that these soils are likely largely
aerobic. While our existing metagenomic data do not allow us to draw conclusions about the
functional differences between sites, we expect that future work in this area will permit direct
evaluation of microbial activity in these soils and thus a more nuanced understanding of their
differences and similarities.
665
This study provides insight into the major predictors of soil microbial communities on the Island
of San Cristóbal, Galápagos. We identified non-uniform correlations between elevation and
environmental variables and microbial patterns at different geological and geographic locations.
Because these volcanic islands and their soils are relatively young compared to continental
soils, our findings suggest that microbial populations may consistently form under similar
environmental conditions regardless of geographic location or relative age of landmass. The
snapshot in time described here demonstrates that the bacterial and fungal soil communities at
these sites are largely influenced by the microclimate and soil characteristics of each site, along
with less dramatic effects by soil depth and plant type. The Galápagos Islands are expected to
experience the impacts of climate change early relative to other areas, and thus are hot-spots to
investigate this global environmental perturbation[112]. Future work investigating how these soil
communities shift over time and in response to changes in the seasons is needed to address
and potentially mitigate the effects of human-induced global changes such as climate change.
679
680
681 **ACKNOWLEDGEMENTS**
682

683 This research was authorized under the agreement between the GSC-USFQ, Galápagos
684 National Park and the Ministry of Environment of Ecuador with reference number MAE – DNB –
685 CM – 2016 – 0041.
686
687 We would like to thank the UNC Center for Galápagos Studies (CGS) for funding provided to
688 E.A.S. and D.R-I. through their Galápagos Seed Grant Program in 2018 and 2019 that allowed
689 us to initiate and carry out these studies. We would also like to thank the administrators, staff,
690 and scientists at the Galápagos Science Center for providing space, assistance, and equipment
691 to carry out this work.
692
This material is based upon work supported by the U.S. Department of Energy, Office of
Science, Office of Workforce Development for Teachers and Scientists, Office of Science
Graduate Student Research (SCGSR) program (fellowship to S.M.Y). The SCGSR program is

administered by the Oak Ridge Institute for Science and Education (ORISE) for the DOE.
ORISE is managed by ORAU under contract number DE-SC0014664. All opinions expressed in
this paper are the author's and do not necessarily reflect the policies and views of DOE, ORAU,
or ORISE.

A portion of this research was performed on a project award
(10.46936/expl.proj.2019.51105/60000139, to E.A.S.) from the Environmental Molecular
Sciences Laboratory, a DOE Office of Science User Facility sponsored by the Biological and
Environmental Research program under Contract No. DE-AC05-76RL01830.

We would like to thank the Wellesley College Undergraduate Student Summer Research
Program for providing funding to support C.M. work on this research.

COMPETING INTERESTS

Authors have no competing interests to report.

MATERIALS AND METHODS

Soil sample collection

Soil samples were collected at three sites on San Cristóbal Island, Galápagos Islands, Ecuador on consecutive days in June 2019. The three collection sites were: Mirador (June 20, 2019; lat: -0.886068, long: -89.539818), Cerro Alto (June 21, 2019; lat: -0.883333, long: -89.516667) and El Junco (June 22, 2019; lat: -0.8966, long: -89.480206). At Mirador, we collected soil surrounding 6 guava and 6 scalasia plants. At Cerro Alto, we collected soil surrounding 6 guava and 6 verbena, and at El Junco, we collected soil surrounding 12 guava and 12 miconia plants. Enhanced sampling was performed at El Junco as there were sufficient numbers of plant species within our 50 m x 50 m plot. Additionally, this site was the most preserved and had limited human impact thus, we wanted to have enough power to determine if invasive plants had unique or altered microbial communities compared to native plants. All equipment and gloves were wiped down between sampling using isopropanol wipes. From each plant we collected roughly 5 g of soil from three depths per plant (rhizosphere, 5 cm below ground, and 15 - 20 cm below ground surface). To collect soil from the rhizosphere, we pulled roots out and shook off soil into a 5 mL conical tube. To collect 5 cm and 15 - 20 cm sample depths, we measured the depth from the surface and collected plugs horizontally from the soil. Collections took less than 4 hours at each site and samples were stored in a cooler with ice packs for transport until returning to the lab, where samples were allotted for microbiological analysis and frozen at -20 °C until further processing.

DNA isolation

After transport from the collection site on ice, soil samples for DNA isolation were aliquoted (~1 g into 1.5 ml Eppendorf tubes) and stored immediately at -20 °C. DNA isolation was performed using Qiagen DNeasy PowerSoil Kit (Cat # 12888-100), according to the manufacturer's instructions with slight modifications: samples were vortexed for 30 min at maximum speed in the PowerBead tubes, and DNA was eluted in 50 µl of Solution C6. Isolated DNA yield and quality was checked using a Spectrophotometer – Thermo Scientific NanoDrop 2000 (Cat #ND-2000). DNA extracts were transferred on ice to the high-throughput sequencing facility at University of North Carolina at Chapel Hill for amplicon sequencing or to EMSL for

746   metagenomic sequencing.
747

Amplicon 16S and ITS rRNA gene sequencing

Members of bacterial and fungal communities were identified by sequencing the V3-V4 region of
the 16S rRNA gene and the ITS region, respectively. Soil samples were sequenced according
to the protocol described by Caporaso et al.[113]. The extracted DNA was amplified with barcoded
primers to enable multiplexed sequencing. We used 341F 5'-CCTACGGGNGGCWGCAG-3'
and 806R 5'-GACTACHVGGGTWTCTAAT-3 for bacteria and ITS-9F 5'-
GAACGCAGCRAAIIGYGA-3' and ITS-4R 5'-TCCTCCGCTTATTGATATGC-3' for fungi. The
16S and ITS libraries were constructed with the same barcodes for each sample and combined
into the same amplicon pool after they were purified using AMPure beads (Beckman Coulter,
Illinois, USA), pooled into a library (100 ng) and quantified by qPCR. Twenty percent of
denatured PhiX was added to the amplicon pool (12 pM) and sequenced on the MiSeq platform
using 300+300 bp paired-end V3 chemistry (Illumina, San Diego, CA). Raw sequence files were
deposited to the NCBI Sequence Read Archive under the BioProject accession number: XXX.
761

Metagenomic sequencing

The template library was generated using Nextera XT Library prep kit (FC-131-1024) according
to the manufacture protocol. Pair-end sequencing with read length of 150bp was done on
NextSeq550 platform using NextSeq 500/550 High Output v2 kit 300 cycles (cat#20024908).
Shotgun metagenomic sequences were analyzed using the bioBakery suite of computational
tools[114]. First, KneadData (v0.7.7) was used to perform quality control of raw sequence reads,
such as read trimming and removal of reads matching a human genome reference. Next,
MetaPhlAn (v3.0.7, using database mpa_v30_CHOCOPhlAn_201901) was used to generate
taxonomic profiles by aligning reads to a reference database of marker genes. Finally, HUMAnN
(v3.0.0a4 with databases) was used to functionally profile the metagenomes[115]. Sequences are
deposited in the NCBI database XXX under the accession number XXX

Processing of the sequence data

We processed sequences in QIIME 2 v2020.8[116], using a modified pipeline workflow[117]. Primers
were removed using the cutadapt plugin[118] and the reads were sorted based on the primers into
either the 16S (bacterial) or ITS (fungal) group. Raw sequence reads from soil samples were
denoised, filtered and clustered into amplicon sequence variants (ASVs) using the Divisive
Amplicon Denoising Algorithm (DADA2) plugin[119]. For the bacterial sequence data, we used the
following parameters: ASVs with a frequency of <0.1% of the mean sample depth were
removed to account for a bleed-through between MiSeq runs, and the remaining reads were
used for subsequent analysis. For the 16S rRNA gene dataset, sequences belonging to
mitochondria and chloroplasts were filtered out. After denoising and filtering, the number of
sequences per sample ranged between 50 and 555,568 sequences for the 16S run and 7 to
105,710 sequences for the ITS run. Samples with fewer than 20,000 sequences were excluded
from the analysis. These included samples M6-A, C3-C, and C1-B for the 16S rRNA gene
amplicon dataset and C1-C and C6-A for the fungal dataset. A multiple-sequence alignment
was created using MAFFT, and FastTree was used to create an unrooted phylogenetic tree,
both with default values[120]. Taxonomy was assigned to the ASVs using a Naïve-Bayes classifier
compared against SILVA v138.99 reference database trained on the 515-926 region of the 16S
rRNA gene[46]. For fungal ASVs, Naïve-Bayes classifier was compared against the UNITE
(ver8_99_s_04.02.2020) ITS database files.
792

To visualize the community composition, we constructed bubble plots and stacked bar plots of
bacterial relative abundances using the ggplot2 package in R v4.0.3[121]. Alpha-diversity was
estimated using Shannon diversity index[122]. Bray-Curtis dissimilarity was used to assess

compositional differences between conditions (sites or plants or soil depth). The clustering of
bacterial communities was visualized via NMDS plots constructed from Bray-Curtis distance
matrices using ggplot2[121]. We used Permutational Multivariate Analysis of Variance
(PERMANOVA) tests to determine whether the microclimate, sampling depth, or the plant
species shaped microbial community structure[123] using the adonis function from the R vegan
package[124]. We used Bray-Curtis dissimilarity with 999 permutations. The strata term was used
to limit permutations within individual sites to account for repeated measurements. We
compared across both experiments, but we also ran PERMANOVAs separately for each of the
microclimates.
805
Functional group characterization and core taxa identification
To predict the fungal guild or functional groups, we used FunGUILD v1.0, an open annotation
tool that predicts ecological guild by parsing fungal OTUs[50]. To visualize these results, we
constructed a stacked bar plot of guild relative abundances using the ggplot2 package in R
v4.0.3. To predict the functional potential of the soil microbiome from 16S rRNA genes, we used
FAPROTAX[49]. FAPROTAX uses experimental evidence, such as culturing, to estimate
metabolic phenotypes based on the 16S rRNA gene. To identify core taxa associated with the
rhizosphere of each plant or site, we used COREMIC[125].
814
815   **Differential abundance**
816   Associations of bacterial and fungal phyla across different sites or plants was evaluated by
817   using DESeq2[126] in MicrobiomeDB[127]. DESeq2 uses an empirical Bayesian approach to fit
818   abundance data to a negative binomial model and converts P values for differential abundance
819   to q values to correct for multiple hypothesis testing.
820
Weather stations
Weather stations were deployed in 2015 at each of the sampling sites (Images of weather
stations from Mirador and Cerro Alto in Supplemental Figure 1). The weather stations collected
data on precipitation, wind speed and wind direction, temperature, and solar radiation. Data
gaps were the result of instrument failure. Due to remote location and difficult terrain, repairs to
instrumentation took time.
827
Soil Analysis
Soil was collected and dried at 56 °C for 72 hours and then moved to room-temp and air-dried
for an additional 2 weeks. Soil was then stored in plastic ziploc bags at room temperature in the
dark until soil analysis was performed. We pooled ~1 g from each of the 144 by site, depth, and
plant type (ex. El Junco guava rhizosphere samples) resulting in 18 samples total. Soil analysis
was performed by the University of Georgia Athens, Agriculture, and Environmental Service
Lab. The Dumas method was used for the analysis of total Carbon and Nitrogen and previously
described[128], briefly, for total organic carbon soil is treated with water and 8% sulfuric acid,
mixed, and incubated for several hours and then placed in the soil dryer to volatilize carbonates
from the sample. Soil (~0.5 g) is loaded into a steel crucible and combusted in an oxygen
atmosphere at 1200°C in a Elementar Vario Max Total Combustion Analyzer. Elemental C and
N are converted into CO_2, NO_x, and N_2. Carbon content was determined using an infrared cell
and N2 was determined using a thermal conductivity cell. Results were reported as a %. To
measure anions USEPA Method 300 was used as previously described () briefly, a Dionex
Model DX-120 ion chromatograph (Sunnyvale, CA, USA) was used with a 25 µl sample loop
injector and AS40 automated sampler employed along with a Dionex IonPac AS14A separation
column (3 × 150 mm) and IonPac AG14A guard column (4 × 50 mm). A Dionex ACRS-500 (2
mm) Anion Self-Regenerating Suppressor was used for conductivity suppression before
detection by a Dionex DS4-1 conductivity cell. Soil was mixed at a 1:1 ratio in deionized water

for an hour and filtered before running the samples. Eluent: 8 mM Na_2CO_3 + 1 mM NaHCO3 (1
ml min−1) was added at a flow rate of 0.48 ml per minute.
849
850
851

REFERENCES

1. Naeem S, Duffy JE, Zavaleta E. The functions of biological diversity in an age of
extinction. Science. 2012;336(6087):1401-1406. Epub 2012/06/16. doi:
10.1126/science.1215855. PMID: 22700920.
2. Butchart SH, Walpole M, Collen B, van Strien A, Scharlemann JP, Almond RE, Baillie
JE, Bomhard B, Brown C, Bruno J, Carpenter KE, Carr GM, Chanson J, Chenery AM,
Csirke J, Davidson NC, Dentener F, Foster M, Galli A, Galloway JN, Genovesi P,
Gregory RD, Hockings M, Kapos V, Lamarque JF, Leverington F, Loh J, McGeoch MA,
McRae L, Minasyan A, Hernandez Morcillo M, Oldfield TE, Pauly D, Quader S, Revenga
C, Sauer JR, Skolnik B, Spear D, Stanwell-Smith D, Stuart SN, Symes A, Tierney M,
Tyrrell TD, Vie JC, Watson R. Global biodiversity: indicators of recent declines. Science.
2010;328(5982):1164-1168. Epub 2010/05/01. doi: 10.1126/science.1187512. PMID:
20430971.
3. Pejchar L, Mooney HA. Invasive species, ecosystem services and human well-being.
Trends Ecol Evol. 2009;24(9):497-504. Epub 2009/07/07. doi:
10.1016/j.tree.2009.03.016. PMID: 19577817.
4. Pimentel D, McNair S, Janecka J, Wightman J, Simmonds C, O'Connell C, Wong E,
Russel L, Zern J, Aquino T, Tsomondo T. Economic and environmental threats of alien
plant, animal, and microbe invasions. Agriculture, Ecosystems & Environment.
2001;84(1):1-20. doi: https://doi.org/10.1016/S0167-8809(00)00178-X.
5. Dulloo M, Kell S, Jones C. Impact and control of invasive alien species on small islands.
International Forestry Review - INT FOR REV. 2002;4:277-285. doi:
10.1505/ifor.4.4.277.40525.
6. Caujapé-Castells J, Tye A, Crawford DJ, Santos-Guerra A, Sakai A, Beaver K, Lobin W,
Vincent Florens FB, Moura M, Jardim R, Gómes I, Kueffer C. Conservation of oceanic
island floras: Present and future global challenges. Perspectives in Plant Ecology,
Evolution and Systematics. 2010;12(2):107-129. doi:
https://doi.org/10.1016/j.ppees.2009.10.001.
7. van Kleunen M, Dawson W, Essl F, Pergl J, Winter M, Weber E, Kreft H, Weigelt P,
Kartesz J, Nishino M, Antonova LA, Barcelona JF, Cabezas FJ, Cárdenas D, Cárdenas-
Toro J, Castaño N, Chacón E, Chatelain C, Ebel AL, Figueiredo E, Fuentes N, Groom
QJ, Henderson L, Inderjit, Kupriyanov A, Masciadri S, Meerman J, Morozova O, Moser
D, Nickrent DL, Patzelt A, Pelser PB, Baptiste MP, Poopath M, Schulze M, Seebens H,
Shu W-s, Thomas J, Velayos M, Wieringa JJ, Pyšek P. Global exchange and
accumulation of non-native plants. Nature. 2015;525(7567):100-103. doi:
10.1038/nature14910.
8. Toral-Granda MV, Causton CE, Jager H, Trueman M, Izurieta JC, Araujo E, Cruz M,
Zander KK, Izurieta A, Garnett ST. Alien species pathways to the Galapagos Islands,
Ecuador. Plos One. 2017;12(9):e0184379. Epub 2017/09/14. doi:
10.1371/journal.pone.0184379. PMID: 28902860 PMCID: PMC5597199.
9. Atwood AR, Sachs JP. Separating ITCZ- and ENSO-related rainfall changes in the
Galápagos over the last 3 kyr using D/H ratios of multiple lipid biomarkers. Earth and
Planetary Science Letters. 2014;404:408-419. doi: 10.1016/j.epsl.2014.07.038.
10. Putnam AE, Broecker WS. Human-induced changes in the distribution of rainfall.
Science Advances. 2017;3(5):e1600871. doi: doi:10.1126/sciadv.1600871.

11. Percy M, Schmitt S, Riveros-Iregui D, Mirus BB. The Galapagos archipelago: A natural laboratory to examine sharp hydroclimatic, geologic and anthropogenic gradients. WIREs Water. 2016;3(4):587-600. doi: 10.1002/wat2.1145.

12. Schmitt SR, Riveros-Iregui DA, Hu J. The role of fog, orography, and seasonality on precipitation in a semiarid, tropical island. Hydrological Processes. 2018;32(18):2792-2805. doi: https://doi.org/10.1002/hyp.13228.

13. DR R, KL C. Shore fishes and biogeographic subdivisions of the Tropical Eastern Pacific. Mar Ecol Prog. 2009. doi: https://doi.org/10.3354/meps07925.

14. Graham EB, Knelman JE, Schindlbacher A, Siciliano S, Breulmann M, Yannarell A, Beman JM, Abell G, Philippot L, Prosser J, Foulquier A, Yuste JC, Glanville HC, Jones DL, Angel R, Salminen J, Newton RJ, Burgmann H, Ingram LJ, Hamer U, Siljanen HM, Peltoniemi K, Potthast K, Baneras L, Hartmann M, Banerjee S, Yu RQ, Nogaro G, Richter A, Koranda M, Castle SC, Goberna M, Song B, Chatterjee A, Nunes OC, Lopes AR, Cao Y, Kaisermann A, Hallin S, Strickland MS, Garcia-Pausas J, Barba J, Kang H, Isobe K, Papaspyrou S, Pastorelli R, Lagomarsino A, Lindstrom ES, Basiliko N, Nemergut DR. Microbes as Engines of Ecosystem Function: When Does Community Structure Enhance Predictions of Ecosystem Processes? Front Microbiol. 2016;7:214. Epub 2016/03/05. doi: 10.3389/fmicb.2016.00214. PMID: 26941732 PMCID: PMC4764795.

15. Graham EB, Knelman JE, Schindlbacher A, Siciliano S, Breulmann M, Yannarell A, Beman JM, Abell G, Philippot L, Prosser J, Foulquier A, Yuste JC, Glanville HC, Jones DL, Angel R, Salminen J, Newton RJ, Bürgmann H, Ingram LJ, Hamer U, Siljanen HMP, Peltoniemi K, Potthast K, Bañeras L, Hartmann M, Banerjee S, Yu R-Q, Nogaro G, Richter A, Koranda M, Castle SC, Goberna M, Song B, Chatterjee A, Nunes OC, Lopes AR, Cao Y, Kaisermann A, Hallin S, Strickland MS, Garcia-Pausas J, Barba J, Kang H, Isobe K, Papaspyrou S, Pastorelli R, Lagomarsino A, Lindström ES, Basiliko N, Nemergut DR. Microbes as Engines of Ecosystem Function: When Does Community Structure Enhance Predictions of Ecosystem Processes? Front Microbiol. 2016;7. doi: 10.3389/fmicb.2016.00214.

16. Gougoulias C, Clark JM, Shaw LJ. The role of soil microbes in the global carbon cycle: tracking the below-ground microbial processing of plant-derived carbon for manipulating carbon dynamics in agricultural systems. J Sci Food Agric. 2014;94(12):2362-2371. Epub 2014/01/16. doi: 10.1002/jsfa.6577. PMID: 24425529 PMCID: PMC4283042.

17. Msimbira LA, Smith DL. The Roles of Plant Growth Promoting Microbes in Enhancing Plant Tolerance to Acidity and Alkalinity Stresses. Frontiers in Sustainable Food Systems. 2020;4. doi: 10.3389/fsufs.2020.00106.

18. Naik K, Mishra S, Srichandan H, Singh PK, Sarangi PK. Plant growth promoting microbes: Potential link to sustainable agriculture and environment. Biocatalysis and Agricultural Biotechnology. 2019;21:101326. doi: https://doi.org/10.1016/j.bcab.2019.101326.

19. Qadri M, Short S, Gast K, Hernandez J, Wong AC-N. Microbiome Innovation in Agriculture: Development of Microbial Based Tools for Insect Pest Management. Frontiers in Sustainable Food Systems. 2020;4. doi: 10.3389/fsufs.2020.547751.

20. Spaepen S, Vanderleyden J, Remans R. Indole-3-acetic acid in microbial and microorganism-plant signaling. Fems Microbiol Rev. 2007;31(4):425-448. doi: 10.1111/j.1574-6976.2007.00072.x.

21. Souza R, Ambrosini A, Passaglia LM. Plant growth-promoting bacteria as inoculants in agricultural soils. Genet Mol Biol. 2015;38(4):401-419. Epub 2015/11/06. doi: 10.1590/s1415-475738420150053. PMID: 26537605 PMCID: PMC4763327.

22. Singh U, Walvekar VA, Sharma S. Microbiome as Sensitive Markers for Risk Assessment of Pesticides. Pesticides in Crop Production2020. p. 89-108.

23.   Gomiero T, Pimentel D, Paoletti MG. Is There a Need for a More Sustainable
Agriculture? Critical Reviews in Plant Sciences. 2011;30(1-2):6-23. doi:
10.1080/07352689.2011.553515.
24.   Hendgen M, Hoppe B, Döring J, Friedel M, Kauer R, Frisch M, Dahl A, Kellner H. Effects
of different management regimes on microbial biodiversity in vineyard soils. Sci Rep-uk.
2018;8(1):9393. doi: 10.1038/s41598-018-27743-0.
25.   Lauber CL, Hamady M, Knight R, Fierer N. Pyrosequencing-Based Assessment of Soil
pH as a Predictor of Soil Bacterial Community Structure at the Continental Scale. Appl
Environ Microb. 2009;75(15):5111-5120. doi: doi:10.1128/AEM.00335-09.
26.   Griffiths RI, Thomson BC, James P, Bell T, Bailey M, Whiteley AS. The bacterial
biogeography of British soils. Environ Microbiol. 2011;13(6):1642-1654. Epub
2011/04/22. doi: 10.1111/j.1462-2920.2011.02480.x. PMID: 21507180.
27.   Fierer N, Jackson RB. The diversity and biogeography of soil bacterial communities.
Proc National Acad Sci. 2006;103(3):626-631. doi: doi:10.1073/pnas.0507535103.
28.   George SF, Fierer N, Levy JS, Adams B. Antarctic Water Tracks: Microbial Community
Responses to Variation in Soil Moisture, pH, and Salinity. Front Microbiol. 2021;12. doi:
10.3389/fmicb.2021.616730.
29.   Zeglin LH, Dahm CN, Barrett JE, Gooseff MN, Fitpatrick SK, Takacs-Vesbach CD.
Bacterial community structure along moisture gradients in the parafluvial sediments of
two ephemeral desert streams. Microb Ecol. 2011;61(3):543-556. Epub 2010/12/15. doi:
10.1007/s00248-010-9782-7. PMID: 21153024.
30.   Van Horn DJ, Van Horn ML, Barrett JE, Gooseff MN, Altrichter AE, Geyer KM, Zeglin
LH, Takacs-Vesbach CD. Factors Controlling Soil Microbial Biomass and Bacterial
Diversity and Community Composition in a Cold Desert Ecosystem: Role of Geographic
Scale. Plos One. 2013;8(6):e66103. doi: 10.1371/journal.pone.0066103.
31.   Xu X, Thornton PE, Post WM. A global analysis of soil microbial biomass carbon,
nitrogen and phosphorus in terrestrial ecosystems. Global Ecology and Biogeography.
2013;22(6):737-749. doi: https://doi.org/10.1111/geb.12029.
32.   Sul WJ, Asuming-Brempong S, Wang Q, Tourlousse DM, Penton CR, Deng Y,
Rodrigues JLM, Adiku SGK, Jones JW, Zhou J, Cole JR, Tiedje JM. Tropical agricultural
land management influences on soil microbial communities through its effect on soil
organic carbon. Soil Biology and Biochemistry. 2013;65:33-38. doi:
https://doi.org/10.1016/j.soilbio.2013.05.007.
33.   Oliverio AM, Bradford MA, Fierer N. Identifying the microbial taxa that consistently
respond to soil warming across time and space. Global Change Biology.
2017;23(5):2117-2129. doi: https://doi.org/10.1111/gcb.13557.
34.   Serna-Chavez HM, Fierer N, van Bodegom PM. Global drivers and patterns of microbial
abundance in soil. Global Ecology and Biogeography. 2013;22(10):1162-1172. doi:
https://doi.org/10.1111/geb.12070.
35.   Pett-Ridge J, Firestone MK. Redox Fluctuation Structures Microbial Communities in a
Wet Tropical Soil. Appl Environ Microb. 2005;71(11):6998-7007. doi:
doi:10.1128/AEM.71.11.6998-7007.2005.
36.   Peay KG, Baraloto C, Fine PVA. Strong coupling of plant and fungal community
structure across western Amazonian rainforests. Isme J. 2013;7(9):1852-1861. doi:
10.1038/ismej.2013.66.
37.   Barberán A, McGuire KL, Wolf JA, Jones FA, Wright SJ, Turner BL, Essene A, Hubbell
SP, Faircloth BC, Fierer N. Relating belowground microbial composition to the
taxonomic, phylogenetic, and functional trait distributions of trees in a tropical forest.
Ecology Letters. 2015;18(12):1397-1405. doi: https://doi.org/10.1111/ele.12536.
38.   Prober SM, Leff JW, Bates ST, Borer ET, Firn J, Harpole WS, Lind EM, Seabloom EW,
Adler PB, Bakker JD, Cleland EE, DeCrappeo NM, DeLorenze E, Hagenah N, Hautier Y,

Hofmockel KS, Kirkman KP, Knops JMH, La Pierre KJ, MacDougall AS, McCulley RL,
Mitchell CE, Risch AC, Schuetz M, Stevens CJ, Williams RJ, Fierer N. Plant diversity
predicts beta but not alpha diversity of soil microbes across grasslands worldwide.
Ecology Letters. 2015;18(1):85-95. doi: https://doi.org/10.1111/ele.12381.
39.   Ehrenfeld JG. Effects of Exotic Plant Invasions on Soil Nutrient Cycling Processes.
Ecosystems. 2003;6(6):503-523. doi: 10.1007/s10021-002-0151-3.
40.   Broz AK, Manter DK, Vivanco JM. Soil fungal abundance and diversity: another victim of
the invasive plant Centaurea maculosa. Isme J. 2007;1(8):763-765. doi:
10.1038/ismej.2007.81.
41.   Knapp DG, Pintye A, Kovács GM. The Dark Side Is Not Fastidious – Dark Septate
Endophytic Fungi of Native and Invasive Plants of Semiarid Sandy Areas. Plos One.
2012;7(2):e32570. doi: 10.1371/journal.pone.0032570.
42.   McLeod ML, Bullington L, Cleveland CC, Rousk J, Lekberg Y. Invasive plant-derived
dissolved organic matter alters microbial communities and carbon cycling in soils. Soil
Biology and Biochemistry. 2021;156:108191. doi:
https://doi.org/10.1016/j.soilbio.2021.108191.
43.   Farrer EC, Birnbaum C, Waryszak P, Halbrook SR, Brady MV, Bumby CR, Candaele H,
Kulick NK, Lee SFH, Schroeder CS, Smith MKH, Wilber W. Plant and microbial impacts
of an invasive species vary across an environmental gradient. Journal of Ecology.
2021;109(5):2163-2176. doi: https://doi.org/10.1111/1365-2745.13629.
44.   Tian B, Pei Y, Huang W, Ding J, Siemann E. Increasing flavonoid concentrations in root
exudates enhance associations between arbuscular mycorrhizal fungi and an invasive
plant. Isme J. 2021;15(7):1919-1930. doi: 10.1038/s41396-021-00894-1.
45.   Urquía D, Gutierrez B, Pozo G, Pozo MJ, Espín A, Torres MdL. Psidium guajava in the
Galapagos Islands: Population genetics and history of an invasive species. Plos One.
2019;14(3):e0203737. doi: 10.1371/journal.pone.0203737.
46.   Bokulich NA, Kaehler BD, Rideout JR, Dillon M, Bolyen E, Knight R, Huttley GA,
Gregory Caporaso J. Optimizing taxonomic classification of marker-gene amplicon
sequences with QIIME 2's q2-feature-classifier plugin. Microbiome. 2018;6(1):90. doi:
10.1186/s40168-018-0470-z.
47.   Schmidt SK, Scow KM. Mycorrhizal Fungi on the Galapagos Islands. Biotropica.
1986;18(3):236-240. doi: 10.2307/2388491.
48.   Bever JD, Schultz PA, Pringle A, Morton JB. Arbuscular Mycorrhizal Fungi: More
Diverse than Meets the Eye, and the Ecological Tale of Why: The high diversity of
ecologically distinct species of arbuscular mycorrhizal fungi within a single community
has broad implications for plant ecology. BioScience. 2001;51(11):923-931. doi:
10.1641/0006-3568(2001)051[0923:Amfmdt]2.0.Co;2.
49.   Louca S, Parfrey LW, Doebeli M. Decoupling function and taxonomy in the global ocean
microbiome. Science. 2016;353(6305):1272-1277. doi: doi:10.1126/science.aaf4507.
50.   Nguyen NH, Song Z, Bates ST, Branco S, Tedersoo L, Menke J, Schilling JS, Kennedy
PG. FUNGuild: An open annotation tool for parsing fungal community datasets by
ecological guild. Fungal Ecology. 2016;20:241-248. doi:
https://doi.org/10.1016/j.funeco.2015.06.006.
51.   Li G, Xu G, Shen C, Tang Y, Zhang Y, Ma K. Contrasting elevational diversity patterns
for soil bacteria between two ecosystems divided by the treeline. Science China Life
Sciences. 2016;59(11):1177-1186. doi: 10.1007/s11427-016-0072-6.
52.   Peay KG, von Sperber C, Cardarelli E, Toju H, Francis CA, Chadwick OA, Vitousek PM.
Convergence and contrast in the community structure of Bacteria, Fungi and Archaea
along a tropical elevation–climate gradient. FEMS Microbiology Ecology. 2017;93(5).
doi: 10.1093/femsec/fix045.
53.   Fierer N, McCain CM, Meir P, Zimmermann MB, Rapp JM, Silman MR, Knight R.

Microbes do not follow the elevational diversity patterns of plants and animals. Ecology. 2011;92 4:797-804.

54. Singh D, Lee-Cruz L, Kim W-S, Kerfahi D, Chun J-H, Adams JM. Strong elevational trends in soil bacterial community composition on Mt. Halla, South Korea. Soil Biology and Biochemistry. 2014;68:140-149. doi: https://doi.org/10.1016/j.soilbio.2013.09.027.

55. Looby CI, Martin PH. Diversity and function of soil microbes on montane gradients: the state of knowledge in a changing world. FEMS Microbiology Ecology. 2020;96(9). doi: 10.1093/femsec/fiaa122.

56. Clague DA, Dalrymple GB. Tectonics, Geochronology, and Origin of the Hawaiian-Emperor Volcanic Chain. A Natural History of the Hawaiian Islands: Selected Readings II: University of Hawaii Press; 2021. p. 5-40.

57. Candra IN, Gerzabek MH, Ottner F, Wriessnig K, Tintner J, Schmidt G, Rechberger MV, Rampazzo N, Zehetner F. Soil development and mineral transformations along a one-million-year chronosequence on the Galápagos Islands. Soil Science Society of America Journal. 2021;85:2077-2099. doi: 10.1002/saj2.20317.

58. Candra IN, Gerzabek MH, Ottner F, Wriessnig K, Tintner J, Schmidt G, Rechberger MV, Rampazzo N, Zehetner F. Soil development and mineral transformations along a one-million-year chronosequence on the Galápagos Islands. Soil Science Society of America Journal. 2021;85(6):2077-2099. doi: https://doi.org/10.1002/saj2.20317.

59. Selmants PC, Adair KL, Litton CM, Giardina CP, Schwartz E. Increases in mean annual temperature do not alter soil bacterial community structure in tropical montane wet forests. Ecosphere. 2016;7(4):e01296. doi: https://doi.org/10.1002/ecs2.1296.

60. Borneman J, Triplett EW. Molecular microbial diversity in soils from eastern Amazonia: evidence for unusual microorganisms and microbial population shifts associated with deforestation. Appl Environ Microb. 1997;63(7):2647-2653. doi: doi:10.1128/aem.63.7.2647-2653.1997.

61. Hartman WH, Richardson CJ, Vilgalys R, Bruland GL. Environmental and anthropogenic controls over bacterial communities in wetland soils. Proc National Acad Sci. 2008;105(46):17842-17847. doi: doi:10.1073/pnas.0808254105.

62. Shen C, Xiong J, Zhang H, Feng Y, Lin X, Li X, Liang W, Chu H. Soil pH drives the spatial distribution of bacterial communities along elevation on Changbai Mountain. Soil Biology and Biochemistry. 2013;57:204-211. doi: https://doi.org/10.1016/j.soilbio.2012.07.013.

63. Wang JT, Cao P, Hu HW, Li J, Han LL, Zhang LM, Zheng YM, He JZ. Altitudinal distribution patterns of soil bacterial and archaeal communities along mt. Shegyla on the Tibetan Plateau. Microb Ecol. 2015;69(1):135-145. Epub 2014/07/31. doi: 10.1007/s00248-014-0465-7. PMID: 25074792.

64. Crits-Christoph A, Robinson CK, Barnum T, Fricke WF, Davila AF, Jedynak B, McKay CP, DiRuggiero J. Colonization patterns of soil microbial communities in the Atacama Desert. Microbiome. 2013;1(1):28. doi: 10.1186/2049-2618-1-28.

65. Neilson JW, Califf K, Cardona C, Copeland A, Treuren Wv, Josephson KL, Knight R, Gilbert JA, Quade J, Caporaso JG, Maier RM, Chu H. Significant Impacts of Increasing Aridity on the Arid Soil Microbiome. Msystems. 2017;2(3):e00195-00116. doi: doi:10.1128/mSystems.00195-16.

66. Angel R, Soares MIM, Ungar ED, Gillor O. Biogeography of soil archaea and bacteria along a steep precipitation gradient. Isme J. 2010;4(4):553-563. doi: 10.1038/ismej.2009.136.

67. Bachar A, Al-Ashhab A, Soares MI, Sklarz MY, Angel R, Ungar ED, Gillor O. Soil microbial abundance and diversity along a low precipitation gradient. Microb Ecol. 2010;60(2):453-461. Epub 2010/08/05. doi: 10.1007/s00248-010-9727-1. PMID: 20683588.

68. Chowdhury TRo, Mitsch WJ, Dick RP. Seasonal methanotrophy across a hydrological
gradient in a freshwater wetland. Ecol Eng. 2014;72:116-124. doi:
10.1016/j.ecoleng.2014.08.015.
69. Balasooriya WK, Denef K, Peters J, Verhoest NEC, Boeckx P. Vegetation composition
and soil microbial community structural changes along a wetland hydrological gradient.
Hydrol Earth Syst Sci. 2008;12(1):277-291. doi: 10.5194/hess-12-277-2008.
70. Lee KC, Caruso T, Archer SDJ, Gillman LN, Lau MCY, Cary SC, Lee CK, Pointing SB.
Stochastic and Deterministic Effects of a Moisture Gradient on Soil Microbial
Communities in the McMurdo Dry Valleys of Antarctica. Front Microbiol. 2018;9. doi:
10.3389/fmicb.2018.02619.
71. Couradeau E, Sasse J, Goudeau D, Nath N, Hazen TC, Bowen BP, Chakraborty R,
Malmstrom RR, Northen TR. Probing the active fraction of soil microbiomes using
BONCAT-FACS. Nat Commun. 2019;10(1):2770. doi: 10.1038/s41467-019-10542-0.
72. Haichar FEZ, Heulin T, Guyonnet JP, Achouak W. Stable isotope probing of carbon flow
in the plant holobiont. Curr Opin Biotechnol. 2016;41:9-13. Epub 2016/03/29. doi:
10.1016/j.copbio.2016.02.023. PMID: 27019410.
73. Van Der Heijden MGA, Bardgett RD, Van Straalen NM. The unseen majority: soil
microbes as drivers of plant diversity and productivity in terrestrial ecosystems. Ecology
Letters. 2008;11(3):296-310. doi: https://doi.org/10.1111/j.1461-0248.2007.01139.x.
74. Schirawski J, Perlin MH. Plant–Microbe Interaction 2017—The Good, the Bad and the
Diverse. International Journal of Molecular Sciences. 2018;19(5):1374. PMID:
doi:10.3390/ijms19051374.
75. Backer R, Rokem JS, Ilangumaran G, Lamont J, Praslickova D, Ricci E, Subramanian S,
Smith DL. Plant Growth-Promoting Rhizobacteria: Context, Mechanisms of Action, and
Roadmap to Commercialization of Biostimulants for Sustainable Agriculture. Frontiers in
Plant Science. 2018;9. doi: 10.3389/fpls.2018.01473.
76. Biological Nitrogen Fixation. Biological Nitrogen Fixation. Frans J. de Bruijn ed: John
Wiley & Sons, Inc 2015. p. i-xxviii.
77. Hough M, McClure A, Bolduc B, Dorrepaal E, Saleska S, Klepac-Ceraj V, Rich V. Biotic
and Environmental Drivers of Plant Microbiomes Across a Permafrost Thaw Gradient.
Front Microbiol. 2020;11. doi: 10.3389/fmicb.2020.00796.
78. Lekberg Y, Waller LP. What drives differences in arbuscular mycorrhizal fungal
communities among plant species? Fungal Ecology. 2016;24:135-138. doi:
https://doi.org/10.1016/j.funeco.2016.05.012.
79. Nunan N, Daniell TJ, Singh BK, Papert A, McNicol JW, Prosser JI. Links between Plant
and Rhizoplane Bacterial Communities in Grassland Soils, Characterized Using
Molecular Techniques. Appl Environ Microb. 2005;71(11):6784-6792. doi:
doi:10.1128/AEM.71.11.6784-6792.2005.
80. Schlatter DC, Bakker MG, Bradeen JM, Kinkel LL. Plant community richness and
microbial interactions structure bacterial communities in soil. Ecology. 2015;96(1):134-
142. Epub 2015/08/04. doi: 10.1890/13-1648.1. PMID: 26236898.
81. Peay KG, Russo SE, McGuire KL, Lim Z, Chan JP, Tan S, Davies SJ. Lack of host
specificity leads to independent assortment of dipterocarps and ectomycorrhizal fungi
across a soil fertility gradient. Ecology Letters. 2015;18(8):807-816. doi:
https://doi.org/10.1111/ele.12459.
82. Zimmerman NB, Vitousek PM. Fungal endophyte communities reflect environmental
structuring across a Hawaiian landscape. Proc National Acad Sci. 2012;109(32):13022-
13027. doi: doi:10.1073/pnas.1209872109.
83. Hu L, Robert CAM, Cadot S, Zhang X, Ye M, Li B, Manzo D, Chervet N, Steinger T, van
der Heijden MGA, Schlaeppi K, Erb M. Root exudate metabolites drive plant-soil
feedbacks on growth and defense by shaping the rhizosphere microbiota. Nat Commun.

1153   2018;9(1):2738. doi: 10.1038/s41467-018-05122-7.

1154 84.   Fierer N, Leff JW, Adams BJ, Nielsen UN, Bates ST, Lauber CL, Owens S, Gilbert JA,
1155   Wall DH, Caporaso JG. Cross-biome metagenomic analyses of soil microbial
1156   communities and their functional attributes. Proc National Acad Sci.
1157   2012;109(52):21390-21395. doi: doi:10.1073/pnas.1215210110.

1158 85.   Janssen PH. Identifying the Dominant Soil Bacterial Taxa in Libraries of 16S rRNA and
1159   16S rRNA Genes. Appl Environ Microb. 2006;72(3):1719-1728. doi:
1160   doi:10.1128/AEM.72.3.1719-1728.2006.

1161 86.   Harris RF. Effect of Water Potential on Microbial Growth and Activity. Water Potential
1162   Relations in Soil Microbiology1981. p. 23-95.

1163 87.   Tripathi BM, Moroenyane I, Sherman C, Lee YK, Adams JM, Steinberger Y. Trends in
1164   Taxonomic and Functional Composition of Soil Microbiome Along a Precipitation
1165   Gradient in Israel. Microb Ecol. 2017;74(1):168-176. Epub 2017/01/12. doi:
1166   10.1007/s00248-017-0931-0. PMID: 28074247.

1167 88.   Naidoo Y, Valverde A, Pierneef RE, Cowan DA. Differences in Precipitation Regime
1168   Shape Microbial Community Composition and Functional Potential in Namib Desert
1169   Soils. Microb Ecol. 2021. Epub 2021/06/10. doi: 10.1007/s00248-021-01785-w. PMID:
1170   34105010.

1171 89.   Gargouri M, Karray F, Chebaane A, Mhiri N, Partida-Martínez LP, Sayadi S, Mliki A.
1172   Increasing aridity shapes beta diversity and the network dynamics of the belowground
1173   fungal microbiome associated with Opuntia ficus-indica. Science of The Total
1174   Environment. 2021;773:145008. doi: https://doi.org/10.1016/j.scitotenv.2021.145008.

1175 90.   Maestre FT, Delgado-Baquerizo M, Jeffries TC, Eldridge DJ, Ochoa V, Gozalo B, Quero
1176   JL, García-Gómez M, Gallardo A, Ulrich W, Bowker MA, Arredondo T, Barraza-Zepeda
1177   C, Bran D, Florentino A, Gaitán J, Gutiérrez JR, Huber-Sannwald E, Jankju M, Mau RL,
1178   Miriti M, Naseri K, Ospina A, Stavi I, Wang D, Woods NN, Yuan X, Zaady E, Singh BK.
1179   Increasing aridity reduces soil microbial diversity and abundance in global drylands. Proc
1180   National Acad Sci. 2015;112(51):15684-15689. doi: doi:10.1073/pnas.1516684112.

1181 91.   Sun S, Li S, Avera BN, Strahm BD, Badgley BD, Löffler FE. Soil Bacterial and Fungal
1182   Communities Show Distinct Recovery Patterns during Forest Ecosystem Restoration.
1183   Appl Environ Microb. 2017;83(14):e00966-00917. doi: doi:10.1128/AEM.00966-17.

1184 92.   Cloutier ML, Murrell E, Barbercheck M, Kaye J, Finney D, García-González I, Bruns MA.
1185   Fungal community shifts in soils with varied cover crop treatments and edaphic
1186   properties. Sci Rep-uk. 2020;10(1):6198. doi: 10.1038/s41598-020-63173-7.

1187 93.   Smith ME, Douhan GW, Rizzo DM. Ectomycorrhizal community structure in a xeric
1188   Quercus woodland based on rDNA sequence analysis of sporocarps and pooled roots.
1189   New Phytol. 2007;174(4):847-863. Epub 2007/05/17. doi: 10.1111/j.1469-
1190   8137.2007.02040.x. PMID: 17504467.

1191 94.   Lanfranco L, Bonfante P, Genre A. The Mutualistic Interaction between Plants and
1192   Arbuscular Mycorrhizal Fungi. Microbiol Spectr. 2016;4(6). Epub 2017/01/15. doi:
1193   10.1128/microbiolspec.FUNK-0012-2016. PMID: 28087942.

1194 95.   Taylor TN, Krings M, Taylor EL. 7 - Glomeromycota. In: Taylor TN, Krings M, Taylor EL,
1195   editors. Fossil Fungi. San Diego: Academic Press; 2015. p. 103-128.

1196 96.   Birhane E, Sterck FJ, Fetene M, Bongers F, Kuyper TW. Arbuscular mycorrhizal fungi
1197   enhance photosynthesis, water use efficiency, and growth of frankincense seedlings
1198   under pulsed water availability conditions. Oecologia. 2012;169(4):895-904. Epub
1199   2012/01/31. doi: 10.1007/s00442-012-2258-3. PMID: 22286084 PMCID: PMC3398253.

1200 97.   Rouphael Y, Franken P, Schneider C, Schwarz D, Giovannetti M, Agnolucci M, Pascale
1201   SD, Bonini P, Colla G. Arbuscular mycorrhizal fungi act as biostimulants in horticultural
1202   crops. Scientia Horticulturae. 2015;196:91-108. doi:
1203   https://doi.org/10.1016/j.scienta.2015.09.002.

1204 98. Evelin H, Giri B, Kapoor R. Contribution of Glomus intraradices inoculation to nutrient
1205  acquisition and mitigation of ionic imbalance in NaCl-stressed Trigonella foenum-
1206  graecum. Mycorrhiza. 2012;22(3):203-217. Epub 2011/06/23. doi: 10.1007/s00572-011-
1207  0392-0. PMID: 21695577.
1208 99. Bowles TM, Barrios-Masias FH, Carlisle EA, Cavagnaro TR, Jackson LE. Effects of
1209  arbuscular mycorrhizae on tomato yield, nutrient uptake, water relations, and soil carbon
1210  dynamics under deficit irrigation in field conditions. Science of The Total Environment.
1211  2016;566-567:1223-1234. doi: https://doi.org/10.1016/j.scitotenv.2016.05.178.
1212 100. Miozzi L, Vaira AM, Catoni M, Fiorilli V, Accotto GP, Lanfranco L. Arbuscular Mycorrhizal
1213  Symbiosis: Plant Friend or Foe in the Fight Against Viruses? Front Microbiol. 2019;10.
1214  doi: 10.3389/fmicb.2019.01238.
1215 101. Abdel-Salam E, Alatar A, El-Sheikh MA. Inoculation with arbuscular mycorrhizal fungi
1216  alleviates harmful effects of drought stress on damask rose. Saudi J Biol Sci.
1217  2018;25(8):1772-1780. Epub 2018/12/29. doi: 10.1016/j.sjbs.2017.10.015. PMID:
1218  30591799 PMCID: PMC6303177.
1219 102. Begum N, Qin C, Ahanger MA, Raza S, Khan MI, Ashraf M, Ahmed N, Zhang L. Role of
1220  Arbuscular Mycorrhizal Fungi in Plant Growth Regulation: Implications in Abiotic Stress
1221  Tolerance. Frontiers in Plant Science. 2019;10. doi: 10.3389/fpls.2019.01068.
1222 103. Maherali H, Oberle B, Stevens PF, Cornwell WK, McGlinn DJ. Mutualism Persistence
1223  and Abandonment during the Evolution of the Mycorrhizal Symbiosis. Am Nat.
1224  2016;188(5):E113-E125. doi: 10.1086/688675. PMID: 27788343.
1225 104. Duchicela J, Bever JD, Schultz PA. Symbionts as Filters of Plant Colonization of Islands:
1226  Tests of Expected Patterns and Environmental Consequences in the Galapagos. Plants.
1227  2020;9(1):74. PMID: doi:10.3390/plants9010074.
1228 105. Thirkell TJ, Charters MD, Elliott AJ, Sait SM, Field KJ. Are mycorrhizal fungi our
1229  sustainable saviours? Considerations for achieving food security. Journal of Ecology.
1230  2017;105(4):921-929. doi: https://doi.org/10.1111/1365-2745.12788.
1231 106. Bender SF, Wagg C, van der Heijden MGA. An Underground Revolution: Biodiversity
1232  and Soil Ecological Engineering for Agricultural Sustainability. Trends in Ecology &
1233  Evolution. 2016;31(6):440-452. doi: https://doi.org/10.1016/j.tree.2016.02.016.
1234 107. Sosa-Hernández MA, Leifheit EF, Ingraffia R, Rillig MC. Subsoil Arbuscular Mycorrhizal
1235  Fungi for Sustainability and Climate-Smart Agriculture: A Solution Right Under Our
1236  Feet? Front Microbiol. 2019;10. doi: 10.3389/fmicb.2019.00744.
1237 108. Louca S, Polz MF, Mazel F, Albright MBN, Huber JA, O'Connor MI, Ackermann M, Hahn
1238  AS, Srivastava DS, Crowe SA, Doebeli M, Parfrey LW. Function and functional
1239  redundancy in microbial systems. Nature Ecology & Evolution. 2018;2(6):936-943. doi:
1240  10.1038/s41559-018-0519-1.
1241 109. Jurburg SD, Salles JF. Functional Redundancy and Ecosystem Function — The Soil
1242  Microbiota as a Case Study In: Lo Y, Blanco JA, Roy S, editors. Biodiversity in
1243  Ecosystems - Linking Structure and Function: IntechOpen; 2015.
1244 110. Awasthi A, Singh M, Soni SK, Singh R, Kalra A. Biodiversity acts as insurance of
1245  productivity of bacterial communities under abiotic perturbations. Isme J.
1246  2014;8(12):2445-2452. doi: 10.1038/ismej.2014.91.
1247 111. Falcão Salles J, Le Roux X, Poly F. Relating Phylogenetic and Functional Diversity
1248  among Denitrifiers and Quantifying their Capacity to Predict Community Functioning.
1249  Front Microbiol. 2012;3. doi: 10.3389/fmicb.2012.00209.
1250 112. Paltán HA, Benitez FL, Rosero P, Escobar-Camacho D, Cuesta F, Mena CF. Climate
1251  and sea surface trends in the Galapagos Islands. Sci Rep-uk. 2021;11(1):14465. doi:
1252  10.1038/s41598-021-93870-w.
1253 113. Caporaso JG, Lauber CL, Walters WA, Berg-Lyons D, Lozupone CA, Turnbaugh PJ,
1254  Fierer N, Knight R. Global patterns of 16S rRNA diversity at a depth of millions of

sequences per sample. Proc National Acad Sci. 2011;108(supplement_1):4516-4522.
doi: doi:10.1073/pnas.1000080107.

114.   McIver LJ, Abu-Ali G, Franzosa EA, Schwager R, Morgan XC, Waldron L, Segata N,
Huttenhower C. bioBakery: a meta'omic analysis environment. Bioinformatics.
2018;34(7):1235-1237. Epub 2017/12/02. doi: 10.1093/bioinformatics/btx754. PMID:
29194469 PMCID: PMC6030947.

115.   Beghini F, McIver LJ, Blanco-Míguez A, Dubois L, Asnicar F, Maharjan S, Mailyan A,
Manghi P, Scholz M, Thomas AM, Valles-Colomer M, Weingart G, Zhang Y, Zolfo M,
Huttenhower C, Franzosa EA, Segata N. Integrating taxonomic, functional, and strain-
level profiling of diverse microbial communities with bioBakery 3. Elife. 2021;10:e65088.
doi: 10.7554/eLife.65088.

116.   Bolyen E, Rideout JR, Dillon MR, Bokulich NA, Abnet CC, Al-Ghalith GA, Alexander H,
Alm EJ, Arumugam M, Asnicar F, Bai Y, Bisanz JE, Bittinger K, Brejnrod A, Brislawn CJ,
Brown CT, Callahan BJ, Caraballo-Rodríguez AM, Chase J, Cope EK, Da Silva R,
Diener C, Dorrestein PC, Douglas GM, Durall DM, Duvallet C, Edwardson CF, Ernst M,
Estaki M, Fouquier J, Gauglitz JM, Gibbons SM, Gibson DL, Gonzalez A, Gorlick K, Guo
J, Hillmann B, Holmes S, Holste H, Huttenhower C, Huttley GA, Janssen S, Jarmusch
AK, Jiang L, Kaehler BD, Kang KB, Keefe CR, Keim P, Kelley ST, Knights D, Koester I,
Kosciolek T, Kreps J, Langille MGI, Lee J, Ley R, Liu Y-X, Loftfield E, Lozupone C,
Maher M, Marotz C, Martin BD, McDonald D, McIver LJ, Melnik AV, Metcalf JL, Morgan
SC, Morton JT, Naimey AT, Navas-Molina JA, Nothias LF, Orchanian SB, Pearson T,
Peoples SL, Petras D, Preuss ML, Pruesse E, Rasmussen LB, Rivers A, Robeson MS,
Rosenthal P, Segata N, Shaffer M, Shiffer A, Sinha R, Song SJ, Spear JR, Swafford AD,
Thompson LR, Torres PJ, Trinh P, Tripathi A, Turnbaugh PJ, Ul-Hasan S, van der Hooft
JJJ, Vargas F, Vázquez-Baeza Y, Vogtmann E, von Hippel M, Walters W, Wan Y, Wang
M, Warren J, Weber KC, Williamson CHD, Willis AD, Xu ZZ, Zaneveld JR, Zhang Y, Zhu
Q, Knight R, Caporaso JG. Reproducible, interactive, scalable and extensible
microbiome data science using QIIME 2. Nature Biotechnology. 2019;37(8):852-857.
doi: 10.1038/s41587-019-0209-9.

117.   Comeau AM, Douglas GM, Langille MGI, Eisen J. Microbiome Helper: a Custom and
Streamlined Workflow for Microbiome Research. Msystems. 2017;2(1):e00127-00116.
doi: doi:10.1128/mSystems.00127-16.

118.   Martin M. Cutadapt removes adapter sequences from high-throughput sequencing
reads. 2011. 2011;17(1):3. Epub 2011-08-02. doi: 10.14806/ej.17.1.200.

119.   Callahan BJ, McMurdie PJ, Rosen MJ, Han AW, Johnson AJA, Holmes SP. DADA2:
High-resolution sample inference from Illumina amplicon data. Nat Methods.
2016;13(7):581-583. doi: 10.1038/nmeth.3869.

120.   Price MN, Dehal PS, Arkin AP. FastTree 2 – Approximately Maximum-Likelihood Trees
for Large Alignments. Plos One. 2010;5(3):e9490. doi: 10.1371/journal.pone.0009490.

121.   Wickham H. ggplot2: Elegant Graphics for Data Analysis: Springer International
Publishing; 2016.

122.   Shannon CE. A Mathematical Theory of Communication. Bell System Technical Journal.
1948;27(3):379-423. doi: https://doi.org/10.1002/j.1538-7305.1948.tb01338.x.

123.   Anderson MJ. Permutational Multivariate Analysis of Variance (PERMANOVA). Wiley
StatsRef: Statistics Reference Online. p. 1-15.

124.   Oksanen J, Blanchet GF, Kindt R, Legendre R, Minchin P, O'Hara RB, Simpson G,
Solymos P, Stevens H, Szöcs E. Vegan: Community ecology package ver2012.

125.   Rodrigues RR, Rodgers NC, Wu X, Williams MA. COREMIC: a web-tool to search for a
niche associated CORE MICrobiome. PeerJ. 2018;6:e4395. doi: 10.7717/peerj.4395.

126.   Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for
RNA-seq data with DESeq2. Genome Biology. 2014;15(12):550. doi: 10.1186/s13059-

014-0550-8.
127.   Oliveira FS, Brestelli J, Cade S, Zheng J, Iodice J, Fischer S, Aurrecoechea C, Kissinger
JC, Brunk BP, Stoeckert CJ, Jr., Fernandes GR, Roos DS, Beiting DP. MicrobiomeDB: a
systems biology platform for integrating, mining and analyzing microbiome experiments.
Nucleic Acids Res. 2018;46(D1):D684-d691. Epub 2017/11/07. doi:
10.1093/nar/gkx1027. PMID: 29106667 PMCID: PMC5753346.
128.   Kirsten WJ. Automatic methods for the simultaneous determination of carbon, hydrogen,
nitrogen, and sulfur, and for sulfur alone in organic and inorganic materials. Analytical
Chemistry. 1979;51(8):1173-1179. doi: 10.1021/ac50044a019.
1315