The RNA content of extracellular vesicles from *PRPF31*[+/−] hiPSC-RPE show potential as

biomarkers of retinal degeneration

Heran Getachew[1], Sudeep Mehrotra[1], Tarandeep Kaur[1], Rosario Fernandez-Godino[1], and Eric A.

Pierce[1*], Marcela Garita-Hernandez[1*]

[1]Ocular Genomics Institute, Department of Ophthalmology, Mass Eye and Ear, Harvard Medical

School, Boston, MA

*Address correspondence to:

Eric A. Pierce Eric_pierce@meei.harvard.edu or Marcela Garita-Hernandez

mgarita@meei.harvard.edu

Ocular Genomics Institute

Mass Eye and Ear

243 Charles St.

Boston, MA 02114

Short title: Cell-free RNA biomarker of *PRPF31*[+/−] RPE degeneration

Abstract

Retinitis pigmentosa (RP), is the most common inherited retinal degeneration (IRD),

leading to vision loss via dysfunction and death of photoreceptor cells and retinal pigment

epithelium (RPE). Mutations in the pre-mRNA processing factor 31 (*PRPF31*) gene are

associated with autosomal dominant RP, impairing RPE function. While adeno-associated virus

(AAV)-mediated gene therapy shows promise for treating IRDs, the slow progression of these

diseases often makes timely measurement of clinical efficacy challenging. Extracellular vesicles

(EVs) are lipid enclosed vesicles secreted by cells and their RNA contents are being explored as

circulating biomarkers of cancer and other diseases. We hypothesize that EV RNAs could serve

as biomarkers of the health status of the neural retina and RPE health. To test this, we used

PRPF31$^{+/+}$ and *PRPF31$^{+/-}$* human-induced pluripotent stem cell (hiPSC)-derived RPE (hi-RPE)

to investigate the RNAs contained in RPE-derived EVs, and how they change in disease. We

also compared the RNA contents of RPE-EVs with the RNAs contained in the hi-RPE cells

themselves. We found that EVs from mutant *PRPF31$^{+/-}$* hi-RPE cells have distinct RNA profiles

compared to those from control cells, suggesting EV RNA contents change during disease and

could serve as biomarkers for retinal degeneration.

Keywords:

PRPF31, Retinitis pigmentosa, HiPSC-derived RPE, disease modeling, Extracellular Vesicles,

RNA-seq, biomarkers

Introduction

Inherited retinal degenerations (IRDs) are significant contributors to vision loss in both

children and adults [1,2]. Among these, rod-cone degeneration, commonly known as retinitis

pigmentosa (RP), is the most prevalent form of IRD, affecting roughly 1 in 2,500 individuals

globally [3]. The pathology of IRD and RP includes degeneration of the retinal pigment epithelium

(RPE) and/or neural retina, which lead to vision loss [4]. Mutations in pre-RNA processing factors

(PRPFs) are among the primary causes of autosomal dominant RP [5,6], with mutations in *PRPF31*

being the most frequent, accounting for approximately 10% of cases [7]. Notably, mutations in

PRPF31 lead to dominant disease through haploinsufficiency [8-12].

Research using mouse models of *PRPF31*-associated RP have shown that

haploinsufficiency of *PRPF31* causes cell autonomous defects in RPE function, including

reduced phagocytosis of photoreceptor outer segments, which is crucial for RPE and retinal

health [13-17]. Similarly, $PRPF31^{+/-}$ mutant human induced pluripotent stem cell (hiPSC) derived

RPE (hiPSC-RPE) cells exhibit reduced phagocytosis function, shorter cilia, and reduced barrier

function of the RPE [16-18]. Gene augmentation therapies have shown promise in addressing the

consequences of haploinsufficiency of *PRPF31* gene [17,19]. Adeno-associated virus (AAV)

mediated gene augmentation therapy was able to restore normal function to $PRPF31^{+/-}$ hiPSC-

RPE cells, suggesting its potential use to treat *PRPF31*-associated RP [17,18]. Gene augmentation

has also been shown to prevent retinal degeneration in a somatic mouse model of *PRPF31*

deficiency [20].

Genetic therapies including AAV – mediated gene therapy have shown great promise for

the treatment of additional genetic forms of IRD [21-25]. Given the slow progression in many IRDs,

central vision can be retained until late stages of disease making challenging to demonstrate the

benefit of treatment during clinical trials. It can take years for improvement in clinical outcome

measures such as visual acuity and visual fields to become detectable [26]. Therefore, surrogate

markers of retinal health could greatly enhance the monitoring of IRD patients during trials. One

promising type of biomarker is RNAs contained within extracellular vesicles (EVs). EVs are

cell-derived membranous structures (40–1,000 nm in diameter) [27,28] that carry a diverse cargo

including lipids, proteins, RNA and DNA [29,30]. Present both locally and systemically, EVs are

believed to facilitate intercellular communication by transferring their protein and RNA between

cells [31-33]. EVs are secreted from nearly all cell types and are found in various biofluids,

including plasma, urine, and the vitreous and aqueous humors of the eye [34]. Their potential as

biomarkers is supported by their ability to mirror the health status of their source cells [29,35-39].

Liquid biopsies of EV contents are proving effective in monitoring tumor burden and

minimal residual disease across various cancers [40]. Early research suggests that the contents of

retina and RPE-derived EVs may reflect retinal health status [41,42], although systematic studies of

RNA contents in these EVs are still lacking. Several studies have explored the potential of RNA

in EVs from the vitreous, aqueous humor, and serum as biomarkers for conditions such as uveal

melanoma, age-related macular degeneration, and glaucoma [43-47]. Additionally, the protein

content of EVs produced by the retinal pigment epithelium (RPE) has also been investigated [41,48].

Research involving RPE cells derived from swine primary cultures and retinal organoids has

demonstrated their ability to release EVs [48]. Notably, proteomic analyses of these EVs have

linked their content to age-related macular degeneration [49]. However, their precise function in

retinal pathology remains largely unexplored.

In order to test the hypothesis that EV RNAs can be used as biomarkers for *PRPF31*-

associated RP as well as other IRDs affecting the RPE, it is crucial to define the RNA profiles of

RPE-derived EVs. In this study, we characterized the RNA profiles of EVs from control

PRPF31$^{+/+}$ and mutant *PRPF31$^{+/-}$* hiPSC-RPE cells. We also compared these profiles with the

RNA transcriptomes of the originating RPE cells. Our findings reveal that the RNA cargo in EVs

mirrors that of the originating RPE cells, with distinct differences reflecting disease perturbation

and regulatory changes in genes associated with RPE de-differentiation and function. These

results support the potential of RNA biomarkers in assessing RPE health status.

Materials and Methods

<u>Generation of hiPSC-derived RPE cell cultures</u>

Wildtype *PRPF31* and mutant *PRPF31$^{+/-}$* hiPSC were previously engineered using CRISPR

Cas9 genome editing [17,50]. These hiPSC were maintained on growth-factor-reduced Matrigel (BD

Biosciences) coated plates and their pluripotency state was confirmed by expression of

pluripotency markers OCT4, NANOG and SSEA4 (**Supplementary Figure 1**). The

differentiation capacity of the lines was assessed with the spontaneous differentiation Embryoid

Body (EB) model, which demonstrated *PRPF31$^{+/-}$* and its isogenic control hiPSC line were

capable of differentiating into the three germ layers: Ectoderm, mesoderm and endoderm

(**Supplementary Figure 1**). For the directed differentiation towards RPE, *PRPF31$^{+/+}$* and

mutant *PRPF31$^{+/-}$* hiPSC were left in culture until an approximate 90% confluence was reached

(Day 0) and differentiated as previously described [17]. Briefly, hiPSC were treated with a

combination of Noggin, Dickkopf-1 (Dkk-1), Insulin Growth Factor 1, Nicotinamide, Activin A,

basic Fibroblast Growth Factor, SU-5402, and CHIR99021 to induce the directed differentiation

of hiPSCs into RPE cells. On day 30, cells were enzymatically digested with TrypLE^{TM} Express

(12604013, Thermo Fisher, Waltham, MA, USA), strained through a 40-mm filter, and seeded at

a density of $1X10^5$ cells/cm^2 onto Matrigel-coated 6-well plates in X-VIVO 10 media. After 60

days in culture, hi-RPE cells were seeded onto non-growth factor reduced Matrigel-coated 24-

mm Transwell® with 0.4-μm pore polyester membrane inserts (3450, Corning, Bedford, MA,

USA) at a density of 700,500 cells per well. The cells were matured for 4 weeks on Transwells®

before conditioned media was collected (**Figure 1**). Apical and basal conditioned media was

collected every other day for 12 weeks and stored at -80°C.

EV isolation

The conditioned media from serum-free RPE cell cultures was thawed on ice and centrifuged

$3000 \times g$ for 15 minutes to remove cells and cell debris. EVs were isolated using ExoQuick-TC

(EXOTC50A-1, System Biosciences, Palo Alto, CA, USA) according to manufacturer

instructions. Briefly, the conditioned was mixed with ExoQuick-TC and incubated at 4°C for at

least 12 hours. Then, ExoQuick-TC/conditioned media mixture was centrifuged at $1500 \times g$ for

30 minutes and the supernatant was discarded. A final spin was performed at $1500 \times g$ for 5

minutes to remove any residual fluid. The EV pellet was resuspended in 100 μL of 1X particle

free phosphate-buffered saline (PBS) and stored at -80°C (**Figure 1A**). The isolated EVs,

ranging in size from 40 to 1000 nm, were characterized as EVs using EM imaging. Since this

study concentrates on the RNA content as potential biomarkers of the isolated EVs, and because

non-vesicular extracellular particles lack RNA, no further separation was deemed necessary in

accordance with the latest Minimal Information for Studies of Extracellular Vesicles

(MISEV) recommendations [51].

Nanoparticle tracking analysis (NTA)

The size distribution and concentration of EVs was determined using a NanoSight NS300

instrument (Malvern Instruments Ltd, Malvern, UK) as previously described (Mass SL 2015).

The EV suspensions were diluted 1:100 in PBS, and 1 mL was used for NTA. Size and diffusion

of nanoparticles was tracked, recorded, and analyzed using NTA software 3.2. Detection

threshold was set for each sample individually in a way to meet manufacturer recommended

quality standards.

<u>Transmission Electron Microscopy TEM)</u>

EV morphologies were visualized with transmission electron microscopy (TEM) following a

procedure adapted from Théry et al [52]. Briefly, the EVs were lyophilized and fixed for 1 hour at

room temperature in 2% paraformaldehyde in 0.1 mol/L sodium phosphate buffer (Electron

Microscopy Sciences, Hatfield, Pa). Five microliters of the exosome were absorbed onto

Formvar/carbon-coated electron microscopy grids (Electron Microscopy Sciences) for 20

minutes. The grids were rinsed 8 times in 0.1 mol/L sodium phosphate buffer and then incubated

in 1% glutaraldehyde in 0.1 mol/L sodium phosphate buffer (Electron Microscopy Services) for

5 minutes. After rinsing 8 times in deionized water, grids were contrasted in uranyl-oxalate

solution (pH 7; Electron Microscopy Services) for 5 minutes and transferred to uranyl acetate-

methyl cellulose solution for 5 minutes. The grids were blotted on filter paper and air-dried

before imaging. The EVs were observed with an FEI Tecnai G2 Spirit transmission electron

microscope (FEI, Hillsboro, Ore) at an accelerating voltage of 100 kV interfaced with an AMT

XR41 digital CCD camera (Advanced Microscopy Techniques, Woburn, Mass) for digital TIFF

file image acquisition.

<u>RNA extraction and purification</u>

Total RNA was isolated from 200µl of EV using Seramir Exosome RNA amplification kit

following manufacturer's instructions. Purified RNA was quantified using Agilent RNA 6000

Pico kit in a Bioanalyzer.

<u>miRNA Sequencing</u>

RNA was isolated from the EVs using the ExoQuick RNA column purification kit (EQ808A-1,

System Biosciences, Palo Alto, CA, USA). Purified RNA was quantified using a Bioanalyzer

(Agilent 2100 Bioanalyzer) and the Agilent RNA 6000 Pico kit (5067-1513, Agilent, Santa

Clara, CA, USA). miRNA libraries were prepared from the isolated EV RNA and a water control

using the TruSeq Small RNA Library Preparation Kit (RS-200-0012, Illumina, San Diego, CA,

USA) using the manufacturer instructions with the modification of performing the adaptor

ligation in the presence of 15% polyethylene glycol in order to reduce bias by driving the ligation

reactions towards completion as determined by the Extracellular RNA Communication

Consortium[53] . Adapter dimers were removed by gel purification and the purified cDNA libraries

were pooled and normalized 2nM. Nuclease free water was added to the samples as a control

during sequencing. The pooled libraries were then sequenced single-read, 50bp (1×50) on an

Illumina MiSeq using version 3 reagents at the Ocular Genomic Institute, MEE, Boston, MA

(USA).

<u>miRNA Sequencing Analysis</u>

Base quality per sequence cycle was checked with FastQC. An average minimum mean base

quality above Q30 (Phred scale) was observed using MultiQC across the read length for all

samples. The sample replicates were sequenced across four different sequencing runs. Each

sequenced run was independently quality checked. The files were merged and quality checked

again for any biases. Illumina TruSeq small RNA adapters were scanned on the 5` and 3` ends

and trimmed using Cutadapt(v.3.4). Post trimming, minimum length of 17 bp was maintained.

On average 1.5 million reads were observed across all the samples. Samples less then 1 million

reads were dropped. Following normalization using edgeR (v3.30), multidimensional scaling

(MDS) plot with biological coefficient of variations (BCV) were further used to identify outlier

samples and were removed from all downstream analysis. The standard miRDeep2 (v2.0.1.2)

was used for the analysis. Briefly, mapper module (mapper.pl) was used for read alignment

against the reference human genome (GRCh38). Identification and quantification of each

miRNA was performed by the quantifier module (quantifier.pl). Precursor and mature miRNAs

sequences of human miRBase (miRNA repository) was used as reference sequence. For novel

miRNA prediction, mature miRNA of closely related species (*Gorilla gorilla, Pongo pygmaeus,*

Pan troglodytes and Pan paniscus) were pooled together for alignment and novel miRNA

discoveries. The counts of miRNA identified and reported in the water sample were used to

penalize the identified miRNA in other sample replicates. Samples with alignment rate of <1%

were further removed. The Trimmed Mean of the M-values (TMM) from the edgeR package was

used for normalized within and across sample replicates. Post normalization, counts per million

(CPM) values were to filter and identify "expressed" miRNA across samples replicates. The

standard settings in the edgeR package used for differential analysis. Gene set enrichment

analysis; Gene Ontology (GO) pathways (KEGG) and diseases was performed using miEAA a

web-based application which performs rich functionality in terms of miRNA categories and

determines if a category is significantly enriched in a given miRNA set with respect to a

reference[54,55].

<u>EV Poly-A RNA Sequencing</u>

cDNA was prepared from EV RNA using the Single Cell RNA kit SMART- *Seq V4* Ultra Low

Input RNA Kit (634890, Takara, Mountain View, CA, USA) according to manufacturer

instructions using 1ng input. Next, poly-A libraries were prepared using the SMARTer

ThruPLEX DNA-seq Kit (R400675, Takara, Mountain View, CA, USA). Each library was

normalized to 1.5 nM, pooled and 1% PhiX was added. Libraries were sequenced for 101 cycles

(2 × 101bp (PE)) using the NovaSeq 6000 (Illumina) at the Ocular Genomic Institute, MEE,

Boston, MA (USA).

RPE poly-A RNA Sequencing

RNA was extracted from hiPSC-RPE cells and Spike-in RNA standards, sequins, were used as a

reference in all samples [56]. Different sets of sequin mixtures were used, one for wildtype

$PRPF31^{+/+}$ and another set for mutant $PRPF31^{+/-}$ samples. Libraries were prepared using the

NEBNext Ultra II Directional RNA Library Prep with Sample Purification

Beads (E7765S, New England Biolabs, Ipswich, MA, USA), NEBNext Poly(A) mRNA

Magnetic Isolation Module (E7490S, New England Biolabs, Ipswich, MA, USA), and NEBNext

Multiplex Oligos for Illumina (E6440S, New England Biolabs, Ipswich, MA, USA). Libraries

were normalized, pooled, and sequenced as described above.

Poly-A RNAseq Analysis of RPE cells and its EVs

Read and sample level QC was performed. Read quality was assessed with FastQC

(https://www.bioinformatics.babraham.ac.uk/projects/fastqc/) and MultiQC. An in-house Perl

script was used to check and filter out reads with presence of adapter and ambiguous character,

'N'. The Bowtie2 aligner was used to identify ribosomal contamination. Reads aligned to rRNA

reference sequences were dropped from all downstream analysis using SAMtools, resulting in

high quality (HQ) reads. The STAR aligner was used to align PE reads to a mouse reference in

two-pass mode within the sample and across replicates for each sample set. Feature Counts from

the Subread package (v2.0.3), was used to generate gene expression matrix with the following

non-default settings: reads must be paired, both the pairs must be mapped, use only uniquely

mapped reads, multi-mapped reads are not counted, chimeric reads were not counted, and strand

specificity turned on. Anaquin was used to further evaluate alignment sensitivity and gene

expression [57]. Here sensitivity indicates the fraction of annotated regions covered by alignment.

The Picard tools (http://broadinstitute.github.io/picard/) and RSeQC were used to calculate mean

fragment length. The approach implemented in Kallisto was used to covert raw reads to TPM

values. An average TPM of the lowest Sequins between test and control samples was calculated

and used as cutoff. DESeq2 was used for differential gene expression analysis. RNA-seq analysis

is described in detail in Mehrotra et al. [58]. g:Profiler (version e101_eg48_p14_baf17f0) with a

custom background gene set was used for gene set enrichment analysis (GSEA), pathway

enrichment and transcription factor binding site analysis.

<u>Reverse transcription and quantitative real-time polymerase chain reaction</u>

hi-RPE monolayers scraped off the plate into a 1.5-ml tube. Cells were washed twice in ice cold

PBS and centrifuged at $13.000 \times g$ for 1 min to obtain a dry pellet and store them at -80C. At the

time of the analysis, the pellet was resuspended in RLT plus buffer and RNA extraction was

performed according to manufacturer's protocol (RNeasy Micro Kit; Qiagen). Reverse

transcription was performed using SuperScript VILO cDNA Synthesis Kit (Life Technologies,

Thermo Fisher Scientific). Quantitative real-time PCRs (RT-qPCR), reactions were performed

with TaqMan® Fast Advanced Master Mix and validated TaqMan® Gene Expression Assays

(Life Technologies, Thermo Fisher Scientific) on QuantStudio™3 Real-Time instrument

(Applied Biosystems) for 40 cycles. Samples were run in quadruplicates, and expression levels

were normalized using 18S as housekeeping gene in the case of poly-A RNAs and hsa-miR26a-

5p for the miRNAs. The differential expression of the poly-A RNA and miRNA were expressed

as fold change (2^ (-$\Delta\Delta$Ct). The relative expression between PRPF31$^{+/-}$ hi-RPE and PRPF31$^{+/+}$

hi-RPE cells was compared by *Students t-test* and p value ≤ 0.05 was considered as significant. A

complete list of the TaqMan Gene Expression and miRNA Assays used are provided in

Supplementary Table S9 and S10.

Results

<u>Derivation and characterization of human RPE from human induced pluripotent stem cells</u>

<u>haploinsufficient for *PRPF31*.</u>

We previously generated hiPSCs harboring a 10 bp deletion in exon 7 of the *PRPF31*

gene using CRISPR/Cas9-mediated genome editing [17] . The resulting *PRPF31*[+/-] mutant and

isogenic wild-type control hiPSCs displayed typical ES cell-like colony morphology and

expressed pluripotency markers OCT4, SSEA4, NANOG, and SOX2 (**Supplementary Figure**

1). Both the *PRPF31*[+/−] mutant and isogenic control hiPSCs did not exhibit any chromosomal

abnormalities and were able to differentiate into the three embryonic germ cell layers evaluated

by the spontaneous *in vitro* differentiation EB model (**Supplementary Figure 1**).

The *PRPF31*[+/-] mutant and isogenic control hiPSC lines were differentiated into RPE using

established methods (**Figure 1A**)[17,59]. At Passage (P) 2, RPE cells differentiated from

PRPF31[+/−] mutant and *PRPF31*[+/+] control hiPSCs display the typical cobblestone morphology

pigmented cells (**Figure 1B-C**). They organized in monolayer polarized cultures and abundantly

expressed the Zonula occludens (ZO1) tight junction associated protein (**Figure 1D-E**). We have

previously shown that these *PRPF31*[+/−] mutant RPE cells produce reduced levels of PRPF31,

demonstrate defects in phagocytosis function, cilia formation and barrier function compared to

RPE from the isogenic control hiPSCs [16,17]. Further, the *PRPF31*[+/-] mutant RPE cells appear to

have irregular shapes as shown by ZO1 staining compared to *PRPF31*[+/+] control RPE cells, as

recently reported for RPE cells differentiated from patient-derived hiPSC lines harboring either a

duplication in exon 8 (c.709_734dup) or a deletion in exon 4 (c.269_273del) in the *PRPF31* gene

[18].

PRPF31 haploinsufficiency does not impact the number or physical properties of EVs produced

by the source hiPSC-RPE cells.

To investigate the effects of *PRPF31* haploinsufficiency and RPE dysfunction on the EVs

produced by RPE cells, 12 wells each of differentiated *PRPF31*$^{+/+}$ and *PRPF31*$^{+/-}$ iPSC-RPE

cells were matured on Transwell® inserts for 4 weeks. Once the hiPSC-RPE were mature, the

apical and basal conditioned media was collected for 12 weeks. EVs were isolated from the

conditioned media (CM) of the RPE monolayers, as described (**Figure 2A**) [60].

Characterization of the isolated EVs by nanoparticle tracking analysis (NTA) showed

they ranged in size from 100nm to 400nm, and EVs from the *PRPF31*$^{+/-}$ mutant and control RPE

cells showed the same size distributions and concentrations (**Figure 2B-E**). Transmission

electron microscopy (TEM) analysis showed the EVs were clear, rounded membrane vesicles as

anticipated (**Figures 2 F-G**) [61,62].

Sequencing of RNAs isolated from EVs produced by *PRPF31*-deficient and control RPE cells.

To study the RNA contents of the EVs produced by the control *PRPF31*$^{+/+}$ and mutant

PRPF31$^{+/-}$ hiPSC-RPE, we isolated EVs from the serum-free CM of 12 wells each of

differentiated *PRPF31*$^{+/+}$ and *PRPF31*$^{+/-}$ iPSC-RPE cells grown in on Transwells as described

above. Total RNA was isolated from the EVs and used to prepare both miRNA and poly-A

RNA libraries from each of the 12 experimental and control cultures. The resulting set of 24

miRNA libraries was multiplexed to reduce batch effects and sequenced on an Illumina MiSeq

instrument. At least 1.5 million reads were obtained for each miRNA library. Post QC, overall 8

mutants and 7 control samples were retained. For mutant samples total high quality (HQ) reads

ranged from 1.5-2.6 million with a median of 1.9 million. For wild-type samples HQ reads

ranged from 1.5-1.9 million with a median of 1.7 million.

Similarly, the set of 24 poly-A RNA libraries were multiplexed to reduced batch effects

and sequenced on an Illumina NovaSeq instrument. Post QC, all 12 mutant and 12 wild-types

samples were retained for differential gene expression analysis. For mutant samples HQ reads

ranged from 46-120 million with a median of 67 million reads. For the wild-type samples, HQ

reads ranged from 64-98 million with a median of 82 million reads.

PRPF31^{+/+} and *PRPF31*^{+/−} EVs have distinct miRNA and poly-A RNA profiles.

To investigate the effects of *PRPF31* haploinsufficiency and RPE dysfunction on the

RNA contents of RPE EVs, we sequenced the miRNAs and poly-A-RNAs contained in EVs

produced by control *PRPF31*^{+/+} and mutant *PRPF31*^{+/−} hiPSC-RPE as described above. Analyses

of these data identified 135 miRNAs within the hiPSC-RPE EVs (**Supplementary Table S1**). To

determine if any of these miRNAs could serve as a biomarker for RPE health, we compared the

miRNA profiles in EVs from *PRPF31*^{+/+} and *PRPF31*^{+/−} hiPSC-RPE. Significant enrichment of

miRNAs was determined using a combination of log fold change and FDR adjusted p-value (q-

value). This differential enrichment analysis showed that 12 miRNAs were enriched in the EVs

derived from the *PRPF31*^{+/−} hiPSC-RPE. In contrast, we found 6 miRNAs were depleted from

the *PRPF31*^{+/−} hiPSC-RPE EVs (**Figure 3**). These miRNAs are also highlighted in

Supplementary Table S4.

Examination of the poly-A RNA-seq reads mapped to the human genome revealed the

presence of 31,776 poly-A RNAs in the hiPSC-RPE EVs (**Supplementary Table S2)**.

Differential enrichment analysis of the poly-A RNA-seq data identified 865 differentially

enriched poly-A RNAs, with 551 enriched in and 314 depleted from the $PRPF31^{+/-}$ hiPSC-RPE

EVs (**Figure 4 and Supplemental Table S3**). Some of the poly-A RNAs were highly specific

for $PRPF31^{+/-}$ or $PRPF31^{+/+}$ hiPSC-RPE EVs, showing more than 1-million-fold enrichment in

the respective EVs (log2 (1,000,000) = 19.9).

<u>The majority of the miRNAs and poly-A RNAs contained in EVs are shared with hiPSC-RPE</u>

<u>cells.</u>

To test the hypothesis that EV RNAs can be used as surrogate biomarkers for the health status of

their source cells, we sequenced and compared the miRNAs and poly-A RNAs of the $PRPF31^{+/+}$

and $PRPF31^{+/-}$ hiPSC-RPE cells from which the EVs studied in the experiments described

above were derived. These comparisons showed that the majority of miRNA and poly-A RNAs

expressed in hiPSC-RPE cells are contained in the RPE-derived EVs. For miRNAs, 135 (64%)

of the 211 miRNAs identified in the source hiPSC-RPE cells were present in the EVs isolated

from the CM of those cells (**Supplementary Table S1 and S7**). EVs produced by the control

$PRPF31^{+/+}$ hiPSC-RPE contained 90% (19,228) of the total 21,375 poly-A RNAs identified in

the hiPSC-RPE (**Figure 5A**). Similarly, EVs produced by the mutant $PRPF31^{+/-}$ hiPSC-RPE

contained 90% (19,247) of the 21,354 total poly-A RNAs identified in the mutant hiPSC-RPE

(**Figure 5B and Supplementary Table S2 and S6**). These results indicate that the EVs

produced by hiPSC-RPE cells contain the great majority of RNAs of their parent cells,

supporting the potential use of EV RNAs as biomarkers. Interestingly, the EVs produced by the

hiPSC-RPE cells also contained many additional poly-A RNAs not identified within their parent

cells. $PRPF31^{+/+}$ EVs contained 11,984 unique poly-A RNAs and $PRPF31^{+/-}$ EVs contained

12,094 unique poly-A RNAs not found in their corresponding hiPSC-RPE cell transcriptomes

(**Figure 5A-B**). The EVs also contained a greater percentage of lncRNAs and pseudogene

mRNAs than the RPE cells (**Figure 5C-D**). Among the identified transcript RNA biotypes

found in RPE cells, the predominant category consisted of protein-coding mRNAs (70%),

followed by long non-coding RNAs (lncRNAs; 20%). In EVs the distribution of protein-coding

mRNAs (54%), and lncRNAs (30%) analysis revealed a higher abundance of the latter. (**Figure**

5C-D and Supplementary Table S2 and S6).

<u>Gene set enrichment analysis (GSEA) of poly-A RNAs and functional over-representation</u>

<u>analysis (ORA) of miRNAs enriched and depleted in the *PRPF31* mutant</u>

To explore the functional significance of the miRNAs and poly-A RNAs identified to be

differentially enriched in the EVs derived from the *PRPF31*$^{+/-}$ mutant hiPSC-RPE cells, we

performed gene set enrichment analysis (GSEA) of poly-A RNAs and functional over-

representation analysis (ORA) of miRNAs. GSEA for the 865 poly-A RNAs found to be

differentially enriched or depleted in the EVs from *PRPF31*$^{+/-}$ mutant hiPSC-RPE cells did not

identify any enrichment of functional groups [63].

We used ORA to evaluate the potential functional significance of the miRNAs found to

be differentially enriched in EVs from *PRPF31*$^{+/-}$ vs. *PRPF31*$^{+/+}$ hiPSC-RPE cells. ORA is used

to evaluate the enrichment of transcripts regulated by miRNAs in functional categories derived

from multiple databases, including gene ontology (GO, miRTarBase GO, miRWalk), target

genes (miRTarBase), pathways (miRWalk, KEGG, miRPathDB), diseases (MNDR) and

biological processes (miRPathD) [55]. ORA found that 11 of the 18 miRNAs significantly enriched

or depleted in the EVs derived from *PRPF31*$^{+/-}$ mutant hiPSC-RPE cells were associated with

eye diseases. Three of these miRNAs were associated specifically with retinal degeneration and

6 were associated with negative regulation of epithelial cell differentiation (**Supplementary**

Figure 4).

Gene ontology analysis revealed that of the 18 miRNAs enriched or depleted in the EVs

derived from *PRPF31*[+/-] mutant hiPSC-RPE cells 6 were enriched in categories associated with

cell de-differentiation or cell death and 6 with retinal function and degeneration (**Figure 6A and**

Supplementary Table S5). For example, miRTarBase analysis identified *CDH1* and *RDH11*

among the top most significantly overrepresented genes and each gene was targeted by a

distinctive set of 6 miRNAs. *CDH1* is as a potential gene target of the differentially regulated

miRNAs: miR-129-5p; miR-193b-3p; miR-34c-5p; miR-9-5p; miR-9-3p; miR-92a-3p. (**Figure**

6B). *CDH1* encodes for E-Cadherin, a structural protein found in the adherens junctions of the

RPE cells which is dysregulated in pathological conditions of the RPE [64,65]. Another distinctive

set of differentially regulated miRNAs (miR-129-5p, miR-378a-3p, miR-34c-5p, miR-375-3p,

miR-449b-5p, miR-449a) targeted *RDH11* gene which encodes Retinol dehydrogenase 11,

related to visual phototransduction and with strong links with other RPE functions such as

phagocytosis [66-68] (**Figure 6C).** The levels of *CDH1* and *RDH11* mRNAs were not altered in the

PRPF31[+/-] hiPSC-RPE cells themselves (**Supplementary Table S8**).

<u>Validation of selected enriched poly-A RNAs and miRNAs by quantitative RT-PCR</u>

We isolated total RNA from *PRPF31*[+/+] control and *PRPF31*[+/−] mutant hiPSC-RPE cells

for qRT-PCR analyses of selected miRNAs and poly-A RNAs identified to be enriched or

depleted in the EVs from the *PRPF31*[+/−] mutant hiPSC-RPE cells in the RNA sequencing

studies. We choose 6 different poly-A RNAs among those differentially enriched in the

PRPF31[+/−] hiPSC-RPE cells (**Supplementary Figure 1A**) compared to the *PRPF31*[+/+] control

based on their read counts and RNA-seq trend. The qRT-PCR results showed that for 5 out 6

poly-A RNAs the levels in the EVs from *PRPF31$^{+/-}$* and *PRPF31$^{+/+}$* hiPSC-RPE cells were

significantly different, and the changes detected were consistent with the RNA-seq data with

regards to enrichment of depletion in the EVs from *PRPF31$^{+/-}$* mutant hiPSC-RPE cells

(**Supplementary Figure 2**). Similar qRT-PCR analyses were performed for 5 miRNAs

identified to be differentially enriched in the EVs from *PRPF31$^{+/-}$* mutant hiPSC-RPE cells from

the RNA-sequencing data. qRT-PCR confirmed the differential enrichment of the miRNAs for 4

out of the 5 selected miRNAs in the *PRPF31$^{+/-}$* heterozygous and control EVs (**Supplementary**

Figure 3).

Discussion

We studied the RNA contents of EVs produced by control *PRPF31$^{+/+}$* and mutant

PRPF31$^{+/-}$ hiPSC-RPE cells to test the hypothesis that EV RNA contents have the potential to

be used as surrogate biomarkers of RPE health. We have found that the RNA cargo in EVs

mirrors that of their originating RPE cells, reflecting disease-induced changes in RPE cell

differentiation and function. Further, by comparing the RNA profiles of EVs produced by

PRPF31$^{+/+}$ and *PRPF31$^{+/-}$* hiPSC-RPE cells, we found 18 miRNAs and 865 poly-A RNAs that

are differentially enriched or depleted in the EVs from *PRPF31$^{+/-}$* mutant hiPSC-RPE cells. This

includes many RNAs that were enriched or depleted at least 1-million-fold compared to EVs

from control cells. Further, miRNAs differentially enriched in EVs from the *PRPF31$^{+/-}$* mutant

hiPSC-RPE were identified as regulators of genes associated with RPE differentiation and

function, supporting the functional relevance of the differentially enriched EV RNAs. Overall,

these findings support the hypothesis that the RNA contents of EVs produced by RPE cells

changes during disease, consistent with the potential use of EV RNAs as biomarkers of RPE

health.

As a first step in assessing the potential use of EV RNAs as biomarkers of RPE health

status, we compared the poly-A RNA profiles of EVs and their source cells. We found the

majority (90%) of the poly-A RNAs detected in the parent hiPSC-RPE cells were contained in

the EVs derived from those cells, supporting the concept that EV RNAs reflect the

transcriptomes of the cells they are derived from [29,35-39]. Of interest, we also found many poly-A

RNAs were detected only in EVs, including additional lncRNAs. The higher abundance of

lncRNA in the EVs is intriguing and suggests specific regulatory mechanisms at play. Long

non-coding RNAs (lncRNAs) are thought to play regulatory roles in multiple biological

processes, including the differentiation and function of RPE cells [69,70]. Dysregulation of

lncRNAs have been associated with age-related macular degeneration (AMD)[71,72]. Other studies

have reported similar differences in RNA profiles and abundance when comparing EVs to their

source cells [73,74]. Although the mechanism of RNA sorting and loading into EVs is currently

unknown, some studies suggest the contents of EVs are loaded by source cells as a mechanism of

controlling intracellular availability of certain RNAs [37,75,76]. Further studies will be needed to

investigate the roles of EV lncRNAs in RPE biology and disease. It is also important to consider

for this study and others, EVs were isolated from pooled CM that was collected for several

weeks, while source cell RNA was isolated from one time point, so differences can also be due

temporal RNA expression differences at the time of sample collection. According to the latest

MISEV recommendations, no method for isolating EVs can completely guarantee the absence of

non-vesicular particles. However, the primary potential contaminants, lipoprotein particles, do

not contain RNA and, therefore, will not contribute to the RNA content in EVs for the purposes

of this study.

Comparisons of the RNA contents of EVs produced by *PRPF31*[+/+] and *PRPF31*[+/−]

hiPSC-RPE cells identified 12 miRNAs that were enriched and 6 miRNAs depleted in the EVs

derived from the *PRPF31*[+/−] hiPSC-RPE. Similarly, differential enrichment analysis of the poly-

A RNA-seq data identified 551 poly-A RNAs enriched in and 314 depleted from the *PRPF31*[+/−]

hiPSC-RPE EVs. Some of the poly-A RNAs were highly specific for *PRPF31*[+/−] or *PRPF31*[+/+]

hiPSC-RPE EVs, supporting their potential use as biomarkers of RPE health and/or disease. The

RNAseq data comes from a large number of replicates (7-8 samples per condition) making these

results statistically robust [58]. Additionally, we have validated the directional change of RNA

concentration for several miRNAs and poly-A RNAs using quantitative RT-PCR as an

orthogonal method.

It has been previously demonstrated that mRNAs and miRNAs can be transferred via

EVs and in turn regulate gene expression of recipient cells [77-81] The data presented here provide a

starting point for future studies of the function(s) of RNAs contained in EVs produced by the

RPE. For example, photoreceptor and RPE cells function together, with coordinated metabolism

and recycling of shed photoreceptor outer segments via RPE phagocytosis [82]. Since overall

retinal function and health depends on these shared processes, it is attractive to hypothesize that

EVs may be used for intercellular signaling between the RPE and photoreceptor cells, however

further studies are needed to support this hypothesis [83].

We focused these initial studies of EV contents on RNAs since with current technologies

better depth of sampling is possible for RNA sequencing than proteomic analysis [84]. Further, EV

RNAs have been reported to be promising biomarkers of cancer and other forms of

neurodegeneration [85-88]. Future studies could also evaluate the protein contents of RPE and

retina derived EVs, as reported [41].

We were especially interested to find that some of the miRNAs differentially enriched in

EVs from the *PRPF31*[+/-] mutant hiPSC-RPE were identified as regulators of genes associated

with RPE differentiation and function. Among the 18 most significantly altered miRNAs

between the EVs produced by *PRPF31*[+/-] and *PRPF31*[+/+] cells, three were linked to retinal

degeneration, and six were connected to the negative regulation of epithelial cell differentiation.

This suggests that the defective differentiation and function of RPE cells caused by *PRPF31*

haploinsufficiency is reflected in the miRNAs content in the EVs produced by the mutant RPE

cells [16,17]. We believe these findings support the hypothesis that EV RNAs can serve as

surrogate biomarkers of retinal and RPE health status, and with further study may provide an

alternative approach to assess the response of patients with IRDs to potential therapies, such as

AAV-mediated gene therapies which show great promise for treating IRDs [89].

Acknowledgements

We are grateful for the contributions of other Ocular Genomics Institute team members,

including Daniela Pignatta and the Ocular Genomics Core for their assistance with the RNA-seq.

We also acknowledge Philip Seifert and the Schepens Eye Research Institute Morphology Core

for their assistance with the TEM imaging. We thank Amanda Fernández-Rodríguez for her

assistance with miRNA sequencing analysis. The research reported was supported by a Stein

Innovation Award from Research to Prevent Blindness (EAP), grants from the National Eye

Institute to EAP (EY012910, EY020902), and a P30 Core Grant for Vision Research

(EY014104).

Abbreviations:

AAV: adeno-associated virus

CM: conditioned media

EB: Embryoid Body

EVs: Extracellular vesicles

GSEA: gene set enrichment analysis

hiPSC: human induced pluripotent stem cell

hiPSC-RPE: hiPSC-derived RPE

IRD: inherited retinal degeneration

lncRNAs: long non-coding RNAs

MISEV: Minimal Information for Studies of Extracellular Vesicles

NTA: Nanoparticle tracking analysis

ORA: over-representation analysis

PRPF31: pre-mRNA processing factor 31

RPE: retinal pigment epithelium

RP: retinitis pigmentosa

TEM: transmission electron microscopy

ZO1: Zonula occludens

Ethics approval and consent to participate

The work reported here has been acknowledged by the Mass General Brighma Instituttional

Review Board as "not human subjects research" (approved application # 2022P000301)

Availability of data and materials

All data supporting the findings of this study are available within the paper and its

supplementary information. RNAseq raw sequence reads were deposisted in NCBIs SRA can be

found in https://www.ncbi.nlm.nih.gov/bioproject/PRJNA824312.

Declaration of interests

The authors have no competing interests or other interests that might be perceived to influence

the results and/or discussion reported in this paper.

Authors' contributions

All authors have made substantial contributions to this manuscript. EAP and MGH conception

EAP HG and MGH design of the work; HG TK SM RFG EAP and MGH the acquisition,

analysis, and interpretation of data; HG EAP and MGH have drafted the work and substantively

revised it. All authors have approved the submitted version

Corresponding authors

Eric A. Pierce Eric_pierce@meei.harvard.edu or Marcela Garita-Hernandez

mgarita@meei.harvard.edu

Figures and Legends

Figure 1. Generation of hiPSC-derived RPE cells. (A) Schematic process of the directed

differentiation of *PRPF31$^{+/+}$* and *PRPF31$^{+/−}$* hiPSC into RPE cells (B–C) Phase contrast images

of *PRPF31$^{+/+}$* and *PRPF31$^{+/-}$* hiPSC-RPE showing the cobblestone morphology of the pigmented

RPE cells. (D-E) Confocal images of *PRPF31$^{+/+}$* and *PRPF31$^{+/-}$* hiPSC-RPE cells

immunoreactive for the tight junction marker ZO1.

Figure 2. Characterization of EVs from PRPF31+/+ and PRPF31+/− iPSC-RPE cells. A-B) Distribution of EVs by size using nanoparticle tracking analysis technology. C) The average size of *PRPF31⁺/⁺* EVs was 247 ± 16 nm and *PRPF31⁺/⁻* EVs 206 ± 26 nm. D) The concentration of *PRPF31⁺/⁺* EVs was $3.8 \times 10^9 \pm 1.2 \times 10^9$ EVs/ml and *PRPF31⁺/⁻* EVs was $2.7 \times 10^9 \pm 1.1 \times 10^9$ EVs/ml. N=4, error bars: standard error of the mean. D) Ultrastructural analysis by TEM of EVs isolated from *PRPF31⁺/⁺* and *PRPF31⁺/⁻* conditioned media; scale bar is 100μm.

A.

Differentially Expressed miRNAs

Figure 3. miRNAs profiling of EVs from PRPF31$^{+/+}$ and PRPF31$^{+/−}$. Volcano plots indicating

the log2 (fold change) of differentially expressed miRNAs on the x-axis and the -log10 (adjusted

P-value <0.05 8:7,Mut:WT) on the y-axis. miRNAs in red are significantly enriched (right) or

depleted (left) in the *PRPF31*$^{+/−}$ hiPSC-RPE EVs.

Figure 4. Comparison of the poly-A RNAs contained in EVs from PRPF31⁺/⁺ and PRPF31⁺/⁻

hiPSC-RPE. Volcano plot indicating the differentially expressed poly-A RNAs in the EVs

isolated from *PRPF31⁺/⁻* compared to the WT expressed as the log2 (fold change) on the x-axis

and the -log10 (adjusted P-value <0.05, Mut:Wt;12:12) on the y-axis. Poly-A RNAs indicated by

red dots were significantly enriched in (right) or depleted from (left) the *PRPF31⁺/⁻* hiPSC-RPE

based on fold change and adjusted P value.

Figure 5. Comparison of RNAs identified in EVs produced by PRPF31$^{+/+}$ and PRPF31$^{+/-}$ hiPSC-RPE cells, and the RNAs contained in the hiPSC-RPE cells (A, B) Venn diagrams showing unique and shared expressed poly-A RNAs among (A) *PRPF31$^{+/+}$* and (B) *PRPF31$^{+/−}$* hi-RPE cells and their EVs. (**C, D**) Relative biotype distribution of dysregulated transcripts in RPE cells and EV poly-A RNAs.

A. Differentially regulated miRNAs significant among *PRPF31*[+/-] vs. *PRPF31*[+/+]

B. Functional over-representation analysis of DEG miRNA in *PRPF31*[+/-]

Figure 6. Functional over-representation analysis (ORA) of enriched and depleted miRNAs in

the PRPF31 mutant. A) ORA analysis of differentially regulated miRNAs in *PRFP31$^{+/-}$*

compared to *PRFP31$^{+/+}$* showed two distinctive subsets of miRNAs strongly associated with de-

differentiation and cell death or associated with retinal function and degeneration. B.

miRTarBase analysis identified *CDH1* and *RDH11* genes as top potential targets of the

distinctive miRNAs subsets, which can explain the defects in the differentiation and function in

the diseased *PRPF31$^{+/-}$* RPE cell.

References

1.   Sahel, J.A., Marazova, K., and Audo, I. (2014). Clinical characteristics and current
therapies for inherited retinal degenerations. Cold Spring Harb Perspect Med *5*, a017111.
10.1101/cshperspect.a017111.

2.   Schneider, N., Sundaresan, Y., Gopalakrishnan, P., Beryozkin, A., Hanany, M., Levanon,
E.Y., Banin, E., Ben-Aroya, S., and Sharon, D. (2021). Inherited retinal diseases: Linking
genes, disease-causing variants, and relevant therapeutic modalities. Prog Retin Eye Res,
101029. 10.1016/j.preteyeres.2021.101029.

3.   Nishiguchi, K.M., and Rivolta, C. (2012). Genes associated with retinitis pigmentosa and
allied diseases are frequently mutated in the general population. PLoS One *7*, e41902.
10.1371/journal.pone.0041902.

4.   Berson, E.L. (1993). Retinitis pigmentosa. The Friedenwald Lecture. Investigative
Ophthalmology & Visual Science *34*, 1659-1676.

5.   Vithana, E.N., Abu-Safieh, L., Allen, M.J., Carey, A., Papaioannou, M., Chakarova, C.,
Al Maghtheh, M., Ebenezer, N.D., Willis, C., Moore, A.T., Bird, A.C., et al. (2001). A
human homolog of yeast pre-mRNA splicing gene, PRP31, underlies autosomal
dominant retinitis pigmentosa on chromosome 19q13.4 (RP11). Molecular Cell *8*, 375-
381.

6.   Tanackovic, G., Ransijn, A., Thibault, P., Abou Elela, S., Klinck, R., Berson, E.L.,
Chabot, B., and Rivolta, C. (2011). PRPF mutations are associated with generalized
defects in spliceosome formation and pre-mRNA splicing in patients with retinitis
pigmentosa. Human Molecular Genetics *20*, 2116-2130. 10.1093/hmg/ddr094.

7.   Daiger, S.P., Bowne, S.J., and Sullivan, L.S. (2014). Genes and Mutations Causing
Autosomal Dominant Retinitis Pigmentosa. Cold Spring Harb Perspect Med *5*.
10.1101/cshperspect.a017129.

8.   Kohn, L., Bowne, S.J., L, S.S., Daiger, S.P., Burstedt, M.S., Kadzhaev, K., Sandgren, O.,
and Golovleva, I. (2009). Breakpoint characterization of a novel approximately 59 kb
genomic deletion on 19q13.42 in autosomal-dominant retinitis pigmentosa with
incomplete penetrance. Eur J Hum Genet *17*, 651-655. 10.1038/ejhg.2008.223.

9.   Hafler, B.P., Comander, J., Weigel DiFranco, C., Place, E.M., and Pierce, E.A. (2016).
Course of Ocular Function in PRPF31 Retinitis Pigmentosa. Seminars in Ophthalmology
31, 49-52. 10.3109/08820538.2015.1114856.

10.   Ruberto, F.P., Balzano, S., Namburi, P., Kimchi, A., Pescini-Gobert, R., Obolensky, A.,
Banin, E., Ben-Yosef, T., Sharon, D., and Rivolta, C. (2021). Heterozygous deletions of
noncoding parts of the PRPF31 gene cause retinitis pigmentosa via reduced gene
expression. Mol Vis *27*, 107-116.

11.   Yang, C., Georgiou, M., Atkinson, R., Collin, J., Al-Aama, J., Nagaraja-Grellscheid, S.,
Johnson, C., Ali, R., Armstrong, L., Mozaffari-Jovin, S., and Lako, M. (2021). Pre-
mRNA Processing Factors and Retinitis Pigmentosa: RNA Splicing and Beyond. Front
Cell Dev Biol *9*, 700276. 10.3389/fcell.2021.700276.

12.   Mousawi, Z., Choukeir, M., Jaffal, L., Karam, L., Assi, A., Ibrahim, J.-N., Chebly, A.,
599   and El Shamieh, S. (2024). A Novel Copy Number Variation in PRPF31 Causes
600   Dominant Rod-Cone Dystrophy By Haploinsufficiency.

601   13.   Bujakowska, K.M., Maubaret, C., Chakarova, C.F., Tanimoto, N., Beck, S.C., Fahl, E.,
602   Humphries, M.M., Kenna, P., Makarov, E., Makarova, O., Paquet-Durand, F., et al.
603   (2009). Study of gene targeted mouse models of splicing factor gene Prpf31 implicated in
604   human autosomal dominant retinitis pigmentosa (RP). Invest Ophthalmol.Vis.Sci.

605   14.   Graziotto, J.J., Farkas, M.H., Bujakowska, K., Deramaudt, B.M., Zhang, Q., Nandrot,
606   E.F., Inglehearn, C.F., Bhattacharya, S.S., and Pierce, E.A. (2011). Three gene-targeted
607   mouse models of RNA splicing factor RP show late-onset RPE and retinal degeneration.
608   Investigative Ophthalmology & Visual Science *52*, 190-198. 10.1167/iovs.10-5194.

609   15.   Farkas, M.H., Lew, D.S., Sousa, M.E., Bujakowska, K., Chatagnon, J., Bhattacharya,
610   S.S., Pierce, E.A., and Nandrot, E.F. (2014). Mutations in Pre-mRNA Processing Factors
611   3, 8, and 31 Cause Dysfunction of the Retinal Pigment Epithelium. Am J Pathol *184*,
612   2641-2652. 10.1016/j.ajpath.2014.06.026.

613   16.   Buskin, A., Zhu, L., Chichagova, V., Basu, B., Mozaffari-Jovin, S., Dolan, D., Droop, A.,
614   Collin, J., Bronstein, R., Mehrotra, S., Farkas, M., et al. (2018). Human iPSC-derived
615   RPE and retinal organoids reveal impaired alternative splicing of genes involved in pre-
616   mRNA splicing in PRPF31 autosomal dominant retinitis pigmentosa. Nature
617   Communication *9*, 4234. 10.1038/s41467-018-06448-y.

618   17.   Brydon, E.M., Bronstein, R., Buskin, A., Lako, M., Pierce, E.A., and Fernandez-Godino,
619   R. (2019). AAV-Mediated Gene Augmentation Therapy Restores Critical Functions in
620   Mutant PRPF31(+/-) iPSC-Derived RPE Cells. Mol Ther Methods Clin Dev *15*, 392-402.
621   10.1016/j.omtm.2019.10.014.

622   18.   Rodrigues, A., Slembrouck-Brec, A., Nanteau, C., Terray, A., Tymoshenko, Y., Zagar,
623   Y., Reichman, S., Xi, Z., Sahel, J.A., Fouquet, S., Orieux, G., et al. (2022). Modeling
624   PRPF31 retinitis pigmentosa using retinal pigment epithelium and organoids combined
625   with gene augmentation rescue. NPJ Regen Med *7*, 39. 10.1038/s41536-022-00235-6.

626   19.   Bronstein, R., Mehrotra, S., Capowski, E.E., Foltz, L., Farkas, M.H., Gamm, D.M.,
627   Clegg, D.O., and Pierce, E.A. (2019). Mutations in splicing factors that cause Retinitis
628   Pigmentosa (RP) do not affect splicing in a global manner. Investigative Ophthalmology
629   & Visual Science *60*, 379-379.

630   20.   Xi, Z., Vats, A., Sahel, J.-A., Chen, Y., and Byrne, L.C. (2022). Gene augmentation
631   prevents retinal degeneration in a CRISPR/Cas9-based mouse model of PRPF31 retinitis
632   pigmentosa. Nature Communications *13*, 7695.

633   21.   Banin, E., Gootwine, E., Obolensky, A., Ezra-Elia, R., Ejzenberg, A., Zelinger, L.,
634   Honig, H., Rosov, A., Yamin, E., and Sharon, D. (2015). Gene augmentation therapy
635   restores retinal function and visual behavior in a sheep model of CNGA3 achromatopsia.
636   Molecular therapy *23*, 1423-1433.

637   22.   Komáromy, A.M., Koehl, K.L., and Park, S.A. (2021). Looking into the future: Gene and
638   cell therapies for glaucoma. Veterinary ophthalmology *24*, 16-33.

23. Maguire, A.M., Simonelli, F., Pierce, E.A., Pugh Jr, E.N., Mingozzi, F., Bennicelli, J.,
Banfi, S., Marshall, K.A., Testa, F., and Surace, E.M. (2008). Safety and efficacy of gene
transfer for Leber's congenital amaurosis. New England Journal of Medicine *358*, 2240-
2248.

24. Bainbridge, J.W., Smith, A.J., Barker, S.S., Robbie, S., Henderson, R., Balaggan, K.,
Viswanathan, A., Holder, G.E., Stockman, A., Tyler, N., Petersen-Jones, S., et al. (2008).
Effect of gene therapy on visual function in Leber's congenital amaurosis. New England
Journal of Medicine *358*, 2231-2239. doi: 10.1056/NEJMoa0802268.

25. Rodrigues, G.A., Shalaev, E., Karami, T.K., Cunningham, J., Slater, N.K., and Rivers,
H.M. (2019). Pharmaceutical development of AAV-based gene therapy products for the
eye. Pharmaceutical research *36*, 29.

26. Calzetti, G., Levy, R.A., Cideciyan, A.V., Garafalo, A.V., Roman, A.J., Sumaroka, A.,
Charng, J., Heon, E., and Jacobson, S.G. (2018). Efficacy Outcome Measures for Clinical
Trials of USH2A Caused by the Common c.2299delG Mutation. Am J Ophthalmol *193*,
114-129. 10.1016/j.ajo.2018.06.017.

27. Xu, J., Wang, Y., Hsu, C.-Y., Gao, Y., Meyers, C.A., Chang, L., Zhang, L., Broderick,
K., Ding, C., and Peault, B. (2019). Human perivascular stem cell-derived extracellular
vesicles mediate bone repair. Elife *8*, e48191.

28. Chronopoulos, A., and Kalluri, R. (2020). Emerging role of bacterial extracellular
vesicles in cancer. Oncogene *39*, 6951-6960.

29. Keerthikumar, S., Chisanga, D., Ariyaratne, D., Al Saffar, H., Anand, S., Zhao, K.,
Samuel, M., Pathan, M., Jois, M., Chilamkurti, N., Gangoda, L., et al. (2016). ExoCarta:
A Web-Based Compendium of Exosomal Cargo. J Mol Biol *428*, 688-692.
10.1016/j.jmb.2015.09.019.

30. Maas, S.L.N., Breakefield, X.O., and Weaver, A.M. (2017). Extracellular Vesicles:
Unique Intercellular Delivery Vehicles. Trends Cell Biol *27*, 172-188.
10.1016/j.tcb.2016.11.003.

31. Valadi, H., Ekstrom, K., Bossios, A., Sjostrand, M., Lee, J.J., and Lotvall, J.O. (2007).
Exosome-mediated transfer of mRNAs and microRNAs is a novel mechanism of genetic
exchange between cells. Nat Cell Biol *9*, 654-659. 10.1038/ncb1596.

32. Ridder, K., Keller, S., Dams, M., Rupp, A.K., Schlaudraff, J., Del Turco, D., Starmann,
J., Macas, J., Karpova, D., Devraj, K., Depboylu, C., et al. (2014). Extracellular vesicle-
mediated transfer of genetic information between the hematopoietic system and the brain
in response to inflammation. PLoS Biol *12*, e1001874. 10.1371/journal.pbio.1001874.

33. Ying, W., Riopel, M., Bandyopadhyay, G., Dong, Y., Birmingham, A., Seo, J.B.,
Ofrecio, J.M., Wollam, J., Hernandez-Carretero, A., Fu, W., Li, P., et al. (2017). Adipose
Tissue Macrophage-Derived Exosomal miRNAs Can Modulate In Vivo and In Vitro
Insulin Sensitivity. Cell *171*, 372-384 e312. 10.1016/j.cell.2017.08.035.

34. Mighty, J., Rubio-Navarro, A., Shi, C., Zhou, J., Flores-Bellver, M., Heissel, S.,
Onwumere, O., Einbond, L., Gharbaran, R., Casper, D.S., Benito-Martin, A., et al.
(2022). Extracellular vesicles of human diabetic retinopathy retinal tissue and urine of

diabetic retinopathy patients are enriched for the junction plakoglo bin protein. Front
Endocrinol (Lausanne) *13*, 1077644. 10.3389/fendo.2022.1077644.

35.   Ni, H., Capodici, J., Cannon, G., Communi, D., Boeynaems, J.-M., Karikó, K., and
Weissman, D. (2002). Extracellular mRNA induces dendritic cell activation by
stimulating tumor necrosis factor-α secretion and signaling through a nucleotide
receptor∗. Journal of Biological Chemistry *277*, 12689-12696.

36.   Hoy, A.M., and Buck, A.H. (2012). Extracellular small RNAs: what, where, why?
Biochemical Society transactions *40*, 886-890.

37.   Kobayashi, M., Salomon, C., Tapia, J., Illanes, S.E., Mitchell, M.D., and Rice, G.E.
(2014). Ovarian cancer cell invasiveness is associated with discordant exosomal
sequestration of Let-7 miRNA and miR-200. Journal of translational medicine *12*, 1-12.

38.   La Marca, V., and Fierabracci, A. (2017). Insights into the diagnostic potential of
extracellular vesicles and their miRNA signature from liquid biopsy as early biomarkers
of diabetic micro/macrovascular complications. International journal of molecular
sciences *18*, 1974.

39.   Klingeborn, M., Dismuke, W.M., Rickman, C.B., and Stamer, W.D. (2017). Roles of
exosomes in the normal and diseased eye. Progress in retinal and eye research *59*, 158-
177.

40.   Yu, W., Hurley, J., Roberts, D., Chakrabortty, S.K., Enderle, D., Noerholm, M.,
Breakefield, X.O., and Skog, J.K. (2021). Exosome-based liquid biopsies in cancer:
opportunities and challenges. Ann Oncol *32*, 466-477. 10.1016/j.annonc.2021.01.074.

41.   Klingeborn, M., Dismuke, W.M., Skiba, N.P., Kelly, U., Stamer, W.D., and Bowes
Rickman, C. (2017). Directional Exosome Proteomes Reflect Polarity-Specific Functions
in Retinal Pigmented Epithelium Monolayers. Sci Rep *7*, 4901. 10.1038/s41598-017-
05102-9.

42.   Zhou, J., Flores-Bellver, M., Pan, J., Benito-Martin, A., Shi, C., Onwumere, O., Mighty,
J., Qian, J., Zhong, X., Hogue, T., Amponsah-Antwi, B., et al. (2021). Human retinal
organoids release extracellular vesicles that regulate gene expression in target human
retinal progenitor cells. Sci Rep *11*, 21128. 10.1038/s41598-021-00542-w.

43.   Ragusa, M., Barbagallo, C., Statello, L., Caltabiano, R., Russo, A., Puzzo, L., Avitabile,
T., Longo, A., Toro, M.D., and Barbagallo, D. (2015). miRNA profiling in vitreous
humor, vitreal exosomes and serum from uveal melanoma patients: Pathological and
diagnostic implications. Cancer biology & therapy *16*, 1387-1396.

44.   Ragusa, M., Caltabiano, R., Russo, A., Puzzo, L., Avitabile, T., Longo, A., Toro, M.D.,
Di Pietro, C., Purrello, M., and Reibaldi, M. (2013). MicroRNAs in vitreus humor from
patients with ocular diseases. Molecular vision *19*, 430.

45.   ElShelmani, H., Wride, M.A., Saad, T., Rani, S., Kelly, D.J., and Keegan, D. (2020).
Identification of novel serum microRNAs in age-related macular degeneration.
Translational Vision Science & Technology *9*, 28-28.

46. Ertekin, S., Yıldırım, Ö., Dinç, E., Ayaz, L., Fidancı, Ş.B., and Tamer, L. (2014).
Evaluation of circulating miRNAs in wet age-related macular degeneration. Molecular
vision *20*, 1057.

47. Tanaka, Y., Tsuda, S., Kunikata, H., Sato, J., Kokubun, T., Yasuda, M., Nishiguchi,
K.M., Inada, T., and Nakazawa, T. (2014). Profiles of extracellular miRNAs in the
aqueous humor of glaucoma patients assessed with a microarray system. Scientific
reports *4*, 5089.

48. Flores-Bellver, M., Mighty, J., Aparicio-Domingo, S., Li, K.V., Shi, C., Zhou, J., Cobb,
H., McGrath, P., Michelis, G., and Lenhart, P. (2021). Extracellular vesicles released by
human retinal pigment epithelium mediate increased polarised secretion of drusen
proteins in response to AMD stressors. Journal of extracellular vesicles *10*, e12165.

49. Hernandez, B.J., Skiba, N.P., Plossl, K., Strain, M., Liu, Y., Grigsby, D., Kelly, U., Cady,
M.A., Manocha, V., Maminishkis, A., Watkins, T., et al. (2023). Polarized Desmosome
and Hemidesmosome Shedding via Small Extracellular Vesicles is an Early Indicator of
Outer Blood-Retina Barrier Dysfunction. J Extracell Biol *2*. 10.1002/jex2.116.

50. Getachew, H., Chinchilla, B., and Fernandez-Godino, R. (2022). Genome Editing of
Induced Pluripotent Stem Cells Using CRISPR/Cas9 Ribonucleoprotein Complexes to
Model Genetic Ocular Diseases. Methods Mol Biol *2549*, 321-334.
10.1007/7651_2021_409.

51. Welsh, J.A., Goberdhan, D.C.I., O'Driscoll, L., Buzas, E.I., Blenkiron, C., Bussolati, B.,
Cai, H., Di Vizio, D., Driedonks, T.A.P., Erdbrugger, U., Falcon-Perez, J.M., et al.
(2024). Minimal information for studies of extracellular vesicles (MISEV2023): From
basic to advanced approaches. J Extracell Vesicles *13*, e12404. 10.1002/jev2.12404.

52. Théry, C., Amigorena, S., Raposo, G., and Clayton, A. (2006). Isolation and
characterization of exosomes from cell culture supernatants and biological fluids. Current
protocols in cell biology *30*, 3.22. 21-23.22. 29.

53. Giraldez, M.D., Spengler, R.M., Etheridge, A., Godoy, P.M., Barczak, A.J., Srinivasan,
S., De Hoff, P.L., Tanriverdi, K., Courtright, A., and Lu, S. (2018). Comprehensive
multi-center assessment of small RNA-seq methods for quantitative miRNA profiling.
Nature biotechnology *36*, 746-757.

54. Backes, C., Khaleeq, Q.T., Meese, E., and Keller, A. (2016). miEAA: microRNA
enrichment analysis and annotation. Nucleic Acids Res *44*, W110-116.
10.1093/nar/gkw345.

55. Kern, F., Fehlmann, T., Solomon, J., Schwed, L., Grammes, N., Backes, C., Van Keuren-
Jensen, K., Craig, D.W., Meese, E., and Keller, A. (2020). miEAA 2.0: integrating multi-
species microRNA enrichment analysis and workflow management systems. Nucleic
Acids Res *48*, W521-W528. 10.1093/nar/gkaa309.

56. Hardwick, S.A., Chen, W.Y., Wong, T., Deveson, I.W., Blackburn, J., Andersen, S.B.,
Nielsen, L.K., Mattick, J.S., and Mercer, T.R. (2016). Spliced synthetic genes as internal
controls in RNA sequencing experiments. Nature methods *13*, 792-798.

57.   Wong, T., Deveson, I.W., Hardwick, S.A., and Mercer, T.R. (2017). ANAQUIN: a
software toolkit for the analysis of spike-in controls for next generation sequencing.
Bioinformatics *33*, 1723-1724. 10.1093/bioinformatics/btx038.

58.   Mehrotra, S., Bronstein, R., Navarro-Gomez, D., Segre, A.V., and Pierce, E.A. (2020).
Evaluating Methods for Differential Gene Expression And Alternative Splicing Using
Internal Synthetic Controls. BioRxiv *238295*,
https://doi.org/10.1101/2020.1108.1105.238295.

59.   Leach, L.L., and Clegg, D.O. (2015). Concise Review: Making Stem Cells Retinal:
Methods for Deriving Retinal Pigment Epithelium and Implications for Patients With
Ocular Disease. Stem Cells *33*, 2363-2373. 10.1002/stem.2010.

60.   Théry, C., Witwer, K.W., Aikawa, E., Alcaraz, M.J., Anderson, J.D., Andriantsitohaina,
R., Antoniou, A., Arab, T., Archer, F., Atkin-Smith, G.K., Ayre, D.C., et al. (2018).
Minimal information for studies of extracellular vesicles 2018 (MISEV2018): a position
statement of the International Society for Extracellular Vesicles and update of the
MISEV2014 guidelines. J Extracell Vesicles *7*, 1535750.
10.1080/20013078.2018.1535750.

61.   Welsh, J.A., Goberdhan, D.C.I., O'Driscoll, L., Buzas, E.I., Blenkiron, C., Bussolati, B.,
Cai, H., Di Vizio, D., Driedonks, T.A.P., Erdbrügger, U., Falcon-Perez, J.M., et al.
(2024). Minimal information for studies of extracellular vesicles (MISEV2023): From
basic to advanced approaches. J Extracell Vesicles *13*, e12404. 10.1002/jev2.12404.

62.   Ekström, K., Crescitelli, R., Pétursson, H.I., Johansson, J., Lässer, C., and Bagge, R.O.
(2022). Characterization of surface markers on extracellular vesicles isolated from
lymphatic exudate from patients with breast cancer. BMC cancer *22*, 1-17.

63.   Maleki, F., Ovens, K., Hogan, D.J., and Kusalik, A.J. (2020). Gene Set Analysis:
Challenges, Opportunities, and Future Research. Front Genet *11*, 654.
10.3389/fgene.2020.00654.

64.   Yang, X., Chung, J.Y., Rai, U., and Esumi, N. (2018). Cadherins in the retinal pigment
epithelium (RPE) revisited: P-cadherin is the highly dominant cadherin expressed in
human and mouse RPE in vivo. PLoS One *13*, e0191279. 10.1371/journal.pone.0191279.

65.   Ghosh, S., Shang, P., Terasaki, H., Stepicheva, N., Hose, S., Yazdankhah, M., Weiss, J.,
Sakamoto, T., Bhutto, I.A., Xia, S., Zigler, J.S., Jr., et al. (2018). A Role for betaA3/A1-
Crystallin in Type 2 EMT of RPE Cells Occurring in Dry Age-Related Macular
Degeneration. Invest Ophthalmol Vis Sci *59*, AMD104-AMD113. 10.1167/iovs.18-
24132.

66.   Kanan, Y., Wicker, L.D., Al-Ubaidi, M.R., Mandal, N.A., and Kasus-Jacobi, A. (2008).
Retinol dehydrogenases RDH11 and RDH12 in the mouse retina: expression levels
during development and regulation by oxidative stress. Invest Ophthalmol Vis Sci *49*,
1071-1078. 10.1167/iovs.07-1207.

67.   Kiser, P.D., Golczak, M., Maeda, A., and Palczewski, K. (2012). Key enzymes of the
retinoid (visual) cycle in vertebrate retina. Biochim Biophys Acta *1821*, 137-151.
10.1016/j.bbalip.2011.03.005.

68. Liu, Y.D., Huang, S.S., Li, M., Lek, M., Song, D.Y., Tan, D.D., Chen, X.Y., Zhang, H.,
Liu, J.Y., Chang, X.Z., and Xiong, H. (2022). A new phenotype of syndromic retinitis
pigmentosa with myopathy is caused by mutations in retinol dehydrogenase 11. Clin
Genet *101*, 448-453. 10.1111/cge.14108.

69. Sun, H.J., Zhang, F.F., Xiao, Q., Xu, J., and Zhu, L.J. (2021). lncRNA MEG3, Acting as
a ceRNA, Modulates RPE Differentiation Through the miR-7-5p/Pax6 Axis. Biochem
Genet *59*, 1617-1630. 10.1007/s10528-021-10072-9.

70. Pan, J., and Zhao, L. (2021). Long non-coding RNA histone deacetylase 4 antisense RNA
1 (HDAC4-AS1) inhibits HDAC4 expression in human ARPE-19 cells with hypoxic
stress. Bioengineered *12*, 2228-2237. 10.1080/21655979.2021.1933821.

71. Chen, X., Jiang, C., Qin, B., Liu, G., Ji, J., Sun, X., Xu, M., Ding, S., Zhu, M., Huang,
G., Yan, B., et al. (2017). LncRNA ZNF503-AS1 promotes RPE differentiation by
downregulating ZNF503 expression. Cell Death Dis *8*, e3046. 10.1038/cddis.2017.382.

72. Li, F., Wen, X., Zhang, H., and Fan, X. (2016). Novel Insights into the Role of Long
Noncoding RNA in Ocular Diseases. Int J Mol Sci *17*, 478. 10.3390/ijms17040478.

73. Jovičić, A., and Gitler, A.D. (2017). Distinct repertoires of microRNAs present in mouse
astrocytes compared to astrocyte-secreted exosomes. PloS one *12*, e0171418.

74. Otake, K., Adachi-Tominari, K., Nagai, H., Saito, M., Sano, O., Hirozane, Y., and Iwata,
H. (2021). Quantitative comparison of the mRNA content of human iPSC-derived motor
neurons and their extracellular vesicles. FEBS open bio *11*, 494-506.

75. de Candia, P., Torri, A., Gorletta, T., Fedeli, M., Bulgheroni, E., Cheroni, C., Marabita,
F., Crosti, M., Moro, M., and Pariani, E. (2013). Intracellular modulation, extracellular
disposal and serum increase of MiR-150 mark lymphocyte activation. PloS one *8*,
e75348.

76. Rashed, M.H., Kanlikilicer, P., Rodriguez-Aguayo, C., Pichler, M., Bayraktar, R.,
Bayraktar, E., Ivan, C., Filant, J., Silva, A., and Aslan, B. (2017). Exosomal miR-940
maintains SRC-mediated oncogenic activity in cancer cells: a possible role for exosomal
disposal of tumor suppressor miRNAs. Oncotarget *8*, 20145.

77. Valadi, H., Ekström, K., Bossios, A., Sjöstrand, M., Lee, J.J., and Lötvall, J.O. (2007).
Exosome-mediated transfer of mRNAs and microRNAs is a novel mechanism of genetic
exchange between cells. Nature cell biology *9*, 654-659.

78. Montecalvo, A., Larregina, A.T., Shufesky, W.J., Beer Stolz, D., Sullivan, M.L.,
Karlsson, J.M., Baty, C.J., Gibson, G.A., Erdos, G., and Wang, Z. (2012). Mechanism of
transfer of functional microRNAs between mouse dendritic cells via exosomes. Blood,
The Journal of the American Society of Hematology *119*, 756-766.

79. Zhou, J., Flores-Bellver, M., Pan, J., Benito-Martin, A., Shi, C., Onwumere, O., Mighty,
J., Qian, J., Zhong, X., and Hogue, T. (2021). Human retinal organoids release
extracellular vesicles that regulate gene expression in target human retinal progenitor
cells. Scientific reports *11*, 21128.

80.   Zhang, Y., Liu, D., Chen, X., Li, J., Li, L., Bian, Z., Sun, F., Lu, J., Yin, Y., Cai, X., Sun,
Q., et al. (2010). Secreted monocytic miR-150 enhances targeted endothelial cell
migration. Mol Cell *39*, 133-144. 10.1016/j.molcel.2010.06.010.

81.   Pegtel, D.M., Cosmopoulos, K., Thorley-Lawson, D.A., van Eijndhoven, M.A.,
Hopmans, E.S., Lindenberg, J.L., de Gruijl, T.D., Würdinger, T., and Middeldorp, J.M.
(2010). Functional delivery of viral miRNAs via exosomes. Proc Natl Acad Sci U S A
107, 6328-6333. 10.1073/pnas.0914843107.

82.   Hurley, J.B. (2021). Retina Metabolism and Metabolism in the Pigmented Epithelium: A
Busy Intersection. Annu Rev Vis Sci *7*, 665-692. 10.1146/annurev-vision-100419-
115156.

83.   Strauss, O. (2005). The retinal pigment epithelium in visual function. Physiol Rev *85*,
845-881. 10.1152/physrev.00021.2004.

84.   Wang, D., Eraslan, B., Wieland, T., Hallstrom, B., Hopf, T., Zolg, D.P., Zecha, J.,
Asplund, A., Li, L.H., Meng, C., Frejno, M., et al. (2019). A deep proteome and
transcriptome abundance atlas of 29 healthy human tissues. Mol Syst Biol *15*, e8503.
10.15252/msb.20188503.

85.   Chen, L., Ge, C., Feng, X., Fu, H., Wang, S., Zhu, J., Linghu, E., and Zheng, X. (2022).
Identification of Combinations of Plasma lncRNAs and mRNAs as Potential Biomarkers
for Precursor Lesions and Early Gastric Cancer. J Oncol *2022*, 1458320.
10.1155/2022/1458320.

86.   Zhu, Z., Chen, Z., Wang, M., Zhang, M., Chen, Y., Yang, X., Zhou, C., Liu, Y., Hong,
L., and Zhang, L. (2022). Detection of plasma exosomal miRNA-205 as a biomarker for
early diagnosis and an adjuvant indicator of ovarian cancer staging. J Ovarian Res *15*, 27.
10.1186/s13048-022-00961-x.

87.   Ratnasari, N., Lestari, P., Renovaldi, D., Raditya Ningsih, J., Qoriansas, N., Wardana, T.,
Hakim, S., Signa Aini Gumilas, N., Indrarti, F., Triwikatmani, C., Bayupurnama, P., et al.
(2022). Potential plasma biomarkers: miRNA-29c, miRNA-21, and miRNA-155 in
clinical progression of Hepatocellular Carcinoma patients. PLoS One *17*, e0263298.
10.1371/journal.pone.0263298.

88.   Nimir, M., Ma, Y., Jeffreys, S.A., Opperman, T., Young, F., Khan, T., Ding, P., Chua,
W., Balakrishnar, B., Cooper, A., De Souza, P., et al. (2019). Detection of AR-V7 in
Liquid Biopsies of Castrate Resistant Prostate Cancer Patients: A Comparison of AR-V7
Analysis in Circulating Tumor Cells, Circulating Tumor RNA and Exosomes. Cells *8*.
10.3390/cells8070688.

89.   Botto, C., Rucli, M., Tekinsoy, M.D., Pulman, J., Sahel, J.A., and Dalkara, D. (2022).
Early and late stage gene therapy interventions for inherited retinal degenerations. Prog
Retin Eye Res *86*, 100975. 10.1016/j.preteyeres.2021.100975.